

Systematic study of Z_c^+ family from quark model's perspective

Chengrong Deng^{a,b*}, Jialun Ping^{c†}, Hongxia Huang^{c‡}, and Fan Wang^{d§}

^a*School of Science, Chongqing Jiaotong University, Chongqing 400074, P.R. China*

^b*Department of Physics and Astronomy, University of California, Los Angeles 90095, USA*

^c*Department of Physics, Nanjing Normal University, Nanjing 210097, P.R. China and*

^d*Department of Physics, Nanjing University, Nanjing 210093, P.R. China*

Inspired by the present experimental status of charged charmonium-like states Z_c^+ , the tetraquark states $[cu][\bar{c}\bar{d}]$ are systematically studied in a color flux-tube model with a multi-body confinement potential. The investigation indicates that charged charmonium-like states $Z_c^+(3900)$ or $Z_c^+(3885)$, $Z_c^+(3930)$, $Z_c^+(4020)$ or $Z_c^+(4025)$, $Z_1^+(4050)$, $Z_2^+(4250)$, and $Z_c^+(4200)$ can be uniformly described as tetraquark states $[cu][\bar{c}\bar{d}]$ with the quantum numbers $n^{2S+1}L_J$ and J^P of 1^3S_1 and 1^+ , 2^3S_1 and 1^+ , 1^5S_2 and 2^+ , 1^3P_1 and 1^- , 1^5D_1 and 1^+ , and 1^3D_1 and 1^+ , respectively. The predicted lowest charged tetraquark state $[cu][\bar{c}\bar{d}]$ with 0^+ and 1^1S_0 has a energy of 3780 ± 10 MeV in the model. The tetraquark states are compact three-dimensional spatial configurations similar to a rugby ball, the higher orbital angular momentum L between the diquark $[cu]$ and antidiquark $[\bar{c}\bar{d}]$, the more prolate of the states. The multibody color flux-tube, a collective degree of freedom, plays an important role in the formation of those charge tetraquark states. However, the two heavier charged states $Z_c^+(4430)$ and $Z_c^+(4475)$ can not be explained as tetraquark states $[cu][\bar{c}\bar{d}]$ in this model approach.

PACS numbers: 14.20.Pt, 12.40.-y

I. INTRODUCTION

Quantum chromodynamics (QCD) is nowadays widely accepted as the fundamental theory to describe hadrons and their interactions. Conventional hadrons are composed of either a valence quark q and an antiquark \bar{q} (mesons) or three valence quarks (baryons) on top of the sea of $q\bar{q}$ pairs and gluons. One of the long standing challenges in hadron physics is to establish and classify genuine multi-quark states other than conventional hadrons because multi-quark states may contain more information about the low-energy QCD than that of conventional hadrons. In the past several years, a charged charmoniumlike Z_c^+ family, $Z_c^+(4430)$, $Z_1^+(4050)$, $Z_2^+(4250)$, $Z_c^+(3900)$, $Z_c^+(3885)$, $Z_c^+(3930)$, $Z_c^+(4020)$, $Z_c^+(4025)$, $Z_c^+(4475)$ and $Z_c^+(4200)$, has been successively observed by experimental Collaborations [1–9]. Obviously, those charged charmoniumlike states go beyond conventional $c\bar{c}$ -meson picture and prefer to tetraquark systems $c\bar{c}u\bar{d}$ due to carrying one charge, which provides a good place for testing various phenomenological research methods of hadron physics. On the theoretical side, a large amount of work has been devoted to describe the internal structure of these charged states, which has been related to meson-meson molecules [11, 12], diquark-antidiquark states [13], hadrocharmonium or Born-Oppenheimer tetraquarks [14], coupled channel cusps [15], and kinematic effects [16].

A systematic understanding of the internal structure of

these charged states may not only contribute to provide new insights to the strong interaction dynamics of multi-quark systems and low-energy QCD but also provide important information on future experimental search for the missing higher orbital excitations in the Z_c^+ family. This is the goal of the present work. In our approach, a phenomenological model, color flux-tube model with a multi-body confinement potential instead of a two-body one in traditional quark model, is employed to explore the properties of excited charged tetraquark states $c\bar{c}u\bar{d}$ systematically. The model has been successfully applied to the ground states of charged tetraquark states $[Qq][\bar{Q}'\bar{q}']$ ($Q, Q' = c, b$ and $q, q' = u, d, s$) in our previous work [17].

This work is organized as follows: the color flux-tube model and the model parameters are given in Sec. II. The numerical results and discussions of the charged tetraquark states are presented in Sec. III. A brief summary is given in the last section.

II. COLOR FLUX-TUBE MODEL AND PARAMETERS

The details of the color flux-tube model basing on traditional quark models and lattice QCD picture can be found in our previous work [18], the prominent characteristics of the model are just presented here. The model Hamiltonian for the state $[cu][\bar{c}\bar{d}]$ is given as follows,

$$H_4 = \sum_{i=1}^4 \left(m_i + \frac{\mathbf{p}_i^2}{2m_i} \right) - T_C + \sum_{i>j}^4 V_{ij} + V_{min}^C + V_{min}^{C,SL},$$

$$V_{ij} = V_{ij}^B + V_{ij}^{B,SL} + V_{ij}^\sigma + V_{ij}^{\sigma,SL} + V_{ij}^G + V_{ij}^{G,SL}. \quad (1)$$

*crdeng@cqjtu.edu.cn

†jlping@njnu.edu.cn, corresponding author

‡hxhuang@njnu.edu.cn

§fgwang@chenwang.nju.edu.cn

T_c is the center-of-mass kinetic energy of the state, \mathbf{p}_i and m_i are the momentum and mass of the i -th quark (antiquark), respectively. The codes of the quarks (antiquarks) c and u (\bar{c} and \bar{d}) are assumed to be 1 and 2 (3 and 4), respectively, their positions are denoted as \mathbf{r}_1 and \mathbf{r}_2 (\mathbf{r}_3 and \mathbf{r}_4).

The quadratic confinement potential, which is believed to be flavor independent, of the tetraquark state with a diquark-antidiquark structure has the following form,

$$V^C = K [(\mathbf{r}_1 - \mathbf{y}_{12})^2 + (\mathbf{r}_2 - \mathbf{y}_{12})^2 + (\mathbf{r}_3 - \mathbf{y}_{34})^2 + (\mathbf{r}_4 - \mathbf{y}_{34})^2 + \kappa_d(\mathbf{y}_{12} - \mathbf{y}_{34})^2], \quad (2)$$

The positions \mathbf{y}_{12} and \mathbf{y}_{34} are the junctions of two Y-shaped color flux-tube structures. The parameter K is the stiffness of a three-dimension flux-tube, $\kappa_d K$ is other compound color flux-tube stiffness. The relative stiffness parameter κ_d of the compound flux-tube is [19]

$$\kappa_d = \frac{C_d}{C_3}, \quad (3)$$

where C_d is the eigenvalue of the Casimir operator associated with the $SU(3)$ color representation d at either end of the color flux-tube, such as $C_3 = \frac{4}{3}$, $C_6 = \frac{10}{3}$, and $C_8 = 3$.

The minimum of the confinement potential V_{min}^C can be obtained by taking the variation of V^C with respect to \mathbf{y}_{12} and \mathbf{y}_{34} , and it can be expressed as

$$V_{min}^C = K \left(\mathbf{R}_1^2 + \mathbf{R}_2^2 + \frac{\kappa_d}{1 + \kappa_d} \mathbf{R}_3^2 \right), \quad (4)$$

The canonical coordinates \mathbf{R}_i have the following forms,

$$\begin{aligned} \mathbf{R}_1 &= \frac{1}{\sqrt{2}}(\mathbf{r}_1 - \mathbf{r}_2), \quad \mathbf{R}_2 = \frac{1}{\sqrt{2}}(\mathbf{r}_3 - \mathbf{r}_4), \\ \mathbf{R}_3 &= \frac{1}{\sqrt{4}}(\mathbf{r}_1 + \mathbf{r}_2 - \mathbf{r}_3 - \mathbf{r}_4), \\ \mathbf{R}_4 &= \frac{1}{\sqrt{4}}(\mathbf{r}_1 + \mathbf{r}_2 + \mathbf{r}_3 + \mathbf{r}_4). \end{aligned} \quad (5)$$

The use of V_{min}^C can be understood here that the gluon field readjusts immediately to its minimal configuration. It is worth emphasizing that the confinement V_{min}^C is a multi-body interaction in a multiquark state rather than the sum of many pairwise confinement interactions,

$$V^C = \sum_{i < j} \lambda_i \cdot \lambda_j r_{ij}^n, \quad (6)$$

in Isgur-Karl quark model and chiral quark model with $n = 1$ or 2.

The central parts of one-boson-exchange V_{ij}^B and σ -meson exchange V_{ij}^σ only occur between u and \bar{d} , and that of one-gluon-exchange V_{ij}^G is universal. V_{ij}^B , V_{ij}^σ and V_{ij}^G

take their standard forms and are listed in the following,

$$V_{ij}^B = V_{ij}^\pi \sum_{k=1}^3 \mathbf{F}_i^k \mathbf{F}_j^k + V_{ij}^K \sum_{k=4}^7 \mathbf{F}_i^k \mathbf{F}_j^k + V_{ij}^\eta (\mathbf{F}_i^8 \mathbf{F}_j^8 \cos \theta_P - \sin \theta_P), \quad (7)$$

$$V_{ij}^\chi = \frac{g_{ch}^2}{4\pi} \frac{m_\chi^3}{12m_i m_j} \frac{\Lambda_\chi^2}{\Lambda_\chi^2 - m_\chi^2} \sigma_i \cdot \sigma_j \times \left(Y(m_\chi r_{ij}) - \frac{\Lambda_\chi^3}{m_\chi^3} Y(\Lambda_\chi r_{ij}) \right), \quad (8)$$

$$V_{ij}^G = \frac{\alpha_s}{4} \lambda_i^c \cdot \lambda_j^c \left(\frac{1}{r_{ij}} - \frac{2\pi\delta(\mathbf{r}_{ij})\sigma_i \cdot \sigma_j}{3m_i m_j} \right), \quad (9)$$

$$V_{ij}^\sigma = -\frac{g_{ch}^2}{4\pi} \frac{\Lambda_\sigma^2 m_\sigma}{\Lambda_\sigma^2 - m_\sigma^2} \left(Y(m_\sigma r_{ij}) - \frac{\Lambda_\sigma}{m_\sigma} Y(\Lambda_\sigma r_{ij}) \right). \quad (10)$$

Where χ stands for π , K and η , $Y(x) = e^{-x}/x$. The symbols \mathbf{F} , $\boldsymbol{\lambda}$ and $\boldsymbol{\sigma}$ are the flavor $SU(3)$, color $SU(3)$ Gell-Mann and spin $SU(2)$ Pauli matrices, respectively. θ_P is the mixing angle between η_1 and η_8 to give the physical η meson. $g_{ch}^2/4\pi$ is the chiral coupling constant. α_s is the running strong coupling constant and takes the following form [20],

$$\alpha_s(\mu_{ij}) = \frac{\alpha_0}{\ln((\mu_{ij}^2 + \mu_0^2)/\Lambda_0^2)}, \quad (11)$$

where μ_{ij} is the reduced mass of two interacting particles q_i (or \bar{q}_i) and q_j (or \bar{q}_j). Λ_0 , α_0 and μ_0 are model parameters. The function $\delta(\mathbf{r}_{ij})$ in V_{ij}^G should be regularized [22],

$$\delta(\mathbf{r}_{ij}) = \frac{1}{4\pi r_{ij} r_0^2(\mu_{ij})} e^{-r_{ij}/r_0(\mu_{ij})}, \quad (12)$$

where $r_0(\mu_{ij}) = \hat{r}_0/\mu_{ij}$, \hat{r}_0 is a model parameter.

The diquark $[cu]$ and antidiquark $[\bar{c}\bar{d}]$ can be considered as compound objects \bar{Q} and Q with no internal orbital excitation, and the angular excitation L are assumed to occur only between Q and \bar{Q} in the present work and the parity of the state $[cu][\bar{c}\bar{d}]$ is therefore simply related to L as $P = (-1)^L$. In this way, the state $[cu][\bar{c}\bar{d}]$ has lower energy than that of the states with additional internal orbital excitation in Q and \bar{Q} . In order to facilitate numerical calculations, the spin-orbit interactions are approximately assumed to just take place between compound objects \bar{Q} and Q , which is consistence with the work [21]. The related interactions can be presented as follows

$$V_{12,34}^{G,LS} \approx \frac{\alpha_s}{4} \lambda_{12}^{\bar{c}} \cdot \lambda_{34}^c \frac{1}{8M_{12}M_{34}} \frac{3}{X^3} \mathbf{L} \cdot \mathbf{S}, \quad (13)$$

$$V_{12,34}^{\sigma,LS} \approx -\frac{g_{ch}^2}{4\pi} \frac{\Lambda_\sigma^2}{\Lambda_\sigma^2 - m_\sigma^2} \frac{m_\sigma^3}{2M_{12}M_{34}} \mathbf{L} \cdot \mathbf{S} \times \left(G(m_\sigma X) - \frac{\Lambda_\sigma^3}{m_\sigma^3} G(\Lambda_\sigma X) \right), \quad (14)$$

$$V_{12,34}^{C,LS} \approx \frac{K}{8M_{12}M_{34}} \frac{\kappa_d}{1 + \kappa_d} \mathbf{L} \cdot \mathbf{S}. \quad (15)$$

where $M_{12} = M_{34} = m_c + m_{u,d}$, $G(x) = Y(x)(\frac{1}{x} + \frac{1}{x^2})$, and S stands for the total spin angular momentum of the tetraquark state $[cu][\bar{c}\bar{d}]$.

The model parameters are determined as follows. The mass parameters m_π , m_K and m_η in the interaction V_{ij}^B are taken their experimental values, namely, $m_\pi = 0.7$ fm $^{-1}$, $m_K = 2.51$ fm $^{-1}$ and $m_\eta = 2.77$ fm $^{-1}$. The cutoff parameters take the values, $\Lambda_\pi = \Lambda_\sigma = 4.20$ fm $^{-1}$ and $\Lambda_\eta = \Lambda_K = 5.20$ fm $^{-1}$, the mixing angle $\theta_P = -15^\circ$ [20]. The mass parameter m_σ in the interaction V_{ij}^σ is determined through the PCAC relation $m_\sigma^2 \approx m_\pi^2 + 4m_{u,d}^2$ [23], $m_{u,d} = 280$ MeV and $m_\sigma = 2.92$ fm $^{-1}$. The chiral coupling constant g_{ch} is determined from the πNN coupling constant through

$$\frac{g_{ch}^2}{4\pi} = \left(\frac{3}{5}\right)^2 \frac{g_{\pi NN}^2}{4\pi} \frac{m_{u,d}^2}{m_N^2} = 0.43. \quad (16)$$

The other adjustable parameters and their errors are determined by fitting the masses of the ground states of mesons using Minuit program, which are shown in Table I. The mass spectrum of the ground states of mesons, which is listed in Table II, can be obtained by solving the two-body Schrödinger equation

$$(H_2 - E_2)\Phi_{IJ}^{Meson} = 0. \quad (17)$$

The mass error of mesons ΔE_2 introduced by the parameter uncertainty Δx_i can be calculated by the formula of error propagation,

$$\Delta H_2 = \sum_{i=1}^8 \left| \frac{\partial H_2}{\partial x_i} \right| \Delta x_i, \quad (18)$$

$$\Delta E_2 \approx \langle \Phi_{IJ}^{Meson} | \Delta H_2 | \Phi_{IJ}^{Meson} \rangle. \quad (19)$$

where x_i and Δx_i represent the i -th adjustable parameter and its error, respectively, which are listed in Table I.

III. NUMERICAL RESULTS AND DISCUSSIONS

Within the framework of the diquark-antidiquark configuration, the wave function of the state $[cu][\bar{c}\bar{d}]$ can be written as a sum of the following direct products of color χ_c , isospin η_I , spin η_s and spatial ϕ terms,

$$\begin{aligned} \Phi_{IM_I J M_J}^{[cu][\bar{c}\bar{d}]} &= \sum_{\alpha} \xi_{\alpha} \left[\left[\left[\phi_{l_a m_a}^G(\mathbf{r}) \chi_{s_a} \right]_{j_a}^{[cu]} \left[\phi_{l_b m_b}^G(\mathbf{R}) \right. \right. \right. \\ &\quad \times \left. \left. \chi_{s_b} \right]_{j_b}^{[\bar{c}\bar{d}]} \right]_{J_{ab}}^{[cu][\bar{c}\bar{d}]} F_{LM}(\mathbf{X}) \Big]_{J M_J}^{[cu][\bar{c}\bar{d}]} \\ &\quad \times \left[\eta_{I_a}^{[cu]} \eta_{I_b}^{[\bar{c}\bar{d}]} \right]_{I M_I}^{[cu][\bar{c}\bar{d}]} \left[\chi_{c_a}^{[cu]} \chi_{c_b}^{[\bar{c}\bar{d}]} \right]_{C W_C}^{[cu][\bar{c}\bar{d}]} \end{aligned} \quad (20)$$

In which \mathbf{r} , \mathbf{R} and \mathbf{X} are relative spatial coordinates,

$$\begin{aligned} \mathbf{r} &= \mathbf{r}_1 - \mathbf{r}_2, \quad \mathbf{R} = \mathbf{r}_3 - \mathbf{r}_4 \\ \mathbf{X} &= \frac{m_1 \mathbf{r}_1 + m_2 \mathbf{r}_2}{m_1 + m_2} - \frac{m_3 \mathbf{r}_3 + m_4 \mathbf{r}_4}{m_3 + m_4}. \end{aligned} \quad (21)$$

TABLE I: Adjustable model parameters. (units: $m_s, m_c, m_b, \mu_0, \Lambda_0$, MeV; K , MeV·fm $^{-2}$; r_0 , MeV·fm; α_0 , dimensionless)

Parameters	x_i	Δx_i	Parameters	x_i	Δx_i
m_s	511.78	0.228	α_0	4.554	0.018
m_c	1601.7	0.441	Λ_0	9.173	0.175
m_b	4936.2	0.451	μ_0	0.0004	0.540
K	217.50	0.230	r_0	35.06	0.156

TABLE II: Ground state meson spectra, unit in MeV.

States	E_2	ΔE_2	PDG	States	E_2	ΔE_2	PDG
π	142	26	139	η_c	2912	5	2980
K	492	20	496	J/Ψ	3102	4	3097
ρ	826	4	775	B^0	5259	5	5280
ω	780	4	783	B^*	5301	4	5325
K^*	974	4	892	B_s^0	5377	5	5366
ϕ	1112	4	1020	B_s^*	5430	4	5416
D^\pm	1867	8	1880	B_c	6261	7	6277
D^*	2002	4	2007	B_c^*	6357	4	...
D_s^\pm	1972	9	1968	η_b	9441	8	9391
D_s^*	2140	4	2112	$\Upsilon(1S)$	9546	5	9460

The other details of the construction of the wave function can be found in our previous work [17]. Subsequently, the converged numerical results can be obtained by solving the four-body Schrödinger equation

$$(H_4 - E_4)\Phi_{IM_I J M_J}^{[cu][\bar{c}\bar{d}]} = 0. \quad (22)$$

with the Rayleigh-Ritz variational principle.

The energies $E_4 \pm \Delta E_4$ of the charged states $[cu][\bar{c}\bar{d}]$ with $n^{2S+1}L_J$ and J^P under the assumptions of $S = 0, \dots, 2$ and $L = 0, \dots, 3$ are systematically calculated and presented in Table III. The mass error of the states ΔE_4 can be calculated just as ΔE_2 , they are around several MeV except for that of the state 1^1S_0 . The spin-orbit interactions are extremely weak, less than 2 MeV, therefore the energies of excited states with the same L and S but different J are almost degenerate, see the energies of the excited states with 1^5D_0 , 1^5D_1 , 1^5D_2 , 1^5D_3 and 1^5D_4 in Table III, which is consistent with the conclusion of the work [24]. Other spin-related interactions are stronger and bring about a larger energy difference than spin-orbital interactions, especially for the ground states with 1^1S_0 , 1^3S_1 and 1^5S_2 . The energy difference among excited states mainly comes from the kinetic energy and confinement potential, which are proportional to the relative orbital excitation L . However, the relative kinetic energy between two clusters $[cu]$ and $[\bar{c}\bar{d}]$ is inversely proportional to $\langle \mathbf{X}^2 \rangle$ while confinement potential is proportional to $\langle \mathbf{X}^2 \rangle$ so that they compete each other to reach an optimum balance.

The rms $\langle \mathbf{r}^2 \rangle^{\frac{1}{2}}$, $\langle \mathbf{R}^2 \rangle^{\frac{1}{2}}$ and $\langle \mathbf{X}^2 \rangle^{\frac{1}{2}}$ stand for the size of the diquark $[cu]$, the antidiquark $[\bar{c}\bar{d}]$ and the distance between the two clusters, respectively, which are also calculated and listed in Table III. One can find that the

TABLE III: The energy $E_4 + \Delta E_4$ and rms $\langle \mathbf{r}^2 \rangle^{\frac{1}{2}}$, $\langle \mathbf{R}^2 \rangle^{\frac{1}{2}}$ and $\langle \mathbf{X}^2 \rangle^{\frac{1}{2}}$ of charged tetraquark states $[cu][\bar{c}\bar{d}]$ with J^P and $n^{2S+1}L_J$, unit of energy: MeV and unit of rms: fm.

J^P	0^+	0^-	0^+	1^+	1^+	1^-	1^-	1^-	1^+	1^+
$n^{2S+1}L_J$	1^1S_0	1^3P_0	1^5D_0	1^3S_1	2^3S_1	1^1P_1	1^3P_1	1^5P_1	1^3D_1	1^5D_1
$E_4 \pm \Delta E_4$	3782 ± 12	4097 ± 8	4274 ± 7	3858 ± 10	3950 ± 10	4075 ± 8	4097 ± 8	4153 ± 7	4235 ± 7	4273 ± 7
$\langle \mathbf{r}^2 \rangle^{\frac{1}{2}}$	0.85	0.96	1.01	0.90	0.92	0.94	0.96	1.00	0.98	1.01
$\langle \mathbf{R}^2 \rangle^{\frac{1}{2}}$	0.85	0.96	1.01	0.90	0.92	0.94	0.96	1.00	0.98	1.01
$\langle \mathbf{X}^2 \rangle^{\frac{1}{2}}$	0.42	0.85	1.12	0.48	0.66	0.85	0.85	0.92	1.10	1.12
J^P	1^-	2^+	2^-	2^-	2^+	2^+	2^+	2^-	2^-	3^-
$n^{2S+1}L_J$	1^5F_1	1^5S_2	1^3P_2	1^5P_2	1^1D_2	1^3D_2	1^5D_2	1^3F_2	1^5F_2	1^5P_3
$E_4 \pm \Delta E_4$	4387 ± 7	4001 ± 7	4096 ± 8	4152 ± 7	4212 ± 8	4235 ± 7	4273 ± 7	4354 ± 7	4387 ± 7	4150 ± 7
$\langle \mathbf{r}^2 \rangle^{\frac{1}{2}}$	1.02	1.03	0.96	1.00	0.95	0.98	1.01	0.99	1.02	1.00
$\langle \mathbf{R}^2 \rangle^{\frac{1}{2}}$	1.02	1.03	0.96	1.00	0.95	0.98	1.01	0.99	1.02	1.00
$\langle \mathbf{X}^2 \rangle^{\frac{1}{2}}$	1.30	0.57	0.85	0.92	1.09	1.10	1.12	1.30	1.30	0.92
J^P	3^+	3^+	3^-	3^-	3^-	4^+	4^-	4^-	5^-	
$n^{2S+1}L_J$	1^3D_3	1^5D_3	1^1F_3	1^3F_3	1^5F_3	1^5D_4	1^3F_4	1^5F_4	1^5F_5	
$E_4 \pm \Delta E_4$	4234 ± 7	4272 ± 7	4332 ± 8	4353 ± 7	4386 ± 7	4274 ± 7	4353 ± 7	4387 ± 7	4387 ± 7	
$\langle \mathbf{r}^2 \rangle^{\frac{1}{2}}$	0.98	1.01	0.96	0.99	1.02	1.01	0.99	1.02	1.02	
$\langle \mathbf{R}^2 \rangle^{\frac{1}{2}}$	0.98	1.01	0.96	0.99	1.02	1.01	0.99	1.02	1.02	
$\langle \mathbf{X}^2 \rangle^{\frac{1}{2}}$	1.10	1.12	1.27	1.30	1.30	1.12	1.30	1.30	1.30	

diquark $[cu]$ and antidiquark $[\bar{c}\bar{d}]$ share the same size in every Z_c^+ state. The sizes of the diquark $[cu]$ and antidiquark $[\bar{c}\bar{d}]$ ($\langle \mathbf{r}^2 \rangle^{\frac{1}{2}}$ and $\langle \mathbf{R}^2 \rangle^{\frac{1}{2}}$) are mainly determined by the total spin S , the relative orbital excitation L of the states has a minor effect on them. However, the sizes do not vary largely with the total spin S , especially for higher orbital excited states. So the diquark $[cu]$ and antidiquark $[\bar{c}\bar{d}]$ are rather rigid against the rotation. For examples, the sizes of the two groups 1^1S_0 - 1^3S_1 - 1^5S_2 and 1^1F_3 - 1^3F_3 - 1^5F_3 changes gradually with the total spin S , 0.85-0.90-1.03 fm and 0.96-0.99-1.02 fm, respectively. And the sizes of the two groups 1^1S_0 - 1^1P_1 - 1^1D_2 - 1^1F_3 and 1^3S_1 - 1^3P_1 - 1^3D_1 - 1^3F_2 vary slightly with relative orbital excitation L , 0.85-0.94-0.95-0.96 fm and 0.90-0.96-0.98-0.99 fm, respectively. On the contrary, the distance between the diquark $[cu]$ and antidiquark $[\bar{c}\bar{d}]$ ($\langle \mathbf{X}^2 \rangle^{\frac{1}{2}}$) changes remarkably with the relative orbital excitation L between the two clusters and is irrelevant to the total spin of the system, see the sizes of 1^3S_1 - 1^3P_1 - 1^3D_1 - 1^3F_2 and 1^1S_0 - 1^3S_1 - 1^5S_2 in Table III. The sizes of the diquark $[cu]$, antidiquark $[\bar{c}\bar{d}]$ and the distance between the two clusters are helpful to understand the changing tendency of energies of charged states Z_c^+ with quantum numbers S and L .

In order to make clear the spatial configuration of charged states $[cu][\bar{c}\bar{d}]$, the distances in four states between any two particles are given in Table IV. The ground state (1^1S_0 and 1^+) of charged tetraquark $[cu][\bar{c}\bar{d}]$ possesses a three-dimensional spatial configuration due to the competition of the confinement and the kinetic energy of the systems [17], which is similar to a rugby ball. The diquark $[cu]$ and antidiquark $[\bar{c}\bar{d}]$ in the ground state have a large overlap because of the small $\langle \mathbf{X}^2 \rangle^{\frac{1}{2}}$, so the pic-

TABLE IV: The average distances $\langle \mathbf{r}_{ij}^2 \rangle^{\frac{1}{2}}$ of the states $[cu][\bar{c}\bar{d}]$ with 1^1S_0 , 1^1P_1 , 1^1D_2 , and 1^1F_3 , $\mathbf{r}_{ij} = \mathbf{r}_i - \mathbf{r}_j$, unit in fm.

$n^{2S+1}L_J$	$\langle \mathbf{r}_{12}^2 \rangle^{\frac{1}{2}}$	$\langle \mathbf{r}_{34}^2 \rangle^{\frac{1}{2}}$	$\langle \mathbf{r}_{24}^2 \rangle^{\frac{1}{2}}$	$\langle \mathbf{r}_{13}^2 \rangle^{\frac{1}{2}}$	$\langle \mathbf{r}_{14}^2 \rangle^{\frac{1}{2}}$	$\langle \mathbf{r}_{23}^2 \rangle^{\frac{1}{2}}$	$\langle \mathbf{X}^2 \rangle^{\frac{1}{2}}$
1^1S_0	0.85	0.85	1.11	0.46	0.85	0.85	0.42
1^1P_1	0.94	0.94	1.41	0.87	1.17	1.17	0.85
1^1D_2	0.95	0.95	1.59	1.11	1.37	1.37	1.09
1^1F_3	0.96	0.96	1.72	1.28	1.52	1.52	1.27

ture of the diquark or antidiquark is not extremely distinct. However, all distances except for the sizes of the diquark and antidiquark ($\langle \mathbf{r}_{12}^2 \rangle^{\frac{1}{2}}$ and $\langle \mathbf{r}_{34}^2 \rangle^{\frac{1}{2}}$) evidently augment with the increasing of the orbital angular momentum L in the excited states, see Table IV, which means that the picture of the diquark or antidiquark is more and more clear with the raising of the orbital angular momentum L . The spatial configuration of the excited states is still similar to a rugby ball, the higher orbital angular momentum L , the more prolate of the shape of the excited states. The multibody color flux-tube basing on lattice QCD picture, a collective degree of freedom, plays an important role in the formation of these charged tetraquark states, it should therefore be the dynamical mechanism of the tetraquark systems.

Next, let us turn to argue the properties of the charged states Z_c^+ observed in experiments and their possible candidates in the color flux-tube model, which are presented in Table IV. It can be seen from the Table IV that the spin and parity of the Z_c^+ (3900) are still unclear up to now. The Z_c^+ (3900) may correspond to the same state as the Z_c^+ (3885) with 1^+ [3]. The charged state $[cu][\bar{c}\bar{d}]$ with 1^+ and 1^3S_1 has a mass of 3858 ± 10 MeV in the

TABLE V: Z_c^+ states observed in experiments and their possible candidates in the color flux-tube model.

State	Experiment		Model			
	Mass, MeV	J^P	Mass, MeV	J^P	$n^{2S+1}L_J$	
$Z_1^+(4050)$ [1]	4051_{-14-41}^{+14+20}	??	4075 ± 8	1^-	1^1P_1	
$Z_2^+(4250)$ [1]	$4248_{-29-35}^{+44+180}$??	4273 ± 7	1^+	1^5D_1	
$Z_c^+(3900)$ [2]	$3899.0_{-3.6-4.9}^{+3.6+4.9}$??	3858 ± 10	1^+	1^3S_1	
$Z_c^+(3885)$ [3]	$3883.9_{-1.5-4.2}^{+1.5+4.2}$	1^+	3858 ± 10	1^+	1^3S_1	
$Z_c^+(3930)$ [4]	3929_{-5-2}^{+5+2}	1^+	3950 ± 10	1^+	2^3S_1	
$Z_c^+(4025)$ [5]	$4026.3_{-2.6-3.7}^{+2.6+3.7}$??	4001 ± 7	2^+	1^5S_2	
$Z_c^+(4020)$ [6]	$4022.9_{-0.8-2.7}^{+0.8+2.7}$??	4001 ± 7	2^+	1^5S_2	
$Z_c^+(4200)$ [7]	4196_{-29-13}^{+36+17}	??	4235 ± 7	1^+	1^3D_1	
$Z_c^+(4475)$ [8]	4475_{-22-11}^{+22+28}	1^+	
$Z_c^+(4430)$ [9]	4433_{-2-4}^{+2+4}	1^+	

color flux-tube model, which is very close to those of the two charged states $Z_c^+(3885)$ and $Z_c^+(3900)$. It can not be excluded that the main component of $Z_c^+(3885)$ and $Z_c^+(3900)$ is the state $[cu][\bar{c}\bar{d}]$ with 1^+ and 1^3S_1 , which is supported by many theoretical work [13]. The radial excited state 2^3S_1 has a mass of 3950 ± 10 MeV, which is extremely close to that of $Z_c^+(3930)$. It is possible to identify $Z_c^+(3930)$ as the tetraquark state $[cu][\bar{c}\bar{d}]$ with 1^+ and 2^3S_1 . In other words, the $Z_c^+(3930)$ is the first radial excited state of the $Z_c^+(3900)$ in the color flux-tube model. The pair $Z_c^+(4020)$ and $Z_c^+(4025)$ show up with a similar mass (slightly above $D^*\bar{D}^*$ threshold). They might therefore be the same resonance, their spin and parity are unclear. QCD sum rule identified the $Z_c^+(4020)$ and $Z_c^+(4025)$ as a tetraquark state $[cu][\bar{c}\bar{d}]$ with 1^+ [25], the same approach also favored a tetraquark state but with different quantum numbers 2^+ and $5S_2$ [26]. In our calculations, the nearest tetraquark state $[cu][\bar{c}\bar{d}]$ to the $Z_c^+(4020)$ or $Z_c^+(4025)$ occupies quantum numbers 2^+ and 1^5S_2 . The tetraquark states $[cu][\bar{c}\bar{d}]$ with 1^- and 1^1P_1 and 1^+ and 1^5D_1 have the energies of 4075 ± 8 MeV and 4273 ± 7 MeV, respectively, which are consistent with those of $Z_1^+(4050)$ and $Z_2^+(4250)$. So the two states may be assigned as the tetraquark states $[cu][\bar{c}\bar{d}]$ with 1^- and 1^1P_1 and 1^+ and 1^5D_1 , respectively, in the color flux-tube model. The newly observed $Z_c^+(4200)$ prefers 1^+ , which can be described as the tetraquark state $[cu][\bar{c}\bar{d}]$ with 1^+ and 1^3D_1 in the color flux-tube model. The study of the three-point function sum rules on this state gives a support to the tetraquark interpretation [27]. Of cause, it seems difficult to rule out other two possibilities of 2^+ and 1^1D_2 and 2^+ and 1^3D_2 in the model. The $Z_c^+(4430)$ is the first charged state, the J^P of the state is determined unambiguously to be 1^+ , the $Z_c^+(4475)$ favors the spin-parity 1^+ over other hypotheses [8]. Due to the heavy of the diquark and antidiquark, the energy of radial excitation between the diquark and antidiquark is too small to make the tetraquark state $[cu][\bar{c}\bar{d}]$ reach the energy above 4400 MeV. The internal excited states of the diquark and/or antidiquark are needed to account for the

heavy charged states, which is left for future. The alternative configuration for the two states may be meson-meson molecular states, which is suggested by several theoretical methods [12].

From the above analysis and Table III, we can see that the most of the low energy theoretical states can be matched with the experimental ones. One of the exception is the state with 0^+ and 1^1S_0 , which has a mass of 3780 ± 10 MeV. The experimental search of the η_c -like charged state will give a crucial test of the present approach. Our calculation also suggests that there are two negative parity states around 4100 MeV. More information on the the states around this energy is expected.

The model assignments of the Z_c^+ states are completed just in term of the proximity to the experimental masses, the more stringent check of the assignment is to study the decay properties of the states. These charged states should eventually decay into several color singlet mesons due to their high energy. In the course of the decay, the color flux-tube structure should break down first which leads to the collapses of the three-dimension spatial configuration, and then through the recombination of the color flux tubes the particles of decay products formed. The decay widths of the charged states $[cu][\bar{c}\bar{d}]$ are determined by the transition probability of the breakdown and recombination of color flux tubes. The calculations are in progress. This decay mechanism is similar to compound nucleus decay and therefore should induce a resonance, which we called it as “color confined, multi-quark resonance” state before [28].

IV. SUMMARY

The charged tetraquark states $[cu][\bar{c}\bar{d}]$ are systematically investigated from the perspective of the color flux-tube model with a four-body confinement potential. The investigation demonstrates that the charged charmoniumlike states $Z_c^+(3900)$ or $Z_c^+(3885)$, $Z_c^+(3930)$, $Z_c^+(4020)$ or $Z_c^+(4025)$, $Z_1^+(4050)$, $Z_2^+(4250)$, and $Z_c^+(4200)$ can be uniformly identified as tetraquark states $[cu][\bar{c}\bar{d}]$ with the quantum numbers 1^3S_1 and 1^+ , 2^3S_1 and 1^+ , 1^5S_2 and 2^+ , 1^3P_1 and 1^- , 1^5D_1 and 1^+ , and 1^3D_1 and 1^+ , respectively, in the color flux-tube model. The predicted lowest charged tetraquark state $[cu][\bar{c}\bar{d}]$ with 0^+ and 1^1S_0 has a mass of 3780 ± 10 MeV in the color flux-tube model. The model predictions would shed light on looking for possible charmoniumlike charged states in the future at the BESIII, LHCb and Belle-II. They favor three-dimensional spatial structures which is similar to a rugby ball, the higher orbital angular momentum L , the more prolate of the shape of the states. Those charged charmoniumlike states may be so-called “color confined, multi-quark resonance”. However, the two heavier charged states $Z_c^+(4430)$ and $Z_c^+(4475)$ can not be described as tetraquark states $[cu][\bar{c}\bar{d}]$ in the color flux-tube model.

The multibody color flux-tube is a collective degree of

freedom, which acts as a dynamical mechanism and plays an important role in the formation and decay of those compact states. Just as colorful organic world because of chemical bonds, multiquark hadron world should be various due to the diversity of color flux-tube structure. The well-defined charged state $Z_c^+(3900)$ and dibaryon d^* resonance have been opening the gate of abundant multiquark hadronic world.

Acknowledgments

C.R. Deng thanks Huanzhong Huang for hosting Deng's visit to UCLA. This research is partly supported by the NSFC under contracts Nos. 11305274, 11175088, 11035006, 11205091, and the Chongqing Natural Science Foundation under Project No. cstc2013jcyjA00014.

-
- [1] R. Mizuk, et al. (Belle Collaboration), Phys. Rev. D **78**, 072004 (2008).
 - [2] M. Ablikim et al. (BES III Collaboration), Phys. Rev. Lett. **110**, 252001 (2013); Z. Liu et al. (Belle Collaboration), Phys. Rev. Lett. **110**, 252002 (2013); T. Xiao, S. Dobbs, A. Tomaradze, and K.K. Seth, Phys. Lett. B **727**, 366 (2013).
 - [3] M. Ablikim et al. (BES III Collaboration), Phys. Rev. Lett. **112**, 022001 (2014);
 - [4] S. Uehara et al. (Belle Collaboration), Phys. Rev. Lett. **96**, 082003 (2006).
 - [5] M. Ablikim et al. (BES III Collaboration), Phys. Rev. Lett. **112**, 132001 (2014).
 - [6] M. Ablikim et al. (BES III Collaboration), Phys. Rev. Lett. **111**, 242001 (2013).
 - [7] K. Chilikin et al. (Belle Collaboration), arXiv: 1408.6457 [hep-ex].
 - [8] K. Chilikin et al. [Belle Collaboration], Phys. Rev. D **88**, 074026 (2013). R. Aaij et al. (LHCb Collaboration), Phys. Rev. Lett. **112**, 222002 (2014).
 - [9] S.K. Choi et al. (Belle Collaboration), Phys. Rev. Lett. **100**, 142001 (2008); R. Aaij et al. (LHCb Collaboration), Phys. Rev. Lett. **112**, 222002 (2014).
 - [10] N. Brambilla, et al. arXiv: 1404.3723 [hep-ph]. A. Esposito, A.L. Guerrieri, F. Piccinini, A. Pilloni, and A.D. Polosa, Int. J. Mod. Phys. A **30**, 1530002 (2015); S.L. Olsen, Front. Phys. **10**, 101401 (2015), and therein.
 - [11] Q. Wang, C. Hanhart, and Q. Zhao, Phys. Rev. Lett. **111**, 132003 (2013);
 - [12] T. Barnes, F.E. Close and E.S. Swanson, Phys. Rev. D **91**, 014004 (2015); J. He, arXiv: 1410.8645v2 [hep-ph].
 - [13] L. Maiani, V. Riquer, R. Faccini, F. Piccinini, A. Pilloni, and A.D. Polosa, Phys. Rev. D **87**, 111102 (2013); J.M. Dias, F.S. Navarra, M. Nielsen, and C. M. Zanetti, Phys. Rev. D **88**, 016004 (2013); E. Braaten, arXiv: 1305.6905 [hep-ph]; Z.G. Wang and T. Huang, Phys. Rev. D **89**, 054019 (2014).
 - [14] I.V. Danilkin, V.D. Orlovsky, and Y.A. Simonov, Phys. Rev. D **85**, 034012 (2012); N. Mahajan, arXiv: 1304.1301 [hep-ph].
 - [15] E.S. Swanson, Phys. Rev. D **91**, 034009 (2015).
 - [16] F.K. Guo, C. Hanhart, Q. Wang, and Q. Zhao, Phys. Rev. D **91**, 051504 (2015).
 - [17] C.R. Deng, J.L. Ping, H.X. Huang, and F. Wang, Phys. Rev. D **90**, 054009 (2014).
 - [18] C.R. Deng, J.L. Ping, F. Wang, and T. Goldman, Phys. Rev. D **82**, 074001 (2010); C.R. Deng, J.L. Ping, H. Wang, P. Zhou, and F. Wang, Phys. Rev. D **86**, 114035 (2012); C.R. Deng, J.L. Ping, Y.C. Yang, and F. Wang, Phys. Rev. D **86**, 014008 (2012); C.R. Deng, J.L. Ping, Y.C. Yang, and F. Wang, Phys. Rev. D **88**, 074007 (2013);
 - [19] G.S. Bail, Phys. Rev. D **62**, 114503 (2000); C. Semay, Eur. Phys. J. A **22**, 353 (2004); N. Cardoso, M. Cardoso, and P. Bicudo, Phys. Lett. B **710**, 343 (2012).
 - [20] J. Vijande, F. Fernandez, and A. Valcarce, J. Phys. G **31**, 481 (2005).
 - [21] M. Cleven, F.K. Guo, C. Hanhart, Q. Wang and Q. Zhao, arXiv: 1505.01771 [hep-ph].
 - [22] J. Weinstein and N. Isgur, Phys. Rev. D **27**, 588 (1983).
 - [23] M.D. Scadron, Phys. Rev. D **26**, 239 (1982).
 - [24] S. Patel, M. Shah, and P.C. Vinodkumara, Eur. Phys. J. A **50**, 131 (2014).
 - [25] Z.G. Wang, Eur. Phys. J. C **74**, 2874 (2014).
 - [26] K.P. Khemchandani, A. M. Torres, M. Nielsen, and F.S. Navarra, Phys. Rev. D **89**, 014029 (2014).
 - [27] W. Chen, T.G. Steele, H.X. Chen, and S.L. Zhu, arXiv: 1501.03863 [hep-ph].
 - [28] F. Wang, J.L. Ping, H.R. Pang, and L.Z. Chen, Nucl. Phys. A **790**, 493c (2007).