

# Pentaquarks and resonances in the $pJ/\psi$ spectrum

V.V. Anisovich<sup>+</sup>, M.A. Matveev<sup>+</sup>, J. Nyiri<sup>\*</sup>, A.V. Sarantsev<sup>+\diamond</sup>,  
A.N. Semenova<sup>+</sup>

<sup>+</sup>*National Research Centre "Kurchatov Institute": Petersburg Nuclear Physics Institute,  
Gatchina, 188300, Russia*

<sup>\diamond</sup> *Helmholtz-Institut für Strahlen- und Kernphysik, Universität Bonn, Germany*

<sup>\*</sup>*Institute for Particle and Nuclear Physics, Wigner RCP, Budapest 1121, Hungary*

## Abstract

We consider exotic baryons with hidden charm as antiquark-diquark-diquark composite systems. Spin and isospin structure of such exotic states is given and masses are estimated. The data for production of pentaquarks in the reaction  $\Lambda_b \rightarrow K^- pJ/\psi$  are discussed. We suggest that the narrow peak in  $pJ/\psi$  spectra at 4450 MeV is antiquark-diquark-diquark state with negative parity,  $5/2^-$  (4450), while the broad bump  $3/2^+$  (4380) is the result of rescatterings in the  $(pJ/\psi)$ -channel. Positions of other pentaquarks with negative parity are estimated.

PACS: 12.40.Yx, 12.39.-x, 14.40.Lb

## 1 Introduction

Data for the decay  $\Lambda_b^0 \rightarrow pJ/\psi K^-$  [1] provide a definite argument for the existence of a pentaquark, a baryon system in  $pJ/\psi$  spectra which has the following quark content:

$$P_{\bar{c}cuud}^+ = \bar{c}(cuud). \quad (1)$$

In terms of the quark-diquark states it can be presented as a three-body systems:

$$\bar{c}(cuud) = \bar{c} \cdot (cu) \cdot (ud) + \text{permutations of the } u, d \text{ quarks}. \quad (2)$$

A diquark is a color triplet member, similar to a quark, and the right-hand side of eq. (2) presents a three-body system with a color structure similar to that in low-lying baryons. It is reasonable to suppose that we face similar color forces in three quark and antiquark-diquark-diquark systems as well. Following this idea we perform a classification of such baryon states and give estimations of their masses. Estimation of diquark masses is given in [2] where diquark-antidiquark states are studied.

The notion of the diquark was introduced by Gell-Mann [3]. Diquarks were discussed for baryon states during a long time, see pioneering papers [4, 5, 6, 7, 8] and conference presentations [9, 10, 11]. The systematization of baryons in terms of the quark-diquark states is presented in [12, 13]. The application of the diquarks to exotic mesons in the sector of heavy diquarks was discussed by Maiani *et al.* [14], Voloshin [15], Ali *et al.* [16].

Pentaquarks built of light-light and light-heavy diquarks present natural extension of multi-quark schemes studied in the last decade for mesons [17, 18] and baryons [19, 20].

## 2 Pentaquarks

In color space we write for the pentaquark:

$$P_{\bar{c}cuud}^+ = \epsilon_{\alpha\beta\gamma} \bar{c}^\alpha (cu)^\beta (ud)^\gamma + \text{permutations of the } u, d \text{ quarks} \quad , \quad (3)$$

$$(cu)^\beta = \epsilon^{\beta\beta'\gamma'} c_{\beta'} u_{\gamma'}, \quad (ud)^\gamma = \epsilon^{\gamma\beta''\gamma''} u_{\beta''} d_{\gamma''},$$

where  $\alpha, \beta, \gamma$  refer to color indices.

We discuss a scheme in which the exotic baryon states are formed by standard QCD-motivated interactions (gluonic exchanges, confinement forces) but in addition with diquarks as constituents.

### 2.1 Spin structure of the pentaquarks

We work with two diquarks: scalar  $S$  and axial-vector  $A$ . In terms of these diquarks the color-flavor wave function of pentaquark reads:

$$P_{\bar{c}(cq)\cdot(q'q'')} = \bar{c}^\alpha \cdot \epsilon_{\alpha\beta\gamma} \begin{vmatrix} S_{(cq)}^\beta \\ A_{(cq)}^\beta \end{vmatrix} \cdot \begin{vmatrix} S_{(q'q'')}^\gamma \\ A_{(q'q'')}^\gamma \end{vmatrix} \quad (4)$$

We have six diquark-diquark states:

$$P_{\bar{c}(cq)\cdot(q'q'')} = \bar{c}^\alpha \cdot \begin{vmatrix} (S_{(cq)} S_{(q'q'')})^\alpha(0^+) \\ (S_{(cq)} A_{(q'q'')})^\alpha(1^+) \\ (A_{(cq)} S_{(q'q'')})^\alpha(1^+) \\ (A_{(cq)} A_{(q'q'')})^\alpha(0^+) \\ (A_{(cq)} A_{(q'q'')})^\alpha(1^+) \\ (A_{(cq)} A_{(q'q'')})^\alpha(2^+) \end{vmatrix} \quad (5)$$

with the spin-parity numbers  $J^P = 0^+, 1^+, 2^+$ .

### 2.2 Isospin structure of the diquarks

We face the following isospin states for the diquarks:

$$\begin{aligned} S_{(cq)}(I_d = 1/2, J_d = 0), & \quad A_{(cq)}(I_d = 1/2, J_d = 1), \\ S_{(q'q'')}(I_d = 0, J_d = 0), & \quad A_{(q'q'')}(I_d = 1, J_d = 1). \end{aligned} \quad (6)$$

So, the isospin-spin sector of the pentaquarks  $P(I, J^P)$  reads:

$$P_{\bar{c} \cdot (cq) \cdot (q'q'')} = \bar{c}^\alpha \cdot \begin{vmatrix} (S_{(cq)} S_{(q'q'')})^\alpha (0^+) \\ (S_{(cq)} A_{(q'q'')})^\alpha (1^+) \\ (A_{(cq)} S_{(q'q'')})^\alpha (1^+) \\ (A_{(cq)} A_{(q'q'')})^\alpha (0^+) \\ (A_{(cq)} A_{(q'q'')})^\alpha (1^+) \\ (A_{(cq)} A_{(q'q'')})^\alpha (2^+) \end{vmatrix} \Rightarrow \begin{matrix} P(\frac{1}{2}, \frac{1}{2}^-) \\ P(\frac{1}{2}, \frac{1}{2}^-), P(\frac{1}{2}, \frac{3}{2}^-), P(\frac{3}{2}, \frac{1}{2}^-), P(\frac{3}{2}, \frac{3}{2}^-), \\ P(\frac{1}{2}, \frac{1}{2}^-), P(\frac{1}{2}, \frac{3}{2}^-) \\ P(\frac{1}{2}, \frac{1}{2}^-), P(\frac{3}{2}, \frac{1}{2}^-) \\ P(\frac{1}{2}, \frac{1}{2}^-), P(\frac{1}{2}, \frac{3}{2}^-), P(\frac{3}{2}, \frac{1}{2}^-), P(\frac{3}{2}, \frac{3}{2}^-) \\ P(\frac{1}{2}, \frac{3}{2}^-), P(\frac{1}{2}, \frac{5}{2}^-), P(\frac{3}{2}, \frac{3}{2}^-), P(\frac{3}{2}, \frac{5}{2}^-) \end{matrix} \quad (7)$$

## 2.3 Pentaquarks, their masses and spins

Mass formula for a diquark-diquark system  $(cq) \cdot (q'q'')$ , is accepted to be the same as for the diquark-antidiquark one [2]. We write

$$M_{(cq) \cdot (q'q'')} = m_{(cq)} + m_{(q'q'')} + J_{(cq) \cdot (q'q'')} (J_{(cq) \cdot (q'q'')} + 1) \Delta \quad (8)$$

with the parameters which were determined in ref. [2]:

$$\Delta = 70 \pm 10 \text{ MeV}, \quad (9)$$

$$m_{S_{(cq)}} = 2000 \pm 50 \text{ MeV}, \quad m_{A_{(cq)}} = 2050 \pm 50 \text{ MeV}.$$

Here we concentrate our attention on non-strange diquarks only, for an expansion of the results to the strange quark sector we need  $m_{S_{(cs)}}$  and  $m_{A_{(cs)}}$ .

For the  $(q'q'')$  diquarks we accept masses found in the analysis of baryons [12, 13]:

$$m_{S_{(q'q'')}} = 650 \pm 50 \text{ MeV}, \quad m_{A_{(q'q'')}} = 750 \pm 50 \text{ MeV}. \quad (10)$$

The mass of the constituent antiquark  $\bar{c}$  is equal to [21, 22]:

$$m_c = 1300 \pm 50 \text{ MeV}. \quad (11)$$

Within these masses (9),(10),(11) we roughly estimate the masses of the pentaquarks as:

$$M_{\bar{c} \cdot (cq) \cdot (q'q'')} \simeq m_{\bar{c}} + M_{(cq) \cdot (q'q'')}. \quad (12)$$

Correspondingly we write a set of the low-laying pentaquark states:

$I = 1/2$	$I = 3/2$
$P_{\bar{c} S_{(cq)} S_{(q'q'')}}^{(\frac{1}{2}, \frac{1}{2}^-)} (3800),$	$P_{\bar{c} S_{(cq)} A_{(q'q'')}}^{(\frac{3}{2}, \frac{1}{2}^-)} (4190), P_{\bar{c} S_{(cq)} A_{(q'q'')}}^{(\frac{3}{2}, \frac{3}{2}^-)} (4190),$
$P_{\bar{c} S_{(cq)} A_{(q'q'')}}^{(\frac{1}{2}, \frac{1}{2}^-)} (4190), P_{\bar{c} S_{(cq)} A_{(q'q'')}}^{(\frac{1}{2}, \frac{3}{2}^-)} (4190),$	$P_{\bar{c} A_{(cq)} S_{(q'q'')}}^{(\frac{1}{2}, \frac{1}{2}^-)} (4140), P_{\bar{c} A_{(cq)} S_{(q'q'')}}^{(\frac{1}{2}, \frac{3}{2}^-)} (4140),$
$P_{\bar{c} A_{(cq)} S_{(q'q'')}}^{(\frac{1}{2}, \frac{1}{2}^-)} (4100),$	$P_{\bar{c} A_{(cq)} A_{(q'q'')}}^{(\frac{3}{2}, \frac{1}{2}^-)} (4100),$
$P_{\bar{c} A_{(cq)} A_{(q'q'')}}^{(\frac{1}{2}, \frac{1}{2}^-)} (4240), P_{\bar{c} A_{(cq)} A_{(q'q'')}}^{(\frac{1}{2}, \frac{3}{2}^-)} (4240),$	$P_{\bar{c} A_{(cq)} A_{(q'q'')}}^{(\frac{3}{2}, \frac{1}{2}^-)} (4240), P_{\bar{c} A_{(cq)} A_{(q'q'')}}^{(\frac{3}{2}, \frac{3}{2}^-)} (4240),$
$P_{\bar{c} A_{(cq)} A_{(q'q'')}}^{(\frac{1}{2}, \frac{3}{2}^-)} (4520), P_{\bar{c} A_{(cq)} A_{(q'q'')}}^{(\frac{1}{2}, \frac{5}{2}^-)} (4520),$	$P_{\bar{c} A_{(cq)} A_{(q'q'')}}^{(\frac{3}{2}, \frac{3}{2}^-)} (4520), P_{\bar{c} A_{(cq)} A_{(q'q'')}}^{(\frac{3}{2}, \frac{5}{2}^-)} (4520).$

(13)

Masses are given in MeV units, an uncertainty in the determination of masses is of the order of  $\pm 150$  MeV.

### 3 Discussion

The state  $P_{\bar{c}A_{(cq)}A_{(q'q'')}}^{(\frac{1}{2}, \frac{5}{2}^-)}(4520 \pm 150)$  is a good candidate to be a state which was observed in [1]:  $5/2^?(4450 \pm 4)$  with a width of  $\Gamma = 39 \pm 24$  MeV. Then the broad state,  $3/2^?(3380 \pm 38)$  with  $\Gamma = 205 \pm 94$  MeV also observed in the  $pJ/\psi$  spectrum, is a positive parity state,  $3/2^+$ .

An opposite classification of states is suggested in [23]:  $3/2^?(3380 \pm 38) \rightarrow 3/2^-(S\text{-wave pentaquark})$  and  $5/2^?(4450 \pm 4) \rightarrow 5/2^+(P\text{-wave pentaquark})$ .

We suppose that the broad bump in the  $3/2^+$ -wave is the result of rescatterings in  $pJ/\psi$ -channel, for example, such as  $\Lambda_b \rightarrow \Lambda(1520)J/\psi \rightarrow K^-(pJ/\psi)$ .

In the suggested scheme the mass interval (4040 – 4500) MeV should contain additionally several resonances with  $J^P = 1/2^-, 3/2^-$ . Some of them may dominantly decay into  $p\eta_c$ -channel, concerning mainly the  $1/2^-$  states.

The masses of pentaquarks with strange diquarks ( $cs$ ), see eq.(9), are estimated quite similar to the non-strange ones.

In conclusion, the scheme of the low-lying pentaquark is suggested on the basis of the study of the tetraquark states [2]. We are convinced that the crossing studies of exotic mesons and baryons give a correct way for investigation of these topics.

The work was supported by grants RSGSS-4801.2012.2., RFBR-13-02-00425 and RSCF-14-22-00281 .

### References

- [1] The LHCb collaboration, arXiv:1507.03414v1 [hep-ex].
- [2] V.V. Anisovich, M.A. Matveev, A.V. Sarantsev, A.N. Semenova *Exotic mesons with hidden charm as diquark-antidiquark states*, arXiv:1507.07232 [hep-ph] (2015).
- [3] M. Gell-Mann, Phys. Lett. **8**, 214 (1964).
- [4] M. Ida and R. Kobayashi, Progr. Theor. Phys. **36**, 846 (1966).
- [5] D.B Lichtenberg and L.J. Tassie, Phys. Rev. **155**, 1601 (1967).
- [6] S. Ono, Progr. Theor. Phys. **48**, 964 (1972).
- [7] V.V. Anisovich, Pis'ma ZhETF **21**, 382 (1975) [JETP Lett. **21**, 174 (1975)];  
V.V. Anisovich, P.E. Volkovitski, and V.I. Povzun, ZhETF **70**, 1613 (1976) [Sov. Phys. JETP **43**, 841 (1976)].
- [8] A. Schmidt and R. Blankenbeckler, Phys. Rev. **D16**, 1318 (1977).
- [9] M. Anselmino and E. Predazzi, eds., *Proceedings of the Workshop on Diquarks*, World Scientific, Singapore (1989).
- [10] K. Goeke, P.Kroll, and H.R. Petry, eds., *Proceedings of the Workshop on Quark Cluster Dynamics* (1992).

- [11] M. Anselmino and E. Predazzi, eds., *Proceedings of the Workshop on Diquarks II*, World Scientific, Singapore (1992).
- [12] A.V. Anisovich, V.V. Anisovich, M.A. Matveev, V.A. Nikonov, A.V. Sarantsev and T.O. Vulfs, Int. J. Mod. Phys. **A25**, 2965 (2010);  
Phys. Atom. Nucl. **74**, 418 (2011) [Phys. Atom. Nucl. **74**, 418 (2011)].
- [13] A.V. Anisovich, V.V. Anisovich, M.A. Matveev, V.A. Nikonov, J. Nyiri, A.V. Sarantsev *Three-particle physics and dispersion relation theory*, World Scientific, Singapore (2013).
- [14] L.Maiani, F.Piccinini, A.D.Polosa, V.Riquer, Phys. Rev. D**71**, 014028 (2005).
- [15] M.B. Voloshin, Phys. Rev. D**84**, 031502 (2011).
- [16] A. Ali, C. Hambrock, W.Wang, Phys. Rev. D**85**, 054011 (2012).
- [17] S. Weinberg, Phys. Rev. Lett. **110**, 261601 (2013).
- [18] S.J. Brodsky, D.S. Hwang and R.F. Lebed Phys. Rev. Lett. **113**, 112001 (2014).
- [19] R.L. Jaffe and F. Wilchek, Phys. Rev. Lett. **91**, 232003 (2003).
- [20] G.C. Rossi and G. Veneziano, Phys. Lett.B**597**, 338 (2004).
- [21] V.V. Anisovich, L.G. Dakhno, M.A. Matveev, V.A. Nikonov and A.V. Sarantsev, Yad. Fiz. **70**, 392 (2007) [Phys. Atom. Nucl. **70**, 364 (2007)].
- [22] A.V. Manohar, C.T. Sachrajda, Phys. Rev. D**66**, 010001-271 (2002).
- [23] L. Maiani, A.D. Polosa, V. Riquer, arXiv:1507.04980v1 [hep-ph] .