

Polarization correlated photons from a positively charged quantum dot

Y. Cao,^{1,2} A. J. Bennett,^{1,*} I. Farrer,³ D. A. Ritchie,³ and A. J. Shields¹

¹*Toshiba Research Europe Limited, Cambridge Research Laboratory,
208 Science Park, Milton Road, Cambridge, CB4 0GZ, U. K.*

²*Department of Physics, Imperial College London,
Prince Consort Road, London SW7 2AZ, U. K.*

³*Cavendish Laboratory, Cambridge University,
J. J. Thomson Avenue, Cambridge, CB3 0HE, U. K.*

(Dated: October 3, 2018)

Polarized cross-correlation spectroscopy on a quantum dot charged with a single hole shows the sequential emission of photons with common circular polarization. This effect is visible without magnetic field, but becomes more pronounced as the field along the quantization axis is increased. We interpret the data in terms of electron dephasing in the X^+ state caused by the Overhauser field of nuclei in the dot. We predict the correlation timescale can be increased by accelerating the emission rate with cavity-QED.

PACS numbers:

Spins in quantum dots (QDs) provide a promising platform for manipulating and storing quantum information in the solid state. Optical measurements have demonstrated spin preparation [1, 2] coherent spin control [3] and electron-spin – photon entanglement [4, 5]. There are also proposals for achieving photon entanglement [6] and non-destructive measurement of photons [7] using charged QDs. However, the time evolution of the carrier spin is unavoidably affected by the $10^4 - 10^5$ nuclei in the dot, all with non-zero spin. One example of the utility of the electron-nuclear interaction is its use in spin pumping the hole into the spin down state in zero external field, by pumping on the spin up state of X^+ [2]. From a fundamental point of view then, the hyperfine interaction provides an interesting system for manipulating a mesoscopic nuclear ensemble and observing its dynamics.

It has now been established that the electron-nuclear hyperfine interaction is dominated by the contact interaction and is isotropic in a QD [8]. The dynamics of this interaction manifests itself in studies of polarized photo-luminescence [9, 10]. Contrastingly, the hole's p -like wave-function has a node at each nucleus leaving the dipole-dipole interaction between the hole and nuclear spins to dominate [11]. This interaction has a strength one order of magnitude below that of the electron [12, 13]. Thus, there has been interest in using the hole-spin as quantum bit with reduced decoherence. Direct measurements of the hole spin relaxation time in a vertical magnetic field, T_1^h , have shown it is hundreds of microseconds [2, 14]. Without applied magnetic field some experiments suggest the hole spin T_1^h time is 13 ns [8]. Several studies have now estimated the hole dephasing time, T_2^h in magnetic field is approximately $1 \mu s$ [3, 17].

We study here the emission from the X^+ state in a dot deterministically charged with a hole using a diode

(Figure 1a). We show photons from this transition display polarization correlation over a timescale one order of magnitude greater than the radiative lifetime when excited by a linearly polarized laser. This time is short relative to some reports of the hole spin polarization lifetime [14] and we show that this is a result of dephasing caused by the electron when the system is excited. We investigate the magnitude of the effect as a function of applied external magnetic field and radiative lifetime.




FIG. 1: (a) Schematic band-structure of the hole-charging diode when the dot contains a single hole. (b) Energy level diagram for the X^+ at zero magnetic field. (c) Spectrum under quasi-resonant excitation at 1.2 V. (d) Emission pattern for an X^+ transition showing $\beta = 0.092$.

The X^+ consists of two holes in a singlet $S = 0$ state and an electron. The zero net-hole-spin ensures the electron-hole anisotropic exchange interaction is absent. In zero magnetic field, the X^+ eigenstates are degenerate and labeled by the spin of the electron (Figure 1b) [15, 16]. Radiative decay from X^+ to the single hole ground state occurs with a change in total angular momentum ± 1 , the polarization of the photon being corre-

*Electronic address: anthony.bennett@crl.toshiba.co.uk

lated with the initial and final spin state. In a dot where the hole is “heavy” ($m_j = \pm 3/2$) only the vertical transitions in Figure 1b are allowed: the detection of a left-handed photon (L) photon ensures the decay occurred by the left hand transition on Figure 1b.

The strain, shape anisotropy and inversion asymmetry in InGaAs/GaAs QDs ensures the optically active transition has a mixed heavy- ($m_j = \pm 3/2$) and light- ($m_j = \pm 1/2$) hole character [18, 19]. The state is given by $\phi_{\pm} = (|\pm 3/2\rangle + \beta|\mp 1/2\rangle)/\sqrt{1 + \beta^2}$, which we denote \uparrow and \downarrow . Recombination of an electron and a mixed heavy/light-hole now results in elliptically polarized photons from X^+ , and β may be determined from the emission pattern (Figure 1d) [18]). Within the sample studied β values from 0.02-0.20 are typical, and for the data shown here $\beta = 0.092$ (Figure 1d).




FIG. 2: (a) Apparatus to measure the polarization correlation from a dot. (b) Circular co- and cross- polarized emission correlation from X^+ at zero external field, $g_{co}(t)$ (black) and $g_{cross}(t)$ (red), respectively. Extracted degree of polarization correlation, $C(t)$ (green). (c) The same measurement made in the linear detection basis.

The diode for controlled charging has a 20 nm GaAs tunnel barrier between the dot and p -contact (Figure 1a). A 75% AlGaAs barrier on the n -side prevents electron charging, so the X^+ dominates at 1.2-1.3 V. Emission from X^+ at an energy of 1349.2 meV is excited quasi-resonantly by a linearly polarized laser at 1317.2 meV. This excitation scheme equally excites both transitions in Figure 1b and the absence of spin pumping ensures there is no build up of nuclear spin polarization, but it does not populate other carrier combinations such as neutral and negatively charged excitons (Figure 1c).

The experiment is shown in Figure 2a. After filtering, the emission is passed to polarization-maintaining fibre-optics which enable 4 simultaneous measurements of correlation in a basis selected by the quarter-wave plate

(QWP) and the polarizing coupler (PBS).

Figure 2b presents correlations recorded at zero external field at an excitation power $\times 10$ below saturation, in the circular basis. Comparing the sum of the two co- and cross-polarized measurements ($g_{co}(t)$ in black and $g_{cross}(t)$ in red, respectively), we see a clear difference. Note that both $g_{co}(t)$ and $g_{cross}(t)$ show a reduced signal within ~ 1 ns of zero-time delay due to the anti-bunched nature of the light. Outside the central ~ 1 ns there is an enhanced probability of the source emitting two photons of the same circular polarization over the case of emitting photon of opposite circular polarization.

The degree of polarization correlation, $C(t)$ is defined as $C(t) = (g_{co}(t) - g_{cross}(t))/(g_{co}(t) + g_{cross}(t))$ from which a least squares fit with a function $C(t) = C_0 * \exp(-|t|/\tau_d)$ extracts the polarization correlation at zero delay C_0 and the timescale, τ_d . Empirically, this function is a good fit to $C(t)$ (Figure 2b). $C_0 = 0.33 \pm 0.01$ and the decay time of the correlation $\tau_d = 9.0 \pm 0.4$ ns. In contrast, measurements in the linear-polarization basis (H/V) show an absence of polarization correlation (Figure 2c).

Non-zero heavy-light hole mixing is an obvious source of reduced polarization correlation. Taking the heavy: light hole oscillator strength of 3:1 [18, 19] we see that recombination of a ϕ_+ hole and an electron in the X^+ level leads to an elliptical photon with state $\propto \sqrt{3}|L\rangle + \beta|R\rangle$, and a ϕ_- hole. Conversely, decay involving a ϕ_- hole and an electron leads to a $\propto \sqrt{3}|R\rangle + \beta|L\rangle$ photon. The measurement in Fig. 2b is in the circular basis so detection of a left-handed photon implies the decay came from ϕ_+ with $3/(3 + \beta^2)$ probability. In the absence of dephasing in the upper or lower states, this reduces the probability of obtaining sequential left-left photo-detections to $(9 + \beta^4)/(3 + \beta^2)^2$. For this QD $\beta = 0.092$, so the probability of co-polarized photon emission is reduced to 0.994. This is higher than we have measured, so we conclude an additional factor must be included.




FIG. 3: A measurement of the timescale of correlation at zero external magnetic field, τ_d , as a function of the normalized intensity of the source. The data is fitted with an inverse relationship.

In fact, the data can be explained by the fast dephasing of the electron spin in the upper state, which dom-

inates any dephasing from the hole spin. A coincidence detection event arises as follows: the transition emits a photon that is detected with circular polarization and the hole spin is left in the corresponding state. Some time later, the system is re-excited to the upper state, where electron spin dephasing occurs during the radiative lifetime of the X^+ state, following which a second photon is emitted from the spontaneous decay. These two photons form a single coincidence in Figure 2b. We stress that our model implicitly assumes the hole spin lifetime is greater than the measured τ_d , though we envisage that future experiments that reduce the effect of electron dephasing it will be necessary to include the contribution of the hole.

Our studies provide four pieces of evidence electron spin dephasing is the factor limiting the polarization correlation. Firstly, the degree of polarization correlation from the X^+ observed in Figure 2b is $1/3$. When excited, the unpaired electron spin evolves through a hyperfine interaction with the nuclei. Only those nuclear field fluctuations in the two directions perpendicular to the spin will cause precession. If this precession is faster than the radiative lifetime of the upper state, its effect is to randomize the spin. The electron spin parallel to the nuclear field is preserved. Thus, the mean spin projection along \mathbf{z} is reduced to $1/3$. Secondly, the timescale over which polarization correlation is observed is inversely proportional to pump rate, as shown in Figure 3. This cannot be explained by dephasing occurring in the ground state. The increased excitation increases the number of times the system is excited between photon detection events, and this increases the rate at which the polarization correlation is lost. Thirdly, there is no polarization correlation in the linear basis (Figure 2c). This is consistent with a dephasing of the electron spin state in a time faster than the X^+ radiative lifetime. Finally, we shall show that change in C_0 with magnetic field is explained by the dynamics of the electron spin.




FIG. 4: Energy level diagram for the X^+ transition with finite z -magnetic field. (b) The degree of correlation, C_0 as a function of external Faraday magnetic field (black points). The calculated variation with field is shown as a red line. (c) Variation in the timescale of correlation, τ_d .

We next discuss the application of a Faraday magnetic

field, which removes the degeneracy of the upper and lower states, shown in Figure 4a. The net field experienced by the spins is the sum of the external field B_{ext} and the internal nuclear field, B_N . This stabilizes the electron spin along \mathbf{z} and causes it to precess about the sum of the two fields, which is predominantly along the \mathbf{z} when the $|B_{ext}| > B_N$. Thus, the application of vertical field increases the value of C_0 as shown in Figure 4b. Figure 4c plots the polarization correlation timescale, τ_d , versus magnetic field at constant laser intensity. This value changes from 9.0 ± 0.4 ns at zero field to 14.5 ± 0.5 ns at 300 mT.

A model of the dephasing of electron spin in QDs was presented by Merkolov, Efros and Rosen [20]. In this framework it is assumed that on timescales below $1 \mu s$ the hyperfine interaction between the electron spin and the nuclei in the dot can be considered semi-classically as a “frozen” magnetic field, of finite variance, but no directional preference. The electron g -factor is assumed isotropic. The time evolution of the electron spin $\mathbf{S}(t)$, (initially along \mathbf{S}_0) is given by:

$$\mathbf{S}(t) = (\mathbf{S}_0 \cdot \mathbf{n})\mathbf{n} + \{\mathbf{S}_0 - (\mathbf{S}_0 \cdot \mathbf{n})\mathbf{n}\} \cos(\omega t) + \{[\mathbf{S}_0 - (\mathbf{S}_0 \cdot \mathbf{n})\mathbf{n}] \times \mathbf{n}\} \sin(\omega t) \quad (1a)$$

$$W(B_N) \propto \exp\left[-\frac{(B_N)^2}{\delta B_N^2}\right] \quad (1b)$$

Where the distribution of nuclear field strengths, $W(B_N)$ is parameterized by the Gaussian width of fluctuations, δB_N . Figure 5a shows how the spin projection along the \mathbf{z} direction S_z varies with the external magnetic field B_{ext}^z [20]. At fields of a few times δB_N the spin-projection along \mathbf{z} is stabilized. This has not eliminated the nuclear spin fluctuations, it has merely overwhelmed them at the cost of increased rate of precession about the net field. Any measurement along an orthogonal polarization direction will reveal the increased precession rate.




FIG. 5: (a) Projection of the electron spin along the z -direction as a function of magnetic field and time, assuming a nuclear field fluctuation of width $\delta B_N = 100$ mT and an electron g -factor of 0.5 [21]. (b) The averaged projection of the z -component of the spin over the radiative lifetime, scaled in terms of the electron spin dephasing time.

From Equation 1a we extract the expected final electron spin projection along \mathbf{z} , S_z , which is equal to the

polarization correlation C_0 (solid line in Figure 4b). The only fitting parameter is the width of the fluctuations in the nuclear-field δB_N , set to 100 mT. The dephasing time for this electron [8] is therefore $T_\Delta = \hbar/g_e\mu_B\delta B_N \sim 200ps$, which is, as expected, much less than the 1 ns radiative lifetime of the upper state. This provides a good fit to the data, reproducing the value of C_0 and width around zero field. The model fits less well at the higher fields. Partly, this can be explained by non-zero β , but the discrepancy requires further investigation.

The extracted δB_N is within the range derived from a spin-noise measurement [22] but is greater than inferred from dephasing of the X_0 state in similar dots [23]. We attribute this to the smaller wave-function extent of the electron when the dot additionally contains two holes. As δB_N scales with $1/\sqrt{V}$ where V is the volume of spins overlapping with the wave-function there is a variation in the effective δB_N between states. This is the same reason the electron g -factor changes in the presence of additional holes [21].

To increase C_0 one could employ a host semiconductor without nuclear spin, a QD of greater volume or reduce the fluctuations in the nuclear field. Alternatively, Figure

5b shows that reducing τ_{rad} to the electron spin-lifetime leads to a significant increase in polarization correlation. This could be achieved by placing the dot into a cavity that equally enhances the radiative decay, independent of polarization.

In conclusion, despite the long hole spin coherence time in quantum dots the emission of polarization-correlated photons from the X^+ state is limited by electron spin dephasing. A significant increase in the polarization correlation time should be achieved by reducing the radiative lifetime of the X^+ state. Additionally, the degree of polarization correlation can be increased by applying an external magnetic field greater than the nuclear field.

I. ACKNOWLEDGEMENTS

We thank the EPSRC for funding the molecular beam epitaxy (MBE) machine and the Controlled Quantum Dynamics Centre for Doctoral Training (CQD-CDT) which supported Y. Cao.

-
- [1] M. Atature, J. Dreiser, A. Badolato, A. Högele, K. Karrai and A. Imamoglu, **312**, 551-553 (2006)
 - [2] B. D. Gerardot, D. Brunner, P. A. Dalgarno, P. Öhberg, S. Seidl, M. Kroner, K. Karrai, N. G. Stoltz, P. M. Petroff, and R. J. Warburton, *Nature*. **451**, 441-444 (2007).
 - [3] K. De Greve, P. L. McMahon, D. Press, T. D. Ladd, D. Bisping, C. Schneider, M. Kamp, L. Worschech, S. Höfling, A. Forchel, and Y. Yamamoto, *Nature Phys.* **7**, 872-878 (2011).
 - [4] K. De Greve, L. Yu, P. L. McMahon, J. S. Pelc, C. M. Natarajan, N. Y. Kim, E. Abe, S. Maier, C. Schneider, M. Kamp, S. Höfling, R. H. Hadfield, A. Forchel, M. M. Fejer, and Y. Yamamoto, *Nature* **491**, 421-425 (2012).
 - [5] W. B. Gao, P. Fallahi, E. Togan, J. Miguel-Sánchez, and A. Imamoglu, *Nature Phys.* **491**, 426-430 (2012).
 - [6] C. Y. Hu, W. J. Munro, and J. G. Rarity, *Phys. Rev. B* **78**, 125318 (2008)
 - [7] D. Witthaut, M. D. Lukin, and A. S. Sørensen, *Euro. Phys. Lett.* **97**, 50007 (2012)
 - [8] B. Urbaszek, X. Marie, T. Amand, O. Krebs, P. Voisin, P. Maletinsky, A. Högele and A. Imamoglu, *Rev. Mod. Phys.* **85**, 79 (2011)
 - [9] S. Cortez, O. Krebs, S. Laurent, M. Senes, X. Marie, P. Voisin, R. Ferreira, G. Bastard, J.-M. Gérard, and T. Amand, *Phys. Rev. Lett.* **89**, 207401 (2002)
 - [10] S. Laurent, M. Senes, O. Krebs, V. K. Kalevich, B. Urbaszek, X. Marie, T. Amand, and P. Voisin, *Phys. Rev. B* **73**, 235302 (2006)
 - [11] C. Testelin, F. Bernardot, B. Eble, and M. Chamarro, *Phys. Rev. B* **79**, 195440 (2009)
 - [12] E. A. Chekhovich, A. B. Krysa, M. S. Skolnick, and A. I. Tartakovskii, *Phys. Rev. Lett.* **106**, 027402 (2011)
 - [13] P. Fallahi, S. T. Yilmaz, and A. Imamoglu, *Phys. Rev. Lett.* **105**, 257802 (2010)
 - [14] D. Heiss, S. Schaeck, H. Huebl, M. Bichler, G. Abstreiter, J. J. Finley, D. V. Bulaev, and D. Loss, *Phys. Rev. B* **76**, 241306(R) (2007)
 - [15] M. Bayer, A. Kuther, A. Forchel, A. Gorbunov, V. B. Timofeev, F. Schäfer, J. P. Reithmaier, T. L. Reinecke, and S. N. Walck, *Phys. Rev. Lett.* **82**, 1748-1751 (1999).
 - [16] M. Bayer, G. Ortner, O. Stern, A. Kuther, A. A. Gorbunov, A. Forchel, P. Hawrylak, S. Fafard, K. Hinzer, T. L. Reinecke, S. N. Walck, J. P. Reithmaier, F. Klopff, and F. Schäfer, *Phys. Rev. B* **65**, 195315 (2002).
 - [17] D. Brunner, B. D. Gerardot, P. A. Dalgarno, G. Wüst, K. Karrai, N. G. Stoltz, P. M. Petroff, and R. J. Warburton, *Science*. **325**, 70-73 (2009)
 - [18] A. V. Koudinov, I. A. Akimov, Yu. G. Kusrayev, and F. Henneberger, *Phys. Rev. B* **70**, 241305(R) (2004)
 - [19] T. Belhadj, T. Amand, A. Kunold, C. Simon, T. Kuroda, M. Abbarchi, T. Mano, K. Sakoda, S. Kunz, X. Marie, and B. Urbaszek, *Appl. Phys. Lett.* **97**, 051111 (2010)
 - [20] I. A. Merkulov, Al. L. Efros, and M. Rosen, *Phys. Rev. B* **65**, 205309 (2002)
 - [21] A. J. Bennett, M. A. Pooley, Y. Cao, N. Sköld, I. Farrer, D. A. Ritchie, and A. J. Shields *Nature Comms.* **4**, 1544 (2013).
 - [22] A. V. Kuhlmann, J. Houel, A. Ludwig, L. Greuter, D. Reuter, A. D. Wieck, M. Poggio, and R. J. Warburton, *Nature Phys.* **9**, 570-575 (2013).
 - [23] A. J. Bennett, M. A. Pooley, R. M. Stevenson, I. Farrer, D. A. Ritchie, and A. J. Shields, *Phys. Rev. B* **84**, 195401 (2011)