

Stancu-variant of generalized Baskakov operators

Nadeem Rao*, Abdul Wafi

Department of Mathematics, Jamia Millia Islamia, New Delhi-110025, India

Abstract

In the present paper, we introduce Stancu-variant of generalized Baskakov operators and study the rate of convergence using modulus of continuity, order of approximation for the derivative of function f . Direct estimate is proved using K-functional and Ditzian-Totik modulus of smoothness. In the last, we have proved Voronovskaya type theorem.

Keywords: Generalized Baskakov operators, Modulus of continuity, Ditzian-Totik modulus of smoothness, Voronovskaya.

1. Introduction

In 1998 V. Mihesan[8] constructed an important generalization of the well known Baskakov operators on $[0, \infty)$ with non-negative constant a independent of n

$$B_n^a(f; x) = \sum_{k=0}^{\infty} W_{n,k}^a(x) f\left(\frac{k}{n}\right), \quad (1)$$

where $f \in C[0, \infty)$ and

$$W_{n,k}^a(x) = e^{-\frac{ax}{1+x}} \frac{p_k(n, a)}{k!} \frac{x^k}{(1+x)^{k+n}}, \quad (2)$$

such that $\sum_{k=0}^{\infty} W_{n,k}^a(x) = 1$ and $p_k(n, a) = \sum_{i=0}^{\infty} \binom{n}{i} (n)_i a^{k-i}$, with $(n)_0 = 1, (n)_i = n(n+1)\dots(n+i-1)$.

In the last decade, many papers were published for generalized Baskakov operators on order of approximation, Voronovskaya type theorem, Kantorovich form, and order of approximation for the derivative of the function([13],[14]). For $f \in C[0, \infty)$, Stancu[10] introduced the sequence of positive linear operators

$$B_n^{\alpha, \beta}(f; x) = \sum_{k=0}^n p_{n,k}(x) f\left(\frac{k+\alpha}{n+\beta}\right),$$

where $p_{n,k}(x) = \binom{n}{k} x^k (1-x)^{n-k}$ and α, β are any two non negative real numbers such that $0 \leq \alpha \leq \beta$. If $\alpha = \beta = 0$, it reduces to so called Bernstein operators. Recently, many researchers ([1],[4],[5],[6],[9],[11],[15]) have introduced the Stancu-variant for different linear positive operators. Motivated by the above development, we are giving a Stancu-variant of the operators (1) as

$$L_{n,a}^{\alpha, \beta}(f; x) = \sum_{k=0}^{\infty} W_{n,k}^a(x) f\left(\frac{k+\alpha}{n+\beta}\right), \quad (3)$$

where $W_{n,k}^a(x)$ defined in (2) and $0 \leq \alpha \leq \beta$. For $\alpha = \beta = 0$, we get the operators (1).

*Corresponding Author

Email addresses: nadeemrao1990@gmail.com (Nadeem Rao), abdulwafi2k2@gmail.com (Abdul Wafi)

2. Approximation properties of $L_{n,a}^{\alpha,\beta}$

To prove the approximation properties of $L_{n,a}^{\alpha,\beta}$, we need the following lemma[13]

Lemma2.1 For $a, x \geq 0, n = 1, 2, \dots$, we have

$$\begin{aligned}
 B_n^a(1; x) &= 1, \\
 B_n^a(t; x) &= x + \frac{ax}{n(1+x)}, \\
 B_n^a(t^2; x) &= \frac{x^2}{n} + \frac{x}{n} + x^2 + \frac{a^2x^2}{n^2(1+x)^2} + \frac{2ax^2}{n(1+x)} + \frac{ax}{n^2(1+x)}, \\
 B_n^a(t^3; x) &= x^3 + \frac{3x^2(1+x)}{n} + \frac{x(1+x)(1+2x)}{n^2} + \frac{3ax^3}{n(1+x)} + \frac{1}{n^2} \left(3ax^2 + \frac{3a^2x^3}{(1+x)^2} + \frac{3ax^2}{(1+x)} \right) \\
 &\quad + \frac{1}{n^3} \left(\frac{ax}{1+x} + \frac{3a^2x^2}{(1+x)^2} + \frac{a^3x^3}{(1+x)^3} \right), \\
 B_n^a(t^4; x) &= x^4 + \frac{6x^3(1+x)}{n} + \frac{x^2(1+x)(7+11x)}{n^2} + \frac{x(1+x)(6x^2+6x+1)}{n^3} + \frac{4ax^4}{n(1+x)} \\
 &\quad + \frac{1}{n^2} \left(\frac{12ax^4}{(1+x)} + \frac{18ax^3}{(1+x)^2} + \frac{18ax^3}{(1+x)^2} + \frac{18a^2x^3}{(1+x)^2} + \frac{14ax^2}{(1+x)} \right) \\
 &\quad + \frac{1}{n^4} \left(\frac{ax}{1+x} + \frac{7a^2x^2}{(1+x)^2} + \frac{6a^3x^3}{(1+x)^3} + \frac{a^4x^4}{(1+x)^4} \right).
 \end{aligned}$$

Next, we prove

Lemma 2.2 Let $a, x \geq 0$ and $n = 1, 2, 3, \dots$. Then for the operators defined in (3), we have

$$\begin{aligned}
 (i) \quad L_{n,a}^{\alpha,\beta}(1; x) &= 1, \\
 (ii) \quad L_{n,a}^{\alpha,\beta}(t; x) &= \frac{n}{n+\beta}x + \frac{a}{n+\beta} \frac{x}{1+x} + \frac{\alpha}{n+\beta}, \\
 (iii) \quad L_{n,a}^{\alpha,\beta}(t^2; x) &= \frac{n^2+n}{(n+\beta)^2}x^2 + \frac{n(1+2\alpha)}{(n+\beta)^2}x + \frac{a^2}{(n+\beta)^2} \frac{x^2}{(1+x)^2} + \frac{2an}{(n+\beta)^2} \frac{x^2}{(1+x)} + \frac{a(1+2\alpha)}{(n+\beta)^2} \frac{x}{1+x} + \frac{\alpha^2}{(n+\beta)^2}, \\
 (iv) \quad L_{n,a}^{\alpha,\beta}(t^3; x) &= \frac{n^3+3n^2+2n}{(n+\beta)^3}x^3 + \frac{n^2(3+3\alpha)+n(3+3\alpha+3a)}{(n+\beta)^3}x^2 + \frac{n(1+3\alpha+3\alpha^2)}{(n+\beta)^3}x + \frac{3an^2}{(n+\beta)^3} \frac{x^3}{(1+x)} \\
 &\quad + \frac{n}{(n+\beta)^3} \left(\frac{3a^2x^3}{(1+x)^2} + \frac{3ax^2}{1+x} + \frac{6a\alpha x^2}{1+x} \right) + \frac{1}{(n+\beta)^3} \left(\frac{ax}{1+x} + \frac{3a^2x^2}{(1+x)^2} + \frac{a^3x^3}{(1+x)^3} \right) \\
 &\quad + \frac{3\alpha a^2x^2}{(1+x)^2} + \frac{3\alpha^2ax}{1+x} + \alpha^3, \\
 (v) \quad L_{n,a}^{\alpha,\beta}(t^4; x) &= \frac{n^4+6n^3+11n^2+6n}{(n+\beta)^4}x^4 + \frac{(6+4\alpha)n^3+(18+12\alpha)n^2+(9+8\alpha)n}{(n+\beta)^4}x^3 + \left(\frac{(7+12\alpha+6\alpha^2)n^2}{(n+\beta)^4} \right. \\
 &\quad + \frac{(7+12\alpha+12\alpha a+6\alpha^2)n}{(n+\beta)^4} \Big) x^2 + \frac{(1+4\alpha+6\alpha^2+4\alpha^3)n}{(n+\beta)^4}x + \frac{4an^3+12an^2+8an}{(n+\beta)^4} \frac{x^4}{1+x} \\
 &\quad + \frac{6a^2n^2+6a^2n}{(n+\beta)^4} \frac{x^4}{(1+x)^2} + \frac{4a^3n}{(n+\beta)^4} \frac{x^4}{(1+x)^3} + \frac{a^4}{(n+\beta)^4} \frac{x^4}{(1+x)^4} + \frac{18an^2+18an}{(n+\beta)^4} \frac{x^3}{(1+x)} \\
 &\quad + \frac{(18a^2+12a^2\alpha)n}{(n+\beta)^4} \frac{x^3}{(1+x)^2} + \frac{6a^3+4\alpha a^3}{(n+\beta)^4} \frac{x^3}{(1+x)^3} + \frac{(12a\alpha^2+12a\alpha+14a)n}{(n+\beta)^4} \frac{x^2}{(1+x)} \\
 &\quad + \frac{7a^2+12a^2\alpha+6a^2\alpha^2}{(n+\beta)^4} \frac{x^2}{(1+x)^2} + \frac{a+4\alpha a+6\alpha^2a+4\alpha^3a}{(n+\beta)^4} \frac{x}{1+x} + \frac{\alpha^4}{(n+\beta)^4}.
 \end{aligned}$$

Proof To prove these identities, we use the lemma(2.1) and linearity property

$$L_{n,a}^{\alpha,\beta}(t; x) = \frac{n}{n+\beta}B_n^a(t; x) + \frac{\alpha}{n+\beta}B_n^a(1; x).$$

In similar manner, we can prove identities (iii), (iv) and (v).

Lemma 2.3 Let $\psi_x^i(t) = (t-x)^i, i = 1, 2, 3, \dots$. For $a, x \geq 0$ and $n = 1, 2, 3, \dots$,

$$\begin{aligned}
L_{n,a}^{\alpha,\beta}(\psi_x^0(t);x) &= 1, \\
L_{n,a}^{\alpha,\beta}(\psi_x^1(t);x) &= \left(\frac{n}{n+\beta} - 1\right)x + \frac{a}{n+\beta} \frac{x}{1+x} + \frac{\alpha}{n+\beta}, \\
L_{n,a}^{\alpha,\beta}(\psi_x^2(t);x) &= \frac{n+\beta^2}{(n+\beta)^2}x^2 + \frac{n-2\alpha\beta}{(n+\beta)^2}x + \frac{a^2}{(n+\beta)^2} \frac{x^2}{(1+x)^2} - \frac{2a\beta}{(n+\beta)^2} \frac{x^2}{(1+x)} + \frac{a(1+2\alpha)}{(n+\beta)^2} \frac{x}{1+x} + \frac{\alpha^2}{(n+\beta)^2}, \\
L_{n,a}^{\alpha,\beta}(\psi_x^4(t);x) &= \frac{(3-12\beta)n^2 + (6+4\beta+2\beta^2+4\beta^3)n + \beta^4}{(n+\beta)^4}x^4 + \left(\frac{(6-12a-12\beta)n^2 + (9+8\alpha-12\beta(1+a+\alpha+\alpha\beta))n}{(n+\beta)^4}\right. \\
&\quad \left. + \frac{(6-12a-12\beta-12\alpha\beta^2)}{(n+\beta)^4}\right)x^3 + \frac{3n^2 + (7-4\beta+12\alpha a-12\alpha\beta+6\alpha^2)n + 6\alpha^2\beta^2}{(n+\beta)^4}x^2 \\
&\quad + \frac{(1+4\alpha+6\alpha^2)n-4\alpha^3\beta}{(n+\beta)^4}x + \frac{12an^2+8an-4a\beta^3}{(n+\beta)^4} \frac{x^4}{(1+x)} + \frac{6a^2n+6a^2\beta^2}{(n+\beta)^4} \frac{x^4}{(1+x)^2} - \frac{4a^3\beta}{(n+\beta)^4} \frac{x^4}{(1+x)^3} \\
&\quad + \frac{a^4}{(n+\beta)^4} \frac{x^4}{(1+x)^4} + \frac{12an^2+18an+6a(1+2\alpha)\beta^2}{(n+\beta)^4} \frac{x^3}{1+x} + \frac{6a^2n-(12a^2+12\alpha a^2)\beta}{(n+\beta)^4} \frac{x^3}{(1+x)^2} \\
&\quad + \frac{(6a^3+4\alpha a^3)}{(n+\beta)^4} \frac{x^3}{(1+x)^3} + \frac{(12a\alpha+8a-6a\alpha^2)n-(6a+18\alpha^2a)\beta}{(n+\beta)^4} \frac{x^2}{1+x} + \frac{7a^2+12a^2\alpha+6a^2\alpha^2}{(n+\beta)^4} \frac{x^2}{(1+x)^2} \\
&\quad + \frac{(a)+4\alpha a+6\alpha^2a+4\alpha^3a}{(n+\beta)^4} \frac{x}{1+x} + \frac{\alpha^4}{(n+\beta)^4}.
\end{aligned}$$

Proof In view of lemma(2.2) and using equalities,

$$\begin{aligned}
L_{n,a}^{\alpha,\beta}(\psi_x(t);x) &= L_{n,a}^{\alpha,\beta}(t;x) - xL_{n,a}^{\alpha,\beta}(1;x), \\
L_{n,a}^{\alpha,\beta}(\psi_x^2(t);x) &= L_{n,a}^{\alpha,\beta}(t^2;x) - 2xL_{n,a}^{\alpha,\beta}(t;x) + x^2L_{n,a}^{\alpha,\beta}(1;x), \\
L_{n,a}^{\alpha,\beta}(\psi_x^4(t);x) &= L_{n,a}^{\alpha,\beta}(t^4;x) - 4xL_{n,a}^{\alpha,\beta}(t^3;x) + 6x^2L_{n,a}^{\alpha,\beta}(t^2;x) + 4x^3L_{n,a}^{\alpha,\beta}(t;x) + x^4L_{n,a}^{\alpha,\beta}(1;x).
\end{aligned}$$

we get the proof of this lemma.

Lemma 2.4 Let $\psi_x^i(t) = (t-x)^i, i = 1, 2, 3, \dots$. For $a, x \geq 0$ and $n = 1, 2, 3, \dots$, we have

$$\begin{aligned}
\lim_{n \rightarrow \infty} nL_{n,a}^{\alpha,\beta}(\psi_x^1(t);x) &= \alpha - \beta x + a \frac{x}{1+x}, \\
\lim_{n \rightarrow \infty} nL_{n,a}^{\alpha,\beta}(\psi_x^2(t);x) &= x^2 + x, \\
\lim_{n \rightarrow \infty} n^2L_{n,a}^{\alpha,\beta}(\psi_x^4(t);x) &= (3-12\beta)x^4 + (6-12a-12\beta)x^3 + 3x^2 + 12a \frac{x^2}{1+x} + 12a \frac{x^3}{1+x}.
\end{aligned}$$

3. The degree of approximation

Theorem 3.1 If $f \in C[0, \infty)$, $x \in [0, \infty)$ and $\omega(f; \delta)$ is the modulus of continuity, then

$$|L_{n,a}^{\alpha,\beta}(f;x) - f(x)| \leq \left\{1 + \sqrt{\gamma_{n,a}^{\alpha,\beta}(x)}\right\} \omega(f; \delta_{n,\beta}),$$

where $\delta_{n,\beta} = (n+\beta)^{-\frac{1}{2}}$ and

$$\gamma_n^{\alpha,\beta}(x) = \frac{n+\beta^2}{n+\beta}x^2 + \frac{n-2\alpha\beta}{n+\beta}x + \frac{a^2}{n+\beta} \frac{x^2}{(1+x)^2} - \frac{2a\beta}{n+\beta} \frac{x^2}{(1+x)} + \frac{a(1+2\alpha)}{n+\beta} \frac{x}{1+x} + \frac{\alpha^2}{n+\beta}.$$

Proof Let $f \in C[0, \infty)$ and $x \geq 0$. Then, using linearity property and monotonicity of the operators defined by (3), we can easily find, for every $\delta > 0$, and $n \in \mathbb{N}$, that

$$\begin{aligned} |L_{n,a}^{\alpha,\beta}(f;x) - f(x)| &\leq \left\{ 1 + \delta_{n,\beta}^{-1} \sqrt{L_{n,a}^{\alpha,\beta}(\psi_x^2;x)} \right\} \omega(f; \delta_{n,\beta}). \\ &\leq \left\{ 1 + \sqrt{\frac{n+\beta^2}{n+\beta}x^2 + \frac{n-2\alpha\beta}{n+\beta}x + \frac{a^2}{n+\beta} \frac{x^2}{(1+x)^2} - \frac{2a\beta}{n+\beta} \frac{x^2}{(1+x)} + \frac{a(1+2\alpha)}{n+\beta} \frac{x}{1+x} + \frac{\alpha^2}{n+\beta}} \right\} \omega(f; \delta_{n,\beta}), \end{aligned}$$

which obtained by using Lemma 2.2 and choosing $\delta_{n,\beta} = (n+\beta)^{-\frac{1}{2}}$. Thus, we arrive at the result.

Remark If we put $\alpha = \beta = 0$, we find the same result given by Miheşan[8]

$$|B_n^a(f;x) - f(x)| \leq \left\{ 1 + \sqrt{x(1+x) + \frac{ax}{n(1+x)} \frac{(a+1)x+1}{(1+x)}} \right\} \omega(f; \delta),$$

where $\delta = \frac{1}{\sqrt{n}}$, which shows that $\delta_{n,\beta} \leq \delta$. Therefore, rate of convergence of $L_{n,a}^{\alpha,\beta}$ is better than B_n^a .

Now, we will find the rate of convergence of operators defined by (3) in terms of modulus of continuity of first derivative of function i.e. $\omega(f'; \delta_{n,\beta}) = \omega_1(f; \delta_{n,\beta})$, which is an improvement over the Theorem 3.1. This type of result was given for Bernstein polynomials by Lorentz ([7], p.p. 21).

Theorem 3.2 Let $f'(x)$ is the continuous derivative over $[0, \infty)$ and $\omega_1(f; \delta_{n,\beta})$ is the modulus of continuity of $f'(x)$. Then, for $a, x \geq 0$, $0 \leq \alpha \leq \beta$, we have

$$|L_{n,a}^{\alpha,\beta}(f;x) - f(x)| \leq \omega_1((n+\beta)^{-1}) \sqrt{L_{n,a}^{\alpha,\beta}(\psi_x^2(t);x)} \left\{ 1 + \sqrt{(n+\beta)} \sqrt{L_{n,a}^{\alpha,\beta}(\psi_x^2(t);x)} \right\}.$$

Proof For $x_1, x_2 \in [a, b]$, we have

$$\begin{aligned} f(x_1) - f(x_2) &= (x_1 - x_2) f'(\xi), \\ &= (x_1 - x_2) f'(x_1) + (x_1 - x_2) [f'(\xi) - f'(x_1)], \end{aligned} \quad (4)$$

where $x_1 < \xi < x_2$. As we know that

$$|(x_1 - x_2)[f'(\xi) - f'(x_1)]| \leq |x_1 - x_2|(\lambda + 1)\omega_1(\delta), \quad \lambda = \lambda(x_1, x_2; \delta).$$

Next, we get

$$|L_{n,a}^{\alpha,\beta}(f;x) - f(x)| = \left| \sum_{n=0}^{\infty} W_{n,k}^a(x) \left\{ f\left(\frac{k+\alpha}{n+\beta}\right) - f(x) \right\} \right|. \quad (5)$$

From (4) and (5), we obtained

$$\begin{aligned} |L_{n,a}^{\alpha,\beta}(f;x) - f(x)| &\leq \left| \sum_{n=0}^{\infty} W_{n,k}^a(x) \left(\frac{k+\alpha}{n+\beta} - x \right) f'(x) \right| + \omega_1(\delta_{n,\beta}) \sum_{k=0}^{\infty} \left| \frac{k+\alpha}{n+\beta} - x \right| (\lambda + 1) W_{n,k}^a(x), \\ &\leq \omega_1(\delta_{n,\beta}) \left\{ \sum_{k=0}^{\infty} \left| \frac{k+\alpha}{n+\beta} - x \right| W_{n,k}^a(x) + \sum_{\lambda \geq 1} \left| \frac{k+\alpha}{n+\beta} - x \right| \lambda \left(x_1, \frac{k+\alpha}{n+\beta}; \delta \right) W_{n,k}^a(x) \right\} \\ &\leq \omega_1(\delta_{n,\beta}) \left\{ \sum_{k=0}^{\infty} \left| \frac{k+\alpha}{n+\beta} - x \right| W_{n,k}^a(x) + \delta^{-1} \sum_{k=0}^{\infty} \left(\frac{k+\alpha}{n+\beta} - x \right)^2 W_{n,k}^a(x) \right\} \\ &\leq \omega_1(\delta_{n,\beta}) \sqrt{L_{n,a}^{\alpha,\beta}(\psi_x^2(t);x)} \left\{ 1 + \delta_{n,\beta}^{-1} \sqrt{L_{n,a}^{\alpha,\beta}(\psi_x^2(t);x)} \right\}. \end{aligned}$$

Taking $\delta_{n,\beta} = (n+\beta)^{-1}$, we get

$$|L_{n,a}^{\alpha,\beta}(f;x) - f(x)| \leq \omega_1((n+\beta)^{-1}) \sqrt{L_{n,a}^{\alpha,\beta}(\psi_x^2(t);x)} \left\{ 1 + \sqrt{(n+\beta)} \sqrt{L_{n,a}^{\alpha,\beta}(\psi_x^2(t);x)} \right\},$$

which is the required result.

4. Direct Estimate

Here we introduced the Ditzian-Totik Modulus of smoothness[3] which is defined as:

$$\begin{aligned}\omega_{\varphi^\lambda}^2(f; \delta) &= \sup_{0 < h \leq \delta} \|\Delta_{h\varphi(x)}^2 f(x)\|, \\ &= \sup_{0 < h \leq \delta} \sup_{x \pm h\varphi^\lambda \in [0, \infty)} |f(x - h\varphi^\lambda(x)) - 2f(x) + f(x + h\varphi^\lambda(x))|,\end{aligned}$$

where $\varphi^2(x) = x(1-x)$. And, Peetre's K-functional is given by

$$K_{\varphi^\lambda}(f, \delta^2) = \inf_g \left(\|f - g\|_{C[0, \infty)} + \delta^2 \|\varphi^{2\lambda} g''\|_{C[0, \infty)} \right), \quad g, g' \in AC_{loc}. \quad (6)$$

The K-functional is equivalent to the modulus of smoothness, i.e.,

$$C^{-1}K_{\varphi^\lambda}(f, \delta^2) \leq \omega_{\varphi^\lambda}^2(f, \delta) \leq CK_{\varphi^\lambda}(f, \delta^2). \quad (7)$$

First result based on Ditzian-Totik modulus of smoothness was given by Ditzian[2] for the Bernstein polynomials as:

$$|B_n(f; x) - f(x)| \leq C\omega_{\varphi^\lambda}^2(f, n^{-\frac{1}{2}}\varphi(x)^{1-\lambda}).$$

Now, we prove the similar result for the operator $L_{n,a}^{\alpha,\beta}$.

Theorem 4.1 For $a, x \geq 0$, and $0 \leq \alpha \leq \beta$, we have

$$|L_{n,a}^{\alpha,\beta}(f; x) - f(x)| \leq C\omega_{\varphi^\lambda}^2(f, (n+\beta)^{-\frac{1}{2}}\varphi(x)^{1-\lambda}) \text{ for large } n.$$

Proof Using (6),(7), we can choose $g_n \equiv g_{n,x,\lambda}$ for fixed x and $\lambda + 1$ such that

$$\|f - g\|_{C[0, \infty)} \leq A\omega_{\varphi^\lambda}^2(f, n^{-\frac{1}{2}}\varphi(x)^{1-\lambda}), \quad (8)$$

$$n^{-1}\varphi(x)^{2-2\lambda} \|\varphi^{2\lambda} g''\|_{C[0, \infty)} \leq B\omega_{\varphi^\lambda}^2(f, n^{-\frac{1}{2}}\varphi(x)^{1-\lambda}). \quad (9)$$

Next

$$\begin{aligned}|L_{n,a}^{\alpha,\beta}(f; x) - f(x)| &\leq |L_{n,a}^{\alpha,\beta}(f - g_n; x) - (f - g_n)(x)| + |L_{n,a}^{\alpha,\beta}(g_n; x) - g_n(x)|, \\ &\leq 2 \|f - g_n\|_{C[0, \infty)} + |L_{n,a}^{\alpha,\beta}(g_n; x) - g_n(x)|.\end{aligned}$$

From (8), we get

$$|L_{n,a}^{\alpha,\beta}(f; x) - f(x)| \leq 2A\omega_{\varphi^\lambda}^2(f, n^{-\frac{1}{2}}\varphi(x)^{1-\lambda}) + |L_{n,a}^{\alpha,\beta}(g_n; x) - g_n(x)|. \quad (10)$$

Now, the last term can be calculated by using Taylor's formula

$$\begin{aligned}
|L_{n,a}^{\alpha,\beta}(g_n(t) - g_n(x); x)| &\leq |g'_n(x)L_{n,a}^{\alpha,\beta}((t-x); x)| + \left| L_{n,a}^{\alpha,\beta} \left(\int_t^x (x-u)g''_n(u)du; x \right) \right| \\
&\leq L_{n,a}^{\alpha,\beta} \left(\frac{|x - \frac{k}{n}|}{\varphi^{2\lambda}(x)} \int_{\frac{k}{n}}^x \varphi^{2\lambda}(u) |g''_n(u)| du; x \right) \\
&\leq \| \varphi^{2\lambda} g''_n \|_{C[0,\infty)} \frac{1}{\varphi^{2\lambda}(x)} L_{n,a}^{\alpha,\beta}((t-x)^2; x) \\
&\leq \| \varphi^{2\lambda} g''_n \|_{C[0,\infty)} \frac{1}{\varphi^{2\lambda}(x)} \left[\frac{n+\beta^2}{(n+\beta)^2} x^2 + \frac{n-2\alpha\beta}{(n+\beta)^2} x + \frac{a^2}{(n+\beta)^2} \frac{x^2}{(1+x)^2} - \frac{2a\beta}{(n+\beta)^2} \frac{x^2}{(1+x)} \right. \\
&\quad \left. + \frac{a(1+2\alpha)}{(n+\beta)^2} \frac{x}{1+x} + \frac{\alpha^2}{(n+\beta)^2} \right] \\
&\leq \| \varphi^{2\lambda} g''_n \|_{C[0,\infty)} \frac{x(1+x)(n+\beta)^{-1}}{\varphi^{2\lambda}(x)} \left[\frac{n+\beta^2}{(n+\beta)} \frac{x}{1+x} + \frac{n-2\alpha\beta}{(n+\beta)} \frac{1}{1+x} + \frac{a^2}{(n+\beta)} \frac{x}{(1+x)^3} \right. \\
&\quad \left. - \frac{2a\beta}{(n+\beta)} \frac{x}{(1+x)^2} + \frac{a(1+2\alpha)}{(n+\beta)} \frac{1}{(1+x)^2} + \frac{\alpha^2}{(n+\beta)x(1+x)} \right] \\
&\leq \| \varphi^{2\lambda} g''_n \|_{C[0,\infty)} \varphi^{2-2\lambda}(x)(n+\beta)^{-1} \left[\frac{n+\beta^2}{(n+\beta)} \frac{x}{1+x} + \frac{n-2\alpha\beta}{(n+\beta)} \frac{1}{1+x} + \frac{a^2}{(n+\beta)} \frac{x}{(1+x)^3} \right. \\
&\quad \left. - \frac{2a\beta}{(n+\beta)} \frac{x}{(1+x)^2} + \frac{a(1+2\alpha)}{(n+\beta)} \frac{1}{(1+x)^2} + \frac{\alpha^2}{(n+\beta)x(1+x)} \right].
\end{aligned}$$

From (9), we have

$$\begin{aligned}
|L_{n,a}^{\alpha,\beta}(g_n(t) - g_n(x); x)| &\leq B\omega_{\varphi}^2(f, (n+\beta)^{-\frac{1}{2}}\varphi(x)^{1-\lambda}) \left[\frac{n+\beta^2}{(n+\beta)} \frac{x}{1+x} + \frac{n-2\alpha\beta}{(n+\beta)} \frac{1}{1+x} + \frac{a^2}{(n+\beta)} \frac{x}{(1+x)^3} \right. \\
&\quad \left. - \frac{2a\beta}{(n+\beta)} \frac{x}{(1+x)^2} + \frac{a(1+2\alpha)}{(n+\beta)} \frac{1}{(1+x)^2} + \frac{\alpha^2}{(n+\beta)x(1+x)} \right]. \quad (11)
\end{aligned}$$

Using (10) and (11), we get

$$\begin{aligned}
|L_{n,a}^{\alpha,\beta}(f(t) - f(x); x)| &\leq M\omega_{\lambda}^2 \left(f, (n+\beta)^{-\frac{1}{2}}\varphi(x)^{1-\lambda} \right) \left[\frac{n+\beta^2}{(n+\beta)} \frac{x}{1+x} + \frac{n-2\alpha\beta}{(n+\beta)} \frac{1}{1+x} + \frac{a^2}{(n+\beta)} \frac{x}{(1+x)^3} \right. \\
&\quad \left. - \frac{2a\beta}{(n+\beta)} \frac{x}{(1+x)^2} + \frac{a(1+2\alpha)}{(n+\beta)} \frac{1}{(1+x)^2} + \frac{\alpha^2}{(n+\beta)x(1+x)} \right]
\end{aligned}$$

where $M = \max(2A, B)$. For a large value of n

$$|L_{n,a}^{\alpha,\beta}(f(t) - f(x); x)| \leq M\omega_{\lambda}^2 \left(f, (n+\beta)^{-\frac{1}{2}}\varphi(x)^{1-\lambda} \right).$$

Asymptotic relation is the study of rate of convergence for at least two times differentiable functions which was given by Voronovskaya [12]. Here, we prove a similar result.

Theorem 4.2 Let $a, x \geq 0$, $0 \leq \alpha \leq \beta$ and $n \in \mathbb{N}$. For $f \in C^2[0, \infty)$, we have

$$\lim_{n \rightarrow \infty} n \{ L_{n,a}^{\alpha,\beta}(f; x) - f(x) \} = \left(\alpha - \beta x + \frac{ax}{1+x} \right) f'(x) + \frac{x^2 + x}{2} f''(x).$$

Proof Let $x, t \in [0, \infty)$, $f \in C^2[0, \infty)$. By Taylor's formula, we have

$$f(t) = f(x) + (t-x)f'(x) + \frac{(t-x)^2}{2} f''(x) + \eta(t, x)(t-x)^2,$$

where the function $\eta(t, x) \in C[0, \infty)$ and $\lim_{t \rightarrow x} \eta(t, x) = 0$. Multiplying both sides by $W_{n,k}^a(x)$ and summing over k , we get

$$L_{n,a}^{\alpha,\beta}(f; x) = f(x)L_{n,a}^{\alpha,\beta}(1; x) + f'(x)L_{n,a}^{\alpha,\beta}(t-x; x) + \frac{f''(x)}{2}L_{n,a}^{\alpha,\beta}((t-x)^2; x) + L_{n,a}^{\alpha,\beta}(\eta(t, x)(t-x); x).$$

Using lemma(2.2), we obtain

$$\lim_{n \rightarrow \infty} n\{L_{n,a}^{\alpha,\beta}(f; x) - f(x)\} = \left(\alpha - \beta x + \frac{ax}{1+x}\right)f'(x) + \frac{x^2+x}{2}f''(x) + \lim_{n \rightarrow \infty} nL_{n,a}^{\alpha,\beta}(\eta(t; x)(t-x)^2; x). \quad (12)$$

Now, the last term can be obtained using Holder's inequality and lemma 2.4

$$nL_{n,a}^{\alpha,\beta}(\eta(t; x)(t-x)^2; x) \leq n^2L_{n,a}^{\alpha,\beta}((t-x)^4; x)L_{n,a}^{\alpha,\beta}(\eta(t; x)^2; x),$$

Let $\varphi(t; x) = \eta^2(t; x)$. Then, $\lim_{t \rightarrow x} \varphi(t; x) = 0$. Therefore,

$$\lim_{n \rightarrow \infty} nL_{n,a}^{\alpha,\beta}(\eta(t; x)(t-x)^2; x) = 0.$$

On substituting this value in equation (12), we get the desired result.

References

References

- [1] Aral A., Acar T., Weighted approximation by new Bernstein-Chlodowsky-Gadjiev operators, Filomat, 2013, **27** (2), 371-380.
- [2] Ditzian Z., Direct estimate for Bernstein polynomials, J. Approx. Theory, 1994, **79** (1), 165-166.
- [3] Ditzian Z., Totik V., Moduli of smoothness, Springer Series in Computational Mathematics, 9. Springer-Verlag, New York, 1987.
- [4] Gupta V., A Stancu variant of beta-Szasz operators, Georgian Math. J., 2014, **21** (1), 75-82.
- [5] Gupta V., Verma D.K., Agrawal P.N., Simultaneous approximation by certain Baskakov-Durrmeyer-Stancu operators, J. Egyptian Math. Soc., 2012, **20** (3), 183-187.
- [6] Gupta V., Agarwal R.P., Convergence Estimates in Approximtion Theory, Springer, Cham, 2014. .
- [7] Lorentz G.G., Mathematical Expositions, No. 8, Bernstein polynomials, University of Toronto Press, Toronto 1953.
- [8] Mihasan V., Uniform approximation with positive linear operators generated by generalised Baskakov method, Automat. Copmut. Appl. Math., 1998, **7** (1), 34-37.
- [9] Paltanea R., Approximation Theory Using Positive Linear Operators, Birkhauser, Boston Inc., Boston, MA, 2004.
- [10] Stancu D.D., On a generalization of the Bernstein polynomials(Romanian), Studia. Univ. Babes-Bolyai Ser. Math.-Phys., 1969, **14** (2), 31-45.
- [11] Sucu S., Ibikli E., Approximation by means of Kantorovich -Stancu type operators, Numer. Funct. Anal. Optim., 2013, **34** (5), 557-575.
- [12] Voronovskaja E., Determination de la forme asymptotique d'approximation des fonction par ploynomes de M. Bernstein, C. R. Acad. Sci. URSS 1932, **79**, 79-85.
- [13] Wafi A., Khatoun S., Convergence and Voronovskaja-type theorems for derivatives of generalized Baskakov operators, Cent. Eur. J. Math., 2008, **6** (2), 325-334.
- [14] Wafi A., Khatoun S., On the order of approximation of functions by generalized Baskakov operators, Indian J. pure appl. Math., 2004, **35** (3), 347-358.
- [15] Wang M., Yu D., Zhou P., On the approximation by operators of Bernstein-Stancu types, Appl. Math. Comput., 2014, **246**, 79-87.