A simple two-sample Bayesian *t*-test for hypothesis testing

Min Wang*

Department of Mathematical Sciences, Michigan Technological University, Houghton, MI, USA

and

Guangying Liu

Department of Mathematics and Statistics, Nanjing Audit University, Nanjing, Jiangsu, China

December 30, 2021

Abstract

In this paper, we propose an explicit closed-form Bayes factor for the problem of two-sample hypothesis testing. The proposed approach can be regarded as a Bayesian version of the pooled-variance t-statistic and has various appealing properties in practical applications. It relies on data only through the t-statistic and can thus be calculated by using an Excel spreadsheet or a pocket calculator. It avoids several undesirable paradoxes, which may be encountered by the previous Bayesian approach of Gönen et al. (2005). Specifically, the proposed approach can be easily taught in an introductory statistics course with an emphasis on Bayesian thinking. Simulated and real data examples are provided for illustrative purposes.

Keywords: Bayes factor; Posterior probability, Pooled-variance *t*-statistic; Prior elicitation; Teaching elementary statistics.

^{*}Corresponding author: minwang@mtu.edu

1 Introduction

In an introductory statistics course, we usually teach students how to conduct a hypothesis test based on independent samples to compare the means of two populations with equal, but unknown variance. Let y_{ij} be random samples drawn from independent and normally distributed populations with means μ_i and common variance σ^2 for $j = 1, \dots, n_i$ and i = 1, 2. We are interested in testing

$$H_0: \mu_1 = \mu_2 \quad \text{versus} \quad H_1: \mu_1 \neq \mu_2.$$
 (1)

Within a frequentist framework, the pooled-variance two-sample t test is commonly used for the above hypothesis testing. The test statistic is given by

$$t = \frac{\bar{y}_1 - \bar{y}_2}{s_p / \sqrt{n_\delta}},\tag{2}$$

where $\bar{y}_i = \sum_{j=1}^{n_i} y_{ij}/n_i$ and

$$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2} \tag{3}$$

is the pooled-variance estimate of σ^2 with $s_i^2 = \sum_{j=1}^{n_i} (y_{ij} - \bar{y}_i)^2 / (n_i - 1)$ for i = 1, 2. Here, $n_{\delta} = (1/n_1 + 1/n_2)^{-1}$ is often called the "effective sample size" in the two-sample experiment. At the α significance level, we obtain the critical value $t_{1-\alpha/2,v}$ or P-value $p = 2P(T \ge |t|)$ with degrees of freedom $v = n_1 + n_2 - 2$, where $t_{1-\alpha/2,v}$ is the $(1 - \alpha/2)$ quantile of T_v distribution and T has the T_v distribution. We reject the null hypothesis H_0 if either $|t| > t_{1-\alpha/2,v}$ or $p < \alpha$; see Weiss (2012).

Bayesian approaches to hypothesis testing have recently received considerable attention and are becoming important in different disciplines, such as sociology (Western, 1999), economics (Fernández et al., 2001), and psychology (Rouder et al., 2009). Many recent studies suggest that we should offer at least one course about Bayesian methods to students at early stages in their mathematics and statistics education; see, for example, Albert (1997), Gönen et al. (2005), Wetzels et al. (2012), Wulff and Robinson (2014), among others. Specifically, as stated by Carlin and Louis (2000), "*The Bayesian approach to statistical design and analysis is emerging as an increasingly effective and practical alternative to the frequentist one.*" Such a course will not only motivate students' interests in Bayesian thinking, but also help them know how to formulate Bayesian methods in simple statistical scenarios, such as the hypothesis testing in (1). More importantly, it will make students ready to use both Bayesian and frequentist ideas.

A natural approach within a Bayesian framework to compare hypotheses is the Bayes factor (ratio of the marginal densities of the two models); see Kass and Raftery (1995). For the hypothesis testing in (1), Gönen et al. (2005) proposed a simple closed-form Bayes factor based on the two-sample t-statistic and it is given by

$$GBF[H_1:H_0](\sigma_a^2) = \left[\frac{1+t^2/\nu}{1+t^2/(\nu(1+n_\delta\sigma_a^2))}\right]^{(\nu+1)/2} (1+n_\delta\sigma_a^2)^{-1/2},$$
(4)

where σ_a^2 is a hyperparameter of the prior that needs to be specified. The choice of prior distributions for deriving the GBF will be stated in detail in the following section. The GBF in (4) shows a close relationship between frequentist and Bayesian ideas and can be easily covered in an elementary statistics course. Note that the choice of σ_a^2 is critical, because it acts as an inverse prior sample size. Specifically, the GBF with fixed σ_a^2 may exhibit some undesirable features, such as Bartlett's paradox and the information paradox; see Liang et al. (2008). These paradoxes will definitely confuse students and even make them struggle when conducting Bayesian data analysis.

In this paper, we specify a hyper-prior for the hyperparameter σ_a^2 to reduce the impact of misspecified hyperparameter values. The prior will still result in an explicit expression of the Bayes factor based on the two-sample *t*-statistic. It is shown that the proposed approach resolves several potential difficulties and paradoxes encountered by the previous approach due to Gönen et al. (2005). We hope that our results will facilitate an intuitive understanding and discussion of the relationship between frequentist and Bayesian ideas, but also shed some light on the importance of hyper-prior specifications to students, teachers, and researchers.

The remainder of this paper is organized as follows. In Section 2, we review the existing Bayes factor of Gönen et al. (2005) and discuss potential difficulties associated with fixed hyperparameter values. In Section 3, we specify a hyper-prior on that hyperparameter, which yields a closed-form expression for the Bayes factor. We investigate the finite sample performance of the two Bayesian procedures in a simulation study (Section 3) and a realdata example (Section 4). Some concluding remarks are given in Section 5, with derivation of the proposed procedure in the appendix.

2 Bayes inference

The Bayesian analysis begins with prior specifications for the unknown parameters. Let $p(\mathbf{Y} \mid \theta_j)$ and π_j be the likelihood function of \mathbf{Y} and the prior probability on hypothesis H_j ($\pi_0 + \pi_1 = 1$) for j = 0, 1, respectively. From Bayes theorem, the posterior probability of H_j is defined as

$$P(H_j \mid \mathbf{Y}) = \frac{\pi_j m_j(\mathbf{Y})}{\pi_0 m_0(\mathbf{Y}) + \pi_1 m_1(\mathbf{Y})}.$$
(5)

The corresponding marginal likelihood of $\mathbf Y$ given H_j is

$$m_j(\mathbf{Y}) = \int p(\mathbf{Y} \mid \theta_j) \pi_j(\theta_j) \, d\theta_j, \tag{6}$$

where $\pi_j(\theta_j)$ is the prior for the unknown parameter θ_j under H_j for j = 0, 1. The posterior probability of H_1 can be expressed as

$$P(H_1 \mid \mathbf{Y}) = \frac{\pi_1 \mathrm{BF}[H_1 : H_0]}{\pi_0 + \pi_1 \mathrm{BF}[H_1 : H_0]} = \left[1 + \frac{\pi_0}{\pi_1} \frac{1}{\mathrm{BF}[H_1 : H_0]}\right]^{-1},\tag{7}$$

where the Bayes factor, $BF[H_1: H_0]$, for comparing H_1 to H_0 is given by

$$BF[H_1:H_0] = \frac{m_1(\mathbf{Y})}{m_0(\mathbf{Y})}.$$
(8)

The hypothesis H_1 (H_0) is more likely to be selected when BF[$H_1 : H_0$] > 1 (< 1). More specifically, Jeffreys (1961) suggested that BF[$H_1 : H_0$] < 0.1, provides "strong" evidence in favor of H_0 , and BF[$H_1 : H_0$] < 0.01, provides "decisive" evidence. Note that the Bayes factor for the null relative to the alternative, denoted by BF[$H_0 : H_1$], is given by

$$BF[H_0: H_1] = \frac{1}{BF[H_1: H_0]}$$

For the hypothesis testing problem in (1), we need to specify appropriate prior distributions for (μ_1, μ_2, σ^2) . Gönen et al. (2005) show that this testing problem can be written in equivalent form as

$$H_0: \delta = \mu_1 - \mu_2 = 0 \quad \text{versus} \quad H_1: \delta \neq 0. \tag{9}$$

Therefore, they advocate a prior for δ/σ^2 , instead of μ , where $\mu = (\mu_1 + \mu_2)/2$. After reparameterization from (μ_1, μ_2, σ^2) to (μ, δ, σ^2) , the suggested priors are given by

$$\pi(\mu, \sigma^2) \propto 1/\sigma^2 \quad \text{and} \quad \delta/\sigma \mid \mu, \sigma^2, \delta \neq 0 \sim N(\lambda, \sigma_a^2),$$
(10)

where λ and σ_a^2 are the hyperparameters that need to be pre-specified. Due to lack of prior knowledge in practice, it is natural to set $\lambda = 0$ to reflect the uncertain direction of an effect. Thus, the case for which $\lambda = 0$ will be of interest to us in what follows. The Bayes factor under the above priors is

$$GBF[H_1:H_0](\sigma_a^2) = \left[\frac{1+t^2/v}{1+t^2/(v(1+n_\delta\sigma_a^2))}\right]^{(v+1)/2} (1+n_\delta\sigma_a^2)^{-1/2},$$
(11)

where $v = n_1 + n_2 - 2$. Note that the Bayes factor depends on the data only through the *t*-statistic and can often be calculated using a pocket calculator. As mentioned in the Introduction, the choice of σ_a^2 is quite critical, and in particular, the Bayes factor with fixed σ_a^2 may lead to several undesirable properties, such as Bartlett's paradox and the information paradox, briefly summarized as follows.

Bartlett's paradox: Because the hyperparameter σ_a^2 reflects the variance of the univariate normal distribution in (10), a large value of σ_a^2 is often chosen to minimize prior information. However, when σ_a^2 becomes sufficiently large, while v is fixed (n_{δ} is also fixed), the GBF tends to 0, indicating that it always favors the null hypothesis, regardless of the information from the data. This phenomenon is often called Bartlett's paradox, which has been studied by Jeffreys (1961) and more recently by Liang et al. (2008).

Information paradox: Suppose that samples are generated under H_1 . In this setting, when v is fixed, the posterior probability of H_1 should be higher than the one for H_0 when the t-statistic goes to infinity. We thus expect that the GBF tends to infinity as the information against H_0 accumulates. However, with a fixed value of σ_a^2 , the GBF becomes a constant $(1 + n_\delta \sigma_a^2)^{v/2}$ as $t \to \infty$. This is referred to as the information paradox.

The two paradoxes may confuse students and even make them struggle about Bayesian data analysis, especially when we introduce basic ideas of Bayesian inference in an elementary level. In this paper, we advocate a hyper-prior for σ_a^2 , which not only alleviates the impacts of misspecified hyperparameter, but also yields an explicit Bayes factor. More im-

portantly, the proposed approach is still a function of the two-sample *t*-statistic and enjoys various appealing properties, as discussed next.

2.1 The hyper-prior for σ_a^2

In this section, we consider a proper prior for σ_a^2 , denoted by $\pi(\sigma_a^2)$. The proposed Bayes factor can be written as

$$PBF[H_1:H_0] = \int_0^\infty \left[\frac{1+t^2/\nu}{1+t^2/(\nu(1+n_\delta\sigma_a^2))} \right]^{(\nu+1)/2} (1+n_\delta\sigma_a^2)^{-1/2} \pi(\sigma_a^2) \, d\sigma_a^2.$$
(12)

The prior for σ_a^2 is assigned to be the Pearson type VI distribution with shape parameters a > -1, b > -1, and scale parameter $\kappa > 0$. Its probability density function (pdf) is

$$\pi(\sigma_a^2) = \frac{\kappa(\kappa\sigma_a^2)^b (1+\kappa\sigma_a^2)^{-a-b-2}}{B(a+1,b+1)} I_{(0,\infty)}(\sigma_a^2),$$
(13)

where $B(\cdot, \cdot)$ is a beta function. This prior has also been used by Wang and Sun (2014) in the one-way random effects model. With the particular choice of $\kappa = n_{\delta}$ and b = (v+1)/2 - a - 5/2, the Bayes factor can be greatly simplified as

$$PBF[H_1:H_0] = \frac{\Gamma(v/2)\Gamma(a+3/2)}{\Gamma((v+1)/2)\Gamma(a+1)} \left(1 + \frac{t^2}{v}\right)^{(v-2a-2)/2},$$
(14)

which is an explicit expression and can thus be easily computed using an Excel spreadsheet or a simple calculator. Such an expression is unavailable for other choices of κ and b. Like the GBF in (11), it can be regarded as a Bayesian version of the *t*-statistic; in addition, our approach enjoys several appealing properties, which are not shared by the GBF. The proof of the theorem is straightforward and is thus omitted here for simplicity.

Theorem 1 In the setting of the information paradox mentioned above, the Bayes factor in (14) tends to infinity when -1 < a < v/2 - 1.

The theorem shows that when -1 < a < v/2 - 1, the specified hyper-prior provides a resolution of the information paradox that arises in the GBF. In the case of minimum sample sizes of the two samples (i.e., $n_1 + n_2 = 3$), we have v = 1, indicating that $a \in (-1, -1/2)$. Of particular note is that when a = -1/2, the asymptotic tail behavior of

$$\pi(\delta/\sigma \mid \mu, \sigma^2, \delta \neq 0) = \int_0^\infty N(\delta/\sigma \mid \lambda, \sigma_a^2) \pi(\sigma_a^2) \, d\sigma_a^2$$

becomes the Cauchy density for sufficiently large δ/σ , which provides a flat tail behavior and diminishes the prior influence of $\pi(\delta/\sigma \mid \mu, \sigma^2, \delta \neq 0)$, especially when *a* is small. Consequently, we recommend $a \in (-1, -1/2]$.

It deserves mentioning that the prior depends on the sample size and that as the sample size increases, the prior has a density in the right tail that behaves like $(\sigma_a^2)^{-a-2}$, leading to a fat tail for small value of a. Furthermore, it can be seen from Figure 1 that a higher prior probability is assigned to the event $\sigma_a^2 > 1$. This phenomenon occurs because the parameter σ_a^2 seems to act as an inverse prior sample size. A small value of σ_a^2 (such as $\sigma_a^2 \to 0$) makes the prior converge to a point mass at $\delta = 0$, and the alternative H_1 may collapse to H_0 . We thus obtain that the Bayes factor (the GBF) tends to 1, indicating that both hypotheses are equal descriptions to the data in the limit.



Figure 1: The hyper-prior for σ_a^2 with $\kappa = n_{\delta}$, a = -3/4, and b = (v+1)/2 - a - 5/2 for different choices of n_1 and n_2 .

To see how the PBF avoids Bartlett's paradox and the information paradox, we consider two simple examples with $n_1 = n_2 = 10$: one with a fixed *t*-statistic, and the other with an increasing value. Suppose that t = 5, providing strong evidence against H_0 . We observe



Figure 2: The Bayes factor as a function of the hyperparameter (left: the GBF; right: the PBF) when $n_1 = n_2 = 10$.

from Figure 2 that the PBF with $a \in (-1, -1/2]$ always rejects H_0 , while the GBF fails to reject H_0 when σ_a becomes large, regardless of the information from the data. Also, it is well-known that the larger the *t*-statistic, the stronger the evidence against H_0 . Figure 3 shows that as the *t*-statistic increases, the PBF grows faster than the GBF, which tends to a constant, even though *t* becomes significantly large. These two examples show that the PBF not only avoids these paradoxes, but also provides a way to enhance students' better understanding of these paradoxes.

3 Simulation study

In this section, we conduct simulation studies and sensitivity analysis to investigate the finite sample performance of the two Bayes factors (GBF and PBF) with various choices of their corresponding hyperparameters. For sample 1, we generate n_1 random variables normally distributed with mean 0 and standard deviation 1. For sample 2, we generate n_2 random variables normally distributed with mean δ and standard deviation 1, where δ ranges from -4 to 4 in increments of 0.1. To assess the sensitivity of the



Figure 3: The Bayes factor as a function of the *t*-statistic (left: the GBF with $\sigma_a = .1$; right: the PBF with a = -.75) when t = 5 and $n_1 = n_2 = 10$.

hyperparameters, we take $\sigma_a = \{0.1, 1/3, 0.5, 1, 1.2, 2, 5\}$ for the GBF in (11) and $a = \{-0.95, -0.9, -0.8, -0.75, -0.7, -0.6, -0.5\}$ for the PBF in (14). For each case, we analyze 10,000 simulated datasets with various choices of n_1 and n_2 . The decision criterion used in this paper is to choose H_1 if the Bayes factor > 1 and H_0 otherwise.

The relative frequencies of rejecting H_0 under the three different choices of sample size are depicted in Figures 4, 5, and 6. Rather than providing exhaustive results based on these simulations, we merely highlight the most important findings from the three figures. (i) The GBF is quite sensitive to the choice of the hyperparameter σ_a , even when the sample size is large. For instance, when $n_1 = n_2 = 100$ and $\delta = -0.3$, the frequency of rejecting H_0 changes from 0.8479 to 0.2843 with σ_a increasing from 0.1 to 5. (ii) The PBF is relatively insensitive to the hyperparameter a, and when the sample size is large, the PBF behaves similarly for all values. (iii) We observe that under H_0 (i.e., $\delta = 0$), the relative frequency of rejecting H_0 varies greatly for the GBF with different choice of σ_a , whereas the PBF is quite stable in terms of different value of a.

We now compare the performance of the two Bayes factors with the P-value based on the *t*-statistic in (2) when $\alpha = 0.05$. Based on the same simulation scheme described



Figure 4: The relative frequency of rejection of H_0 under different procedures (left: the GBF; right: the PBF) when $n_1 = n_2 = 10$.



Figure 5: The relative frequency of rejection of H_0 under different procedures (left: the GBF; right: the PBF) when $n_1 = n_2 = 30$.



Figure 6: The relative frequency of rejection of H_0 under different procedures (left: the GBF; right: the PBF) when $n_1 = n_2 = 100$.

above, we consider the GBF with $\sigma_a = 1/3$, suggested by Gönen et al. (2005) and the PBF with a = -3/4. Figure 7 depicts the numerical findings with different sample sizes. We observe that the PBF and the P-value have similar performances, whereas they are significantly different from the GBF. As expected, when the sample size becomes large, the three procedures behave very similarly. In addition, the PBF has a faster decreasing rate to zero than the two other methods, in terms of the relative frequency of rejecting H_0 . Thus, we may conclude that the PBF is consistent under H_0 when the sample size approaches infinity. This property is not shared by the two other methods under consideration.

4 A real-data application

We compare the performance of the two Bayes factors via a real-data example available at The Data and Story Library. (http://lib.stat.cmu.edu/DASL/Datafiles/Calcium.html). The data consist of the blood pressure measurements for 21 African-American men: 10 of the men took calcium supplements and 11 took placebos. We are interested in testing if increasing calcium intake reduces blood pressure. The pooled-variance t-statistic is 1.634,



with different sample sizes. Figure 7: The relative frequency of rejection of H_0 under the three considered procedures

with the two-sided P-value of 0.1187. The positive t-statistic indicates intake of calcium is beneficial for reducing blood pressure, and the P-value shows that the null hypothesis that calcium has no effect is more likely at the 5% significance level.

To fully specify the Bayesian approach, we need to choose appropriate priors for the unknown parameters. Due to lack of prior knowledge, we consider $\pi_0 = \pi_1 = 1/2$. Therefore, for decision-making, the hypothesis H_1 is more likely to be selected if $P(H_1 | \mathbf{Y}) > 1/2$, or equivalently, the value of the Bayes factor is larger than 1.

σ_a	1/10	1/3	1/2	1	1.5	2	5
$\operatorname{GBF}[H_1:H_0]$	1.307	1.264	1.358	1.193	0.934	0.746	0.321
$P(H_1 \mid \mathbf{Y})$	0.509	0.558	0.576	0.544	0.483	0.427	0.243

Table 1: Numerical summaries of the GBF with different choice of σ_a .

Gönen et al. (2005) analyze this dataset by using the GBF with $\sigma_a = 1/3$ and obtain that the null hypothesis is less likely because $P(H_1 | \mathbf{Y}) = 0.558$. From a practical viewpoint, we shall be interested in a sensitivity analysis of the hyperparameter σ_a . Numerical results are reported in Table 1. We observe that as σ_a increases, the GBF decreases. When $\sigma_a > 1$, the GBF tends to favor H_0 , whereas it tends to reject H_0 when $\sigma_a \leq 1$. The corresponding posterior probability changes from 0.509 (against H_0) to 0.243 (against H_1) when σ_a changes from 1/10 to 5. This observation shows that the GBF is quite sensitive to the choice of σ_a and that different choice of σ_a may lead to a contradiction in a decisionmarking process. We now employ the PBF with different values of $a \in (-1, -1/2]$. It can be seen from Table 2 that the PBF is quite robust to the choice of a and leads to the same decision. In addition, the conclusion based on the PBF is coincident with the one based on the two-sided P-value.

a	-9/10	-4/5	-3/4	-7/10	-3/5	-1/2
$\operatorname{PBF}[H_1:H_0]$	0.177	0.316	0.375	0.429	0.534	0.606
$P(H_1 \mid \mathbf{Y})$	0.150	0.240	0.273	0.300	0.344	0.377

Table 2: Numerical summaries based on the PBF with different choice of a.

5 Concluding remarks

In this paper, we propose an explicit closed-form Bayes factor for testing the difference between two means from two separate groups of subjects. The proposed approach enjoys several appealing properties. It relies on data only through the classical *t*-statistic and can thus be easily calculated using a simple calculator. It avoids several undesirable properties encountered by the approach due to Gönen et al. (2005). More importantly, it can be easily taught in elementary statistics with an emphasis on Bayesian thinking. We hope that the results of this paper will not only facilitate an intuitive understanding of the relationship between frequentist and Bayesian ideas, but also shed some light on the importance of hyper-prior specifications to students, educators, and researchers.

6 Appendix

Derivation of equation (14): When we consider the Pearson type VI distribution with $\kappa = n_{\delta}$, the Bayes factor in (12) can be expressed as

$$PBF[H_1:H_0] = \frac{n_{\delta}}{B(a+1,b+1)} \int_0^\infty \left[\frac{1+t^2/\nu}{1+t^2/(\nu(1+n_{\delta}\sigma_a^2))}\right]^{(\nu+1)/2} (n_{\delta}\sigma_a^2)^b (1+n_{\delta}\sigma_a^2)^{-a-b-5/2} d\sigma_a^2.$$

With the transformation $\tau = n_{\delta}\sigma_a^2$ and b = (v+1)/2 - a - 5/2, it follows

$$PBF[H_1:H_0] = \frac{1}{B(a+1,b+1)} \int_0^\infty \left[\frac{1+t^2/v}{1+t^2/(v(1+\tau))} \right]^{(v+1)/2} \tau^b (1+\tau)^{-a-b-5/2} d\tau$$
$$= \frac{(1+t^2/v)^{(v+1)/2}}{B(a+1,b+1)} \int_0^\infty \left[1+\frac{t^2}{v} \frac{1}{1+\tau} \right]^{-(v+1)/2} \tau^b (1+\tau)^{-a-b-5/2} d\tau$$
$$= \frac{(1+t^2/v)^{(v+1)/2}}{B(a+1,b+1)} \int_0^\infty \left[1+\tau+\frac{t^2}{v} \right]^{-(v+1)/2} \tau^b (1+\tau)^{(v+1)/2-a-b-5/2} d\tau$$
$$= \frac{(1+t^2/v)^{(v+1)/2}}{B(a+1,b+1)} \int_0^\infty \left[1+\tau+\frac{t^2}{v} \right]^{-(v+1)/2} \tau^b d\tau \text{ since } b = (v+1)/2 - a - 5/2$$
$$= \frac{1}{B(a+1,b+1)} \int_0^\infty \left[1+\frac{\tau}{1+t^2/v} \right]^{-(v+1)/2} \tau^b d\tau.$$

With the transformation $x = \tau/(1 + t^2/v)$, it follows

$$PBF[H_1:H_0] = \frac{(1+t^2/v)^{b+1}}{B(a+1,b+1)} \int_0^\infty (1+x)^{-(v+1)/2} x^b \, dx$$
$$= \frac{B(b+1,(v+1)/2-b-1)}{B(a+1,b+1)} \left(1+\frac{t^2}{v}\right)^{b+1}$$
$$= \frac{\Gamma(v/2)\Gamma(a+3/2)}{\Gamma((v+1)/2)\Gamma(a+1)} \left(1+\frac{t^2}{v}\right)^{(v-2a-2)/2},$$

because of b = (v+1)/2 - a - 5/2. This completes the proof.

References

- Albert, J. (1997). Teaching Bayes' rule: A data-oriented approach. The American Statistician 51(3), pp. 247–253.
- Carlin, B. and T. Louis (2000). Bayes and Empirical Bayes Methods for Data Analysis (2nd ed.). New York, NY: Chapman and Hall.
- Fernández, C., E. Ley, and M. F. J. Steel (2001). Benchmark priors for Bayesian model averaging. Journal of Econometrics 100(2), 381–427.
- Gönen, M., W. O. Johnson, Y. Lu, and P. H. Westfall (2005). The Bayesian two-sample t test. The American Statistician 59(3), 252–257.
- Jeffreys, H. (1961). *Theory of Probability*. Statistics and Computing, 3rd edn. London: Oxford University Press.
- Kass, R. E. and A. E. Raftery (1995). Bayes factors. Journal of the American Statistical Association 90(430), 773–795.
- Liang, F., R. Paulo, G. Molina, M. A. Clyde, and J. O. Berger (2008). Mixtures of g priors for Bayesian variable selection. *Journal of the American Statistical Association 103* (481), 410–423.
- Rouder, J., P. Speckman, D. Sun, R. Morey, and G. Iverson (2009). Bayesian t tests for accepting and rejecting the null hypothesis. *Psychonomic Bulletin & Review 16*(2), 225–237.

- Wang, M. and X. Sun (2014). Bayes factor consistency for one-way random effects model. Communications in Statistics - Theory and Methods 43(23), 5072–5090.
- Weiss, N. (2012). Introductory Statistics. Pearson Education, Inc., Boston, MA.
- Western, B. (1999). Bayesian analysis for sociologists: An introduction. Sociological Methods & Research 28(1), 7–34.
- Wetzels, R., R. P. P. P. Grasman, and E.-J. Wagenmakers (2012). A default Bayesian hypothesis test for ANOVA designs. *The American Statistician* 66(2), 104–111.
- Wulff, S. S. and T. J. Robinson (2014). What is the probability you are a Bayesian? *Journal* of Statistics Education 22(2), 1–22.