

Fast and Simple PCA via Convex Optimization

Dan Garber

Technion - Israel Inst. of Tech.

dangar@tx.technion.ac.il

Elad Hazan

Princeton University

ehazan@cs.princeton.edu

January 18, 2022

Abstract

The problem of principle component analysis (PCA) is traditionally solved by spectral or algebraic methods. We show how PCA could be formulated as a sequence of *convex* optimization problems. This gives rise to a new efficient method for computing the PCA based on recent advances in stochastic methods for convex optimization. In particular, we present running times that improve over the current state-of-the-art.

1 Introduction

Since its introduction by Pearson [10] and Hotelling [5], the principle component analysis technique of finding the subspace of largest variance in the data has become ubiquitous in unsupervised learning, feature generation and data visualization.

For data given as a set of n vectors in $\mathbb{R}^d$, $\mathbf{x}_1, \dots, \mathbf{x}_n$, denote by $\mathbf{X}$ the normalized covariance matrix $\mathbf{X} = \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i^\top$. The PCA method finds the k -dimensional subspace such that the projection of the data onto the subspace has largest variance. Formally, let $\mathbf{W} \in \mathbb{R}^{d \times k}$ be an orthogonal projection matrix, PCA could be formalized as the following optimization problem.

$$\max_{\mathbf{W} \in \mathbb{R}^{d \times k}, \mathbf{W}^T \mathbf{W} = \mathbf{I}} \|\mathbf{XW}\|_F^2, \quad (\text{PCA})$$

where $\|\cdot\|_F$ is the Frobenius norm. Note that the above optimization problem is an inherently non-convex optimization problem even for $k = 1$. For simplicity, henceforth we focus on the problem of finding with high precision only the leading principal component, i.e. on the case $k = 1$.

Finding the leading principal component could be carried out using matrix factorization techniques in time $O(nd^2 + d^3)$, namely by explicitly computing the matrix $\mathbf{X}$ and then computing its singular value decomposition (SVD). However this requires super-linear time and potentially $O(d^2)$ space.

Since super-linear times are prohibitive for large scale machine learning problems, approximation methods have been the focus of recent literature. Iterative eigenvector methods, such as the Lanczos method or power method, can be applied without explicitly

computing the matrix $\mathbf{X}$. These latter methods require the data to be well-conditioned, and the spectral gap, i.e., the distance between largest and second largest eigenvalues of $\mathbf{X}$, to be bounded away from zero.

If we let $\delta > 0$ denote this spectral gap, then the Power and Lanczos methods requires roughly $\delta^{-\gamma}$ passes over the entire data to compute the leading PC which amounts to $\tilde{O}((\lambda_1/\delta)^\gamma nd)$ total time, where $\gamma = 1$ for the Power method and $\gamma = 1/2$ for the Lanczos method.

Thus, iterative methods replace the expensive super-linear operations of matrix factorization by linear-time operations, but trade that for several passes over the data. Depending on the spectral gap of $\mathbf{X}$, the latter methods may also be prohibitively expensive. This motivates the study of methods that retain the best from both worlds: linear time per iteration, as well as doing only a small number (i.e., at most logarithmic in the natural parameters of the problem) of passes over the entire data.

In the convex optimization world, such hybrid results with simple iterations coupled with few passes over the data were obtained in [6, 8, 7]. Recently Shamir [13, 14] made headway in the non-convex spectral world by suggesting a stochastic optimization method for the problem that is based Oja’s randomized Power method [9, 2] with an application of the variance reduction principle demonstrated in [6].

Shamir’s algorithm runs in total time of $\tilde{O}(d(\delta^{-2} + n))$, assuming the availability of a unit vector $\mathbf{w}$ such that $(\mathbf{w}^\top \mathbf{u})^2 = \Omega(1)$, called a “warm start”. Finding such a warm start vector $\mathbf{w}$ could take as much as $\tilde{O}(\delta^{-2.5}d)$ time using existing methods¹, hence the total running time becomes $\tilde{O}(\delta^{-2.5}d + nd)$. Quite remarkably, Shamir’s result separates the dependency on the gap δ and the data size n .

Analyzing the Power method in case of stochastic updates, as done in [2, 13], is much more intricate than analyzing stochastic gradient algorithms for convex optimization, and indeed both the analysis of Oja’s algorithm in [9] and Shamir’s algorithm in [13, 14] are quite elaborate.

In this paper we continue the search for efficient algorithms for PCA and present new algorithms whose significance is two-folded. First, in terms of running time, their worst case running time starting from a “cold-start” is equivalent or better than that of the algorithm of Shamir initialized with a “warm-start”. Second, it is conceptually simpler: we show that computing the leading PC to arbitrary precision is equivalent to solving a small number of well-conditioned convex optimization problems. The latter allows us to immediately apply recent algorithmic advances in stochastic convex optimization, such as the results in [6, 8, 7, 4, 3], to obtain the improved running time.

1.1 Problem setting and main results

Assume we are given a set of n vectors in $\mathbb{R}^d$, $\mathbf{x}_1, \dots, \mathbf{x}_n$, where $n > d$ and let us denote by $\mathbf{X}$ the normalized covariance matrix $\mathbf{X} = \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i^\top$. Assume further that $\mathbf{X}$ has an eigengap of at least δ , i.e. $\lambda_1(\mathbf{X}) - \lambda_2(\mathbf{X}) \geq \delta$, and w.l.o.g. that $\|\mathbf{x}_i\| \leq 1$ for all $i \in [n]$. Our goal is to find a vector $\mathbf{w}$ such that

$$(\mathbf{w}^\top \mathbf{u})^2 \geq 1 - \epsilon,$$

¹Finding a “warm start” could be carried out either by applying iterative methods such as Power and Lanczos methods to the entire data or by applying them only to a small random sample of the data. Since we want to be overall competitive with these methods, we focus on the second option.

, where $\mathbf{u}$ is the leading eigenvector of $\mathbf{X}$ and ϵ is the desired accuracy parameter.

In the rest of the paper we denote the eigenvalues of X in descending order by $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_d$ and by $\mathbf{u} = \mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_d$ the corresponding eigenvectors.

We prove the following theorems. In both theorems we assume we are given an estimate $\hat{\delta}$ such that

$$c_1\delta \leq \hat{\delta} \leq c_2\delta$$

for some universal constants c_1, c_2 which satisfy $c_2 - c_1 = \Theta(1)$. Note that finding such a $\hat{\delta}$ could be done in time that is logarithmic in $1/\delta$.

Theorem 1.1. *Fix $\epsilon > 0, p > 0$. There exists an algorithm that finds with probability at least $1 - p$ a unit vector $\mathbf{w}$ such that $(\mathbf{w}^\top \mathbf{u})^2 \geq 1 - \epsilon$, in total time $\tilde{O}\left(\frac{d}{\delta^2} + nd\right)$.*

Theorem 1.2. *Fix $\epsilon > 0, p > 0$. Assume that $\delta = O(1/\sqrt{n})$. There exists an algorithm that finds with probability at least $1 - p$ a unit vector $\mathbf{w}$ such that $(\mathbf{w}^\top \mathbf{u})^2 \geq 1 - \epsilon$, in total time $\tilde{O}\left(\frac{n^{3/4}d}{\sqrt{\delta}}\right)$.*

Throughout this work we use the notation $\tilde{O}(\cdot)$ to hide poly-logarithmic dependencies on $d, \epsilon^{-1}, p^{-1}, \delta^{-1}$.

Method	Complexity
VRPCA [13, 14]	$\frac{d}{\delta^{5/2}} + nd$
Lanczos	$\sqrt{\frac{\lambda_1}{\delta}}nd$
Theorem 1.1	$\frac{d}{\delta^2} + nd$
Theorem 1.2 ($\delta < n^{-1/2}$)	$\frac{n^{3/4}d}{\sqrt{\delta}}$

Table 1: Comparison with previous results. Note that the result of [13, 14] apply in general only from a “warm-start”. Finding such a warm-start could be carried out efficiently using a sample size of roughly $1/\delta^2$ matrices $\mathbf{x}_i \mathbf{x}_i^\top$ and applying the Lanczos method to this sample.

The rest of this paper is organized as follows. In Section 2 we give relevant preliminaries on classical methods for the computation of the largest eigenvector and related tools, and also preliminaries on convex optimization. In Section 3 we present the core idea for our fast algorithms: a *shrinking inverse power method* algorithm that computes the leading eigenvector after computing only a poly-logarithmic number of matrix-vector products. Based on this algorithm, in Section 4 we present our convex optimization-based eigenvector algorithms that requires to solve only a poly-logarithmic number of smooth and strongly-convex unconstrained optimization problems in order to compute the largest eigenvector. Finally, in Section 5 we combine the result of Section 4 with recent fast stochastic methods for convex optimization to prove Theorems 1.1, 1.2.

2 Preliminaries

2.1 Classical algorithms for the leading eigenvector problem

2.1.1 The Power Method

Our main algorithmic tool for proving the convergence of our method is the classical *Power Method* for approximating the leading eigenvalue and eigenvector of a positive semidefinite matrix.

Algorithm 1 POWER METHOD

```

1: Input: a positive definite matrix  $\mathbf{M}$ 
2: Let  $\mathbf{w}_0$  to be a random unit vector
3: for  $t = 1, 2, \dots$  do
4:    $\mathbf{w}_t \leftarrow \frac{\mathbf{M}\mathbf{w}_{t-1}}{\|\mathbf{M}\mathbf{w}_{t-1}\|}$ 
5: end for

```

In our analysis we will require the following theorem that gives guarantees on the approximation error of Algorithm 1. Both parts of the theorem follows essentially from the same analysis. Part one upper bounds the number of iterations of the algorithm required to achieve a crude approximation to the leading eigenvalue, and part two upper bounds the number of iterations required to achieve a high-precision approximation for the leading eigenvector.

Theorem 2.1. *Let $\mathbf{M}$ be a positive definite matrix and denote its eigenvalues in descending order by $\lambda_1, \lambda_2, \dots, \lambda_d$ and let $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_d$ denote the corresponding eigenvectors. Define $\delta = \lambda_1 - \lambda_2$ and $\kappa = \frac{\lambda_1}{\delta}$. Fix an error tolerance $\epsilon > 0$ and failure probability $p > 0$. Define:*

$$T_{\text{crude}}^{\text{PM}} = \lceil \frac{1}{\epsilon} \ln \left(\frac{18d}{p^2 \epsilon} \right) \rceil, \quad T_{\text{acc}}^{\text{PM}} = \lceil \frac{\kappa}{2} \ln \left(\frac{9d}{p^2 \epsilon} \right) \rceil.$$

Then, with probability $1 - p$ it holds that

1. (crude regime) $\forall t \geq T_{\text{crude}}^{\text{PM}}(\epsilon, p): \mathbf{w}_t^\top \mathbf{M} \mathbf{w}_t \geq (1 - \epsilon) \lambda_1$.
2. (accurate regime) $\forall t \geq T_{\text{acc}}^{\text{PM}}(\kappa, \epsilon, p): (\mathbf{w}_t^\top \mathbf{u}_1)^2 \geq 1 - \epsilon$.

In both cases, the success probability depends only on the random variable $(\mathbf{w}_0^\top \mathbf{u}_1)^2$.

A proof is given in the appendix for completeness.

2.1.2 The Inverse Power Method and Conditioning

As seen in Theorem 2.1 in Subsection 2.1, for a given matrix $\mathbf{X}$, the convergence rate of the *Power Method* algorithms is strongly connected with the condition number $\kappa(\mathbf{X}) = \frac{\lambda_1(\mathbf{X})}{\delta(\mathbf{X})}$ which can be quite large. Consider now the following matrix,

$$\mathbf{M}^{-1} := (\lambda \mathbf{I} - \mathbf{X})^{-1},$$

where λ is a parameter.

Note that if $\mathbf{X}$ is positive semidefinite and $\lambda > \lambda_1(\mathbf{X})$, then $\mathbf{M}^{-1}$ is positive definite. Furthermore, the eigenvectors of $\mathbf{M}^{-1}$ are the same as those of $\mathbf{X}$ and its eigenvalue are (in descending order) $\lambda_i(\mathbf{M}^{-1}) = \frac{1}{\lambda - \lambda_i(\mathbf{X})}$. Thus, if our goal is to compute the leading eigenvector of $\mathbf{X}$ we might as well compute the leading eigenvector of $\mathbf{M}^{-1}$.

The following lemma shows that with a careful choice for the parameter λ , we can make the condition number $\frac{\lambda_1(\mathbf{M}^{-1})}{\delta(\mathbf{M}^{-1})}$ to be much smaller than that of the original matrix $\mathbf{X}$.

Lemma 2.1 (Inverse Conditioning). *Fix a scalar $a > 0$. Let $\mathbf{M}^{-1} = (\lambda \mathbf{I} - \mathbf{X})^{-1}$ such that $\lambda_1(\mathbf{X}) + a\delta(\mathbf{X}) \geq \lambda > \lambda_1(\mathbf{X})$. It holds that*

$$\frac{\lambda_1(\mathbf{M}^{-1})}{\delta(\mathbf{M}^{-1})} \leq 1 + a.$$

Proof. We denote by $\lambda_1, \lambda_2, \delta$ the values $\lambda_1(\mathbf{X}), \lambda_2(\mathbf{X})$ and $\delta(\mathbf{X})$ respectively.

It holds that

$$\lambda_1(\mathbf{M}^{-1}) = \frac{1}{\lambda - \lambda_1}, \quad \lambda_2(\mathbf{M}^{-1}) = \frac{1}{\lambda - \lambda_2}.$$

Thus we have that

$$\begin{aligned} \frac{\lambda_1(\mathbf{M}^{-1})}{\delta(\mathbf{M}^{-1})} &= \frac{1}{\lambda - \lambda_1} \left(\frac{1}{\lambda - \lambda_1} - \frac{1}{\lambda - \lambda_2} \right)^{-1} \\ &= \frac{1}{\lambda - \lambda_1} \left(\frac{1}{\lambda - \lambda_1} - \frac{1}{\lambda - \lambda_1 + \delta} \right)^{-1} = \frac{1}{\lambda - \lambda_1} \left(\frac{\delta}{(\lambda - \lambda_1)(\lambda - \lambda_1 + \delta)} \right)^{-1} \\ &= \frac{\lambda - \lambda_1 + \delta}{\delta} \leq 1 + \frac{a\delta}{\delta} = 1 + a, \end{aligned}$$

and the lemma follows. $\square$

Of course, a problem with the above suggested approach is that we don't know how to set the parameter λ to be close enough to $\lambda_1(\mathbf{X})$. The following simple lemma shows that by approximating the largest eigenvalue of $(\lambda \mathbf{I} - \mathbf{X})^{-1}$, we can derive bounds on the suboptimality gap $\lambda - \lambda_1(\mathbf{X})$, which in turn can be used to better tune λ .

Lemma 2.2. *Fix a positive semidefinite matrix $\mathbf{X}$ and denote the largest eigenvalue by λ_1 . Let $\lambda > \lambda_1$ and consider a unit vector $\mathbf{w}$ such that*

$$\mathbf{w}^\top (\lambda \mathbf{I} - \mathbf{X})^{-1} \mathbf{w} \geq (1 - \epsilon) \lambda_1 ((\lambda \mathbf{I} - \mathbf{X})^{-1}),$$

for some $\epsilon \in (0, 1]$.

Denote $\Delta = \frac{1 - \epsilon}{\mathbf{w}^\top (\lambda \mathbf{I} - \mathbf{X})^{-1} \mathbf{w}}$. Then it holds that

$$(1 - \epsilon)(\lambda - \lambda_1) \leq \Delta \leq \lambda - \lambda_1.$$

Proof. According to our assumption on $\mathbf{w}$ it holds that

$$\lambda_1 ((\lambda \mathbf{I} - \mathbf{X})^{-1}) \geq \mathbf{w}^\top (\lambda \mathbf{I} - \mathbf{X})^{-1} \mathbf{w} \geq (1 - \epsilon) \lambda_1 ((\lambda \mathbf{I} - \mathbf{X})^{-1}).$$

Thus by our definition of Δ and the fact that $\lambda_1 ((\lambda \mathbf{I} - \mathbf{X})^{-1}) = \frac{1}{\lambda - \lambda_1}$, it holds that

$$(1 - \epsilon)(\lambda - \lambda_1) \leq \Delta \leq \lambda - \lambda_1,$$

and the lemma follows. $\square$

2.2 Matrix Inversion via Convex Optimization

2.2.1 Smoothness and strong convexity of continuous functions

Definition 2.1 (smooth function). *We say that a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is β smooth if for all $x, y \in \mathbb{R}^d$ it holds that*

$$\|\nabla f(x) - \nabla f(y)\| \leq \beta \|x - y\|.$$

Definition 2.2 (strongly convex function). *We say that a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is α -strongly convex if for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$ it holds that*

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) + \frac{\alpha}{2} \|\mathbf{x} - \mathbf{y}\|^2.$$

The above definition combined with first order optimality conditions imply that for a α -strongly convex function f , if $\mathbf{x}^* = \arg \min_{\mathbf{x} \in \mathbb{R}^d} f(\mathbf{x})$, then for any $\mathbf{x} \in \mathbb{R}^d$ it holds that

$$f(\mathbf{x}) - f(\mathbf{x}^*) \geq \frac{\alpha}{2} \|\mathbf{x} - \mathbf{x}^*\|^2. \quad (1)$$

Lemma 2.3 (smoothness and strong convexity of quadratic functions). *Consider the function*

$$f(\mathbf{x}) = \frac{1}{2} \mathbf{x}^\top \mathbf{M} \mathbf{x} + \mathbf{b}^\top \mathbf{x},$$

where $\mathbf{M} \in \mathbb{R}^{d \times d}$ is symmetric and $\mathbf{b} \in \mathbb{R}^d$. If, $\mathbf{M} \succeq 0$ then $f(\mathbf{x})$ is $\lambda_1(\mathbf{M})$ -smooth and $\lambda_d(\mathbf{M})$ -strongly convex, where λ_1, λ_d denote the largest and smallest eigenvalues of $\mathbf{M}$ respectively. Otherwise, $f(\mathbf{x})$ is $\|\mathbf{M}\|$ -smooth.

2.2.2 Matrix inversion via convex optimization

In order to apply the Inverse Method discussed in the previous subsection, we need to apply Power Method steps (Algorithm 1) to the matrix $(\lambda \mathbf{I} - \mathbf{X})^{-1}$. Denote $\mathbf{M} = \lambda \mathbf{I} - \mathbf{X}$. Thus on each iteration of the Inverse Method we need to compute a matrix-vector product of the form $\mathbf{M}^{-1} \mathbf{w}$. This requires in general to solve a linear system of equations. However, it could be also approximated arbitrarily well using convex optimization. Consider the following optimization problem:

$$\min_{\mathbf{z} \in \mathbb{R}^d} \{F(\mathbf{z}) := \frac{1}{2} \mathbf{z}^\top \mathbf{M} \mathbf{z} - \mathbf{w}^\top \mathbf{z}\}. \quad (2)$$

By the first order optimality condition, we have that an optimal solution for Problem (2) - $\mathbf{z}^*$ satisfies that $\nabla F(\mathbf{z}^*) = \mathbf{M} \mathbf{z}^* - \mathbf{w} = 0$, meaning,

$$\mathbf{z}^* = \mathbf{M}^{-1} \mathbf{w}.$$

Note that under the assumption that $\lambda > \lambda(\mathbf{X})$ (as stated Subsection 2.1.2) it holds that $\mathbf{M}$ is positive definite and hence invertible.

Most importantly, note that under this assumption on λ it further follows from Lemma 2.3 that $F(\mathbf{z})$ is $\lambda_d(\mathbf{M}) = (\lambda - \lambda_1(\mathbf{X}))$ -strongly convex and $\lambda_1(\mathbf{M}) = (\lambda - \lambda_d(\mathbf{X}))$ -smooth and thus can be solved very efficiently via algorithms for convex minimization.

Since by using algorithms for convex minimization we can only find an approximated-minimizer of $F(\mathbf{z})$, we must discuss the effect of the approximation error on the convergence of the proposed algorithms. As it will turn out, the approximation error that we will care about is the distance $\|\mathbf{z} - \mathbf{z}^*\|$ where $\mathbf{z}$ is an approximated minimizer of f . The following lemma, which follows directly from the strong convexity of f and Eq. (1), ties between the approximation error of a point $\mathbf{z}$ with respect to the function f and the distance to the optimal solution $\mathbf{z}^*$.

Lemma 2.4. *Given a positive semidefinite matrix $\mathbf{X}$, a vector $\mathbf{w}$, a scalar λ such that $\lambda > \lambda_1(\mathbf{X})$ and an error tolerance ϵ , let $\mathbf{M} = \lambda\mathbf{I} - \mathbf{X}$, and denote by $\mathbf{z}^*$ the minimizer of $F(\mathbf{z})$ - as defined in Eq. (2). Then, for any $\mathbf{z}$ it holds that*

$$\|\mathbf{z} - \mathbf{z}^*\| \leq \sqrt{\frac{2(F(\mathbf{z}) - F(\mathbf{z}^*))}{\lambda - \lambda_1(\mathbf{X})}}.$$

2.2.3 Fast stochastic gradient methods for smooth and strongly convex optimization

In this subsection we briefly survey recent development in stochastic optimization algorithms for convex optimization which we leverage in our analysis in order to get the fast rates for PCA.

Consider an optimization problem of the following form

$$\min_{\mathbf{z}} \{F(\mathbf{z}) := \frac{1}{n} \sum_{i=1}^n f_i(\mathbf{z})\} \quad (\text{P})$$

where we assume that each f_i is convex and β -smooth and that the sum $F(\mathbf{z})$ is σ -strongly convex.

We are going to show that the PCA problem could be reduced to solving a series of convex optimization problems that takes the form of Problem (P) (this is actually not a precise statement since in our case each function f_i won't be convex on its own and we will need to address this issue). Thus we are interested in fast algorithms for solving Problem (P).

The standard gradient descent method can solve Problem (P) to ϵ precision in $O((\beta/\sigma) \log(1/\epsilon))$ iterations where each iteration requires to compute the gradient of $F(\mathbf{z})$. Thus the overall time becomes $\tilde{O}(\frac{\beta}{\sigma} n T_g)$ where we denote by T_g the time to evaluate the gradient direction of any of the functions f_i . The dependence on the condition number $\frac{\beta}{\sigma}$ could be dramatically improved without increasing significantly the per-iteration complexity, by using Nesterov's accelerated methods that requires $O(\sqrt{\beta/\sigma} \log(1/\epsilon))$ iterations, and overall time of $\tilde{O}(\sqrt{\beta/\sigma} n T_g)$.

However, both methods could be quite computationally expensive when both $\frac{\beta}{\sigma}$ and n are large.

Another alternative is to use stochastic gradient descent, which on each iteration t , performs a gradient improvement step based on a single function f_{i_t} where i_t is chosen uniformly at random from $[n]$. This single random gradient serves as an estimator for the full gradient. The benefit of this method is that each iteration is extremely cheap -

only requires to evaluate the gradient of a single function. However the convergence rate of this method is roughly $1/(\sigma\epsilon)$ which is ineffective when ϵ is very small. The intuitive reason for the slow convergence is the large variance of the gradient estimator.

Each of the methods mentioned above, the deterministic gradient descent, and the stochastic one has its own caveats. The deterministic gradient method does not consider the special structure of Problem (P) which is given by a sum of functions, and on the other hand, the stochastic gradient method does not exploit the fact that the sum of functions is finite.

Recently, a new family of stochastic gradient descent-based methods was devised, which is tailored to tackling Problem (P) [11, 12, 6, 8, 7, 4, 3]. Roughly speaking, these methods apply cheap stochastic gradient descent update steps, but use the fact the the objective function is given in the form a finite sum, to construct a gradient estimator with reduced variance.

For instance, the SVRG algorithm presented in [6], requires $O(\log(1/\epsilon))$ iterations to reach ϵ accuracy, where each iteration requires computing a single full gradient of $F(\mathbf{z})$ and roughly $O(\beta/\sigma)$ cheap stochastic gradient updates. Thus the total running time becomes $O((\beta/\sigma + n)T_g \log(1/\epsilon))$.

The following theorem summarizes the application of SVRG to solving Problem (P). For details see [6].

Theorem 2.2 (Convergence of SVRG). *Fix $\epsilon > 0$. Assume each $f_i(\mathbf{z})$ is β -smooth and $F(\mathbf{z})$ is σ -strongly convex. Then the SVRG Algorithm detailed in [6] finds in total time $O((n + \beta/\sigma)T_g \log(1/\epsilon))$ a vector $\hat{\mathbf{z}} \in \mathbb{R}^d$ such that*

$$\mathbf{E}[F(\hat{\mathbf{z}})] - \min_{\mathbf{z} \in \mathbb{R}^d} F(\mathbf{z}) \leq \epsilon.$$

3 The Basic Approach: a Shrinking Inverse Power Algorithm

The goal of this section is to present a Power-Method-based algorithm that requires to compute an overall **poly-logarithmic** number of matrix-vector products in order to find an ϵ approximation for the leading eigenvector of a given positive semidefinite matrix $\mathbf{X}$.

We prove the following theorem.

Theorem 3.1. *Assume $\hat{\delta}$ satisfies that $\hat{\delta} \in [\frac{\delta}{2}, 2\delta]$. There exists an implementation for Algorithm 2 that requires computing at most*

$$O\left(\log(d/p) \log(\delta^{-1}) + \log\left(\frac{d}{p\epsilon}\right)\right) = \tilde{O}(1)$$

matrix-vector products of the form $\mathbf{M}^{-1}\mathbf{w}$, where $\mathbf{M}$ is one of the matrices computed during the run of the algorithm ($\mathbf{M}_s$ or $\mathbf{M}_f$) and $\mathbf{w}$ is some vector, such that with probability at least $1 - p$ it holds that the output of the algorithm, the vector $\mathbf{w}_f$, satisfies:

$$(\mathbf{w}_f^\top \mathbf{u})^2 \geq 1 - \epsilon.$$

Algorithm 2 SHRINKING INVERSE POWER METHOD

- 1: Input: matrix $\mathbf{X} \in \mathbb{R}^{n \times n}$ such that $X \succeq 0$, $\lambda_1(\mathbf{X}) \leq 1$, an estimate $\hat{\delta}$ for $\delta(\mathbf{X})$, accuracy parameter $\epsilon \in (0, 1)$
- 2: $\lambda_{(0)} \leftarrow 1 + \hat{\delta}$
- 3: $s \leftarrow 0$
- 4: **repeat**
- 5: $s \leftarrow s + 1$
- 6: Let $\mathbf{M}_s = (\lambda_{(s-1)}\mathbf{I} - \mathbf{X})$
- 7: Apply the POWER METHOD (Algorithm 1) to the matrix $\mathbf{M}_s^{-1}$ to find a unit vector $\mathbf{w}_s$ such that

$$\mathbf{w}_s^\top \mathbf{M}_s^{-1} \mathbf{w}_s \geq \frac{1}{2} \lambda_1(\mathbf{M}_s^{-1})$$

- 8: $\Delta_s \leftarrow \frac{1}{2} \cdot \frac{1}{\mathbf{w}_s^\top \mathbf{M}_s^{-1} \mathbf{w}_s}$
- 9: $\lambda_{(s)} \leftarrow \lambda_{(s-1)} - \frac{\Delta_s}{2}$
- 10: **until** $\Delta_s \leq \hat{\delta}$
- 11: $\lambda_{(f)} \leftarrow \lambda_{(s)}$
- 12: Let $\mathbf{M}_f = (\lambda_{(f)}\mathbf{I} - \mathbf{X})$
- 13: Apply the POWER METHOD (Algorithm 1) to the matrix $\mathbf{M}_f^{-1}$ to find a unit vector $\mathbf{w}_f$ such that

$$(\mathbf{w}_f^\top \mathbf{u})^2 \geq 1 - \epsilon$$

- 14: **return** $\mathbf{w}_f$
-

In order to prove Theorem 3.1 we need a few simple Lemmas.

First, it is important that throughout the run of Algorithm 2, the matrices $\mathbf{M}_s$ and the matrix $\mathbf{M}_f$ will be positive definite (and as a results so are their inverses). This following lemma shows that this is indeed the case.

Lemma 3.1. *For all $s \geq 0$ it holds that $\lambda_{(s)} > \lambda_1$.*

Proof. The proof is by a simple induction. It claim clearly holds for $s = 0$ since by our assumption $\lambda_1 \leq 1$. Suppose now that the claim holds for some iteration s . According to Lemma 2.2 it holds that the value Δ_{s+1} computed on iteration $s + 1$ satisfies that

$$\Delta_{s+1} \leq \lambda_{(s)} - \lambda_1.$$

Hence, according to the algorithms, it holds that

$$\lambda_{(s+1)} = \lambda_{(s)} - \frac{\Delta_{s+1}}{2} \geq \lambda_{(s)} - \frac{\lambda_{(s)} - \lambda_1}{2} = \frac{\lambda_{(s)} + \lambda_1}{2} > \lambda_1,$$

where the last inequality follows from the induction hypothesis. □

The following lemma bounds the number of iterations of the loop in Algorithm 2.

Lemma 3.2. *The repeat-until loop is executed at most $O(\log(\hat{\delta}^{-1}))$ times.*

Proof. Fix an iteration s of the loop. By applying Lemma 2.2 with respect to the unit vector $\mathbf{w}_s$ we have that

$$\Delta_s \geq \frac{1}{2}(\lambda_{(s-1)} - \lambda_1).$$

By the update rule of the algorithm it follows that

$$\begin{aligned} \lambda_{(s)} - \lambda_1 &= \lambda_{(s-1)} - \frac{\Delta_s}{2} - \lambda_1 \geq (\lambda_{(s-1)} - \lambda_1) - \frac{1}{4}(\lambda_{(s-1)} - \lambda_1) \\ &= \frac{3}{4}(\lambda_{(s-1)} - \lambda_1). \end{aligned}$$

Thus, after at most $T = \lceil \log_{3/4} \left(\frac{\hat{\delta}}{\lambda_{(0)} - \lambda_1} \right) \rceil = O(\log(\hat{\delta}^{-1}))$ (using our choice of $\lambda_{(0)}$) iterations, we arrive at a value $\lambda_{(T)}$ which satisfies $\lambda_{(T)} - \lambda_1 \leq \hat{\delta}$. By Lemma 2.2 it follows that in the following iteration it will hold that $\Delta_{T+1} \leq \hat{\delta}$ and the loop will end. Hence, the overall number of iterations is at most $T + 1 = O(\log(\hat{\delta}^{-1}))$. $\square$

Finally, the following Lemma gives approximation guarantees of the estimate $\lambda_{(f)}$.

Lemma 3.3. *Suppose that all executions of the Power Method in Algorithm 2 are successful. Then it holds that*

$$\lambda_1 + \frac{3\hat{\delta}}{2} \geq \lambda_{(f)} \geq \lambda_1 + \frac{\hat{\delta}}{4}. \quad (3)$$

Proof. Denote by s_f the last iteration of the loop in Algorithm 2, and note that using this notation we have that $\lambda_{(f)} = \lambda_{(s_f)}$ and that $\Delta_{s_f} \leq \hat{\delta}$. Using Lemma 2.2, we thus have that

$$\lambda_{(f)} - \lambda_1 = \lambda_{(s_f-1)} - \frac{\Delta_{s_f}}{2} - \lambda_1 \leq 2\Delta_{s_f} - \frac{\Delta_{s_f}}{2} = \frac{3}{2}\Delta_{s_f} \leq \frac{3}{2}\hat{\delta},$$

which gives the first part of the lemma.

For the second part, using Lemma 2.2 again, we have that

$$\begin{aligned} \lambda_{(f)} - \lambda_1 &= \lambda_{(s_f-1)} - \frac{\Delta_{s_f}}{2} - \lambda_1 \geq \lambda_{(s_f-1)} - \lambda_1 - \frac{1}{2}(\lambda_{(s_f-1)} - \lambda_1) \\ &= \frac{1}{2}(\lambda_{(s_f-1)} - \lambda_1). \end{aligned} \quad (4)$$

In case $s_f = 1$, then by our choice of $\lambda_{(0)}$ we have that $\lambda_{(0)} - \lambda_1 \geq \hat{\delta}$, and the lemma follows. Otherwise, by unfolding Eq. (4) one more time, we have that

$$\lambda_{(f)} - \lambda_1 \geq \frac{1}{4}(\lambda_{(s_f-2)} - \lambda_1) \geq \frac{\Delta_{(s_f-1)}}{4} > \frac{\hat{\delta}}{4},$$

where the second inequality follows from Lemma 2.2 and the last inequality follows from the stopping condition of the loop. $\square$

We can now prove Theorem 3.1.

Proof. First a note regarding the success probability of the invocations of the *Power Method* algorithm in Algorithm 2: since, as stated in Theorem 2.1, the success of the PM algorithm depends only on the magnitude of $(\mathbf{w}_0^\top \mathbf{u})^2$, and all matrices $\mathbf{M}_s, \mathbf{M}_f$ in Algorithm 2 have the same leading eigenvector, as long as all invocations use the same random initial vector and number of steps that guarantees success with probability $1 - p$, they all succeed together with probability at least $1 - p$.

Let us now assume that all executions of the *Power Method* algorithm in Algorithm 1 were successful.

According to Lemma 3.2, the loop is executed at most $O(\log(\hat{\delta}^{-1})) = O(\log(\delta^{-1}))$ times (following our assumption on $\hat{\delta}$). Each iteration s of the loop requires to invoke the *Power Method* to approximate $\lambda_1(\mathbf{M})$ up to a factor of $1/2$ which according to Theorem 2.1, requires computing $T_{crude}^{\text{PM}}(1/2, p) = O(\log(d/p))$ matrix-vector products, in order to succeed with probability at least $1 - p$. Thus the overall number of matrix-vector products computed during the loop is $O(\log(d/p) \log(\delta^{-1}))$.

According to lemma 3.3 it holds that $\lambda_{(f)} - \lambda_1 \leq \frac{3}{2}\hat{\delta} \leq 3\delta$. Thus, according to Lemma 2.1, we have that

$$\kappa(\mathbf{M}_f^{-1}) = \frac{\lambda_1(\mathbf{M}_f^{-1})}{\delta(\mathbf{M}_f^{-1})} \leq 4 = O(1).$$

Thus, in the final invocation of the PM algorithm, it requires at most $T_{acc}^{\text{PM}}(4, \epsilon, p) = O\left(\log\left(\frac{d}{p\epsilon}\right)\right)$ matrix-vector products to compute $\mathbf{w}_f$ as desired with probability at least $1 - p$.

Thus the overall number of matrix-vector products is

$$O\left(\log(d/p) \log(\delta^{-1}) + \log\left(\frac{d}{p\epsilon}\right)\right).$$

□

4 A Convex Optimization-based Eigenvector Algorithm

In this section we present our algorithm for approximating the largest eigenvector of a given matrix based on convex optimization. The algorithm is based on Algorithm 2 from the previous section, but replaces explicit computation of products between vectors and inverse matrices, with solving convex optimization problems, as detailed in Subsection 2.2.

Towards this end, we assume that we are given access to an algorithm - $\mathcal{A}$ for solving problems of the following structure:

$$\min_{\mathbf{z} \in \mathbb{R}^d} \{F_{\mathbf{w}, \lambda}(\mathbf{z}) := \frac{1}{2} \mathbf{z}^\top (\lambda \mathbf{I} - \mathbf{X}) \mathbf{z} - \mathbf{w}^\top \mathbf{z}\}, \quad (5)$$

where $\mathbf{X}$ is positive definite, $\lambda > \lambda_1(\mathbf{X})$ and $\mathbf{w}$ is some vector. Note that under these conditions, the function $F_{\mathbf{w}, \lambda}(\mathbf{z})$ is strongly convex. Note also that the minimizer of $F_{\mathbf{w}, \lambda}(\mathbf{z}) - \mathbf{z}^*$ is given by $\mathbf{z}^* = (\lambda \mathbf{I} - \mathbf{X})^{-1} \mathbf{w}$, and thus solving Problem (5) is equivalent to computing a product between a vector and an inverse matrix.

There are a few issues with our approach that require delicate care however: 1) we need to pay close attention that the convex optimization problems are well-conditioned and 2) since now we use a numerical procedure to compute the matrix-vector products, we have approximation errors that we need to consider.

Algorithm 3 LEADING EIGENVECTOR VIA CONVEX OPTIMIZATION

```

1: Input: matrix  $\mathbf{X} \in \mathbb{R}^{n \times n}$  such that  $\mathbf{X} \succeq 0$ ,  $\lambda_1(\mathbf{X}) \leq 1$ , an estimate  $\hat{\delta}$  for  $\delta(\mathbf{X})$ ,
   accuracy parameter  $\epsilon \in (0, 1)$ , failure probability parameters  $p$ 
2:  $\lambda_{(0)} \leftarrow 1 + \hat{\delta}$ 
3: Set:  $m_1 \leftarrow T_{crude}^{\text{PM}}(1/4, p)$ ,  $m_2 \leftarrow T_{acc}^{\text{PM}}(3, \epsilon/2, p)$ 
4: Set:  $\tilde{\epsilon} \leftarrow \dots$ 
5: Let  $\hat{\mathbf{w}}_0$  be a random unit vector
6:  $s \leftarrow 0$ 
7: repeat
8:    $s \leftarrow s + 1$ 
9:   Let  $\mathbf{M}_s = (\lambda_{(s-1)} \mathbf{I} - \mathbf{X})$ 
10:  for  $t = 1 \dots m_1 + 1$  do
11:    Apply Algorithm  $\mathcal{A}$  to find a vector  $\hat{\mathbf{w}}_t$  such that  $\|\hat{\mathbf{w}}_t - \mathbf{M}_s^{-1} \hat{\mathbf{w}}_{t-1}\| \leq \tilde{\epsilon}$ 
12:  end for
13:   $\mathbf{w}_s \leftarrow \frac{\hat{\mathbf{w}}_{m_1}}{\|\hat{\mathbf{w}}_{m_1}\|}$ 
14:  Apply Algorithm  $\mathcal{A}$  to find a vector  $\mathbf{v}_s$  such that  $\|\mathbf{v}_s - \mathbf{M}_s^{-1} \mathbf{w}_s\| \leq \tilde{\epsilon}$ 
15:   $\Delta_s \leftarrow \frac{1}{2} \cdot \frac{1}{\mathbf{w}_s^\top \mathbf{v}_s - \tilde{\epsilon}}$ 
16:   $\lambda_{(s)} \leftarrow \lambda_{(s-1)} - \frac{\Delta_s}{2}$ 
17: until  $\Delta_s \leq \hat{\delta}$ 
18:  $\lambda_{(f)} \leftarrow \lambda_{(s)}$ 
19: Let  $\mathbf{M}_f = (\lambda_{(f)} \mathbf{I} - \mathbf{X})$ 
20: for  $t = 1 \dots m_2$  do
21:  Apply Algorithm  $\mathcal{A}$  to find a vector  $\hat{\mathbf{w}}_t$  such that  $\|\hat{\mathbf{w}}_t - \mathbf{M}_f^{-1} \hat{\mathbf{w}}_{t-1}\| \leq \tilde{\epsilon}$ 
22: end for
23: return  $\mathbf{w}_f \leftarrow \frac{\hat{\mathbf{w}}_{m_2}}{\|\hat{\mathbf{w}}_{m_2}\|}$ 

```

4.1 Power Method with noisy updates

Lemma 4.1 (Power Method with approximated matrix-vector products). *Let $\mathbf{M}$ be a positive definite matrix with largest eigenvalue λ_1 and smallest eigenvalue λ_d . Fix an accuracy parameter $\epsilon > 0$. Let $\mathbf{w}$ be an arbitrary unit vector and consider the two sequences $\{\mathbf{w}_t^*\}_{t=0}^\infty$ and $\{\mathbf{w}_t\}_{t=0}^\infty$ defined as follows:*

$$\begin{aligned}
\mathbf{w}_0^* &= \hat{\mathbf{w}}_0^* = \mathbf{w}, \quad \forall t \geq 1 : \hat{\mathbf{w}}_t^* \leftarrow \mathbf{M} \hat{\mathbf{w}}_{t-1}^*, \quad \mathbf{w}_t^* \leftarrow \frac{\hat{\mathbf{w}}_t^*}{\|\hat{\mathbf{w}}_t^*\|}, \\
\mathbf{w}_0 &= \hat{\mathbf{w}}_0 = \mathbf{w}, \quad \forall t \geq 1 : \hat{\mathbf{w}}_t \text{ satisfies that } \|\hat{\mathbf{w}}_t - \mathbf{M} \hat{\mathbf{w}}_{t-1}\| \leq \epsilon, \quad \mathbf{w}_t \leftarrow \frac{\hat{\mathbf{w}}_t}{\|\hat{\mathbf{w}}_t\|}.
\end{aligned}$$

Denote:

$$\Gamma(\mathbf{M}, t) := \frac{2}{\lambda_d^t} \begin{cases} t & \text{if } \lambda_1 = 1 \\ \frac{\lambda_1^t - 1}{\lambda_1 - 1} & \text{if } \lambda_1 \neq 1 \end{cases}, \quad \hat{\Gamma}(\mathbf{M}, t) := \begin{cases} t & \text{if } \lambda_1 = 1 \\ \frac{\lambda_1^t - 1}{\lambda_1 - 1} & \text{if } \lambda_1 \neq 1 \end{cases}.$$

Then it holds that for all $t \geq 0$,

$$\|\hat{\mathbf{w}}_t - \hat{\mathbf{w}}_t^*\| \leq \epsilon \cdot \hat{\Gamma}(\mathbf{M}, t)$$

and

$$\|\mathbf{w}_t - \mathbf{w}_t^*\| \leq \epsilon \cdot \Gamma(\mathbf{M}, t)$$

Proof. First, observe that:

$$\begin{aligned} \|\hat{\mathbf{w}}_{t+1} - \hat{\mathbf{w}}_{t+1}^*\| &= \|\hat{\mathbf{w}}_{t+1} - \mathbf{M}\hat{\mathbf{w}}_t + \mathbf{M}\hat{\mathbf{w}}_t - \hat{\mathbf{w}}_{t+1}^*\| \leq \|\hat{\mathbf{w}}_{t+1} - \mathbf{M}\hat{\mathbf{w}}_t\| + \|\mathbf{M}\hat{\mathbf{w}}_t - \mathbf{M}\hat{\mathbf{w}}_t^*\| \\ &\leq \epsilon + \lambda_1 \cdot \|\hat{\mathbf{w}}_t - \hat{\mathbf{w}}_t^*\|. \end{aligned}$$

In case $\lambda_1 = 1$, we clearly have that

$$\|\hat{\mathbf{w}}_{t+1} - \hat{\mathbf{w}}_{t+1}^*\| \leq (t+1)\epsilon + \|\hat{\mathbf{w}}_0 - \hat{\mathbf{w}}_0^*\| = (t+1)\epsilon. \quad (6)$$

Otherwise, in case $\lambda_1 \neq 1$, by a simple algebraic manipulation we have that

$$\|\hat{\mathbf{w}}_{t+1} - \hat{\mathbf{w}}_{t+1}^*\| + \frac{\epsilon}{\lambda_1 - 1} \leq \lambda_1 \cdot \|\hat{\mathbf{w}}_t - \hat{\mathbf{w}}_t^*\| + \frac{\epsilon\lambda_1}{\lambda_1 - 1} = \lambda_1 \left(\|\hat{\mathbf{w}}_t - \hat{\mathbf{w}}_t^*\| + \frac{\epsilon}{\lambda_1 - 1} \right).$$

It thus follows that for all $t \geq 0$,

$$\|\hat{\mathbf{w}}_{t+1} - \hat{\mathbf{w}}_{t+1}^*\| \leq \lambda_1^{t+1} \left(\|\hat{\mathbf{w}}_0 - \hat{\mathbf{w}}_0^*\| + \frac{\epsilon}{\lambda_1 - 1} \right) - \frac{\epsilon}{\lambda_1 - 1} = \frac{\epsilon}{\lambda_1 - 1} (\lambda_1^{t+1} - 1). \quad (7)$$

We now have that for all $t \geq 1$ it holds that

$$\begin{aligned} \|\mathbf{w}_t - \mathbf{w}_t^*\| &= \left\| \frac{\hat{\mathbf{w}}_t}{\|\hat{\mathbf{w}}_t\|} - \frac{\hat{\mathbf{w}}_t^*}{\|\hat{\mathbf{w}}_t^*\|} \right\| = \left\| \frac{\hat{\mathbf{w}}_t}{\|\hat{\mathbf{w}}_t\|} - \frac{\hat{\mathbf{w}}_t}{\|\hat{\mathbf{w}}_t^*\|} + \frac{\hat{\mathbf{w}}_t}{\|\hat{\mathbf{w}}_t^*\|} - \frac{\hat{\mathbf{w}}_t^*}{\|\hat{\mathbf{w}}_t^*\|} \right\| \\ &\leq \left\| \frac{\hat{\mathbf{w}}_t}{\|\hat{\mathbf{w}}_t^*\|} - \frac{\hat{\mathbf{w}}_t^*}{\|\hat{\mathbf{w}}_t^*\|} \right\| + \left\| \frac{\hat{\mathbf{w}}_t}{\|\hat{\mathbf{w}}_t\|} - \frac{\hat{\mathbf{w}}_t}{\|\hat{\mathbf{w}}_t^*\|} \right\| \\ &= \frac{1}{\|\hat{\mathbf{w}}_t^*\|} \cdot \|\hat{\mathbf{w}}_t - \hat{\mathbf{w}}_t^*\| + \|\hat{\mathbf{w}}_t\| \cdot \left| \frac{1}{\|\hat{\mathbf{w}}_t\|} - \frac{1}{\|\hat{\mathbf{w}}_t^*\|} \right| \\ &= \frac{1}{\|\hat{\mathbf{w}}_t^*\|} \cdot \|\hat{\mathbf{w}}_t - \hat{\mathbf{w}}_t^*\| + \frac{|\|\hat{\mathbf{w}}_t^*\| - \|\hat{\mathbf{w}}_t\||}{\|\hat{\mathbf{w}}_t^*\|} \leq \frac{2\|\hat{\mathbf{w}}_t - \hat{\mathbf{w}}_t^*\|}{\|\hat{\mathbf{w}}_t^*\|}, \end{aligned}$$

where the last inequality follows from the triangle inequality.

Now, since $\hat{\mathbf{w}}_t^* = \mathbf{M}^t \mathbf{w}$ and $\mathbf{M} \succeq \lambda_d \mathbf{I}$ it follows that

$$\|\hat{\mathbf{w}}_t^*\| \geq \lambda_d^t \cdot \|\mathbf{w}\| = \lambda_d^t.$$

Combining this with Eq. (7) and (6), we have that for all $t \geq 1$,

$$\|\mathbf{w}_t - \mathbf{w}_t^*\| \leq \frac{2\epsilon}{\lambda_d^t} \begin{cases} t & \text{if } \lambda_1 = 1; \\ \frac{\lambda_1^t - 1}{\lambda_1 - 1} & \text{if } \lambda_1 \neq 1. \end{cases}$$

Thus the lemma follows. $\square$

The following corollary is a consequence of the convergence result for the Power Method (Theorem 2.1) and Lemma 4.1.

Theorem 4.1 (Convergence of the Power Method with noisy updates). *Fix $\epsilon > 0$ and $p > 0$. Let $\mathbf{M}$ be a positive definite matrix with largest eigenvalue λ_1 and eigengap $\delta > 0$. Consider a sequence of vectors $\{\mathbf{w}_t\}_{t=0}^\tau$ where $\mathbf{w}_0$ is a random unit vector, and for all $\tau \geq t \geq 1$ it holds that $\|\mathbf{w}_t - \mathbf{M}\mathbf{w}_{t-1}\| \leq \tilde{\epsilon} := \frac{\epsilon}{4\Gamma(\mathbf{M}, \tau)}$. Then the following two guarantees hold:*

1. *If $\tau \geq T_{\text{crude}}^{\text{PM}}(\epsilon/2, p)$ then with probability at least $1 - p$ it holds that*

$$\mathbf{w}_\tau^\top \mathbf{M} \mathbf{w}_\tau \geq (1 - \epsilon) \lambda_1.$$

2. *If $\tau \geq T_{\text{acc}}^{\text{PM}}(\kappa, \epsilon/2, p)$ then with probability at least $1 - p$ it holds that*

$$(\mathbf{w}_\tau^\top \mathbf{u}_1)^2 \geq 1 - \epsilon,$$

where $\mathbf{u}_1$ is the largest eigenvector of $\mathbf{M}$.

In both cases, the probability of success depends only on the random variable $(\mathbf{w}_0^\top \mathbf{u}_1)^2$.

Proof. For the first item, note that

$$\mathbf{w}_\tau^\top \mathbf{M} \mathbf{w}_\tau = \mathbf{w}_\tau^\top \mathbf{M} \mathbf{w}_\tau^* + (\mathbf{w}_\tau^\top \mathbf{M} \mathbf{w}_\tau - \mathbf{w}_\tau^{*\top} \mathbf{M} \mathbf{w}_\tau^*).$$

Since $\mathbf{M}$ is positive definite, the function $f(\mathbf{w}) = \mathbf{w}^\top \mathbf{M} \mathbf{w}$ is convex and we have that

$$\begin{aligned} |\mathbf{w}_\tau^\top \mathbf{M} \mathbf{w}_\tau - \mathbf{w}_\tau^{*\top} \mathbf{M} \mathbf{w}_\tau^*| &\leq 2 \max\{(\mathbf{M} \mathbf{w}_\tau)^\top (\mathbf{w}_\tau - \mathbf{w}_\tau^*), (\mathbf{M} \mathbf{w}_\tau^*)^\top (\mathbf{w}_\tau^* - \mathbf{w}_\tau)\} \\ &\leq 2 \lambda_1(\mathbf{M}) \cdot \|\mathbf{w}_\tau - \mathbf{w}_\tau^*\|. \end{aligned}$$

Thus we have that

$$\begin{aligned} \mathbf{w}_\tau^\top \mathbf{M} \mathbf{w}_\tau &\geq \mathbf{w}_\tau^{*\top} \mathbf{M} \mathbf{w}_\tau^* - 2 \lambda_1(\mathbf{M}) \cdot \|\mathbf{w}_\tau - \mathbf{w}_\tau^*\| \\ &\geq \mathbf{w}_\tau^{*\top} \mathbf{M} \mathbf{w}_\tau^* - \frac{\epsilon}{2} \lambda_1(\mathbf{M}), \end{aligned}$$

where the last inequality follows from Lemma 4.1 and our choice of $\tilde{\epsilon}$.

Now, by our choice of τ and Theorem 2.1, we have that with probability at least $1 - p$ it holds that

$$\mathbf{w}_\tau^\top \mathbf{M} \mathbf{w}_\tau \geq (1 - \epsilon/2) \lambda_1(\mathbf{M}) - \frac{\epsilon}{2} \lambda_1(\mathbf{M}) = (1 - \epsilon) \lambda_1(\mathbf{M}).$$

For the second item, note that

$$\begin{aligned} (\mathbf{w}_\tau^\top \mathbf{u}_1)^2 &= (\mathbf{w}_\tau^{*\top} \mathbf{u}_1 + (\mathbf{w}_\tau - \mathbf{w}_\tau^*)^\top \mathbf{u}_1)^2 \geq (\mathbf{w}_\tau^{*\top} \mathbf{u}_1)^2 - 2 \|\mathbf{w}_\tau - \mathbf{w}_\tau^*\| \\ &\geq (\mathbf{w}_\tau^{*\top} \mathbf{u}_1)^2 - \frac{\epsilon}{2}, \end{aligned}$$

where the last inequality follows again from Lemma 4.1 and our choice of $\tilde{\epsilon}$.

Again, by our definition of τ and Theorem 2.1, we have that with probability at least $1 - p$ it holds that

$$(\mathbf{w}_\tau^\top \mathbf{u}_1)^2 \geq 1 - \epsilon/2 - \epsilon/2 = 1 - \epsilon.$$

□

4.2 Convergence of Algorithm 3

The key step in the analysis of Algorithm 3 is to show that if all numerical computations are carried out with sufficiently small error, then Algorithm 3 successfully simulates the Power Method-based Algorithm 2. A main ingredient in Algorithm 2 is the computations of the values Δ_s which are used in turn to update the estimates $\lambda_{(s)}$. Our use of the values Δ_s in Algorithm 2 was through the approximation guarantees they provided for the gap $\lambda_{(s-1)} - \lambda_1$, i.e., $\frac{1}{2}(\lambda_{(s-1)} - \lambda_1) \leq \Delta_s \leq \lambda_{(s-1)} - \lambda_1$. The following lemma shows that with a correct choice for the numerical error parameter $\tilde{\epsilon}$, these guarantees also hold for the values Δ_s computed in Algorithm 3.

Lemma 4.2. *Suppose that $\hat{\delta} \in [\frac{\delta}{2}, \frac{3\delta}{4}]$ and that $\tilde{\epsilon} \leq \frac{1}{16} \left(\frac{\hat{\delta}}{8}\right)^{m_1+1}$. Then with probability at least $1 - p$ it holds that for any s , the update of the variables $\Delta_s, \lambda_{(s)}$ in Algorithm 3 is equivalent to that in Algorithm 2. In particular, for all $s \geq 1$ it holds that*

$$\frac{1}{2}(\lambda_{(s-1)} - \lambda_1) \leq \Delta_s \leq \lambda_{(s-1)} - \lambda_1.$$

Proof. For clarity we refer by $\Delta_s, \lambda_{(s)}$ to the values computed in Algorithm 3 and by $\tilde{\Delta}_s, \tilde{\lambda}_{(s)}$ to the corresponding values computed in Algorithm 2 from Section 3. The proof of the lemma is by induction on s . For the base case $s = 0$, clearly $\lambda_{(0)} = \tilde{\lambda}_{(0)}$ and both $\Delta_0, \tilde{\Delta}_0$ are undefined. Consider now some $s \geq 1$.

Suppose for now that $\tilde{\epsilon} \leq \min\{\frac{1}{16\Gamma(\mathbf{M}_s^{-1}, m_1)}, \frac{\lambda_1(\mathbf{M}_s^{-1})}{8}\}$. Then it follows from Theorem 4.1 and our choice of m_1 that with probability at least $1 - p$ that

$$\mathbf{w}_s^\top \mathbf{M}_s^{-1} \mathbf{w}_s \geq \frac{3}{4} \lambda_1(\mathbf{M}_s^{-1}). \quad (8)$$

By our definition of the vector $\mathbf{v}_s$ and the Cauchy-Schwartz inequality it holds that

$$\mathbf{w}_s^\top \mathbf{v}_s = \mathbf{w}_s^\top \mathbf{M}_s^{-1} \mathbf{w}_s + \mathbf{w}_s^\top (\mathbf{v}_s - \mathbf{M}_s^{-1} \mathbf{w}_s) \in [\mathbf{w}_s^\top \mathbf{M}_s^{-1} \mathbf{w}_s - \tilde{\epsilon}, \mathbf{w}_s^\top \mathbf{M}_s^{-1} \mathbf{w}_s + \tilde{\epsilon}]. \quad (9)$$

Thus, combining Eq. (8) and (9), we have that

$$\mathbf{w}_s^\top \mathbf{v}_s - \tilde{\epsilon} \in [\mathbf{w}_s^\top \mathbf{M}_s^{-1} \mathbf{w}_s - 2\tilde{\epsilon}, \mathbf{w}_s^\top \mathbf{M}_s^{-1} \mathbf{w}_s] \subseteq [3\lambda_1(\mathbf{M}_s^{-1})/4 - 2\tilde{\epsilon}, \lambda_1(\mathbf{M}_s^{-1})].$$

By our choice of $\tilde{\epsilon}$ it follows that

$$\mathbf{w}_s^\top \mathbf{v}_s - \tilde{\epsilon} \in [\lambda_1(\mathbf{M}_s^{-1})/2, \lambda_1(\mathbf{M}_s^{-1})].$$

Thus, the computation of the value of Δ_s , and as a result the computation of $\lambda_{(s)}$ is identical to that of $\tilde{\Delta}_s, \tilde{\lambda}_{(s)}$, and the claim follows.

It only remains to give an explicit bound on $\tilde{\epsilon}$. For this we need to upper bound $\Gamma(\mathbf{M}_s^{-1}, m_1)$ for all values of s . Recall that from the results of Section 3 it follows that the values $\lambda_{(s)}$ are monotonically non-increasing and lower-bounded by $\lambda_{(f)} \geq \lambda_1 + \hat{\delta}/4$. By our assumption of $\hat{\delta}$ we have that

$$\begin{aligned} \lambda_1(\mathbf{M}_s^{-1}) &= \frac{1}{\lambda_{(s-1)} - \lambda_1} \geq \frac{1}{\lambda_{(0)} - \lambda_1} = \frac{1}{1 + \hat{\delta} - \lambda_1} \geq \frac{1}{1 + \hat{\delta} - \lambda_1} \geq \frac{1}{1 + \frac{3\delta}{4} - \delta} \\ &> 1 + \frac{\delta}{4} \geq 1 + \frac{\hat{\delta}}{3}, \end{aligned} \quad (10)$$

where the third inequality holds since by definition of δ it follows that $\lambda_1 = \lambda_2 + \delta \geq \delta$.

Fix a natural number t . By the definition of $\Gamma(\mathbf{M}, t)$ (see Lemma 4.1) we have that

$$\begin{aligned}\Gamma(\mathbf{M}_s^{-1}, t) &= \frac{2}{\lambda_d(\mathbf{M}_s^{-1})} \cdot \frac{\lambda_1(\mathbf{M}_s^{-1})^t - 1}{\lambda_1(\mathbf{M}_s^{-1}) - 1} \leq \frac{2}{\lambda_1(\mathbf{M}_s^{-1}) - 1} \left(\frac{\lambda_1(\mathbf{M}_s^{-1})}{\lambda_d(\mathbf{M}_s^{-1})} \right)^t \\ &= \frac{6}{\hat{\delta}} \left(\frac{\lambda_{(s-1)} - \lambda_d}{\lambda_{(s-1)} - \lambda_1} \right)^t \leq \frac{6}{\hat{\delta}} \left(\frac{\lambda_{(0)}}{\lambda_{(f)} - \lambda_1} \right)^t \\ &\leq \frac{6}{\hat{\delta}} \left(\frac{1 + \hat{\delta}}{\hat{\delta}/4} \right)^t = \frac{6}{\hat{\delta}} \left(4 + \frac{4}{\hat{\delta}} \right)^t < \left(\frac{8}{\hat{\delta}} \right)^{t+1}.\end{aligned}\tag{11}$$

Thus it follows that we can take $\tilde{\epsilon} \leq \min\left\{\frac{1}{16} \left(\frac{\hat{\delta}}{8}\right)^{m_1+1}, \frac{1+\hat{\delta}/3}{8}\right\} = \frac{1}{16} \left(\frac{\hat{\delta}}{6}\right)^{m_1+1}$.

□

Theorem 4.2 (Convergence of Algorithm 3). *Suppose that $\hat{\delta} \in [\delta/2, 2\delta]$. Fix $\epsilon > 0$. Let $m_1 \geq T_{\text{crude}}^{\text{PM}}(1/8, p)$ and $m_2 \geq T_{\text{acc}}^{\text{PM}}(3, \epsilon/2, p)$. Suppose that $\tilde{\epsilon}$, satisfies that*

$$\tilde{\epsilon} \leq \min\left\{\frac{1}{16} \left(\frac{\hat{\delta}}{8}\right)^{m_1+1}, \frac{\epsilon}{4} \left(\frac{\hat{\delta}}{8}\right)^{m_2+1}\right\}.$$

Then, with probability at least $1 - p$ it holds that the output of Algorithm 3, the unit vector $\mathbf{w}_f$ satisfies that

$$(\mathbf{w}_f^\top \mathbf{u}_1)^2 \geq 1 - \epsilon,$$

and the total number of calls to the convex optimization oracle $\mathcal{A}$ is

$$O\left(\log(d/p) \log(\delta^{-1}) + \log\left(\frac{d}{p\epsilon}\right)\right)$$

Proof. By our choice of $\tilde{\epsilon}$ and Lemma 4.2 it follows that we can invoke the results of Section 3. By Lemma 3.2, we can upper bound the number of iterations made by the *repeat-until* loop by $O(\log(\delta^{-1}))$. Since each iteration of the loop requires $m_1 + 1$ calls to the optimization oracle $\mathcal{A}$, the overall number of calls to $\mathcal{A}$ during the loop is $O(m_1 \log(\delta^{-1}))$.

By Lemma 3.3 we have that the final estimate $\lambda_{(f)}$ satisfies that $\lambda_{(f)} - \lambda_1 \leq \frac{3\hat{\delta}}{2} \leq \frac{9\delta}{8} < 2\delta$. Thus, by Lemma 2.1 we have that

$$\kappa(\mathbf{M}_f^{-1}) = \frac{\lambda_1(\mathbf{M}_f^{-1})}{\delta(\mathbf{M}_f^{-1})} \leq 3.\tag{12}$$

Suppose now that $\tilde{\epsilon} \leq \frac{\epsilon}{4\Gamma(\mathbf{M}_f^{-1}, m_2)}$. By our choice of m_2 and Eq. (12), it follows from Theorem 4.1 that with probability at least $1 - p$ indeed $(\mathbf{w}_f^\top \mathbf{u})^2 \geq 1 - \epsilon$.

The number of calls to the oracle $\mathcal{A}$ in this final part is m_2 . Thus overall number of calls to $\mathcal{A}$ in Algorithm 3 is $O(m_1 \log(\delta^{-1}) + m_2)$.

It remains to lower-bound the term $\frac{\epsilon}{4\Gamma(\mathbf{M}_f^{-1}, m_2)}$.

Since the values $\lambda_{(s)}$ are decreasing as a function of s , it follows that $\lambda_{(f)} \leq \lambda_{(0)}$. We can continue as in Eq. (10), (11) and get that $\Gamma(\mathbf{M}_f^{-1}, m_2) \leq \left(\frac{8}{\hat{\delta}}\right)^{m_2+1}$, which agrees with the bound on $\tilde{\epsilon}$ stated in the theorem.

□

In order to analyze the arithmetic complexity of Algorithm 3 using a specific implementation for the optimization oracle $\mathcal{A}$, it is not only important to bound the number of calls to $\mathcal{A}$ (as done in Theorem 4.2), but to also bound important parameters of the optimization problem (5) that naturally arise when considering the arithmetic complexity of different implementations for $\mathcal{A}$. For this issue we have the following lemma.

The following lemma follows directly from the discussion in Section 3 and the assumptions stated in Theorem 4.2.

Lemma 4.3. *Let $\lambda, \mathbf{w}$ be such that during the run of Algorithm 3, the optimization oracle $\mathcal{A}$ is applied to the minimization of the function*

$$F(\mathbf{z}) = \frac{1}{2} \mathbf{z}^\top (\lambda \mathbf{I} - \mathbf{X}) \mathbf{z} - \mathbf{w}^\top \mathbf{z}.$$

Then, under the conditions stated in Theorem 4.2 it holds that

1. $F(\mathbf{z})$ is $\Omega(\delta)$ -strongly convex.
2. for all $i \in [n]$ it holds that the function $f_i(\mathbf{z}) = \frac{1}{2} \mathbf{z}^\top (\lambda \mathbf{I} - \mathbf{x}_i \mathbf{x}_i^\top) \mathbf{z} - \mathbf{w}^\top \mathbf{z}$ is $1 + \hat{\delta} = O(1)$ -smooth.
3. $\log(\|\mathbf{z}^*\|) = \tilde{O}(1)$, where $\mathbf{z}^*$ is the global minimizer of $F(\mathbf{z})$.

5 Putting it all together: Fast PCA via SVRG

In this section we prove Theorems 1.1, 1.2.

Following the convex optimization-based eigenvector algorithm presented in the previous section - Algorithm 3, we consider the application of the SVRG algorithm presented in [6] to the minimization of the following objective function.

$$F_{\mathbf{w}, \lambda}(\mathbf{z}) := \frac{1}{2} \mathbf{z}^\top (\lambda \mathbf{I} - \mathbf{X}) \mathbf{z} - \mathbf{w}^\top \mathbf{z} = \frac{1}{n} \sum_{i=1}^n \left(\frac{1}{2} \mathbf{z}^\top (\lambda \mathbf{I} - \mathbf{x}_i \mathbf{x}_i^\top) \mathbf{z} - \mathbf{w}^\top \mathbf{z} \right). \quad (13)$$

Note that for $\lambda > \lambda_1(\mathbf{X})$, the function $F_{\mathbf{w}, \lambda}(\mathbf{z})$ is always $(\lambda - \lambda_1(\mathbf{X}))$ -strongly-convex and that for every $i \in [n]$, the function

$$f_i(\mathbf{z}) = \frac{1}{2} \mathbf{z}^\top (\lambda \mathbf{I} - \mathbf{x}_i \mathbf{x}_i^\top) \mathbf{z} - \mathbf{w}^\top \mathbf{z},$$

is $\max\{\lambda, \|\mathbf{x}_i\|^2 - \lambda\}$ smooth. However, $f_i(\mathbf{z})$ need not be convex. Hence the SVRG theorem from [6] could not be directly applied to minimizing (13). However, we prove in the appendix that the SVRG method still converges but with a slightly worse dependence on the condition number.

Below we give an explicit implementation of the the SVRG algorithm for minimizing (13).

The following theorems are proven in the appendix.

Theorem 5.1. *Fix $\epsilon > 0, p > 0$. There exists a choice of η, m such that Algorithm 4 finds with probability at least $1 - p$ an ϵ -approximated minimizer of (13) in overall time*

$$\tilde{O} \left(nd + \frac{1}{(\lambda - \lambda_1)^2} d \right).$$

Algorithm 4 SVRG FOR PCA

```
1: Input:  $\lambda \in \mathbb{R}, \mathbf{X} = \frac{1}{n} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i^\top, \mathbf{w}, \eta, m, T$ .  
2:  $\tilde{\mathbf{z}}_0 \leftarrow \vec{0}$   
3: for  $s = 1, \dots, T$  do  
4:    $\tilde{\mathbf{z}} \leftarrow \tilde{\mathbf{z}}_{s-1}$   
5:    $\tilde{\mu} \leftarrow (\lambda \mathbf{I} - \mathbf{X})\tilde{\mathbf{z}} - \mathbf{w}_{t-1}$   
6:    $\mathbf{z}_0 \leftarrow \tilde{\mathbf{z}}$   
7:   for  $t = 1, 2, \dots, m$  do  
8:     Randomly pick  $i_t \in [n]$   
9:      $\mathbf{z}_t \leftarrow \mathbf{z}_{t-1} - \eta ((\lambda \mathbf{I} - \mathbf{x}_{i_t} \mathbf{x}_{i_t}^\top)(\mathbf{z}_{t-1} - \tilde{\mathbf{z}}) + \tilde{\mu})$   
10:  end for  
11:   $\tilde{\mathbf{z}}_s \leftarrow \frac{1}{m} \sum_{t=0}^{m-1} \mathbf{z}_t$   
12: end for  
13: return  $\mathbf{z}_T$ 
```

Based on the recent acceleration framework of [4] we also have the following result (the proof is given in the appendix).

Theorem 5.2. *Fix $\epsilon > 0, p > 0$. There exists an accelerated version of Algorithm 4 that finds with probability at least $1 - p$ an ϵ -approximated minimizer of (13) in overall time*

$$\tilde{O}\left(\frac{n^{3/4}d}{\sqrt{\lambda - \lambda_1}}\right).$$

5.1 Proving the main theorems

The proof of Theorem 1.1 follows from the bounds in Theorem 4.2, Lemma 4.3 and the above Theorem 5.1.

The proof of Theorem 1.2 follows from the bounds in Theorem Theorem 4.2, Lemma 4.3 and the above Theorem 5.2

A Convergence of the Power Method

We first restate the theorem and then prove it.

Theorem A.1. *Let $\mathbf{M}$ be a positive definite matrix and denote its eigenvalues in descending order by $\lambda_1, \lambda_2, \dots, \lambda_d$ and let $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_d$ denote the corresponding eigenvectors. Denote $\delta = \lambda_1 - \lambda_2$ and $\kappa = \frac{\lambda_1}{\delta}$. Fix an error tolerance $\epsilon > 0$ and failure probability $p > 0$. Define:*

$$T_{crude}^{\text{PM}} = \lceil \frac{1}{\epsilon} \ln \left(\frac{18d}{p^2 \epsilon} \right) \rceil, \quad T_{acc}^{\text{PM}} = \lceil \frac{\kappa}{2} \ln \left(\frac{9d}{p^2 \epsilon} \right) \rceil.$$

Then, with probability $1 - p$ it holds that

1. (crude regime) $\forall t \geq T_{crude}^{\text{PM}}(\epsilon, p): \mathbf{w}_t^\top \mathbf{M} \mathbf{w}_t \geq (1 - \epsilon) \lambda_1$.

2. (accurate regime) $\forall t \geq T_{acc}^{PM}(\kappa, \epsilon, p): (\mathbf{w}_t^\top \mathbf{u}_1)^2 \geq 1 - \epsilon$.

In both cases, the success probability depends only on the random variable $(\mathbf{w}_0^\top \mathbf{u}_1)^2$.

Proof. For the first part of the theorem, denote $k = \max\{i \in [d] \mid \lambda_i \geq (1 - \epsilon/2)\lambda_1\}$.

Using simple algebraic manipulations we have that

$$\begin{aligned}
\mathbf{w}_t^\top \mathbf{M} \mathbf{w}_t &= \frac{\mathbf{w}_{t-1}^\top \mathbf{M}^3 \mathbf{w}_{t-1}}{\|\mathbf{M} \mathbf{w}_{t-1}\|^2} = \dots = \frac{\mathbf{w}_0^\top \mathbf{M}^{2t+1} \mathbf{w}_0}{\|\mathbf{M}^t \mathbf{w}_0\|^2} = \frac{\mathbf{w}_0^\top \mathbf{M}^{2t+1} \mathbf{w}_0}{\mathbf{w}_0^\top \mathbf{M}^{2t} \mathbf{w}_0} \\
&= \frac{\sum_{i=1}^d \lambda_i^{2t+1} (\mathbf{w}_0^\top \mathbf{u}_i)^2}{\sum_{i=1}^d \lambda_i^{2t} (\mathbf{w}_0^\top \mathbf{u}_i)^2} = \frac{\sum_{i=1}^d \lambda_i \left(\frac{\lambda_i}{\lambda_1}\right)^{2t} (\mathbf{w}_0^\top \mathbf{u}_i)^2}{\sum_{i=1}^d \left(\frac{\lambda_i}{\lambda_1}\right)^{2t} (\mathbf{w}_0^\top \mathbf{u}_i)^2} \\
&\geq \frac{\sum_{i=1}^k \lambda_1 (1 - \epsilon/2) \left(\frac{\lambda_i}{\lambda_1}\right)^{2t} (\mathbf{w}_0^\top \mathbf{u}_i)^2}{\sum_{i=1}^k \left(\frac{\lambda_i}{\lambda_1}\right)^{2t} (\mathbf{w}_0^\top \mathbf{u}_i)^2 + \sum_{i=k+1}^d \left(\frac{\lambda_i}{\lambda_1}\right)^{2t} (\mathbf{w}_0^\top \mathbf{u}_i)^2} \\
&\geq \lambda_1 (1 - \epsilon/2) \cdot \frac{\sum_{i=1}^k \left(\frac{\lambda_i}{\lambda_1}\right)^{2t} (\mathbf{w}_0^\top \mathbf{u}_i)^2}{\sum_{i=1}^k \left(\frac{\lambda_i}{\lambda_1}\right)^{2t} (\mathbf{w}_0^\top \mathbf{u}_i)^2 + \sum_{i=k+1}^d (1 - \epsilon/2)^{2t} (\mathbf{w}_0^\top \mathbf{u}_i)^2} \\
&= \lambda_1 (1 - \epsilon/2) \cdot \frac{1}{1 + \frac{\sum_{i=k+1}^d (1 - \epsilon/2)^{2t} (\mathbf{w}_0^\top \mathbf{u}_i)^2}{\sum_{i=1}^k \left(\frac{\lambda_i}{\lambda_1}\right)^{2t} (\mathbf{w}_0^\top \mathbf{u}_i)^2}} \\
&\geq \lambda_1 (1 - \epsilon/2) \cdot \frac{1}{1 + (1 - \epsilon/2)^{2t} (\mathbf{w}_0^\top \mathbf{u}_1)^{-2}} \\
&\geq \lambda_1 (1 - \epsilon/2) \cdot \frac{1}{1 + (\mathbf{w}_0^\top \mathbf{u}_1)^{-2} e^{-\epsilon t}}.
\end{aligned}$$

Thus, for $t \geq \frac{1}{\epsilon} \ln \left(\frac{2}{\epsilon (\mathbf{w}_0^\top \mathbf{u}_1)^2} \right)$ we have that

$$\mathbf{w}_t^\top \mathbf{M} \mathbf{w}_t \geq \lambda_1 \frac{1 - \epsilon/2}{1 + \epsilon/2} > (1 - \epsilon) \lambda_1.$$

Part 1 of the theorem now follows, since $\mathbf{w}_0$ is a random unit vector and thus according to Lemma 5 in [1] it holds that with probability at least $1 - p$, $(\mathbf{w}_0^\top \mathbf{u}_1)^2 \geq \frac{p^2}{9d}$.

For the second part of the Theorem, we have very similarly that

$$\begin{aligned}
(\mathbf{w}_t^\top \mathbf{u})^2 &= \frac{((\mathbf{M}^t \mathbf{w}_0)^\top \mathbf{u})^2}{\|\mathbf{M}^t \mathbf{w}_0\|^2} = \frac{(\mathbf{w}_0^\top \mathbf{M}^t \mathbf{u})^2}{\mathbf{w}_0^\top \mathbf{M}^{2t} \mathbf{w}_0} = \frac{(\lambda_1^t \mathbf{w}_0^\top \mathbf{u})^2}{\sum_{i=1}^d \lambda_i^{2t} (\mathbf{w}_0^\top \mathbf{u}_i)^2} \\
&\geq \frac{1}{1 + (\mathbf{w}_0^\top \mathbf{u})^{-2} \left(\frac{\lambda_i}{\lambda_1}\right)^{2t}} \geq \frac{1}{1 + (\mathbf{w}_0^\top \mathbf{u})^{-2} \left(1 - \frac{\delta}{\lambda_1}\right)^{2t}} \\
&\geq \frac{1}{1 + (\mathbf{w}_0^\top \mathbf{u})^{-2} e^{-2\kappa^{-1}t}}.
\end{aligned}$$

Thus for $t \geq \frac{\kappa}{2} \ln \left(\frac{1}{(\mathbf{w}_0^\top \mathbf{u})^2 \epsilon} \right)$ we have that $(\mathbf{w}_t^\top \mathbf{u})^2 \geq 1 - \epsilon$. The second part of the Theorem follows since again, with probability at least $1 - p$ we have that $(\mathbf{w}_0^\top \mathbf{u}_1)^2 \geq \frac{p^2}{9d}$. $\square$

B SVRG for Convex Functions given by Sums of Non-convex Functions and its Acceleration

Suppose we want to minimize a function $F(x)$ that admits the following structure

$$F(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n f_i(\mathbf{x}), \quad (14)$$

where each f_i is β smooth, i.e.

$$\|\nabla f_i(\mathbf{x}) - \nabla f_i(\mathbf{y})\| \leq \beta \|\mathbf{x} - \mathbf{y}\| \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^d,$$

and $F(\mathbf{x})$ is σ -strongly convex, i.e.

$$F(\mathbf{y}) \leq F(\mathbf{x}) + (\mathbf{y} - \mathbf{x})^\top \nabla F(\mathbf{x}) + \frac{\sigma}{2} \|\mathbf{x} - \mathbf{y}\|^2 \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^d.$$

Algorithm 5 SVRG

```

1: Input:  $\tilde{\mathbf{x}}_0, \eta, m$ 
2: for  $s = 1, 2, \dots$  do
3:    $\tilde{\mathbf{x}} \leftarrow \tilde{\mathbf{x}}_{s-1}$ 
4:    $\tilde{\mu} \leftarrow \nabla F(\tilde{\mathbf{x}})$ 
5:    $\mathbf{x}_0 \leftarrow \tilde{\mathbf{x}}$ 
6:   for  $t = 1, 2, \dots, m$  do
7:     Randomly pick  $i_t \in [n]$ 
8:      $\mathbf{x}_t \leftarrow \mathbf{x}_{t-1} - \eta (\nabla f_{i_t}(\mathbf{x}_{t-1}) - \nabla f_{i_t}(\tilde{\mathbf{x}}) + \tilde{\mu})$ 
9:   end for
10:   $\tilde{\mathbf{x}}_s \leftarrow \frac{1}{m} \sum_{t=0}^{m-1} \mathbf{x}_t$ 
11: end for
```

Theorem B.1. *Suppose that each function $f_i(\mathbf{x})$ in the objective (14) is β -smooth and that $F(\mathbf{x})$ is σ -strongly convex. Then for $\eta = \frac{\sigma}{7\beta^2}$ and $m \geq \frac{1}{2\eta^2\beta^2}$ it holds that*

$$\mathbf{E}[\|\tilde{\mathbf{x}}_s - \mathbf{x}^*\|^2] \leq 2^{-s} \|\tilde{\mathbf{x}}_0 - \mathbf{x}^*\|^2.$$

Proof. We begin by analyzing the reduction in error on a single epoch s and then apply the result recursively. Let us fix an iteration $t \in [m]$ of the inner loop in epoch s . In the sequel we denote by $\mathbf{E}_t[\cdot]$ the expectation with respect to the random choice of i_t (i.e., the expectation is conditioned on all randomness introduced up to the t th iteration of the inner loop during epoch s). Define

$$\mathbf{v}_t = \nabla f_{i_t}(\mathbf{x}_{t-1}) - \nabla f_{i_t}(\tilde{\mathbf{x}}) + \tilde{\mu}.$$

Note that $\mathbf{E}_t[\mathbf{v}_t] = \nabla F(\mathbf{x}_{t-1})$ and thus $\mathbf{v}_t$ is an unbiased estimator for $\nabla F(\mathbf{x}_{t-1})$. We continue to upper bound the variance of $\mathbf{v}_t$ in terms of the distance of $\mathbf{x}_{t-1}$ and $\tilde{\mathbf{x}}$ from

$\mathbf{x}^*$.

$$\begin{aligned}
\mathbf{E}_t[\|\mathbf{v}_t\|^2] &\leq 2\mathbf{E}_t[\|\nabla f_{i_t}(\mathbf{x}_{t-1}) - \nabla f_{i_t}(\mathbf{x}^*)\|^2] + 2\mathbf{E}_t[\|\nabla f_{i_t}(\tilde{\mathbf{x}}) - \nabla f_{i_t}(\mathbf{x}^*) - \nabla F(\tilde{\mathbf{x}})\|^2] \\
&= 2\mathbf{E}_t[\|\nabla f_{i_t}(\mathbf{x}_{t-1}) - \nabla f_{i_t}(\mathbf{x}^*)\|^2] + 2\mathbf{E}_t[\|\nabla f_{i_t}(\tilde{\mathbf{x}}) - \nabla f_{i_t}(\mathbf{x}^*)\|^2] \\
&\quad - 4\nabla F(\tilde{\mathbf{x}})^\top (\nabla F(\tilde{\mathbf{x}}) - \nabla F(\mathbf{x}^*)) + 2\|\nabla F(\tilde{\mathbf{x}})\|^2 \\
&\leq 2\mathbf{E}_t[\|\nabla f_{i_t}(\mathbf{x}_{t-1}) - \nabla f_{i_t}(\mathbf{x}^*)\|^2] + 2\mathbf{E}_t[\|\nabla f_{i_t}(\tilde{\mathbf{x}}) - \nabla f_{i_t}(\mathbf{x}^*)\|^2] \\
&\leq 2\beta^2(\|\mathbf{x}_{t-1} - \mathbf{x}^*\|^2 + \|\tilde{\mathbf{x}} - \mathbf{x}^*\|^2),
\end{aligned}$$

where the first inequality follows from $(a+b)^2 \leq 2a^2 + 2b^2$, the first equality follows since $\mathbf{E}_t[f_{i_t}(\tilde{\mathbf{x}})] = \nabla F(\tilde{\mathbf{x}})$ (same goes for $\mathbf{x}^*$), the second inequality follows since $\nabla F(\mathbf{x}^*) = 0$, and the third inequality follows from smoothness of f_{i_t} .

We now have that,

$$\begin{aligned}
\mathbf{E}_t[\|\mathbf{x}_t - \mathbf{x}^*\|^2] &= \|\mathbf{x}_{t-1} - \mathbf{x}^*\|^2 - 2\eta(\mathbf{x}_{t-1} - \mathbf{x}^*)^\top \mathbf{E}_t[\mathbf{v}_t] + \eta^2 \mathbf{E}_t[\|\mathbf{v}_t\|^2] \\
&\leq \|\mathbf{x}_{t-1} - \mathbf{x}^*\|^2 - 2\eta(\mathbf{x}_{t-1} - \mathbf{x}^*)^\top \nabla F(\mathbf{x}_{t-1}) + 2\eta^2 \beta^2(\|\mathbf{x}_{t-1} - \mathbf{x}^*\|^2 + \|\tilde{\mathbf{x}} - \mathbf{x}^*\|^2) \\
&\leq \|\mathbf{x}_{t-1} - \mathbf{x}^*\|^2 - 2\eta\sigma\|\mathbf{x}_{t-1} - \mathbf{x}^*\|^2 + 2\eta^2 \beta^2(\|\mathbf{x}_{t-1} - \mathbf{x}^*\|^2 + \|\tilde{\mathbf{x}} - \mathbf{x}^*\|^2),
\end{aligned}$$

where the second inequality follow from convexity and strong-convexity of F .

Thus we have that,

$$\mathbf{E}[\|\mathbf{x}_t - \mathbf{x}^*\|^2] - \mathbf{E}[\|\mathbf{x}_{t-1} - \mathbf{x}^*\|^2] \leq 2\eta(\eta\beta^2 - \sigma)\mathbf{E}[\|\mathbf{x}_{t-1} - \mathbf{x}^*\|^2] + 2\eta^2 \beta^2 \mathbf{E}[\|\tilde{\mathbf{x}} - \mathbf{x}^*\|^2].$$

Summing over all iterations of the inner loop on epoch s we have

$$\mathbf{E}[\|\mathbf{x}_m - \mathbf{x}^*\|^2] - \mathbf{E}[\|\mathbf{x}_0 - \mathbf{x}^*\|^2] \leq 2\eta(\eta\beta^2 - \sigma) \sum_{t=1}^m \mathbf{E}[\|\mathbf{x}_{t-1} - \mathbf{x}^*\|^2] + 2m\eta^2 \beta^2 \mathbf{E}[\|\tilde{\mathbf{x}} - \mathbf{x}^*\|^2].$$

Rearranging and using $\mathbf{x}_0 = \tilde{\mathbf{x}}$ we have that,

$$2\eta(\sigma - \eta\beta^2) \sum_{t=1}^m \mathbf{E}[\|\mathbf{x}_t - \mathbf{x}^*\|^2] \leq (1 + 2m\eta^2 \beta^2) \mathbf{E}[\|\tilde{\mathbf{x}} - \mathbf{x}^*\|^2].$$

Using $\tilde{\mathbf{x}} = \tilde{\mathbf{x}}_{s-1}$ and $\tilde{\mathbf{x}}_s = \frac{1}{m} \sum_{t=0}^{m-1} \mathbf{x}_t$ we have that,

$$\mathbf{E}[\|\tilde{\mathbf{x}}_s - \mathbf{x}^*\|^2] \leq \frac{1 + 2m\eta^2 \beta^2}{2\eta m(\sigma - \eta\beta^2)} \mathbf{E}[\|\tilde{\mathbf{x}}_{s-1} - \mathbf{x}^*\|^2].$$

Plugging the values of η, m gives the theorem. $\square$

B.1 Acceleration of Algorithm 5

We now discuss, how using the recent generic acceleration framework of [4], we can further accelerate Algorithm 5. Note that Algorithm 5 requires to compute overall $\tilde{O}(n+m) = \tilde{O}\left(\frac{\beta^2}{\sigma^2} + n\right)$ gradients of functions from the set $\{f_i(\mathbf{x})\}_{i=1}^n$. Using the framework of [4], this quantity could be dramatically reduced.

On a very high-level, the framework of [4] applies a convex optimization algorithm in an almost black-box fashion in order to simulate an algorithm known as the *Accelerated*

Proximal-point algorithm. That is, it uses the convex optimization algorithm to find an approximated global minimizer of the modified function:

$$\tilde{F}(\mathbf{x}) = \frac{1}{n} \sum_{i=1}^n f_i(\mathbf{x}) + \frac{\kappa}{2} \|\mathbf{x} - \mathbf{y}\|^2, \quad (15)$$

where $\kappa, \mathbf{y}$ are parameters.

Note that the SVRG algorithm (5) could be directly applied to minimize (15) by considering the set of functions $\tilde{f}_i(\mathbf{x}) = f_i(\mathbf{x}) + \frac{\kappa}{2} \|\mathbf{x} - \mathbf{y}\|^2$ for all $i \in [n]$. It clearly holds that $\tilde{F}(\mathbf{x}) = \sum_{i=1}^n \tilde{f}_i(\mathbf{x})$. Not also that now $\tilde{F}(\mathbf{x})$ is $\sigma + \kappa$ strongly convex and for each $i \in [n]$ it holds that $\tilde{f}_i(\mathbf{x})$ is $\beta + \kappa$ smooth.

The following Theorem (rephrased for our needs) is proven in [4].

Theorem B.2. *Fix the parameter κ . There exists an acceleration scheme for Algorithm 5 that finds an ϵ approximated solution to the objective in (14) after approximately minimizing $\tilde{O}\left(\sqrt{\frac{\sigma+\kappa}{\sigma}}\right)$ instances of (15).*

Plugging Theorems B.1, B.2 and the proprieties of $\tilde{F}(\mathbf{x})$ we have that Algorithm 5 could be applied to finding an ϵ minimizer of (14) that requires computing at most

$$\tilde{O}\left(\sqrt{\frac{\sigma+\kappa}{\sigma}}\left(n + \left(\frac{\beta+\kappa}{\sigma+\kappa}\right)^2\right)\right)$$

gradient vectors of functions from the set $\{f_i(\mathbf{x})\}_{i=1}^n$.

By optimizing the above bound with respect to κ we arrive at the following theorem.

Theorem B.3. *Assume that the gradient vector of each function $f_i(\mathbf{x})$ could be computed in $O(d)$ time. Assume further that $\beta = \Omega(\sqrt{n}\sigma)$. Algorithm 5 combined with the acceleration framework of [4], find an ϵ -approximated minimizer of (14) in total time $\tilde{O}\left(\sqrt{\frac{\beta}{\sigma}}n^{3/4}d\right)$.*

References

- [1] Sanjeev Arora, Satish Rao, and Umesh V. Vazirani. Expander flows, geometric embeddings and graph partitioning. *Journal of the ACM*, 56(2), 2009.
- [2] Akshay Balsubramani, Sanjoy Dasgupta, and Yoav Freund. The fast convergence of incremental pca. In *Advances in Neural Information Processing Systems 26*, pages 3174–3182. 2013.
- [3] Roy Frostig, Rong Ge, Sham Kakade, and Aaron Sidford. Un-regularizing: approximate proximal point and faster stochastic algorithms for empirical risk minimization. In *Proceedings of the 32nd International Conference on Machine Learning, ICML 2015, Lille, France, 6-11 July 2015*, pages 2540–2548, 2015.
- [4] Julien Mairal Hongzhou Lin and Zaid Harchaoui. A universal catalyst for first-order optimization. *CoRR*, abs/1506.02186, 2015.

- [5] H. Hotelling. Analysis of a complex of statistical variables into principal components. *J. Educ. Psych.*, 24, 1933.
- [6] Rie Johnson and Tong Zhang. Accelerating stochastic gradient descent using predictive variance reduction. In *Advances in Neural Information Processing Systems 26: 27th Annual Conference on Neural Information Processing Systems.*, pages 315–323, 2013.
- [7] Jakub Konečný and Peter Richtárik. Semi-stochastic gradient descent methods. *CoRR*, abs/1312.1666, 2013.
- [8] Mehrdad Mahdavi, Lijun Zhang, and Rong Jin. Mixed optimization for smooth functions. In *Advances in Neural Information Processing Systems 26*, pages 674–682. 2013.
- [9] Erkki Oja. Simplified neuron model as a principal component analyzer. *Journal of mathematical biology*, 15(3):267–273, 1982.
- [10] K. Pearson. On lines and planes of closest fit to systems of points in space. *Philosophical Magazine*, 2(6):559–572, 1901.
- [11] Nicolas Le Roux, Mark W. Schmidt, and Francis R. Bach. A stochastic gradient method with an exponential convergence rate for finite training sets. In *Advances in Neural Information Processing Systems 25: 26th Annual Conference on Neural Information Processing Systems.*, pages 2672–2680, 2012.
- [12] Shai Shalev-Shwartz and Tong Zhang. Stochastic dual coordinate ascent methods for regularized loss minimization. *CoRR*, abs/1209.1873, 2012.
- [13] Ohad Shamir. A stochastic PCA algorithm with an exponential convergence rate. *CoRR*, abs/1409.2848, 2014.
- [14] Ohad Shamir. Fast stochastic algorithms for svd and pca: Convergence properties and convexity. *CoRR*, abs/1507.08788, 2015.