

Semantics for a Logic of Presuppositions

by X.Y. Newberry

Abstract

In 1952 P. F. Strawson proposed a logic of presuppositions. It is an interpretation of Aristotelian logic, i.e. of the logic of the traditional syllogism. In 1981 Richard Diaz published a monograph in which he presented *truth-relevant logic*. This paper shows that truth-relevant logic is but a propositional version of the logic of presuppositions. A semantics of the logic of presuppositions is developed using truth-relevant logic. The semantics is then further extended to polyadic propositions and some consequences discussed.

1. Introduction

The purpose of this paper is to establish a foundation for a logic where vacuous sentences, such as $\forall x(x \neq x \rightarrow x = x+1)$, are not derivable. In such a logic Gödel's self-referential sentence is not derivable [it is vacuous], but $\sim(Ex)Prf(x, g)$ might be. Here g is the Gödel number of Gödel's sentence. And $Prf(x, z)$ means that x is the Gödel number of a sequence that is a proof of the sentence with Gödel number z . (Newberry, 2015)

In 1981 M. Richard Diaz (1981) published a monograph titled *Topics in The Logic of Relevance*. He proposed a new type of relevance logic called *truth-relevant logic*. In this logic the paradox of material implication $(P \& \sim P) \rightarrow Q$ is *not* truth-relevant, but neither are its classical equivalents such as $(P \vee \sim P) \vee Q$.

In 1952 P. F. Strawson introduced the Logic of Presuppositions (Strawson, 1952.) It conforms to Aristotle's logic of the traditional syllogism. In this logic, sentences with empty subjects are not considered true. For example the sentence "All John's children are asleep" would be considered neither true nor false if John had no children. This paper attempts to develop the semantics of the Logic of Presuppositions based on truth-relevant logic of Richard Diaz. The philosophical justification of this project can be found in Newberry (2014.)

2. Propositional Calculus

There are compound formulae of propositional logic such that their value can be determined by a subset of their variables. Consider for example $S = P \vee \sim P \vee Q$. When $|P|$ is 1 then $|S| = 1$ regardless of the value of Q . When $|P| = 0$ then $|S| = 1$ also regardless of the value Q . The set of variables occurring in S is $\{P, Q\}$. We say that the subset $\{P\}$ is truth-determining for S ; for *all the valuations* of $\{P\}$, i.e. $|P| = 1$ or $|P| = 0$, we can determine the value of S regardless of the other variables. (Here ' $|P|$ ' means the Boolean value of P .) We use the *Boolean* values 1/0 because we will consider as true only a proper subset of the compound formulas that evaluate as Boolean 1. In case of atomic sentences $T = 1, F = 0$, in case of compound sentences, only t-relevant sentences are considered *true* when they evaluate as 1.

In truth relevant logic the truth tables are constructed from the following elementary truth tables (Diaz, p.66.)

v		1	x	0	&		1	x	0	→		1	x	0
1		1	1	1	1		1	x	0	1		1	x	0
x		1	x	x	x		x	x	0	x		1	x	x
0		1	x	0	0		0	0	0	0		1	1	1

Table 2.1

These tables happen to be identical with Kleene's "strong" tables. But Diaz's objective is to identify irrelevant variables, i.e. variables that do not make any contribution to the Boolean value of the compound sentence.

Definition 2.1: A set of propositional variables is *truth determining* for a proposition S iff the truth value of S may be determined as 1 or 0 (using the matrices above) on all assignments of 1, 0 to the set.

That is, it may be the case that we assign x to some variable, and yet we obtain a definite 1 or 0 once the entire compound formula is evaluated. It is an indication that the variable does not contribute anything to the 'output.' In this event, said variable is deemed *redundant*.

Another example: $P \rightarrow (Q \rightarrow P)$ (2.1)

If $|P| = 1$ then $|(Q \rightarrow P)| = 1$ regardless of the value of Q , but then the entire formula (2.1) evaluates as 1 (regardless of the value of Q .) Suppose $|P| = 0$. Then (2.1) is true regardless of the value of Q . In both cases (i.e. $|P| = 1$ or $|P| = 0$) the value of (2.1) can be determined without any knowledge of the value of Q . Thus $\{P\}$ is truth-determining for (2.1). This is not the case for Q . For assume $|Q| = 1$. Then the value of $(Q \rightarrow P)$ cannot be determined without knowing the value of P . And without knowing the value of $(Q \rightarrow P)$, the value of (2.1) cannot be determined. (Diaz, 1981, pp. 66, 67)

Definition 2.2: Let P_1, P_2, \dots, P_n be all the variables occurring in S , P_i is *truth-redundant* in S iff there is a truth determining set for S that does not contain P_i .

Definition 2.3: S is *truth-relevant* if it contains no truth-redundant variables.

Of special interest is the set of tautologies that are also *t-relevant*.

These definitions are due to Diaz (1981, pp. 65-67). We can equivalently rephrase 2.2 as follows:

Definition 2.4: Let P_1, P_2, \dots, P_n be all the variables occurring in S , P_i is *truth-redundant* in S iff the value of S can be determined regardless of the value of P_i .

3. Monadic Sentences

3.1 Definitions

Definition 3.1.1: Let S be a first order monadic sentence. Let P_1, P_2, \dots, P_n be all the atomic predicates occurring in S , P_i is *truth-redundant* in S iff the truth value of S can be determined regardless of the interpretation of P_i .

Definition 3.1.2: S is *truth-relevant* if it contains no truth-redundant atomic predicates.

We will use the term *satisfied* in the same sense as in classical logic. But we will say that a sentence is *true* iff it is *satisfied and t-relevant*.

The extension of the concept of truth relevance to monadic sentences is natural and straightforward. It does not present any major challenges. We will illustrate by examples.

3.2 “All John’s children are asleep”

Let us consider the sentence “All John’s children are asleep” when John has no children (Strawson’s example.) That is, consider the sentence $\forall x(Jx \rightarrow Sx)$ and an interpretation D such that:

$|D|$ = the set of three children: Alex, Betty, Cindy; that is

$|D| = \{a, b, c\}$

Jx : x is John's child

$\sim Ja, \sim Jb, \sim Jc$

Sx : x is asleep

$Sa, Sb, \sim Sc$

I.e. Alex and Betty are asleep and John has no children.

The Boolean table of $Jx \rightarrow Sx$ looks as follows:

Row	Jx	Sx	$Jx \rightarrow Sx$
1	0	0	1
2	0	1	1
3	1	0	0
4	1	1	1

Table 2.1

We will now substitute constants corresponding to the elements of $|D|$ one by one thus successively obtaining $Ja \rightarrow Sa$, $Jb \rightarrow Sb$, $Jc \rightarrow Sc$, the corresponding Boolean values being:

	Jx	Sx	$Jx \rightarrow Sx$	Row of Table 1
$Ja \rightarrow Sa$	0	1	1	2
$Jb \rightarrow Sb$	0	1	1	2
$Jc \rightarrow Sc$	0	0	1	1

Table 2.2

We observe that Jx is always 0, and subsequently the truth value of the implication is always 1 regardless of the value of Sx . That is, given the interpretation D , the value of $\forall x(Jx \rightarrow Sx)$ can be determined without knowing the interpretation of Sx . And since Sx is not a t-relevant atomic predicate, the sentence is not t-relevant and we consider it as $\sim T$ & $\sim F$. It may seem contradictory that we claim that a value determination can be effected without knowing a predicate's interpretation while at the same time considering a given interpretation. But, in this case, it is the interpretation of J that causes the interpretation of S to be irrelevant.

3.3 Valid Sentences

Consider $\forall x(Px \rightarrow (Qx \rightarrow Px))$. It is a valid sentence, and these evaluate as 1 under any interpretation. Yet even in this case there is a sense in which the interpretation of Qx is

irrelevant. Pick an arbitrary a : $Pa \rightarrow (Qa \rightarrow Pa)$. Under some interpretations $|Pa| = 1$, and under some $|Pa| = 0$. It turns out that we could not care less what the interpretation of Qa would or would not be. If $|Pa| = 1$ then $|(Qa \rightarrow Pa)| = 1$ regardless of the value of Qa , but then the entire formula evaluates as 1 (regardless of the value of Qa .) Suppose $|Pa| = 0$. Then the value of the entire sentence is 1 regardless of the value of Qa . In either case (i.e. $|Pa| = 1$ or $|Pa| = 0$) the value of the sentence can be determined without any knowledge of the value of Qa . Thus $\{Px\}$ is truth-determining for $\forall x(Px \rightarrow (Qx \rightarrow Px))$.

Let us look at prototypical cases more systematically.

3.4 Basic Cases

Case 1: $\forall x(Px \vee Qx)$

Consider $\forall x(Px \vee Qx)$, where P, Q are atomic predicates. The formula can also be written as $\sim \exists x(\sim Px \ \& \ \sim Qx)$. Let us examine the scenarios below. The diagrams depict the extents of the (atomic) predicates Px, Qx in some finite domain.

P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P
Q	Q	Q	Q

Figure 3.4.1

P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P
.

Figure 3.4.2

P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P	P
Q	Q	Q	Q	Q	Q	Q	Q	Q	Q	Q	Q	Q	Q	Q	Q	Q	Q	Q	Q

Figure 3.4.3

In all these cases the value of $\forall x(Px \vee Qx)$ can be determined without knowing the interpretation of Qx . The truth determining subset of $\forall x(Px \vee Qx)$ is $\{Px\}$. In the last scenario there are in fact two truth determining subsets $\{Px\}$ and $\{Qx\}$. It is apparent that the formula will be t-relevant only if $\exists x \sim Px$ and $\exists x \sim Qx$. P.F. Strawson called these conditions *presuppositions*, albeit in a slightly different context.

The negation of the above formula is $\exists x(\sim Px \ \& \ \sim Qx)$. The conditions of relevance ought to be the same, i.e. $\exists x \sim Px$, $\exists x \sim Qx$, and we observe that these are satisfied automatically.

Consider $\forall x(x \leq 5 \vee x \geq 5)$. The presuppositions $\exists x(x > 5)$, $\exists x(x < 5)$ are met; the formula is t-relevant. It is also satisfied and hence true. This cannot be said for example about $\forall x(x = x \vee x \geq 5)$. The required presupposition $\exists x(x \neq x)$ is not met. The sentence is $\sim T$ & $\sim F$.

Case 2: $\exists x(Px \vee Qx)$

The truth value of this sentence can be determined from $\exists x(Px)$ [or $\exists x(Qx)$.]

Therefore $\exists x(Px \vee Qx)$ is never t-relevant.

The negation is $\forall x(\sim Px \ \& \ \sim Qx)$, i.e. a clean board. This sentence is always t-relevant [when satisfied.]

It follows from the aforesaid that $\forall x(Px \vee Qx) \not\models \exists x(Px \vee Qx)$. More about this below.

It turns out that the only interesting case is the first one. Let us put $Fx =_{\text{def}} \sim Px$ and $Gx =_{\text{def}} \sim Qx$, and consider the ‘existential’ version $\sim \exists x(Fx \ \& \ Gx)$. The presuppositions become $\exists xFx$, $\exists xGx$. Then a typical interpretation when the sentence is satisfied and t-relevant (i.e. true) is as follows:

F F F F F F G G G G G G G

Figure 3.4.4

Under the interpretation of F and G below (Fig 3.4.5) it would be satisfied but not t -relevant (i.e. $\sim T$ & $\sim F$.)

[illegible]

Figure 3.4.5

And under the next interpretation (Fig. 3.4.6) it would be false.

F F F F F F F F F F F F F
. G G G G G G G G G G G

Figure 3.4.6

Our findings are summarized in the following table.

	Universal	Existential	Presuppositions	Negation	Presuppositions
1	$\forall x(Px \vee Qx)$	$\sim \exists x(\sim Px \ \& \ \sim Qx)$	$\exists x \sim Px \ \& \ \exists x \sim Qx$	$\exists x(\sim Px \ \& \ \sim Qx)$	automatic
2	$\forall x(Px \ \& \ Qx)$	$\sim \exists x(\sim Px \vee \sim Qx)$	automatic	$\exists x(\sim Px \vee \sim Qx)$	$\exists x Px \ \& \ \exists x Qx$
3		$\sim \exists x(Fx \ \& \ Gx)$	$\exists x Fx \ \& \ \exists x Gx$	$\exists x(Fx \ \& \ Gx)$	automatic
4		$\sim \exists x(Fx \vee Gx)$	automatic	$\exists x(Fx \vee Gx)$	$\exists x \sim Fx \ \& \ \exists x \sim Gx$

Table 3.4.1

Consider for example the sentence $\forall x(Fx \vee Gx)$, $Fx =_{def} x > 6$, $Gx =_{def} x < 8$, i.e.

$$\forall x(x > 6 \vee x < 8) \quad (3.4.1)$$

We claim that this sentence is true (that implies t-relevant.) The presuppositions are $\exists x(x \leq 6) \ \& \ \exists x(x \geq 8)$, and they are both satisfied. Nevertheless no substitution instance is true. For example none of the sentences below are t-relevant.

$$5 > 6 \vee 5 < 8 \quad (3.4.2)$$

$$7 > 6 \vee 7 < 8 \quad (3.4.3)$$

$$9 > 6 \vee 9 < 8 \quad (3.4.4)$$

The redundant predicates are grayed out. All the above sentences are valid but not t-relevant. The sentence (3.4.3) has two truth-determining subsets $\{7 > 6\}$, $\{7 < 8\}$.

So $\forall x(x < 6 \vee x > 8) \not\models (5 < 6 \vee 5 > 8)$. The only way to “instantiate” (3.4.1) is to pick one disjunct or the other. For example when we want to substitute 5 for x we have to pick either $5 > 6$ or $5 < 8$. (Choose wisely.)

Does this make sense? Let us briefly recapitulate how we got here. Presumably the classical definitions of *satisfaction* and *validity* do make sense. I think that my definition of an atomic predicate *relevance* also makes sense. And the result is what it is. Still, does it

make sense? When some property applies to all then it must apply to one, right? Nobody is denying that this is the case with atomic predicates. Indeed $\forall x Fx \models \exists x Fx$. If all apples are round then there exists a round apple. But here we are talking about compound ‘properties’, and while $\forall x(Fx \vee Gx)$ talks about a *collection*, $Fa \vee Ga$ talks about an *individual*. You may say “All apples are green or shiny”, but you would not say about any particular one “This apple is green or shiny.” It makes sense to say about a collection that its members are this or that. But it does not make sense to say about one particular thing that it is this *or* that. It is one or the other or both! Just pick one and see. (Fig. 4.2.3)



Figure 4.2.3

In t-relevant logic the *or* statement can be used only in case of uncertainty. For example you can say “It has rained or somebody has watered the lawn.” You are pretty sure it is one or the other, but you do not know which one. The *or* connective cannot be used once a determination has been made or in case of necessary truths (tautologies, valid sentences, analytic sentences etc.) In particular it cannot be used about individual objects in mathematics. This is in accordance with the definitions of the logical connectives in Newberry (2019.)

$$|P \vee Q| = T \text{ iff } (P + Q) \cdot \diamondsim P \cdot \diamondsim Q$$

$$|P \vee Q| = F \text{ iff } \sim(P + Q) \cdot \diamondsim P \cdot \diamondsim Q$$

$$|P \& Q| = T \text{ iff } P \cdot Q \cdot \diamond P \cdot \diamond Q$$

$$|P \& Q| = F \text{ iff } \sim(P \cdot Q) \cdot \diamond P \cdot \diamond Q$$

where ‘+’, ‘•’ are a boolean operators, ‘ \diamond ’ means *possible*.

4. Polyadic Sentences

4.1 Definitions

In classical logic we have

Definition 4.1.1:

$$(\forall x)(\forall y)U_{xy} \quad (4.1.1)$$

is *true* iff for all a_i **in the range of x**

$$(\forall y)U_{a_i y} \quad (4.1.2)$$

is true. What about our t-relevant logic? If we apply our classical definition “in the range of x ” we find ourselves quantifying over sentences that are $\sim(T \vee F)$. Nothing good can possibly come out of that. In fact in our view those sentences are devoid of any meaning. The most sensible thing is to leave them out. We will therefore modify the above definition as follows:

Definition 4.1.2:

$$(\forall x)(\forall y)U_{xy} \quad (4.1.3)$$

is *true* iff for all a_i *such that* $(\forall y)U_{a_i y}$ *is t-relevant*

$$(\forall y)U_{a_i y} \quad (4.1.4)$$

is true.

The situation is completely symmetrical with respect to y . The definition implies that there has to be at least one pair $\langle a, b \rangle$ such that $\forall x Uxb$ is true (and hence relevant) and $\forall y Uay$ is true (and hence relevant.) If no such pair exists the universally quantified dyadic sentence is $\sim(T \vee F)$.

Now we need to generalize the definition of t-relevance to polyadic sentences. Hence the following definition:

Definition 4.1.1 A first order sentence $Q_1x_1Q_2x_2\dots Q_nx_nAx_1x_2\dots x_n$ with n variables in prenex normal form is *t-relevant* iff there is an n -tuple $\langle c_1, \dots, c_n \rangle$ such that $Q_1x_1Ax_1c_2\dots c_n$ is t-relevant [per definition 3.1.2], and $Q_2x_2Ac_1x_2\dots c_n$ is t-relevant, and
...
 $Q_nx_nAc_1c_2\dots x_n$ is t-relevant.

That is to say, there has to be at least one t-relevant sentence in each ‘dimension’, and they all have to intersect at one point. In the above notation Q_i means \forall or \exists .

Now consider $\exists x \exists y Uxy$. It says that there is an a such that $\exists y Uay$. It means that the given $\exists y Uay$ has to be t-relevant. It implies that there is an a such that $\exists y Uay$ is t-relevant. But the quantifiers commute. So the same reasoning applies to $\exists y \exists x Uxy$. That is, there has to be a b such that $\exists x Uxb$ is t-relevant. Putting the two together we have: $\exists x \exists y Uxy$ is t-relevant iff there exists a pair $\langle a, b \rangle$ such that

$\exists y Uay$ is t-relevant, and

$\exists x Uxb$ is t-relevant.

By extrapolation, a first order sentence $\exists x_1 \exists x_2 \dots \exists x_n A x_1 x_2 \dots x_n$ with n variables in prenex normal form is *t-relevant* iff there is an n -tuple $\langle c_1, \dots, c_n \rangle$ such that

$\exists x_1 A x_1 c_2 \dots c_n$ is t-relevant [per definition 3.1.2], and

$\exists x_2 A c_1 x_2 \dots c_n$ is t-relevant, and

...

$\exists x_n A c_1 c_2 \dots x_n$ is t-relevant.

We know that $\exists x \forall y Uxy$ implies $\forall x \exists y Uxy$. Therefore given a prenex sentence, we can push all the existential quantifiers inside and obtain

$\forall x_1 \forall x_2 \dots \forall x_m \exists x_{m+1} \exists x_{m+2} \dots \exists x_{m+n} A x_1 x_2 \dots x_m x_{m+1} x_{m+2} \dots x_{m+n}$

This will be t-relevant iff there is an n -tuple $\langle c_1, \dots, c_n \rangle$ such that

$\forall x_1 \exists x_{m+1} \exists x_{m+2} \dots \exists x_{m+n} A x_1 c_2 \dots c_m x_{m+1} x_{m+2} \dots x_{m+n}$ is t-relevant, and

$\forall x_2 \exists x_{m+1} \exists x_{m+2} \dots \exists x_{m+n} A c_1 x_2 \dots c_m x_{m+1} x_{m+2} \dots x_{m+n}$ is t-relevant, and

...

$\forall x_n \exists x_{m+1} \exists x_{m+2} \dots \exists x_{m+n} A c_1 c_2 \dots x_m x_{m+1} x_{m+2} \dots x_{m+n}$ is t-relevant.

By putting it all together we obtain definition 4.1.

The reason we need an expanded definition for polyadic sentences is the following.

Consider $\forall x \forall y Uxy$. It means that for all a_i it is the case that $\forall y Ua_i y$ ²⁾. But it could be that for $b \neq c$ the sentences $\forall y Uby$ and $\forall y Ucy$ are both non-relevant while having *different* redundant primitive predicates. We will illustrate by examples.

² For the sake of brevity we assume that each object has a name, and each constant denotes.

4.2 Examples

Example 1: $\sim \exists x \exists y (Fxy \ \& \ Gxy)$

b	y																		x																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																													
		</

Figure 4.2.1

The instance $\sim\exists y(Fay \ \& \ Gay)$ is t-relevant as the presuppositions $\exists yFay \ \& \ \exists yGy$ are satisfied (Table 3.4.1, line 3.) The instance $\sim\exists x(Fxb \ \& \ Gxb)$ is *not*. Therefore there is *no* pair $\langle a, b \rangle$ such that both $\sim\exists y(Fay \ \& \ Gay)$ and $\sim\exists x(Fxb \ \& \ Gxb)$ are t-relevant.

Example 2: $\sim\exists x\exists y(Fxy \ \& \ Gxy)$

Here is a different example.: $\sim\exists x\exists y(Fxy \ \& \ Gxy) = \forall x\forall y(\sim Fxy \vee \sim Gxy)$,

$Fxy =_{\text{def}} x + y < 8$, $Gxy =_{\text{def}} x + y > 8$,

i.e. $\sim\exists x\exists y(x + y < 8 \ \& \ x + y > 8) = \forall x\forall y(x + y \geq 8 \vee x + y \leq 8)$.




Figure 4.2.2

In this scenario the formula is t-relevant. Not all substitution instances are t-relevant. By substituting any number from the interval $[1,7]$ for x we obtain a t-relevant formula. For example substitute 3:

$$\forall y(\mathbf{3} + y \geq 8 \vee \mathbf{3} + y \leq 8)$$

The sentence is t-relevant as the presuppositions $\exists y\sim(\mathbf{3} + y \geq 8)$ and $\exists y\sim(\mathbf{3} + y \leq 8)$ are both satisfied. Substitute 13:

$$\forall y(\mathbf{13} + y \geq 8 \vee \mathbf{13} + y \leq 8)$$

This sentence is not t-relevant as the presuppositions $\exists y \sim (13 + y \geq 8)$ i.e. $\exists y (13 + y < 8)$ is *not* satisfied. Note that no substitution instance of both x and y is t-relevant. E.g.

$$6 + 7 \geq 8 \vee 6 + 7 < 8$$

The presuppositions are $6 + 7 < 8$ & $6 + 7 \geq 8$. There is no chance the former could be satisfied. There are no two natural numbers such that their sum is simultaneously less and greater than 8.

Most of the sentences $\sim \exists y (Fay \& Gay)$ in the diagram are not t-relevant, hence $\sim T \& \sim F$. We only want to quantify over the t-relevant sentences, i.e. when $b = [1, 7]$. But according to figure 4.2.2 we are doing that, and we can simply disregard that we are also quantifying over the non-relevant ones. Nevertheless the question arises if it could be that all the t-relevant instances were satisfied while $\sim \exists x \exists y (Fxy \& Gxy)$ was not. The answer is no because any counterexamples to $\sim \exists x \exists y (Fxy \& Gxy)$ must necessarily be t-relevant.

Theorem 4.2.1

Let $\sim \exists x \exists y (Fxy \& Gxy)$ be a sentence with atomic predicates F, G . If all the t-relevant instances $\sim \exists y (Faiy \& Gaiy)$ and $\sim \exists x (Fxb_i \& Gxb_i)$ are satisfied then $\sim \exists x \exists y (Fxy \& Gxy)$ is satisfied. (*Satisfied* means *evaluates as Boolean 1*.)

Proof: Assume the opposite. Then $\exists x \exists y (Fxy \& Gxy)$. So there is some $\langle c, d \rangle$, say $\langle 11, 12 \rangle$ (Fig. 4.2.2), such that

$$\exists y (Fcy \& Gcy) \text{ and } \exists x (Fxd \& Gxd) \quad (4.2.1)$$

Both of these sentences are t-relevant, and so are their negations

$$\sim \exists y (Fcy \& Gcy) \text{ and } \sim \exists x (Fxd \& Gxd) \quad (4.2.2)$$

These are t-relevant instances of $\sim \exists x \exists y (Fxy \& Gxy)$. If such $\langle c, d \rangle$ do exist then (4.2.2) are t-relevant albeit false. But this contradicts the assumption that all the t-relevant instances are satisfied. QED.

Example 3: $\sim\exists x\exists y(Lxy \ \& \ My)$

Let $Lxy =_{def} (x + y \leq 8)$, $My =_{def} (y = 12)$. We obtain $\sim\exists x\exists y(x + y \leq 8 \ \& \ y = 12)$.

The above formula is not t-relevant according to Figure 4.4.1 below.




Figure 4.4.1

The sentence $\sim\exists x\exists y(x + y \leq 8 \ \& \ y = 12)$ bears a strong resemblance to Gödel's sentence. Let

$$\sim(\exists x)(\exists z)(Prf(x, z) \ \& \ Diag(y, z)) \quad (U)$$

be a formula with one free variable y , where $Diag(y, z)$ means that y is the Gödel number of a formula with one free variable iff z is the Gödel number number of the formula obtained from y by substituting (the numeral of) the Gödel number of y for the free variable in y . And $Prf(x, z)$ means that x is the Gödel number of a sequence that is a proof of the sentence with Gödel number z .

Let the constant k be the Gödel number of (U). We substitute k for the free variable y in (U), and obtain

$$\sim(\exists x)(\exists z)(Prf(x,z) \ \& \ Diag(k,z)) \quad (G)$$

This is Gödel's self-referential sentence. Then graphically the situation will look much like Figure 4.4.1 in the sense that the pairs $\langle x,y \rangle$ of natural numbers picked by $Prf()$ and $Diag()$ respectively will not intersect. That means that G will *not* be t-relevant even if $\sim\exists x(Prf(x, \ulcorner G \urcorner))$ is true and provable!

The sentence G is $\sim T$ & $\sim F$ in the logic of presuppositions since $\sim(\exists x)Prf(x, \ulcorner G \urcorner)$ contradicts the *presupposition* $(\exists x)Prf(x, \ulcorner G \urcorner)$. This renders $Diag(k,z)$ redundant. However Prf and $Diag$ are compound predicates while in t-relevant logic the relevance attribute applies only to atomic predicates.

Let us take a closer look. Note that $\ulcorner U(y) \urcorner$ is a specific natural number. When we expand $Diag(\ulcorner U(y) \urcorner, z)$ then $\ulcorner U(y) \urcorner$ will appear in some predicate $v = \ulcorner U(y) \urcorner$ for some variable v . This atomic predicate will be redundant.

Bibliography

Diaz, M.R. (1981) *Topics in the Logic of Relevance*, Munich, Germany: Philosophia Verlag.

Newberry, X.Y. (2014) Meaning, Presuppositions, Truth-relevance, Gödel's Sentence and the Liar Paradox, <https://philpapers.org/archive/NEWMPT.pdf>

Newberry, X.Y. (2015) The Diagonal Lemma Fails in Aristotelian Logic, *The Reasoner* Volume 10, Number 1, <http://arxiv.org/abs/1510.0439>

Newberry, X.Y. (2019) Truth-relevant Logic – Propositional Calculus, <https://arxiv.org/abs/2204.10887>

Strawson, P.F. (1952) *Introduction to Logical Theory*, London: Methuen.

Copyright © X.Y. Newberry 2019 - 2023

For non-commercial purposes this document may be freely copied and distributed.

