

Anomalous weak amplification with arbitrarily orthogonal postselections: A further consideration on “A classical postselected weak amplification scheme and thermal light cross-Kerr effect” and “Can anomalous amplification be attained without postselection?”

Tao Wang* and Rui Zhang†

¹ College of Physics, Jilin University, Changchun 130012, People's Republic of China and
² College of Physics, Tonghua Normal University, Tonghua 134000, People's Republic of China

(Dated: July 21, 2022)

Nearly thirty years ago the possibility of anomalous weak amplification (AWA) was revealed by Aharonov, Albert and Vaidman [1]. Recently two papers presents two AWA schemes which are beyond the traditional proposal given by them [9, 10]. At the first glance the two papers seems very different. Ref. [9] discusses the thermal light cross-Kerr effect and finds only postselection can give the amplification effect without interference and Ref. [10] shows that only weak interaction can give the amplification effect without postselection. Here the relationships between the two papers are pointed out and a generalized framework for AWA via postselecting a pair of orthogonal final state are shown.

PACS numbers: 42.50.Wk, 42.65.Hw, 03.65.Ta

In 1988 anomalous weak amplification was proposed by Aharonov, Albert, and Vaidman [1]. Weakly measuring a quantum system given the initial state and a final state can attain a prominent amplification result. The inconceivable effect was exploited to observe many tiny physical effects [2, 3]. An endeavor to amplify the faint gravitational wave effect via AWA and optomechanics was made in [4, 5]. Reviews on this strange idea can be found in [6, 7].

This AWA scheme consists of three steps. *i)* the quantum system to be measured and the meter system to measure are prepared respectively. For simplicity, we choose $|\psi_i\rangle = \frac{1}{\sqrt{2}}(e^{i\theta}|0\rangle + |1\rangle)$ for the quantum system, and $\varphi(q)$ for the meter system. θ is a small phase shift. *ii)* the quantum system weakly interacts with the meter system via $U = e^{-ig\hat{A}\hat{p}}$, and after interaction, the two systems are weakly entangled together, and the state for the whole system is $\frac{1}{\sqrt{2}}(e^{i\theta}|0\rangle e^{ig\hat{p}}\varphi(q) + |1\rangle e^{-ig\hat{p}}\varphi(q))$. Here \hat{q} and \hat{p} are conjugate variables. And $\hat{A}|0\rangle = -|0\rangle$, $\hat{A}|1\rangle = +|1\rangle$. *iii)* a final state is particularly selected, which is nearly orthogonal to the initial state. Here we choose $|\psi_{f,1}\rangle = \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle)$, so $\langle\psi_f|\psi_i\rangle = \frac{1}{2}(e^{i\theta} - 1) \approx \frac{\theta}{2}i$ [8]. The final state of the meter system is $\frac{1}{2P}(e^{i\theta}e^{ig\hat{p}}\varphi(q) - e^{-ig\hat{p}}\varphi(q))$, where P is the detection probability and is very small. Here destructive interference occurs and we have an amplification effect.

Generally speaking, postselection, weak interaction, weak entanglement and destructive interference are all regard as the indispensable components for AWA. However Ref. [9, 10] presents another edition for AWA [11].

We exploit $\varphi(p)$ instead of $\varphi(q)$. The key thing here is to measure the physical quantity appearing in the interaction instead of its conjugate one. The measurement quantity communicates with the Hamiltonian of the system. *If there is no postselection, it is a conserved quantity, and can not change at all. So its shift must be an artifact of postselection.*

After interaction, the whole system is $\frac{1}{\sqrt{2}}(e^{i\theta}e^{igp}|0\rangle + e^{-igp}|1\rangle)\varphi(p)$, so a relative phase appears in the quantum system, and the entanglement between the quantum system and meter system disappears. The meter is still $\varphi(p)$. When the final state $|\psi_{f,1}\rangle$ is selected, the probability for measure p is $P_1(p, \theta) = |\frac{1}{2}(e^{i\theta}e^{igp} - e^{-igp})\varphi(p)|^2 = \frac{1}{2}(1 - \cos(\theta + 2gp))|\varphi(p)|^2$. We can see the destructive interference of the probe system is only a part of the whole story. In this new perspective postselection plays a more classical role.

Ref. [10] goes further. They choose both of the final orthogonal states $|\psi_{f,2}\rangle = \frac{1}{\sqrt{2}}(|0\rangle + i|1\rangle)$ and $|\psi_{f,3}\rangle = \frac{1}{\sqrt{2}}(|0\rangle - i|1\rangle)$ for an almost balanced detection, which is an extension of the joint weak measurements in Ref [12]. That is, they give up the traditional postselection, and exploit new postselection. This is the reason why they claim “without postselection”. This change is a significant progress for AWA. Thus the detection probability for p at two ports are respectively $P_2(p, \theta) = \frac{1}{2}(1 + \sin(\theta + 2gp))|\varphi(p)|^2$ and $P_3(p, \theta) = \frac{1}{2}(1 - \sin(\theta + 2gp))|\varphi(p)|^2$. The sum and difference of $P_2(p, \theta)$ and $P_3(p, \theta)$ are

$$P_+(p, \theta) = P_2(p, \theta) + P_3(p, \theta) = |\varphi(p)|^2, \quad (1)$$

and

$$P_-(p, \theta) = P_2(p, \theta) - P_3(p, \theta) = |\varphi(p)|^2 \sin(\theta + 2gp). \quad (2)$$

The amplification effect results from the difference signal

*suiyueqiaoqiao@163.com

†572262643@qq.com

(2), which can be remarkably different from the sum one (1).

The results in [9, 10] are beyond our expectation. Now we have two AWA schemes. From the view in [10], we can easily improve the traditional AWA. We also detect the bright port $|\psi_{f,4}\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$ and its detection probability for p is $P_4(p, \theta) = \frac{1}{2}(1 + \cos(\theta + 2gp))|\varphi(p)|^2$. So $P_+(p, \theta) = P_1(p, \theta) + P_4(p, \theta) = |\varphi(p)|^2$ and an amplification signal can be defined as

$$\begin{aligned} P'_-(p, \theta) &= P_+(p, \theta) - (P_1(p, \theta) - P_4(p, \theta)) \\ &= |\varphi(p)|^2(1 - \cos(\theta + 2gp)). \end{aligned} \quad (3)$$

In this improved scheme all of the information are used.

We can generalize the idea in [10] to arbitrarily orthogonal postselections, $|\psi_{f,+}\rangle = \frac{1}{\sqrt{2}}(|0\rangle + e^{i\chi}|1\rangle)$ and $|\psi_{f,-}\rangle = \frac{1}{\sqrt{2}}(|0\rangle - e^{i\chi}|1\rangle)$, where $\chi \in [0, \pi]$. So the detection probability for p are

$$\begin{aligned} Pr_{\pm}(p, \theta) &= \frac{1}{2}(1 \pm \cos(\theta + 2gp - \chi))|\varphi(p)|^2 \\ &= \frac{1}{2}(1 \pm \cos(\theta + 2gp) \cos(\chi) \\ &\quad \mp \sin(\theta + 2gp) \sin(\chi))|\varphi(p)|^2, \end{aligned} \quad (4)$$

thus amplification signal can be defined as

$$\begin{aligned} Pr(p, \theta) &= P_+(p, \theta) - (Pr_+(p, \theta) - Pr_-(p, \theta)) \\ &= (1 - \cos(\theta + 2gp) \cos(\chi) \\ &\quad + \sin(\theta + 2gp) \sin(\chi))|\varphi(p)|^2 \\ &= (1 - \cos(\theta + 2gp - \chi))|\varphi(p)|^2. \end{aligned} \quad (5)$$

This is the general result for AWA.

-
- [1] Y. Aharonov, D. Z. Albert, and L. Vaidman, Phys. Rev. Lett. **60**, 1351 (1988).
 - [2] O. Hosten and P. Kwait, Science **319**, 787 (2008).
 - [3] P. B. Dixon, D. J. Starling, A. N. Jordan, and J. C. Howell, Phys. Rev. Lett. **102**, 173601 (2009).
 - [4] G. Li, T. Wang, S. Xu, H. S. Song, arXiv:1507.00850.
 - [5] G. Li, M. Y. Ye, H. S. Song, arXiv:1508.05705.
 - [6] Y. Shikano and A. Hosoya, J. Phys. A: Math. Theor. **43**, 025304 (2010).
 - [7] J. Dressel, M. Malik, F. M. Miatto, A. N. Jordan, and R. W. Boyd, Rev. Mod. Phys. **86**, 307 (2014).
 - [8] Surely we can also define a real result.
 - [9] T. Wang, L. H. Jiang, G. Li, C. B. Fu and X. M. Su, arXiv:1509.02250v2.
 - [10] J. Martínez-Rincón, W. T. Liu, G. I. Viza, and J. C. Howell, arXiv:1509.04810v1.
 - [11] Ref. [9] discussed thermal light cross-Kerr effect. The initial states of the quantum system and the meter system are $|\psi_i\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$ and the thermal state. The weak interaction is $e^{i\phi_0 \hat{n}_q \hat{n}_m}$, where ϕ_0 is the cross-Kerr phase shift per photon, $\hat{n}_q(\hat{n}_m)$ is the number operator for the quantum (meter) system. This is a special case for the new perspective. A general statement is given in Ref. [10]. However the meaning behind the new perspective is not pointed out.
 - [12] G. Strübi and C. Bruder, Phys. Rev. Lett. **110**, 083605 (2013).