

Inert Doublet Model in the light of LHC and astrophysical data

Agnieszka Ilnicka^{a,b}, Maria Krawczyk^c, Tania Robens^d

^a *Institute of Physics, University of Zürich
Winterthurststrasse 190, CH-8057 Zürich*

^b *Physics Department, ETH Zürich
Otto-Stern-Weg 5, CH - 8093 Zürich*

^c *Faculty of Physics, University of Warsaw
ul. Pasteura 5, 02-093 Warsaw*

^d *TU Dresden, Institut für Kern- und Teilchenphysik
Zellescher Weg 19, D-01069 Dresden*

E-mail:

ailnicka@physik.uzh.ch, maria.krawczyk@fuw.edu.pl, Tania.Robens@tu-dresden.de

We investigate the parameter space of the Inert Doublet Model, which is a straightforward extension of the SM in the scalar sector. We apply a set of constraints both from the theoretical and experimental side to extract and determine allowed regions of parameter space. These constraints put strong limits on both masses and couplings of the new particles. We also present a set of benchmarks for the current LHC run. This work is based on [1, 2].

*The European Physical Society Conference on High Energy Physics
22–29 July 2015
Vienna, Austria*

*Speaker.

1. Introduction

The Inert Doublet Model (IDM) is one of the most straightforward extensions of the Standard Model. In this model, the scalar sector is augmented by a second complex doublet, and an exact Z_2 symmetry is imposed on the Lagrangian. The first doublet corresponds to the SM Higgs doublet with the Higgs particle, and is responsible for electroweak symmetry breaking (EWSB) in a standard way. The second doublet, called dark doublet, contains a stable dark matter candidate. The IDM was studied in context of LHC phenomenology, both with respect to the Higgs boson discovery [3, 4] as well as dark matter discovery, the latter e.g. in the two lepton + MET channel [5, 6]. Moreover the model offers also rich cosmological phenomenology, for a review of references see [2].

The discovery of a Higgs boson in 2012 basically fixes the particle content of the first doublet in the IDM, in analogy to the scalar sector of the SM.¹ In the light of LHC run II knowledge about regions of parameter space which are in agreement with all current constraints is imminent in order to correctly determine open search channels and their experimental signatures. We address this need by presenting a complete survey on the model's parameter space including a wide range of constraints, coming from theoretical bounds as well as collider and astrophysical data. We additionally provide a set of benchmark points and planes for the current LHC run.

2. The Model

The scalar sector of IDM consists of two doublets of complex scalar fields, which we label ϕ_S for the SM-like doublet, and ϕ_D - for the dark doublet. After EWSB, only ϕ_S acquires a nonzero vacuum expectation value (v). The Z_2 - symmetric potential reads:

$$V = -\frac{1}{2} \left[m_{11}^2 (\phi_S^\dagger \phi_S) + m_{22}^2 (\phi_D^\dagger \phi_D) \right] + \frac{\lambda_1}{2} (\phi_S^\dagger \phi_S)^2 + \frac{\lambda_2}{2} (\phi_D^\dagger \phi_D)^2 + \lambda_3 (\phi_S^\dagger \phi_S) (\phi_D^\dagger \phi_D) + \lambda_4 (\phi_S^\dagger \phi_D) (\phi_D^\dagger \phi_S) + \frac{\lambda_5}{2} \left[(\phi_S^\dagger \phi_D)^2 + (\phi_D^\dagger \phi_S)^2 \right], \quad (2.1)$$

with the Z_2 transformation being defined by $\phi_S \rightarrow \phi_S, \phi_D \rightarrow -\phi_D, SM \rightarrow SM$. Due to this symmetry the lightest particle of the dark sector is stable. In total, the dark sector contains 4 new particles: H, A and H^\pm . We here choose H to be the dark matter (DM) candidate².

The Higgs boson data and electroweak precision observables fixes the SM-like Higgs mass M_h and v , and we are left with 5 free parameters, for which we take:

$$M_H, M_A, M_{H^\pm}, \lambda_2, \lambda_{345}, \quad (2.2)$$

where the $\lambda_{345} = \lambda_3 + \lambda_4 + \lambda_5$ describes coupling between SM-like Higgs and DM particle H and λ_2 corresponds to self-couplings of the dark scalars.

¹ After its discovery, several analyses studied the impact of this discovery as well as the signal strength measurements on the particles parameter space [7, 6, 8, 9].

² A priori, any of the new scalars can function as a dark matter candidate. However, we neglect the choice of a charged dark matter candidate, as these are strongly constrained [10]. Choosing A instead of H changes the meaning of λ_5 , but not the overall phenomenology of the model, cf. [2].

3. Scan Procedure

3.1 Constraints

Here we enumerate the constraints which were included in our analysis [2].

Theoretical constraints:

- positivity: potential is bounded from below at tree level
- perturbative unitarity of $2 \rightarrow 2$ scalar scattering matrix
- perturbativity of all couplings, (we chose 4π as upper limit)
- condition to be in the inert vacuum [11, 12].

Experimental constraints

- Mass of the SM-like Higgs boson h set to $M_h = 125.1$ GeV in agreement with results from the LHC experiments [13]
- Total width of the h obey an upper limit $\Gamma_{\text{tot}} \leq 22$ MeV [14, 15]
- Bounds provided by the total width of the electroweak gauge bosons:

$$M_{A,H} + M_H^\pm \geq m_W, M_A + M_H \geq m_Z, 2M_H^\pm \geq m_Z$$

- Bound on the lower mass of $M_H^\pm \geq 70$ GeV [16].
- Agreement with the current null-searches from the LEP, Tevatron, and LHC experiments using HiggsBounds [17, 18, 19]
- Agreement within 2σ for the 125 GeV Higgs signal strength measurements using HiggsSignals [20]
- 2σ agreement with electroweak precision observables, parameterized through the (correlated) electroweak oblique parameters S, T, U [21, 22, 23, 24].
- Upper limit of the H^+ lifetime $\tau \leq 10^{-7} s$, leading to $\Gamma_{\text{tot}} \geq 6.58 \times 10^{-18}$ GeV.
- Upper limit on relic density within 2σ from measurement of the Planck experiment [25]: $\Omega_c h^2 \leq 0.1241$
- Respect direct detection limits from dark matter nucleon scattering: the most stringent bounds are provided by the LUX experiment [26]
- Obey exclusions from recasted SUSY LEP and LHC analyses [27, 28]

3.2 Scan setup

We have performed a flat scan over the input parameters as given in eq. (2.2), where we have run up 1 TeV with the mass of the DM candidate. We confronted these points with the above constraints (sec. 3.1), where for each point we memorised the exclusion criteria if applicable. Our scan is performed in several steps: first we check the theoretical constraints as well as S, T, U parameters and total widths using 2HDMC [29]. In a second step, points which pass these bounds were confronted with null searches and Higgs signal strength measurements using HiggsBounds and HiggsSignals, respectively [17, 18, 19, 20]. In the final step of the scan, the calculation of Dark Matter observables, i.e. the total relic density as well as the direct detection cross section, was performed using MicrOmegas [30] and confronted with Planck and LUX measurements [25, 26]. For points which are in agreement with all bounds, we provide cross sections for pair-production of dark scalars, where we employed MadGraph [31], using the IDM UFO model presented in [32].

4. Allowed Parameter Space of IDM

In this section, we present the results of our scans and we emphasise the source of the strongest bounds, following the order of checks as discussed above. In the left panel of Figure 1 we show the region of parameter space where the dark scalar mass M_H is smaller than the SM like Higgs mass of 125 GeV. This region is strongly constrained by the combination of the 125 GeV Higgs width and the signal strength, leaving a very narrow allowed stripe of λ_{345} , with absolute values $\lesssim 0.02$. Also astrophysical data pose important constraints in this region. Relic density requires the mass of the DM candidate to values $\gtrsim 45$ GeV. In addition, LUX measurements narrow down the allowed values for the λ_{345} coupling. In the right panel of Figure 1, we display the whole region of mass values for M_H up to 1 TeV. Here, especially LUX data limit the models parameter space, reflecting the dependance of the direct detection cross section on these parameters.




Figure 1: Dark matter mass vs DM-SM coupling planes: (left) a zoom into the low mass region, with $M_H \leq 100$ GeV showing the importance of constraints from LHC and astrophysical measurements; (right) the general case with M_H up to 1 TeV, where bounds from direct detection are dominant.

The interplay of all constraints, presented in the left panel of Figure 2, leads to the strict mass hierarchy

$$M_H < M_A \leq M_{H+}.$$

Also clearly visible is the preference for more degenerate masses in the dark sector and the more relaxed parameter space for high masses.




Figure 2: (left) The exclusion plot in M_A vs M_{H^+} plane. (right) The plot presents leading contribution to DM relic density in the DM mass vs relic density plane. (Exclusion from LUX not included).

In right panel of Figure 2 we present the leading contributions to the DM relic density. The relic density produced by IDM particles can be large for very low or relatively high DM masses. The dominant channel for most of the parameter space for mass of H above 100 GeV is annihilation into vector bosons.

5. Benchmarks

For the points allowed by all constraints the leading order cross section dark scalars pair-production³ was calculated. The dominating channel leading to visible collider signatures is HA production⁴, and Figure 3 shows its dependence on the masses and coupling of dark matter. Figure 3 shows that the production cross section is mainly driven by kinematics, and especially by particles masses. This follows from the fact that the dominant production mechanism, namely production via Z mediation in s-channel, only depends on kinematics and the SM electroweak couplings, but not on couplings of the extended scalar sector which are absent in the SM.

From the presented points, five were chosen as benchmarks [1, 33, 2], see Table 5. While benchmarks I and II are exceptional points in a sense that the allowed parameter space is extremely constrained in the low mass region, benchmarks III to V are more typical, as these parts of the parameter space are more highly populated. Furthermore, for scenario IV the production cross sections for HA and H^+H^- have similar order of magnitude.

6. Conclusions

The Inert Doublet Model is a very promising extension of the SM in the scalar sector. Its parameter space is subject to several theoretical and experimental constraints. In a flat scan, high,

³Due to the Z_2 symmetry dark sector particles are always produced in pairs.

⁴As most DM models at colliders, the IDM will always lead to signatures including missing transverse energy.




Figure 3: The planes of allowed points, with HA production cross sections (in pb) at a 13 TeV LHC.

BP	BP I	BP II	BP III	BP IV	BP V
$M_H [GeV]$	57.5	85.5	128.0	363.0	311.0
$M_A [GeV]$	113.0	111.0	134.0	374.0	415.0
$M_{H^\pm} [GeV]$	123.0	140.0	176.0	374.0	447.0
$ \lambda_{345} $	[0.002;0.015]	[0;0.015]	[0;0.05]	[0,0.25]	[0;0.19]
$ \lambda_2 $	[0,4.2]	[0,4.2]	[0,4.2]	[0,4.2]	[0,4.2]
$\sigma(pp \rightarrow HA)[pb]$	0.371(4)	0.226 (2)	0.0765 (7)	0.00122(1)	0.00129 (1)
$\sigma(pp \rightarrow H^+H^-)[pb]$	0.097 (1)	0.0605 (9)	0.0259 (3)	0.00124 (1)	0.000553 (7)
$BR(H^+ \rightarrow HW^+)$	0.99	0.96	0.66	1	0.99

Table 1: Benchmark points for dark scalars pair production at LHC run 2.

nearly degenerate masses of the dark particles are favoured, leaving however some viable parameter space for low dark matter masses ($M_H < \frac{M_h}{2}$). The pair production of dark particles at the LHC is mainly determined by their masses and regions with sizeable cross sections are subject to much more severe limits. The current LHC run will hopefully allow to provide more insight into this model, either by strengthening the above constraints or by means of a possible discovery.

Acknowledgement

This work is supported by the 7th Framework Programme of the European Commission through the Initial Training Network HiggsTools PITN-GA-2012-316704. This work was also partly supported by the Polish grant NCN OPUS 2012/05/B/ST2/03306 (2012-2016).

References

- [1] Agnieszka Ilnicka, Maria Krawczyk, and Tania Robens. Constraining the Inert Doublet Model. In *2nd Toyama International Workshop on Higgs as a Probe of New Physics (HPNP2015) Toyama, Japan, February 11-15, 2015*, 2015, 1505.04734.
- [2] Agnieszka Ilnicka, Maria Krawczyk, and Tania Robens. The Inert Doublet Model in the light of LHC and astrophysical data – An Update –. 2015, 1508.01671.
- [3] Riccardo Barbieri, Lawrence J. Hall, and Vyacheslav S. Rychkov. Improved naturalness with a heavy Higgs: An Alternative road to LHC physics. *Phys.Rev.*, D74:015007, 2006, hep-ph/0603188.

- [4] Qing-Hong Cao, Ernest Ma, and G. Rajasekaran. Observing the Dark Scalar Doublet and its Impact on the Standard-Model Higgs Boson at Colliders. *Phys.Rev.*, D76:095011, 2007, 0708.2939.
- [5] Ethan Dolle, Xinyu Miao, Shufang Su, and Brooks Thomas. Dilepton Signals in the Inert Doublet Model. *Phys.Rev.*, D81:035003, 2010, 0909.3094.
- [6] Michael Gustafsson, Sara Rydbeck, Laura Lopez-Honorez, and Erik Lundstrom. Status of the Inert Doublet Model and the Role of multileptons at the LHC. *Phys.Rev.*, D86:075019, 2012, 1206.6316.
- [7] Bogumila Swiezewska and Maria Krawczyk. Diphoton rate in the inert doublet model with a 125 GeV Higgs boson. *Phys.Rev.*, D88(3):035019, 2013, 1212.4100.
- [8] Abdesslam Arhrib, Yue-Lin Sming Tsai, Qiang Yuan, and Tzu-Chiang Yuan. An Updated Analysis of Inert Higgs Doublet Model in light of the Recent Results from LUX, PLANCK, AMS-02 and LHC. *JCAP*, 1406:030, 2014, 1310.0358.
- [9] Maria Krawczyk, Dorota Sokolowska, Pawel Swaczyna, and Bogumila Swiezewska. Constraining Inert Dark Matter by $R_{\gamma\gamma}$ and WMAP data. *JHEP*, 1309:055, 2013, 1305.6266.
- [10] Leonid Chuzhoy and Edward W. Kolb. Reopening the window on charged dark matter. *JCAP*, 0907:014, 2009, 0809.0436.
- [11] Bogumila Swiezewska. Yukawa independent constraints for two-Higgs-doublet models with a 125 GeV Higgs boson. *Phys.Rev.*, D88(5):055027, 2013, 1209.5725.
- [12] Michael Gustafsson. The Inert Doublet Model and Its Phenomenology. *PoS*, CHARGED2010:030, 2010, 1106.1719.
- [13] Georges Aad et al. Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS Experiments. *Phys. Rev. Lett.*, 114:191803, 2015, 1503.07589.
- [14] Vardan Khachatryan et al. Constraints on the Higgs boson width from off-shell production and decay to Z-boson pairs. *Phys.Lett.*, B736:64, 2014, 1405.3455.
- [15] Georges Aad et al. Constraints on the off-shell Higgs boson signal strength in the high-mass ZZ and WW final states with the ATLAS detector. *Eur. Phys. J.*, C75(7):335, 2015, 1503.01060.
- [16] Aaron Pierce and Jesse Thaler. Natural Dark Matter from an Unnatural Higgs Boson and New Colored Particles at the TeV Scale. *JHEP*, 0708:026, 2007, hep-ph/0703056.
- [17] Philip Bechtle, Oliver Brein, Sven Heinemeyer, Georg Weiglein, and Karina E. Williams. HiggsBounds: Confronting Arbitrary Higgs Sectors with Exclusion Bounds from LEP and the Tevatron. *Comput.Phys.Commun.*, 181:138–167, 2010, 0811.4169.
- [18] Philip Bechtle, Oliver Brein, Sven Heinemeyer, Georg Weiglein, and Karina E. Williams. HiggsBounds 2.0.0: Confronting Neutral and Charged Higgs Sector Predictions with Exclusion Bounds from LEP and the Tevatron. *Comput.Phys.Commun.*, 182:2605–2631, 2011, 1102.1898.
- [19] Philip Bechtle, Oliver Brein, Sven Heinemeyer, Oscar Stal, Tim Stefaniak, et al. HiggsBounds – 4: Improved Tests of Extended Higgs Sectors against Exclusion Bounds from LEP, the Tevatron and the LHC. *Eur.Phys.J.*, C74(3):2693, 2014, 1311.0055.
- [20] Philip Bechtle, Sven Heinemeyer, Oscar Stal, Tim Stefaniak, and Georg Weiglein. *HiggsSignals*: Confronting arbitrary Higgs sectors with measurements at the Tevatron and the LHC. *Eur.Phys.J.*, C74(2):2711, 2014, 1305.1933.
- [21] Guido Altarelli and Riccardo Barbieri. Vacuum polarization effects of new physics on electroweak processes. *Phys. Lett. B*, 253:161, 1991.

- [22] Michael E. Peskin and Tatsu Takeuchi. A New constraint on a strongly interacting Higgs sector. *Phys.Rev.Lett.*, 65:964–967, 1990.
- [23] Michael E. Peskin and Tatsu Takeuchi. Estimation of oblique electroweak corrections. *Phys.Rev.*, D46:381–409, 1992.
- [24] I. Maksymyk, C.P. Burgess, and David London. Beyond S, T and U. *Phys.Rev.*, D50:529–535, 1994, hep-ph/9306267.
- [25] P.A.R. Ade et al. Planck 2015 results. XIII. Cosmological parameters. 2015, 1502.01589.
- [26] D.S. Akerib et al. First results from the LUX dark matter experiment at the Sanford Underground Research Facility. *Phys.Rev.Lett.*, 112(9):091303, 2014, 1310.8214.
- [27] Erik Lundstrom, Michael Gustafsson, and Joakim Edsjo. The Inert Doublet Model and LEP II Limits. *Phys.Rev.*, D79:035013, 2009, 0810.3924.
- [28] Genevieve Belanger, Beranger Dumont, Andreas Goudelis, Bjorn Herrmann, Sabine Kraml, and Dipan Sengupta. Dilepton constraints in the Inert Doublet Model from Run 1 of the LHC. *Phys. Rev.*, D91(11):115011, 2015, 1503.07367.
- [29] David Eriksson, Johan Rathsman, and Oscar Stal. 2HDMC: Two-Higgs-Doublet Model Calculator Physics and Manual. *Comput.Phys.Commun.*, 181:189–205, 2010, 0902.0851.
- [30] G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov. micrOMEGAs_3: A program for calculating dark matter observables. *Comput. Phys. Commun.*, 185:960–985, 2014, 1305.0237.
- [31] Johan Alwall, Michel Herquet, Fabio Maltoni, Olivier Mattelaer, and Tim Stelzer. MadGraph 5 : Going Beyond. *JHEP*, 1106:128, 2011, 1106.0522.
- [32] A. Goudelis, B. Herrmann, and O. Stal. Dark matter in the Inert Doublet Model after the discovery of a Higgs-like boson at the LHC. *JHEP*, 1309:106, 2013, 1303.3010.
- [33] A. Ilnicka, M. Krawczyk, and T. Robens. Inert Doublet Model benchmarks for the 13 TeV run of the LHC. Submitted to the Higgs Cross Section Working Group, April 2015.