

THE ROLE OF THE PAULI-LUBAŃSKI VECTOR FOR THE DIRAC, WEYL, PROCA, MAXWELL, AND FIERZ-PAULI EQUATIONS

SERGEY I. KRYUCHKOV, NATHAN A. LANFEAR, AND SERGEI K. SUSLOV

ABSTRACT. We analyze basic relativistic wave equations for the classical fields, such as Dirac's equation, Weyl's two-component equation for massless neutrinos, and the Proca, Maxwell, and Fierz-Pauli equations, from the viewpoint of the Pauli-Lubański vector and the Casimir operators of the Poincaré group. In general, in this group-theoretical approach, the above wave equations arise in certain overdetermined forms, which can be reduced to the conventional ones by a Gaussian elimination. A connection between the spin of a particle/field and consistency of the corresponding overdetermined system is emphasized in the massless case.

1. INTRODUCTION

All physically interesting representations of the proper orthochronous inhomogeneous Lorentz group (known nowadays as the Poincaré group) were classified by Wigner [70] and, since then, this approach has been utilized for the mathematical description of mass and spin of an elementary particle. To this end, the Pauli-Lubański pseudo-vector is introduced,

$$w_\mu = \frac{1}{2} e_{\mu\nu\sigma\tau} p^\nu M^{\sigma\tau}, \quad p_\mu w^\mu = 0, \quad (1.1)$$

where p_μ is the relativistic linear momentum operator and $M^{\sigma\tau}$ are the angular momentum operators, or generators of the proper orthochronous Lorentz group, with summation over repeated indices. (We use Einstein summation convention unless stated otherwise.) The mass and spin of a particle are defined in terms of two quadratic invariants (Casimir operators of the Poincaré group) as follows

$$p^2 = p_\mu p^\mu = m^2, \quad w^2 = w_\mu w^\mu = -m^2 s(s+1), \quad (1.2)$$

when $m > 0$ (see, for example, [3], [4], [5], [10], [36], [37], [38], [54], [55], [56], [58], [60] and the references therein; throughout the article we use the standard notations in the Minkowski space-time \mathbb{R}^4 and the natural units $c = \hbar = 1$).

For the massless fields, when $m = 0$, one gets $w^2 = p^2 = pw = 0$, and the Pauli-Lubański vector should be proportional to p :¹

$$w_\mu = \lambda p_\mu \quad (1.3)$$

Date: January 22, 2016.

1991 Mathematics Subject Classification. Primary 35Q41, 35Q61, 35L05; Secondary 35N05, 35Q75.

Key words and phrases. Poincaré group, Pauli-Lubański vector, Dirac's equation, Weyl's equations for massless neutrinos, Proca and Maxwell's equations, Fierz-Pauli equations, linearized Einstein equations.

¹This assumption was made by Bargmann and Wigner [4] for the massless limit of the spinor wave equation for particles with an arbitrary integer or half-integer spin proposed by Dirac [13] (see also [17], [18], [31], [48] and the references therein). The pseudo-vector (1.1) was introduced, in a slightly different notation, by Eqs. (4.a)–(4.b) of Ref. [4].

(acting on common eigenstates [45], [55]). The number λ is sometimes called the helicity of the representation and the value $s = |\lambda|$ is called the spin of a particle with zero mass [10], [55], [56], [58]. Although the concept of helicity is discussed in most textbooks on quantum field theory, a practical implementation of this definition of the spin of a massless particle deserves a certain clarification. As is shown in our previous article [32], in the case of the electromagnetic field in vacuum, the constant λ in the latter equation is fixed, otherwise violating the classical Maxwell equations. Thus, for the photon, or a harmonic circular classical electromagnetic wave², the latter equation allows one to introduce the field equations and spin, but not the helicity, when a certain choice of eigenstates is required. A similar situation occurs in the case of Weyl's equation for massless neutrinos. (In Ref. [32], we do not discuss the equation for a graviton, another massless spin-2 particle; it will be analyzed elsewhere; cf. [28], [42], [44], [49] and our discussion in section 7.)

The theory of relativistic-invariant wave equations is studied, from different perspectives, in numerous classical accounts [4], [5], [7], [12], [13], [17], [18], [23], [31], [39], [48], [52], [68] (see also [8], [36], [37], [38] and the references therein). Nonetheless, in our opinion, the importance of the Pauli-Lubański vector for conventional relativistic equations, which allows one to derive all of them directly from the postulated transformation law of the corresponding classical field in pure group-theoretical terms, is not fully appreciated. In this article, we would like to start from Dirac's relativistic electron, or any free relativistic particle with a nonzero mass and spin 1/2, which can be described by a bispinor wave function. Our analysis shows that an analog of the linear operator relation (1.3) takes the form,

$$w_\mu = \frac{1}{2} (p_\mu + m\gamma_\mu) \gamma_5, \quad (1.4)$$

provided that the Dirac equation, $(\gamma^\mu p_\mu - m) \psi = 0$, holds, when the corresponding overdetermined system of equations is consistent. This automatically implies that $s = 1/2$, in the covariant form, by definition (1.2). (We were not able to find the operator relation (1.4) in the extensive literature on Dirac's equation.)

In the rest of the article, a similar program is utilized, in a systematic way and from first principles, for other familiar relativistic wave equations. Once again, we postulate the transformation law of the field in question (a law of nature) and, with the help of the corresponding Lorentz generators, evaluate the action of the Pauli-Lubański vector on the field in order to compute, eventually, not one, but both Casimir operators (1.2). If a linear relation, similar to (1.3) or (1.4), does exist, one obtains an overdetermined PDE system, which can be reduced to the corresponding relativistic wave equation by a matrix version of Gaussian elimination [22]. We show that this approach allows one to derive equations of motion for the most useful classical fields, including the Weyl, Proca, Fierz-Pauli, and Maxwell equations in vacuum, as a statement of consistency for the original overdetermined systems. At the moment, we shall not discuss relativistic wave equations for particles with an arbitrary spin, such as the Bargmann-Wigner equations, Majorana equations, and/or the (first order) Duffin-Kemmer equations which also describe spin-0 and spin-1 fields (see, for example, [5], [9], [24] and [59] for more details; the case of the Klein-Gordon equation, or the relativistic Schrödinger equation [57], is, of course, obvious).

We have also entirely concentrated on four dimensions. A more general group-theoretical approach to the relativistic wave equations, which allows one to include higher dimensions and spins, is formulated in [5] with the help of induced representations of the semi-direct products of separable, locally compact groups. Spinors in arbitrary dimensions are also discussed in [60].

²Multiple meanings of the word "photon" are analyzed in [30].

2. DIRAC EQUATION

In this section, for the reader's convenience, we summarize basic facts about Dirac's equation and then discuss its relation with the Pauli-Lubański vector (1.4). To this end, a familiar bispinor representation of the proper orthochronous Lorentz group $SO_+(1, 3)$ is used.

2.1. Gamma matrices, bispinors, and transformation laws. We shall use the following Dirac matrices: $\gamma^\mu = (\gamma^0, \boldsymbol{\gamma})$, $\gamma_\mu = g_{\mu\nu}\gamma^\nu = (\gamma^0, -\boldsymbol{\gamma})$, and $\gamma_5 = -\gamma^5 = i\gamma^0\gamma^1\gamma^2\gamma^3$, where

$$\boldsymbol{\gamma} = \begin{pmatrix} 0 & \boldsymbol{\sigma} \\ -\boldsymbol{\sigma} & 0 \end{pmatrix}, \quad \gamma^0 = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad \gamma^5 = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \quad (2.1)$$

and $\boldsymbol{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ are the standard 2×2 Pauli matrices [47], [62]. The familiar anticommutation relations,

$$\gamma^\mu\gamma^\nu + \gamma^\nu\gamma^\mu = 2g^{\mu\nu}, \quad \gamma^\mu\gamma^5 + \gamma^5\gamma^\mu = 0 \quad (\mu, \nu = 0, 1, 2, 3), \quad (2.2)$$

hold. (Most of the results here will not depend on a particular choice of gamma matrices, but it is always useful to have an example in mind.) The four-vector notation, $x^\mu = (t, \mathbf{r})$, $\partial_\mu = \partial/\partial x^\mu$, and $\partial^\alpha = g^{\alpha\mu}\partial_\mu$ in natural units $c = \hbar = 1$ with the standard metric, $g_{\mu\nu} = g^{\mu\nu} = \text{diag}(1, -1, -1, -1)$, in the Minkowski space-time \mathbb{R}^4 are utilized throughout the article [6], [9], [10], [42], [59].

In this notation, the transformation law of a bispinor wave function³,

$$\psi(x) = \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \\ \psi_4 \end{pmatrix} \in \mathbb{C}^4, \quad (2.3)$$

under a proper Lorentz transformation, is given by

$$\psi'(x') = S_\Lambda \psi(x), \quad x' = \Lambda x, \quad (2.4)$$

together with the rule,

$$S_\Lambda^{-1} \gamma^\mu S_\Lambda = \Lambda^\mu_\nu \gamma^\nu, \quad (2.5)$$

for the sake of covariance of the celebrated Dirac equation,

$$i\gamma^\mu \partial_\mu \psi - m\psi = 0. \quad (2.6)$$

As is well known, a general solution of the latter matrix equation has the form

$$S = S_\Lambda = \exp\left(-\frac{1}{4}\theta_{\mu\nu}\Sigma^{\mu\nu}\right), \quad \theta_{\mu\nu} = -\theta_{\nu\mu}, \quad (2.7)$$

$$\Sigma^{\mu\nu} = (\gamma^\mu\gamma^\nu - \gamma^\nu\gamma^\mu)/2$$

(with summation over every two repeated indices; see, for example, [27], [43]) and, in turn,

$$S_\Lambda^{-1} \Sigma^{\mu\nu} S_\Lambda = \Lambda^\mu_\sigma \Lambda^\nu_\tau \Sigma^{\sigma\tau}. \quad (2.8)$$

³The relativistic wave equation for a massive spin 1/2 particle was proposed by Dirac [12], when only tensor representations of the Lorentz group were known. Thus, the problem of covariance of Dirac's equation gave rise to a new class of representations of the Lorentz group, namely, the spinor representations [5].

In explicit form,

$$\begin{aligned}\Sigma^{\mu\nu} &= \frac{1}{2}(\gamma^\mu\gamma^\nu - \gamma^\nu\gamma^\mu) = \begin{pmatrix} 0 & \alpha_q \\ -\alpha_p & -ie_{pqr}\Sigma_r \end{pmatrix} \\ &= \begin{pmatrix} 0 & \alpha_1 & \alpha_2 & \alpha_3 \\ -\alpha_1 & 0 & -i\Sigma_3 & i\Sigma_2 \\ -\alpha_2 & i\Sigma_3 & 0 & -i\Sigma_1 \\ -\alpha_3 & -i\Sigma_2 & i\Sigma_1 & 0 \end{pmatrix},\end{aligned}\tag{2.9}$$

where, by definition,

$$\Sigma = \begin{pmatrix} \boldsymbol{\sigma} & 0 \\ 0 & \boldsymbol{\sigma} \end{pmatrix}, \quad \boldsymbol{\alpha} = \begin{pmatrix} 0 & \boldsymbol{\sigma} \\ \boldsymbol{\sigma} & 0 \end{pmatrix}.\tag{2.10}$$

Their familiar product identities,

$$\begin{aligned}\Sigma_p\Sigma_q &= ie_{pqr}\Sigma_r + \delta_{pq}, & \alpha_p\alpha_q &= ie_{pqr}\Sigma_r + \delta_{pq}, \\ \alpha_p\Sigma_q &= \Sigma_p\alpha_q = ie_{pqr}\alpha_r + \delta_{pq}\gamma^5,\end{aligned}\tag{2.11}$$

hold.

Setting $\mathbf{n} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ for each of the unit vectors in the directions of the mutually orthogonal coordinate axes, one can write in compact form:

$$S_R = e^{i\theta(\mathbf{n}\cdot\boldsymbol{\Sigma})/2} = \cos\frac{\theta}{2} + i(\mathbf{n}\cdot\boldsymbol{\Sigma})\sin\frac{\theta}{2}, \quad (\mathbf{n}\cdot\boldsymbol{\Sigma})^2 = I\tag{2.12}$$

and

$$S_L = e^{-\vartheta(\mathbf{n}\cdot\boldsymbol{\alpha})/2} = \cosh\frac{\vartheta}{2} - (\mathbf{n}\cdot\boldsymbol{\alpha})\sinh\frac{\vartheta}{2}, \quad (\mathbf{n}\cdot\boldsymbol{\alpha})^2 = I\tag{2.13}$$

with $\tanh\vartheta = v$, in the cases of rotations and boosts, respectively [5], [43].

The important dual four-tensor identities,

$$ie_{\mu\nu\sigma\tau}\Sigma^{\sigma\tau} = 2\gamma_5\Sigma_{\mu\nu}, \quad ie^{\mu\nu\sigma\tau}\gamma_5\Sigma_{\mu\nu} = 2\Sigma^{\sigma\tau},\tag{2.14}$$

can be directly verified. Here, $e^{\mu\nu\sigma\tau} = -e_{\mu\nu\sigma\tau}$ and $e_{0123} = +1$ is the Levi-Civita symbol [10], [21].

For the conjugate bispinor,

$$\bar{\psi}(x) = \psi^\dagger(x)\gamma^0, \quad \bar{\psi}'(x') = \bar{\psi}(x)S_\Lambda^{-1}, \quad x' = \Lambda x,\tag{2.15}$$

the Dirac equation (2.6) takes the form

$$i\partial_\mu\bar{\psi}\gamma^\mu + m\bar{\psi} = 0.\tag{2.16}$$

(For more details see classical accounts [1], [6], [9], [16], [27], [43], [50], [55], [58], [59], [66].)

Examples. In particular, for the boost in the plane (x^0, x^1) , when

$$S_L = e^{-(\vartheta/2)\Sigma^{01}} = e^{-(\vartheta/2)\alpha_1} = \cosh\frac{\vartheta}{2} - \alpha_1\sinh\frac{\vartheta}{2}\tag{2.17}$$

with $\tanh\vartheta = v$, one can easily verify by matrix multiplication that

$$e^{(\vartheta/2)\alpha_1} \begin{pmatrix} \gamma^0 \\ \gamma^1 \\ \gamma^2 \\ \gamma^3 \end{pmatrix} e^{-(\vartheta/2)\alpha_1} = \begin{pmatrix} \cosh\vartheta & -\sinh\vartheta & 0 & 0 \\ -\sinh\vartheta & \cosh\vartheta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \gamma^0 \\ \gamma^1 \\ \gamma^2 \\ \gamma^3 \end{pmatrix}.\tag{2.18}$$

In a similar fashion, for the rotation in the plane (x^1, x^2) :

$$S_R = e^{-(\theta/2)\Sigma^{12}} = e^{i(\theta/2)\Sigma_3} = \cos \frac{\theta}{2} + i\Sigma_3 \sin \frac{\theta}{2} \quad (2.19)$$

and

$$e^{-i(\theta/2)\Sigma_3} \begin{pmatrix} \gamma^0 \\ \gamma^1 \\ \gamma^2 \\ \gamma^3 \end{pmatrix} e^{i(\theta/2)\Sigma_3} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \theta & \sin \theta & 0 \\ 0 & -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \gamma^0 \\ \gamma^1 \\ \gamma^2 \\ \gamma^3 \end{pmatrix}. \quad (2.20)$$

(See [5], [43] for further details.)

2.2. Generators and commutators. In the fundamental representation of the proper orthochronous Lorentz group, we shall choose the following six 4×4 real-valued matrices ($\alpha, \beta = 0, 1, 2, 3$ are fixed with no summation):

$$\begin{aligned} \Lambda(\theta_{\alpha\beta}) &= \exp(-\theta_{\alpha\beta} m^{\alpha\beta}), & m^{\alpha\beta} &= -m^{\beta\alpha}, \\ (m^{\alpha\beta})^\mu_\nu &= g^{\alpha\mu} \delta^\beta_\nu - g^{\beta\mu} \delta^\alpha_\nu \end{aligned} \quad (2.21)$$

for the corresponding one-parameter subgroups of rotations and boosts [10], [32], [42], [58]. Then, differentiation of a particular expression (for the corresponding tensor operator [5]),

$$e^{\theta_{\alpha\beta}\Sigma^{\alpha\beta}/2} \gamma^\mu e^{-\theta_{\alpha\beta}\Sigma^{\alpha\beta}/2} = \left(e^{-\theta_{\alpha\beta}m^{\alpha\beta}} \right)^\mu_\nu \gamma^\nu, \quad (2.22)$$

at $\theta_{\alpha\beta} = 0$ results in

$$[\Sigma^{\alpha\beta}, \gamma^\mu] := \Sigma^{\alpha\beta} \gamma^\mu - \gamma^\mu \Sigma^{\alpha\beta} = 2(g^{\beta\mu} \gamma^\alpha - g^{\alpha\mu} \gamma^\beta), \quad (2.23)$$

which can be independently verified with the help of (2.2).

In a similar fashion, the action of four-angular momentum operators⁴,

$$M^{\alpha\beta} = x^\beta \partial^\alpha - x^\alpha \partial^\beta, \quad \partial^\alpha = g^{\alpha\kappa} \partial_\kappa, \quad (2.24)$$

on Dirac's bispinors (2.3) can be derived directly from the transformation law as follows

$$\begin{aligned} M^{\alpha\beta} \psi &:= - \left[\frac{d}{d\theta_{\alpha\beta}} \psi'(\Lambda^\mu_\nu(\theta_{\alpha\beta}) x^\nu) \right] \Big|_{\theta_{\alpha\beta}=0} \\ &= - \left(\frac{d}{d\theta_{\alpha\beta}} e^{-\theta_{\alpha\beta}\Sigma^{\alpha\beta}/2} \right) \Big|_{\theta_{\alpha\beta}=0} \psi(x) = \frac{1}{2} \Sigma^{\alpha\beta} \psi \end{aligned} \quad (2.25)$$

(see also [27] for a slightly different derivation). A familiar commutator,

$$[\Sigma^{\alpha\beta}, \Sigma^{\sigma\tau}] = 2(g^{\beta\sigma} \Sigma^{\alpha\tau} - g^{\beta\tau} \Sigma^{\alpha\sigma} + g^{\alpha\tau} \Sigma^{\beta\sigma} - g^{\alpha\sigma} \Sigma^{\beta\tau}), \quad (2.26)$$

can be readily verified with the help of (2.23) and/or independently derived from (2.8). These results are independent of our choice of the gamma matrices representation.

⁴We follow [32]. Traditionally, $M^{\alpha\beta} \rightarrow -iM^{\alpha\beta}$; see, for example, [10].

2.3. Balance conditions and energy-momentum tensors. We shall use a familiar notation for the partial derivatives [10],

$$\mathcal{D}^p \psi(x) := \frac{\partial^{p_0+p_1+p_2+p_3}}{\partial x_0^{p_0} \partial x_1^{p_1} \partial x_2^{p_2} \partial x_3^{p_3}} \psi(x_0, x_1, x_2, x_3), \quad \mathcal{D}^0 \psi(x) := \psi(x) \quad (2.27)$$

where $p = (p_0, p_1, p_2, p_3)$ is an ordered set of non-negative integers $p_\mu \geq 0$. It follows from the Dirac equations (2.6) and (2.16) that

$$\partial_\mu [(\mathcal{D}^p \bar{\psi}(x)) \gamma^\mu (\mathcal{D}^q \psi(x))] = 0, \quad (2.28)$$

or, for a finite multi-sum,

$$\partial_\mu \left[\sum_{p,q} c_{p,q} \mathcal{D}^p \bar{\psi} \gamma^\mu \mathcal{D}^q \psi \right] = 0, \quad (2.29)$$

which can be thought of as a “master” differential balance condition set.

Indeed, in view of $\mathcal{D}^r \partial_\mu = \partial_\mu \mathcal{D}^r$, one gets

$$i \gamma^\mu \partial_\mu (\mathcal{D}^q \psi) = m (\mathcal{D}^q \psi), \quad i \partial_\mu (\mathcal{D}^p \bar{\psi}) \gamma^\mu = -m (\mathcal{D}^p \bar{\psi}). \quad (2.30)$$

Let us multiply the first (second) equation by $\mathcal{D}^p \bar{\psi} (\mathcal{D}^q \psi)$ from the left (right) and add the results. Then

$$\begin{aligned} i \partial_\mu (\mathcal{D}^p \bar{\psi} \gamma^\mu \mathcal{D}^q \psi) &= i \partial_\mu (\mathcal{D}^p \bar{\psi}) \gamma^\mu \mathcal{D}^q \psi + i \mathcal{D}^p \bar{\psi} \gamma^\mu \partial_\mu (\mathcal{D}^q \psi) \\ &= m (-\mathcal{D}^p \bar{\psi} \mathcal{D}^q \psi + \mathcal{D}^p \bar{\psi} \mathcal{D}^q \psi) \equiv 0. \end{aligned}$$

Among important special cases of (2.28) are the following important identities:

$$\partial_\mu j^\mu(x) = 0, \quad j^\mu(x) = \bar{\psi}(x) \gamma^\mu \psi(x), \quad (2.31)$$

corresponding to the total charge conservation, and

$$\partial_\mu [\bar{\psi}(x) \gamma^\mu \partial_\nu \psi(x)] = 0, \quad \partial_\mu [\partial_\nu \bar{\psi}(x) \gamma^\mu \psi(x)] = 0. \quad (2.32)$$

Therefore, one can introduce the energy-momentum tensor, such that $\partial_\mu T^\mu_\nu(x) = 0$, in two different forms

$$T^\mu_\nu := i \bar{\psi} \gamma^\mu \partial_\nu \psi, \quad T^\mu_\mu = m \bar{\psi} \psi \quad (2.33)$$

and/or

$$T^\mu_\nu := \frac{i}{2} [\bar{\psi} \gamma^\mu (\partial_\nu \psi) - (\partial_\nu \bar{\psi}) \gamma^\mu \psi], \quad T^\mu_\mu = m \bar{\psi} \psi. \quad (2.34)$$

As is well-known, all quantities of physical interest can be derived from the energy-momentum tensor [9], [58].

2.4. Variants of Dirac’s equation. In view of (2.6) and (2.2),

$$i \gamma^\mu \gamma^\nu \partial_\nu \psi = m \gamma^\mu \psi, \quad (2.35)$$

$$\gamma^\mu \gamma^\nu = \frac{1}{2} (\gamma^\mu \gamma^\nu - \gamma^\nu \gamma^\mu) + \frac{1}{2} (\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu) = \Sigma^{\mu\nu} + g^{\mu\nu},$$

and, as a result, we obtain an overdetermined but very convenient form of Dirac’s system:

$$i (\Sigma^{\mu\nu} + g^{\mu\nu}) \partial_\nu \psi = m \gamma^\mu \psi. \quad (2.36)$$

On the one hand, in a 3D “vector” form,

$$i\partial_0\psi = (-i\boldsymbol{\alpha} \cdot \nabla + m\beta)\psi, \quad \beta = \gamma^0 \quad (\hbar = c = 1, \quad \partial_0 = \partial/\partial t) \quad (2.37)$$

and

$$-i\partial_0\boldsymbol{\alpha}\psi + (\nabla \times \boldsymbol{\Sigma})\psi = (i\nabla + m\boldsymbol{\gamma})\psi. \quad (2.38)$$

It is worth noting that the latter vectorial equation in our overdetermined system (2.37)–(2.38) can be obtained by matrix multiplication, from the first one, in view of familiar relations:

$$\alpha_p\alpha_q = ie_{pqr}\Sigma_r + \delta_{pq}, \quad \boldsymbol{\alpha}\beta = -\boldsymbol{\gamma}. \quad (2.39)$$

On the other hand, by letting $p_\mu = i\partial_\mu$, one gets

$$(\Sigma^{\mu\nu} + g^{\mu\nu})p_\nu\psi = m\gamma^\mu\psi \quad (2.40)$$

and applying the momentum operator p_μ to the both sides:

$$p^2\psi = (\Sigma^{\mu\nu} + g^{\mu\nu})p_\mu p_\nu\psi = m\gamma^\mu p_\mu\psi, \quad (2.41)$$

in view of $\Sigma^{\mu\nu}p_\mu p_\nu = 0$. If $p^2\psi = m^2\psi$, we derive, once again, that $\gamma^\mu p_\mu\psi = m\psi$. Therefore both forms of the Dirac system, (2.6) and (2.36), are equivalent and every component of the bispinor (2.3) does satisfy the d’Alembert equation,

$$(\partial^\mu\partial_\mu + m^2)\psi = (\partial_t^2 - \Delta + m^2)\psi = (\square + m^2)\psi = 0, \quad (2.42)$$

as required by (1.2).

In a similar fashion, for the conjugate bispinor (2.15) one can obtain

$$i\partial_\nu\bar{\psi}(\Sigma^{\mu\nu} - g^{\mu\nu}) = m\bar{\psi}\gamma^\mu \quad (2.43)$$

and our equations (2.36) and (2.43) results in the following balance relation:

$$i\partial_\nu(\bar{\psi}\Sigma^{\mu\nu}\psi) + i[\bar{\psi}\partial^\mu\psi - (\partial^\mu\bar{\psi})\psi] = 2m\bar{\psi}\gamma^\mu\psi, \quad (2.44)$$

which, in turn, implies a familiar conservation law, $\partial_\mu j^\mu(x) = 0$, for the four-current, $j^\mu(x) = \bar{\psi}(x)\gamma^\mu\psi(x)$, in view of $\partial_\mu\partial_\nu(\bar{\psi}\Sigma^{\mu\nu}\psi) \equiv 0$ and $\partial_\mu[\bar{\psi}\partial^\mu\psi - (\partial^\mu\bar{\psi})\psi] \equiv 0$. (The latter equation gives a differential balance condition on its own.)

The following identity,

$$i\partial_\nu(\bar{\psi}\Sigma^{\mu\nu}\gamma^\lambda\psi) = 2i\bar{\psi}\gamma^\mu\partial^\lambda\psi - i[\bar{\psi}\gamma^\lambda\partial^\mu\psi - (\partial^\mu\bar{\psi})\gamma^\lambda\psi], \quad (2.45)$$

can be obtained with the help of (2.36), (2.43) and (2.23). The latter shows how the difference between two forms of the energy-momentum tensor (2.33) and (2.34) can be written as the four-divergence of a given tensor. Moreover, one can write

$$T^{\mu\nu} := i\bar{\psi}\gamma^\mu\partial^\nu\psi = mg^{\mu\nu}\bar{\psi}\psi + m\bar{\psi}\Sigma^{\mu\nu}\psi - i\bar{\psi}\gamma^\mu\Sigma^{\nu\lambda}\partial_\lambda\psi \quad (2.46)$$

and

$$\begin{aligned} T^{\mu\nu} &:= \frac{i}{2}[\bar{\psi}\gamma^\mu(\partial^\nu\psi) - (\partial^\nu\bar{\psi})\gamma^\mu\psi] \\ &= mg^{\mu\nu}\bar{\psi}\psi - \frac{i}{2}[\bar{\psi}\gamma^\mu\Sigma^{\nu\lambda}\partial_\lambda\psi + (\partial_\lambda\bar{\psi})\Sigma^{\nu\lambda}\gamma^\mu\psi] \end{aligned} \quad (2.47)$$

in view of (2.36) and (2.43).

2.5. Covariance and transformation of generators. The relativistic invariance of the Dirac equation is a fundamental consequence of (2.4)–(2.6); see for example, [6], [9], [27], [43], [59] and [58]. Covariance of system (2.36) can be derived, in a similar fashion, by invoking (2.8). The details are left to the reader (see also section 5.3).

It is worth noting that from the four-tensor character of $\Sigma^{\mu\nu}$, in (2.8), follow the transformation laws for the generators of rotations and boosts. Let us assume that the velocity vector \mathbf{v} , for going over to a moving frame of reference, has the direction of one of the coordinate axes, say $\{\mathbf{e}_a\}_{a=1,2,3}$. Consider also “orthogonal decompositions”, $\Sigma = \Sigma_{\parallel} + \Sigma_{\perp}$ and $\alpha = \alpha_{\parallel} + \alpha_{\perp}$, in the corresponding parallel and perpendicular directions, respectively. Then, under the Lorentz transformation,

$$S_{\Lambda}^{-1} \Sigma_{\parallel} S_{\Lambda} = \Sigma_{\parallel}, \quad S_{\Lambda}^{-1} \alpha_{\parallel} S_{\Lambda} = \alpha_{\parallel} \quad (2.48)$$

and

$$S_{\Lambda}^{-1} \Sigma_{\perp} S_{\Lambda} = \frac{\Sigma_{\perp} - i(\mathbf{v} \times \alpha)}{\sqrt{1 - v^2}}, \quad S_{\Lambda}^{-1} \alpha_{\perp} S_{\Lambda} = \frac{\alpha_{\perp} - i(\mathbf{v} \times \Sigma)}{\sqrt{1 - v^2}}, \quad (2.49)$$

by analogy with the transformations of electromagnetic fields in classical electrodynamics [32], [33], [59], [61].

The corresponding invariants [43] are given by

$$I_1 = \Sigma_{\mu\nu} \Sigma^{\mu\nu} = -2(\alpha^2 + \Sigma^2) = -12 \quad (2.50)$$

and

$$I_2 = e_{\mu\nu\sigma\tau} \Sigma^{\mu\nu} \Sigma^{\sigma\tau} = -8i\alpha \cdot \Sigma = -2i\Sigma^{\mu\nu} \Sigma_{\mu\nu} \gamma_5 = -2iI_1 \gamma_5 = 24i\gamma_5, \quad (2.51)$$

in view of the first identity (2.14). The invariant nature of the helicity and relativistic rotation of the particle spin [43] can be naturally explained from these transformations.

Example. If $\mathbf{v} = v\mathbf{e}_1$, we set $\Sigma_{\parallel} = \Sigma_1$, $\Sigma_{\perp} = \{\Sigma_2, \Sigma_3\}$ and $\alpha_{\parallel} = \alpha_1$, $\alpha_{\perp} = \{\alpha_2, \alpha_3\}$ for the boost S_L given by (2.17). By the transformation law (2.48), one gets $S^{-1}\Sigma_1 S = \Sigma_1$, $S^{-1}\alpha_1 S = \alpha_1$, which is evident. In view of (2.49), the following matrix identities,

$$S^{-1}\Sigma_2 S = \frac{\Sigma_2 + iv\alpha_3}{\sqrt{1 - v^2}}, \quad S^{-1}\Sigma_3 S = \frac{\Sigma_3 - iv\alpha_2}{\sqrt{1 - v^2}} \quad (2.52)$$

and

$$S^{-1}\alpha_2 S = \frac{\alpha_2 + iv\Sigma_3}{\sqrt{1 - v^2}}, \quad S^{-1}\alpha_3 S = \frac{\alpha_3 - iv\Sigma_2}{\sqrt{1 - v^2}}, \quad (2.53)$$

hold. Indeed, in the first relation,

$$\begin{aligned} S^{-1}\Sigma_2 S &= \left(\cosh \frac{\vartheta}{2} + \alpha_1 \sinh \frac{\vartheta}{2} \right) \Sigma_2 \left(\cosh \frac{\vartheta}{2} - \alpha_1 \sinh \frac{\vartheta}{2} \right) \\ &= \Sigma_2 \cosh^2 \frac{\vartheta}{2} + (\alpha_1 \Sigma_2 - \Sigma_2 \alpha_1) \cosh \frac{\vartheta}{2} \sinh \frac{\vartheta}{2} - \alpha_1 \Sigma_2 \alpha_1 \sinh^2 \frac{\vartheta}{2} \\ &= \Sigma_2 \cosh \vartheta + i\alpha_3 \sinh \vartheta = \frac{\Sigma_2 + iv\alpha_3}{\sqrt{1 - v^2}}, \end{aligned}$$

provided $\tanh \vartheta = v$. Verifications of the remaining identities are similar.

2.6. Hamiltonian and energy balance. The energy-momentum (density) four-vector is given by $T^{0\nu}$ and, in view of (2.46) and (2.37), one gets

$$\begin{aligned} T^{00} &= i\bar{\psi}\gamma^0\partial^0\psi = m\bar{\psi}\psi - i\bar{\psi}\gamma^0\Sigma^{0p}\partial_p\psi \\ &= \psi^\dagger(-i\boldsymbol{\alpha}\cdot\nabla + m\beta)\psi = \psi^\dagger H\psi. \end{aligned} \quad (2.54)$$

Here,

$$i\partial_0\psi = H\psi, \quad H = -i\boldsymbol{\alpha}\cdot\nabla + m\beta, \quad (2.55)$$

which presents a familiar Hamiltonian form of the Dirac equation. The differential balance equation take the form

$$\partial_0(\psi^\dagger H\psi) + \text{div}(\psi^\dagger \boldsymbol{\alpha} H\psi) = 0, \quad (2.56)$$

where $\boldsymbol{\alpha}H = \nabla \times \boldsymbol{\Sigma} - i\nabla - m\boldsymbol{\gamma}$ in view of (2.39).

2.7. The Pauli-Lubański vector and Dirac's equation for a free particle. One can easily see that equation (2.36) is related to the Pauli-Lubański vector in view of the dual identities (2.14). Indeed,

$$\Sigma^{\mu\nu} = \frac{i}{2}e^{\mu\nu\sigma\tau}(\gamma_5\Sigma_{\sigma\tau}), \quad \gamma_5\Sigma_{\sigma\tau} = \Sigma_{\sigma\tau}\gamma_5, \quad (2.57)$$

and

$$\frac{i}{2}e^{\mu\nu\sigma\tau}\gamma_5\Sigma_{\sigma\tau}p_\nu\psi = -g^{\mu\nu}(p_\nu - m\gamma_\nu)\psi.$$

By “index manipulations”,

$$\begin{aligned} \frac{1}{2}e_{\mu\nu\sigma\tau}(i\partial^\nu)(-i\Sigma^{\sigma\tau})\psi &= \frac{1}{2}e_{\mu\nu\sigma\tau}\partial^\nu\Sigma^{\sigma\tau}\psi \\ &= \gamma_5(i\partial_\mu - m\gamma_\mu)\psi = (i\partial_\mu + m\gamma_\mu)\gamma_5\psi \end{aligned} \quad (2.58)$$

with the help of familiar properties of the gamma matrices, namely, $\gamma_5^2 = I$ and $\gamma_5\gamma_\mu = -\gamma_\mu\gamma_5$. As a result, we arrive at the following equations,

$$\frac{1}{2}e_{\mu\nu\sigma\tau}\partial^\nu(\gamma^\sigma\gamma^\tau\psi) = (i\partial_\mu + m\gamma_\mu)\gamma_5\psi, \quad (2.59)$$

with summation over any two repeated indices.

The latter equation can also be obtained in view of (2.25), by letting $p_\mu = i\partial_\mu$ in operator relation (1.4). Once again, our goal is to emphasize that both overdetermined systems (2.36) and (2.59), which are related to the Pauli-Lubański vector, are equivalent to Dirac's equation in vacuum (2.6).

In components, by (2.59), for the rotations $\boldsymbol{\Sigma}$ and boosts $\boldsymbol{\alpha}$ the following standard equations hold

$$i(\nabla \cdot \boldsymbol{\Sigma})\psi = (i\partial_0 + m\gamma_0)\gamma_5\psi \quad (2.60)$$

and

$$i\partial_0\boldsymbol{\Sigma}\psi - (\nabla \times \boldsymbol{\alpha})\psi = (i\nabla - m\boldsymbol{\gamma})\gamma_5\psi, \quad (2.61)$$

respectively. Once again, this system is overdetermined and by a proper matrix multiplication, each of the four equations (2.60)–(2.61) can be reduced to a single Dirac's equation. We leave the details to the reader.

2.8. Relativistic definition of spin for Dirac particles. In view of (1.4) and (1.2), one gets

$$\begin{aligned} 4w_\mu w^\mu &= (p_\mu + m\gamma_\mu) \gamma_5 (p^\mu + m\gamma^\mu) \gamma_5 \\ &= (p_\mu + m\gamma_\mu) \gamma_5^2 (p^\mu - m\gamma^\mu) \\ &= p_\mu p^\mu - m^2 \gamma_\mu \gamma^\mu = -3m^2, \end{aligned}$$

where $s(s+1) = 3/4$, or $s = 1/2$, in covariant form.

On the other hand, introducing the familiar generators $\mathbf{M} = (i/2) \boldsymbol{\Sigma}$ and $\mathbf{N} = (1/2) \boldsymbol{\alpha}$ for the rotations and boosts, respectively, one gets

$$\mathbf{N}^2 = -\mathbf{M}^2 = 3/4, \quad \mathbf{M} \cdot \mathbf{N} = i(3/4) \gamma^5. \quad (2.62)$$

In view of $\boldsymbol{\Sigma} \gamma^5 = \gamma^5 \boldsymbol{\Sigma} = \boldsymbol{\alpha}$, in the complex space of the bispinors under consideration, we arrive at

$$\mathbf{M} \pm i \mathbf{N} = \frac{i}{2} (1 \mp \gamma_5) \boldsymbol{\Sigma} \quad (2.63)$$

and the Casimir operators of the proper orthochronous Lorentz group are given by $(\mathbf{M} \pm i \mathbf{N})^2 / 4 = -3(1 \mp \gamma_5)^2 / 16 = -3(1 \mp \gamma_5) / 8$. Here, $\mathbf{M}^2 = -s(s+1) = -3/4$, which implies, once again, that the spin is equal to $1/2$ (we have chosen real-valued boost generators; see also [2], [4], [5], [10], [23], [54], [58] and [66] for more details on the Lorentz group representations).

Miscellaneous. In addition,

$$\frac{1}{2} e_{\mu\nu\sigma\tau} \partial^\nu (\gamma^\mu \gamma^\sigma \gamma^\tau \psi) = 3i \gamma^5 \gamma_\nu \partial^\nu \psi = 3m \gamma^5 \psi, \quad (2.64)$$

in view of familiar relations:

$$\gamma^5 = \frac{i}{4!} e_{\mu\nu\sigma\tau} \gamma^\mu \gamma^\nu \gamma^\sigma \gamma^\tau, \quad \gamma^5 \gamma_\mu = \frac{i}{3!} e_{\mu\nu\sigma\tau} \gamma^\nu \gamma^\sigma \gamma^\tau \quad (2.65)$$

(see, for example, [43]).

2.9. Massless limit. Under the transformation,

$$\begin{aligned} \gamma^\mu \rightarrow \gamma'^\mu &= U \gamma^\mu U^{-1}, \quad \psi \rightarrow \psi' = \begin{pmatrix} \phi \\ \chi \end{pmatrix} = U \psi, \\ U &= \frac{1}{\sqrt{2}} \begin{pmatrix} I & I \\ I & -I \end{pmatrix} = U^{-1}, \end{aligned} \quad (2.66)$$

Dirac's equation (2.6) takes a familiar block form,

$$i \begin{pmatrix} 0 & \partial_0 - \boldsymbol{\sigma} \cdot \nabla \\ \partial_0 + \boldsymbol{\sigma} \cdot \nabla & 0 \end{pmatrix} \begin{pmatrix} \phi \\ \chi \end{pmatrix} = m \begin{pmatrix} \phi \\ \chi \end{pmatrix} \quad (2.67)$$

(see, for example, [16], [43], and [50]). As $m \rightarrow 0$, this system decouples,

$$\partial_0 \phi + (\boldsymbol{\sigma} \cdot \nabla) \phi = 0, \quad \partial_0 \chi - (\boldsymbol{\sigma} \cdot \nabla) \chi = 0, \quad (2.68)$$

resulting in Weyl's two-component equations for massless neutrinos.

3. WEYL EQUATION FOR MASSLESS NEUTRINOS

The complex unimodular matrix group $SL(2, \mathbb{C})$ is a two-fold universal covering group of the Lorentz group $SO(1, 3)$ (see, for example, [5], [10], [23], [54]). In this section, we shall use this connection in order to analyze the two-component spinor field associated with Weyl's equation.

3.1. Rotations, boosts, and their generators. Let us consider the fundamental representation of $SL(2, \mathbb{C})$, namely, we take a spinor field,

$$\phi(x) = \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix} \in \mathbb{C}^2, \quad (3.1)$$

and postulate the transformation law under the proper orthochronous Lorentz group as follows

$$\phi'(x') = S_\Lambda \phi(x), \quad x' = \Lambda x, \quad S_\Lambda = \exp\left(\frac{1}{4}\theta_{\mu\nu}\Sigma^{\mu\nu}\right). \quad (3.2)$$

Explicitly, these transformations include rotations,

$$S_R = e^{i\theta(\mathbf{n}\cdot\boldsymbol{\sigma})/2} = \cos\frac{\theta}{2} + i(\mathbf{n}\cdot\boldsymbol{\sigma})\sin\frac{\theta}{2}, \quad (\mathbf{n}\cdot\boldsymbol{\sigma})^2 = I \quad (3.3)$$

about the coordinate axes $\mathbf{n} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$, and boosts,

$$S_L = e^{-\vartheta(\mathbf{n}\cdot\boldsymbol{\sigma})/2} = \cosh\frac{\vartheta}{2} - (\mathbf{n}\cdot\boldsymbol{\sigma})\sinh\frac{\vartheta}{2}, \quad \mathbf{n} = \frac{\mathbf{v}}{v} \quad (3.4)$$

in the directions $\mathbf{n} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$, respectively, when the familiar relations,

$$v = \tanh \vartheta, \quad \cosh \vartheta = \frac{1}{\sqrt{1-v^2}}, \quad \sinh \vartheta = \frac{v}{\sqrt{1-v^2}} \quad (c=1), \quad (3.5)$$

hold. (Here, $\sigma_1, \sigma_2, \sigma_3$ are the Pauli matrices with the products given by $\sigma_p\sigma_q = ie_{pqr}\sigma_r + \delta_{pq}$ and $\{\mathbf{e}_a\}_{a=1,2,3}$ is an orthonormal basis in \mathbb{R}^3 .)

The action of the generators $M^{\alpha\beta} = x^\beta\partial^\alpha - x^\alpha\partial^\beta$ takes the form

$$\begin{aligned} M^{\alpha\beta}\phi(x) &:= -\left(\frac{d}{d\theta_{\alpha\beta}}\phi'(\Lambda x)\right)\Big|_{\theta_{\alpha\beta}=0} \\ &= -\left(\frac{dS}{d\theta_{\alpha\beta}}\right)\Big|_{\theta_{\alpha\beta}=0}\phi(x) = -\frac{1}{2}\Sigma^{\alpha\beta}\phi(x) \end{aligned} \quad (3.6)$$

for the corresponding one-parameter subgroups: $S_\Lambda = \exp(\theta_{\alpha\beta}\Sigma^{\alpha\beta}/2)$ ($\alpha, \beta = 0, 1, 2, 3$ are fixed with no summation).

In block form, the generators of this spinor representation are given by⁵

$$\Sigma^{\alpha\beta} = -\Sigma^{\beta\alpha} = \begin{pmatrix} 0 & -\sigma_q \\ \sigma_p & ie_{pqr}\sigma_r \end{pmatrix} = \begin{pmatrix} 0 & -\sigma_1 & -\sigma_2 & -\sigma_3 \\ \sigma_1 & 0 & i\sigma_3 & -i\sigma_2 \\ \sigma_2 & -i\sigma_3 & 0 & i\sigma_1 \\ \sigma_3 & i\sigma_2 & -i\sigma_1 & 0 \end{pmatrix} \quad (3.7)$$

⁵Another choice of the generators in the transformation laws (3.2)–(3.4), corresponds to $\vartheta \rightarrow -\vartheta$; see, for example, [58].

and the following self-duality identity holds

$$e_{\mu\nu\sigma\tau}\Sigma^{\sigma\tau} = 2i\Sigma_{\mu\nu} = 2ig_{\mu\sigma}g_{\nu\tau}\Sigma^{\sigma\tau}, \quad (3.8)$$

where

$$\Sigma_{\alpha\beta} = g_{\alpha\mu}g_{\beta\nu}\Sigma^{\mu\nu} = \begin{pmatrix} 0 & \sigma_q \\ -\sigma_p & ie_{pqr}\sigma_r \end{pmatrix} = \begin{pmatrix} 0 & \sigma_1 & \sigma_2 & \sigma_3 \\ -\sigma_1 & 0 & i\sigma_3 & -i\sigma_2 \\ -\sigma_2 & -i\sigma_3 & 0 & i\sigma_1 \\ -\sigma_3 & i\sigma_2 & -i\sigma_1 & 0 \end{pmatrix} \quad (3.9)$$

stated here for the reader's convenience.

3.2. The Pauli-Lubański vector and Weyl's equation. Letting $\lambda = -1/2$ in (1.3), one gets

$$e_{\mu\nu\sigma\tau}(i\partial^\nu)(-iM^{\sigma\tau})\phi = -\frac{1}{2}e_{\mu\nu\sigma\tau}\partial^\nu\Sigma^{\sigma\tau}\phi = -i\partial_\mu\phi \quad (3.10)$$

by (3.6). With the help of (3.8), we finally arrive at the following overdetermined system:

$$\Sigma^{\mu\nu}\partial_\nu\phi = \partial^\mu\phi, \quad (3.11)$$

which takes the explicit form,

$$\begin{aligned} -(\boldsymbol{\sigma} \cdot \nabla)\phi &= \partial_0\phi, \\ \partial_0\boldsymbol{\sigma}\phi + i(\nabla \times \boldsymbol{\sigma})\phi &= -\nabla\phi. \end{aligned} \quad (3.12)$$

The latter vectorial equation can be obtained from the first one by matrix multiplication with the help of familiar products of the Pauli matrices. Moreover, in (1.3), only the value $\lambda = -1/2$ results in a consistent system, defining the spin as $s = |\lambda| = 1/2$.

Thus, the relativistic two-component Weyl equations for a massless particle with the spin $1/2$, namely, $\partial_0\phi + (\boldsymbol{\sigma} \cdot \nabla)\phi = 0$, can be derived from the representation theory of the Poincaré group with the aid of the Pauli-Lubański vector⁶.

3.3. Covariance. Equations (3.11) are covariant under a proper Lorentz transformation. In view of the laws (3.2) one gets

$$S_\Lambda^{-1}\Sigma^{\sigma\tau}S_\Lambda = \Lambda^\sigma{}_\mu\Lambda^\tau{}_\nu\Sigma^{\mu\nu} \quad (3.13)$$

and

$$\frac{1}{2}[\Sigma^{\alpha\beta}, \Sigma^{\gamma\delta}] = g^{\alpha\gamma}\Sigma^{\beta\delta} - g^{\alpha\delta}\Sigma^{\beta\gamma} + g^{\beta\delta}\Sigma^{\alpha\gamma} - g^{\beta\gamma}\Sigma^{\alpha\delta}. \quad (3.14)$$

(cf. Sections 5.2–5.3).

On the second hand, with the help of the Pauli-Lubański vector, from (3.10) one can get,

$$\frac{1}{2}e_{\mu\nu\sigma\tau}\partial^\nu\Sigma^{\sigma\tau}\phi' = i\partial'_\mu\phi', \quad (3.15)$$

in the new system of coordinates, when $\phi'(x') = S\phi(x)$ and $x' = \Lambda x$. Let us multiply both sides of the latter equation by $S^{-1}\Lambda^\mu{}_\lambda$ from the left and then use (3.13) together with the two convenient transformations,

$$S\partial_\lambda\phi = \Lambda^\mu{}_\lambda\partial'_\mu\phi', \quad \partial'^\nu\phi' = \Lambda^\nu{}_\kappa S\partial^\kappa\phi, \quad (3.16)$$

⁶Some authors, see for example, [55], [56], suggest that an empirical condition is required in order to quantize the value of spin for a massless particle by (1.3). As we have just demonstrated, for Weyl's equation, it is a misconception. Indeed, the full relativistic analysis automatically includes the quantization rule of the corresponding spin and helicity as a consistency condition of the overdetermined system (3.11).

which can be readily verified with the aid of

$$(\Lambda^{-1})^\lambda{}_\nu \Lambda^\nu{}_\tau = \delta^\lambda{}_\tau, \quad (\Lambda^{-1})^\lambda{}_\nu = g^{\lambda\sigma} \Lambda^\mu{}_\sigma g_{\mu\nu}. \quad (3.17)$$

As a result,

$$(\det \Lambda) e_{\lambda\kappa\rho\chi} \partial^\kappa \Sigma^{\rho\chi} \phi = i \partial_\lambda \phi, \quad (3.18)$$

in view of the following determinant identity [43]:

$$e_{\mu\nu\sigma\tau} \Lambda^\mu{}_\lambda \Lambda^\nu{}_\kappa \Lambda^\sigma{}_\rho \Lambda^\tau{}_\chi = (\det \Lambda) e_{\lambda\kappa\rho\chi}. \quad (3.19)$$

This consideration reveals the pseudo-vector character of the Pauli-Lubański operator.

Note. Weyl's equation (3.12) was originally introduced in [68] and then quickly rejected [48], being “resurrected” only after the discovery of parity violation in beta decay [53], [72]. (The experimentally detected oscillation among the different flavors of neutrinos leads us to believe that they are not massless after all [29], [40], [51].)

Let \mathbf{v} , the velocity vector of the moving frame of reference, lie along one of the coordinate axes, say $\{\mathbf{e}_a\}_{a=1,2,3}$. Also, let $\boldsymbol{\sigma} = \boldsymbol{\sigma}_\parallel + \boldsymbol{\sigma}_\perp$ be the corresponding “orthogonal decomposition” in parallel and perpendicular directions, respectively. These components transform as,

$$S_\Lambda^{-1} \boldsymbol{\sigma}_\parallel S_\Lambda = \boldsymbol{\sigma}_\parallel, \quad S_\Lambda^{-1} \boldsymbol{\sigma}_\perp S_\Lambda = \frac{\boldsymbol{\sigma}_\perp - i(\mathbf{v} \times \boldsymbol{\sigma})}{\sqrt{1-v^2}}, \quad (3.20)$$

under a Lorentz transformation, thus resembling the transformations of electromagnetic fields in classical electrodynamics [32], [33], [41], [59], [61].

Example. Let $\mathbf{v} = v\mathbf{e}_1$. Then $\boldsymbol{\sigma}_\parallel = \sigma_1$, $\boldsymbol{\sigma}_\perp = \{\sigma_2, \sigma_3\}$ and

$$S_L = e^{-(\vartheta/2)\sigma_1} = \cosh \frac{\vartheta}{2} - \sigma_1 \sinh \frac{\vartheta}{2}, \quad S_L^{-1} = e^{(\vartheta/2)\sigma_1} = \cosh \frac{\vartheta}{2} + \sigma_1 \sinh \frac{\vartheta}{2}. \quad (3.21)$$

By the transformation law (3.20), one should get $S^{-1}\sigma_1 S = \sigma_1$, which is obvious, and

$$S^{-1}\sigma_2 S = \frac{\sigma_2 + iv\sigma_3}{\sqrt{1-v^2}}, \quad S^{-1}\sigma_3 S = \frac{\sigma_3 - iv\sigma_2}{\sqrt{1-v^2}} \quad (3.22)$$

for the corresponding Lorentz boost. Let us directly verify, for instance, the first relation. Indeed,

$$\begin{aligned} S^{-1}\sigma_2 S &= \left(\cosh \frac{\vartheta}{2} + \sigma_1 \sinh \frac{\vartheta}{2} \right) \sigma_2 \left(\cosh \frac{\vartheta}{2} - \sigma_1 \sinh \frac{\vartheta}{2} \right) \\ &= \sigma_2 \cosh^2 \frac{\vartheta}{2} + (\sigma_1 \sigma_2 - \sigma_2 \sigma_1) \cosh \frac{\vartheta}{2} \sinh \frac{\vartheta}{2} - \sigma_1 \sigma_2 \sigma_1 \sinh^2 \frac{\vartheta}{2} \\ &= \sigma_2 \cosh \vartheta + i\sigma_3 \sinh \vartheta = \frac{\sigma_2 + iv\sigma_3}{\sqrt{1-v^2}}, \end{aligned}$$

provided that $\tanh \vartheta = v$. The proof of the last identity is similar.

As is well-known, under spatial rotations the set of three Pauli matrices $\boldsymbol{\sigma}$ transform as a $3D$ vector. For instance,

$$e^{-i(\theta/2)\sigma_3} \begin{pmatrix} \sigma_1 \\ \sigma_2 \\ \sigma_3 \end{pmatrix} e^{i(\theta/2)\sigma_3} = \begin{pmatrix} \cos \theta & \sin \theta & 0 \\ -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \sigma_1 \\ \sigma_2 \\ \sigma_3 \end{pmatrix}. \quad (3.23)$$

3.4. An alternative derivation. Denoting, $\sigma^\mu = (\sigma_0 = I, \sigma_1, \sigma_2, \sigma_3)$, one can rewrite Weyl's equation in a more familiar form [50], [58]:

$$\sigma^\mu \partial_\mu \phi = 0. \quad (3.24)$$

Then, under the Lorentz transformations,

$$S_\Lambda^\dagger \sigma^\mu S_\Lambda = \Lambda^\mu_\nu \sigma^\nu \quad (3.25)$$

and

$$(\Sigma^{\alpha\beta})^\dagger \sigma^\mu + \sigma^\mu \Sigma^{\alpha\beta} = 2(g^{\beta\mu} \sigma^\alpha - g^{\alpha\mu} \sigma^\beta), \quad (3.26)$$

which also implies the covariance⁷.

Examples. In particular, one can easily verify that

$$e^{-(\vartheta/2)\sigma_1} \begin{pmatrix} \sigma_0 \\ \sigma_1 \\ \sigma_2 \\ \sigma_3 \end{pmatrix} e^{-(\vartheta/2)\sigma_1} = \begin{pmatrix} \cosh \vartheta & -\sinh \vartheta & 0 & 0 \\ -\sinh \vartheta & \cosh \vartheta & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \sigma_0 \\ \sigma_1 \\ \sigma_2 \\ \sigma_3 \end{pmatrix}, \quad (3.27)$$

with $\tanh \vartheta = v$, and

$$e^{-i(\theta/2)\sigma_3} \begin{pmatrix} \sigma_0 \\ \sigma_1 \\ \sigma_2 \\ \sigma_3 \end{pmatrix} e^{i(\theta/2)\sigma_3} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \theta & \sin \theta & 0 \\ 0 & -\sin \theta & \cos \theta & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \sigma_0 \\ \sigma_1 \\ \sigma_2 \\ \sigma_3 \end{pmatrix} \quad (3.28)$$

for the corresponding boost and rotation, respectively.

4. PROCA EQUATION

The fundamental four-vector representation of the proper orthochronous Lorentz group $SO_+(1, 3)$ is related to the relativistic wave equation for a massive particle with spin 1.

4.1. Massive vector field. The relativistic equation of motion for a real or complex four-vector field $A^\mu = (A^0, \mathbf{A})$ with a positive mass $m > 0$ can be derived in a natural way with the help of the Pauli-Lubański vector. By definition,

$$\begin{aligned} w_\mu A^\alpha &= \frac{1}{2} e_{\mu\nu\sigma\tau} \partial^\nu M^{\sigma\tau} A^\alpha = \frac{1}{2} e_{\mu\nu\sigma\tau} \partial^\nu \left((m^{\sigma\tau})^\alpha_\rho A^\rho \right) \\ &= \frac{1}{2} e_{\mu\nu\sigma\tau} \partial^\nu (g^{\sigma\alpha} A^\tau - g^{\tau\alpha} A^\sigma) = -g^{\alpha\nu} e_{\mu\nu\sigma\tau} \partial^\sigma A^\tau, \end{aligned} \quad (4.1)$$

where the matrix form of the generators (2.21) has been used. The standard decomposition,

$$\partial^\mu A^\nu = \frac{1}{2} (\partial^\mu A^\nu - \partial^\nu A^\mu) + \frac{1}{2} (\partial^\mu A^\nu + \partial^\nu A^\mu),$$

followed by the dual tensor relation,

$$F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu, \quad e_{\mu\nu\sigma\tau} F^{\sigma\tau} = -2G_{\mu\nu}, \quad (4.2)$$

gives the explicit action of the Pauli-Lubański operator on the four-vector field:

$$w_\mu A^\alpha = g^{\alpha\nu} G_{\mu\nu} = g_{\mu\nu} G^{\nu\alpha}, \quad w_\mu A_\alpha = G_{\mu\alpha}. \quad (4.3)$$

⁷Relations of this spinor representation with Maxwell's equations are discussed in section 6.2.

It is worth noting that this results in a second rank four-tensor.

In a similar fashion, for the squared operator,

$$w^2 A^\alpha = w^\mu (w_\mu A^\alpha) = \frac{1}{2} e_{\mu\nu\sigma\tau} \partial^\nu (M^{\sigma\tau} G^{\mu\alpha}). \quad (4.4)$$

But, in view of (4.7) of [32],

$$M^{\sigma\tau} G^{\mu\alpha} = g^{\sigma\mu} G^{\tau\alpha} - g^{\tau\mu} G^{\sigma\alpha} + g^{\sigma\alpha} G^{\mu\tau} - g^{\tau\alpha} G^{\mu\sigma}, \quad (4.5)$$

and with the help of a companion dual tensor identity, $e^{\mu\nu\sigma\tau} G_{\sigma\tau} = 2F^{\mu\nu}$, one gets

$$w^2 A^\alpha = g^{\sigma\alpha} \partial^\nu (e_{\nu\sigma\mu\tau} G^{\mu\tau}) = 2g^{\sigma\alpha} \partial^\nu (F_{\nu\sigma}) = 2\partial_\nu F^{\nu\alpha} = -2m^2 A^\alpha$$

as a consequence of (1.2). As a result, we have arrived at the Proca equation for a vector particle with a finite mass [52],

$$\partial_\nu F^{\nu\mu} + m^2 A^\mu = 0, \quad F^{\mu\nu} = \partial^\mu A^\nu - \partial^\nu A^\mu, \quad (4.6)$$

directly from the representation theory of the Poincaré group. In view of $w^2 = -m^2 s(s+1) = -2m^2$, one concludes that the spin of the particle is equal to one.

Moreover,

$$\partial_\mu A^\mu = 0, \quad \square A^\mu = -m^2 A^\mu, \quad m > 0, \quad (4.7)$$

in view of

$$0 \equiv \partial_\mu \partial_\nu F^{\nu\mu} = -m^2 \partial_\mu A^\mu \quad (4.8)$$

and

$$\partial_\nu F^{\nu\mu} = \partial_\nu (\partial^\nu A^\mu - \partial^\mu A^\nu) = \partial_\nu \partial^\nu A^\mu = -m^2 A^\mu. \quad (4.9)$$

The massless case of the Proca equation, $m = 0$, reveals a gauge invariance. If $A_\mu \rightarrow A'_\mu = A_\mu + \partial_\mu f$, then $F'_{\mu\nu} = \partial_\mu (A_\nu + \partial_\nu f) - \partial_\nu (A_\mu + \partial_\mu f) = F_{\mu\nu}$.

4.2. An alternative “bispinor” derivation. Let us consider a second rank bispinor of the form⁸ $Q = A_\lambda \gamma^\lambda = A^\lambda \gamma_\lambda$, where $\{\gamma^\lambda\}_{\lambda=0,1,2,3}$ are the standard gamma matrices. We shall use the following transformation law for a proper Lorentz transformation,

$$Q'(x') = S_\Lambda Q(x) S_\Lambda^{-1}, \quad x' = \Lambda x, \quad (4.10)$$

where the matrix S_Λ is given by (2.7), as in the case of Dirac’s equation. Then A^λ must be a four-vector. Indeed,

$$\begin{aligned} Q' &= A_\lambda \delta_\sigma^\lambda (S \gamma^\sigma S^{-1}) = (g^{\tau\lambda} A_\lambda \Lambda^\nu_\tau) g_{\mu\nu} [\Lambda^\mu_\sigma (S \gamma^\sigma S^{-1})] = A''^\nu \gamma_\nu, \\ A'^\nu &= \Lambda^\nu_\tau A^\tau, \end{aligned} \quad (4.11)$$

in view of an “inversion” of (2.5), $\Lambda^\mu_\sigma (S \gamma^\sigma S^{-1}) = \gamma^\mu$, and the familiar property: $A_\mu B^\mu = \text{invariant}$ or

$$g_{\mu\nu} \Lambda^\mu_\sigma \Lambda^\nu_\tau g^{\tau\lambda} = \delta_\sigma^\lambda. \quad (4.12)$$

In this “bispinor representation”, the action of generators of the corresponding one-parameter subgroups is given by

$$M^{\alpha\beta} Q := - \left(\frac{d}{d\theta_{\alpha\beta}} Q'(\Lambda x) \right) \Big|_{\theta_{\alpha\beta}=0} = \frac{1}{2} (\Sigma^{\alpha\beta} Q - Q \Sigma^{\alpha\beta}). \quad (4.13)$$

⁸This is the so-called “Feynman slash” notation from quantum electrodynamics [16].

By letting $Q = \gamma^\lambda A_\lambda$, one gets:

$$M^{\alpha\beta}Q = (g^{\beta\lambda}\gamma^\alpha - g^{\alpha\lambda}\gamma^\beta) A_\lambda,$$

in view of the familiar commutator (2.23). As a result, we obtain $w_\mu Q = G_{\mu\tau}\gamma^\tau$ for the action of the Pauli-Lubański operator on the bispinor Q , which also follows from (4.3). One gets, in a similar fashion, that

$$w^2 Q = \frac{1}{2} e_{\mu\nu\sigma\tau} \partial^\nu (M^{\sigma\tau} G^{\mu\lambda}) \gamma_\lambda, \quad (4.14)$$

where equation (4.5) holds, once again, in view of the transformation law of the four-tensor $G^{\mu\lambda}$. As a result, Proca's equation follows.

4.3. Maxwell's equations vs Proca equation. For the real-valued vector potential A^μ and $m = 0$, the Proca equation (4.6) is reduced to the Maxwell equations in vacuum. Indeed, in view of the dual relation,

$$6\partial_\nu G^{\mu\nu} = -e^{\mu\nu\sigma\tau} (\partial_\nu F_{\sigma\tau} + \partial_\sigma F_{\tau\nu} + \partial_\tau F_{\nu\sigma}) = 0, \quad (4.15)$$

both pairs of Maxwell's equations can be written together in the following complex form

$$\partial_\nu Q^{\mu\nu} = 0, \quad Q^{\mu\nu} = F^{\mu\nu} - \frac{i}{2} e^{\mu\nu\sigma\tau} F_{\sigma\tau}, \quad (4.16)$$

with the help of a self-dual complex four-tensor [8], [32]:

$$2iQ^{\mu\nu} = e^{\mu\nu\sigma\tau} Q_{\sigma\tau}, \quad e_{\mu\nu\sigma\tau} Q^{\sigma\tau} = 2iQ_{\mu\nu}. \quad (4.17)$$

The corresponding overdetermined system of Maxwell's equations in vacuum, which is related to the Pauli-Lubański vector, is investigated in [32]; see equation (5.2) there and section 5.5 below. (Two different spinor forms of Maxwell's equation will be discussed in section 6.)

5. COMPLEX VECTOR FIELD

Finally, we would like to discuss the fundamental representation for the complex orthogonal group $SO(3, \mathbb{C})$ in connection with Maxwell's equations in vacuum.

5.1. Vector covariant form. As is well-known, the transformation laws of the complex electromagnetic field $\mathbf{F}(\mathbf{r}, t) = \mathbf{E} + i\mathbf{H} \in \mathbb{C}^3$ in vacuum can be written in terms of $SO(3, \mathbb{C})$ rotations. In addition to the standard rotations of the frame of reference, the Lorentz transformations are equivalent to rotations through imaginary angles thus preserving the relativistic invariant $\mathbf{F}^2 = \mathbf{E}^2 + \mathbf{H}^2 + 2i\mathbf{E} \cdot \mathbf{H}$, which can be thought of as a “complex length” of this vector [34], [41]. In these transformations, $\mathbf{F}'(x') = S_\Lambda \mathbf{F}(x)$, $x' = \Lambda x$, (or $F'_p(x') = a_{pq} F_q(x)$ for a given complex orthogonal matrix), one can choose $S_R = e^{-\omega(\mathbf{n} \cdot \mathbf{s})}$ and $S_L = e^{-iv(\mathbf{n} \cdot \mathbf{s})}$ for the rotations and boosts, respectively. Here, $\mathbf{n} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$, when $\{\mathbf{e}_a\}_{a=1,2,3}$ is an orthonormal basis in \mathbb{R}^3 , and s_1, s_2, s_3 are the real-valued spin matrices [62]:

$$s_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \quad s_2 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad s_3 = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad (5.1)$$

such that $[s_p, s_q] = s_p s_q - s_q s_p = e_{pqr} s_r$ and $s_1^2 + s_2^2 + s_3^2 = -2$. (In this representation the matrices $\mathbf{M} = \mathbf{s}$ and $\mathbf{N} = i\mathbf{s}$ obey the commutation law of the generators of the proper orthochronous Lorentz group.)

It can be directly verified that the corresponding generators,

$$\Sigma^{\alpha\beta} = -\Sigma^{\beta\alpha} = \begin{pmatrix} 0 & i s_q \\ -i s_p & e_{pqr} s_r \end{pmatrix} = \begin{pmatrix} 0 & i s_1 & i s_2 & i s_3 \\ -i s_1 & 0 & s_3 & -s_2 \\ -i s_2 & -s_3 & 0 & s_1 \\ -i s_3 & s_2 & -s_1 & 0 \end{pmatrix}, \quad (5.2)$$

form a self-dual four-tensor

$$2i\Sigma^{\mu\nu} = e^{\mu\nu\sigma\tau}\Sigma_{\sigma\tau}, \quad e_{\mu\nu\sigma\tau}\Sigma^{\sigma\tau} = 2i\Sigma_{\mu\nu}. \quad (5.3)$$

As a result, in this realization, $M^{\alpha\beta}\mathbf{F} = m^{\alpha\beta}\mathbf{F}$ and, in view of (1.3) and (5.3), we arrive at the set of overdetermined equations:

$$(\Sigma^{\mu\nu} + g^{\mu\nu})\partial_\nu\mathbf{F} = 0, \quad (5.4)$$

where

$$S_\Lambda^{-1}\Sigma^{\mu\nu}S_\Lambda = \Lambda^\mu_\sigma\Lambda^\nu_\tau\Sigma^{\sigma\tau}, \quad x'^\mu = \Lambda^\mu_\sigma x^\sigma, \quad \mathbf{F}'(x') = S_\Lambda\mathbf{F}(x) \quad (5.5)$$

under a proper Lorentz transformation. Once again, the latter equations, that, as we shall see later, determine the complete dynamics of the electromagnetic field in vacuum, are obtained here by a pure group-theoretical consideration up to an undetermined sign of the fixed constant in (1.3).

5.2. Commutators. For a one-parameter transformation in the plane (α, β) , when

$$\Lambda(\theta_{\alpha\beta}) = \exp(-\theta_{\alpha\beta} m^{\alpha\beta}), \quad (m^{\alpha\beta})^\mu_\nu = g^{\alpha\mu}\delta^\beta_\nu - g^{\beta\mu}\delta^\alpha_\nu \quad (5.6)$$

and

$$S_\Lambda = \exp(-\theta_{\alpha\beta}\Sigma^{\alpha\beta}), \quad \theta_{\alpha\beta} = -\theta_{\beta\alpha} \quad (5.7)$$

($\alpha, \beta = 0, 1, 2, 3$ are fixed), one gets

$$e^{\theta_{\alpha\beta}\Sigma^{\alpha\beta}}\Sigma^{\mu\nu}e^{-\theta_{\alpha\beta}\Sigma^{\alpha\beta}} = \Lambda^\mu_\sigma(\theta_{\alpha\beta})\Lambda^\nu_\tau(\theta_{\alpha\beta})\Sigma^{\sigma\tau}. \quad (5.8)$$

The differentiation $(d/d\theta_{\alpha\beta})|_{\theta_{\alpha\beta}=0}$, results in a familiar law,

$$\begin{aligned} [\Sigma^{\alpha\beta}, \Sigma^{\mu\nu}] &:= \Sigma^{\alpha\beta}\Sigma^{\mu\nu} - \Sigma^{\mu\nu}\Sigma^{\alpha\beta} \\ &= g^{\alpha\nu}\Sigma^{\beta\mu} - g^{\beta\nu}\Sigma^{\alpha\mu} + g^{\alpha\mu}\Sigma^{\beta\nu} - g^{\beta\mu}\Sigma^{\alpha\nu}, \end{aligned} \quad (5.9)$$

which can be directly verified in components with the help of commutators of the spin matrices (5.1) that form the four-tensor (5.2).

5.3. Lorentz invariance. With the help of (5.4)–(5.5), under a Lorentz transformation,

$$S^{-1}[(\Sigma'^{\mu\nu} + g'^{\mu\nu})\partial'_\nu\mathbf{F}'(x')] = 0, \quad (5.10)$$

where $\Sigma'^{\mu\nu} = \Sigma^{\mu\nu} = \text{inv}$ and $g'^{\mu\nu} = g^{\mu\nu} = \text{inv}$. Then

$$\begin{aligned} 0 &= [S^{-1}(\Sigma^{\mu\nu} + g^{\mu\nu})S]S^{-1}\partial'_\nu\mathbf{F}'(x') \\ &= \Lambda^\mu_\sigma(\Sigma^{\sigma\tau} + g^{\sigma\tau})S^{-1}[\Lambda^\nu_\tau\partial'_\nu\mathbf{F}'(x')] \\ &= \Lambda^\mu_\sigma(\Sigma^{\sigma\tau} + g^{\sigma\tau})S^{-1}S\partial_\tau\mathbf{F}(x) \end{aligned} \quad (5.11)$$

by $\Lambda^\nu_\tau\partial'_\nu\mathbf{F}'(x') = \partial_\tau\mathbf{F}(x)$. Thus, our equation $(\Sigma^{\mu\nu} + g^{\mu\nu})\partial_\nu\mathbf{F}(x) = 0$ preserves its covariant form for all real and complex rotations $S \in SO(3, \mathbb{C})$.

5.4. Vector covariant form vs traditional form of Maxwell's equations. A vector form of (5.4) is given by

$$(\nabla \cdot \mathbf{s}) \mathbf{F} = i\partial_0 \mathbf{F} \quad (5.12)$$

and

$$\partial_0 \mathbf{s} \mathbf{F} + i(\nabla \times \mathbf{s}) \mathbf{F} = i\nabla \mathbf{F}. \quad (5.13)$$

As it has been pointed out in [32], equation (5.12) implies the complex Maxwell equation, $\text{curl } \mathbf{F} = i\partial_0 \mathbf{F}$. The first component of (5.13) is given by $\partial_0 s_1 \mathbf{F} + i(\partial_2 s_3 - \partial_3 s_2) \mathbf{F} = i\partial_1 \mathbf{F}$, or

$$\begin{aligned} \partial_0 \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} F_1 \\ F_2 \\ F_3 \end{pmatrix} + i\partial_2 \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \begin{pmatrix} F_1 \\ F_2 \\ F_3 \end{pmatrix} \\ - i\partial_3 \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix} \begin{pmatrix} F_1 \\ F_2 \\ F_3 \end{pmatrix} = i\partial_1 \begin{pmatrix} F_1 \\ F_2 \\ F_3 \end{pmatrix} \end{aligned}$$

and

$$\begin{aligned} -i\partial_2 F_2 - i\partial_3 F_3 &= i\partial_1 F_1, \\ -\partial_0 F_3 + i\partial_2 F_1 &= i\partial_1 F_2, \\ \partial_0 F_2 + i\partial_3 F_1 &= i\partial_1 F_3. \end{aligned}$$

As a result, $\text{div } \mathbf{F} = 0$ and $(\text{curl } \mathbf{F})_{2,3} = i\partial_0 (\mathbf{F})_{2,3}$. (Cyclic permutations of the spatial indices cover the two remaining cases.)

5.5. An alternative form of Maxwell's equations. Complex covariant form of Maxwell's equations, which was introduced in [35] (see also [32]), can be presented as follows

$$(\partial_0, \partial_1, \partial_2, \partial_3) \begin{pmatrix} 0 & -F_1 & -F_2 & -F_3 \\ F_1 & 0 & iF_3 & -iF_2 \\ F_2 & -iF_3 & 0 & iF_1 \\ F_3 & iF_2 & -iF_1 & 0 \end{pmatrix} = (j_0, j_1, j_2, j_3), \quad (5.14)$$

or $\partial Q = j$, by matrix multiplication. Here, $\partial = (\partial_0, \partial_1, \partial_2, \partial_3)$, $j = (j_0, j_1, j_2, j_3)$, and $\mathbf{J} = (J_1, J_2, J_3)$ such that

$$\begin{aligned} Q &= \mathbf{F} \cdot \mathbf{J} = F_1 \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \end{pmatrix} \\ &+ F_2 \begin{pmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & -i \\ 1 & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{pmatrix} + F_3 \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & i & 0 \\ 0 & -i & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}. \end{aligned} \quad (5.15)$$

These matrices are transformed as a dual complex vector in \mathbb{C}^3 ,

$$\Lambda J_p \Lambda^T = a_{qp} J_q, \quad (5.16)$$

under a proper Lorentz transformation. In infinitesimal form,

$$m^{\alpha\beta} \mathbf{J} + \mathbf{J} (m^{\alpha\beta})^T = (\Sigma^{\alpha\beta})^T \mathbf{J}, \quad (5.17)$$

which gives an alternative representation of the group $SO(3, \mathbb{C})$ in a subspace of complex 4×4 matrices. (Details are left to the reader.)

As a result, $\Lambda Q \Lambda^T = Q'$ and, in “new” coordinates, $\partial' Q' = j'$, provided that $\partial' = \partial(\Lambda^{-1})$ and $j' = j\Lambda^T$.

6. ON SPINOR FORMS OF MAXWELL’S EQUATIONS

In conclusion, the complex matrix group $SL(2, \mathbb{C})$ has a representation of the proper orthochronous Lorentz group $SO_+(1, 3)$ by the second rank spinors.

6.1. Spinor covariant form. The complex electromagnetic field in vacuum, $\mathbf{F} = \mathbf{E} + i\mathbf{H}$, can also be written in a familiar form of the following 2×2 matrix:

$$Q = \boldsymbol{\sigma} \cdot \mathbf{F} = \sigma_1 F_1 + \sigma_2 F_2 + \sigma_3 F_3 = \begin{pmatrix} F_3 & F_1 - iF_2 \\ F_1 + iF_2 & -F_3 \end{pmatrix}, \quad (6.1)$$

where $\sigma_1, \sigma_2, \sigma_3$ are the standard Pauli matrices. The corresponding transformation law,

$$Q'(x') = S_\Lambda Q(x) S_\Lambda^{-1}, \quad x' = \Lambda x, \quad S_\Lambda = \exp\left(\frac{1}{4}\theta_{\mu\nu}\Sigma^{\mu\nu}\right), \quad (6.2)$$

under a proper Lorentz transformation preserves two invariants $\text{tr}Q = 0$ and $\det Q = -\mathbf{F}^2$. Here, $S_\Lambda \sigma_p S_\Lambda^{-1} = a_{pq} \sigma_q$ and $F'_p(x') = a_{pq} F_q(x)$, with $a_{qr} a_{qs} = \delta_{rs}$ for a given complex orthogonal 3×3 matrix (see section 5.1).

In this “spinor” representation, one gets

$$M^{\alpha\beta} Q = - \left(\frac{d}{d\theta_{\alpha\beta}} Q'(\Lambda x) \right) \Big|_{\theta_{\alpha\beta}=0} = -\frac{1}{2} (\Sigma^{\alpha\beta} Q - Q \Sigma^{\alpha\beta}) \quad (6.3)$$

for generators of the one-parameter subgroups. Here, as in the case of Weyl’s equation, the matrices $\Sigma^{\alpha\beta}$ are given by (3.7), but now, in view of (6.3), equation (1.3) with $\lambda = -1$ takes the form,

$$\frac{1}{2} (\Sigma_{\mu\nu} \partial^\nu Q - \partial^\nu Q \Sigma_{\mu\nu}) = \partial_\mu Q, \quad (6.4)$$

when the self-duality property (3.8) is applied.

Equations (6.4), obtained with the aid of the Pauli-Lubański vector, are equivalent to the system of complex Maxwell equations in vacuum,

$$\text{div } \mathbf{F} = 0, \quad \text{curl } \mathbf{F} = i\partial_0 \mathbf{F}. \quad (6.5)$$

Indeed, when $\mu = 0$, with the help of (3.9) one gets

$$-\frac{1}{2} (\sigma_p \partial_p Q - \partial_p Q \sigma_p) = \partial_0 Q,$$

or

$$-(\sigma_p \sigma_q - \sigma_q \sigma_p) \partial_p F_q = 2\partial_0 (\sigma_r F_r),$$

which gives the second complex Maxwell equation (6.5) in view of the commutation relation, $[\sigma_p, \sigma_q] = 2ie_{pqr}\sigma_r$.

When $\mu = p = 1, 2, 3$, in a similar fashion,

$$2\partial_p Q = \partial_0 Q \sigma_p - \sigma_p \partial_0 Q + ie_{pqr} (\partial_q Q \sigma_r - \sigma_r \partial_q Q), \quad (6.6)$$

and letting $Q = F_s \sigma_s$, we obtain

$$2\partial_p (F_s \sigma_s) = (\sigma_s \sigma_p - \sigma_p \sigma_s) \partial_0 F_s + ie_{pqr} (\sigma_s \sigma_r - \sigma_r \sigma_s) \partial_q F_s.$$

Evaluation of the commutators,

$$\sigma_s \partial_p F_s = ie_{spl} \sigma_l \partial_0 F_s + e_{pqr} e_{slr} \sigma_l \partial_q F_s, \quad (6.7)$$

and a familiar identity,

$$e_{pqr} e_{slr} = \begin{vmatrix} \delta_{ps} & \delta_{pl} \\ \delta_{qs} & \delta_{ql} \end{vmatrix} = \delta_{ps} \delta_{ql} - \delta_{pl} \delta_{qs}, \quad (6.8)$$

result in the system of Maxwell's equations (6.5).

6.2. Traditional spinor form of Maxwell's equations. Equations (6.4), that are obtained here with the help of the Pauli-Lubański vector, give an alternative (vacuum) version of the spinor form of Maxwell's equations,

$$(\partial_0 + \boldsymbol{\sigma} \cdot \nabla) (\boldsymbol{\sigma} \cdot \mathbf{F}) = j_0 + \boldsymbol{\sigma} \cdot \mathbf{j}, \quad (6.9)$$

or, explicitly,

$$\begin{pmatrix} \partial_0 + \partial_3 & \partial_1 - i\partial_2 \\ \partial_1 + i\partial_2 & \partial_0 - \partial_3 \end{pmatrix} \begin{pmatrix} F_3 & F_1 - iF_2 \\ F_1 + iF_2 & -F_3 \end{pmatrix} = \begin{pmatrix} j_0 + j_3 & j_1 - ij_2 \\ j_1 + ij_2 & j_0 - j_3 \end{pmatrix}, \quad (6.10)$$

which was originally established in [35] (see also [54]).

Let $Q = \boldsymbol{\sigma} \cdot \mathbf{F}$ and $\mathcal{D} = \sigma^\mu \partial_\mu$, $\mathcal{J} = \sigma^\mu j_\mu$, when $\mathcal{D}Q = \mathcal{J}$. Then $S_\Lambda Q S_{\Lambda^{-1}} = Q'$ and, in view of (3.17) and (3.25), one gets

$$S_{\Lambda^{-1}}^\dagger \mathcal{D} S_{\Lambda^{-1}} = \mathcal{D}', \quad S_{\Lambda^{-1}}^\dagger \mathcal{J} S_{\Lambda^{-1}} = \mathcal{J}' \quad (6.11)$$

under a proper Lorentz transformation. In “new” coordinates, equation (6.10) should take a compact form, $\mathcal{D}'Q' = \mathcal{J}'$. Thus

$$\left(S_{\Lambda^{-1}}^\dagger \mathcal{D} S_{\Lambda^{-1}} \right) S_\Lambda Q S_{\Lambda^{-1}} = S_{\Lambda^{-1}}^\dagger \mathcal{J} S_{\Lambda^{-1}}, \quad (6.12)$$

or, $S_{\Lambda^{-1}}^\dagger (\mathcal{D}Q = \mathcal{J}) S_{\Lambda^{-1}}$, as a short proof of the covariance of Maxwell's equations.

7. MASSIVE SYMMETRIC FOUR-TENSOR FIELD

7.1. Group-theoretical derivation. The relativistic wave equation for a massive particle of spin two, described by a real or complex symmetric four-tensor field $A^{\mu\nu}(x) = A^{\nu\mu}(x)$ (see [17], [19], [20]), can be obtained in a way that is similar to our study of the Proca equation in section 4. Once again,

$$\begin{aligned} w_\mu A^{\alpha\beta} &= \frac{1}{2} e_{\mu\nu\sigma\tau} \partial^\nu (M^{\sigma\tau} A^{\alpha\beta}) \\ &= -g^{\alpha\nu} e_{\mu\nu\sigma\tau} \partial^\sigma A^{\tau\beta} - g^{\beta\nu} e_{\mu\nu\sigma\tau} \partial^\sigma A^{\tau\alpha} \end{aligned} \quad (7.1)$$

and

$$e_{\mu\nu\sigma\tau} \partial^\sigma A^{\tau\beta} = \frac{1}{2} e_{\mu\nu\sigma\tau} F^{\sigma\tau\beta} = -G_{\mu\nu}{}^\beta, \quad (7.2)$$

where, by definition, $F^{\sigma\tau\beta} = \partial^\sigma A^{\tau\beta} - \partial^\tau A^{\sigma\beta}$ and $e_{\mu\nu\sigma\tau} F^{\sigma\tau\beta} = -2G_{\mu\nu}{}^\beta$. Thus

$$w_\mu A^{\alpha\beta} = g^{\alpha\nu} G_{\mu\nu}{}^\beta + g^{\beta\nu} G_{\mu\nu}{}^\alpha, \quad (7.3)$$

or

$$w^\mu A^{\alpha\beta} = G^{\mu\alpha\beta} + G^{\mu\beta\alpha}, \quad w_\mu A_{\alpha\beta} = G_{\mu\alpha\beta} + G_{\mu\beta\alpha} \quad (7.4)$$

and

$$w^2 A^{\alpha\beta} = w_\mu (w^\mu A^{\alpha\beta}) = w_\mu G^{\mu\alpha\beta} + w_\mu G^{\mu\beta\alpha}. \quad (7.5)$$

In a similar fashion, one can show that

$$\begin{aligned} w_\mu G^{\mu\alpha\beta} &= 2\partial_\nu F^{\nu\alpha\beta} + \partial_\nu F^{\nu\beta\alpha} \\ &\quad + \partial^\alpha (F^{\beta\sigma\tau} g_{\sigma\tau}) - g^{\alpha\beta} \partial_\nu (F^{\nu\sigma\tau} g_{\sigma\tau}), \end{aligned} \quad (7.6)$$

as a result of an elementary but rather tedious four-tensor algebra calculation with the help of a familiar relation

$$e^{\mu\nu\sigma\tau} e_{\mu\kappa\lambda\rho} = - \begin{vmatrix} \delta_\kappa^\nu & \delta_\lambda^\nu & \delta_\rho^\nu \\ \delta_\kappa^\sigma & \delta_\lambda^\sigma & \delta_\rho^\sigma \\ \delta_\kappa^\tau & \delta_\lambda^\tau & \delta_\rho^\tau \end{vmatrix}. \quad (7.7)$$

For a massive spin-2 field, the second Casimir operator is equal to $w^2 A^{\alpha\beta} = -6m^2 A^{\alpha\beta}$, and we obtain

$$\begin{aligned} &\partial_\nu F^{\nu\alpha\beta} + \partial_\nu F^{\nu\beta\alpha} - \frac{2}{3} g^{\alpha\beta} \partial_\nu (F^{\nu\sigma\tau} g_{\sigma\tau}) \\ &\quad + \frac{1}{3} \partial^\alpha (F^{\beta\sigma\tau} g_{\sigma\tau}) + \frac{1}{3} \partial^\beta (F^{\alpha\sigma\tau} g_{\sigma\tau}) = -2m^2 A^{\alpha\beta}, \end{aligned} \quad (7.8)$$

where

$$\begin{aligned} F^{\nu\alpha\beta} &= \partial^\nu A^{\alpha\beta} - \partial^\alpha A^{\nu\beta} = -F^{\alpha\nu\beta}, \\ F^{\nu\sigma\tau} g_{\sigma\tau} &= \partial^\nu A - \partial_\tau A^{\nu\tau}, \quad A = g_{\sigma\tau} A^{\sigma\tau}, \end{aligned} \quad (7.9)$$

and $F^{\alpha\beta\gamma} + F^{\beta\gamma\alpha} + F^{\gamma\alpha\beta} = 0$ provided $A^{\alpha\beta} = A^{\beta\alpha}$.

In terms of potentials, one gets

$$\begin{aligned} &\partial^2 A^{\alpha\beta} - \frac{2}{3} (\partial^\alpha \partial_\nu A^{\nu\beta} + \partial^\beta \partial_\nu A^{\nu\alpha}) + \frac{1}{3} \partial^\alpha \partial^\beta A \\ &\quad - \frac{1}{3} g^{\alpha\beta} (\partial^2 A - \partial_\mu \partial_\nu A^{\mu\nu}) = -m^2 A^{\alpha\beta}, \end{aligned} \quad (7.10)$$

with $\partial_\mu \partial_\nu A^{\mu\nu} = 0$, if $m > 0$, in view of

$$0 \equiv w^2 (\partial_\mu \partial_\nu A^{\mu\nu}) = \partial_\mu \partial_\nu (w^2 A^{\mu\nu}) = -6m^2 \partial_\mu \partial_\nu A^{\mu\nu}.$$

The same results can be derived by differentiation from (7.8) or, independently, with the help of an operator identity

$$w^2 = -\frac{1}{2} \partial^2 (M^{\sigma\tau} M_{\sigma\tau}) - \partial_\mu \partial^\nu (M^{\mu\sigma} M_{\sigma\nu}).$$

Moreover, equation (7.10) is reduced to

$$2 (\partial^\alpha \partial_\nu A^{\nu\beta} + \partial^\beta \partial_\nu A^{\nu\alpha}) - \partial^\alpha \partial^\beta A + g^{\alpha\beta} \partial^2 A = 0 \quad (7.11)$$

with the help of the first Casimir operator, $\partial^2 A^{\alpha\beta} = -m^2 A^{\alpha\beta}$. Multiplication of (7.11) by $g_{\alpha\beta}$ with contraction over two repeated indices results in $\partial^2 A = 0$. Then $A = 0$, due to

$$0 = \partial^2 A = \partial^2 (g_{\mu\nu} A^{\mu\nu}) = g_{\mu\nu} \partial^2 (A^{\mu\nu}) = -m^2 A, \quad m > 0.$$

Thus, the system of wave equations for a massive spin-2 particle has the form

$$\partial^2 A^{\mu\nu} + m^2 A^{\mu\nu} = 0, \quad \partial^\alpha (\partial_\nu A^{\nu\beta}) + \partial^\beta (\partial_\nu A^{\nu\alpha}) = 0, \quad (7.12)$$

subject to $A = g_{\mu\nu} A^{\mu\nu} = \partial_\mu \partial_\nu A^{\mu\nu} = 0$. It is worth noting, once again, that we have derived these equations by using the Pauli-Lubański vector and the relativistic definition of mass and spin in terms of Casimir operators of the Poincaré group.

In the massless limit $m \rightarrow 0$, instead of (7.12), one has $\partial^2 A^{\mu\nu} = 0$ and $\partial^2 A = 0$ subject to

$$\partial^\alpha \left(\partial_\nu A^{\nu\beta} - \frac{1}{4} \partial^\beta A \right) + \partial^\beta \left(\partial_\nu A^{\nu\alpha} - \frac{1}{4} \partial^\alpha A \right) = \frac{1}{2} g^{\alpha\beta} \partial_\mu \partial_\nu A^{\mu\nu}, \quad (7.13)$$

in a similar fashion. Moreover, if $m = 0$, equation (7.8) is invariant under a familiar gauge transformation $A_{\alpha\beta} \rightarrow A'_{\alpha\beta} + \partial_\alpha f_\beta + \partial_\beta f_\alpha$ provided that $\partial^\nu (\partial_\nu f_\alpha - \partial_\alpha f_\nu) = 0$ (Maxwell's equations in vacuum).

7.2. An alternative gauge condition. In view of the following identity,

$$\begin{aligned} & \partial^\alpha \left(\partial_\nu A^{\nu\beta} - \frac{1}{4} \partial^\beta A \right) + \partial^\beta \left(\partial_\nu A^{\nu\alpha} - \frac{1}{4} \partial^\alpha A \right) \\ &= \partial^\alpha \partial_\nu A^{\nu\beta} + \partial^\beta \partial_\nu A^{\nu\alpha} - \frac{1}{2} \partial^\alpha \partial^\beta A, \end{aligned}$$

one can impose another condition,

$$4\partial_\nu A^{\mu\nu} - \partial^\mu A = 0, \quad (7.14)$$

in order to simplify (7.10):

$$\partial^2 A^{\mu\nu} - \frac{1}{4} g^{\mu\nu} \partial^2 A = -m^2 A^{\mu\nu}. \quad (7.15)$$

Moreover, by contraction,⁹

$$0 \equiv \partial^2 A - \frac{1}{4} g_{\mu\nu} g^{\mu\nu} \partial^2 A = -m^2 A, \quad A = g_{\sigma\tau} A^{\sigma\tau} = 0,$$

if $m > 0$. As a result, we obtain

$$\partial^2 A^{\mu\nu} + m^2 A^{\mu\nu} = 0, \quad \partial_\nu A^{\mu\nu} = \partial_\nu A^{\nu\mu} = 0. \quad (7.16)$$

These equations were originally introduced by Fierz and Pauli [17], [19], [20] with the help of a Lagrangian approach. It is worth noting that our equations (7.12) are necessary and sufficient with the relativistic definition of mass and spin-2 of the field in question, whereas traditional equations (7.16) give only sufficient conditions.

⁹Condition $A = 0$ is not required in the massless limit.

7.3. Fierz-Pauli vs Maxwell's equations. When $m = 0$, one gets

$$\partial_\nu F^{\mu\nu\alpha} = \partial_\nu (\partial^\mu A^{\nu\alpha} - \partial^\nu A^{\mu\alpha}) = \partial^\mu (\partial_\nu A^{\nu\alpha}) - \partial^2 A^{\mu\alpha} = 0, \quad (7.17)$$

subject to (7.16). In addition,

$$\partial_\lambda F_{\sigma\tau\alpha} + \partial_\sigma F_{\tau\lambda\alpha} + \partial_\tau F_{\lambda\sigma\alpha} = 0, \quad (7.18)$$

which follows from definition. These facts allow one to represent the massless Fierz-Pauli equations in terms of the third rank field tensor, somewhat similar to classical electrodynamics. Indeed, by analogy with Maxwell's equations, we obtain

$$\partial_\nu F^{\mu\nu\alpha} = 0, \quad \partial_\nu G^{\mu\nu\alpha} = 0 \quad (7.19)$$

in view of $2G^{\mu\nu\alpha} = -e^{\mu\nu\sigma\tau} F_{\sigma\tau}{}^\alpha$ and

$$\partial_\nu G^{\mu\nu\alpha} = -\frac{1}{6} e^{\mu\nu\sigma\tau} (\partial_\nu F_{\sigma\tau}{}^\alpha + \partial_\sigma F_{\tau\nu}{}^\alpha + \partial_\tau F_{\nu\sigma}{}^\alpha) = 0$$

(for every fixed $\alpha = 0, 1, 2, 3$).

Finally, both pairs of these equations can be combined together in the following complex form

$$\partial_\nu Q^{\mu\nu\alpha} = 0, \quad Q^{\mu\nu\alpha} = F^{\mu\nu\alpha} - \frac{i}{2} e^{\mu\nu\sigma\tau} F_{\sigma\tau}{}^\alpha, \quad (7.20)$$

with the help of a self-dual complex four-tensor:

$$2iQ^{\mu\nu\alpha} = e^{\mu\nu\sigma\tau} Q_{\sigma\tau}{}^\alpha, \quad e_{\mu\nu\sigma\tau} Q^{\sigma\tau\alpha} = 2iQ_{\mu\nu}{}^\alpha. \quad (7.21)$$

The covariant field “energy-momentum” tensor and the corresponding differential balance equation,

$$\frac{\partial}{\partial x^\nu} \left(Q_{\mu\lambda\sigma}^* Q^{\lambda\nu\sigma} + Q_{\mu\lambda\sigma} Q^{\lambda\nu\sigma*} \right) = 0, \quad (7.22)$$

can be derived in a complete analogy with complex electrodynamics [32], [33].

7.4. Fierz-Pauli vs linearized Einstein's equations. In general relativity, the linearized equations for a weak gravitational field [14], [25], namely,

$$\begin{aligned} \partial_\mu \partial^\sigma h_{\sigma\nu} + \partial_\nu \partial^\sigma h_{\sigma\mu} - \partial_{\mu\nu} h \\ - \partial^2 h_{\mu\nu} - g_{\mu\nu} (\partial_\sigma \partial_\tau h^{\sigma\tau} - \partial^2 h) = 0, \end{aligned} \quad (7.23)$$

describe small deviations from the flat Minkowski metric, $g_{\mu\nu} = \text{diag}(1, -1, -1, -1)$, on the pseudo-Riemannian manifold subject to a gauge condition

$$2\partial^\nu h_{\mu\nu} - \partial_\mu h = 0, \quad h = g^{\sigma\tau} h_{\sigma\tau} \quad (7.24)$$

(see, for example, [11], [15], [20], [21], [26], [28], [34], [42], [46], [63], [64], [65], [67], [69], [71], [73], and the references therein for more details).

Our calculations have shown that linearized Einstein's equations (7.23) do not coincide with the massless limit of the spin-2 particle wave equation (7.10). But they can be reduced to the massless case of the Fierz-Pauli equations (7.16) in view of an additional condition (7.24) on a certain solution set. In the literature, this fact is usually interpreted as spin-2 for a graviton although, from the group-theoretical point of view, this massless limit yet requires certain analysis of helicity, say similar to the one in electrodynamics [32], which will be discuss elsewhere.

8. SUMMARY

In this article, we analyze kinematics of the fundamental relativistic wave equations, in a traditional way, from the viewpoint of the representation theory of the Poincaré group. In particular, the importance of the Pauli-Lubański pseudo-vector is emphasized here not only for the covariant definition of spin and helicity of a given field but also for the derivation of the corresponding equation of motion from first principles. In this consistent group-theoretical approach, the resulting wave equations occur, in general, in certain overdetermined forms, which can be reduced to the standard ones by a matrix version of Gaussian elimination.

Although, mathematically, all representations of the Poincaré group are locally equivalent [4], their explicit realizations in conventional linear spaces of four-vectors and tensors, spinors and bispinors, etc. are quite different from the viewpoint of physics. This is why, as the reader can see in the table below, the corresponding relativistic wave equations are so different.

<i>Classical field</i>	<i>Transformation law (a law of nature)</i>	<i>Wave equation</i>
Bispinor (2.3)	$\psi'(x') = S_\Lambda \psi(x), x' = \Lambda x$; see (2.7) and (2.21)	Dirac (2.6)
Spinor (3.1)	$\psi'(x') = S_\Lambda \psi(x), x' = \Lambda x$; see (3.2) and (2.21)	Weyl (3.24)
Four-vector	$A'^\mu(x') = \Lambda^\mu_\nu A^\nu(x)$; see (2.21)	Proca (4.6)
“Feynman slash”	$Q'(x') = S_\Lambda Q(x) S_\Lambda^{-1}, x' = \Lambda x$; see (4.10)	Proca (4.6)
Four-tensor (4.16)	$Q'^{\mu\nu}(x') = \Lambda^\mu_\sigma \Lambda^\nu_\tau Q^{\sigma\tau}(x), x' = \Lambda x$; see (2.21)	Maxwell (4.16, 5.14)
Complex 3D vector	$\mathbf{F}'(x') = S_\Lambda \mathbf{F}(x), x' = \Lambda x$; see Sect. 5.1, (2.21)	Maxwell (6.5)
Complex matrix (6.1)	$Q'(x') = S_\Lambda Q(x) S_\Lambda^{-1}, x' = \Lambda x$; see (6.2)	Maxwell (6.5)
Symmetric four-tensor	$A'^{\mu\nu}(x') = \Lambda^\mu_\sigma \Lambda^\nu_\tau A^{\sigma\tau}(x), x' = \Lambda x$	Fierz-Pauli (7.12)

Acknowledgements. This research was partially supported by NSF grants DMS # 1535833 and DMS # 1151618. We are indebted to Prof. Dr. Albert Boggess, Prof. Dr. Mark P Faifman, Prof. Dr. Gerald A. Goldin, Prof. Dr. John Klauder, Prof. Dr. Margarita A. Man’ko, Prof. Dr. Vladimir I. Man’ko, Prof. Dr. Svetlana Roudenko, and Prof. Dr. Igor N. Toptygin for their valuable comments and encouragement. Comments from the referees, which have helped to improve the presentation, are much appreciated.

REFERENCES

- [1] Akhiezer A and Berestetskii V B 1965 *Quantum Electrodynamics* (New York: Interscience).
- [2] Bargmann V 1947 Irreducible unitary representations of the Lorentz group *Ann. Math.* **48** 568–640.
- [3] Bargmann V 1954 On unitary ray representations of continuous groups *Ann. Math.* **59** 1–46.
- [4] Bargmann V and Wigner E 1948 Group theoretical discussion of relativistic wave equations *Proc. Nat. Acad. Sci. U.S.A.* **34** 211–223.
- [5] Barut A O and Raczka R 1986 *Theory of Group Representations and Applications* 2nd edn (Singapore: World Scientific).
- [6] Berestetskii V B, Lifshitz E M and Pitaevskii L P 1971 *Relativistic Quantum Theory* (Oxford: Pergamon).
- [7] Bhabha H J 1945 Relativistic wave equations for the elementary particles *Rev. Mod. Phys.* **17** 200–216.
- [8] Białynicki-Birula I and Białynicki-Birula Z 1975 *Quantum Electrodynamics* (Oxford, New York: Pergamon and PWN–Polish Scientific Publishers).
- [9] Bogoliubov N N and Shirkov D V 1980 *Introduction to the Theory of Quantized Fields* 3rd edn (New York: Wiley).

- [10] Bogolubov N N, Logunov A A, Oksak A I and Todorov I T 1990 *General Principles of Quantum Field Theory* (Dordrecht: Kluwer).
- [11] Damour T 2007 General relativity today *Gravitation and Experiment - Poincaré Seminar 2006* ed T Damour, B Duplantier and V Rivasseau (Basel: Birkhäuser) pp 1–49.
- [12] Dirac P A M 1928 The quantum theory of the electron *Proc. Roy. Soc. London* **A117** 610–624.
- [13] Dirac P A M 1936 Relativistic wave equations *Proc. Roy. Soc. London* **A155** 447–459.
- [14] Einstein A 1916 Näherungsweise Integration der Feldgleichungen der Gravitation *Sitzungsber. Kön. Preuß. Akad. Wiss. zu Berlin* **1** 688–696.
- [15] Faddeev L D 1982 The energy problem in Einstein’s theory of gravitation *Phys. Usp.* **25** 130–142.
- [16] Feynman R P 1998 *The Theory of Fundamental Processes* (Cambridge: Perseus).
- [17] Fierz M 1939 Über die relativistische Theorie kräftefreier Teilchen mit beliebigem Spin *Helv. Phys. Acta* **12** 3–37.
- [18] Fierz M 1940 Über den Drehimpuls von Teilchen mit Ruhemasse null und beliebigem Spin *Helv. Phys. Acta* **13** 45–60.
- [19] Fierz M and Pauli W 1939 Über die relativistische Feldgleichungen von Teilchen mit beliebigem Spin in elektromagnetischen Feld *Helv. Phys. Acta* **12** 297–300.
- [20] Fierz M and Pauli W 1939 On relativistic wave equations for particles of arbitrary spin in an electromagnetic field *Proc. Roy. Soc. London* **A173** 211–232.
- [21] Fock V A 1965 *The Theory of Space, Time and Gravitation* (New York: Pergamon).
- [22] Gantmacher F R 2000 *The Theory of Matrices* vol 1 (Providence, Rhode Island: AMS Chelsea).
- [23] Gel’fand I M, Minlos R A and Shapiro Z Ya 1963 *Representations of the Rotation and Lorentz Groups and their Applications* (New York: Pergamon).
- [24] Greiner W 2000 *Relativistic Quantum Mechanics: Wave Equations* 3rd edn (Berlin, Heidelberg, New York: Springer).
- [25] Hilbert D 1924 Die Grundlagen der Physik (Zweite Mitteilung) *Math. Ann.* **92** 1–32.
- [26] Iorio L 2015 Editorial for the special issue 100 years of chronogeometro-dynamics: the status of the Einstein’s theory of gravitation in its centennial year *Universe* **1** 38–81.
- [27] Itzykson C and Zuber J-B 2005 *Quantum Field Theory* (New York: Dover).
- [28] Khriplovich I B 2005 *General Relativity* (New York: Springer).
- [29] Kajita T 2015 Discovery of atmospheric neutrino oscillations *Nobel Lecture*, December 8; http://www.nobelprize.org/nobel_prizes/physics/laureates/2015/kajita-lecture.html
- [30] Klyshko D. N. 1994 Quantum optics: quantum, classical, and metaphysical aspects *Phys.-Usp.* **37** 1097–1123; see also: *Annals of the New York Academy of Sciences* vol. 755 *Fundamental Problems in Quantum Theory* April 1995 pp 13–27.
- [31] Kramers H A, Belinfante F J and Lubański J K 1941 Über freie Teilchen mit nichtverschwindender Masse und beliebiger Spinquantenzahl *Physica* **8** 597–627.
- [32] Kryuchkov S I, Lanfear N A and Suslov S K 2015 The Pauli-Lubański vector, complex electrodynamics, and photon helicity *Physica Scripta* **90** 074065 (8pp)
- [33] Kryuchkov S I, Lanfear N A and Suslov S K 2015 Complex Form of Classical and Quantum Electrodynamics *in preparation*.
- [34] Landau L D and Lifshitz E M 1975 *The Classical Theory of Fields* 4th edn (Oxford: Butterworth-Heinemann).
- [35] Laporte O and Uhlenbeck G E 1931 Applications of spinor analysis to the Maxwell and Dirac equations *Phys. Rev.* **37** 1380–1397.
- [36] Lubański J K 1941 Sur le spin des particules élémentaires *Physica* **8** 44–52.
- [37] Lubański J K 1942 Sur la théorie des particules élémentaires de spin quelconque. I *Physica* **9** 310–324.
- [38] Lubański J K 1942 Sur la théorie des particules élémentaires de spin quelconque. II *Physica* **9** 325–338.
- [39] Majorana E 1932 Teoria relativistica di particelle con momento intrinseco arbitrario *Nuovo Cimento* **9** 335–344; reprinted in English: Relativistic theory of particles with arbitrary intrinsic angular momentum, Majorana E 2006 *Scientific Papers* ed J F Bassani (Berlin, Heidelberg, New York: Springer) pp 150–159.
- [40] McDonald A B 2015 The Sunbury Neutrino Observatory: Observation of flavor change for solar neutrinos *Nobel Lecture*, December 8; http://www.nobelprize.org/nobel_prizes/physics/laureates/2015/mcdonald-lecture.html
- [41] Minkowski H 1908, Die Grundlagen für die elektromagnetischen Vorgänge in bewegten Körpern *Nachr. König. Ges. Wiss. Göttingen, math.-phys. Kl.* 53–111.

- [42] Misner C W, Thorne K S and Wheeler J A 1973 *Gravitation* (San Francisco: Freeman).
- [43] Moskalev A N 2006 *Relativistic Field Theory* (St. Petersburg: PIYaF RAN) (in Russian).
- [44] Mukhanov V and Winitzki S 2007 *Introduction to Quantum Effects in Gravity* (Cambridge: Cambridge University Press).
- [45] Naber G L 2012 *The Geometry of Minkowski Spacetime: An Introduction to the Mathematics of the Special Theory of Relativity* 2nd edn (Berlin, Heidelberg, New York: Springer).
- [46] Parker E and Taubes C H 1982 On Witten's proof of the positive energy theorem *Comm. Math. Phys.* **84** 223–238.
- [47] Pauli W 1936 Contributions mathématiques à la théorie des matrices de Dirac *Ann. Inst. H. Poincaré* **6** 109–136.
- [48] Pauli W 1941 Relativistic field theories of elementary particles *Rev. Mod. Phys.* **13** 203–232.
- [49] Pauli W 1958 *Theory of Relativity* (Oxford: Pergamon).
- [50] Peskin M E and Schroeder D V 1995 *An Introduction to Quantum Field Theory* (Boulder: Westview Press).
- [51] Pontecorvo B M 1983 Pages in the development of neutrino physics *Phys. Usp.* **26** 1087–1108.
- [52] Proca A 1936 Sur la théorie ondulatoire des électrons positifs et négatifs *J. Phys. Radium* **7** 347–353.
- [53] Roy A 2001 Discovery of parity violation: breakdown of a symmetry principle *Resonance* **6** 32–43.
- [54] Rumer Yu B and Fet A I 1977 *Group Theory and Quantized Fields* (Moscow: Nauka) (in Russian).
- [55] Ryder L H 1996, *Quantum Field Theory* 2nd edn (Cambridge: Cambridge University Press).
- [56] Scheck F 2007 *Quantum Physics* (Berlin, Heidelberg, New York: Springer).
- [57] Schiff L I 1968 *Quantum Mechanics* 3rd edn (New York: McGraw-Hill).
- [58] Schweber S S 1961 *An Introduction to Relativistic Quantum Field Theory* (Evanston, Illinois; Elmsford, New York: Row, Peterson and Company).
- [59] Schweber S S, Bethe H A and Hoffmann D F 1955 *Mesons and Fields* vol 1 (Evanston, Illinois; Elmsford, New York: Row, Peterson and Company).
- [60] Sohnius M F 1985 Introducing supersymmetry *Phys. Rep.* **128** 39–204.
- [61] Toptygin I N 2014 *Foundations of Classical and Quantum Electrodynamics* (Weinheim: Wiley-VCH).
- [62] Varshalovich D A, Moskalev A N and Khersonskii V K 1988 *Quantum Theory of Angular Momentum* (Singapore: World Scientific).
- [63] Vizgin V P and Smorodinskii Ya A 1979 From the equivalence principle to the equations of gravitation *Phys. Usp.* **22** 489–513.
- [64] Wald R N 1984 *General Relativity* (Chicago: University of Chicago Press).
- [65] Weinberg S 1972 *Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity* (New York: Wiley).
- [66] Weinberg S 1998 *The Quantum Theory of Fields* vol 1 (Cambridge: Cambridge University Press).
- [67] Weinberg S 2008 *Cosmology* (Oxford: Oxford University Press).
- [68] Weyl H 1929 Gravitation and the electron *Proc. Nat. Acad. Sci. U.S.A.* **15** 323–334.
- [69] Weyl H 1944 How far can one get with a linear field theory of gravitation in flat space-time? *Amer. J. Math.* **66** 591–604.
- [70] Wigner E 1939 On unitary representations of the inhomogeneous Lorentz group *Ann. Math.* **40** 149–204.
- [71] Witten E 1981 A new proof of the positive energy theorem *Comm. Math. Phys.* **80** 381–402.
- [72] Wu C S, Ambler E, Hayward R W, Hoppes D D and Hudson R P 1957 Experimental test of parity conservation in beta decay *Phys. Rev.* **105** 1413–1415.
- [73] Zakharov V D 1973 *Gravitational waves in Einstein's theory* (New York: Wiley).

SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, ARIZONA STATE UNIVERSITY, TEMPE, AZ 85287–1804, U.S.A.

E-mail address: `sergeykryuchkov@yahoo.com`

SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, ARIZONA STATE UNIVERSITY, TEMPE, AZ 85287–1804, U.S.A.

E-mail address: `nlanfear@asu.edu`

SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES, ARIZONA STATE UNIVERSITY, TEMPE, AZ 85287–1804, U.S.A.

E-mail address: `sks@asu.edu`

URL: `http://hahn.la.asu.edu/~suslov/index.html`