

AN INDEX THEOREM FOR CR MANIFOLDS WITH S^1 ACTION

JIH-HSIN CHENG, CHIN-YU HSIAO, AND I-HSUN TSAI

ABSTRACT. Let $(X, T^{1,0}X)$ be a compact CR manifold of dimension $2n + 1$ with a transversal CR locally free S^1 action and let E be a rigid CR vector bundle over X . For every $m \in \mathbb{Z}$, let $H_{b,m}^j(X, E)$ be the m -th S^1 Fourier coefficient of the j -th $\bar{\partial}_b$ Kohn-Rossi cohomology group with values in E . In this paper, we prove that the Euler characteristic $\sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, E)$ can be computed in terms of the tangential Chern character of E , the tangential Todd class of $T^{1,0}X$, and the Chern polynomial of the Levi curvature of X . As applications, we can produce many CR functions on a weakly pseudoconvex CR manifold with such an S^1 action and many CR sections on some class of CR manifolds. In some cases, we can reinterpret Kawasaki's Hirzebruch-Riemann-Roch formula for a complex orbifold with an orbifold holomorphic line bundle by an integral over a smooth CR manifold.

CONTENTS

1. Introduction and statement of the results	2
1.1. Introduction and Motivation	2
1.2. Main theorem	3
1.3. Applications	5
1.4. Kawasaki's Hirzebruch-Riemann-Roch and Grauert-Riemenschneider criterion for orbifold line bundles	8
1.5. Examples	10
1.6. Proof of Theorem 1.2	12
1.7. The idea of the proof of Theorem 1.3	14
2. Preliminaries	20
2.1. Some standard notations	21
2.2. Set up and terminology	21
2.3. Tangential De-Rham cohomology group, Tangential Chern character and Tangential Todd class	23
2.4. BRT trivializations and rigid Hermitian metrics	24
3. Hodge theory for $\square_{b,m}^{(q)}$	26
4. Modified Kohn Laplacians	30
5. Asymptotic expansions for the heat kernels of the modified Kohn Laplacians	38
5.1. Heat kernels of the modified Kodaira Laplacians on BRT trivializations	38
5.2. Heat kernels of the modified Kohn Laplacians	43
6. Local density for the Euler characteristic $\sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, E)$	48
References	50

The second-named author was partially supported by Taiwan Ministry of Science of Technology projects 103-2115-M-001-001, 104-2628-M-001-003-MY2 and the Golden-Jade fellowship of Kenda Foundation.

1. INTRODUCTION AND STATEMENT OF THE RESULTS

1.1. Introduction and Motivation. Let $(X, T^{1,0}X)$ be a compact CR manifold of dimension $2n+1$ and let $\bar{\partial}_b : \Omega^{0,q}(X) \rightarrow \Omega^{0,q+1}(X)$ be the tangential Cauchy-Riemann operator. For a smooth function u , we say u is CR if $\bar{\partial}_b u = 0$. In CR geometry, it is crucial to be able to produce many CR functions. When X is strongly pseudoconvex and dimension of X is greater than or equal to five, it is well-known that the space of global smooth CR functions $H_b^0(X)$ is infinite dimensional. When X is weakly pseudoconvex or the Levi form of X has negative eigenvalues, the space of global CR functions could be trivial. In general, it is very difficult to determine when the space $H_b^0(X)$ is large. Since $\bar{\partial}_b^2 = 0$, we have $\bar{\partial}_b$ -complex: $\dots \rightarrow \Omega^{0,q-1}(X) \rightarrow \Omega^{0,q}(X) \rightarrow \Omega^{0,q+1}(X) \rightarrow \dots$ and we can define cohomology group: $H_b^q(X) := \frac{\text{Ker } \bar{\partial}_b : \Omega^{0,q}(X) \rightarrow \Omega^{0,q+1}(X)}{\text{Im } \bar{\partial}_b : \Omega^{0,q-1}(X) \rightarrow \Omega^{0,q}(X)}$. As in complex geometry, to understand the space $H_b^0(X)$, we could

try to establish CR Hirzebruch-Riemann-Roch theorem for the Euler characteristic $\sum_{j=0}^n (-1)^j \dim H_b^j(X)$

and vanishing theorems for $H_b^j(X)$, $j \geq 1$. But the difficulty comes from the fact that $\dim H_b^j(X)$ could be infinite for some j . Let's see some examples. If X is strongly pseudoconvex of dimension ≥ 5 , it is well-known that $\bar{\partial}_b : \Omega^{0,0}(X) \rightarrow \Omega^{0,1}(X)$ is not hypoelliptic but $\bar{\partial}_b : \Omega^{0,q}(X) \rightarrow \Omega^{0,q+1}(X)$ is hypoelliptic, for $q \geq 1$. We have $\dim H_b^0(X) = \infty$ and $\dim H_b^q(X) < \infty$, for $q \geq 1$. If the Levi form of X has exactly one negative, $n-1$ positive eigenvalues on X and $n > 3$, it is well-known that $\dim H_b^1(X) = \infty$, $\dim H_b^{n-1}(X) = \infty$ and $\dim H_b^q(X) < \infty$, for $q \notin \{1, n-1\}$. In general, we don't know how to define the Euler characteristic $\sum_{j=0}^n (-1)^j \dim H_b^j(X)$. On the other hand, even if $\bar{\partial}_b$ is

hypoelliptic, it is still difficult to calculate local index formula for $\bar{\partial}_b$ operator. We are lead to ask the following question:

Question 1.1. Can we establish some kind of CR Hirzebruch-Riemann-Roch theorem (not the usual one) on some class of CR manifolds?

To answer this question, let's come back to complex geometry case. Consider a compact complex manifold M of dimension n and let $(L, h^L) \rightarrow M$ be a holomorphic line bundle over M , where h^L denotes a Hermitian fiber metric of L . Let $(L^*, h^{L^*}) \rightarrow M$ be the dual bundle of (L, h^L) and put $X = \{v \in L^*; |v|_{h^{L^*}}^2 = 1\}$. We call X the circle bundle of (L^*, h^{L^*}) . It is clear that X is a compact CR manifold of dimension $2n+1$. Given a local holomorphic frame s of L on an open subset $U \subset M$ we define the associated local weight of h^L by $|s(x)|_{h^L}^2 = e^{-2\phi(x)}$, $\phi \in C^\infty(U, \mathbb{R})$. The CR manifold X is equipped with a natural S^1 action. Locally X can be represented in local holomorphic coordinates $(z, \lambda) \in \mathbb{C}^{n+1}$, where λ is the fiber coordinate, as the set of all (z, λ) such that $|\lambda|^2 e^{2\phi(z)} = 1$, where ϕ is a local weight of h^L . The S^1 action on X is given by $e^{-i\theta} \circ (z, \lambda) = (z, e^{-i\theta} \lambda)$, $e^{-i\theta} \in S^1$, $(z, \lambda) \in X$. Let $T \in C^\infty(X, TX)$ be the real vector field induced by the S^1 action, that is, $Tu = \frac{\partial}{\partial \theta}(u(e^{-i\theta} \circ x))|_{\theta=0}$, $u \in C^\infty(X)$. We can check that $[T, C^\infty(X, T^{1,0}X)] \subset C^\infty(X, T^{1,0}X)$ and $\mathbb{C}T(x) \oplus T_x^{1,0}X \oplus T_x^{0,1}X = \mathbb{C}T_x X$ (we say that the S^1 action is CR and transversal). For every $m \in \mathbb{Z}$ and $q = 0, 1, 2, \dots, n$, put

$$\begin{aligned} \Omega_m^{0,q}(X) &:= \{u \in \Omega^{0,q}(X); Tu = -imu\} \\ &= \left\{ u \in \Omega^{0,q}(X); u(e^{-i\theta} \circ x) = e^{-im\theta} u(x), \forall \theta \in [0, 2\pi[\right\}. \end{aligned}$$

Since $\bar{\partial}_b T = T \bar{\partial}_b$, we have $\bar{\partial}_{b,m} : \Omega_m^{0,q}(X) \rightarrow \Omega_m^{0,q+1}(X)$ and we could consider the cohomology group: $H_{b,m}^q(X) := \frac{\text{Ker } \bar{\partial}_{b,m} : \Omega_m^{0,q}(X) \rightarrow \Omega_m^{0,q+1}(X)}{\text{Im } \bar{\partial}_{b,m} : \Omega_m^{0,q-1}(X) \rightarrow \Omega_m^{0,q}(X)}$. The following theorem can be seen as the starting point of this work.

Theorem 1.2. Let $\Omega^{0,q}(M, L^m)$ be the space of smooth sections of $(0, q)$ forms of M with values in L^m and let $H^q(M, L^m)$ be the q -th $\bar{\partial}$ Dolbeault cohomology group with value in L^m , where L^m is the m -th power of L . For every $q = 0, 1, 2, \dots, n$, and every $m \in \mathbb{Z}$, there is a bijection map $A_m^{(q)} : \Omega_m^{0,q}(X) \rightarrow$

$\Omega^{0,q}(M, L^m)$ such that $A_m^{(q+1)}\bar{\partial}_{b,m} = \bar{\partial}A_m^{(q)}$ on $\Omega_m^{0,q}(X)$. Hence, $\Omega_m^{0,q}(X) \cong \Omega^{0,q}(M, L^m)$ and $H_{b,m}^q(X) \cong H^q(M, L^m)$. In particular $\dim H_{b,m}^q(X) < \infty$, for every m and every q and $\sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X) = \sum_{j=0}^n (-1)^j \dim H^j(M, L^m)$.

Theorem 1.2 is probably well-known. Since it is difficult to indicate a precise reference, we will give a proof of Theorem 1.2 in Section 1.6 for the convenience of the reader.

Hirzebruch-Riemann-Roch Theorem or Atiyah-Singer index Theorem tells us that

$$(1.1) \quad \sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X) = \sum_{j=0}^n (-1)^j \dim H^j(M, L^m) = \int_M \text{Td}(T^{1,0}M) \text{ch}(L^m),$$

where $\text{Td}(T^{1,0}M)$ denotes the Todd class of $T^{1,0}M$ the holomorphic tangent bundle of M and $\text{ch}(L^m)$ denotes the Chern character of L^m . Let's reformulate (1.1) in terms of characteristic classes on X :

$$(1.2) \quad \sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X) = \frac{1}{2\pi} \int_X \text{Td}_b(T^{1,0}X) \wedge e^{-m \frac{d\omega_0}{2\pi}} \wedge \omega_0,$$

where $\text{Td}_b(T^{1,0}X)$ denotes the tangential Todd class of $T^{1,0}X$ and $e^{-m \frac{d\omega_0}{2\pi}}$ denotes the Chern polynomial of the Levi curvature and ω_0 is the uniquely determined global real 1-form (see Section 2.2 and Section 2.3 for the precise definitions). The goal of this work is to generalize (1.2) to any CR manifold with S^1 action. It turns out that to establish (1.1) on any CR manifold with S^1 action, we need to calculate local density function for the Euler characteristic $\sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X)$ and this is a non-trivial generalization. The above Euler characteristic has an index interpretation related to $\bar{\partial}_b + \bar{\partial}_b^*$ (see (3.23), (3.24) and (3.25)).

There is another index theory of geometric significance, developed by Charles Epstein. He studied so called relative index of a pair of CR structures through their Szegő projectors in a series of papers (see [11], [12], [13], [14] and [15]). On the other hand, Erik van Erp derived an index formula for subelliptic operators on a contact manifold (see [16], [17]).

1.2. Main theorem. We now formulate the main results. We refer to Section 2.2 and Section 2.3 for some notations and terminology used here.

Let $(X, T^{1,0}X)$ be a compact CR manifold with a transversal CR locally free S^1 action $e^{-i\theta}$, where $T^{1,0}X$ is a CR structure of X . Let $T \in C^\infty(X, TX)$ be the real vector field induced by the S^1 action and let $\omega_0 \in C^\infty(X, T^*X)$ be the global real one form determined by $\langle \omega_0, T \rangle = 1$, $\langle \omega_0, u \rangle = 0$, for every $u \in T^{1,0}X \oplus T^{0,1}X$.

Let E be a rigid CR vector bundle over X (see Definition 2.4) and let $\bar{\partial}_b : \Omega^{0,q}(X, E) \rightarrow \Omega^{0,q+1}(X, E)$ be the tangential Cauchy-Riemann operator with values in E . For every $u \in \Omega^{0,q}(X, E)$, we can define $Tu \in \Omega^{0,q}(X, E)$ and we have $T\bar{\partial}_b = \bar{\partial}_b T$. For $m \in \mathbb{Z}$, put

$$\begin{aligned} \Omega_m^{0,q}(X, E) &:= \{u \in \Omega^{0,q}(X, E); Tu = -imu\} \\ &= \left\{ u \in \Omega^{0,q}(X, E); (e^{-i\theta})^* u = e^{-im\theta} u, \forall \theta \in [0, 2\pi[\right\}, \end{aligned}$$

where $(e^{-i\theta})^*$ denotes the pull-back map by $e^{-i\theta}$ (see (2.3)). Since $T\bar{\partial}_b = \bar{\partial}_b T$, we have

$$\bar{\partial}_{b,m} := \bar{\partial}_b : \Omega_m^{0,q}(X, E) \rightarrow \Omega_m^{0,q+1}(X, E).$$

Thus, for every $m \in \mathbb{Z}$, we have $\bar{\partial}_{b,m}$ -complex:

$$\bar{\partial}_{b,m} : \cdots \rightarrow \Omega_m^{0,q-1}(X, E) \rightarrow \Omega_m^{0,q}(X, E) \rightarrow \Omega_m^{0,q+1}(X, E) \rightarrow \cdots$$

and we can consider q -th $\bar{\partial}_{b,m}$ Kohn-Rossi cohomology group:

$$H_{b,m}^q(X, E) := \frac{\text{Ker } \bar{\partial}_{b,m} : \Omega_m^{0,q}(X, E) \rightarrow \Omega_m^{0,q+1}(X, E)}{\text{Im } \bar{\partial}_{b,m} : \Omega_m^{0,q-1}(X, E) \rightarrow \Omega_m^{0,q}(X, E)}.$$

When E is trivial, for every $m \in \mathbb{Z}$, we denote $\Omega_m^{0,q}(X) := \Omega_m^{0,q}(X, E)$ and $H_{b,m}^q(X) := H_{b,m}^q(X, E)$. We will prove in Theorem 3.7 that without any Levi curvature assumption, $\dim H_{b,m}^q(X, E) < \infty$, for every $m \in \mathbb{Z}$ and every $q = 0, 1, 2, \dots, n$.

For $x \in X$, we say that the period of x is $\frac{2\pi}{\ell}$, $\ell \in \mathbb{N}$, if $e^{-i\theta} \circ x \neq x$, for every $0 < \theta < \frac{2\pi}{\ell}$ and $e^{-i\frac{2\pi}{\ell}} \circ x = x$. For each $\ell \in \mathbb{N}$, put

$$(1.3) \quad X_\ell = \left\{ x \in X; \text{ the period of } x \text{ is } \frac{2\pi}{\ell} \right\}$$

and let $p = \min \{ \ell \in \mathbb{N}; X_\ell \neq \emptyset \}$. The main result of this work is the following

Theorem 1.3. *Suppose $(X, T^{1,0}X)$ is a compact, connected CR manifold with a transversal CR locally free S^1 action. Then, X_p is dense in X and*

$$(1.4) \quad \begin{aligned} & \sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, E) \\ &= \sum_{s=1}^p e^{\frac{2\pi(s-1)}{p}mi} \frac{1}{2\pi} \int_X \text{Td}_b(T^{1,0}X) \wedge \text{ch}_b(E) \wedge e^{-m\frac{d\omega_0}{2\pi}} \wedge \omega_0, \end{aligned}$$

where $\text{Td}_b(T^{1,0}X)$ denotes the tangential Todd class of $T^{1,0}X$ and $\text{ch}_b(E)$ denotes the tangential Chern character of E .

We refer the reader to Section 2.3 for the precise meanings of $\text{Td}_b(T^{1,0}X)$ and $\text{ch}_b(E)$.

Remark 1.4. We note that there is topological obstruction for a CR manifold to admit a transversal CR S^1 action. For instance, a compact strictly pseudoconvex CR 3-manifold must have even first Betti number if admitting a transversal CR S^1 action. The reason is that such a manifold must have free pseudohermitian torsion (see [25]). On the other hand, Alan Weinstein showed that vanishing pseudohermitian torsion implies even first Betti number (see the Appendix in [6]). In this paper, we only consider that the S^1 action is transversal and locally free. It should be mentioned that the S^1 action on a circle bundle is globally free. We give some examples:

Example I: $X = \left\{ (z_1, z_2, z_3) \in \mathbb{C}^3; |z_1|^2 + |z_2|^2 + |z_3|^2 + |z_1^2 + z_2|^4 + |z_2^3 + z_3|^6 = 1 \right\}$. Then X admits a transversal CR locally free S^1 action: $e^{-i\theta} \circ (z_1, z_2, z_3) = (e^{-i\theta} z_1, e^{-2i\theta} z_2, e^{-6i\theta} z_3)$. It is clear that this S^1 action is not globally free.

Example II: Let X be a compact orientable Seifert 3-manifold. Kamishima and Tsuboi [23] proved that X is a compact CR manifold with a transversal CR locally free S^1 action. Moreover, X is a circle bundle but with singular base.

In Section 1.5, we collect more examples.

Remark 1.5. (I) The number $\sum_{s=1}^p e^{\frac{2\pi(s-1)}{p}mi}$ is equal to 0 or p .

(II) X_p is dense in X is a famous result in topology (see Proposition 1.24 in Meinrenken [29] and Duistermaat-Heckman [9]). We prove this result by computing local density function of the Euler characteristic $\sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, E)$.

Remark 1.6. Let $p_1 = p < p_2 < p_3 < \dots < p_N$ be the periods of X with respect to the given S^1 action

$$\begin{aligned} S^1 \times X &\rightarrow X, \\ (e^{-i\theta}, x) &\rightarrow e^{-i\theta} \circ x. \end{aligned}$$

We claim that one has $p_1 = p$ divides every others. The reason is that, the subgroup \mathbb{Z}_p acts trivially on the dense open subset which is this p -stratum or p_1 -stratum, hence it acts trivially on the whole X by continuity and density of this stratum. Hence, the fixed groups q , $q = p_j$, $j = 2, \dots, N$, of any other stratum must contain \mathbb{Z}_p . Namely \mathbb{Z}_p is a subgroup of \mathbb{Z}_q . Hence, $\frac{q}{p} \in \mathbb{N}$. The claim follows.

By the above fact, one may re-normalize the given S^1 -action by lifting, so as to make the new S^1 -action have $p_1 = 1$. More precise, define

$$\begin{aligned} S^1 \times X &\rightarrow X, \\ (e^{-i\theta}, x) &\rightarrow e^{-i\theta} \diamond x := e^{-i\frac{\theta}{p}} \circ x. \end{aligned}$$

The new S^1 -action $(e^{-i\theta}, \diamond)$ has $p_1 = 1$. Let $\tilde{\omega}_0$ be the global real one form with respect to $(e^{-i\theta}, \diamond)$ and for every $m \in \mathbb{Z}$, let $\tilde{H}_{b,m}^q(X, E)$ be the m -th S^1 Fourier coefficient of the q -th Kohn-Rossi cohomology group with respect to $(e^{-i\theta}, \diamond)$. It is straightforward to see that

$$(1.5) \quad \begin{aligned} \tilde{\omega}_0 &= p\omega_0, \\ \tilde{H}_{b,m}^q(X, E) &= H_{b,pm}^q(X, E), \quad \forall m \in \mathbb{Z}, \quad \forall q = 0, 1, 2, \dots, n. \end{aligned}$$

From (1.5) and Theorem 1.3, we see that the index formulas can be transformed to each other: the one before the above reduction and the one after the reduction. The point is that, the old one shall have a sum of exponential terms, and the new one not.

1.3. Applications.

1.3.1. *Applications in CR geometry.* In CR geometry, it is crucial to be able to produce many CR functions or CR sections. Put

$$H_b^0(X, E) = \{u \in C^\infty(X, E); \bar{\partial}_b u = 0\}.$$

Kohn asked the following question

Question 1.7. Let X be a compact weakly pseudoconvex CR manifold. When is the space $H_b^0(X, E)$ large?

See the discussion after Definition 2.2 for the meanings of weakly pseudoconvex and strongly pseudoconvex CR manifolds.

In [26], Lempert proved that a three dimensional compact strongly pseudoconvex CR manifold X with a locally free transversal CR S^1 -action can be CR embedded into \mathbb{C}^N . In [10], Epstein proved that a three dimensional compact strongly pseudoconvex CR manifold X with a global free transversal CR S^1 action can be embedded into \mathbb{C}^N by the positive Fourier coefficients. The embeddability of X by positive Fourier coefficients is related to the behavior of the S^1 action on X . For example, suppose that we can find $f_1 \in H_{b,m}^0(X), \dots, f_{d_m} \in H_{b,m}^0(X)$ and $g_1 \in H_{b,m_1}^0(X), \dots, g_{h_{m_1}} \in H_{b,m_1}^0(X)$ such that the map

$$\Phi_{m,m_1} : x \in X \rightarrow (f_1(x), \dots, f_{d_m}(x), g_1(x), \dots, g_{h_{m_1}}(x)) \in \mathbb{C}^{d_m+h_{m_1}}$$

is a CR embedding. The S^1 action on X induces naturally a S^1 action on $\Phi_{m,m_1}(X)$ and this S^1 action on $\Phi_{m,m_1}(X)$ is simply given by the following:

$$e^{-i\theta} \circ (z_1, \dots, z_{d_m}, z_{d_m+1}, \dots, z_{d_m+h_{m_1}}) = (e^{-im\theta} z_1, \dots, e^{-im\theta} z_{d_m}, e^{-im_1\theta} z_{d_m+1}, \dots, e^{-im_1\theta} z_{d_m+h_{m_1}}).$$

Thus, if we could embed such CR manifold by positive Fourier coefficients, we can describe the S^1 action explicitly. To study the embedding theorem of such CR manifold by positive Fourier coefficients, it is crucial to be able to know

Question 1.8. When $\dim H_{b,m}^0(X, E) \approx m^n$ for m large?

Combining our index theorems with some vanishing theorems, we could answer Question 1.7 and Question 1.8. From Theorem 1.3, we deduce

Corollary 1.9. *Suppose we have the same assumption as in Theorem 1.3. Then,*

$$(1.6) \quad \begin{aligned} & \sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, E) \\ &= r \sum_{s=1}^p e^{\frac{2\pi(s-1)}{p}mi} \frac{m^n}{n!(2\pi)^{n+1}} \int_X (-d\omega_0)^n \wedge \omega_0 + O(m^{n-1}), \end{aligned}$$

where r denotes the complex rank of the vector bundle E .

We can repeat the proof of Theorem 2.1 in [22] with minor change and get

Theorem 1.10. *Suppose we have the same assumption as in Theorem 1.3 and assume that X is weakly pseudoconvex. Then, $\dim H_{b,m}^j(X, E) = o(m^n)$, for every $j = 1, 2, \dots, n$.*

From Corollary 1.9 and Theorem 1.10, we obtain

Theorem 1.11. *Suppose we have the same assumption as in Theorem 1.3 and assume that X is weakly pseudoconvex. Then for $m \gg 1$,*

$$\begin{aligned} & \dim H_{b,m}^0(X, E) \\ &= r \sum_{s=1}^p e^{\frac{2\pi(s-1)}{p}mi} \frac{m^n}{n!(2\pi)^{n+1}} \int_X (-d\omega_0)^n \wedge \omega_0 + o(m^n), \end{aligned}$$

where r denotes the complex rank of the vector bundle E .

In particular, if the Levi form is strongly pseudoconvex at some point of X , then $\dim H_{b,pm}^0(X) = O(m^n)$, for $m \gg 1$ and hence, $\dim H_b^0(X, E) = \infty$.

From Theorem 1.11, we answer Question 1.7 and Question 1.8.

In some applications, we need to sum over m . Given a CR manifold X , when X can be CR embedded into complex space is an important problem in CR geometry. When X is strongly pseudoconvex and dimension of X is greater than or equal to five, a classical theorem of L. Boutet de Monvel [3] asserts that X can be globally CR embedded into \mathbb{C}^N , for some $N \in \mathbb{N}$. When X is not strongly pseudoconvex, the space of global CR functions could be even trivial. Since many interesting examples live in the projective space (e. g. the quadric $\{|z\} \in \mathbb{C}\mathbb{P}^{N-1}; |z_1|^2 + \dots + |z_q|^2 - |z_{q+1}|^2 - \dots - |z_N|^2 = 0\}$), it is thus natural to consider a setting analogous to the Kodaira embedding theorem and ask if X can be embedded into the projective space by means of CR sections of a CR line bundle. Inspired by Kodaira, we consider L^k the k -th power of a CR line bundle $L \rightarrow X$. If the dimension of the space $H_b^0(X, L^k)$ of CR sections of L^k is large, when $k \rightarrow \infty$, one should find many CR sections to embed X into projective space. The following question is asked by Henkin and Marinescu [28, p.47-48]

Question 1.12. When is $\dim H_b^0(X, L^k) = O(k^{n+1})$ for k large?

We now assume that L is a rigid CR line bundle with a rigid Hermitian fiber metric h^L (see Definition 2.4 and Definition 2.9 for the precise meanings of rigid line bundle and rigid Hermitian metric) and let $\mathcal{R}^L \in \Omega_0^2(X)$ be the curvature of L induced by h^L . If s is a local trivializing section of L on an open subset $D \subset X$, we have $|s(x)|_{h^L}^2 = e^{-2\phi(x)}$, $\forall x \in D$, where $\phi \in C^\infty(D, \mathbb{R})$ with $T\phi = 0$. Then, $\mathcal{R}^L = 2\partial_b\bar{\partial}_b\phi \in \Omega_0^2(X)$. For $k > 0$, let (L^k, h^{L^k}) be the k -th power of (L, h^L) .

Theorem 1.13. *With the notations above, for k large, we have*

$$(1.7) \quad \begin{aligned} & \sum_{m \in \mathbb{Z}, |m| \leq k\delta} \sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, L^k \otimes E) \\ &= r(2\pi)^{-n-1} \frac{1}{n!} k^{n+1} \int_X \int_{[-\delta, \delta]} (i\mathcal{R}_x^L - sd\omega_0(x))^n \wedge \omega_0(x) ds + o(k^{n+1}), \end{aligned}$$

where $\delta > 0$ and r denotes the complex rank of the vector bundle E .

Proof. From Theorem 1.3, we can check that

$$\begin{aligned}
(1.8) \quad & \sum_{m \in \mathbb{Z}, |m| \leq k\delta} \sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, L^k \otimes E) \\
&= r(2\pi)^{-n-1} \sum_{m \in \mathbb{Z}, |m| \leq k\delta} \sum_{s=1}^p e^{\frac{2\pi(s-1)}{p}mi} \frac{1}{n!} \int_X (ik\mathcal{R}_x^L - md\omega_0(x))^n \wedge \omega_0(x) + o(k^n) \\
&= r(2\pi)^{-n-1} \sum_{m \in \mathbb{Z}, |m| \leq k\delta} \sum_{s=1}^p e^{\frac{2\pi(s-1)}{p}mi} \frac{k^n}{n!} \int_X (i\mathcal{R}_x^L - \frac{m}{k}d\omega_0(x))^n \wedge \omega_0(x) + o(k^n).
\end{aligned}$$

Note that $\sum_{s=1}^p e^{\frac{2\pi(s-1)}{p}mi} = p$ if m is divided by p and $\sum_{s=1}^p e^{\frac{2\pi(s-1)}{p}mi} = 0$ if m is not divided by p . From this observation and (1.8), we get

$$\begin{aligned}
(1.9) \quad & \sum_{m \in \mathbb{Z}, |m| \leq k\delta} \sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, L^k \otimes E) \\
&= r(2\pi)^{-n-1} \sum_{\ell \in \mathbb{Z}, |p\ell| \leq k\delta} \frac{p}{n!} k^n \int_X (i\mathcal{R}_x^L - \frac{p\ell}{k}d\omega_0(x))^n \wedge \omega_0(x) + o(k^n).
\end{aligned}$$

It is clear that the Riemann sum $\sum_{\ell \in \mathbb{Z}, |p\ell| \leq k\delta} \frac{p}{k} \int_X (i\mathcal{R}_x^L - \frac{p\ell}{k}d\omega_0(x))^n \wedge \omega_0(x)$ converges to

$$\int_X \int_{[-\delta, \delta]} (i\mathcal{R}_x^L - sd\omega_0(x))^n \wedge \omega_0(x) ds$$

as $k \rightarrow \infty$, and hence

$$\begin{aligned}
(1.10) \quad & \sum_{\ell \in \mathbb{Z}, |p\ell| \leq k\delta} \frac{p}{n!} k^n \int_X (i\mathcal{R}_x^L - \frac{p\ell}{k}d\omega_0(x))^n \wedge \omega_0(x) \\
&= \frac{1}{n!} k^{n+1} \int_X \int_{[-\delta, \delta]} (i\mathcal{R}_x^L - sd\omega_0(x))^n \wedge \omega_0(x) ds + o(k^{n+1}).
\end{aligned}$$

Combining (1.10) with (1.9), (1.7) follows. \square

Definition 1.14. We say that (L, h^L) is positive at $p \in X$ if \mathcal{R}_p^L is a positive Hermitian quadratic form over $T_p^{1,0}X$. We say that (L, h^L) is semi-positive if for any $x \in X$ there exists a constant $\delta > 0$ such that $\mathcal{R}_x^L - sid\omega_0(x)$ is a semi positive Hermitian quadratic form over $T_x^{1,0}X$ for any $|s| < \delta$.

We can repeat the proof of Theorem 1.24 in [21] with minor change and get

Theorem 1.15. Assume that (L, h^L) is a semi-positive CR line bundle over X . Then, for $\delta > 0$, δ small, we have

$$\sum_{m \in \mathbb{Z}, |m| \leq k\delta} \dim H_{b,m}^j(X, L^k \otimes E) = o(k^{n+1}), \quad j = 1, 2, \dots, n.$$

From Theorem 1.13 and Theorem 1.15, we get

Theorem 1.16. Assume that (L, h^L) is semi-positive. Then, for $\delta > 0$, δ small, we have

$$\begin{aligned}
(1.11) \quad & \sum_{m \in \mathbb{Z}, |m| \leq k\delta} \dim H_{b,m}^0(X, L^k \otimes E) \\
&= r(2\pi)^{-n-1} \frac{1}{n!} k^{n+1} \int_X \int_{[-\delta, \delta]} (i\mathcal{R}_x^L - sd\omega_0(x))^n \wedge \omega_0(x) ds + o(k^{n+1}),
\end{aligned}$$

where r denotes the complex rank of the vector bundle E .

In particular, if (L, h^L) is positive at some point of X , then $\dim H_b^0(X, L^k \otimes E) = O(k^{n+1})$.

From Theorem 1.16, we have answered Question 1.12.

1.4. Kawasaki's Hirzebruch-Riemann-Roch and Grauert-Riemenschneider criterion for orbifold line bundles. Kawasaki's Hirzebruch-Riemann-Roch formula, introduced by Kawasaki [24], is the Hirzebruch-Riemann-Roch formula for orbifolds. Comparing with Kawasaki's formula, we get a simpler Hirzebruch-Riemann-Roch formula for some class of orbifold line bundles from our main result Theorem 1.3. Moreover, from Theorem 1.11, we establish Grauert-Riemenschneider criterion for orbifold line bundles.

Let M be a manifold and let G be a compact Lie group. Assume that M admits a G -action:

$$\begin{aligned} G \times M &\rightarrow M, \\ (g, x) &\rightarrow g \circ x. \end{aligned}$$

We suppose that the action G on M is locally free, that is, for every point $x \in M$, the stabilizer group $G_x = \{g \in G; g \circ x = x\}$ of x is a finite subgroup of G . If G is an action of this type the quotient space

$$(1.12) \quad X := M/G$$

is an orbifold. A theorem of Satake [30] says that the converse is true: every orbifold has a presentation of the form (1.12). We assume that M is a compact connected complex manifold with complex structure $T^{1,0}M$. G induces an action on $\mathbb{C}TM$:

$$(1.13) \quad \begin{aligned} G \times \mathbb{C}TM &\rightarrow \mathbb{C}TM, \\ (g, u) &\rightarrow g^*u, \end{aligned}$$

where $g^* = (g^{-1})_*$, $(g^{-1})_*$ denotes the push-forward by g^{-1} on $\mathbb{C}TM$. We suppose that G is holomorphic, that is $g^*(T^{1,0}M) \subset T^{1,0}M$, for every $g \in G$. Let $T^{0,1}M := \overline{T^{1,0}M}$. Put $\mathbb{C}T(M/G) := \mathbb{C}TM/G$, $T^{0,1}(M/G) := T^{0,1}M/G$, $T^{1,0}(M/G) := T^{1,0}M/G$. Assume that $T^{0,1}(M/G) \cap T^{1,0}(M/G) = \{0\}$. Then, $T^{1,0}(M/G)$ is a complex structure on M/G and M/G is a complex orbifold. Suppose

$$\dim_{\mathbb{C}} T^{1,0}(M/G) = n.$$

Let L be a G -invariant holomorphic line bundle over M , that is, for every transition function h of L on an open set $U \subset M$, we have $h(g \circ x) = h(x)$, for every $g \in G$, $x \in U$ with $g \circ x \in U$. Suppose that L admits a locally free G -action:

$$(1.14) \quad \begin{aligned} G \times L &\rightarrow L, \\ (g, x) &\rightarrow g \circ x \end{aligned}$$

with $\pi(g \circ x) = g \circ (\pi(x))$, for every $g \in G$, where $\pi : L \rightarrow M$ denotes the natural projection, and where the action of G on L is linear on the fibers of L , that is, for every $g \in G$, every $z \in M$, we have $g \circ (s(z) \otimes \lambda) = s_1(g \circ z) \otimes \rho(g, z)\lambda$, for every $\lambda \in \mathbb{C}$, where s and s_1 are local sections of L defined near z and $g \circ z$ respectively and $\rho(g, z) \in \mathbb{C}$ depends smoothly on z and g . Then, L/G is an orbifold holomorphic line bundle over M/G . For every $m \in \mathbb{N}$, let L^m be the m -th power of L . Then, the G -action on L induces a locally free G -action on L^m :

$$(1.15) \quad \begin{aligned} G \times L^m &\rightarrow L^m, \\ (g, x) &\rightarrow g \circ x \end{aligned}$$

with $\pi_m(g \circ x) = g \circ (\pi_m(x))$, for every $g \in G$, where $\pi_m : L^m \rightarrow M$ denotes the natural projection, and where the action of G on L^m is linear on the fibers of L^m . Then, L^m/G is also an orbifold holomorphic line bundle over M/G . Now, we fix $m \in \mathbb{Z}$. Let $T^{*0,q}M$ denote the bundle of $(0, q)$ forms on M . Since G is holomorphic, G induces a natural action on $L^m \otimes T^{*0,q}M$:

$$(1.16) \quad \begin{aligned} G \times (L^m \otimes T^{*0,q}M) &\rightarrow L^m \otimes T^{*0,q}M, \\ (g, u) &\rightarrow g^*u. \end{aligned}$$

For every $q = 0, 1, 2, \dots, n$, put

$$(1.17) \quad \Omega^{0,q}(M/G, L^m/G) := \{u \in \Omega^{0,q}(M, L^m); g^*u = u, \quad \forall g \in G\},$$

where $\Omega^{0,q}(M, L^m)$ denotes the space of smooth sections with values in $T^{*0,q}M \otimes L^m$. The Cauchy-Riemann operator $\bar{\partial} : \Omega^{0,q}(M, L^m) \rightarrow \Omega^{0,q+1}(M, L^m)$ is G -invariant and we have $\bar{\partial}$ -complex:

$$\bar{\partial} : \dots \rightarrow \Omega^{0,q-1}(M/G, L^m/G) \rightarrow \Omega^{0,q}(M/G, L^m/G) \rightarrow \Omega^{0,q+1}(M/G, L^m/G) \rightarrow \dots$$

and we can consider q -th Dolbeault cohomology group:

$$H^q(M/G, L^m/G) := \frac{\text{Ker } \bar{\partial} : \Omega^{0,q}(M/G, L^m/G) \rightarrow \Omega^{0,q+1}(M/G, L^m/G)}{\text{Im } \bar{\partial} : \Omega^{0,q-1}(M/G, L^m/G) \rightarrow \Omega^{0,q}(M/G, L^m/G)}.$$

Let L^* be the dual bundle of L . Then, L^* is also a G -invariant holomorphic line bundle and L^* admits a locally free G -action:

$$\begin{aligned} G \times L^* &\rightarrow L^*, \\ (g, x) &\rightarrow g \circ x \end{aligned}$$

with $\pi^*(g \circ x) = g \circ (\pi^*(x))$, for every $g \in G$, where $\pi^* : L^* \rightarrow M$ denotes the natural projection, and where the action of G on L^* is linear on the fibers of L^* . Then, L^*/G is an orbifold holomorphic line bundle over M/G . Let $\text{Tot}(L^*)$ be the space of all non-zero vectors of L^* . Assume that $\text{Tot}(L^*)/G$ is a smooth manifold. Take any G -invariant Hermitian fiber metric h^{L^*} on L^* , set $\tilde{X} = \{v \in L^*; |v|_{h^{L^*}} = 1\}$ and put $X = \tilde{X}/G$. Since $\text{Tot}(L^*)$ is a smooth manifold, $X = \tilde{X}/G$ is a smooth manifold. The natural S^1 action on \tilde{X} induces a locally free S^1 action $e^{-i\theta}$ on X . Moreover, we can check that X is a CR manifold and the S^1 action on X is CR and transversal. We will use the same notations as before. We can repeat the proof of Theorem 1.2 with minor change and deduce that for every $q = 0, 1, 2, \dots, n$, and every $m \in \mathbb{Z}$, we have

$$(1.18) \quad H^q(M/G, L^m/G) \cong H_{b,m}^q(X), \quad \dim H^q(M/G, L^m/G) = \dim H_{b,m}^q(X).$$

We pause and introduce some notations. For every $x \in \text{Tot}(L^*)$ and $g \in G$, put $N(g, x) = 1$ if $g \notin G_x$ and $N(g, x) = \inf \{\ell \in \mathbb{N}; g^\ell = I\}$ if $g \in G_x$. Set

$$p = \inf \{N(g, x); x \in \text{Tot}(L^*), g \in G, g \neq I\},$$

where I denotes the identity element of G . It is straightforward to see that X_p is an open and dense subset of X , where X_p is given by (1.3). From Theorem 1.3 and (1.18), we deduce

Theorem 1.17. *With the notations used above and recall that we work with the assumptions that M is connected and $\text{Tot}(L^*)/G$ is smooth. Then, for every $m \in \mathbb{Z}$, we have*

$$(1.19) \quad \begin{aligned} &\sum_{j=0}^n (-1)^j \dim H^j(M/G, L^m/G) \\ &= \sum_{s=1}^p e^{\frac{2\pi(s-1)}{p}mi} \frac{1}{2\pi} \int_X \text{Td}_b(T^{1,0}X) \wedge e^{-m\frac{d\omega_0}{2\pi}} \wedge \omega_0. \end{aligned}$$

From Theorem 1.11 and (1.18), we establish Grauert-Riemenschneider criterion for orbifold line bundles

Theorem 1.18. *With the notations and assumptions used above, assume that there is a G -invariant Hermitian fiber metric h^L on L such that the associated curvature R^L is semi-positive and positive at some point of M . Then, $\dim H^0(M/G, L^{pm}/G) = O(m^n)$, for $m \gg 1$.*

In Section 1.5, we construct an orbifold holomorphic line bundle over a singular complex orbifold such that the assumptions of Theorem 1.17 and Theorem 1.18 are fulfilled (see the discussion after Corollary 1.20).

1.5. Examples. In this section, some examples of CR manifolds with locally free S^1 action are collected.

We first review the construction of generalized Hopf manifolds introduced by Brieskorn and Van de Ven [4].

Let $a = (a_1, \dots, a_{n+2}) \in \mathbb{N}^{n+2}$, let $z = (z_1, \dots, z_{n+2})$ be the standard coordinates of \mathbb{C}^{n+2} and let $M(a)$ be the affine algebraic variety given by the equation

$$\sum_{j=1}^{n+2} z_j^{a_j} = 0.$$

If some $a_j = 1$, the variety $M(a)$ is non-singular. Otherwise $M(a)$ has exactly one singular point, namely $0 = (0, \dots, 0)$. Put $\widetilde{M}(a) := M(a) - \{0\}$. Now we define a holomorphic \mathbb{C} -action on $\widetilde{M}(a)$ by

$$t \circ (z_1, \dots, z_{n+2}) = (e^{\frac{t}{a_1}} z_1, \dots, e^{\frac{t}{a_{n+2}}} z_{n+2}), \quad t \in \mathbb{C}, \quad (z_1, \dots, z_{n+2}) \in \widetilde{M}(a).$$

It is easy to see that the \mathbb{Z} -action on $\widetilde{M}(a)$ is globally free. The equivalence class of $(z_1, \dots, z_{n+2}) \in \mathbb{C}^{n+2}$ with respect to the \mathbb{Z} -action is denoted by $(z_1, \dots, z_{n+2}) + \mathbb{Z}$ and hence

$$H(a) := \widetilde{M}(a)/\mathbb{Z} = \left\{ (z_1, \dots, z_{n+2}) + \mathbb{Z}; (z_1, \dots, z_{n+2}) \in \widetilde{M}(a) \right\}$$

is a compact complex manifold of complex dimension $n + 1$. We call $H(a)$ a Hopf manifold. Let Γ_a be the discrete subgroup of \mathbb{C} , generated by 1 and $2\pi\alpha i$, where α is the least common multiple of a_1, a_2, \dots, a_{n+2} . Consider the complex 1-torus $T_a = \mathbb{C}/\Gamma_a$. $H(a)$ admits a natural T_a -action. Put $V(a) := H(a)/T_a$. By Holmann [18], $V(a)$ is a complex orbifold. Let $\pi_a : H(a) \rightarrow V(a)$ be the natural projection. For $m_1, m_2, \dots, m_d \in \mathbb{N}$, let $[m_1, \dots, m_d]$ denote the least common multiple of m_1, \dots, m_d . The following is well-known (see the discussion before Proposition 4 in [4])

Theorem 1.19. *Let $p = (z_1, \dots, z_{n+2}) + \mathbb{Z} \in H(a)$. Assume that there are exactly k coordinates z_{j_1}, \dots, z_{j_k} all different from zero, $k \geq 2$. Then, $V(a)$ is non-singular at $\pi_a(p)$ if and only if*

$$\frac{[a_1, \dots, a_{n+2}]}{[a_{j_1}, \dots, a_{j_k}]} = \prod_{\ell \notin \{j_1, \dots, j_k\}} \frac{[a_1, \dots, a_{n+2}]}{[a_1, \dots, a_{\ell-1}, a_{\ell+1}, \dots, a_{n+2}]}$$

From Theorem 1.19, we deduce

Corollary 1.20. *Assume $n \geq 2$ and $(a_1, a_2, \dots, a_{n+2}) = (4b_1, 4b_2, 2b_3, 2b_4, \dots, 2b_{n+2})$, where $b_j \in \mathbb{Z}$, b_j is odd, $j = 1, \dots, n + 2$. Let $p = (0, 0, 1, i, 0, 0, \dots, 0) + \mathbb{Z} \in H(a)$. Then, $V(a)$ is singular at $\pi_a(p)$.*

Put

$$(1.20) \quad X := \{(z_1, \dots, z_{n+2}) \in \mathbb{C}^{n+2}; z_1^{a_1} + z_2^{a_2} + \dots + z_{n+2}^{a_{n+2}} = 0, \\ |z_1|^{2a_1} + |z_2|^{2a_2} + |z_3|^{2a_3} + \dots + |z_{n+2}|^{2a_{n+2}} = 1\}.$$

It is easy to see that X is a compact weakly pseudoconvex CR manifold of dimension $2n + 1$ with CR structure $T^{1,0}X := T^{1,0}\mathbb{C}^{n+2} \cap CTX$, where $T^{1,0}\mathbb{C}^{n+2}$ denotes the standard complex structure on \mathbb{C}^{n+2} . Let α be the least common multiple of a_1, \dots, a_{n+2} . Consider the following S^1 action on X :

$$(1.21) \quad S^1 \times X \rightarrow X, \\ e^{-i\theta} \circ (z_1, \dots, z_{n+2}) \rightarrow (e^{-i\frac{\alpha}{a_1}\theta} z_1, \dots, e^{-i\frac{\alpha}{a_{n+2}}\theta} z_{n+2}).$$

It is not difficult to check that the S^1 action is well-defined, locally free, CR and transversal. Moreover, it is straightforward to see that the quotient X/S^1 is equal to $V(a)$. Hence, X/S^1 is a complex orbifold. Moreover, from Corollary 1.20, we see that X/S^1 is singular if $n \geq 2$ and $(a_1, \dots, a_{n+2}) = (4b_1, 4b_2, 2b_3, 2b_4, \dots, 2b_{n+2})$, where $b_j \in \mathbb{Z}$, b_j is odd, $j = 1, 2, \dots, n + 2$. We now show that $(X, T^{1,0}X)$

is CR-isomorphic to a circle bundle associated with an orbifold line bundle over $X/S^1 = V(a)$. Let $L = (\widetilde{M(a)} \times \mathbb{C})/\equiv$, where $(z_1, \dots, z_{n+2}, \lambda) \equiv (\tilde{z}_1, \dots, \tilde{z}_{n+2}, \tilde{\lambda})$ if

$$\begin{aligned}\tilde{z}_j &= e^{\frac{m}{a_j}} z_j, \quad j = 1, \dots, n+2, \\ \tilde{\lambda} &= e^m \lambda,\end{aligned}$$

where $m \in \mathbb{Z}$. We can check that \equiv is an equivalence relation and L is a holomorphic line bundle over $H(a)$. The equivalence class of $(z_1, \dots, z_{n+2}, \lambda) \in \widetilde{M(a)} \times \mathbb{C}$ is denoted by $[(z_1, \dots, z_{n+2}, \lambda)]$. The complex 1-torus T_a action on L is given by the following:

$$(1.22) \quad \begin{aligned}T_a \times L &\rightarrow L, \\ (t + i\theta) \circ [(z_1, \dots, z_{n+2}, \lambda)] &\rightarrow [(e^{\frac{t+i\theta}{a_1}} z_1, \dots, e^{\frac{t+i\theta}{a_{n+2}}} z_{n+2}, e^{t-i\frac{\theta}{\alpha}} \lambda)],\end{aligned}$$

where α is the least common multiple of a_1, \dots, a_{n+2} . It is easy to see that the torus action (1.22) is well-defined and L/T_a is an orbifold line bundle over $H(a)/T_a = V(a)$. Let $\tau : L \rightarrow L/T_a$ be the natural projection and for $[(z_1, \dots, z_{n+2}, \lambda)] \in L$, we write $\tau([(z_1, \dots, z_{n+2}, \lambda)]) = [z_1, \dots, z_{n+2}, \lambda] + T_a$. It is easy to check that the pointwise norm

$$|[(z_1, \dots, z_{n+2}, \lambda)] + T_a|_{h^{L/T_a}}^2 := |\lambda|^2 \left(|z_1|^{2a_1} + |z_2|^{2a_2} + |z_3|^{2a_3} + \dots + |z_{2n+2}|^{2a_{n+2}} \right)^{-1}$$

is well-defined as a Hermitian fiber metric on L/T_a . The circle bundle $C(L/T_a)$ with respect to $(L/T_a, h^{L/T_a})$ is given by

$$(1.23) \quad \begin{aligned}C(L/T_a) &:= \left\{ v \in L/T_a; |v|_{h^{L/T_a}}^2 = 1 \right\} \\ &= \left\{ [(z_1, \dots, z_{n+2}, \lambda)] + T_a; |\lambda|^2 = |z_1|^{2a_1} + |z_2|^{2a_2} + |z_3|^{2a_3} + \dots + |z_{2n+2}|^{2a_{n+2}} \right\}.\end{aligned}$$

It is easy to see that $C(L/T_a)$ is a smooth CR manifold with CR structure

$$T^{1,0}C(L/T_a) := T^{1,0}L/T_a \cap \mathbb{C}TC(L/T_a),$$

where $T^{1,0}C(L/T_a)$ denotes the complex structure on L/T_a . Moreover, the orbifold line bundle $L/T_a \rightarrow V(a)$ satisfies the assumptions in Theorem 1.17 and Theorem 1.18.

The circle bundle $C(L/T_a)$ admits a nature S^1 action:

$$e^{-i\theta} \circ ([z_1, \dots, z_{n+2}, \lambda] + T_a) = [z_1, \dots, z_{n+2}, e^{-i\theta} \lambda] + T_a, \quad [z_1, \dots, z_{n+2}, \lambda] + T_a \in C(L/T_a).$$

Let $\Phi : C(L/T_a) \rightarrow X$ be the smooth map defined as follows. For every $[z_1, \dots, z_{n+2}, \lambda] + T_a \in C(L/T_a)$, there is a unique $(\hat{z}_1, \dots, \hat{z}_{n+2}) \in X$ such that

$$[z_1, \dots, z_{n+2}, \lambda] + T_a = [(\hat{z}_1, \dots, \hat{z}_{n+2}, 1)] + T_a.$$

Then, $\Phi([z_1, \dots, z_{n+2}, \lambda] + T_a) := (\hat{z}_1, \dots, \hat{z}_{n+2}) \in X$. It is straightforward to check that Φ is a CR embedding, globally one to one, onto, and the inverse $\Phi^{-1} : X \rightarrow C(L/T_a)$ is also a CR embedding. Moreover, we have

$$e^{-i\theta} \circ \Phi(x) = \Phi(e^{-i\theta} \circ x), \quad \forall x \in C(L/T_a).$$

We now consider another example. For simplicity, from now on, we assume that $a_1 = 1$. Then,

$$M(a) = \{(z_1, \dots, z_{n+2}) \in \mathbb{C}^{n+2}; z_1 = -z_2^{a_2} - \dots - z_{n+2}^{a_{n+2}}\}.$$

Fix $q = 2, 3, \dots, n+2$. Put

$$(1.24) \quad X_q := \left\{ (z_1, \dots, z_{n+2}) + \mathbb{Z} \in H(a); -|z_2|^{2a_2} - \dots - |z_q|^{2a_q} + |z_{q+1}|^{2a_{q+1}} + \dots + |z_{n+2}|^{2a_{n+2}} = 0 \right\}.$$

It is not difficult to check that X_q is a compact CR manifold of dimension $2n + 1$ with CR structure $T^{1,0}X_q := T^{1,0}H(a) \cap CTX_q$, where $T^{1,0}H(a)$ denotes the natural complex structure on $H(a)$. Let \tilde{a} be the least common multiple of a_1, \dots, a_q . Consider the following S^1 action on X_q :

$$(1.25) \quad \begin{aligned} & S^1 \times X_q \rightarrow X_q, \\ & e^{-i\theta} \circ ((z_1, \dots, z_{n+2}) + \mathbb{Z}) \\ & \rightarrow (e^{-i\tilde{a}\theta}(-z_2^{a_2} - \dots - z_q^{a_q}) - z_{q+1}^{a_{q+1}} - \dots - z_{n+2}^{a_{n+2}}, e^{-i\frac{\tilde{a}}{a_2}\theta} z_2, \dots, e^{-i\frac{\tilde{a}}{a_q}\theta} z_q, z_{q+1}, \dots, z_{n+2}) + \mathbb{Z}. \end{aligned}$$

It is straightforward to check that the S^1 action is well-defined, locally free, CR and transversal. Moreover, the CR manifold X_q is not pseudoconvex.

Similarly, put

$$(1.26) \quad \begin{aligned} \widetilde{X}_q := \{ & (z_1, \dots, z_{n+2}) + \mathbb{Z} \in H(a); -|z_2^{a_2} + z_3^{a_3}|^2 - |z_3|^{2a_3} - \dots - |z_q|^{2a_q} \\ & + |z_{q+1}|^{2a_{q+1}} + \dots + |z_{n+2}|^{2a_{n+2}} = 0\}. \end{aligned}$$

It is not difficult to check that \widetilde{X}_q is a compact CR manifold of dimension $2n + 1$ with CR structure $T^{1,0}\widetilde{X}_q := T^{1,0}H(a) \cap CT\widetilde{X}_q$. Let \tilde{a} be the least common multiple of a_1, \dots, a_q . Consider the following S^1 action on \widetilde{X}_q :

$$(1.27) \quad \begin{aligned} & S^1 \times \widetilde{X}_q \rightarrow \widetilde{X}_q, \\ & e^{-i\theta} \circ ((z_1, \dots, z_{n+2}) + \mathbb{Z}) \\ & \rightarrow (e^{-i\tilde{a}\theta}(-z_2^{a_2} - \dots - z_q^{a_q}) - z_{q+1}^{a_{q+1}} - \dots - z_{n+2}^{a_{n+2}}, e^{-i\frac{\tilde{a}}{a_2}\theta} z_2, \dots, e^{-i\frac{\tilde{a}}{a_q}\theta} z_q, z_{q+1}, \dots, z_{n+2}) + \mathbb{Z}. \end{aligned}$$

It is straightforward to check that the S^1 action is well-defined, locally free, CR and transversal. Let $f(x) \in C^\infty(\widetilde{X}_q)$ with $Tf = 0$, where T is the global real vector field induced by the S^1 action (1.27). Let $Z_1, \dots, Z_n \in C^\infty(\widetilde{X}_q, T^{1,0}\widetilde{X}_q)$ be a basis for $T^{1,0}\widetilde{X}_q$. Put

$$H^{1,0}\widetilde{X}_q := \{Z_j + Z_j(f)T; j = 1, 2, \dots, n\}.$$

We can check that $H^{1,0}\widetilde{X}_q$ is a CR structure and the S^1 action (1.27) is locally free, CR and transversal with respect to this new CR structure $H^{1,0}\widetilde{X}_q$.

1.6. Proof of Theorem 1.2. In this section, we will prove Theorem 1.2 and we will use the same notations as in Theorem 1.2. Let s be a local trivializing section of L defined on some open set U of M , $|s|_h^2 = e^{-2\phi}$. Let $z = (z_1, \dots, z_n)$ be holomorphic coordinates on U . We identify U with an open set of \mathbb{C}^n . We have the local diffeomorphism:

$$(1.28) \quad \tau : U \times]-\varepsilon_0, \varepsilon_0[\rightarrow X, (z, \theta) \mapsto e^{-\phi(z)} s^*(z) e^{-i\theta}, \quad 0 < \varepsilon_0 \leq \pi.$$

Put $D = U \times]-\varepsilon_0, \varepsilon_0[$. We will identify D with some open set in X and we call D a canonical coordinate patch with respect to local trivializing section s and (z, θ) canonical coordinates of D . It is convenient to work with the local coordinates (z, θ) . In terms of these coordinates, it is straightforward to see that $T = \frac{\partial}{\partial \theta}$. Recall that T is the global real vector field induced by the S^1 action. Moreover, it is straightforward to check that on (z, θ) coordinates, we have

$$(1.29) \quad \begin{aligned} T^{1,0}X &= \left\{ \frac{\partial}{\partial z_j} - i \frac{\partial \phi}{\partial z_j}(z) \frac{\partial}{\partial \theta}; j = 1, 2, \dots, n \right\}, \\ T^{0,1}X &= \left\{ \frac{\partial}{\partial \bar{z}_j} + i \frac{\partial \phi}{\partial \bar{z}_j}(z) \frac{\partial}{\partial \theta}; j = 1, 2, \dots, n \right\}, \end{aligned}$$

and

$$(1.30) \quad T^{*1,0}X = \{dz_j; j = 1, 2, \dots, n\}, \quad T^{*0,1}X = \{d\bar{z}_j; j = 1, 2, \dots, n\}.$$

Let $f(z) \in \Omega^{0,q}(D)$. From (1.30), we may identify f with an element in $\Omega^{0,q}(U)$. Let $A_m^{(q)} : \Omega_m^{0,q}(X) \rightarrow \Omega^{0,q}(M, L^m)$ be the map defined as follows. Let $u \in \Omega_m^{0,q}(X)$. $A_m^{(q)}u$ is an element in $\Omega^{0,q}(M, L^m)$ such that on a canonical coordinate patch $D = U \times]-\varepsilon_0, \varepsilon_0[$ with respect to local trivializing section s , if we write $u(z, \theta) = e^{-im\theta}\hat{u}(z)$ on D , where $\hat{u}(z) \in \Omega^{0,q}(U)$, then on U , $A_m^{(q)}u = s^m(z)e^{m\phi(z)}\hat{u}(z) \in \Omega^{0,q}(U, L^m)$. We now check that $A_m^{(q)}$ is well-defined. Let s and s_1 be local trivializing sections of L on an open set U . Let $(z, \theta) \in \mathbb{C}^n \times \mathbb{R}$ and $(z, \eta) \in \mathbb{C}^n \times \mathbb{R}$ be canonical coordinates of D with respect to s and s_1 respectively, where $D = U \times]-\varepsilon_0, \varepsilon_0[$, $0 < \varepsilon_0, \varepsilon_1 < \pi$. Fix $u \in \Omega_m^{0,q}(X)$. Set $|s|_{hL}^2 = e^{-2\phi}$ and $|s_1|_{hL}^2 = e^{-2\phi_1}$. We write

$$\begin{aligned} u &= e^{-im\theta}\hat{u}(z) \quad \text{on } D, \\ u &= e^{-im\eta}\hat{u}_1(z) \quad \text{on } D. \end{aligned}$$

From the construction of $A_m^{(q)}$, to check that $A_m^{(q)}$ is well-defined, we need to check that

$$(1.31) \quad s^m(z)e^{m\phi(z)}\hat{u}(z) = s_1^m(z)e^{m\phi_1(z)}\hat{u}_1(z), \quad \forall z \in U.$$

Let $s_1 = gs$, where $g \in C^\infty(U)$ is a non-zero holomorphic function. We have

$$|s_1|_{hL}^2 = e^{-2\phi_1} = |g|^2 |s|_{hL}^2 = e^{2\log|g| - 2\phi}.$$

Thus,

$$(1.32) \quad \phi_1 = \phi - \log|g|.$$

We claim that

$$(1.33) \quad \text{If } (z, \theta) = (z, \eta), \text{ then } e^{-i\theta} \left(\frac{g(z)}{\bar{g}(z)}\right)^{\frac{1}{2}} = e^{-i\eta}.$$

Proof of the claim. From (1.32), we have

$$\begin{aligned} s^*(z)e^{-i\theta - \phi(z)} &= s_1^*(z)g(z)e^{-i\theta - \phi(z)} \\ (1.34) \quad &= s_1^*(z)g(z)e^{-i\theta - \phi_1(z) - \log|g(z)|} \\ &= s_1^*(z) \left(\frac{g(z)}{\bar{g}(z)}\right)^{\frac{1}{2}} e^{-i\theta - \phi_1(z)}. \end{aligned}$$

Since $(z, \theta) = (z, \eta)$, we have

$$(1.35) \quad s^*(z)e^{-i\theta - \phi(z)} = s_1^*(z)e^{-i\eta - \phi_1(z)}.$$

From (1.34) and (1.35), we deduce that $e^{-i\theta} \left(\frac{g(z)}{\bar{g}(z)}\right)^{\frac{1}{2}} = e^{-i\eta}$. The claim follows. \square

From (1.33), we have

$$e^{-im\theta}\hat{u}(z) = e^{-im\eta} \left(\frac{g(z)}{\bar{g}(z)}\right)^{-\frac{m}{2}} \hat{u}(z) = e^{-im\eta}\hat{u}_1(z).$$

Thus,

$$(1.36) \quad \left(\frac{g(z)}{\bar{g}(z)}\right)^{-\frac{m}{2}} \hat{u}(z) = \hat{u}_1(z).$$

From (1.32) and (1.36), we have

$$\begin{aligned} s^m(z)e^{m\phi(z)}\hat{u}(z) &= s_1^m(z)g^{-m}(z)e^{m\phi_1(z) + m\log|g(z)|}\hat{u}(z) \\ (1.37) \quad &= s_1^m(z) \left(\frac{\bar{g}(z)}{g(z)}\right)^{\frac{m}{2}} e^{m\phi_1(z)}\hat{u}(z) \\ &= s_1^m(z)e^{m\phi_1(z)}\hat{u}_1(z). \end{aligned}$$

Hence, (1.31) holds and $A_m^{(q)} : \Omega_m^{0,q}(X) \rightarrow \Omega^{0,q}(M, L^m)$ is well-defined. Moreover, it is easy to check that $A_m^{(q)} : \Omega_m^{0,q}(X) \rightarrow \Omega^{0,q}(M, L^m)$ is a bijection. We omit the detail.

We now prove that $\bar{\partial}A_m^{(q)} = A_m^{(q+1)}\bar{\partial}_b$. Let $D = U \times] - \varepsilon_0, \varepsilon_0[$, $0 < \varepsilon_0 < \pi$, be a canonical coordinate patch with respect to local trivializing section s and let (z, θ) be canonical coordinates of D . Let $u \in \Omega_m^{0,q}(X)$. On D , we write $u(z, \theta) = e^{-im\theta}\hat{u}(z)$, where $\hat{u}(z) \in \Omega^{0,q}(U)$. In view of (1.29) and (1.30), we can check that on D ,

$$(1.38) \quad \bar{\partial}_b u = \bar{\partial}_b(e^{-im\theta}\hat{u}) = \sum_{j=1}^n e^{-im\theta} d\bar{z}_j \wedge \left(\frac{\partial \hat{u}}{\partial \bar{z}_j}(z) + m \frac{\partial \phi}{\partial \bar{z}_j}(z) \hat{u}(z) \right).$$

From (1.38) and by the definition of $A_m^{(q+1)}$, it is easy to see that

$$(1.39) \quad \begin{aligned} A_m^{(q+1)}(\bar{\partial}_b u) &= s^m(z) e^{m\phi(z)} \sum_{j=1}^n d\bar{z}_j \wedge \left(\frac{\partial \hat{u}}{\partial \bar{z}_j}(z) + m \frac{\partial \phi}{\partial \bar{z}_j}(z) \hat{u}(z) \right) \\ &= s^m(z) \bar{\partial}(e^{m\phi(z)} \hat{u}(z)) \quad \text{on } U. \end{aligned}$$

Thus, $\bar{\partial}A_m^{(q)} = A_m^{(q+1)}\bar{\partial}_b$. Theorem 1.2 follows.

1.7. The idea of the proof of Theorem 1.3. In this section, we will give an outline of the idea of the proof of Theorem 1.3. We refer to Section 2.2 and Section 2.3 for some notations and terminology used here. To prove Theorem 1.3, we need to construct heat kernel for Kohn Laplacian associated to m -th S^1 Fourier coefficient.

1.7.1. Global difficulties. For simplicity, we assume that E is trivial and X is CR Kähler, that is, there is a closed form $\Theta \in C^\infty(X, T^{*1,1}X)$ such that $\Theta(Z, \bar{Z}) > 0$, for all $Z \in C^\infty(X, T^{1,0}X)$. Θ induces a Hermitian metric $\langle \cdot | \cdot \rangle$ on $\mathbb{C}TX$. Let $\bar{\partial}_b^*$ be the adjoint of $\bar{\partial}_b$ with respect to $(\cdot | \cdot)$. We fix $m \in \mathbb{Z}$. Then, $\bar{\partial}_{b,m}^* = \bar{\partial}_b^* : \Omega_m^{0,q+1}(X) \rightarrow \Omega_m^{0,q}(X)$. Consider

$$\begin{aligned} D_{b,m} &:= \bar{\partial}_{b,m} + \bar{\partial}_{b,m}^* : \Omega_m^{0,+}(X) \rightarrow \Omega_m^{0,-}(X), \\ D_{b,m} &:= \bar{\partial}_{b,m} + \bar{\partial}_{b,m}^* : \Omega_m^{0,-}(X) \rightarrow \Omega_m^{0,+}(X), \end{aligned}$$

and let

$$\begin{aligned} \square_{b,m}^+ &:= D_{b,m}^2 : \Omega_m^{0,+}(X) \rightarrow \Omega_m^{0,+}(X), \\ \square_{b,m}^- &:= D_{b,m}^2 : \Omega_m^{0,-}(X) \rightarrow \Omega_m^{0,-}(X), \end{aligned}$$

where $\Omega_m^{0,+}(X)$ and $\Omega_m^{0,-}(X)$ are given by (3.21). Let $L^{2,+}(X)$, $L^{2,-}(X)$, $L_m^{2,+}(X)$ and $L_m^{2,-}(X)$ be as in the discussion after (3.22). We extend $\square_{b,m}^+$ and $\square_{b,m}^-$ to $L_m^{2,+}(X)$ and $L_m^{2,-}(X)$ in the standard way and we will show in Theorem 3.5 that $\text{Spec } \square_{b,m}^+$ and $\text{Spec } \square_{b,m}^-$ are discrete subsets of $[0, \infty[$ and for every $\nu \in \text{Spec } \square_{b,m}^+$, $\mu \in \text{Spec } \square_{b,m}^-$, ν and μ are eigenvalues of $\square_{b,m}^+$ and $\square_{b,m}^-$ respectively. For every $\nu \in \text{Spec } \square_{b,m}^+$, let $\{f_1^\nu, \dots, f_{d_\nu}^\nu\}$ be an orthonormal frame for the eigenspace of $\square_{b,m}^+$ with eigenvalue ν . The heat kernel $e^{-t\square_{b,m}^+}(x, y)$ is given by

$$(1.40) \quad e^{-t\square_{b,m}^+}(x, y) = \sum_{\nu \in \text{Spec } \square_{b,m}^+} \sum_{j=1}^{d_\nu} e^{-\nu t} f_j^\nu(x) \wedge (f_j^\nu(y))^\dagger,$$

where $f_j^\nu(x) \wedge (f_j^\nu(y))^\dagger$ denotes the linear map:

$$\begin{aligned} f_j^\nu(x) \wedge (f_j^\nu(y))^\dagger &: \oplus_{q \in \{0,1,2,\dots,n\}} T_y^{*0,q} X \rightarrow T_x^{*0,+} X, \\ u(y) &\in \oplus_{q \in \{0,1,2,\dots,n\}} T_y^{*0,q} X \rightarrow f_j^\nu(x) \langle u(y) | f_j^\nu(y) \rangle \in T_x^{*0,+} X. \end{aligned}$$

Similarly, we can define $e^{-t\bar{\square}_{b,m}^-}(x, y)$ as (1.40). We will show in Theorem 4.10 (see also Remark 4.11) that we have *CR McKean-Singer type formula*: for every $t > 0$,

$$(1.41) \quad \sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X) = \int_X \left(\text{Tr} e^{-t\bar{\square}_{b,m}^+}(x, x) - \text{Tr} e^{-t\bar{\square}_{b,m}^-}(x, x) \right) dv_X.$$

By using *CR McKean-Singer type formula*, the proof of Theorem 1.3 is reduced to determine the small t behaviour of the function $\left(\text{Tr} e^{-t\bar{\square}_{b,m}^+}(x, x) - \text{Tr} e^{-t\bar{\square}_{b,m}^-}(x, x) \right)$. Let $e^{-t\bar{\square}_{b,m}^+} : \Omega^{0,+}(X) \rightarrow \Omega_m^{0,+}(X)$ be the operator with distribution kernel $e^{-t\bar{\square}_{b,m}^+}(x, y)$. From (1.40), it is easy to see that $e^{-t\bar{\square}_{b,m}^+}$ satisfies

$$(1.42) \quad \frac{\partial e^{-t\bar{\square}_{b,m}^+}}{\partial t} + \bar{\square}_{b,m}^+ e^{-t\bar{\square}_{b,m}^+} = 0$$

and

$$(1.43) \quad e^{-t\bar{\square}_{b,m}^+}|_{t=0} = Q_m^+,$$

where $Q_m^+ : L^{2,+}(X) \rightarrow L_m^{2,+}(X)$ is the orthogonal projection. The main *difficulty* is that the initial condition (1.43) is a *projection* rather than an *identity operator*. This is indeed a new point beyond the tradition, arising from the situation that we deal with part of the L^2 space (eigenspaces) rather than the whole L^2 space. This initial condition (1.43) causes that the heat kernels $e^{-t\bar{\square}_{b,m}^+}(x, y)$ and $e^{-t\bar{\square}_{b,m}^-}(x, y)$ do not have *standard expansions* with respect to t . For the better understanding, let's assume that X is an orbifold circle bundle of an orbifold line bundle L over a Kähler orbifold M . Then, X is a CR manifold with a locally free S^1 action. We can repeat the proof of Theorem 1.2 (see Section 1.6) and conclude that there is a bijection map

$$\begin{aligned} A_m^+ &: \Omega_m^{0,+}(X) \rightarrow \Omega^{0,+}(M, L^m), \\ A_m^- &: \Omega_m^{0,-}(X) \rightarrow \Omega^{0,-}(M, L^m), \end{aligned}$$

such that $A_m^+ \bar{\partial}_b = \bar{\partial} A_m^+$. Let $\bar{\square}_m^+$ be the Kodaira Laplacian with values in $T^{*,0,+}M \otimes L^m$ and let $e^{-t\bar{\square}_m^+}$ be the associated heat operator. Consider $B_m(t) := (A_m^+)^{-1} \circ e^{-t\bar{\square}_m^+} \circ A_m^+$. We can check that $B'_m(t) + \bar{\square}_{b,m}^+ B_m(t) = 0$ and $B_m(0) = I$ on $\Omega_m^{0,+}(X)$. But $B_m(t)$ is *not* the heat operator $e^{-t\bar{\square}_{b,m}^+}$. A trivial understanding is that, they are defined on different spaces, as $B_m(t)$ is on $\Omega_m^{0,+}(X)$ while the heat kernel $e^{-t\bar{\square}_{b,m}^+}$ is on the whole $\Omega^{0,+}(X)$. Moreover, it is straightforward to check that

$$(1.44) \quad e^{-t\bar{\square}_{b,m}^+} = B_m(t) \circ Q_m^+ = Q_m^+ \circ B_m(t) \circ Q_m^+.$$

Let $B_m(t, x, y)$ and $e^{-t\bar{\square}_{b,m}^+}(x, y)$ be the distribution kernels of $B_m(t)$ and $e^{-t\bar{\square}_{b,m}^+}$ respectively. From (1.44), we can check that (see (2.5) and (4.26))

$$(1.45) \quad e^{-t\bar{\square}_{b,m}^+}(x, y) = \frac{1}{(2\pi)^2} \int B_m(t, e^{-i\theta} \circ x, e^{-iu} \circ y) e^{im(\theta-u)} d\theta du.$$

For $x \notin X_p$, from (1.45), it is difficult to understand the asymptotic behavior of $e^{-t\bar{\square}_{b,m}^+}(x, x)$ from the asymptotic expansion of $B_m(t)$. For $x \in X_p$, from (1.45), $e^{-t\bar{\square}_{b,m}^+}(x, x)$ admits a standard expansion as $t \rightarrow 0^+$:

$$(1.46) \quad e^{-t\bar{\square}_{b,m}^+}(x, x) \sim t^{-n} a_n^+(x) + t^{-(n-1)} a_{n-1}^+(x) + \dots$$

Another problem is that the expansion (1.46) converges only *locally uniformly* on X_p . It seems to us that even in orbifold circle bundle cases, to understand the asymptotic behavior of the heat operator $e^{-t\bar{\square}_{b,m}^+}$, we still need to work directly on CR manifold X . In this paper, we give a construction which is independent of the use of orbifold geometry and more adapted to CR geometry. Our choice of cut-off functions for the patching of local heat kernels is adapted to BRT trivializations for our CR manifolds. By this construction it becomes natural that our computation of local density need not

introduce orbifold contributions. It does not seem obvious how Kawasaki's result can bypass the orbifold contributions in the case where the complex orbifold has a circle bundle whose total space is globally smooth, unless the complex orbifold is itself smooth.

In view of global difficulties above, it is not immediate that solely using the global argument can arrive at a detailed understanding of the heat kernel. Moreover, our CR manifold X is not necessary to be an orbifold circle bundle of a complex orbifold. We thus turn our attention to work on it locally. The main ingredient here is *BRT trivialization*, which is first treated by Baouendi, Rothschild and Treves [1] in a more general context.

1.7.2. *Transition to local situation.* Let $B := (D, (z, \theta), \varphi)$ be a BRT trivialization (see Theorem 2.8). We may assume that $D = U \times]-\varepsilon, \varepsilon[$, where $\varepsilon > 0$ and U is an open set of \mathbb{C}^n . Consider $L \rightarrow U$ be a trivial line bundle with non-trivial Hermitian fiber metric $|1|_{h^L}^2 = e^{-2\varphi}$, where $\varphi \in C^\infty(D, \mathbb{R})$ is as in Theorem 2.8. Let $(L^m, h^{L^m}) \rightarrow U$ be the m -th power of (L, h^L) . Θ induces a Kähler form Θ_U on the complex manifold U and let $\langle \cdot, \cdot \rangle$ be the Hermitian metric on CTU induced by Θ_U and let $(\cdot, \cdot)_m$ be the L^2 inner product on $\Omega^{0,*}(U, L^m)$ induced by $\langle \cdot, \cdot \rangle$ and h^{L^m} . Let

$$\bar{\partial} : \Omega^{0,q}(U, L^m) \rightarrow \Omega^{0,q+1}(U, L^m), \quad q = 0, 1, 2, \dots, n-1,$$

be the Cauchy-Riemann operator and let

$$\bar{\partial}^{*,m} : \Omega^{0,q+1}(U, L^m) \rightarrow \Omega^{0,q}(U, L^m), \quad q = 0, 1, 2, \dots, n-1,$$

be the formal adjoint of $\bar{\partial}$ with respect to $(\cdot, \cdot)_m$. Put

$$(1.47) \quad \begin{aligned} D_{B,m} &:= \bar{\partial} + \bar{\partial}^{*,m} : \Omega^{0,+}(U, L^m) \rightarrow \Omega^{0,-}(U, L^m), \\ D_{B,m} &:= \bar{\partial} + \bar{\partial}^{*,m} : \Omega^{0,-}(U, L^m) \rightarrow \Omega^{0,+}(U, L^m). \end{aligned}$$

Put

$$(1.48) \quad \begin{aligned} \square_{B,m}^+ &:= D_{B,m}^2 : \Omega^{0,+}(U, L^m) \rightarrow \Omega^{0,+}(U, L^m), \\ \square_{B,m}^- &:= D_{B,m}^2 : \Omega^{0,-}(U, L^m) \rightarrow \Omega^{0,-}(U, L^m). \end{aligned}$$

Let $u \in \Omega_m^{0,+}(X)$, $v \in \Omega_m^{0,-}(X)$. On D , we write $u(z, \theta) = e^{-im\theta} \tilde{u}(z)$, $v(z, \theta) = e^{-im\theta} \tilde{v}(z)$, $\tilde{u}(z) \in \Omega^{0,+}(U, L^m)$, $\tilde{v}(z) \in \Omega^{0,-}(U, L^m)$. Then, we will show in Lemma 5.1 that

$$(1.49) \quad \begin{aligned} e^{-m\varphi} \square_{B,m}^+ (e^{m\varphi} \tilde{u}) &= e^{im\theta} \square_{b,m}^+ (u), \\ e^{-m\varphi} \square_{B,m}^- (e^{m\varphi} \tilde{v}) &= e^{im\theta} \square_{b,m}^- (v). \end{aligned}$$

From (1.49), we can guess that the heat kernel $e^{-t\square_{b,m}^+}(x, y)$ locally should be

$$(1.50) \quad e^{-m\varphi(z) - im\theta} e^{-t\square_{B,m}^+}(x, y) e^{m\varphi(w) + im\eta}.$$

By using the BRT trivializations we localize the construction and are inspired to obtain a *localize* heat kernel on these particular charts. We would like to patch up these local heat kernels. We will see that the standard construction for the heat kernel does not work in our case. Assume that $X = D_1 \cup D_2 \cup \dots \cup D_N$, where $B_j := (D_j, (z, \theta), \varphi_j)$ is a BRT trivialization, for each j . We may assume that for each j , $D_j = U_j \times]-\delta_j, \delta_j[\subset \mathbb{C}^n \times \mathbb{R}$, $\delta_j > 0$, $\tilde{\delta}_j > 0$, U_j is an open set in \mathbb{C}^n . Let $\chi_j, \tilde{\chi}_j \in C_0^\infty(D_j)$, $j = 1, 2, \dots, N$. Put

$$(1.51) \quad \begin{aligned} \mathcal{A}_m(t) &= \sum_{j=1}^N \chi_j(x) \left(e^{-m\varphi_j(z) - im\theta} e^{-t\square_{B_j,m}^+}(z, w) e^{m\varphi_j(w) + im\eta} \right) \tilde{\chi}_j(y), \\ \mathcal{P}_m(t) &= Q_m^+ \circ \mathcal{A}_m(t) \circ Q_m^+. \end{aligned}$$

We hope that $\mathcal{P}_m(0) = Q_m^+$ and $\mathcal{P}'_m(t) + \square_{b,m}^+ \mathcal{P}_m(t)$ is small as $t \rightarrow 0^+$.

1.7.3. *Local difficulties.* A necessary condition for $\mathcal{P}_m(0) = Q_m^+$ is

$$(1.52) \quad \sum_{j=1}^N \chi_j(x) \int \tilde{\chi}_j(z, \eta) d\eta = 1.$$

Unfortunately, it is unclear to see whether (1.52) can be enough to reach the desired heat kernel. A reasonable choice for the cut-off functions χ_j , $\tilde{\chi}_j$ is the following: take $\chi_j(z, \theta) \in C_0^\infty(D_j)$, $j = 1, 2, \dots, N$, with $\sum_{j=1}^N \chi_j = 1$ on X and for each $j = 1, 2, \dots, N$, take $\tau_j(z) \in C_0^\infty(U_j)$ with $\tau_j(z) = 1$ if $(z, \theta) \in \text{Supp } \chi_j$ and take $\sigma_j \in C_0^\infty(\cdot - \delta_j, \tilde{\delta}_j)$ with $\int \sigma_j(\eta) d\eta = 1$. Then, $\chi_j(x)$, $\tilde{\chi}_j(y) = \tau_j(w)\sigma_j(\eta)$, $j = 1, 2, \dots, N$, satisfy (1.52). We can check that $\mathcal{P}_m(0) = Q_m^+$ and

$$(1.53) \quad \mathcal{P}'_m(t) + \square_{b,m}^+ \mathcal{P}_m(t) = Q_m^+ \circ \mathcal{R}_m(t) \circ Q_m^+,$$

where

$$(1.54) \quad \mathcal{R}_m(t) = \sum_{j=1}^N \sum_{\ell=1}^k L_{\ell,j} \left(\chi_j(x) e^{-m\varphi_j(z) - im\theta} \right) P_{\ell,j} \left(e^{-t\square_{B_j,m}^+}(z, w) \right) e^{m\varphi_j(w) + im\eta} \tilde{\chi}_j(y),$$

where $L_{\ell,j}$ is a partial differential operator of order ≤ 2 and ≥ 1 , for all ℓ, j , $P_{\ell,j}$ is a partial differential operator of order 1 acting on x , for all ℓ, j . Since $\left| e^{-t\square_{B_j,m}^+}(z, w) \right| \sim e^{-\frac{|z-w|^2}{t}}$, it could happen that

$$(1.55) \quad P_{\ell,j} \left(e^{-t\square_{B_j,m}^+}(z, w) \right) \sim e^{-\frac{|z-w|^2}{t}} \frac{|z-w|}{t}.$$

From (1.55), we want $\mathcal{P}'_m(t) + \square_{b,m}^+ \mathcal{P}_m(t)$ to be small, we need

$$(1.56) \quad L_{\ell,j} \left(\chi_j(x) e^{-m\varphi_j(z) - im\theta} \right) e^{m\varphi_j(w) + im\eta} \tilde{\chi}_j(y) = 0 \quad \text{if } z \text{ is close to } w \text{ } (|z-w| \lesssim \sqrt{t}).$$

Since χ_j is not constant on $\text{Supp } \tilde{\chi}_j$, it is difficult to see if (1.56) holds. To overcome the difficulty, instead of the original equation, we make use of an *adjoint* version of the original equation. It turns out that the aforementioned difficulty can be avoided, so that a global heat kernel can be obtained by the process of patching in this new context. Fix $j = 1, 2, \dots, N$. There is $A_{B_j,+}(t, z, w) \in C^\infty(\mathbb{R}_+ \times U_j \times U_j, (T_w^{*0,+}U \boxtimes T_z^{*0,+}U))$ such that

$$(1.57) \quad \begin{aligned} \lim_{t \rightarrow 0^+} A_{B_j,+}(t) &= I \text{ in } \mathcal{D}'(U, T^{*0,+}U), \\ A'_{B_j,+}(t)u + A_{B_j,+}(t)(\square_{B_j,m}^+ u) &= 0, \quad \forall u \in \Omega_0^{0,+}(U), \quad \forall t > 0, \end{aligned}$$

and $A_{B_j,+}(t, z, w)$ admits an asymptotic expansion as $t \rightarrow 0^+$ (see (5.17)). Put

$$(1.58) \quad H_j(t, x, y) = \chi_j(x) e^{-m\varphi_j(z) - im\theta} A_{B_j,+}(t, z, w) e^{m\varphi_j(w) + im\eta} \tilde{\chi}_j(y).$$

It should be noticed that there is no guarantee that $H_j(t)$ still stay in the eigenspace $\Omega_m^{0,+}(X)$. We thus set

$$(1.59) \quad \Gamma(t) := \sum_{j=1}^N Q_m^+ \circ H_j(t) \circ Q_m^+ : \Omega_m^{0,+}(X) \rightarrow \Omega_m^{0,+}(X).$$

In Section 5.2, we will prove that $\Gamma(t)$ is a *successive approximation* for the adjoint of $e^{-t\square_{b,m}^+}$ (see Theorem 5.8) and since $e^{-t\square_{b,m}^+}$ is self-adjoint, $\Gamma(t)$ is a *successive approximation* for $e^{-t\square_{b,m}^+}$. More precisely, we have (see Theorem 5.10)

$$(1.60) \quad \begin{aligned} \left\| e^{-t\square_{b,m}^+}(x, y) - \Gamma(t, x, y) \right\|_{C^0(X \times X)} &\leq e^{-\frac{\epsilon_0}{t}}, \quad \forall t \in (0, \epsilon), \\ \Gamma(t, x, y) &\sim t^{-n} a_n^+(t, x, y) + t^{-n+1} a_{n-1}^+(t, x, y) + \dots \quad \text{as } t \rightarrow 0^+, \end{aligned}$$

where $\epsilon > 0$, $\epsilon_0 > 0$ are constants and there is a constant $C > 0$ such that $|a_s^+(t, x, y)| \leq C$, for every $t > 0$, $(x, y) \in X \times X$ and every $s = n, n-1, \dots$. Hence,

$$(1.61) \quad e^{-t\Box_{b,m}^+} \sim t^{-n} a_n^+(t, x, y) + t^{-n+1} a_{n-1}^+(t, x, y) + \dots \quad \text{as } t \rightarrow 0^+.$$

Similarly, we have

$$(1.62) \quad e^{-t\Box_{b,m}^-} \sim t^{-n} a_n^-(t, x, y) + t^{-n+1} a_{n-1}^-(t, x, y) + \dots \quad \text{as } t \rightarrow 0^+.$$

From (1.41), (1.61) and (1.62), we conclude that

$$(1.63) \quad \sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, E) = \lim_{t \rightarrow 0^+} \int_X \sum_{\ell=0}^n t^{-\ell} \left(\text{Tr } a_\ell^+(t, x, x) - \text{Tr } a_\ell^-(t, x, x) \right) dv_X(x).$$

It is indeed that we have already had a heat kernel which is put in the disguise of the spectral geometry (4.21). But we found no argument in the literature to claim that (4.21) shall have the asymptotic estimates as (1.61) and (1.62), which are essential for computation of the local density needed by our proof of the index theorem. The somewhat lengthy part of our reconstruction of the heat kernel beyond its spectral realization becomes indispensable as far as our approach is concerned.

1.7.4. Completion by evaluating of local density and by using Spin^c structure. From (1.63), to complete the proof of Theorem 1.3, we need to understand the small t behaviour of the local density $\sum_{\ell=0}^n t^{-\ell} \left(\text{Tr } a_\ell^+(t, x, x) - \text{Tr } a_\ell^-(t, x, x) \right)$. Let's come back to local situation. Fix $x_0 \in X_p$. Let $B_j = (D_j, (z, \theta), \varphi_j)$, $j = 1, 2, \dots, N$, be BRT trivializations as before. Fix $j = 1, 2, \dots, N$. Assume that $e^{-i\theta_0} \circ x_0 \in D_j$ for some $\theta_0 \in]-\pi, \pi]$ and suppose that $e^{-i\theta_0} \circ x_0 = (z_j, 0) \in D_j$. By using some properties of BRT trivializations, our choice of cut-off functions and some tricks, we will show in Theorem 6.1 that

$$(1.64) \quad \begin{aligned} & (Q_m^+ \circ H_j \circ Q_m^+)(t, x_0, x_0) \\ &= \left(\sum_{s=1}^p e^{im\frac{2\pi}{p}(s-1)} \right) \left(\sum_{k=1}^p e^{-im\frac{2\pi}{p}(k-1)} \right) \int_{-\frac{\pi}{p}}^{\frac{\pi}{p}} \chi_j(e^{-i\theta} \circ x_0) A_{B_j,+}(t, z_j, z_j) d\theta, \end{aligned}$$

where $A_{B_j,+}(t, z, w)$ is as in (1.57). Hence

$$(1.65) \quad \begin{aligned} & \sum_{\ell=0}^n t^{-\ell} \left(\text{Tr } a_\ell^+(t, x, x) - \text{Tr } a_\ell^-(t, x, x) \right) \\ &= \left(\sum_{s=1}^p e^{im\frac{2\pi}{p}(s-1)} \right) \left(\sum_{k=1}^p e^{-im\frac{2\pi}{p}(k-1)} \right) \\ & \quad \times \sum_{j=1}^N \int_{-\frac{\pi}{p}}^{\frac{\pi}{p}} \chi_j(e^{-i\theta} \circ x_0) \left(\text{Tr } A_{B_j,+}(t, z_j, z_j) - \text{Tr } A_{B_j,-}(t, z_j, z_j) \right) d\theta + O(t). \end{aligned}$$

By using the *rescaling technique* in [2] and [8], we can show that for each $j = 1, 2, \dots, N$, we have

$$(1.66) \quad \begin{aligned} & \left(\text{Tr } A_{B_j,+}(t, z, z) - \text{Tr } A_{B_j,-}(t, z, z) \right) dv_{U_j}(z) \\ &= [\text{Td}(\nabla^{T^{1,0}U_j}, T^{1,0}U_j) \wedge \text{ch}(\nabla^{L^m}, L^m)]_{2n}(z) + O(t), \quad \forall z \in U_j, \end{aligned}$$

where dv_{U_j} is the induced volume form on U_j , $A_{B_j,+}(t, z, w)$ is as in (1.57), $\text{Td}(\nabla^{T^{1,0}U_j}, T^{1,0}U_j)$ denotes the representative of the Todd class of $T^{1,0}U_j$ induced by the given Hermitian metric on $T^{1,0}U_j$ and $\text{ch}(\nabla^{L^m}, L^m)$ denotes the representative of the Chern character induced by the Hermitian metric

h^{L^m} . In Section 2.3, we will introduce *tangential characteristic classes*, *tangential Chern character* and *tangential Todd class* on CR manifolds with S^1 action and we have

$$(1.67) \quad \frac{[\mathrm{Td}(\nabla^{T^{1,0}U_j}, T^{1,0}U_j) \wedge \mathrm{ch}(\nabla^{L^m}, L^m)]_{2n}(z_j)}{dv_{U_j}(z_j)} = \frac{[\mathrm{Td}_b(\nabla^{T^{1,0}X}, T^{1,0}X) \wedge e^{-m\frac{d\omega_0}{2\pi}} \wedge \omega_0]_{2n+1}(x_0)}{dv_X(x_0)},$$

where $\mathrm{Td}_b(\nabla^{T^{1,0}X}, T^{1,0}X)$ denotes the representative of the tangential Todd class of $T^{1,0}X$ induced by the given Hermitian metric (see (2.11)). From (1.65), (1.66) and (1.67), we deduce that

$$(1.68) \quad \begin{aligned} & \sum_{\ell=0}^n t^{-\ell} \left(\mathrm{Tr} a_\ell^+(t, x, x) - \mathrm{Tr} a_\ell^-(t, x, x) \right) dv_X(x) \\ &= \sum_{s=1}^p e^{\frac{2\pi(s-1)}{p}mi} \left[\mathrm{Td}_b(\nabla^{T^{1,0}X}, T^{1,0}X) \wedge e^{-m\frac{d\omega_0}{2\pi}} \wedge \omega_0 \right]_{2n+1}(x) + O(t), \quad \forall x \in X_p. \end{aligned}$$

By continuity argument, we see that (1.68) holds for all $x \in \overline{X_p}$. In Theorem 6.2, we will show that $\overline{X_p} = X$ (recall that X is connected) and hence (1.68) holds for all $x \in X$. From this observation, (1.68) and (1.63), we get Theorem 1.3 when X is CR Kähler.

When X is not CR Kähler, we still have (1.60), (1.61), (1.62) and (1.63). The *problem* is that the local operator $\square_{B_j, m}^+$ in (1.57) is not induced by Kähler form, it is difficult to understand small t behaviour of $A_{B_j, +}(t, z, z)$ and (1.66) is not true in general. To overcome this difficulty, we introduce some kind of CR Spin^c Dirac operator:

$$\tilde{D}_{b, m} = \bar{\partial}_b + \bar{\partial}_b^* + \text{zero order term}$$

on CR manifolds with S^1 action and modified Kohn Laplacians $\tilde{\square}_{b, m}^+ = \tilde{D}_{b, m}^* \tilde{D}_{b, m}$, $\tilde{\square}_{b, m}^- = \tilde{D}_{b, m} \tilde{D}_{b, m}^*$. We will show in Theorem 4.7 and Theorem 4.10 that we have *homotopy invariance* for the index of $\bar{\partial}_b + \bar{\partial}_b^*$ and *McKean-Singer formula* for the modified Kohn Laplacians: for every $t > 0$,

$$(1.69) \quad \sum_{j=0}^n (-1)^j \dim H_{b, m}^j(X, E) = \int_X \left(\mathrm{Tr} e^{-t\tilde{\square}_{b, m}^+}(x, x) - \mathrm{Tr} e^{-t\tilde{\square}_{b, m}^-}(x, x) \right) dv_X.$$

Let $B := (D, (z, \theta), \varphi)$ be a BRT trivialization. Let $u \in \Omega_m^{0,+}(X)$, $v \in \Omega_m^{0,-}(X)$. On D , we write $u(z, \theta) = e^{-im\theta} \tilde{u}(z)$, $v(z, \theta) = e^{-im\theta} \tilde{v}(z)$, $\tilde{u}(z) \in \Omega^{0,+}(U, L^m)$, $\tilde{v}(z) \in \Omega^{0,-}(U, L^m)$. We will show in Lemma 5.1 that

$$(1.70) \quad \begin{aligned} e^{-m\varphi} \tilde{\square}_{B, m}^+(e^{m\varphi} \tilde{u}) &= e^{im\theta} \tilde{\square}_{b, m}^+(u), \\ e^{-m\varphi} \tilde{\square}_{B, m}^-(e^{m\varphi} \tilde{v}) &= e^{im\theta} \tilde{\square}_{b, m}^-(v), \end{aligned}$$

where $\tilde{\square}_{B, m}^+ = D_{B, m}^* D_{B, m} : \Omega^{0,+}(U, L^m) \rightarrow \Omega^{0,+}(U, L^m)$, $\tilde{\square}_{B, m}^- = D_{B, m} D_{B, m}^* : \Omega^{0,+}(U, L^m) \rightarrow \Omega^{0,+}(U, L^m)$, where $D_{B, m} : \Omega^{0,+}(U, L^m) \rightarrow \Omega^{0,-}(U, L^m)$ is the Spin^c Dirac operator with respect to the Chern connection on L^m induced by h^{L^m} and Clifford connection on $\Lambda(T^{*0,1}U)$ induced by the given Hermitian metric on $\Lambda(T^{*0,1}U)$ (see Definition 4.1).

Let $B_j = (D_j, (z, \theta), \varphi_j)$, $j = 1, 2, \dots, N$, be BRT trivializations as before. Fix $j = 1, 2, \dots, N$. As (1.57), there is $\tilde{A}_{B_j, +}(t, z, w) \in C^\infty(\mathbb{R}_+ \times U_j \times U_j, (T_w^{*0,+}U \boxtimes T_z^{*0,+}U))$ such that

$$(1.71) \quad \begin{aligned} \lim_{t \rightarrow 0^+} \tilde{A}_{B_j, +}(t) &= I \text{ in } \mathcal{D}'(U, T^{*0,+}U), \\ \tilde{A}'_{B_j, +}(t)u + \tilde{A}_{B_j, +}(t)(\tilde{\square}_{B_j, m}^+ u) &= 0, \quad \forall u \in \Omega_0^{0,+}(U), \quad \forall t > 0, \end{aligned}$$

and $\tilde{A}_{B_j,+}(t, z, w)$ admits an asymptotic expansion as $t \rightarrow 0^+$ (see (5.17)). Put

$$(1.72) \quad \begin{aligned} \tilde{H}_j(t, x, y) &= \chi_j(x) e^{-m\varphi_j(z) - im\theta} \tilde{A}_{B_j,+}(t, z, w) e^{m\varphi_j(w) + im\eta} \tilde{\chi}_j(y), \\ \tilde{\Gamma}(t) &= \sum_{j=1}^N Q_m \circ \tilde{H}_j(t) \circ Q_m. \end{aligned}$$

We can repeat the method above and deduce that

$$(1.73) \quad \begin{aligned} \left\| e^{-t\tilde{\square}_{b,m}^+}(x, y) - \tilde{\Gamma}(t, x, y) \right\|_{C^0(X \times X)} &\leq e^{-\frac{\epsilon_0}{t}}, \quad \forall t \in (0, \epsilon), \\ \tilde{\Gamma}(t, x, y) &\sim t^{-n} \tilde{a}_n^+(t, x, y) + t^{-n+1} \tilde{a}_{n-1}^+(t, x, y) + \dots \quad \text{as } t \rightarrow 0^+, \\ e^{-t\tilde{\square}_{b,m}^+} &\sim t^{-n} \tilde{a}_n^+(t, x, y) + t^{-n+1} \tilde{a}_{n-1}^+(t, x, y) + \dots \quad \text{as } t \rightarrow 0^+, \\ e^{-t\tilde{\square}_{b,m}^-} &\sim t^{-n} \tilde{a}_n^-(t, x, y) + t^{-n+1} \tilde{a}_{n-1}^-(t, x, y) + \dots \quad \text{as } t \rightarrow 0^+, \end{aligned}$$

where $\epsilon > 0$, $\epsilon_0 > 0$ are constants and there is a constant $C > 0$ such that $|\tilde{a}_s^+(t, x, y)| \leq C$, for every $t > 0$, $(x, y) \in X \times X$ and every $s = n, n-1, \dots$. From (1.69) and (1.72), we deduce that

$$(1.74) \quad \sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, E) = \lim_{t \rightarrow 0^+} \int_X \sum_{\ell=0}^n t^{-\ell} \left(\text{Tr } \tilde{a}_\ell^+(t, x, x) - \text{Tr } \tilde{a}_\ell^-(t, x, x) \right) dv_X(x).$$

The *point* is that since *Lichnerowicz formulas* holds for $\tilde{\square}_{B,m}^+$ and $\tilde{\square}_{B,m}^-$, we can apply *rescaling technique* in [2] and [8] and show that for each $j = 1, 2, \dots, N$, we have

$$(1.75) \quad \begin{aligned} &\left(\text{Tr } \tilde{A}_{B_j,+}(t, z, z) - \text{Tr } \tilde{A}_{B_j,-}(t, z, z) \right) dv_{U_j}(z) \\ &= [\text{Td}(\nabla^{T^{1,0}U_j}, T^{1,0}U_j) \wedge \text{ch}(\nabla^{L^m}, L^m)]_{2n}(z) + O(t), \quad \forall z \in U_j. \end{aligned}$$

From (1.75), we can repeat the method above and conclude that

$$(1.76) \quad \begin{aligned} &\sum_{\ell=0}^n t^{-\ell} \left(\text{Tr } \tilde{a}_\ell^+(t, x, x) - \text{Tr } \tilde{a}_\ell^-(t, x, x) \right) dv_X(x) \\ &= \sum_{s=1}^p e^{\frac{2\pi(s-1)}{p} mi} \left[\text{Td}_b(\nabla^{T^{1,0}X}, T^{1,0}X) \wedge e^{-m \frac{d\omega_0}{2\pi}} \wedge \omega_0 \right]_{2n+1}(x) + O(t), \end{aligned}$$

for every $t > 0$ and $x \in X$. From (1.74) and (1.76), Theorem 1.3 follows.

The layout of this paper is as follows. In Section 2.1 and Section 2.2, we collect some notations, definitions, terminology and statements we use throughout. In Section 2.3, we introduce tangential De-Rham cohomology group, tangential Chern character and tangential Todd class on CR manifolds with S^1 action. In Section 2.4, we recall classical result of Baouendi-Rothschild-Treves [1] which plays an important role in our construction of the heat kernel. We also proved that for a rigid vector bundle F over X , there are rigid Hermitian metric and rigid connection on F . In Section 3, we establish Hodge theory for Kohn Laplacian in the L^2 space of m -th S^1 Fourier coefficient. In Section 4, we introduce CR Spin^c Dirac operator $\tilde{D}_{b,m}$, modified Kohn Laplacians $\tilde{\square}_{b,m}^+$, $\tilde{\square}_{b,m}^-$ and we prove (1.69). In Section 5, we construct approximate heat kernels for the operators $e^{-t\tilde{\square}_{b,m}^+}$, $e^{-t\tilde{\square}_{b,m}^-}$ and we prove that $e^{-t\tilde{\square}_{b,m}^+}(x, y)$, $e^{-t\tilde{\square}_{b,m}^-}(x, y)$ admit asymptotic expansions as (1.73). In Section 6, we prove (1.76) and we finish the proof of Theorem 1.3.

2. PRELIMINARIES

2.1. Some standard notations. We use the following notations: $\mathbb{N} = \{1, 2, \dots\}$, $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$, \mathbb{R} is the set of real numbers, $\overline{\mathbb{R}}_+ := \{x \in \mathbb{R}; x \geq 0\}$. For a multiindex $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$ we set $|\alpha| = \alpha_1 + \dots + \alpha_n$. For $x = (x_1, \dots, x_n)$ we write

$$x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}, \quad \partial_{x_j} = \frac{\partial}{\partial x_j}, \quad \partial_x^\alpha = \partial_{x_1}^{\alpha_1} \dots \partial_{x_n}^{\alpha_n} = \frac{\partial^{|\alpha|}}{\partial x^\alpha},$$

$$D_{x_j} = \frac{1}{i} \partial_{x_j}, \quad D_x^\alpha = D_{x_1}^{\alpha_1} \dots D_{x_n}^{\alpha_n}, \quad D_x = \frac{1}{i} \partial_x.$$

Let $z = (z_1, \dots, z_n)$, $z_j = x_{2j-1} + ix_{2j}$, $j = 1, \dots, n$, be coordinates of \mathbb{C}^n . We write

$$z^\alpha = z_1^{\alpha_1} \dots z_n^{\alpha_n}, \quad \bar{z}^\alpha = \bar{z}_1^{\alpha_1} \dots \bar{z}_n^{\alpha_n},$$

$$\partial_{z_j} = \frac{\partial}{\partial z_j} = \frac{1}{2} \left(\frac{\partial}{\partial x_{2j-1}} - i \frac{\partial}{\partial x_{2j}} \right), \quad \partial_{\bar{z}_j} = \frac{\partial}{\partial \bar{z}_j} = \frac{1}{2} \left(\frac{\partial}{\partial x_{2j-1}} + i \frac{\partial}{\partial x_{2j}} \right),$$

$$\partial_z^\alpha = \partial_{z_1}^{\alpha_1} \dots \partial_{z_n}^{\alpha_n} = \frac{\partial^{|\alpha|}}{\partial z^\alpha}, \quad \partial_{\bar{z}}^\alpha = \partial_{\bar{z}_1}^{\alpha_1} \dots \partial_{\bar{z}_n}^{\alpha_n} = \frac{\partial^{|\alpha|}}{\partial \bar{z}^\alpha}.$$

Let X be a C^∞ orientable paracompact manifold. We let TX and T^*X denote the tangent bundle of X and the cotangent bundle of X respectively. The complexified tangent bundle of X and the complexified cotangent bundle of X will be denoted by $\mathbb{C}TX$ and $\mathbb{C}T^*X$ respectively. We write $\langle \cdot, \cdot \rangle$ to denote the pointwise duality between T^*X and TX . We extend $\langle \cdot, \cdot \rangle$ bilinearly to $\mathbb{C}T^*X \times \mathbb{C}TX$. For $u \in \mathbb{C}T^*X$, $v \in \mathbb{C}TX$, we also write $u(v) := \langle u, v \rangle$.

Let E be a C^∞ vector bundle over X . The fiber of E at $x \in X$ will be denoted by E_x . Let F be another vector bundle over X . We write $E \boxtimes F$ to denote the vector bundle over $X \times X$ with fiber over $(x, y) \in X \times X$ consisting of the linear maps from E_x to F_y .

Let $Y \subset X$ be an open set. The spaces of smooth sections of E over Y and distribution sections of E over Y will be denoted by $C^\infty(Y, E)$ and $\mathcal{D}'(Y, E)$ respectively. Let $\mathcal{E}'(Y, E)$ be the subspace of $\mathcal{D}'(Y, E)$ whose elements have compact support in Y . For $m \in \mathbb{R}$, we let $H^m(Y, E)$ denote the Sobolev space of order m of sections of E over Y . Put

$$H_{\text{loc}}^m(Y, E) = \{u \in \mathcal{D}'(Y, E); \varphi u \in H^m(Y, E), \varphi \in C_0^\infty(Y)\},$$

$$H_{\text{comp}}^m(Y, E) = H_{\text{loc}}^m(Y, E) \cap \mathcal{E}'(Y, E).$$

2.2. Set up and terminology. Let $(X, T^{1,0}X)$ be a compact CR manifold of dimension $2n + 1$, $n \geq 1$, where $T^{1,0}X$ is a CR structure of X . That is $T^{1,0}X$ is a subbundle of rank n of the complexified tangent bundle $\mathbb{C}TX$, satisfying $T^{1,0}X \cap T^{0,1}X = \{0\}$, where $T^{0,1}X = \overline{T^{1,0}X}$, and $[\mathcal{V}, \mathcal{V}] \subset \mathcal{V}$, where $\mathcal{V} = C^\infty(X, T^{1,0}X)$. We assume that X admits an S^1 action: $S^1 \times X \rightarrow X$. We write $e^{-i\theta}$ to denote the S^1 action. Let $T \in C^\infty(X, TX)$ be the global real vector field induced by the S^1 action given by $(Tu)(x) = \frac{\partial}{\partial \theta} (u(e^{-i\theta} \circ x))|_{\theta=0}$, $u \in C^\infty(X)$.

Definition 2.1. We say that the S^1 action $e^{-i\theta}$ is CR if $[T, C^\infty(X, T^{1,0}X)] \subset C^\infty(X, T^{1,0}X)$ and the S^1 action is transversal if for each $x \in X$, $\mathbb{C}T(x) \oplus T_x^{1,0}X \oplus T_x^{0,1}X = \mathbb{C}T_xX$. Moreover, we say that the S^1 action is locally free if $T \neq 0$ everywhere.

We assume throughout that $(X, T^{1,0}X)$ is a CR manifold with a transversal CR locally free S^1 action $e^{-i\theta}$ and we let T be the global vector field induced by the S^1 action. Let $\omega_0 \in C^\infty(X, T^*X)$ be the global real one form determined by $\langle \omega_0, u \rangle = 0$, for every $u \in T^{1,0}X \oplus T^{0,1}X$ and $\langle \omega_0, T \rangle = 1$.

Definition 2.2. For $p \in X$, the Levi form \mathcal{L}_p is the Hermitian quadratic form on $T_p^{1,0}X$ given by $\mathcal{L}_p(U, \bar{V}) = -\frac{1}{2i} \langle d\omega_0(p), U \wedge \bar{V} \rangle$, $U, V \in T_p^{1,0}X$.

If the Levi form \mathcal{L}_p is semi-positive definite (positive definite), we say that X is weakly pseudoconvex (strongly pseudoconvex) at p . If the Levi form is semi-positive definite (positive definite) at every point of X , we say that X is weakly pseudoconvex (strongly pseudoconvex).

Denote by $T^{*1,0}X$ and $T^{*0,1}X$ the dual bundles of $T^{1,0}X$ and $T^{0,1}X$ respectively. Define the vector bundle of (p, q) forms by $T^{*p,q}X = \Lambda^p(T^{*1,0}X) \wedge \Lambda^q(T^{*0,1}X)$. Put

$$T^{*0,+}X := \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is even}} T^{*0,j}X, \quad T^{*0,-}X := \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is odd}} T^{*0,j}X.$$

Let $D \subset X$ be an open subset. Let $\Omega^{p,q}(D)$ denote the space of smooth sections of $T^{*p,q}X$ over D and let $\Omega_0^{p,q}(D)$ be the subspace of $\Omega^{p,q}(D)$ whose elements have compact support in D . Similarly, if E is a vector bundle over D , then we let $\Omega^{p,q}(D, E)$ denote the space of smooth sections of $T^{*p,q}X \otimes E$ over D and let $\Omega_0^{p,q}(D, E)$ be the subspace of $\Omega^{p,q}(D, E)$ whose elements have compact support in D . Put

$$(2.1) \quad \begin{aligned} \Omega^{0,+}(X, E) &:= \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is even}} \Omega^{0,j}(X, E), \\ \Omega^{0,-}(X, E) &:= \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is odd}} \Omega^{0,j}(X, E). \end{aligned}$$

Fix $\theta_0 \in]-\pi, \pi[$, θ_0 small. Let

$$de^{-i\theta_0} : \mathbb{C}T_x X \rightarrow \mathbb{C}T_{e^{-i\theta_0}x} X$$

denote the differential map of $e^{-i\theta_0} : X \rightarrow X$. By the transversal property of the S^1 action, we can check that

$$(2.2) \quad \begin{aligned} de^{-i\theta_0} : T_x^{1,0}X &\rightarrow T_{e^{-i\theta_0}x}^{1,0}X, \\ de^{-i\theta_0} : T_x^{0,1}X &\rightarrow T_{e^{-i\theta_0}x}^{0,1}X, \\ de^{-i\theta_0}(T(x)) &= T(e^{-i\theta_0}x). \end{aligned}$$

Let $(e^{-i\theta_0})^* : \Lambda^r(\mathbb{C}T^*X) \rightarrow \Lambda^r(\mathbb{C}T^*X)$ be the pull-back map by $e^{-i\theta_0}$, $r = 0, 1, \dots, 2n+1$. From (2.2), it is easy to see that for every $p, q = 0, 1, \dots, n$,

$$(2.3) \quad (e^{-i\theta_0})^* : T_{e^{-i\theta_0}x}^{*p,q}X \rightarrow T_x^{*p,q}X.$$

Let $u \in \Omega^{p,q}(X)$. Define

$$(2.4) \quad Tu := \frac{\partial}{\partial \theta} ((e^{-i\theta})^* u)|_{\theta=0} \in \Omega^{p,q}(X).$$

(See also (2.14).) For every $\theta \in \mathbb{R}$ and every $u \in C^\infty(X, \Lambda^r(\mathbb{C}T^*X))$, we write $u(e^{-i\theta} \circ x) := (e^{-i\theta})^* u(x)$. It is clear that for every $u \in C^\infty(X, \Lambda^r(\mathbb{C}T^*X))$, we have

$$(2.5) \quad u(x) = \sum_{m \in \mathbb{Z}} \frac{1}{2\pi} \int_{-\pi}^{\pi} u(e^{-i\theta} \circ x) e^{im\theta} d\theta.$$

Let $\bar{\partial}_b : \Omega^{0,q}(X) \rightarrow \Omega^{0,q+1}(X)$ be the tangential Cauchy-Riemann operator. From the transversal property of the S^1 action, it is straightforward to see that (see also (2.15))

$$T\bar{\partial}_b = \bar{\partial}_b T \quad \text{on } \Omega^{0,q}(X).$$

Definition 2.3. Let $D \subset U$ be an open set. We say that a function $u \in C^\infty(D)$ is rigid if $Tu = 0$. We say that a function $u \in C^\infty(X)$ is Cauchy-Riemann (CR for short) if $\bar{\partial}_b u = 0$. We call u a rigid CR function if $\bar{\partial}_b u = 0$ and $Tu = 0$.

Definition 2.4. Let F be a complex vector bundle over X . We say that F is rigid (CR) if X can be covered with open sets U_j with trivializing frames $\{f_j^1, f_j^2, \dots, f_j^r\}$, $j = 1, 2, \dots$, such that the corresponding transition matrices are rigid (CR). The frames $\{f_j^1, f_j^2, \dots, f_j^r\}$, $j = 1, 2, \dots$, are called rigid frames.

From now on, let E be a rigid CR vector bundle over X . We also write $\bar{\partial}_b$ to denote the tangential Cauchy-Riemann operator acting on forms with values in E :

$$\bar{\partial}_b : \Omega^{0,q}(X, E) \rightarrow \Omega^{0,q+1}(X, E).$$

Since E is rigid, we can also define Tu for every $u \in \Omega^{0,q}(X, E)$ and we have

$$(2.6) \quad T\bar{\partial}_b = \bar{\partial}_b T \quad \text{on } \Omega^{0,q}(X, E).$$

For every $m \in \mathbb{Z}$, let

$$(2.7) \quad \Omega_m^{0,q}(X, E) := \{u \in \Omega^{0,q}(X, E); Tu = -imu\}$$

and put $\bar{\partial}_{b,m} := \bar{\partial}_b : \Omega_m^{0,q}(X, E) \rightarrow \Omega_m^{0,q+1}(X, E)$. For every $m \in \mathbb{Z}$, we have $\bar{\partial}_{b,m}$ -complex: $\cdots \rightarrow \Omega_m^{0,q-1}(X, E) \rightarrow \Omega_m^{0,q}(X, E) \rightarrow \Omega_m^{0,q+1}(X, E) \rightarrow \cdots$. Define

$$H_{b,m}^q(X, E) := \frac{\text{Ker } \bar{\partial}_{b,m} : \Omega_m^{0,q}(X, E) \rightarrow \Omega_m^{0,q+1}(X, E)}{\text{Im } \bar{\partial}_{b,m} : \Omega_m^{0,q-1}(X, E) \rightarrow \Omega_m^{0,q}(X, E)}.$$

We call $H_{b,m}^q(X, E)$ the m -th S^1 Fourier coefficient of the q -th $\bar{\partial}_b$ Kohn-Rossi cohomology group. We will prove in Theorem 3.7 that $\dim H_{b,m}^q(X, E) < \infty$, for every $m \in \mathbb{Z}$ and every $q = 0, 1, 2, \dots, n$. When E is trivial, for every $m \in \mathbb{Z}$, we denote $\Omega_m^{0,q}(X) := \Omega_m^{0,q}(X, E)$ and $H_{b,m}^q(X) := H_{b,m}^q(X, E)$.

2.3. Tangential De-Rham cohomology group, Tangential Chern character and Tangential Todd class. For every $r = 0, 1, 2, \dots, 2n$, put $\Omega_0^r(X) = \{u \in \oplus_{p+q=r} \Omega^{p,q}(X); Tu = 0\}$ and set $\Omega_0^*(X) = \oplus_{r=0}^{2n} \Omega_0^r(X)$. Since $Td = dT$ (see (2.6)), we have d -complex:

$$d : \cdots \rightarrow \Omega_0^{r-1}(X) \rightarrow \Omega_0^r(X) \rightarrow \Omega_0^{r+1}(X) \rightarrow \cdots$$

and we define r -th tangential De-Rham cohomology group: $H_{b,0}^r(X) := \frac{\text{Ker } d : \Omega_0^r(X) \rightarrow \Omega_0^{r+1}(X)}{\text{Im } d : \Omega_0^{r-1}(X) \rightarrow \Omega_0^r(X)}$. Put $H_{b,0}^*(X) = \oplus_{r=0}^{2n} H_{b,0}^r(X)$.

Let F be a rigid complex vector bundle over X of rank r . We will show in Theorem 2.11 that there is a connection ∇ of F such that for any rigid local frame $f = (f_1, f_2, \dots, f_r)$ of F on an open set $D \subset X$, the connection matrix $\theta(\nabla, f) = (\theta_{j,k})_{j,k=1}^r$ satisfies $\theta_{j,k} \in \Omega_0^1(D)$, for every $j, k = 1, \dots, r$. We call such ∇ rigid connection. Let $\Theta(\nabla, F) \in C^\infty(X, \Lambda^2(\mathbb{C}T^*X) \otimes \text{End}(F))$ be the associated curvature. Let $h(z) = \sum_{j=0}^{\infty} a_j z^j$, $a_j \in \mathbb{R}$, for every j , be a real power series on $z \in \mathbb{C}$. Set

$$H(\Theta(\nabla, F)) = \text{Tr} \left(h \left(\frac{i}{2\pi} \Theta(\nabla, F) \right) \right).$$

It is clear that $H(\Theta(\nabla, F)) \in \Omega_0^*(X)$. The following is well-known (see Theorem B.5.1. in Ma-Marinescu [27]).

Theorem 2.5. $H(\Theta(\nabla, F))$ is a closed differential form.

We need

Theorem 2.6. Let ∇' be another rigid connection on F . Then, $H(\Theta(\nabla, F)) - H(\Theta(\nabla', F)) = dA$, for some $A \in \Omega_0^*(X)$.

Proof. For each $t \in [0, 1]$, put $\nabla_t = (1-t)\nabla + t\nabla'$. Then, ∇_t is a rigid connection on F . Put

$$Q_t = \frac{i}{2\pi} \text{Tr} \left(\frac{\partial \nabla_t}{\partial t} h \left(\frac{i}{2\pi} \Theta(\nabla_t, F) \right) \right).$$

Since ∇_t is rigid, it is easy to see that

$$(2.8) \quad Q_t \in \Omega_0^*(X).$$

It is well-known that (see Remark B.5.2. in Ma-Marinescu [27])

$$(2.9) \quad H(\Theta(\nabla, F)) - H(\Theta(\nabla', F)) = d \int_0^1 Q_t dt.$$

From (2.8) and (2.9), the theorem follows. \square

From Theorem 2.5 and Theorem 2.6, we see that the tangential De-Rham cohomology class

$$[H(\Theta(\nabla, F))] \in H_{b,0}^*(X)$$

does not depend on the choice of rigid connection ∇ . Put

$$(2.10) \quad \text{ch}_b(\nabla, F) := H(\Theta(\nabla, F)) \in \Omega_0^*(X),$$

where $h(z) = e^z$ and set

$$(2.11) \quad \text{Td}_b(\nabla, F) = e^{H(\Theta(\nabla, F))} \in \Omega_0^*(X),$$

where $h(z) = \log(\frac{z}{1-e^{-z}})$. We can now introduce tangential Todd class and tangential Chern character

Definition 2.7. Tangential Chern character of F is given by

$$\text{ch}_b(F) := [\text{ch}_b(\nabla, F)] \in H_{b,0}^*(X),$$

and tangential Todd class of F is given by

$$\text{Td}_b(F) = [\text{Td}_b(\nabla, F)] \in H_{b,0}^*(X).$$

Baouendi-Rothschild-Treves [1] proved that $T^{1,0}X$ is a rigid complex vector bundle over X . Thus, we can define tangential Todd class of $T^{1,0}X$ and tangential Chern character of $T^{1,0}X$.

Put $\det\left(\frac{i\Theta(\nabla, F)}{2\pi}t + I\right) = \sum_{j=0}^r \hat{c}_j(\nabla, F)t^j$. It is easy to check that $\hat{c}_j(\nabla, F) \in \Omega_0^{2j}(D)$. Since for any matrix A , we have $\det A = e^{\text{Tr}(\log A)}$. By taking $h(z) = \log(1+z)$, we can check that for each $j = 0, 1, \dots, r$, $\hat{c}_j(\nabla, F)$ is a closed differential form on X and its tangential De-Rham cohomology class $[\hat{c}_j(\nabla, F)] \in H_{b,0}^{2j}(X)$ does not depend on the choice of rigid connection D . Put $\hat{c}_j(F) = [\hat{c}_j(\nabla, F)] \in H_{b,0}^{2j}(X)$. We call $\hat{c}_j(F)$ j -th tangential Chern class of F , and $\hat{c}(F) = 1 + \sum_{j=1}^r \hat{c}_j(F) \in H_{b,0}^*(X)$ is called tangential total Chern class of F .

2.4. BRT trivialisations and rigid Hermitian metrics. In this work, we need the following result due to Baouendi-Rothschild-Treves [1].

Theorem 2.8. For every point $x_0 \in X$, we can find local coordinates $x = (x_1, \dots, x_{2n+1}) = (z, \theta) = (z_1, \dots, z_n, \theta)$, $z_j = x_{2j-1} + ix_{2j}$, $j = 1, \dots, n$, $x_{2n+1} = \theta$, defined in some small neighborhood $D = \{(z, \theta) : |z| < \delta, -\varepsilon_0 < \theta < \varepsilon_0\}$ of x_0 , $\delta > 0$, $0 < \varepsilon_0 < \pi$, such that $(z(x_0), \theta(x_0)) = (0, 0)$ and

$$(2.12) \quad \begin{aligned} T &= \frac{\partial}{\partial \theta} \\ Z_j &= \frac{\partial}{\partial z_j} - i \frac{\partial \varphi}{\partial z_j}(z) \frac{\partial}{\partial \theta}, j = 1, \dots, n \end{aligned}$$

where $Z_j(x)$, $j = 1, \dots, n$, form a basis of $T_x^{1,0}X$, for each $x \in D$ and $\varphi(z) \in C^\infty(D, \mathbb{R})$ independent of θ . We call $(D, (z, \theta), \varphi)$ BRT trivialization.

By using BRT trivialization, we get another way to define $Tu, \forall u \in \Omega^{0,q}(X)$. Let $(D, (z, \theta), \varphi)$ be a BRT trivialization. It is clear that

$$\{d\bar{z}_{j_1} \wedge \dots \wedge d\bar{z}_{j_q}, 1 \leq j_1 < \dots < j_q \leq n\}$$

is a basis for $T_x^{*0,q}X$, for every $x \in D$. Let $u \in \Omega^{0,q}(X)$. On D , we write

$$(2.13) \quad u = \sum_{j_1 < \dots < j_q} u_{j_1 \dots j_q} d\bar{z}_{j_1} \wedge \dots \wedge d\bar{z}_{j_q}.$$

Then on D we can check that

$$(2.14) \quad Tu = \sum_{j_1 < \dots < j_q} (Tu_{j_1 \dots j_q}) d\bar{z}_{j_1} \wedge \dots \wedge d\bar{z}_{j_q}$$

and Tu is independent of the choice of BRT trivializations. Note that on BRT trivialization $(D, (z, \theta), \varphi)$, we have

$$(2.15) \quad \bar{\partial}_b = \sum_{j=1}^n d\bar{z}_j \wedge \left(\frac{\partial}{\partial \bar{z}_j} + i \frac{\partial \varphi}{\partial \bar{z}_j}(z) \frac{\partial}{\partial \theta} \right).$$

Let F be a rigid vector bundle over X . We need

Definition 2.9. Let $\langle \cdot | \cdot \rangle_F$ be a Hermitian metric on F . We say that $\langle \cdot | \cdot \rangle_F$ is a rigid Hermitian metric if for every rigid local frames f_1, \dots, f_r of F , we have $T\langle f_j | f_k \rangle_F = 0$, for every $j, k = 1, 2, \dots, r$.

We need

Theorem 2.10. *There is a rigid Hermitian metric $\langle \cdot | \cdot \rangle_F$ on F .*

Proof. Fix $p \in X$ and let $(D, (z, \theta), \varphi)$ be a BRT trivialization defined in some neighbourhood of p such that $(z(p), \theta(p)) = (0, 0)$. Suppose that (z, θ) is defined on $\{z \in \mathbb{C}^{n-1} : |z| < \delta\} \times \{\theta \in \mathbb{R} : |\theta| < \delta\}$, for some $\delta > 0$. Put

$$A := \{\lambda \in [-\pi, \pi] : \text{There is a local rigid trivializing frames } f = (f_1, \dots, f_r) \text{ defined on the set } \\ \{e^{-i\theta} \circ (z, 0); |z| < \varepsilon, \theta \in [-\pi, \lambda + \varepsilon]\}, \text{ for some } 0 < \varepsilon < \delta\}.$$

It is clear that A is a non-empty open set in $[-\pi, \pi]$. We claim that A is closed. Let λ_0 be a limit point of A . Consider the point $e^{-i\lambda_0} \circ (0, 0)$. For some $\varepsilon_1 > 0$, ε_1 small, there are local rigid trivializing frames $\hat{f} = (\hat{f}_1, \dots, \hat{f}_r)$ defined on $\{e^{-i\theta} \circ (z, 0); |z| < \varepsilon_1, \lambda_0 - \varepsilon_1 < \theta_1 < \lambda_0 + \varepsilon_1\}$. Since λ_0 is a limit point of A , we can find local rigid trivializing frames $\tilde{f} = (\tilde{f}_1, \dots, \tilde{f}_r)$ defined on $\{e^{-i\theta} \circ (z, 0); |z| < \varepsilon_2, \theta \in [-\pi, \lambda_0 - \frac{\varepsilon_1}{2}]\}$, for some $\varepsilon_2 > 0$. Now, $\tilde{f} = g\hat{f}$ on

$$\left\{ e^{-i\theta} \circ (z, 0); |z| < \varepsilon_0, \theta \in (\lambda_0 - \varepsilon_1, \lambda_0 - \frac{\varepsilon_1}{2}) \right\}$$

for some rigid CR $r \times r$ matrix g , where $\varepsilon_0 = \min\{\varepsilon_1, \varepsilon_2\}$. Since g is independent of θ , g is well-defined on $\{e^{-i\theta} \circ (z, 0); |z| < \varepsilon_0, \theta \in \mathbb{R}\}$. Put $f = \tilde{f}$ on $\{e^{-i\theta} \circ (z, 0); |z| < \varepsilon_0, \theta \in [-\pi, \lambda_0 - \frac{\varepsilon_1}{2}]\}$ and $f = g\hat{f}$ on $\{e^{-i\theta} \circ (z, 0); |z| < \varepsilon_0, \theta \in [\lambda_0 - \frac{\varepsilon_1}{2}, \lambda_0 + \varepsilon_1]\}$. It is straightforward to check that f is well-defined as local rigid trivializing frames on

$$\left\{ e^{-i\theta} \circ (z, 0); |z| < \varepsilon_0, \theta \in [-\pi, \lambda_0 + \varepsilon_1] \right\}.$$

Thus, $\lambda_0 \in A$ and hence $A = [-\pi, \pi]$.

From the discussion above, we see that we can find local rigid trivializations W_1, \dots, W_N such that $X = \bigcup_{j=1}^N W_j$ and $W_t \subset \bigcup_{-\pi \leq \theta \leq \pi} e^{-i\theta} W_t$, $t = 1, \dots, N$. Take any Hermitian metric $\langle \cdot, \cdot \rangle_F$ on F . Let $\langle \cdot | \cdot \rangle_F$ be the Hermitian metric on F defined as follows. For each $j = 1, 2, \dots, N$, let h_j^1, \dots, h_j^r be local rigid trivializing frames on W_j . Put $\langle h_j^s(x) | h_j^t(x) \rangle_F = \frac{1}{2\pi} \int_{-\pi}^{\pi} \langle h_j^s(e^{-iu} \circ x), h_j^t(e^{-iu} \circ x) \rangle_F du$, $s, t = 1, 2, \dots, r$. Since F is rigid, it is easy to see that $\langle \cdot | \cdot \rangle_F$ is well-defined as a rigid Hermitian metric on F . \square

Now, we can prove

Theorem 2.11. *There is a rigid connection on F .*

Proof. From Theorem 2.10, there is a rigid Hermitian metric $\langle \cdot | \cdot \rangle_F$ on F . Let ∇ be the connection on F induced by $\langle \cdot | \cdot \rangle_F$. It is not difficult to check that ∇ is rigid. \square

3. HODGE THEORY FOR $\square_{b,m}^{(q)}$

From now on, we take a rigid Hermitian metric $\langle \cdot | \cdot \rangle_E$ on E and take a rigid Hermitian metric $\langle \cdot | \cdot \rangle$ on $\mathbb{C}TX$ such that $T^{1,0}X \perp T^{0,1}X$, $T \perp (T^{1,0}X \oplus T^{0,1}X)$, $\langle T | T \rangle = 1$. This is always possible, see Theorem 2.10 and Theorem 9.2 in [20]. The Hermitian metric $\langle \cdot | \cdot \rangle$ on $\mathbb{C}TX$ induces by duality a Hermitian metric on $\mathbb{C}T^*X$ and also on the bundles of $(0, q)$ forms $T^{*0,q}X$, $q = 0, 1, \dots, n$. We shall also denote all these induced metrics by $\langle \cdot | \cdot \rangle$. For every $v \in T^{*0,q}X$, we write $|v|^2 := \langle v | v \rangle$. The Hermitian metrics on $T^{*0,q}X$ and E induce Hermitian metrics on $T^{*0,q}X \otimes E$, $q = 0, 1, \dots, n$. We shall denote these induced metrics by $\langle \cdot | \cdot \rangle_E$. For $f \in \Omega^{0,q}(X, E)$, we denote the pointwise norm $|f(x)|_E^2 := \langle f(x) | f(x) \rangle_E$. We denote by $dv_X = dv_X(x)$ the volume form on X induced by the fixed Hermitian metric $\langle \cdot | \cdot \rangle$ on $\mathbb{C}TX$. Then we get natural global L^2 inner products $(\cdot | \cdot)_E$, $(\cdot | \cdot)$ on $\Omega^{0,q}(X, E)$ and $\Omega^{0,q}(X)$ respectively. We denote by $L^2(X, T^{*0,q}X \otimes E)$ and $L^2(X, T^{*0,q}X)$ the completions of $\Omega^{0,q}(X, E)$ and $\Omega^{0,q}(X)$ with respect to $(\cdot | \cdot)_E$, $(\cdot | \cdot)$ respectively. Similarly, for each $m \in \mathbb{Z}$, we denote by $L_m^2(X, T^{*0,q}X \otimes E)$ and $L_m^2(X, T^{*0,q}X)$ the completions of $\Omega_m^{0,q}(X, E)$ and $\Omega_m^{0,q}(X)$ with respect to $(\cdot | \cdot)_E$ and $(\cdot | \cdot)$ respectively. We extend $(\cdot | \cdot)_E$ and $(\cdot | \cdot)$ to $L^2(X, T^{*0,q}X \otimes E)$ and $L^2(X, T^{*0,q}X)$ in the standard way respectively. For $f \in \Omega^{0,q}(X, E)$, we denote $\|f\|_E^2 := (f | f)_E$. Similarly, for $f \in \Omega^{0,q}(X)$, we denote $\|f\|^2 := (f | f)$.

Let

$$\bar{\partial}_b^* : \Omega^{0,q+1}(X, E) \rightarrow \Omega^{0,q}(X, E)$$

be the formal adjoint of $\bar{\partial}_b$ with respect to $(\cdot | \cdot)_E$. Since $\langle \cdot | \cdot \rangle_E$ is rigid, we can check that

$$(3.1) \quad \begin{aligned} T\bar{\partial}_b^* &= \bar{\partial}_b^*T \quad \text{on } \Omega^{0,q}(X, E), \quad q = 1, 2, \dots, n, \\ \bar{\partial}_b^* &: \Omega_m^{0,q+1}(X, E) \rightarrow \Omega_m^{0,q}(X, E), \quad \forall m \in \mathbb{Z}. \end{aligned}$$

Put

$$(3.2) \quad \square_b^{(q)} := \bar{\partial}_b \bar{\partial}_b^* + \bar{\partial}_b^* \bar{\partial}_b : \Omega^{0,q}(X, E) \rightarrow \Omega^{0,q}(X, E).$$

From (2.6) and (3.1), we have

$$(3.3) \quad \begin{aligned} T\square_b^{(q)} &= \square_b^{(q)}T \quad \text{on } \Omega^{0,q}(X, E), \quad q = 0, 1, \dots, n, \\ \square_b^{(q)} &: \Omega_m^{0,q}(X, E) \rightarrow \Omega_m^{0,q}(X, E), \quad \forall m \in \mathbb{Z}. \end{aligned}$$

The following follows from Kohn's L^2 estimates (see [5, Theorem 8.4.2])

Theorem 3.1. *For every $s \in \mathbb{N}_0$, there is a constant $C_s > 0$ such that*

$$(3.4) \quad \|u\|_{s+1} \leq C_s \left(\left\| \square_b^{(q)} u \right\|_s + \|Tu\|_s + \|u\|_s \right), \quad \forall u \in \Omega^{0,q}(X, E),$$

where $\|\cdot\|_s$ denotes the usual Sobolev norm of order s on X .

We will write $\square_{b,m}^{(q)}$ to denote the restriction of $\square_b^{(q)}$ on the space $\Omega_m^{0,q}(X, E)$. From Theorem 3.1, we deduce

Theorem 3.2. *Fix $m \in \mathbb{Z}$. For every $s \in \mathbb{N}_0$, there is a constant $C_s > 0$ such that*

$$(3.5) \quad \|u\|_{s+1} \leq C_s \left(\left\| \square_{b,m}^{(q)} u \right\|_s + \|u\|_s \right), \quad \forall u \in \Omega_m^{0,q}(X, E),$$

where $\|\cdot\|_s$ denotes the usual Sobolev norm of order s on X .

For every $m \in \mathbb{Z}$, we extend $\square_{b,m}^{(q)}$ to $L_m^2(X, T^{*0,q}X \otimes E)$ by

$$(3.6) \quad \square_{b,m}^{(q)} : \text{Dom } \square_{b,m}^{(q)} \subset L_m^2(X, T^{*0,q}X \otimes E) \rightarrow L_m^2(X, T^{*0,q}X \otimes E),$$

where $\text{Dom } \square_{b,m}^{(q)} := \{u \in L_m^2(X, T^{*0,q}X \otimes E); \square_{b,m}^{(q)} u \in L_m^2(X, T^{*0,q}X \otimes E)\}$, where for any $u \in L_m^2(X, T^{*0,q}X \otimes E)$, $\square_{b,m}^{(q)} u$ is defined in the sense of distribution.

Lemma 3.3. *We have $\text{Dom } \square_{b,m}^{(q)} = L_m^2(X, T^{*0,q}X \otimes E) \cap H^2(X, T^{*0,q}X \otimes E)$.*

Proof. It is clear that $L_m^2(X, T^{*0,q}X \otimes E) \cap H^2(X, T^{*0,q}X \otimes E) \subset \text{Dom } \square_{b,m}^{(q)}$. We only need to prove that $\text{Dom } \square_{b,m}^{(q)} \subset L_m^2(X, T^{*0,q}X \otimes E) \cap H^2(X, T^{*0,q}X \otimes E)$. Let $u \in \text{Dom } \square_{b,m}^{(q)}$. Put $v = \square_{b,m}^{(q)}u \in L_m^2(X, T^{*0,q}X \otimes E)$. We have $(\square_{b,m}^{(q)} - T^2)u = v + m^2u \in L_m^2(X, T^{*0,q}X \otimes E)$. Since $(\square_{b,m}^{(q)} - T^2)$ is elliptic, we conclude that $u \in H^2(X, T^{*0,q}X \otimes E)$. The lemma follows. \square

We can prove

Theorem 3.4. $\square_{b,m}^{(q)} : \text{Dom } \square_{b,m}^{(q)} \subset L_m^2(X, T^{*0,q}X \otimes E) \rightarrow L_m^2(X, T^{*0,q}X \otimes E)$ is self-adjoint.

Proof. Let $(\square_{b,m}^{(q)})^* : \text{Dom } (\square_{b,m}^{(q)})^* \subset L_m^2(X, T^{*0,q}X \otimes E) \rightarrow L_m^2(X, T^{*0,q}X \otimes E)$ be the Hilbert space adjoint of $\square_{b,m}^{(q)}$. Let $v \in \text{Dom } (\square_{b,m}^{(q)})^*$. Then, by the definition of the Hilbert space adjoint of $\square_{b,m}^{(q)}$, it is easy to see that $\square_{b,m}^{(q)}v \in L_m^2(X, T^{*0,q}X \otimes E)$ and hence $v \in \text{Dom } \square_{b,m}^{(q)}$ and $\square_{b,m}^{(q)}v = (\square_{b,m}^{(q)})^*v$.

From Lemma 3.3, we can check that

$$(3.7) \quad (\square_{b,m}^{(q)}g | f)_E = (g | \square_{b,m}^{(q)}f)_E, \quad \forall g, f \in \text{Dom } \square_{b,m}^{(q)}.$$

From (3.7), we deduce that $\text{Dom } \square_{b,m}^{(q)} \subset \text{Dom } (\square_{b,m}^{(q)})^*$ and $\square_{b,m}^{(q)}u = (\square_{b,m}^{(q)})^*u$, for all $u \in \text{Dom } \square_{b,m}^{(q)}$. The theorem follows. \square

Let $\text{Spec } \square_{b,m}^{(q)} \subset [0, \infty[$ denote the spectrum of $\square_{b,m}^{(q)}$. For any $\lambda > 0$, put

$$\Pi_{m, \leq \lambda}^{(q)} := E_m^{(q)}([0, \lambda]),$$

where $E_m^{(q)}$ denotes the spectral measure for $\square_{b,m}^{(q)}$ (see Section 2 in Davies [7], for the precise meaning of spectral measure). We need

Theorem 3.5. $\text{Spec } \square_{b,m}^{(q)}$ is a discrete subset of $[0, \infty[$. Any $\nu \in \text{Spec } \square_{b,m}^{(q)}$, ν is an eigenvalue of $\square_{b,m}^{(q)}$ and the eigenspace

$$\mathcal{E}_{m, \nu}^q(X, E) := \{u \in \text{Dom } \square_{b,m}^{(q)}; \square_{b,m}^{(q)}u = \nu u\}$$

is finite dimensional with $\mathcal{E}_{m, \nu}^q(X, E) \subset \Omega_m^{0,q}(X, E)$.

Proof. Fix $\lambda > 0$. We claim that $\text{Spec } \square_{b,m}^{(q)} \cap [0, \lambda]$ is discrete. If not, we can find $f_j \in \text{Rang } E_m^{(q)}([0, \lambda])$, $j = 1, 2, \dots$, with $(f_j | f_k)_E = \delta_{j,k}$, $j, k = 1, 2, \dots$. Note that

$$(3.8) \quad \left\| \square_{b,m}^{(q)}f_j \right\|_E \leq \lambda \|f_j\|_E, \quad j = 1, 2, \dots$$

From (3.8), we have

$$(3.9) \quad \left\| (\square_{b,m}^{(q)} - T^2)f_j \right\|_E \leq (\lambda + m^2) \|f_j\|_E, \quad j = 1, 2, \dots$$

Since $\square_{b,m}^{(q)} - T^2$ is a second order elliptic operator, there is a constant $C_m > 0$ independent of j such that

$$(3.10) \quad \|f_j\|_2 \leq C_m, \quad j = 1, 2, \dots,$$

where $\|\cdot\|_2$ denotes the usual Sobolev norm of order 2. From (3.10), we can apply Rellich's theorem and find subsequence $\{f_{j_s}\}_{s=1}^\infty$, $1 \leq j_1 < j_2 < \dots$ such that $f_{j_s} \rightarrow f$ in $L_m^2(X, T^{*0,q}X \otimes E)$. Since $(f_j | f_k)_E = \delta_{j,k}$, $j, k = 1, 2, \dots$, we get a contradiction. Thus, $\text{Spec } \square_{b,m}^{(q)} \cap [0, \lambda]$ is discrete. Hence $\text{Spec } \square_{b,m}^{(q)}$ is a discrete subset of $[0, \infty[$.

Let $r \in \text{Spec } \square_{b,m}^{(q)}$. Since $\text{Spec } \square_{b,m}^{(q)}$ is discrete, $\square_{b,m}^{(q)} - r$ has L^2 closed range. If $\square_{b,m}^{(q)} - r$ is injective, then $\text{Range } (\square_{b,m}^{(q)} - r) = L_m^2(X, T^{*0,q}X \otimes E)$ and

$$(\square_{b,m}^{(q)} - r)^{-1} : L_m^2(X, T^{*0,q}X \otimes E) \rightarrow L_m^2(X, T^{*0,q}X \otimes E)$$

is continuous. We get a contradiction. Hence r is an eigenvalue of $\text{Spec } \square_{b,m}^{(q)}$.

For any $\nu \in \text{Spec } \square_{b,m}^{(q)}$, put

$$\mathcal{E}_{m,\nu}^q(X, E) := \{u \in \text{Dom } \square_{b,m}^{(q)}; \square_{b,m}^{(q)} u = \nu u\}.$$

We can repeat the argument before and conclude that $\mathcal{E}_{m,\nu}^q(X, E)$ is finite dimensional. Let $u \in \mathcal{E}_{m,\nu}^q(X, E)$. Then, $\square_{b,m}^{(q)} u = \nu u$ and hence

$$(3.11) \quad (\square_{b,m}^{(q)} - T^2)u = (\nu + m^2)u.$$

From (3.11), we deduce that $u \in \Omega_m^{0,q}(X, E)$. Thus, $\mathcal{E}_{m,\nu}^q(X, E) \subset \Omega_m^{0,q}(X, E)$. The theorem follows. \square

Let $N_m^{(q)} : L_m^2(X, T^{*0,q}X \otimes E) \rightarrow \text{Dom } \square_{b,m}^{(q)}$ be the partial inverse of $\square_{b,m}^{(q)}$ and let

$$\Pi_m^{(q)} : L_m^2(X, T^{*0,q}X \otimes E) \rightarrow \text{Ker } \square_{b,m}^{(q)}$$

be the orthogonal projection. We have

$$(3.12) \quad \begin{aligned} \square_{b,m}^{(q)} N_m^{(q)} + \Pi_m^{(q)} &= I \text{ on } L_m^2(X, T^{*0,q}X \otimes E), \\ N_m^{(q)} \square_{b,m}^{(q)} + \Pi_m^{(q)} &= I \text{ on } \text{Dom } \square_{b,m}^{(q)}. \end{aligned}$$

We need

Lemma 3.6. *We have $N_m^{(q)} : \Omega_m^{0,q}(X, E) \rightarrow \Omega_m^{0,q}(X, E)$.*

Proof. Let $u \in \Omega_m^{0,q}(X, E)$ and put $N_m^{(q)} u = v \in L_m^2(X, T^{*0,q}X \otimes E)$. From (3.12), we have $(I - \Pi_m^{(q)})u = \square_{b,m}^{(q)} v$. Hence,

$$(3.13) \quad (\square_{b,m}^{(q)} - T^2)v = (I - \Pi_m^{(q)})u + m^2v.$$

In view of Theorem 3.5, we see that

$$(3.14) \quad (I - \Pi_m^{(q)})u \in \Omega_m^{0,q}(X, E).$$

Note that $\square_{b,m}^{(q)} - T^2$ is elliptic. From this observation, (3.13) and (3.14), we can repeat the standard technique in elliptic partial differential operator and conclude that $v \in \Omega_m^{0,q}(X, E)$. The lemma follows. \square

Now, we can prove

Theorem 3.7. *For every $q \in \{0, 1, 2, \dots, n\}$ and every $m \in \mathbb{Z}$, we have*

$$(3.15) \quad \text{Ker } \square_{b,m}^{(q)} = \mathcal{E}_{m,0}^q(X, E) \cong H_{b,m}^q(X, E).$$

In particular, $\dim H_{b,m}^q(X, E) < \infty$, for every $m \in \mathbb{Z}$ and every $q = 0, 1, 2, \dots, n$.

Proof. Consider the map

$$\begin{aligned} \tau_m^q : \text{Ker } \bar{\partial}_{b,m} \cap \Omega_m^{0,q}(X, E) &\rightarrow \text{Ker } \square_{b,m}^{(q)}, \\ u &\rightarrow \Pi_m^{(q)} u. \end{aligned}$$

It is clear that τ_m^q is surjective. Put $M_m^q := \left\{ \bar{\partial}_{b,m} u; u \in \Omega_m^{0,q-1}(X, E) \right\}$. We now prove that

$$(3.16) \quad \text{Ker } \tau_m^q = M_m^q.$$

It is clear that $M_m^q \subset \text{Ker } \tau_m^q$. Let $u \in \text{Ker } \tau_m^q$. From (3.12), we have

$$(3.17) \quad \begin{aligned} u &= \square_{b,m}^{(q)} N_m^{(q)} u + \Pi_m^{(q)} u \\ &= (\bar{\partial}_b \bar{\partial}_b^* + \bar{\partial}_b^* \bar{\partial}_b) N_m^{(q)} u. \end{aligned}$$

We claim that

$$(3.18) \quad \bar{\partial}_b^* \bar{\partial}_b N_m^{(q)} u = 0.$$

We have

$$(3.19) \quad \begin{aligned} (\bar{\partial}_b^* \bar{\partial}_b N_m^{(q)} u | \bar{\partial}_b^* \bar{\partial}_b N_m^{(q)} u)_E &= (\bar{\partial}_b^* \bar{\partial}_b \square_{b,m}^{(q)} N_m^{(q)} u | N_m^{(q)} u)_E = (\bar{\partial}_b^* \bar{\partial}_b (I - \Pi_m^{(q)}) u | N_m^{(q)} u)_E \\ &= (\bar{\partial}_b^* \bar{\partial}_b u | N_m^{(q)} u)_E = 0. \end{aligned}$$

From (3.19), the claim (3.18) follows. From (3.18) and (3.17), we deduce that

$$(3.20) \quad u = \bar{\partial}_b \bar{\partial}_b^* N_m^{(q)} u.$$

From Lemma 3.6, we know that $\bar{\partial}_b^* N_m^{(q)} u \in \Omega_m^{0,q-1}(X, E)$. From this observation and (3.20), we get that $u \in M_m^q$. (3.16) follows. From (3.16) and τ_m^q is surjective, the theorem follows. \square

For every $m \in \mathbb{Z}$, put

$$(3.21) \quad \begin{aligned} \Omega_m^{0,+}(X, E) &:= \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is even}} \Omega_m^{0,j}(X, E), \\ \Omega_m^{0,-}(X, E) &:= \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is odd}} \Omega_m^{0,j}(X, E). \end{aligned}$$

Similarly, put

$$(3.22) \quad \begin{aligned} \Omega^{0,+}(X, E) &:= \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is even}} \Omega^{0,j}(X, E), \\ \Omega^{0,-}(X, E) &:= \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is odd}} \Omega^{0,j}(X, E). \end{aligned}$$

Let $L^{2,+}(X, E)$, $L^{2,-}(X, E)$, $L_m^{2,+}(X, E)$, $L_m^{2,-}(X, E)$ be the completions of $\Omega^{0,+}(X, E)$, $\Omega^{0,-}(X, E)$, $\Omega_m^{0,+}(X, E)$ and $\Omega_m^{0,-}(X, E)$ with respect to $(\cdot | \cdot)_E$ respectively. Let

$$(3.23) \quad D_{b,m} := \bar{\partial}_b + \bar{\partial}_b^* : \Omega_m^{0,+}(X, E) \rightarrow \Omega_m^{0,-}(X, E).$$

We extend $D_{b,m}$ to $L_m^{2,+}(X, E)$:

$$D_{b,m} : \text{Dom } D_{b,m} \subset L_m^{2,+}(X, E) \rightarrow L_m^{2,-}(X, E),$$

where $\text{Dom } D_{b,m} = \left\{ u \in L_m^{2,+}(X, E); D_{b,m} u \in L_m^{2,-}(X, E) \right\}$. Let

$$D_{b,m}^* : \text{Dom } D_{b,m}^* \subset L_m^{2,-}(X, E) \rightarrow L_m^{2,+}(X, E)$$

be the Hilbert space adjoint of $D_{b,m}$ with respect to $(\cdot | \cdot)_E$. From Theorem 3.5 and Theorem 3.7, it is not difficult to see that (we omit the proof)

Theorem 3.8. *We have*

$$(3.24) \quad \begin{aligned} \text{Ker } D_{b,m} &= \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is even}} \text{Ker } \square_{b,m}^{(j)} \subset \Omega_m^{0,+}(X, E), \\ \text{Ker } D_{b,m}^* &= \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is odd}} \text{Ker } \square_{b,m}^{(j)} \subset \Omega_m^{0,-}(X, E). \end{aligned}$$

Put $\text{ind } D_{b,m} := \dim \text{Ker } D_{b,m} - \dim \text{Ker } D_{b,m}^*$. Then,

$$(3.25) \quad \sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, E) = \text{ind } D_{b,m}.$$

4. MODIFIED KOHN LAPLACIANS

To calculate a local density of $\sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, E)$, we have to modify the standard Kohn Laplacian.

We recall briefly some basic definitions and properties of Clifford connection and Spin^c Dirac operator. For more details about Clifford connection and Spin^c Dirac operator, see Chapter 1 in [27] and [8].

Let $B := (D, (z, \theta), \varphi)$ be a BRT trivialization. We may assume that $D = U \times]-\varepsilon, \varepsilon[$, where $\varepsilon > 0$ and U is an open set of \mathbb{C}^n . Let $T^{1,0}U$ denote the holomorphic tangent bundle of U and put $T^{0,1}U := \overline{T^{1,0}U}$. Let $\langle \cdot, \cdot \rangle$ be the Hermitian metric on $\mathbb{C}TU$ given by

$$\left\langle \frac{\partial}{\partial z_j}, \frac{\partial}{\partial z_k} \right\rangle = \left\langle \frac{\partial}{\partial z_j} - i \frac{\partial \varphi}{\partial z_j}(z) \frac{\partial}{\partial \theta} \mid \frac{\partial}{\partial z_k} - i \frac{\partial \varphi}{\partial z_k}(z) \frac{\partial}{\partial \theta} \right\rangle, \quad j, k = 1, 2, \dots, n.$$

$\langle \cdot, \cdot \rangle$ induces Hermitian metrics on $T^{*0,q}U$ bundle of $(0, q)$ forms on U , $q = 0, 1, \dots, n$, we shall also denote the Hermitian metrics by $\langle \cdot, \cdot \rangle$. Let g^{TU} be the Riemannian metric on TU induced by $\langle \cdot, \cdot \rangle$. Let $\{w_j\}_{j=1}^n$ be a local orthonormal frame of $T^{1,0}U$ with respect to $\langle \cdot, \cdot \rangle$ with dual frame $\{\bar{w}^j\}_{j=1}^n$. Then,

$$(4.1) \quad e_{2j-1} = \frac{1}{\sqrt{2}}(w_j + \bar{w}_j) \text{ and } e_{2j} = \frac{i}{\sqrt{2}}(w_j - \bar{w}_j), \quad j = 1, 2, \dots, n,$$

form an orthonormal frame of TU . We fix this notation throughout this section and use it without further notice.

For any $v \in TU$ with decomposition $v = v^{(1,0)} + v^{(0,1)} \in T^{1,0}U \oplus T^{0,1}U$, let $\bar{v}^{(1,0),*} \in T^{*0,1}U$ be the metric dual of $v^{(1,0)}$ with respect to $\langle \cdot, \cdot \rangle$. That is,

$$\bar{v}^{(1,0),*}(u) = \langle v^{(1,0)}, \bar{u} \rangle, \quad \forall u \in T^{0,1}U.$$

Then

$$(4.2) \quad c(v) = \sqrt{2}(\bar{v}^{(1,0),*} \wedge -i_{v^{(0,1)}})$$

defines the Clifford action v on $\Lambda(T^{*0,1}U) := \bigoplus_{q=0}^n T^{*0,q}U$, where \wedge and i denote the exterior and interior product respectively.

Let ∇^{TU} be the Levi-Civita connection on TU with respect to g^{TU} . Let ∇^{\det} be the Chern connection on $\det(T^{1,0}U)$ induced by $\langle \cdot, \cdot \rangle$, where $\det(T^{1,0}U)$ denotes the determinant line bundle of $T^{1,0}U$. Let $\Gamma^{TU} \in T^*U \otimes \text{End } TU$, $\Gamma^{\det} \in T^*U$ be the connection forms of ∇^{TU} , ∇^{\det} associated to the frames $\{e_j\}_{j=1}^{2n}$ and $w_1 \wedge \dots \wedge w_n$. That is,

$$(4.3) \quad \begin{aligned} \nabla_{e_j}^{TU} e_\ell &= \Gamma^{TU}(e_j) e_\ell, \quad j, \ell = 1, 2, \dots, 2n, \\ \nabla^{\det}(w_1 \wedge \dots \wedge w_n) &= \Gamma^{\det} w_1 \wedge \dots \wedge w_n. \end{aligned}$$

The Clifford connection ∇^{Cl} on $\Lambda(T^{*0,1}U)$ is defined for the frame

$$\{\bar{w}^{j_1} \wedge \dots \wedge \bar{w}^{j_q}; 1 \leq j_1 < \dots < j_q \leq n\}$$

by the local formula

$$(4.4) \quad \nabla^{\text{Cl}} = d + \frac{1}{4} \sum_{j,\ell=1}^{2n} \langle \Gamma^{TU} e_j, e_\ell \rangle c(e_j) c(e_\ell) + \frac{1}{2} \Gamma^{\det}.$$

By Proposition 1.3.1 in [27], ∇^{Cl} defines a Hermitian connection on $\Lambda(T^{*0,1}X)$.

Since E is rigid, we can consider E as a holomorphic vector bundle over U . We may assume that E is trivial on U . We will use the same notations as before. Let $\Omega^{0,q}(U, E)$ be the space of $(0, q)$ forms on

U with values in E , $q = 0, 1, 2, \dots, n$. Put

$$\begin{aligned}\Omega^{0,+}(U, E) &:= \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is even}} \Omega^{0,j}(U, E), \\ \Omega^{0,-}(U, E) &:= \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is odd}} \Omega^{0,j}(U, E), \\ \Omega^{0,*}(U, E) &:= \Omega^{0,+}(U, E \otimes L^m) \oplus \Omega^{0,-}(U, E).\end{aligned}$$

Since the Hermitian fiber metric $\langle \cdot | \cdot \rangle_E$ is rigid, we can consider $\langle \cdot | \cdot \rangle_E$ as a Hermitian fiber metric on the holomorphic line bundle E over U . Let ∇^E be the Chern connection on E induced by $\langle \cdot | \cdot \rangle_E$. We still denote by ∇^{Cl} the connection on $\Lambda(T^{*0,1}U) \otimes E$ induced by ∇^{Cl} and ∇^E .

Definition 4.1. The Spin^c Dirac operator D_B is defined by

$$(4.5) \quad D_B = \sum_{j=1}^{2n} c(e_j) \nabla_{e_j}^{\text{Cl}} : \Omega^{0,*}(U, E) \rightarrow \Omega^{0,*}(U, E).$$

It is well-known (see Proposition 1.3.1 and equation (1.3.1) in [27]) that D_B is formally self-adjoint and

$$(4.6) \quad \begin{aligned}D_B : \Omega^{0,+}(U, E) &\rightarrow \Omega^{0,-}(U, E), \\ D_B : \Omega^{0,-}(U, E) &\rightarrow \Omega^{0,+}(U, E).\end{aligned}$$

Let

$$\bar{\partial} : \Omega^{0,q}(U, E) \rightarrow \Omega^{0,q+1}(U), \quad q = 0, 1, 2, \dots, n-1,$$

be the Cauchy-Riemann operator. Let (\cdot, \cdot) be the L^2 inner product on $\Omega^{0,q}(U, E)$ induced by $\langle \cdot | \cdot \rangle_E$ and $\langle \cdot | \cdot \rangle_E$, $q = 0, 1, 2, \dots, n$, and let

$$\bar{\partial}^* : \Omega^{0,q+1}(U, E) \rightarrow \Omega^{0,q}(U, E), \quad q = 0, 1, 2, \dots, n-1,$$

be the formal adjoint of $\bar{\partial}$ with respect to (\cdot, \cdot) . By Theorem 1.4.5 in [27], we have

$$(4.7) \quad \begin{aligned}D_B &= \bar{\partial} + \bar{\partial}^* + A_B : \Omega^{0,+}(U, E) \rightarrow \Omega^{0,-}(U, E), \\ D_B &= \bar{\partial} + \bar{\partial}^* + A_B : \Omega^{0,-}(U, E) \rightarrow \Omega^{0,+}(U, E),\end{aligned}$$

where

$$\begin{aligned}A_B : \Omega^{0,+}(U, E) &\rightarrow \Omega^{0,-}(U, E), \\ A_B : \Omega^{0,-}(U, E) &\rightarrow \Omega^{0,+}(U, E),\end{aligned}$$

is a smooth zero order operator. It is clear that $A_B = A_B(z)$, that is A_B is independent of θ . Let $u \in \Omega^{0,q}(D, E)$, $q = 0, 1, 2, \dots, n$. Assume that $u = u(z)$, i.e. u is independent of θ . By using (2.13), from now on, we identify such u with an element in $\Omega^{0,q}(U, E)$. We need

Lemma 4.2. Let $B = (D, (z, \theta), \varphi)$ and $\tilde{B} = (D, (w, \eta), \tilde{\varphi})$ be two BRT trivializations. We assume that $D = U \times]-\varepsilon, \varepsilon[$, where $\varepsilon > 0$ and U is an open set of \mathbb{C}^n . Let $A_B, A_{\tilde{B}} : \Omega^{0,\pm}(U, E) \rightarrow \Omega^{0,\mp}(U, E)$ be the operators given by (4.7). Fix $m \in \mathbb{Z}$. Let $u \in \Omega_m^{0,\pm}(X, E)$. On D , we write $u = e^{-im\theta} v(z) = e^{-im\eta} \tilde{v}(w)$, $v(z), \tilde{v}(w) \in \Omega^{0,\pm}(U, E)$. Then,

$$(4.8) \quad e^{-im\theta} A_B(v(z)) = e^{-im\eta} A_{\tilde{B}}(\tilde{v}(w)) \quad \text{on } D.$$

Proof. It is straightforward to check that

$$(4.9) \quad \begin{aligned}w &= (w_1, \dots, w_n) = H(z) = (H_1(z), \dots, H_n(z)), \quad H_j(z) \in C^\infty, \quad \bar{\partial} H_j(z) = 0, \quad \forall j, \\ \eta &= \theta + G(z), \quad G(z) \in C^\infty.\end{aligned}$$

Since $\bar{\partial} H(z) = 0$, it is not difficult to see that

$$(4.10) \quad \begin{aligned}D_B &= D_{\tilde{B}} \quad \text{on } \Omega^{0,\pm}(U, E), \\ A_B &= A_{\tilde{B}} \quad \text{on } \Omega^{0,\pm}(U, E).\end{aligned}$$

Let $u \in \Omega_m^{0,\pm}(X, E)$. On D , write $u = e^{-im\theta}v(z) = e^{-im\eta}\tilde{v}(w)$, $v(z), \tilde{v}(w) \in \Omega^{0,\pm}(U, E)$. From (4.9), we can check that

$$(4.11) \quad v(z) = e^{-imG(z)}\tilde{v}(w).$$

From (4.11) and (4.10), we have

$$e^{-im\theta}A_B(v(z)) = e^{-im\theta}A_B(e^{-imG(z)}\tilde{v}(w)) = e^{-im\theta}A_{\tilde{B}}(e^{-imG(z)}\tilde{v}(w)) = e^{-im\eta}A_{\tilde{B}}(\tilde{v}(w)).$$

The lemma follows. \square

We can now introduce

Definition 4.3. For every $m \in \mathbb{Z}$, let $A_m : \Omega_m^{0,\pm}(X, E) \rightarrow \Omega_m^{0,\mp}(X, E)$ be the linear operator defined as follows. Let $u \in \Omega_m^{0,\pm}(X, E)$. Then, $v := A_m u$ is an element in $\Omega_m^{0,\mp}(X, E)$ such that for every BRT trivialization $B := (D, (z, \theta), \varphi)$, $D = U \times]-\varepsilon, \varepsilon[$, $\varepsilon > 0$, U is an open set in \mathbb{C}^n , we have $v|_D = e^{-im\theta}A_B(\tilde{u})(z)$, where $u = e^{-im\theta}\tilde{u}(z)$ on D , $\tilde{u} \in \Omega^{0,\pm}(U, E)$, and A_B is given by (4.7).

In view of Lemma 4.2, we see that Definition 4.3 is well-defined.

Until further notice, we fix $m \in \mathbb{Z}$. Consider

$$(4.12) \quad \begin{aligned} \tilde{D}_{b,m} &= \bar{\partial}_b + \bar{\partial}_b^* + A_m : \Omega_m^{0,+}(X, E) \rightarrow \Omega_m^{0,-}(X, E), \\ \tilde{D}_{b,m} &= \bar{\partial}_b + \bar{\partial}_b^* + A_m : \Omega_m^{0,-}(X, E) \rightarrow \Omega_m^{0,+}(X, E). \end{aligned}$$

Let

$$(4.13) \quad \tilde{D}_{b,m}^* : \Omega_m^{0,-}(X, E) \rightarrow \Omega_m^{0,+}(X, E),$$

be the formal adjoint of $\tilde{D}_{b,m} : \Omega_m^{0,+}(X, E) \rightarrow \Omega_m^{0,-}(X, E)$. Then,

$$\tilde{D}_{b,m}^* = \tilde{D}_{b,m} : \Omega_m^{0,-}(X, E) \rightarrow \Omega_m^{0,+}(X, E).$$

Put

$$(4.14) \quad \begin{aligned} \tilde{\square}_{b,m}^+ &:= \tilde{D}_{b,m}^* \tilde{D}_{b,m} = \tilde{D}_{b,m}^2 : \Omega_m^{0,+}(X, E) \rightarrow \Omega_m^{0,+}(X, E), \\ \tilde{\square}_{b,m}^- &:= \tilde{D}_{b,m} \tilde{D}_{b,m}^* = \tilde{D}_{b,m}^2 : \Omega_m^{0,-}(X, E) \rightarrow \Omega_m^{0,-}(X, E). \end{aligned}$$

We extend $\tilde{\square}_{b,m}^+$ and $\tilde{\square}_{b,m}^-$ to $L_m^{2,+}(X, E)$ and $L_m^{2,-}(X, E)$ by

$$(4.15) \quad \begin{aligned} \tilde{\square}_{b,m}^+ &: \text{Dom } \tilde{\square}_{b,m}^+ \subset L_m^{2,+}(X, E) \rightarrow L_m^{2,+}(X, E), \\ \tilde{\square}_{b,m}^- &: \text{Dom } \tilde{\square}_{b,m}^- \subset L_m^{2,-}(X, E) \rightarrow L_m^{2,-}(X, E), \end{aligned}$$

where

$$\begin{aligned} \text{Dom } \tilde{\square}_{b,m}^+ &:= \{u \in L_m^{2,+}(X, E); \tilde{\square}_{b,m}^+ u \in L_m^{2,+}(X, E)\}, \\ \text{Dom } \tilde{\square}_{b,m}^- &:= \{u \in L_m^{2,-}(X, E); \tilde{\square}_{b,m}^- u \in L_m^{2,-}(X, E)\}. \end{aligned}$$

For every $s \in \mathbb{R}$, put

$$\begin{aligned} H^{s,+}(X, E) &:= \bigoplus_{j \in \{0,1,2,\dots,q\}, j \text{ is even}} H^s(X, T^{*0,j} X \otimes E), \\ H^{s,-}(X, E) &:= \bigoplus_{j \in \{0,1,2,\dots,q\}, j \text{ is odd}} H^s(X, T^{*0,j} X \otimes E). \end{aligned}$$

We can repeat the proofs of Lemma 3.3 and Theorem 3.4 and conclude that

$$(4.16) \quad \begin{aligned} \text{Dom } \tilde{\square}_{b,m}^+ &= L_m^{2,+}(X, E) \cap H^{2,+}(X, E), \\ \text{Dom } \tilde{\square}_{b,m}^- &= L_m^{2,-}(X, E) \cap H^{2,-}(X, E), \\ \tilde{\square}_{b,m}^+ &\text{ and } \tilde{\square}_{b,m}^- \text{ are self-adjoint.} \end{aligned}$$

Let $\text{Spec } \tilde{\square}_{b,m}^+ \subset [0, \infty[$ and $\text{Spec } \tilde{\square}_{b,m}^- \subset [0, \infty[$ denote the spectrum of $\tilde{\square}_{b,m}^+$ and $\tilde{\square}_{b,m}^-$ respectively. We can repeat the proof of Theorem 3.5 with minor change and deduce that

Theorem 4.4. $\text{Spec } \tilde{\square}_{b,m}^+$ and $\text{Spec } \tilde{\square}_{b,m}^-$ are discrete subsets of $[0, \infty[$. For any $\mu \in \text{Spec } \tilde{\square}_{b,m}^+$, μ is an eigenvalue of $\tilde{\square}_{b,m}^+$ and the eigenspace

$$\tilde{\mathcal{E}}_{m,\nu}^+(X, E) := \{u \in \text{Dom } \tilde{\square}_{b,m}^+; \tilde{\square}_{b,m}^+ u = \nu u\}$$

is finite dimensional with $\tilde{\mathcal{E}}_{m,\nu}^+(X, E) \subset \Omega_m^{0,+}(X, E)$. Similarly, for any $\mu \in \text{Spec } \tilde{\square}_{b,m}^-$, μ is an eigenvalue of $\tilde{\square}_{b,m}^-$ and the eigenspace

$$\tilde{\mathcal{E}}_{m,\nu}^-(X, E) := \{u \in \text{Dom } \tilde{\square}_{b,m}^-; \tilde{\square}_{b,m}^- u = \nu u\}$$

is finite dimensional with $\tilde{\mathcal{E}}_{m,\nu}^-(X, E) \subset \Omega_m^{0,-}(X, E)$.

We need

Theorem 4.5. We have $\text{Spec } \tilde{\square}_{b,m}^+ \cap]0, \infty[= \text{Spec } \tilde{\square}_{b,m}^- \cap]0, \infty[$ and for every $\mu \in \text{Spec } \tilde{\square}_{b,m}^+$, $\mu \neq 0$, we have $\dim \tilde{\mathcal{E}}_{m,\mu}^+(X, E) = \dim \tilde{\mathcal{E}}_{m,\mu}^-(X, E)$.

Proof. Let $\mu \in \text{Spec } \tilde{\square}_{b,m}^+$, $\mu \neq 0$. Then, there is a function $f \in \Omega_m^{0,+}(X, E)$, $f \neq 0$, such that $\tilde{\square}_{b,m}^+ f = \mu f$. Consider $\tilde{D}_{b,m} f = g \in \Omega_m^{0,-}(X, E)$. Since $\mu \neq 0$, we can check that $g \neq 0$ and $\tilde{\square}_{b,m}^- g = \mu g$. Thus, $\mu \in \text{Spec } \tilde{\square}_{b,m}^-$. We have proved that

$$(4.17) \quad \text{Spec } \tilde{\square}_{b,m}^+ \cap]0, \infty[\subset \text{Spec } \tilde{\square}_{b,m}^- \cap]0, \infty[.$$

Conversely, let $\nu \in \text{Spec } \tilde{\square}_{b,m}^-$, $\nu \neq 0$. Then, there is a function $h \in \Omega_m^{0,-}(X, E)$, $h \neq 0$, such that $\tilde{\square}_{b,m}^- h = \nu h$. Consider $\tilde{D}_{b,m}^* h = \ell \in \Omega_m^{0,+}(X, E)$. Since $\nu \neq 0$, we can check that $\ell \neq 0$ and $\tilde{\square}_{b,m}^+ \ell = \nu \ell$. Thus, $\nu \in \text{Spec } \tilde{\square}_{b,m}^+$. We have proved that

$$(4.18) \quad \text{Spec } \tilde{\square}_{b,m}^- \cap]0, \infty[\subset \text{Spec } \tilde{\square}_{b,m}^+ \cap]0, \infty[.$$

From (4.17) and (4.18), we deduce that $\text{Spec } \tilde{\square}_{b,m}^+ \cap]0, \infty[= \text{Spec } \tilde{\square}_{b,m}^- \cap]0, \infty[$.

Let $\mu \in \text{Spec } \tilde{\square}_{b,m}^+$, $\mu \neq 0$. Consider the map:

$$F_\mu : \tilde{\mathcal{E}}_{m,\mu}^+(X, E) \rightarrow \tilde{\mathcal{E}}_{m,\mu}^-(X, E), \\ f \rightarrow \tilde{D}_{b,m} f.$$

It is clear that F_μ is injective. Let $g \in \tilde{\mathcal{E}}_{m,\mu}^-(X, E)$. Then, $\frac{1}{\mu} \tilde{D}_{b,m}^* g \in \tilde{\mathcal{E}}_{m,\mu}^+(X, E)$, $\frac{1}{\mu} \tilde{D}_{b,m}^* g \neq 0$ and $F_\mu(\frac{1}{\mu} \tilde{D}_{b,m}^* g) = g$. Thus, F_μ is bijective. Hence, $\dim \tilde{\mathcal{E}}_{m,\mu}^+(X, E) = \dim \tilde{\mathcal{E}}_{m,\mu}^-(X, E)$. The theorem follows. \square

We pause and introduce some notations. Let F be a complex vector bundle over X of rank r with a Hermitian metric $\langle \cdot | \cdot \rangle_F$. Let $A(x, y) \in C^\infty(X \times X, F_y \otimes F_x)$. Let $\{f_1, \dots, f_r\}$ be an orthonormal frame of F with respect to $\langle \cdot | \cdot \rangle_F$ on an open set $D \subset X$. On $D \times D$, we write $A(x, y) = (A_{j,k}(x, y))_{j,k=1}^r$, $A_{j,k}(x, y) \in C^\infty(D \times D)$, $j, k = 1, \dots, r$, in the sense that fixing $y_0 \in D$, for every $u = \sum_{j=1}^r u_j \otimes f_j(y_0) \in F_{y_0}$, $u_j \in \mathbb{C}$, $j = 1, 2, \dots, r$, we have

$$A(x, y_0)u = \sum_{j=1}^r (A_{j,1}(x, y_0)u_1 + \dots + A_{j,r}(x, y_0)u_r) \otimes f_j(x) \in F_x.$$

Let $u \in C_0^\infty(D, F)$. We write $u = \sum_{j=1}^r u_j \otimes f_j$, $u_j \in C_0^\infty(D)$, $j = 1, 2, \dots, r$. Define

$$(4.19) \quad \int A(x, y)u(y)dv_X(y) := \sum_{j=1}^r \int (A_{j,1}(x, y)u_1(y) + \dots + A_{j,r}(x, y)u_r(y))dv_X(y) \otimes f_j(x) \in C^\infty(D, F).$$

By using partition of unity, we can define $\int A(x, y)u(y)dv_X(y)$, for every $u \in C^\infty(X, F)$.

For every $\nu \in \text{Spec } \tilde{\square}_{b,m}^+$ and $\mu \in \text{Spec } \tilde{\square}_{b,m}^-$, let

$$(4.20) \quad \begin{aligned} P_{m,\nu}^+ &: L^{2,+}(X, E) \rightarrow \tilde{\mathcal{E}}_{m,\nu}^+(X, E), \\ P_{m,\mu}^- &: L^{2,+}(X, E) \rightarrow \tilde{\mathcal{E}}_{m,\mu}^-(X, E), \end{aligned}$$

be the orthogonal projections with respect to $(\cdot | \cdot)_E$ and let $P_{m,\nu}^+(x, y) \in C^\infty(X \times X, (T_y^{*0,+} X \otimes E_y) \boxtimes (T_x^{*0,+} X \otimes E_x))$, $P_{m,\mu}^-(x, y) \in C^\infty(X \times X, (T_y^{*0,-} X \otimes E_y) \boxtimes (T_x^{*0,-} X \otimes E_x))$ be the distribution kernels of $P_{m,\nu}^+$ and $P_{m,\mu}^-$ respectively. The heat kernels of $\tilde{\square}_{b,m}^+$ and $\tilde{\square}_{b,m}^-$ are given by

$$(4.21) \quad \begin{aligned} e^{-t\tilde{\square}_{b,m}^+}(x, y) &= P_{m,0}^+(x, y) + \sum_{\nu \in \text{Spec } \tilde{\square}_{b,m}^+, \nu > 0} e^{-\nu t} P_{m,\nu}^+(x, y), \\ e^{-t\tilde{\square}_{b,m}^-}(x, y) &= P_{m,0}^-(x, y) + \sum_{\mu \in \text{Spec } \tilde{\square}_{b,m}^-, \mu > 0} e^{-\mu t} P_{m,\mu}^-(x, y). \end{aligned}$$

Let $e^{-t\tilde{\square}_{b,m}^+} : \Omega^{0,+}(X, E) \rightarrow \Omega_m^{0,+}(X, E)$ and $e^{-t\tilde{\square}_{b,m}^-} : \Omega^{0,-}(X, E) \rightarrow \Omega_m^{0,-}(X, E)$ be the continuous operators given by

$$(4.22) \quad \begin{aligned} e^{-t\tilde{\square}_{b,m}^+} &: \Omega^{0,+}(X, E) \rightarrow \Omega_m^{0,+}(X, E), \\ u &\rightarrow \int e^{-t\tilde{\square}_{b,m}^+}(x, y) u(y) dv_X(y), \end{aligned}$$

$$(4.23) \quad \begin{aligned} e^{-t\tilde{\square}_{b,m}^-} &: \Omega^{0,-}(X, E) \rightarrow \Omega_m^{0,-}(X, E), \\ u &\rightarrow \int e^{-t\tilde{\square}_{b,m}^-}(x, y) u(y) dv_X(y). \end{aligned}$$

Let

$$(4.24) \quad \begin{aligned} Q_m^+ &: L^{2,+}(X, E) \rightarrow L_m^{2,+}(X, E), \\ Q_m^- &: L^{2,-}(X, E) \rightarrow L_m^{2,-}(X, E) \end{aligned}$$

be the orthogonal projections. From elementary Fourier analysis and (2.5), it is easy to see that

$$(4.25) \quad \begin{aligned} Q_m^+ &: \Omega^{0,+}(X, E) \rightarrow \Omega_m^{0,+}(X, E), \\ Q_m^- &: \Omega^{0,-}(X, E) \rightarrow \Omega_m^{0,-}(X, E), \end{aligned}$$

and

$$(4.26) \quad \begin{aligned} Q_m^+ u &= \frac{1}{2\pi} \int_{-\pi}^{\pi} u(e^{-i\theta} \circ x) e^{im\theta} d\theta, \quad \forall u \in \Omega^{0,+}(X, E), \\ Q_m^- u &= \frac{1}{2\pi} \int_{-\pi}^{\pi} u(e^{-i\theta} \circ x) e^{im\theta} d\theta, \quad \forall u \in \Omega^{0,-}(X, E). \end{aligned}$$

From (4.21), (4.22) and (4.23), it is straightforward to check that

$$(4.27) \quad \begin{aligned} \left(\frac{\partial}{\partial t} + \tilde{\square}_{b,m}^+\right)(e^{-t\tilde{\square}_{b,m}^+} u) &= 0, \quad \forall u \in \Omega^{0,+}(X, E), \quad \forall t > 0, \\ \lim_{t \rightarrow 0^+} (e^{-t\tilde{\square}_{b,m}^+} u) &= Q_m^+ u, \quad \forall u \in \Omega^{0,+}(X, E), \end{aligned}$$

and

$$(4.28) \quad \begin{aligned} \left(\frac{\partial}{\partial t} + \tilde{\square}_{b,m}^-\right)(e^{-t\tilde{\square}_{b,m}^-} u) &= 0, \quad \forall u \in \Omega^{0,-}(X, E), \quad \forall t > 0, \\ \lim_{t \rightarrow 0^+} (e^{-t\tilde{\square}_{b,m}^-} u) &= Q_m^- u, \quad \forall u \in \Omega^{0,-}(X, E). \end{aligned}$$

For every $\nu \in \text{Spec } \tilde{\square}_{b,m}^+$ and $\mu \in \text{Spec } \tilde{\square}_{b,m}^-$, let $\{f_1^\nu, \dots, f_{d_\nu}^\nu\}$ and $\{g_1^\mu, \dots, g_{d_\mu}^\mu\}$ be orthonormal frames for $\tilde{\mathcal{E}}_{m,\nu}^+(X, E)$ and $\tilde{\mathcal{E}}_{m,\mu}^-(X, E)$ respectively, where $d_\nu = \dim \tilde{\mathcal{E}}_{m,\nu}^+(X, E)$, $d_\mu = \dim \tilde{\mathcal{E}}_{m,\mu}^-(X, E)$. Define

$$(4.29) \quad \begin{aligned} \text{Tr } P_{m,\nu}^+(x, x) &:= \sum_{j=1}^{d_\nu} |f_j^\nu(x)|_E^2 \in C^\infty(X), \\ \text{Tr } P_{m,\mu}^-(x, x) &:= \sum_{j=1}^{d_\mu} |g_j^\mu(x)|_E^2 \in C^\infty(X). \end{aligned}$$

Let $e_1(x), \dots, e_d(x)$ and $f_1(x), \dots, f_{d_1}(x)$ be orthonormal frames of $T_x^{*0,+}X \otimes E_x$ and $T_x^{*0,-}X \otimes E_x$ respectively. We can check that

$$\begin{aligned} \text{Tr } P_{m,\nu}^+(x, x) &= \sum_{j=1}^d \langle P_{m,\nu}^+(x, x) e_j(x) | e_j(x) \rangle_E, \\ \text{Tr } P_{m,\mu}^-(x, x) &= \sum_{j=1}^{d_1} \langle P_{m,\mu}^-(x, x) f_j(x) | f_j(x) \rangle_E. \end{aligned}$$

Note that

$$(4.30) \quad \begin{aligned} \dim \tilde{\mathcal{E}}_{m,\nu}^+(X, E) &= \int_X \text{Tr } P_{m,\nu}^+(x, x) dv_X(x), \\ \dim \tilde{\mathcal{E}}_{m,\mu}^-(X, E) &= \int_X \text{Tr } P_{m,\mu}^-(x, x) dv_X(x). \end{aligned}$$

Put

$$(4.31) \quad \begin{aligned} \text{Tr } e^{-t\tilde{\square}_{b,m}^+}(x, x) &:= \text{Tr } P_{m,0}^+(x, x) + \sum_{\nu \in \text{Spec } \tilde{\square}_{b,m}^+, \nu > 0} e^{-\nu t} \text{Tr } P_{m,\nu}^+(x, x), \\ \text{Tr } e^{-t\tilde{\square}_{b,m}^-}(x, x) &:= \text{Tr } P_{m,0}^-(x, x) + \sum_{\nu \in \text{Spec } \tilde{\square}_{b,m}^-, \nu > 0} e^{-\nu t} \text{Tr } P_{m,\nu}^-(x, x). \end{aligned}$$

From (4.30) and (4.31), we deduce that for every $t > 0$,

$$(4.32) \quad \begin{aligned} \int_X \text{Tr } e^{-t\tilde{\square}_{b,m}^+}(x, x) dv_X(x) &= \dim \tilde{\mathcal{E}}_{m,0}^+(X, E) + \sum_{\nu \in \text{Spec } \tilde{\square}_{b,m}^+, \nu > 0} e^{-\nu t} \dim \tilde{\mathcal{E}}_{m,\nu}^+(X, E), \\ \int_X \text{Tr } e^{-t\tilde{\square}_{b,m}^-}(x, x) dv_X(x) &= \dim \tilde{\mathcal{E}}_{m,0}^-(X, E) + \sum_{\nu \in \text{Spec } \tilde{\square}_{b,m}^-, \nu > 0} e^{-\nu t} \dim \tilde{\mathcal{E}}_{m,\nu}^-(X, E). \end{aligned}$$

From Theorem 4.5 and (4.32), we conclude that

$$(4.33) \quad \int_X \left(\text{Tr } e^{-t\tilde{\square}_{b,m}^+}(x, x) - \text{Tr } e^{-t\tilde{\square}_{b,m}^-}(x, x) \right) dv_X(x) = \dim \text{Ker } \tilde{\square}_{b,m}^+ - \dim \text{Ker } \tilde{\square}_{b,m}^-.$$

We extend $\tilde{D}_{b,m}$ to $L_m^{2,+}(X, E)$:

$$\tilde{D}_{b,m} : \text{Dom } \tilde{D}_{b,m} \subset L_m^{2,+}(X, E) \rightarrow L_m^{2,-}(X, E),$$

where $\text{Dom } \tilde{D}_{b,m} = \{u \in L_m^{2,+}(X, E); D_{b,m}u \in L_m^{2,-}(X, E)\}$. We also write $\tilde{D}_{b,m}^*$ to denote the Hilbert space adjoint of $\tilde{D}_{b,m}$. It is not difficult to see that

$$(4.34) \quad \begin{aligned} \text{Ker } \tilde{D}_{b,m} &= \text{Ker } \tilde{\square}_{b,m}^+ \subset \Omega_m^{0,+}(X, E), \\ \text{Ker } \tilde{D}_{b,m}^* &= \text{Ker } \tilde{\square}_{b,m}^- \subset \Omega_m^{0,-}(X, E). \end{aligned}$$

Put $\text{ind } \widetilde{D}_{b,m} := \dim \text{Ker } \widetilde{D}_{b,m} - \dim \text{Ker } \widetilde{D}_{b,m}^*$. From (4.34) and (4.33), we deduce

Theorem 4.6. *For every $t > 0$, we have*

$$(4.35) \quad \text{ind } \widetilde{D}_{b,m} = \int_X \left(\text{Tr } e^{-t\widetilde{\square}_{b,m}^+}(x, x) - \text{Tr } e^{-t\widetilde{\square}_{b,m}^-}(x, x) \right) dv_X(x).$$

We have

Theorem 4.7. *We have $\text{ind } D_{b,m} = \text{ind } \widetilde{D}_{b,m}$.*

To prove Theorem 4.7, we need some preparations. For every $t \in [0, 1]$, put $L_t = \overline{\partial}_{b,m} + \overline{\partial}_{b,m}^* + tA_m : \Omega_m^{0,+}(X, E) \rightarrow \Omega_m^{0,-}(X, E)$. We extend L_t to $L_m^{2,+}(X, E)$:

$$L_t : \text{Dom } L_t \subset L_m^{2,+}(X, E) \rightarrow L_m^{2,-}(X, E),$$

where $\text{Dom } L_t = \left\{ u \in L_m^{2,+}(X, E); L_t u \in L_m^{2,-}(X, E) \right\}$. We also write L_t^* to denote the Hilbert space adjoint of L_t . It is clear that $L_0 = D_{b,m}$ and $L_1 = \widetilde{D}_{b,m}$. Let $H_m^{1,+}(X, E)$ be the completion of $\Omega_m^{0,+}(X, E)$ with respect to the Hermitian inner product

$$Q(u, v) = (u | v)_E + (\overline{\partial}_b u | \overline{\partial}_b v)_E + (\overline{\partial}_b^* u | \overline{\partial}_b^* v)_E.$$

Then, $H_m^{1,+}(X, E)$ is a Hilbert space and

$$(4.36) \quad H_m^{1,+}(X, E) = \text{Dom } L_t, \quad \forall t \in \mathbb{R}.$$

Consider $H_0 := H_m^{1,+}(X, E) \oplus \text{Ker } L_0^*$ and $H_1 = L_m^{2,-}(X, E) \oplus \text{Ker } L_0$. Let $(\cdot | \cdot)_{H_0}$ and $(\cdot | \cdot)_{H_1}$ be inner products on H_0 and H_1 respectively given by

$$\begin{aligned} ((f_1, g_1) | (f_2, g_2))_{H_0} &= Q(f_1, f_2) + (g_1 | g_2)_E, \quad (f_1, g_1), (f_2, g_2) \in H_m^{1,+}(X, E) \oplus \text{Ker } L_0^*, \\ ((\tilde{f}_1, \tilde{g}_1) | (\tilde{f}_2, \tilde{g}_2))_{H_1} &= (\tilde{f}_1 | \tilde{f}_2)_E + (\tilde{g}_1 | \tilde{g}_2)_E, \quad (\tilde{f}_1, \tilde{g}_1), (\tilde{f}_2, \tilde{g}_2) \in L_m^{2,-}(X, E) \oplus \text{Ker } L_0. \end{aligned}$$

With $(\cdot | \cdot)_{H_0}$ and $(\cdot | \cdot)_{H_1}$, H_0 and H_1 are Hilbert spaces. Let $\|\cdot\|_{H_0}$ and $\|\cdot\|_{H_1}$ denote the corresponding norms. Let $P_{\text{Ker } L_0}$ denote the orthogonal projection onto $\text{Ker } L_0$ with respect to $(\cdot | \cdot)_E$. Let $A_t : H_0 \rightarrow H_1$ be the linear map defined as follows. Let $(u, v) \in H_0 = H_m^{1,+}(X, E) \oplus \text{Ker } L_0^*$. Then,

$$A_t(u, v) = (L_t u + v, P_{\text{Ker } L_0} u) \in H_1 = L_m^{2,-}(X, E) \oplus \text{Ker } L_0.$$

Lemma 4.8. *There is a $r > 0$ such that $A_t : H_0 \rightarrow H_1$ is invertible, for every $0 \leq t \leq r$.*

Proof. We first claim that

$$(4.37) \quad A_0 \text{ is invertible.}$$

If $A_0(u, v) = 0$, for some $(u, v) \in H_0$. Then,

$$(4.38) \quad L_0 u = -v \in \text{Ker } L_0^*$$

and

$$(4.39) \quad P_{\text{Ker } L_0} u = 0.$$

From (4.38), we have

$$(4.40) \quad (L_0 u | L_0 u)_E = -(L_0 u | v)_E = -(u | L_0^* v)_E = 0.$$

Thus, $u \in \text{Ker } L_0$. From this and (4.39), we obtain $u = 0$ and hence $v = 0$. We have proved that A_0 is injective. We now prove that A_0 is surjective. Let $(a, b) \in H_1 = L_m^{2,-}(X, E) \oplus \text{Ker } L_0$. Since $L_0 : \text{Dom } L_0 \rightarrow L_m^{2,-}(X, E)$ has L^2 closed range, we have

$$(4.41) \quad a = L_0 \alpha + \beta, \quad \alpha \in H_m^{1,+}(X, E), \quad \alpha \perp \text{Ker } L_0, \quad \beta \in (\text{Rang } L_0)^\perp = \text{Ker } L_0^*.$$

From (4.41), we can check that

$$A_0(\alpha + b, \beta) = (L_0(\alpha + b) + \beta, P_{\text{Ker } L_0}(\alpha + b)) = (L_0 \alpha + \beta, b) = (a, b).$$

Thus, A_0 is surjective. The claim (4.37) follows.

Let $A_0^{-1} : H_1 \rightarrow H_0$ be the inverse of A_0 . From open mapping theorem, A_0^{-1} is continuous. It is clear that $A_t = A_0 + R_t$, where $R_t : H_0 \rightarrow H_1$ is continuous and there is a constant $c > 0$ such that $\|R_t u\|_{H_1} \leq ct \|u\|_{H_0}$, for every $u \in H_0$. Put

$$\begin{aligned} H_t &= I - A_0^{-1}R_t + (A_0^{-1}R_t)^2 - (A_0^{-1}R_t)^3 + \cdots, \\ \tilde{H}_t &= I - R_t A_0^{-1} + (R_t A_0^{-1})^2 - (R_t A_0^{-1})^3 + \cdots. \end{aligned}$$

Since A_0^{-1} is continuous, $H_t : H_0 \rightarrow H_0$ and $\tilde{H}_t : H_1 \rightarrow H_1$ are well-defined as continuous maps, for $t \geq 0$, t small. Moreover, we can check that $A_t \circ (H_t \circ A_0^{-1}) = I$ on H_1 and $(A_0^{-1} \circ \tilde{H}_t) \circ A_t = I$ on H_0 , for $t \geq 0$, t small. Thus, there is a $r > 0$ such that $A_t : H_0 \rightarrow H_1$ is invertible, for every $0 \leq t \leq r$. The lemma follows. \square

For every $t \in [0, 1]$, let

$$L_t^* : \text{Dom } L_t^* \subset L_m^{2,-}(X, E) \rightarrow L_m^{2,+}(X, E)$$

be the L^2 space adjoint of L_t with respect to $(\cdot | \cdot)_E$. It is clear that for every $t \in [0, 1]$, $\text{Ker } L_t \subset \Omega_m^{0,+}(X, E)$, $\text{Ker } L_t^* \subset \Omega_m^{0,-}(X, E)$ and $\dim \text{Ker } L_t < \infty$, $\dim \text{Ker } L_t^* < \infty$. Put $\text{ind } L_t := \dim \text{Ker } L_t - \dim \text{Ker } L_t^*$. We have

Lemma 4.9. *There is a $r_0 > 0$ such that $\text{ind } L_t = \text{ind } L_0$, for every $0 \leq t \leq r_0$.*

Proof. Let $r > 0$ be as in Lemma 4.8. We first claim that

$$(4.42) \quad \text{ind } L_0 \leq \text{ind } L_t, \quad \forall 0 \leq t \leq r.$$

Fix $0 \leq t \leq r$. Let

$$B : \text{Ker } L_t^* \oplus \text{Ker } L_0 \rightarrow \text{Ker } L_t \oplus \text{Ker } L_0^*$$

be the linear operator defined as follows. Let $(a, b) \in \text{Ker } L_t^* \oplus \text{Ker } L_0$. By Lemma 4.8,

$$A_t : H_m^{1,+}(X, E) \oplus \text{Ker } L_0^* \rightarrow L_m^{2,-}(X, E) \oplus \text{Ker } L_0$$

is invertible. There is a unique $(u, v) \in H_m^{1,+}(X, E) \oplus \text{Ker } L_0^* = \text{Dom } L_t \oplus \text{Ker } L_0^*$ such that $A_t(u, v) = (a, b)$. Let $P_{\text{Ker } L_t} : L_m^{2,+}(X, E) \rightarrow \text{Ker } L_t$ be the orthogonal projection with respect to $(\cdot | \cdot)_E$. Define $B(a, b) := (P_{\text{Ker } L_t} u, v) \in \text{Ker } L_t \oplus \text{Ker } L_0^*$. We now show that B is injective. Assume $B(a, b) = (0, 0)$ for some $(a, b) \in \text{Ker } L_t^* \oplus \text{Ker } L_0$. Let $(u, v) \in H_m^{1,+}(X, E) \oplus \text{Ker } L_0^*$ with $A_t(u, v) = (a, b)$. Then, $B(a, b) = (P_{\text{Ker } L_t} u, v) = (0, 0)$. Thus, $P_{\text{Ker } L_t} u = 0$ and $v = 0$. From the definition of A_t , we have

$$A_t(u, v) = A_t(u, 0) = (L_t u, P_{\text{Ker } L_0} u) = (a, b) \in \text{Ker } L_t^* \oplus \text{Ker } L_0.$$

Thus, $a = L_t u \in \text{Ker } L_t^*$. Hence, $(a | a)_E = (a | L_t u)_E = (L_t^* a | u)_E = 0$. We conclude that $u \in \text{Ker } L_t$ and $u = P_{\text{Ker } L_t} u = 0$. We obtain $(a, b) = (L_t u, P_{\text{Ker } L_0} u) = (0, 0)$ and hence B is injective. Since B is injective, we have

$$(4.43) \quad \dim \text{Ker } L_t^* + \dim \text{Ker } L_0 \leq \dim \text{Ker } L_t + \dim \text{Ker } L_0^*.$$

From (4.43), the claim (4.42) follows.

Similarly, we can repeat the procedure above and deduce that there is a $r_1 > 0$ such that

$$(4.44) \quad \text{ind } L_0^* \leq \text{ind } L_t^*, \quad \forall 0 \leq t \leq r_1.$$

It is not difficult to check that $\text{ind } L_t^* = -\text{ind } L_t$, for every $t \in [0, 1]$. From this observation and (4.44), we conclude that

$$(4.45) \quad \text{ind } L_0 \geq \text{ind } L_t, \quad \forall 0 \leq t \leq r_1.$$

From (4.45) and (4.42), the lemma follows. \square

Proof of Theorem 4.7. Let

$$I_0 := \{r \in [0, 1]; \text{ there is a } \varepsilon > 0 \text{ such that } \text{ind } L_t = \text{ind } L_0, \forall t \in (r - \varepsilon, r + \varepsilon) \cap [0, 1]\}.$$

From the definition of I_0 and Lemma 4.9, it is clear that I_0 is a non-empty open set in $[0, 1]$. Let $r_j \in I_0$, $j = 1, 2, \dots$, with $\lim_{j \rightarrow \infty} r_j = r_\infty \in [0, 1]$. We can repeat the proof of Lemma 4.9 and deduce that there is a $\varepsilon_0 > 0$ such that

$$(4.46) \quad \text{ind } L_t = \text{ind } L_{r_\infty}, \quad \forall t \in (r_\infty - \varepsilon_0, r_\infty + \varepsilon_0).$$

Since for j large, $r_j \in (r_\infty - \varepsilon_0, r_\infty + \varepsilon_0)$, we have

$$(4.47) \quad \text{ind } L_{r_j} = \text{ind } L_{r_\infty}, \quad \text{for } j \text{ large.}$$

Note that $\text{ind } L_{r_j} = \text{ind } L_0$, for each j . From this observation, (4.46) and (4.47), we deduce that

$$\text{ind } L_t = \text{ind } L_0, \quad \forall t \in (r_\infty - \varepsilon_0, r_\infty + \varepsilon_0).$$

Thus, $r_\infty \in I_0$ and hence I_0 is closed in $[0, 1]$. We conclude that $I_0 = [0, 1]$. Theorem 4.7 follows. \square

From Theorem 4.7, Theorem 4.6 and (3.25), we get

Theorem 4.10. *Fix $m \in \mathbb{Z}$. For every $t > 0$, we have*

$$(4.48) \quad \sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, E) = \int_X \left(\text{Tr } e^{-t\tilde{\square}_{b,m}^+}(x, x) - \text{Tr } e^{-t\tilde{\square}_{b,m}^-}(x, x) \right) dv_X(x).$$

Remark 4.11. Let $e^{-t\tilde{\square}_{b,m}^+}$ and $e^{-t\tilde{\square}_{b,m}^-}$ be the heat operators of $\square_{b,m}^+$ and $\square_{b,m}^-$ respectively. As Theorem 4.10, we have

$$\sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, E) = \int_X \left(\text{Tr } e^{-t\tilde{\square}_{b,m}^+}(x, x) - \text{Tr } e^{-t\tilde{\square}_{b,m}^-}(x, x) \right) dv_X(x).$$

When X is not CR Kähler, it is difficult to calculate $\text{Tr } e^{-t\tilde{\square}_{b,m}^+}(x, x) - \text{Tr } e^{-t\tilde{\square}_{b,m}^-}(x, x)$ and therefore we consider modified Kohn Laplacians $\tilde{\square}_{b,m}^+$, $\tilde{\square}_{b,m}^-$. It should be noticed that when X is CR Kähler, $\tilde{\square}_{b,m}^+ = \square_{b,m}^+$, $\tilde{\square}_{b,m}^- = \square_{b,m}^-$.

5. ASYMPTOTIC EXPANSIONS FOR THE HEAT KERNELS OF THE MODIFIED KOHN LAPLACIANS

5.1. Heat kernels of the modified Kodaira Laplacians on BRT trivializations. From now on, we fix $m \in \mathbb{Z}$. Let $B := (D, (z, \theta), \varphi)$ be a BRT trivialization. We may assume that $D = U \times] - \varepsilon, \varepsilon[$, where $\varepsilon > 0$ and U is an open set of \mathbb{C}^n . Since E is rigid, as in Section 4, we can consider E as a holomorphic vector bundle over U . We may assume that E is trivial on U . We will use the same notations as in Section 4. Consider $L \rightarrow U$ be a trivial line bundle with non-trivial Hermitian fiber metric $|\cdot|_{h^L}^2 = e^{-2\varphi}$. Let $(L^m, h^{L^m}) \rightarrow U$ be the m -th power of (L, h^L) . Let $\Omega^{0,q}(U, E \otimes L^m)$ be the space of $(0, q)$ forms on U with values in $E \otimes L^m$, $q = 0, 1, 2, \dots, n$. Put

$$\begin{aligned} \Omega^{0,+}(U, E \otimes L^m) &:= \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is even}} \Omega^{0,j}(U, E \otimes L^m), \\ \Omega^{0,-}(U, E \otimes L^m) &:= \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is odd}} \Omega^{0,j}(U, E \otimes L^m). \end{aligned}$$

Since L is trivial, from now on, we identify $\Omega^{0,q}(U, E)$ with $\Omega^{0,q}(U, E \otimes L^m)$, $q = 0, 1, 2, \dots, n$. Since the Hermitian fiber metric $\langle \cdot | \cdot \rangle_E$ is rigid, we can consider $\langle \cdot | \cdot \rangle_E$ as a Hermitian fiber metric on the holomorphic vector bundle E over U . Let $\langle \cdot, \cdot \rangle$ be the Hermitian metric on $\mathbb{C}TU$ given by

$$\left\langle \frac{\partial}{\partial z_j}, \frac{\partial}{\partial z_k} \right\rangle = \left\langle \frac{\partial}{\partial z_j} - i \frac{\partial \varphi}{\partial z_j}(z) \frac{\partial}{\partial \theta} \mid \frac{\partial}{\partial z_k} - i \frac{\partial \varphi}{\partial z_k}(z) \frac{\partial}{\partial \theta} \right\rangle, \quad j, k = 1, 2, \dots, n.$$

$\langle \cdot, \cdot \rangle$ induces Hermitian metrics on $T^{*0,q}U$ bundle of $(0, q)$ forms on U , $q = 0, 1, \dots, n$. We shall also denote the Hermitian metrics by $\langle \cdot, \cdot \rangle$. The Hermitian metrics on $T^{*0,q}U$ and E induce Hermitian

metrics on $T^{*0,q}U \otimes E$, $q = 0, 1, \dots, n$. We shall also denote these induced metrics by $\langle \cdot | \cdot \rangle_E$. Let (\cdot, \cdot) be the L^2 inner product on $\Omega^{0,q}(U, E)$ induced by $\langle \cdot | \cdot \rangle_E$, $q = 0, 1, 2, \dots, n$. Similarly, let $(\cdot, \cdot)_m$ be the L^2 inner product on $\Omega^{0,q}(U, E \otimes L^m)$ induced by $\langle \cdot | \cdot \rangle_E$ and h^{L^m} , $q = 0, 1, 2, \dots, n$.

Let

$$\bar{\partial} : \Omega^{0,q}(U, E \otimes L^m) \rightarrow \Omega^{0,q+1}(U, E \otimes L^m), \quad q = 0, 1, 2, \dots, n-1,$$

be the Cauchy-Riemann operator. Let

$$\bar{\partial}^{*,m} : \Omega^{0,q+1}(U, E \otimes L^m) \rightarrow \Omega^{0,q}(U, E \otimes L^m), \quad q = 0, 1, 2, \dots, n-1,$$

be the formal adjoint of $\bar{\partial}$ with respect to $(\cdot, \cdot)_m$. Put

$$(5.1) \quad D_{B,m} := \bar{\partial} + \bar{\partial}^{*,m} + A_B : \Omega^{0,+}(U, E \otimes L^m) \rightarrow \Omega^{0,-}(U, E \otimes L^m),$$

where $A_B : \Omega^{0,+}(U, E \otimes L^m) \rightarrow \Omega^{0,-}(U, E \otimes L^m)$ is as in (4.7). Let

$$(5.2) \quad D_{B,m}^* : \Omega^{0,-}(U, E \otimes L^m) \rightarrow \Omega^{0,+}(U, E \otimes L^m)$$

be the formal adjoint of $D_{B,m}$ with respect to $(\cdot, \cdot)_m$. Put

$$(5.3) \quad \begin{aligned} \tilde{\square}_{B,m}^+ &:= D_{B,m}^* D_{B,m} : \Omega^{0,+}(U, E \otimes L^m) \rightarrow \Omega^{0,+}(U, E \otimes L^m), \\ \tilde{\square}_{B,m}^- &:= D_{B,m} D_{B,m}^* : \Omega^{0,-}(U, E \otimes L^m) \rightarrow \Omega^{0,-}(U, E \otimes L^m). \end{aligned}$$

We need

Lemma 5.1. *Let $u \in \Omega_m^{0,+}(X, E)$, $v \in \Omega_m^{0,-}(X, E)$. On D , we write $u(z, \theta) = e^{-im\theta} \tilde{u}(z)$, $v(z, \theta) = e^{-im\theta} \tilde{v}(z)$, $\tilde{u}(z) \in \Omega^{0,+}(U, E)$, $\tilde{v}(z) \in \Omega^{0,-}(U, E)$. Then,*

$$(5.4) \quad \begin{aligned} e^{-m\varphi} \tilde{\square}_{B,m}^+ (e^{m\varphi} \tilde{u}) &= e^{im\theta} \tilde{\square}_{b,m}^+ (u), \\ e^{-m\varphi} \tilde{\square}_{B,m}^- (e^{m\varphi} \tilde{v}) &= e^{im\theta} \tilde{\square}_{b,m}^- (v). \end{aligned}$$

Proof. From (2.15), it is easy to see that for every $g \in \Omega^{0,q}(U, E \otimes L^m)$, we have

$$(5.5) \quad e^{im\theta} \bar{\partial}_b (e^{-im\theta} g(z)) = \bar{\partial}_m g(z),$$

where

$$(5.6) \quad \bar{\partial}_m = \bar{\partial} + m \sum_{j=1}^n \frac{\partial \varphi(z)}{\partial \bar{z}_j} d\bar{z}_j \wedge : \Omega^{0,q}(U, E \otimes L^m) \rightarrow \Omega^{0,q+1}(U, E \otimes L^m).$$

From (5.5) and (5.6), we have for every $h \in \Omega^{0,q+1}(U, E \otimes L^m)$,

$$(5.7) \quad e^{im\theta} \bar{\partial}_b^* (e^{-im\theta} h(z)) = \bar{\partial}_m^* h(z),$$

where

$$(5.8) \quad \bar{\partial}_m^* = \bar{\partial}^* + m \sum_{j=1}^n \frac{\partial \varphi(z)}{\partial z_j} (d\bar{z}_j)^{\wedge,*} : \Omega^{0,q+1}(U, E \otimes L^m) \rightarrow \Omega^{0,q}(U, E \otimes L^m).$$

Here $\bar{\partial}^* : \Omega^{0,q+1}(U, E \otimes L^m) \rightarrow \Omega^{0,q}(U, E \otimes L^m)$ is the formal adjoint of $\bar{\partial}$ with respect to (\cdot, \cdot) and $(d\bar{z}_j)^{\wedge,*} : T^{*0,q+1}U \rightarrow T^{*0,q}U$ is the adjoint of $(d\bar{z}_j) \wedge$. That is, $\langle d\bar{z}_j \wedge \alpha, \beta \rangle = \langle \alpha, (d\bar{z}_j)^{\wedge,*} \beta \rangle$, for every $\alpha \in T^{*0,q}U$, $\beta \in T^{*0,q+1}U$. Moreover, it is straightforward to check that

$$(5.9) \quad \bar{\partial}^{*,m} = \bar{\partial}^* + 2m \sum_{j=1}^n \frac{\partial \varphi(z)}{\partial z_j} (d\bar{z}_j)^{\wedge,*} = \bar{\partial}_m^* + m \sum_{j=1}^n \frac{\partial \varphi(z)}{\partial z_j} (d\bar{z}_j)^{\wedge,*}.$$

From (5.7), (5.8) and (5.9), we can check that for every $g \in \Omega^{0,q+1}(U, E \otimes L^m)$,

$$(5.10) \quad e^{-m\varphi} \bar{\partial}^{*,m} (e^{m\varphi} g) = \bar{\partial}_m^* g = e^{im\theta} \bar{\partial}_b^* (e^{-im\theta} g)$$

and hence

$$(5.11) \quad e^{-m\varphi} \bar{\partial} (e^{m\varphi} h) = e^{im\theta} \bar{\partial}_b (e^{-im\theta} h),$$

for every $h \in \Omega^{0,q}(U, E \otimes L^m)$. Let $u \in \Omega_m^{0,+}(X, E)$. On D , we write $u(z, \theta) = e^{-im\theta}\tilde{u}(z)$, $\tilde{u}(z) \in \Omega^{0,+}(U, E)$. From (5.10) and (5.11), we conclude that

$$(5.12) \quad e^{-m\varphi}(\bar{\partial} + \bar{\partial}^{*,m})(e^{m\varphi}\tilde{u}) = e^{im\theta}(\bar{\partial}_b + \bar{\partial}_b^*)(u).$$

Moreover, from the definition of the zero order operator $A_m : \Omega_m^{0,+}(X, E) \rightarrow \Omega_m^{0,-}(X, E)$ (see Definition 4.3), it is clear that

$$(5.13) \quad e^{-m\varphi}A_B(e^{m\varphi}\tilde{u}) = e^{im\theta}A_m(u).$$

From (5.11), (5.12) and (5.13), we get

$$e^{-m\varphi}D_{B,m}(e^{m\varphi}\tilde{u}) = e^{im\theta}\tilde{D}_{b,m}(u)$$

and hence

$$e^{-m\varphi}\tilde{\square}_{B,m}^+(e^{m\varphi}\tilde{u}) = e^{im\theta}\tilde{\square}_{b,m}^+(u).$$

Let $v \in \Omega_m^{0,-}(X, E)$. On D , we write $v(z, \theta) = e^{-im\theta}\tilde{v}(z)$, $\tilde{v}(z) \in \Omega^{0,-}(U, E)$. We can repeat the procedure above and deduce that

$$e^{-m\varphi}\tilde{\square}_{B,m}^-(e^{m\varphi}\tilde{v}) = e^{im\theta}\tilde{\square}_{b,m}^-(v).$$

The lemma follows. \square

We introduce some notations. Let M be a C^∞ orientable paracompact manifold and let F be a vector bundle over M . We need

Definition 5.2. Let $A(t, x) \in C^\infty(\mathbb{R}_+ \times M, F)$. We write

$$A(t, x) \sim t^k b_{-k}(x, t) + t^{k+1} b_{-k+1}(x, t) + t^{k+2} b_{-k+2}(x, t) + \cdots \quad \text{as } t \rightarrow 0^+,$$

$$b_j(x, t) \in C^\infty(M, F), b_j(x, t) \text{ is a } t\text{-dependent smooth function, } j = -k, -k+1, -k+2, \dots,$$

where $k \in \mathbb{Z}$, if for every compact set $K \Subset M$, every $m \in \mathbb{N}$ and every $N_0 > 0$, there are $C_{m,K,N_0} > 0$ and $\varepsilon_0 > 0$ independent of t such that

$$\left| A(t, x) - \sum_{j=0}^{N_0} t^{k+j} b_{-k+j}(x, t) \right|_{C^m(K)} \leq C_{m,K,N_0} t^{N_0+1}, \quad \forall 0 < t < \varepsilon_0.$$

Put

$$T^{*0,+}U := \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is even}} T^{*0,j}(U),$$

$$T^{*0,-}U := \bigoplus_{j \in \{0,1,\dots,n\}, j \text{ is odd}} T^{*0,j}(U).$$

Let $z, w \in U$ and let $T(z, w) \in (T_w^{*0,+}U \otimes E_w) \boxtimes (T_z^{*0,+}U \otimes E_z)$. We write $|T(z, w)|$ to denote the standard pointwise matrix norm of $T(z, w)$ induced by $\langle \cdot, \cdot \rangle$ and $\langle \cdot | \cdot \rangle_E$. Let $\Omega_0^{0,+}(U, E)$ and $\Omega_0^{0,-}(U, E)$ be the subspaces of $\Omega^{0,+}(U, E)$ and $\Omega^{0,-}(U, E)$ whose elements have compact support in U respectively. Let dv_U be the volume form on U induced by $\langle \cdot, \cdot \rangle$. Let $u \in \Omega_0^{0,+}(U, E)$. We define the integral $\int T(z, w)u(w)dv_U(w)$ as in (4.19). Let $G(t, z, w) \in C^\infty(\mathbb{R}_+ \times U \times U, (T_w^{*0,+}U \otimes E_w) \boxtimes (T_z^{*0,+}U \otimes E_z))$. We write $G(t)$ to denote the continuous operator

$$G(t) : \Omega_0^{0,+}(U, E) \rightarrow \Omega_0^{0,+}(U, E),$$

$$u \rightarrow \int G(t, z, w)u(w)dv_U(w)$$

and we write $G'(t)$ to denote the continuous operator

$$G'(t) : \Omega_0^{0,+}(U, E) \rightarrow \Omega_0^{0,+}(U, E),$$

$$u \rightarrow \int \frac{\partial G(t, z, w)}{\partial t} u(w)dv_U(w).$$

We consider the heat operators of $\tilde{\square}_{B,m}^+$ and $\tilde{\square}_{B,m}^-$. By using the Dirichlet heat kernel construction (see [19]), we deduce the following two theorems

Theorem 5.3. *There is $A_{B,+}(t, z, w) \in C^\infty(\mathbb{R}_+ \times U \times U, (T_w^{*0,+}U \otimes E_w) \boxtimes (T_z^{*0,+}U \otimes E_z))$ such that*

$$(5.14) \quad \begin{aligned} \lim_{t \rightarrow 0^+} A_{B,+}(t) &= I \text{ in } \mathcal{D}'(U, T^{*0,+}U \otimes E), \\ A'_{B,+}(t)u + A_{B,+}(t)(\tilde{\square}_{B,m}^+ u) &= 0, \quad \forall u \in \Omega_0^{0,+}(U, E), \quad \forall t > 0, \end{aligned}$$

and $A_{B,+}(t, z, w)$ satisfies the following: (I) For every compact set $K \Subset U$ and every $\alpha_1, \alpha_2, \beta_1, \beta_2 \in \mathbb{N}_0^n$, every $\gamma \in \mathbb{N}_0$, there are constants $C_{\gamma, \alpha_1, \alpha_2, \beta_1, \beta_2, K} > 0$ and $\varepsilon_0 > 0$ independent of t such that

$$(5.15) \quad \left| \partial_t^\gamma \partial_z^{\alpha_1} \partial_{\bar{z}}^{\alpha_2} \partial_w^{\beta_1} \partial_{\bar{w}}^{\beta_2} A_{B,+}(t, z, w) \right| \leq C_{\gamma, \alpha_1, \alpha_2, \beta_1, \beta_2, K} e^{-\varepsilon_0 \frac{|z-w|^2}{t}}, \quad \forall (t, z, w) \in \mathbb{R}_+ \times K \times K.$$

(II) Let $g \in \Omega_0^{0,+}(U, E)$. For every $\alpha_1, \alpha_2 \in \mathbb{N}_0^n$ and every compact set $K \Subset U$, there is a $C_{\alpha_1, \alpha_2, K} > 0$ independent of t such that

$$(5.16) \quad \begin{aligned} &\sup \{ |\partial_z^{\alpha_1} \partial_{\bar{z}}^{\alpha_2} (A_{B,+}(t)g)(z)| ; z \in K \} \\ &\leq C_{\alpha_1, \alpha_2, K} \sum_{\beta_1, \beta_2 \in \mathbb{N}_0^n, |\beta_1| + |\beta_2| \leq |\alpha_1| + |\alpha_2|} \sup \left\{ \left| \partial_z^{\beta_1} \partial_{\bar{z}}^{\beta_2} g(z) \right| ; z \in U \right\}. \end{aligned}$$

(III) $A_{B,+}(t, z, w)$ admits an asymptotic expansion:

$$(5.17) \quad \begin{aligned} A_{B,+}(t, z, w) &= e^{-\frac{h_+(z, w)}{t}} K_{B,+}(t, z, w), \\ K_{B,+}(t, z, w) &\sim t^{-n} b_n^+(z, w) + t^{-n+1} b_{n-1}^+(z, w) + \cdots + b_0^+(z, w) + t b_{-1}^+(z, w) + \cdots \text{ as } t \rightarrow 0^+, \\ b_s^+(z, w) &\in C^\infty(U \times U, (T_w^{*0,+}U \otimes E_w) \boxtimes (T_z^{*0,+}U \otimes E_z)), \quad s = n, n-1, n-2, \dots, \end{aligned}$$

where $h_+(z, w) \in C^\infty(U \times U, \mathbb{R}_+)$ with $h_+(z, z) = 0$ for every $z \in U$ and for every compact set $K \Subset U$, there is a constant $C_K > 1$ such that $\frac{1}{C_K} |z-w|^2 \leq h_+(z, w) \leq C_K |z-w|^2$.

Theorem 5.4. *There is $A_{B,-}(t, z, w) \in C^\infty(\mathbb{R}_+ \times U \times U, (T_w^{*0,-}U \otimes E_w) \boxtimes (T_z^{*0,-}U \otimes E_z))$ such that*

$$(5.18) \quad \begin{aligned} \lim_{t \rightarrow 0^+} A_{B,-}(t)u &= u \text{ in } \mathcal{D}'(U, T^{*0,-}U \otimes E), \\ A'_{B,-}(t)u + A_{B,-}(t)(\tilde{\square}_{B,m}^- u) &= 0, \quad \forall u \in \Omega_0^{0,-}(U, E), \quad \forall t > 0, \end{aligned}$$

and $A_{B,-}(t, z, w)$ satisfies the following: (I) For every compact set $K \Subset U$ and every $\alpha_1, \alpha_2, \beta_1, \beta_2 \in \mathbb{N}_0^n$, every $\gamma \in \mathbb{N}_0$, there are constants $C_{\gamma, \alpha_1, \alpha_2, \beta_1, \beta_2, K} > 0$ and $\varepsilon_0 > 0$ independent of t such that

$$(5.19) \quad \left| \partial_t^\gamma \partial_z^{\alpha_1} \partial_{\bar{z}}^{\alpha_2} \partial_w^{\beta_1} \partial_{\bar{w}}^{\beta_2} A_{B,-}(t, z, w) \right| \leq C_{\gamma, \alpha_1, \alpha_2, \beta_1, \beta_2, K} e^{-\varepsilon_0 \frac{|z-w|^2}{t}}, \quad \forall (t, z, w) \in \mathbb{R}_+ \times K \times K.$$

(II) Let $g \in \Omega_0^{0,-}(U, E)$. For every $\alpha_1, \alpha_2 \in \mathbb{N}_0^n$ and every compact set $K \Subset U$, there is a $C_{\alpha_1, \alpha_2, K} > 0$ independent of t such that

$$(5.20) \quad \begin{aligned} &\sup \{ |\partial_z^{\alpha_1} \partial_{\bar{z}}^{\alpha_2} (A_{B,-}(t)g)(z)| ; z \in K \} \\ &\leq C_{\alpha_1, \alpha_2, K} \sum_{\beta_1, \beta_2 \in \mathbb{N}_0^n, |\beta_1| + |\beta_2| \leq |\alpha_1| + |\alpha_2|} \sup \left\{ \left| \partial_z^{\beta_1} \partial_{\bar{z}}^{\beta_2} g(z) \right| ; z \in U \right\}. \end{aligned}$$

(III) $A_{B,-}(t, z, w)$ admits an asymptotic expansion:

$$(5.21) \quad \begin{aligned} A_{B,-}(t, z, w) &= e^{-\frac{h_-(z, w)}{t}} K_{B,-}(t, z, w), \\ K_{B,-}(t, z, w) &\sim t^{-n} b_n^-(z, w) + t^{-n+1} b_{n-1}^-(z, w) + \cdots + b_0^-(z, w) + t b_{-1}^-(z, w) + \cdots \text{ as } t \rightarrow 0^+, \\ b_s^-(z, w) &\in C^\infty(U \times U, (T_w^{*0,-}U \otimes E_w) \boxtimes (T_z^{*0,-}U \otimes E_z)), \quad s = n, n-1, n-2, \dots, \end{aligned}$$

where $h_-(z, w) \in C^\infty(U \times U, \mathbb{R}_+)$ with $h_-(z, z) = 0$ for every $z \in U$ and for every compact set $K \Subset U$, there is a constant $C_K > 1$ such that $\frac{1}{C_K} |z - w|^2 \leq h_-(z, w) \leq C_K |z - w|^2$.

Let $e_1(z), \dots, e_d(z)$ and $f_1(z), \dots, f_{d_1}(z)$ be orthonormal frames of $T_z^{*0,+}U \otimes E_z$ and $T_z^{*0,-}U \otimes E_z$ with respect to $\langle \cdot | \cdot \rangle_E$ respectively. Fix $j = n, n-1, \dots$. Let

$$b_j^+(z, z) \in C^\infty(U, \text{End}(T_z^{*0,+}U \otimes E_z)), \quad b_j^-(z, z) \in C^\infty(U, \text{End}(T_z^{*0,-}U \otimes E_z))$$

be as in (5.17) and (5.21) respectively. Define

$$(5.22) \quad \text{Tr } b_j^+(z, z) = \sum_{s=1}^d \langle b_j^+(z, z) e_s(z) | e_s(z) \rangle_E, \quad \text{Tr } b_j^-(z, z) = \sum_{s=1}^{d_1} \langle b_j^-(z, z) f_s(z) | f_s(z) \rangle_E.$$

From Theorem 1.3.5 and Theorem 1.4.5 in [27], we have Lichnerowicz formulas for $\tilde{\square}_{B,m}^+$ and $\tilde{\square}_{B,m}^-$. Thus, we can apply the rescaling technique in [2] and [8] and calculate $\text{Tr } b_j^+(z) - \text{Tr } b_j^-(z)$, $j = n, n-1, \dots, 0$. The calculation is the same as in the standard case and therefore we omit the detail. To state the result precisely, we introduce some notations. Let ∇^{TU} be the Levi-Civita connection on $\mathbb{C}TU$ with respect to $\langle \cdot, \cdot \rangle$. Let $P_{T^{1,0}U}$ be the natural projection from $\mathbb{C}TU$ onto $T^{1,0}U$. Then, $\nabla^{T^{1,0}U} := P_{T^{1,0}U} \nabla^{TU}$ is a connection on $T^{1,0}U$. Let $\nabla^{E \otimes L^m}$ be the connection on $E \otimes L^m \rightarrow U$ induce by $\langle \cdot, \cdot \rangle_E$ and h^{L^m} . Let $\Theta(\nabla^{T^{1,0}U}, T^{1,0}U) \in C^\infty(U, \Lambda^2(\mathbb{C}T^*U) \otimes \text{End}(T^{1,0}U))$ and $\Theta(\nabla^{E \otimes L^m}, E \otimes L^m) \in C^\infty(U, \Lambda^2(\mathbb{C}T^*U) \otimes \text{End}(E \otimes L^m))$ be the curvatures induced by $\nabla^{T^{1,0}U}$ and $\nabla^{E \otimes L^m}$ respectively. As in complex geometry, put

$$\begin{aligned} \text{Td}(\nabla^{T^{1,0}U}, T^{1,0}U) &= e^{\text{Tr}(h(\frac{i}{2\pi} \Theta(\nabla^{T^{1,0}U}, T^{1,0}U)))}, \quad h(z) = \log\left(\frac{z}{1-e^{-z}}\right), \\ \text{ch}(\nabla^{E \otimes L^m}, E \otimes L^m) &= \text{Tr}\left(\tilde{h}\left(\frac{i}{2\pi} \Theta(\nabla^{E \otimes L^m}, E \otimes L^m)\right)\right), \quad \tilde{h}(z) = e^z. \end{aligned}$$

We have

(5.23)

$$\left(\text{Tr } b_j^+(z, z) - \text{Tr } b_j^-(z, z)\right) = 0, \quad \forall z \in U, \quad j = n, n-1, \dots, 1,$$

$$\left(\text{Tr } b_0^+(z, z) - \text{Tr } b_0^-(z, z)\right) dv_U(z) = \left[\text{Td}(\nabla^{T^{1,0}U}, T^{1,0}U) \wedge \text{ch}(\nabla^{E \otimes L^m}, E \otimes L^m)\right]_{2n}(z), \quad \forall z \in U,$$

where $\left[\text{Td}(\nabla^{T^{1,0}U}, T^{1,0}U) \wedge \text{ch}(\nabla^{E \otimes L^m}, E \otimes L^m)\right]_{2n}$ denotes the $2n$ forms part of

$$\text{Td}(\nabla^{T^{1,0}U}, T^{1,0}U) \wedge \text{ch}(\nabla^{E \otimes L^m}, E \otimes L^m).$$

Let ∇^{TX} be the Levi-Civita connection on TX with respect to $\langle \cdot | \cdot \rangle$. Let $P_{T^{1,0}X}$ be the natural projection from $\mathbb{C}TX$ onto $T^{1,0}X$. Then, $\nabla^{T^{1,0}X} := P_{T^{1,0}X} \nabla^{TX}$ is a connection on $T^{1,0}X$. Let ∇^E be the connection on E induced by $\langle \cdot | \cdot \rangle_E$. Since $\langle \cdot | \cdot \rangle$ and $\langle \cdot | \cdot \rangle_E$ are rigid, we can check that $\nabla^{T^{1,0}X}$ and ∇^E are rigid. Moreover, it is straightforward to check that

$$(5.24) \quad \begin{aligned} \text{Td}(\nabla^{T^{1,0}U}, T^{1,0}U)(z) &= \text{Td}_b(\nabla^{T^{1,0}X}, T^{1,0}X)(z, \theta), \quad \forall (z, \theta) \in D, \\ \text{ch}(\nabla^{E \otimes L^m}, E \otimes L^m)(z) &= \left(\text{ch}_b(\nabla^E, E) \wedge e^{-m \frac{d\omega_0}{2\pi}}\right)(z, \theta), \quad \forall (z, \theta) \in D, \end{aligned}$$

and

$$(5.25) \quad \begin{aligned} &\left[\text{Td}(\nabla^{T^{1,0}U}, T^{1,0}U) \wedge \text{ch}(\nabla^{E \otimes L^m}, E \otimes L^m)\right]_{2n}(z) \wedge d\theta \\ &= \left[\text{Td}_b(\nabla^{T^{1,0}X}, T^{1,0}X) \wedge \text{ch}_b(\nabla^E, E) \wedge e^{-m \frac{d\omega_0}{2\pi}} \wedge \omega_0\right]_{2n+1}(z, \theta), \quad \forall (z, \theta) \in D, \end{aligned}$$

where $\left[\text{Td}_b(\nabla^{T^{1,0}X}, T^{1,0}X) \wedge \text{ch}_b(\nabla^E, E) \wedge e^{-m \frac{d\omega_0}{2\pi}} \wedge \omega_0\right]_{2n+1}$ denotes the $2n+1$ forms part of

$$\text{Td}_b(\nabla^{T^{1,0}X}, T^{1,0}X) \wedge \text{ch}_b(\nabla^E, E) \wedge e^{-m \frac{d\omega_0}{2\pi}} \wedge \omega_0.$$

From (5.23), (5.25) and note that $dv_U \wedge d\theta = dv_X$ on D , we get

Theorem 5.5. *With the notations above, we have*

$$(5.26) \quad \begin{aligned} & \left(\operatorname{Tr} b_j^+(z, z) - \operatorname{Tr} b_j^-(z, z) \right) = 0, \quad \forall z \in U, \quad j = n, n-1, \dots, 1, \\ & \left(\operatorname{Tr} b_0^+(z, z) - \operatorname{Tr} b_0^-(z, z) \right) dv_X(z, \theta) \\ & = \left[\operatorname{Td}_b(\nabla^{T^{1,0}X}, T^{1,0}X) \wedge \operatorname{ch}_b(\nabla^E, E) \wedge e^{-m \frac{d\omega_0}{2\pi}} \wedge \omega_0 \right]_{2n+1}(z, \theta), \quad \forall (z, \theta) \in D. \end{aligned}$$

5.2. Heat kernels of the modified Kohn Laplacians. From now on, we fix $m \in \mathbb{Z}$. Assume that $X = D_1 \cup D_2 \cup \dots \cup D_N$, where $B_j := (D_j, (z, \theta), \varphi_j)$ is a BRT trivialization, for each j . We may assume that for each j , $D_j = U_j \times]-\delta_j, \tilde{\delta}_j[\subset \mathbb{C}^n \times \mathbb{R}$, $\delta_j > 0$, $\tilde{\delta}_j > 0$, U_j is an open set in \mathbb{C}^n . Let $\chi_j \in C_0^\infty(D_j)$, $j = 1, 2, \dots, N$, with $\sum_{j=1}^N \chi_j = 1$ on X . Fix $j = 1, 2, \dots, N$. Put

$$K_j = \left\{ z \in U_j; \text{ there is a } \theta \in]-\delta_j, \tilde{\delta}_j[\text{ such that } \chi_j(z, \theta) \neq 0 \right\}.$$

Let $\tau_j(z) \in C_0^\infty(U_j)$ with $\tau_j \equiv 1$ on some neighborhood W_j of K_j . Let $\sigma_j \in C_0^\infty(]-\delta_j, \tilde{\delta}_j[)$ with $\int \sigma_j(\theta) d\theta = 1$. Let $A_{B_j, +}(t, z, w) \in C^\infty(\mathbb{R}_+ \times U_j \times U_j, (T_w^{*0,+} U_j \otimes E_w) \boxtimes (T_z^{*0,+} U_j \otimes E_z))$ be as in Theorem 5.3. Put

$$(5.27) \quad H_j(t, x, y) = \chi_j(x) e^{-m\varphi_j(z) - im\theta} A_{B_j, +}(t, z, w) e^{m\varphi_j(w) + im\eta} \tau_j(w) \sigma_j(\eta),$$

where $x = (z, \theta)$, $y = (w, \eta) \in \mathbb{C}^n \times \mathbb{R}$. Let $H_j(t)$ be the continuous operator

$$(5.28) \quad \begin{aligned} & H_j(t) : \Omega^{0,+}(X, E) \rightarrow \Omega^{0,+}(X, E), \\ & u \rightarrow \int \chi_j(x) e^{-m\varphi_j(z) - im\theta} A_{B_j, +}(t, z, w) e^{m\varphi_j(w) + im\eta} \tau_j(w) \sigma_j(\eta) u(y) dv_X(y). \end{aligned}$$

Consider

$$(5.29) \quad \Gamma(t) := \sum_{j=1}^N Q_m \circ H_j(t) \circ Q_m : \Omega^{0,+}(X, E) \rightarrow \Omega_m^{0,+}(X, E)$$

and let $\Gamma(t, x, y) \in C^\infty(\mathbb{R}_+ \times X \times X, (T_y^{*0,+} X \otimes E_y) \boxtimes (T_x^{*0,+} X \otimes E_x))$ be the distribution kernel of $\Gamma(t)$. Let $h_{j,+}(z, w)$, $A_{B_j, +}(t, z, w)$, $K_{B_j, +}$ and $b_{j,s}^+$, $s = n, n-1, \dots$, be as in (5.17). Let $\hat{\sigma}_j \in C_0^\infty(]-\delta_j, \tilde{\delta}_j[)$ such that $\hat{\sigma}_j = 1$ on some neighbourhood of $\operatorname{Supp} \sigma_j$ and $\hat{\sigma}_j(\theta) = 1$ if $(z, \theta) \in \operatorname{Supp} \chi_j$. Put

$$(5.30) \quad \begin{aligned} & \hat{h}_{j,+}(x, y) = \hat{\sigma}_j(\theta) h_{j,+}(z, w) \hat{\sigma}_j(\eta) \in C_0^\infty(D_j), \quad x = (z, \theta), \quad y = (w, \eta), \\ & \hat{H}_j(t, x, y) = \chi_j(x) e^{-m\varphi_j(z) - im\theta} K_{B_j, +}(t, z, w) e^{m\varphi_j(w) + im\eta} \tau_j(w) \sigma_j(\eta), \\ & \hat{b}_{j,s}^+(x, y) = \chi_j(x) e^{-m\varphi_j(z) - im\theta} b_{j,s}^+(z, w) e^{m\varphi_j(w) + im\eta} \tau_j(w) \sigma_j(\eta), \quad s = n, n-1, \dots \end{aligned}$$

It is not difficult to check that

$$(5.31) \quad \begin{aligned} \Gamma(t, x, y) &= \frac{1}{(2\pi)^2} \sum_{j=1}^N \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} H_j(t, e^{-i\theta} \circ x, e^{-iu} \circ y) e^{im(\theta-u)} d\theta du \\ &= \frac{1}{(2\pi)^2} \sum_{j=1}^N \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} e^{-\frac{\hat{h}_{j,+}(e^{-i\theta} \circ x, e^{-iu} \circ y)}{t}} \hat{H}_j(t, e^{-i\theta} \circ x, e^{-iu} \circ y) e^{im(\theta-u)} d\theta du \\ &\sim t^{-n} a_n^+(t, x, y) + t^{-n+1} a_{n-1}^+(t, x, y) + \dots \quad \text{as } t \rightarrow 0^+, \end{aligned}$$

where

$$(5.32) \quad \begin{aligned} & a_s^+(t, x, y) \\ &= \frac{1}{(2\pi)^2} \sum_{j=1}^N \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} e^{-\frac{\hat{h}_{j,+}(e^{-i\theta} \circ x, e^{-iu} \circ y)}{t}} \hat{b}_{j,s}^+(e^{-i\theta} \circ x, e^{-iu} \circ y) e^{im(\theta-u)} d\theta du, \quad s = n, n-1, n-2, \dots \end{aligned}$$

Lemma 5.6. *We have*

$$\lim_{t \rightarrow 0^+} \Gamma(t)u = Q_m u \text{ on } \mathcal{D}'(X, T^{*0,+} X \otimes E),$$

for every $u \in \Omega^{0,+}(X, T^{*0,+} X \otimes E)$.

Proof. Let $u \in \Omega^{0,+}(X, E)$. Fix $j = 1, 2, \dots, N$. On D_j , set $Q_m u = e^{-im\eta} v_j(w)$, $v_j(w) \in \Omega^{0,+}(U_j, E)$. From (5.28), we have

$$\begin{aligned} & \lim_{t \rightarrow 0^+} H_j(t) Q_m u \\ &= \lim_{t \rightarrow 0^+} \int \chi_j(x) e^{-m\varphi_j(z) - im\theta} A_{B_j,+}(t, z, w) e^{m\varphi_j(w) + im\eta} \tau_j(w) \sigma_j(\eta) e^{-im\eta} v_j(w) dv_{U_j}(w) d\eta \\ &= \lim_{t \rightarrow 0^+} \int \chi_j(x) e^{-m\varphi_j(z) - im\theta} A_{B_j,+}(t, z, w) e^{m\varphi_j(w)} \tau_j(w) v_j(w) dv_{U_j}(w) \\ &= \chi_j(x) e^{-m\varphi_j(z) - im\theta} e^{m\varphi_j(z)} \tau_j(z) v_j(z) \\ &= \chi e^{-im\theta} v_j = \chi_j Q_m u. \end{aligned}$$

Thus,

$$\lim_{t \rightarrow 0^+} \Gamma(t)u = \lim_{t \rightarrow 0^+} \sum_{j=1}^N Q_m \circ H_j(t) \circ Q_m u = \sum_{j=1}^N Q_m \chi_j Q_m u = Q_m u.$$

The lemma follows. □

We need

Lemma 5.7. *We have*

$$\Gamma'(t)u + \Gamma(t) \tilde{\square}_{b,m}^+ u = R(t)u, \quad \forall u \in \Omega^{0,+}(X, E),$$

where $R(t) : \Omega^{0,+}(X, E) \rightarrow \Omega_m^{0,+}(X, E)$ is a continuous operator with distribution kernel $R(t, x, y) \in C^\infty(\mathbb{R}_+ \times X \times X, (T_y^{*0,+} X \otimes E_y) \boxtimes (T_x^{*0,+} X \otimes E_x))$ and $R(t, x, y)$ satisfies the following: For every $\ell \in \mathbb{N}_0$, there is a $\varepsilon_0 > 0$ independent of t such that

$$(5.33) \quad \|R(t, x, y)\|_{C^\ell(X \times X)} \leq C_\ell e^{-\frac{\varepsilon_0}{t}}, \quad \forall t \in \mathbb{R}_+.$$

Proof. Let $u \in \Omega_m^{0,+}(X, E)$. Fix $j = 1, 2, \dots, N$. On D_j , set $u = e^{-im\eta}v_j(w)$, $v_j(w) \in \Omega^{0,+}(U_j, E)$. From (5.4), (5.14), we can check that

$$\begin{aligned}
& H'_j(t)u + H_j(t)\tilde{\square}_{b,m}^+ u \\
&= \int \chi_j(x)e^{-m\varphi_j(z)-im\theta} A'_{B_j,+}(t, z, w)e^{m\varphi_j(w)+im\eta}\tau_j(w)\sigma_j(\eta)u(y)dv_X(y) \\
&\quad + \int \chi_j(x)e^{-m\varphi_j(z)-im\theta} A_{B_j,+}(t, z, w)e^{m\varphi_j(w)+im\eta}\tau_j(w)\sigma_j(\eta)(\tilde{\square}_{b,m}^+ u)(y)dv_X(y) \\
&= \int \chi_j(x)e^{-m\varphi_j(z)-im\theta} A'_{B_j,+}(t, z, w)\tau_j(w)e^{m\varphi_j(w)}v_j(w)dv_{U_j}(w) \\
(5.34) \quad & + \int \chi_j(x)e^{-m\varphi_j(z)-im\theta} A_{B_j,+}(t, z, w)\tau_j(w)(\tilde{\square}_{B_j,m}^+(e^{m\varphi_j}v_j))(w)dv_{U_j}(w) \\
&= \int \chi_j(x)e^{-m\varphi_j(z)-im\theta} A'_{B_j,+}(t, z, w)\tau_j(w)e^{m\varphi_j(w)}v_j(w)dv_{U_j}(w) \\
&\quad + \int \chi_j(x)e^{-m\varphi_j(z)-im\theta} A_{B_j,+}(t, z, w)(\tilde{\square}_{B_j,m}^+(\tau_j e^{m\varphi_j}v_j))(w)dv_{U_j}(w) \\
&\quad + \int \chi(x)R_j(t, x, w)v_j(w)dv_{U_j}(w) \\
&= \int \chi(x)R_j(t, x, w)v_j(w)dv_{U_j}(w) = \int \chi(x)R_j(t, x, w)e^{im\eta}\sigma_j(\eta)u(y)dv_X(y),
\end{aligned}$$

where $R_j(t, x, w) \in C_0^\infty(\mathbb{R}_+ \times D_j \times U_j, (T^{*0,+}X \otimes E) \boxtimes (T^{*0,+}X \otimes E))$. Since $\tau_j(z) = 1$ if (z, θ) in some small neighborhood of χ_j , we can check that $R_j(t, x, w) = 0$ if (x, w) is in some small neighborhood of (z, z) . From this observation and (5.15), we conclude that for every $\ell \in \mathbb{N}_0$, there is a $\varepsilon > 0$ independent of t such that

$$(5.35) \quad \|R_j(t, x, w)\|_{C^\ell(X \times X)} \leq C_\ell e^{-\frac{\varepsilon}{t}}, \quad \forall t \in \mathbb{R}_+.$$

Put $\tilde{R}(t, x, y) := \sum_{j=1}^N R_j(t, x, w)e^{im\eta}\sigma_j(\eta) \in C^\infty(\mathbb{R}_+ \times X \times X, (T^{*0,+}X \otimes E) \boxtimes (T^{*0,+}X \otimes E))$ and set

$$R(t, x, y) = \frac{1}{(2\pi)^2} \sum_{j=1}^N \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} \tilde{R}(t, e^{-i\theta} \circ x, e^{-iu} \circ y) e^{im(\theta-u)} d\theta du.$$

Let $R(t) : \Omega^{0,+}(X, E) \rightarrow \Omega^{0,+}(X, E)$ be the continuous operator with distribution kernel $R(t, x, y)$. From (5.35), it is easy to see that $R(t, x, y)$ satisfies (5.33). Moreover, from (5.34), we see that $\Gamma'(t)u + \Gamma(t)\tilde{\square}_{b,m}^+ u = R(t)u$, $\forall u \in \Omega_m^{0,+}(X, E)$. The lemma follows. \square

Let $\Gamma^*(t) : \Omega^{0,+}(X, E) \rightarrow \Omega^{0,+}(X, E)$ and $R^*(t) : \Omega^{0,+}(X, E) \rightarrow \Omega^{0,+}(X, E)$ be the adjoints of $\Gamma(t)$ and $R(t)$ with respect to $(\cdot | \cdot)_E$ respectively. From Lemma 5.6 and Lemma 5.7, we deduce that

Theorem 5.8. *With the notations above, we have*

$$\lim_{t \rightarrow 0^+} \Gamma^*(t)u = Q_m u \text{ on } \mathcal{D}'(X, T^{*0,+}X \otimes E),$$

for every $u \in \Omega^{0,+}(X, T^{*0,+}X \otimes E)$, and

$$\frac{\partial \Gamma^*(t)}{\partial t} u + \tilde{\square}_{b,m}^+ \Gamma^*(t)u = R^*(t)u, \quad \forall u \in \Omega^{0,+}(X, E),$$

where $R^*(t) : \Omega^{0,+}(X, E) \rightarrow \Omega_m^{0,+}(X, E)$ is a continuous operator with distribution kernel $R^*(t, x, y) \in C^\infty(\mathbb{R}_+ \times X \times X, (T_y^{*0,+}X \otimes E_y) \boxtimes (T_x^{*0,+}X \otimes E_x))$ and $R^*(t, x, y)$ satisfies the following: For every $\ell \in \mathbb{N}_0$, there is a $\varepsilon_0 > 0$ independent of t such that

$$(5.36) \quad \|R^*(t, x, y)\|_{C^\ell(X \times X)} \leq C_\ell e^{-\frac{\varepsilon_0}{t}}, \quad \forall t \in \mathbb{R}_+.$$

We introduce some notations. Let $A(t), B(t) : \Omega^{0,+}(X, E) \rightarrow \Omega^{0,+}(X, E)$ be continuous operators with distribution kernels $A(t, x, y), B(t, x, y) \in C^\infty(\mathbb{R}_+ \times X \times X, (T_y^{*0,+} X \otimes E_y) \boxtimes (T_x^{*0,+} X \otimes E_x))$. Let $(A\sharp B)(t) : \Omega^{0,+}(X, E) \rightarrow \Omega^{0,+}(X, E)$ be the continuous operator with distribution kernel

$$\begin{aligned} (A\sharp B)(t, x, y) &:= \int_0^t \int_X A(t-s, x, z) B(s, z, y) dv_X(z) ds \\ &\in C^\infty(\mathbb{R}_+ \times X \times X, (T_y^{*0,+} X \otimes E_y) \boxtimes (T_x^{*0,+} X \otimes E_x)). \end{aligned}$$

Let $C(t) : \Omega^{0,+}(X, E) \rightarrow \Omega^{0,+}(X, E)$ be a continuous operator with distribution kernel $C(t, x, y) \in C^\infty(\mathbb{R}_+ \times X \times X, (T_y^{*0,+} X \otimes E_y) \boxtimes (T_x^{*0,+} X \otimes E_x))$. It is straightforward to check that $((A\sharp B)\sharp C)(t) = (A\sharp(B\sharp C))(t)$. We write

$$(A\sharp B\sharp C)(t) := (A\sharp(B\sharp C))(t).$$

Similarly, let $A_j(t) : \Omega^{0,+}(X, E) \rightarrow \Omega^{0,+}(X, E)$ be a continuous operator with distribution kernel $A_j(t, x, y) \in C^\infty(\mathbb{R}_+ \times X \times X, (T_y^{*0,+} X \otimes E_y) \boxtimes (T_x^{*0,+} X \otimes E_x))$, $j = 1, 2, \dots, N$. We write

$$(A_1\sharp A_2\sharp \dots \sharp A_N)(t) = (A_1\sharp(A_2\sharp A_3\sharp \dots \sharp A_N))(t).$$

We need

Theorem 5.9. Fix $\ell \in \mathbb{N}$, $\ell \geq 2$. Then, there is a $\epsilon > 0$ such that the sequence

$$(5.37) \quad \Lambda(t) := \Gamma^*(t) - (\Gamma^*\sharp R^*)(t) + (\Gamma^*\sharp R^*\sharp R^*)(t) - \dots$$

converges in $C^\ell((0, \epsilon) \times X \times X, (T^{*0,+} X \otimes E) \boxtimes (T^{*0,+} X \otimes E))$ and

$$(5.38) \quad \begin{aligned} \Lambda(t) : \Omega^{0,+}(X, E) &\rightarrow C^\ell(X, T^{*0,+} X \otimes E) \cap L_m^{2,+}(X, E), \\ \lim_{t \rightarrow 0^+} \Lambda(t)u &= Q_m u \text{ on } \mathcal{D}'(X, T^{*0,+} X \otimes E), \quad \forall u \in \Omega^{0,+}(X, T^{*0,+} X \otimes E), \\ \Lambda'(t)u + \tilde{\square}_{b,m}^+ \Lambda(t)u &= 0, \quad \forall u \in \Omega^{0,+}(X, E). \end{aligned}$$

Let $\Lambda(t, x, y) \in C^\ell((0, \epsilon) \times X \times X, (T_y^{*0,+} X \otimes E_y) \boxtimes (T_x^{*0,+} X \otimes E_x))$ be the distribution kernel of $\Lambda(t)$. Then, there is a $\epsilon_0 > 0$ independent of t such that

$$(5.39) \quad \|\Lambda(t, x, y) - \Gamma^*(t, x, y)\|_{C^\ell(X \times X)} \leq e^{-\frac{\epsilon_0}{t}}, \quad \forall t \in (0, \epsilon_0).$$

Proof. Fix $\ell \in \mathbb{N}$, $\ell \geq 2$. From (5.36), it is not difficult to see that there are $1 > \delta_0, \delta_1 > 0$ such that

$$(5.40) \quad \|R^*(t, x, y)\|_{C^\ell(X \times X)} \leq \min \left\{ \frac{1}{2} \frac{1}{\int_X dv_X}, e^{-\frac{\delta_1}{t}} \right\}, \quad \forall t \in (0, \delta_0).$$

From (5.40), we can check that for all $t \in (0, \delta_0)$,

$$(5.41) \quad \begin{aligned} \|(R^*\sharp R^*)(t, x, y)\|_{C^\ell(X \times X)} &\leq \frac{1}{2} e^{-\frac{\delta_1}{t}}, \quad \|(R^*\sharp R^*\sharp R^*)(t, x, y)\|_{C^\ell(X \times X)} \leq \frac{1}{2^2} e^{-\frac{\delta_1}{t}}, \\ \|(R^*\sharp R^*\sharp R^*\sharp R^*)(t, x, y)\|_{C^\ell(X \times X)} &\leq \frac{1}{2^3} e^{-\frac{\delta_1}{t}}, \quad \|(R^*\sharp R^*\sharp R^*\sharp R^*\sharp R^*)(t, x, y)\|_{C^\ell(X \times X)} \leq \frac{1}{2^4} e^{-\frac{\delta_1}{t}}, \dots \end{aligned}$$

From the construction of $\Gamma^*(t)$, (5.16) and (5.41), we conclude that for all $t \in (0, \delta_0)$,

$$(5.42) \quad \begin{aligned} \|(\Gamma^*\sharp R^*\sharp R^*)(t, x, y)\|_{C^\ell(X \times X)} &\leq \frac{C_1}{2} e^{-\frac{\delta_1}{t}}, \quad \|(\Gamma^*\sharp R^*\sharp R^*\sharp R^*)(t, x, y)\|_{C^\ell(X \times X)} \leq \frac{C_1}{2^2} e^{-\frac{\delta_1}{t}}, \\ \|(\Gamma^*\sharp R^*\sharp R^*\sharp R^*\sharp R^*)(t, x, y)\|_{C^\ell(X \times X)} &\leq \frac{C_1}{2^3} e^{-\frac{\delta_1}{t}}, \dots, \end{aligned}$$

where $C_1 > 0$ is a constant. From (5.42), it is easy to check that the sequence (5.37) converges in $C^\ell((0, \epsilon) \times X \times X, (T^{*0,+} X \otimes E) \boxtimes (T^{*0,+} X \otimes E))$ and (5.38), (5.39) hold. \square

Now, we can prove

Theorem 5.10. For every $\ell \in \mathbb{N}$, $\ell \geq 2$, there are $\epsilon_0 > 0$ and $\epsilon > 0$ independent of t such that

$$(5.43) \quad \left\| e^{-t\tilde{\square}_{b,m}^+}(x, y) - \Gamma(t, x, y) \right\|_{C^\ell(X \times X)} \leq e^{-\frac{\epsilon_0}{t}}, \quad \forall t \in (0, \epsilon),$$

where $e^{-t\tilde{\square}_{b,m}^+}(x, y)$ is as in (4.21). In particular,

(5.44)

$$e^{-t\tilde{\square}_{b,m}^+}(x, y) \sim t^{-n}a_n^+(t, x, y) + t^{-n+1}a_{n-1}^+(t, x, y) + \cdots + a_0^+(t, x, y) + ta_{-1}^+(t, x, y) + \cdots \quad \text{as } t \rightarrow 0^+,$$

$$a_s^+(t, x, y) = \frac{1}{(2\pi)^2} \sum_{j=1}^N \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} e^{-\frac{\hat{h}_{j,+}(e^{-i\theta} \circ x, e^{-iu} \circ y)}{t}} \hat{b}_{j,s}^+(e^{-i\theta} \circ x, e^{-iu} \circ y) e^{im(\theta-u)} d\theta du$$

$$\in C^\infty(\mathbb{R}_+ \times X \times X, (T_y^{*0,+} X \otimes E_y) \boxtimes (T_x^{*0,+} X \otimes E_x)), \quad s = n, n-1, n-2, \dots$$

Proof. Fix $\ell \in \mathbb{N}$, $\ell \geq 2$ and let $\Lambda(t) : \Omega^{0,+}(X, E) \rightarrow C^\ell(X, T^{*0,+} X \otimes E) \cap L_m^{2,+}(X, E)$ be as in Theorem 5.9 with distribution kernel $\Lambda(t, x, y) \in C^\ell((0, \epsilon) \times X \times X, (T_y^{*0,+} X \otimes E_y) \boxtimes (T_x^{*0,+} X \otimes E_x))$, where $\epsilon > 0$ is a constant dependent on ℓ . Let $f, g \in \Omega^{0,+}(X, E)$. From (4.27) and (5.38), it is not difficult to see that for every $0 < t < \epsilon$,

$$(5.45) \quad \begin{aligned} 0 &= \int_0^t \frac{\partial}{\partial s} \left((\Lambda(t-s)f | e^{-s\tilde{\square}_{b,m}^+} g)_E \right) ds \\ &= (Q_m f | e^{-t\tilde{\square}_{b,m}^+} g)_E - (\Lambda(t)f | Q_m g)_E \\ &= (f | e^{-t\tilde{\square}_{b,m}^+} g)_E - (\Lambda(t)f | g)_E \\ &= (e^{-t\tilde{\square}_{b,m}^+} f | g)_E - (\Lambda(t)f | g)_E. \end{aligned}$$

From (5.45), we conclude that $e^{-t\tilde{\square}_{b,m}^+}(x, y) = \Lambda(t, x, y)$ in $C^\ell((0, \epsilon) \times X \times X, (T_y^{*0,+} X \otimes E_y) \boxtimes (T_x^{*0,+} X \otimes E_x))$. From this, (5.39) and note that $e^{-t\tilde{\square}_{b,m}^+}$ is self-adjoint, the theorem follows. \square

We can repeat the proof of Theorem 5.10 and deduce that

Theorem 5.11. We have

(5.46)

$$e^{-t\tilde{\square}_{b,m}^-}(x, y) \sim t^{-n}a_n^-(t, x, y) + t^{-n+1}a_{n-1}^-(t, x, y) + \cdots + a_0^-(t, x, y) + ta_{-1}^-(t, x, y) + \cdots \quad \text{as } t \rightarrow 0^+,$$

$$a_s^-(t, x, y) = \frac{1}{(2\pi)^2} \sum_{j=1}^N \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} e^{-\frac{\hat{h}_{j,-}(e^{-i\theta} \circ x, e^{-iu} \circ y)}{t}} \hat{b}_{j,s}^-(e^{-i\theta} \circ x, e^{-iu} \circ y) e^{im(\theta-u)} d\theta du$$

$$\in C^\infty(\mathbb{R}_+ \times X \times X, (T_y^{*0,-} X \otimes E_y) \boxtimes (T_x^{*0,-} X \otimes E_x)), \quad s = n, n-1, n-2, \dots$$

Let $e_1(x), \dots, e_d(x)$ and $f_1(x), \dots, f_{d_1}(x)$ be orthonormal frames of $T_x^{*0,+} X \otimes E_x$ and $T_x^{*0,-} X \otimes E_x$ with respect to $\langle \cdot | \cdot \rangle_E$ respectively. Fix $\ell = n, n-1, \dots$, let

$$a_\ell^+(t, x, x) \in C^\infty(\mathbb{R}_+ \times X, \text{End}(T_x^{*0,+} X \otimes E_x)), \quad a_\ell^-(t, x, x) \in C^\infty(\mathbb{R}_+ \times X, \text{End}(T_x^{*0,-} X \otimes E_x))$$

be as in (5.44) and (5.46) respectively. Define

$$(5.47) \quad \text{Tr } a_\ell^+(t, x, x) = \sum_{s=1}^d \langle a_\ell^+(t, x, x) e_s(x) | e_s(x) \rangle_E, \quad \text{Tr } a_\ell^-(t, x, x) = \sum_{s=1}^{d_1} \langle a_\ell^-(t, x, x) f_s(x) | f_s(x) \rangle_E.$$

From Theorem 4.10, (5.44) and (5.46), we conclude that

Theorem 5.12. We have

$$(5.48) \quad \sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, E) = \lim_{t \rightarrow 0^+} \int_X \sum_{\ell=0}^n t^{-\ell} \left(\text{Tr } a_\ell^+(t, x, x) - \text{Tr } a_\ell^-(t, x, x) \right) dv_X(x).$$

6. LOCAL DENSITY FOR THE EULER CHARACTERISTIC $\sum_{j=0}^n (-1)^j \dim H_{b,m}^j(X, E)$

Put $X_\ell = \{x \in X; \text{the period of } x \text{ is } \frac{2\pi}{\ell}\}$ and let $p = \min \{\ell \in \mathbb{N}; X_\ell \neq \emptyset\}$. It is easy to see that X_p is an open subset of X . The goal of this section is to prove the following

Theorem 6.1. *Let $x_0 \in \overline{X_p}$. Then, for every $t > 0$,*

$$(6.1) \quad \begin{aligned} & \sum_{\ell=0}^n t^{-\ell} \left(\text{Tr } a_\ell^+(t, x_0, x_0) - \text{Tr } a_\ell^-(t, x_0, x_0) \right) dv_X(x_0) \\ &= \sum_{s=1}^p e^{\frac{2\pi(s-1)}{p} mi} \left[\text{Td}_b(\nabla^{T^{1,0}X}, T^{1,0}X) \wedge \text{ch}_b(\nabla^E, E) \wedge e^{-m \frac{d\omega_0}{2\pi}} \wedge \omega_0 \right]_{2n+1}(x_0), \end{aligned}$$

where $a_\ell^+(t, x, x) \in C^\infty(\mathbb{R}_+ \times X, \text{End}(T_x^{*0,+}X \otimes E_x))$, $a_\ell^-(t, x, x) \in C^\infty(\mathbb{R}_+ \times X, \text{End}(T_x^{*0,-}X \otimes E_x))$ are as in (5.44) and (5.46).

Proof. Fix $x_0 \in X_p$. Let $B_j = (D_j, (z, \theta), \varphi_j)$, $\chi_j(x)$, σ_j and τ_j , $j = 1, 2, \dots, N$, be as in the beginning of Section 5.2. We will use the same notations as in Section 5.2. Fix $j = 1, 2, \dots, N$. Assume that $e^{-i\theta_0} \circ x_0 \in D_j$ for some $\theta_0 \in]-\pi, \pi]$ and suppose that $e^{-i\theta_0} \circ x_0 = (z_0, 0) \in D_j$. Let $\gamma > 0$ be a small constant so that $\{z \in \mathbb{C}^n; |z - z_0| < \gamma\} \subset W_j$ and the canonical coordinates (z, θ) can be defined on

$$(6.2) \quad \hat{D}_0 := \left\{ (z, \theta) \in \mathbb{C}^n \times \mathbb{R}; |z - z_0| < \gamma, \theta \in \left[-\frac{\pi}{p}, \frac{\pi}{p}\right] \right\}$$

(this is always possible, see Lemma 1.17 in [22]), where $W_j \subset U_j$ is as in the beginning of Section 5.2. Let $\psi(x) = \psi(z, \theta) \in C_0^\infty(\hat{D}_0)$ with $\int \psi(x) dv_X(x) = 1$. For every $\varepsilon > 0$, put $\psi_\varepsilon(x) := \varepsilon^{-(2n+1)} \psi(z_0 + \frac{z-z_0}{\varepsilon}, \frac{\theta}{\varepsilon})$. Then,

$$(6.3) \quad f(e^{-i\theta_0} \circ x_0) = \lim_{\varepsilon \rightarrow 0^+} \int f(x) \psi_\varepsilon(x) dv_X(x), \quad f \in C^\infty(X).$$

From (6.3), we have

$$(6.4) \quad \begin{aligned} & \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} H_j(t, e^{-i\theta} \circ x_0, e^{-iu} \circ x_0) e^{im(\theta-u)} d\theta du \\ &= \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} H_j(t, e^{-i\theta} \circ e^{-i\theta_0} \circ x_0, e^{-iu} \circ e^{-i\theta_0} \circ x_0) e^{im(\theta-u)} d\theta du \\ &= \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} \left(\lim_{\delta \rightarrow 0^+} \lim_{\varepsilon \rightarrow 0^+} \int H_j(t, e^{-i\theta} \circ x, e^{-iu} \circ y) e^{im(\theta-u)} \psi_\varepsilon(x) \psi_\delta(y) dv_X(x) dv_X(y) \right) d\theta du. \end{aligned}$$

Fix $\theta, u \in [-\pi, \pi]$, from (5.27), we have

$$(6.5) \quad \begin{aligned} & \lim_{\delta \rightarrow 0^+} \lim_{\varepsilon \rightarrow 0^+} \int H_j(t, e^{-i\theta} \circ x, e^{-iu} \circ y) e^{im(\theta-u)} \psi_\varepsilon(x) \psi_\delta(y) dv_X(x) dv_X(y) \\ &= \lim_{\delta \rightarrow 0^+} \lim_{\varepsilon \rightarrow 0^+} \int H_j(t, x, y) e^{im(\theta-u)} \psi_\varepsilon(e^{i\theta} \circ x) \psi_\delta(e^{iu} \circ y) dv_X(x) dv_X(y) \\ &= \lim_{\delta \rightarrow 0^+} \lim_{\varepsilon \rightarrow 0^+} \int H_j(t, (z, \xi), (w, \eta)) e^{im(\theta-u)} \varepsilon^{-(2n+1)} \psi\left(z_0 + \frac{z-z_0}{\varepsilon}, \frac{\xi-\theta}{\varepsilon}\right) \\ & \quad \times \delta^{-(2n+1)} \psi\left(z_0 + \frac{w-z_0}{\delta}, \frac{\eta-u}{\delta}\right) dv_X(z, \xi) dv_X(w, \eta) \\ &= H_j(t, (z_0, \theta), (z_0, u)) e^{im(\theta-u)}, \end{aligned}$$

where $x = (z, \xi)$ and $y = (w, \eta)$. From (5.27), (6.4) and (6.5), we have

$$\begin{aligned}
& \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} H_j(t, e^{-i\theta} \circ x_0, e^{-iu} \circ x_0) e^{im(\theta-u)} d\theta du \\
&= \left(\sum_{s=1}^p e^{im\frac{2\pi}{p}(s-1)} \right) \left(\sum_{k=1}^p e^{-im\frac{2\pi}{p}(k-1)} \right) \int_{-\frac{\pi}{p}}^{\frac{\pi}{p}} \int_{-\frac{\pi}{p}}^{\frac{\pi}{p}} H_j(t, (z_0, \theta), (z_0, u)) e^{im(\theta-u)} d\theta du \\
(6.6) \quad &= \left(\sum_{s=1}^p e^{im\frac{2\pi}{p}(s-1)} \right) \left(\sum_{k=1}^p e^{-im\frac{2\pi}{p}(k-1)} \right) \int_{-\frac{\pi}{p}}^{\frac{\pi}{p}} \chi_j(z_0, \theta) K_{B_{j,+}}(t, z_0, z_0) d\theta \\
&= \left(\sum_{s=1}^p e^{im\frac{2\pi}{p}(s-1)} \right) \left(\sum_{k=1}^p e^{-im\frac{2\pi}{p}(k-1)} \right) \int_{-\frac{\pi}{p}}^{\frac{\pi}{p}} \chi_j(e^{-i\theta} \circ x_0) K_{B_{j,+}}(t, z_0, z_0) d\theta.
\end{aligned}$$

From (5.30), (5.31), (5.32) and (6.6), we conclude that

$$\begin{aligned}
(6.7) \quad & \sum_{\ell=0}^n t^{-\ell} \text{Tr } a_{\ell}^{+}(t, x_0, x_0) \\
&= \frac{1}{(2\pi)^2} \left(\sum_{s=1}^p e^{im\frac{2\pi}{p}(s-1)} \right) \left(\sum_{k=1}^p e^{-im\frac{2\pi}{p}(k-1)} \right) \sum_{\ell=0}^n t^{-\ell} \int_{-\frac{\pi}{p}}^{\frac{\pi}{p}} \left(\sum_{j=1}^N \text{Tr } b_{j,\ell}^{+}(z_0, z_0) \chi_j(e^{-i\theta} \circ x_0) \right) d\theta.
\end{aligned}$$

We can repeat the proof of (6.7) and get that

$$\begin{aligned}
(6.8) \quad & \sum_{\ell=0}^n t^{-\ell} \text{Tr } a_{\ell}^{-}(t, x_0, x_0) \\
&= \frac{1}{(2\pi)^2} \left(\sum_{s=1}^p e^{im\frac{2\pi}{p}(s-1)} \right) \left(\sum_{k=1}^p e^{-im\frac{2\pi}{p}(k-1)} \right) \sum_{\ell=0}^n t^{-\ell} \int_{-\frac{\pi}{p}}^{\frac{\pi}{p}} \left(\sum_{j=1}^N \text{Tr } b_{j,\ell}^{-}(z_0, z_0) \chi_j(e^{-i\theta} \circ x_0) \right) d\theta.
\end{aligned}$$

From (6.7), (6.8), Theorem 5.5 and notice that $\sum_{j=1}^N \chi_j(e^{-i\theta} \circ x_0) = 1$, for every $\theta \in [-\frac{\pi}{p}, \frac{\pi}{p}]$, we have

$$\begin{aligned}
& \sum_{\ell=0}^n t^{-\ell} \left(\text{Tr } a_{\ell}^{+}(t, x_0, x_0) - \text{Tr } a_{\ell}^{-}(t, x_0, x_0) \right) dv_X(x_0) \\
&= \frac{1}{(2\pi)^2} \left(\sum_{j=1}^p e^{im\frac{2\pi}{p}(j-1)} \right) \left(\sum_{k=1}^p e^{-im\frac{2\pi}{p}(k-1)} \right) \frac{2\pi}{p} \\
&\quad \times \frac{\left[\text{Td}_b(\nabla^{T^{1,0}X}, T^{1,0}X) \wedge \text{ch}_b(\nabla^E, E) \wedge e^{-m\frac{d\omega_0}{2\pi}} \wedge \omega_0 \right]_{2n+1}(x_0)}{dv_X(x_0)}.
\end{aligned}$$

We have proved that (6.1) holds for every $x \in X_p$. Since the left side and the right side of (6.1) are continuous functions on X , we conclude that (6.1) holds for every $x \in \overline{X_p}$ and the theorem follows. \square

Now, we can prove

Theorem 6.2. *Recall that we work with the assumption that X is connected. Then, X_p is dense in X .*

Proof. Assume that $\overline{X_p} \neq X$. For $\ell > p$, $\ell \in \mathbb{N}$, put $\widetilde{X}_{\ell} = \{x \in X \setminus \overline{X_p}; \text{ the period of } x \text{ is } \frac{2\pi}{\ell}\}$. Assume that

$$X = \overline{X_p} \cup \widetilde{X}_{\ell_1} \cup \widetilde{X}_{\ell_2} \cup \cdots \cup \widetilde{X}_{\ell_N},$$

where $p < \ell_1 < \ell_2 < \cdots < \ell_N$. Since $\overline{X_p}$ and $\widetilde{X}_{\ell_2} \cup \cdots \cup \widetilde{X}_{\ell_N}$ are closed sets, \widetilde{X}_{ℓ_1} is an open set. We claim that

$$(6.9) \quad \overline{\widetilde{X}_{\ell_1}} \cap \overline{X_p} = \emptyset,$$

where $\overline{\widehat{X}_{\ell_1}}$ denotes the closure of \widehat{X}_{ℓ_1} in X . If $\overline{\widehat{X}_{\ell_1}} \cap \overline{X_p} \neq \emptyset$. Let $x_0 \in \overline{\widehat{X}_{\ell_1}} \cap \overline{X_p}$. It is straightforward to see that we can find rigid Hermitian metric $\langle \cdot | \cdot \rangle$ on $\mathcal{C}TX$ and rigid Hermitian metric $\langle \cdot | \cdot \rangle_E$ on E such that

$$(6.10) \quad \frac{\left[\text{Td}_b(\nabla^{T^{1,0}X}, T^{1,0}X) \wedge \text{ch}_b(\nabla^E, E) \wedge e^{-m \frac{d\omega_0}{2\pi}} \wedge \omega_0 \right]_{2n+1}(x_0)}{dv_X(x_0)} \neq 0.$$

Since $x_0 \in \overline{\widehat{X}_{\ell_1}}$ and \widehat{X}_{ℓ_1} is open, we can repeat the proof of Theorem 6.1 and conclude that

$$(6.11) \quad \begin{aligned} & \sum_{\ell=0}^n t^{-\ell} \left(\text{Tr } a_{\ell}^+(t, x_0, x_0) - \text{Tr } a_{\ell}^-(t, x_0, x_0) \right) dv_X(x_0) \\ &= \sum_{s=1}^{\ell_1} e^{\frac{2\pi(s-1)}{\ell_1} mi} \left[\text{Td}_b(\nabla^{T^{1,0}X}, T^{1,0}X) \wedge \text{ch}_b(\nabla^E, E) \wedge e^{-m \frac{d\omega_0}{2\pi}} \wedge \omega_0 \right]_{2n+1}(x_0). \end{aligned}$$

From (6.10), (6.11) and (6.1), we get a contradiction. The claim (6.9) follows.

Assume that $\overline{X_p} \cup \overline{\widehat{X}_{\ell_1}} \neq X$. For $\ell > \ell_1$, $\ell \in \mathbb{N}$, put

$$\hat{X}_{\ell} = \left\{ x \in X \setminus (\overline{X_p} \cup \overline{\widehat{X}_{\ell_1}}); \text{ the period of } x \text{ is } \frac{2\pi}{\ell} \right\}.$$

Assume that $X = \overline{X_p} \cup \overline{\widehat{X}_{\ell_1}} \cup \hat{X}_{k_1} \cup \hat{X}_{k_2} \cup \dots \cup \hat{X}_{k_M}$, where $p < \ell_1 < k_1 < k_2 < \dots < k_M$. We can repeat the proof of (6.9) with minor change and get that

$$\overline{\hat{X}_{k_1}} \cap (\overline{X_p} \cup \overline{\widehat{X}_{\ell_1}}) = \emptyset.$$

Continuing in this way, we can find closed sets C_1, \dots, C_K , with $\overline{X_p} \cap (\bigcup_{j=1}^K C_j) = \emptyset$ such that

$$X = \overline{X_p} \cup C_1 \cup \dots \cup C_K.$$

Hence, $\overline{X_p}$ is an open set and also a closed set. Since X is connected, we get a contradiction. Thus, $X = \overline{X_p}$. \square

Proof of Theorem 1.3. From Theorem 5.12, Theorem 6.1 and Theorem 6.2, we get Theorem 1.3. \square

REFERENCES

- [1] M.-S. Baouendi and L.-P. Rothschild and F.-Treves, *CR structures with group action and extendability of CR functions*, Invent. Math., **83** (1985), 359–396.
- [2] N. Berline, E. Getzler and M. Vergne, *Heat Kernels and Dirac Operators*, volume 298 of Grundlehren der mathematischen Wissenschaften. Springer-Verlag, Berlin, Heidelberg, New York, 1992.
- [3] L. Boutet de Monvel, *Intégration des équations de Cauchy-Riemann induites formelles*, Séminaire Goulaouic-Lions-Schwartz 1974–1975; Équations aux dérivées partielles linéaires et non linéaires, Centre Math., École Polytech., Paris, 1975, Exp. no. 9, pp. 13.
- [4] E. Brieskorn and A. Van de Ven, *Some complex structures on products of homotopy spheres*, Topology, vol. 7(1968), 389–393.
- [5] S.-C. Chen and M.-C. Shaw, *Partial differential equations in several complex variables*, AMS/IP Studies in Advanced Mathematics, 19, American Mathematical Society, Providence, RI; International Press, Boston, MA, 2001.
- [6] S.-S. Chern and R.-S. Hamilton, *On Riemannian metrics adapted to three-dimensional contact manifolds*, Lecture Notes in Math., **1111**, 279–305.
- [7] E.-B. Davies, *Spectral Theory and Differential Operators*, Cambridge Stud. Adv. Math., vol. **42**, (1995).
- [8] J.-J. Duistermaat, *The heat kernel Lefschetz fixed point formula for the spin-c Dirac operator*, Reprint of the 1996 edition. Modern Birkhäuser Classics. Birkhuser/Springer, New York, 2011. xii+247 pp.
- [9] J.-J. Duistermaat and G.-J. Heckman, *On the Variation in the Cohomology of the Symplectic Form of the Reduced Phase Space*, Invent. Math., **69**(1982), 259-268.
- [10] C.-L. Epstein, *CR-structures on three dimensional circle bundles*, Invent. Math., **109**(1992), 351-403.
- [11] C.-L. Epstein, *A relative index on the space of embeddable CR-structures. I*, Ann. of Math. (2) **147**(1998), no.1, 1-59.
- [12] C.-L. Epstein, *A relative index on the space of embeddable CR-structures. II*, Ann. of Math. (2) **147**(1998), no.1, 61-91.

- [13] C.-L. Epstein, *Subelliptic boundary conditions for $Spin^c$ -Dirac operators, gluing, relative indices, and tame Fredholm pairs*, Proc. Natl. Acad. Sci. USA **103** (2006), no. 42, 15364–15369.
- [14] C.-L. Epstein, *Subelliptic $Spin^c$ Dirac operators. III. The Atiyah-Weinstein conjecture*, Ann. of Math. (2) **168**(2008), no.1, 299-365.
- [15] C.-L. Epstein, *Addendum to: Subelliptic $Spin^c$ Dirac operators, III*, Ann. of Math. **176** (2012), no. 2, 1373–1380.
- [16] E.-V. Erp, *The Atiyah-Singer index formula for subelliptic operators on contact manifolds. Part I*, Ann. of Math. **171**(2010), 1647–1681.
- [17] E.-V. Erp, *The Atiyah-Singer index formula for subelliptic operators on contact manifolds. Part II*, Ann. of Math. **171**(2010), 1683–1706.
- [18] H. Holmann, *Quotientenräume komplexer Mannigfaltigkeiten nach komplexen Lieschen Automorphismengruppen*, Math. Ann., **139**(1960), 383-402.
- [19] Alexander Grigoryan and Laurent Saloff-Coste, *Dirichlet heat kernel in the exterior of a compact set*, Comm. Pure Appl. Math., **55**(2002), 93–133.
- [20] C-Y. Hsiao, *Szegö kernel asymptotics for high power of CR line bundles and Kodaira embedding theorems on CR manifolds*, 116 pages, preprint available at arXiv:1401.6647.
- [21] C-Y. Hsiao and Xiaoshan Li, *Szegö kernel asymptotics and Morse inequalities on CR manifolds with S^1 action*, preprint available at arXiv:1502.02365.
- [22] C-Y. Hsiao and Xiaoshan Li, *Morse inequalities for positive Fourier coefficients on CR manifolds with S^1 action*, preprint available at arXiv:1506.06459.
- [23] Y. Kamishima and T. Tsuboi, *CR-structure on Seifert manifolds*, Invent. Math., **104**(1991), 149-164.
- [24] T. Kawasaki, *The Riemann-Roch theorem for complex V-manifolds*, Osaka J. Math., **16**(1)(1979), 151-159.
- [25] J.-M. Lee, *Pseudo-Einstein structures on CR manifolds*, Amer. J. Math., **110** (1988), 157–178.
- [26] L. Lempert, *On three dimensional Cauchy-Riemann manifolds*, J. Amer. Math. Soc. **5** (1992), 1–50.
- [27] X. Ma and G. Marinescu, *Holomorphic Morse inequalities and Bergman kernels*, Progress in Math., vol. 254, Birkhäuser, Basel, 2007, 422 pp.
- [28] G. Marinescu, *Asymptotic Morse Inequalities for Pseudoconcave Manifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. **23** (1996), no. 1, 27–55.
- [29] E. Meinrenken, *Group actions on manifolds*, Lecture Notes, University of Toronto, Spring 2003.
- [30] I. Satake, *On a generalization of the notion of manifold*, Proc. Nat. Acad. Sci. U.S.A. **42**(1956) 359-363.

INSTITUTE OF MATHEMATICS, ACADEMIA SINICA AND NATIONAL CENTER FOR THEORETICAL SCIENCES, 6F, ASTRONOMY-MATHEMATICS BUILDING, NO.1, SEC.4, ROOSEVELT ROAD, TAIPEI 10617, TAIWAN

E-mail address: cheng@math.sinica.edu.tw

INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, 6F, ASTRONOMY-MATHEMATICS BUILDING, NO.1, SEC.4, ROOSEVELT ROAD, TAIPEI 10617, TAIWAN

E-mail address: chsiao@math.sinica.edu.tw or chinyu.hsiao@gmail.com

DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN UNIVERSITY, TAIPEI, TAIWAN

E-mail address: ihtsai@math.ntu.edu.tw