

# A SHORT NOTE ON ADDITIVE FUNCTIONS ON RIEMANNIAN CO-COMPACT COVERINGS

MINH KHA

ABSTRACT. The main purpose of this note is to provide a topological approach to defining additive functions on Riemannian co-compact normal coverings.

## 1. INTRODUCTION

Let  $X$  be a connected smooth Riemannian manifold of dimension  $n$  equipped with an isometric, properly discontinuous, free, and co-compact action of a discrete deck group  $G$ . Notice that the deck group  $G$  is finitely generated due to the Švarc-Milnor lemma and hence,  $\text{Hom}(G, \mathbb{R})$  is finite dimensional. Furthermore, the orbit space  $M := X/G$  is a compact Riemannian manifold when equipped with the metric pushed down from  $X$ . The action of an element  $g \in G$  on  $x \in X$  is denoted by  $g \cdot x$ . Let  $\pi$  be the covering map from  $X$  onto  $M$ . Thus,  $\pi(g \cdot x) = \pi(x)$  for any  $(g, x) \in G \times X$ .

Following [6], we define the class of additive functions on the covering  $X$  as follows:

### Definition 1.1.

- A real smooth function  $u$  on  $X$  is said to be **additive** if there is a homomorphism  $\alpha : G \rightarrow \mathbb{R}$  such that

$$u(g \cdot x) = u(x) + \alpha(g), \quad \text{for all } (g, x) \in G \times X.$$

We denote by  $\mathcal{A}(X)$  the space of all additive functions on  $X$ .

- A map  $h$  from  $X$  to  $\mathbb{R}^m$  ( $m \in \mathbb{N}$ ) is called a vector-valued additive function on  $X$  if every component of  $h$  belongs to  $\mathcal{A}(X)$ .

We remark that additive functions on co-compact covers appeared in various results such as studying the structure of positive  $G$ -multiplicative type solutions [1, 6], describing the off-diagonal long time asymptotics of the heat kernel [3] and the Green's function asymptotics of periodic elliptic operators [2] on a noncompact abelian cover of a compact Riemannian manifold.

A direct construction of additive functions on  $X$  can be found in either [4, Section 3] or [6, Remark 2.6]. However, this construction depends on the choice of a fundamental domain for the base  $M$  in  $X$ . A more invariant approach to defining additive functions on covers was mentioned briefly in [1, 4]. Our goal in this note is to present the full details of this approach for any co-compact covering.

## 2. ADDITIVE FUNCTIONS ON CO-COMPACT NORMAL COVERINGS

We begin with the following notion (see [1, 4]):

**Definition 2.1.** Let  $H_{DR}^1(M), H_{DR}^1(X)$  be De Rham cohomologies of  $M$  and  $X$ , respectively. We denote by  $\Omega^1(M; G)$  the image in  $H_{DR}^1(M)$  of the set of all closed differential 1-forms  $\omega$  on  $M$  (modulo the exact ones) such that their lifts  $\omega$  to  $X$  are exact. In other words,  $\Omega^1(M; G)$  is the kernel of the homomorphism

$$\pi^* : H_{DR}^1(M) \rightarrow H_{DR}^1(X),$$

where  $\pi^*$  is the induced homomorphism of the covering map  $\pi : X \rightarrow M$ .

By De Rham's theorem,  $\Omega^1(M; G)$  is a finite dimensional vector space. Indeed, more is true:

**Lemma 2.2.**  $\Omega^1(M, G) \cong \text{Hom}(G, \mathbb{R})$ .

*Proof.* By Hurewicz's theorem (see e.g., [5]), the homologies  $H_1(M)$  and  $H_1(X)$  are isomorphic to the abelianizations of the fundamental groups  $\pi_1(M)$  and  $\pi_1(X)$ , respectively. Therefore, we can identify De Rham cohomologies  $H_{DR}^1(M)$  and  $H_{DR}^1(X)$  with  $\text{Hom}(\pi_1(M), \mathbb{R})$  and  $\text{Hom}(\pi_1(X), \mathbb{R})$ , correspondingly. Since  $X$  is a normal covering of  $M$ ,  $\pi_1(X)$  is a normal subgroup of  $\pi_1(M)$  and moreover, the sequence

$$0 \rightarrow \pi_1(X) \rightarrow \pi_1(M) \rightarrow G \rightarrow 0$$

is exact. Because  $\text{Hom}(\cdot, \mathbb{R})$  is a contravariant exact functor, we deduce the exactness of the following sequence of vector spaces:

$$0 \rightarrow \text{Hom}(G, \mathbb{R}) \rightarrow H_{DR}^1(M) \rightarrow H_{DR}^1(X) \rightarrow 0.$$

Hence,  $\Omega^1(M, G)$  is isomorphic to  $\text{Hom}(G, \mathbb{R})$ . □

Fixing a base point  $x_0 \in X$ . For any closed 1-form  $\omega$  on  $M$  such that its lift to  $X$  is exact, there exists a unique function  $f_\omega \in C^\infty(X, \mathbb{R})$  such that  $\pi^*\omega = df_\omega$  and  $f_\omega(x_0) = 0$ . Equivalently,

$$f_\omega(x) = \int_{x_0}^x \pi^*\omega, \quad \forall x \in X.$$

**Lemma 2.3.** *For such 1-form  $\omega$ , we have:*

- i) *Fix any  $g \in G$ , then  $f_\omega(g \cdot x) - f_\omega(x)$  is independent of  $x \in X$ .*
- ii) *If  $\pi^*\omega = 0$  then  $\omega = 0$ .*

*Proof.*

- i) For each  $g \in G$ , let  $L_g$  be the diffeomorphism of  $X$  that maps  $x$  to  $g \cdot x$ . Since  $\pi \circ L_g = \pi$ , we get  $df_\omega = \pi^*\omega = L_g^*\pi^*\omega = L_g^*df_\omega = dL_g^*f_\omega = d(f_\omega \circ L_g)$ . Thus,  $d(f_\omega - f_\omega \circ L_g) = 0$  and so,  $f_\omega \circ L_g - f_\omega$  is constant since  $X$  is connected.
- ii) Fix any point  $p \in M$ . We pick an evenly covered open subset  $U$  of  $M$  such that it contains  $p$ . Then there is a smooth local section  $\sigma : U \rightarrow X$ , i.e.,  $\pi \circ \sigma = \text{id}_U$  (see e.g., [5, Proposition 4.36]). Hence,  $\omega(p) = \sigma^*\pi^*\omega(p) = 0$ . □

On  $\mathcal{A}(X)$ , we introduce an equivalent relation  $\sim$  as follows:  $f_1 \sim f_2$  in  $\mathcal{A}(X)$  if and only if  $f_1 - f_2 = f \circ \pi$  for some function  $f \in C^\infty(M, \mathbb{R})$ .

By Lemma 2.3 (i), the map  $\omega \mapsto f_\omega$  induces the following linear map

$$(1) \quad \begin{aligned} \Lambda : \Omega^1(M, G) &\rightarrow \mathcal{A}(X)/\sim \\ [\omega] &\mapsto [f_\omega], \end{aligned}$$

where  $[\omega]$ ,  $[f_\omega]$  are the equivalent classes of  $\omega$ ,  $f_\omega$  in  $\Omega^1(M, G)$  and  $\mathcal{A}(X)/\sim$ , correspondingly. We now claim that  $[\omega] = 0$  if and only if  $[f_\omega] = 0$ , and hence  $\Lambda$  is an injective linear map. Indeed, due to Lemma 2.3 (ii), the condition that  $\omega$  is exact is equivalent to  $\pi^*\omega = d(f \circ \pi)$  for some  $f \in C^\infty(M, \mathbb{R})$ . But this is the same as  $df_\omega = d(f \circ \pi)$ , or  $[f_\omega] = 0$ .

Consider an additive function  $f$  on  $X$ . According to the definition, there exists a unique group homomorphism  $\ell_f : G \rightarrow \mathbb{R}$  such that  $f(g \cdot x) = f(x) + \ell_f(g)$  for any  $g \in G$ ,  $x \in X$ . Then the map  $f \mapsto \ell_f$  induces the linear map

$$(2) \quad \begin{aligned} \Upsilon : \mathcal{A}(X)/\sim &\rightarrow \text{Hom}(G, \mathbb{R}) \\ [f] &\mapsto \ell_f, \end{aligned}$$

which is injective.

Then the composition  $\Upsilon \circ \Lambda$  is also injective. By Lemma 2.2,  $\dim_{\mathbb{R}} \Omega^1(M, G) = \dim_{\mathbb{R}} \text{Hom}(G, \mathbb{R}) < \infty$ . These facts together imply that the linear maps  $\Upsilon$  and  $\Lambda$  are isomorphism. We conclude:

**Theorem 2.4.** *The three vector spaces  $\Omega^1(M, G)$ ,  $\mathcal{A}(X)/\sim$  and  $\text{Hom}(G, \mathbb{R})$  are isomorphic to each other.*

In particular, we obtain:

**Corollary 2.5.** *Assume that  $G = \mathbb{Z}^d$ . Then there is a smooth  $\mathbb{R}^d$ -valued function  $h$  on  $X$  such that for any  $(g, x) \in \mathbb{Z}^d \times X$ ,*

$$(3) \quad h(g \cdot x) = h(x) + g.$$

The following standard proposition says that given any additive function  $u$  on  $X$ , then one can pick a harmonic additive function  $f$  such that  $f - u$  is  $G$ -periodic.

**Proposition 2.6.** *For any  $\ell$  in  $\text{Hom}(G, \mathbb{R})$ , there exists a unique (modulo a real constant) **harmonic** function  $f$  on  $X$  such that for any  $(g, x) \in G \times X$ , we have*

$$(4) \quad f(g \cdot x) = f(x) + \ell(g).$$

*Proof.* First, we show the existence part. Due to Theorem 2.4, let  $\tilde{f}$  be a function on  $X$  satisfying  $\tilde{f}(g \cdot x) = \tilde{f}(x) + \ell(g)$  for any  $(g, x) \in G \times X$ . We recall the isomorphism  $\Lambda$  defined in (1). We put  $\alpha := \Lambda^{-1}([\tilde{f}]) \in \Omega^1(M, G)$ . By the Hodge theorem, there exists a unique harmonic 1-form  $\omega$  on  $M$  such that  $[\omega] = \alpha$  in  $H_{DR}^1(M)$ . Let  $f$  be a smooth function such that  $f \in \mathcal{A}(X)$  and  $\pi^*\omega = df$ . Then  $f$  satisfies (4) since  $[f] = [\tilde{f}]$  in  $\mathcal{A}(X)/\sim$ . Thus, it is sufficient to show that  $f$  is harmonic on  $X$ . We denote by  $\delta_X$ ,  $\Delta_X$  and  $\delta_M$ ,  $\Delta_M$  the codifferential and Laplace-Beltrami operators on  $X$  and  $M$ , respectively. Since the covering map  $\pi$  is a local isometry between  $X$  and  $M$ , its pullback  $\pi^*$  intertwines the codifferential operators, i.e.,

$$\delta_X \pi^* = \pi^* \delta_M.$$

Since  $\Delta_M \omega = 0$ , it follows that  $\delta_M \omega = 0$ . Thus,

$$\Delta_X f = \delta_X df = \delta_X \pi^* \omega = \pi^* \delta_M \omega = 0.$$

For the uniqueness part, let  $f_1$  and  $f_2$  be any two harmonic functions on  $X$  such that (4) holds for each of these functions. Since  $f_1 - f_2$  is  $G$ -periodic, it can be pushed down to a real function  $f$  on  $M$ . Moreover,  $\pi^* \Delta_M f = \Delta_X \pi^* f = \Delta_X (f_1 - f_2) = 0$ . Therefore,  $f$  must be constant since it is a harmonic function on a compact, connected Riemannian manifold  $M$ . Thus,  $f_1 - f_2$  is constant.  $\square$

**Corollary 2.7.** *Fixing a base point  $x_0$  in  $X$ . Then to each  $\alpha \in \text{Hom}(G, \mathbb{R})$ , there exists a unique harmonic function  $f_\alpha$  defined on  $X$  such that  $f_\alpha(x_0) = 0$  and  $\Upsilon([f_\alpha]) = \alpha$ , where  $\Upsilon$  is introduced in (2). Consequently,*

$$\mathcal{A}(X) = \bigsqcup_{\alpha \in \text{Hom}(G, \mathbb{R})} \{f_\alpha + \varphi \mid \varphi \text{ is periodic}\}.$$

**Remark 2.8.** When  $G = \mathbb{Z}^d$ , the Albanese pseudo-metric  $d_G$  introduced in [3, Section 2] is actually the pseudo-distance arising from **any** harmonic vector-valued additive function  $h$  satisfying (3), i.e.,  $d_G(x, y) = |h(x) - h(y)|$  for any  $x, y \in X$ .

### 3. ACKNOWLEDGEMENTS

The author was partially supported by the NSF grant DMS-1517938.

### REFERENCES

- [1] S. Agmon, *On positive solutions of elliptic equations with periodic coefficients in  $\mathbf{R}^n$ , spectral results and extensions to elliptic operators on Riemannian manifolds*, Differential equations (Birmingham, Ala., 1983), 1984, pp. 7–17.
- [2] M. Kha, *Green's function asymptotics of periodic elliptic operators on abelian coverings of compact manifolds*. arXiv:1511.00276, preprint.
- [3] M. Kotani and T. Sunada, *Albanese maps and off diagonal long time asymptotics for the heat kernel*, Comm. Math. Phys. **209** (2000), no. 3, 633–670.
- [4] P. Kuchment and Y. Pinchover, *Liouville theorems and spectral edge behavior on abelian coverings of compact manifolds*, Trans. Amer. Math. Soc. **359** (2007), no. 12, 5777–5815.
- [5] J. M. Lee, *Introduction to smooth manifolds*, Second Edition, Graduate Texts in Mathematics, vol. 218, Springer, New York, 2013.
- [6] V. Ya. Lin and Y. Pinchover, *Manifolds with group actions and elliptic operators*, Mem. Amer. Math. Soc. **112** (1994), no. 540, vi+78.

M.K., DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, USA

*E-mail address:* `kha@math.tamu.edu`