

Hypothesis Testing of Matrix Graph Model with Application to Brain Connectivity Analysis

Yin Xia

Department of Statistics and Operations Research,

University of North Carolina at Chapel Hill, Chapel Hill, NC 27514, U.S.A.

Email: xiayin@email.unc.edu

and

Lexin Li

Division of Biostatistics,

University of California at Berkeley, Berkeley, CA 94720, U.S.A.

Email: lexinli@berkeley.edu

Abstract

Brain connectivity analysis is now at the foreground of neuroscience research. A connectivity network is characterized by a graph, where nodes represent neural elements such as neurons and brain regions, and links represent statistical dependences that are often encoded in terms of partial correlations. Such a graph is inferred from matrix-valued neuroimaging data such as electroencephalography and functional magnetic resonance imaging. There have been a good number of successful proposals for sparse precision matrix estimation under normal or matrix normal distribution; however, this family of solutions do not offer a statistical significance

quantification for the estimated links. In this article, we adopt a matrix normal distribution framework and formulate the brain connectivity analysis as a precision matrix hypothesis testing problem. Based on the separable spatial-temporal dependence structure, we develop oracle and data-driven procedures to test the global hypothesis that all spatial locations are conditionally independent, which are shown to be particularly powerful against the sparse alternatives. In addition, simultaneous tests for identifying conditional dependent spatial locations with false discovery rate control are proposed in both oracle and data-driven settings. Theoretical results show that the data-driven procedures perform asymptotically as well as the oracle procedures and enjoy certain optimality properties. The empirical finite-sample performance of the proposed tests is studied via simulations, and the new tests are applied on a real electroencephalography data analysis.

Key Words: Connectivity analysis; False discovery rate; Gaussian graphical model; Matrix-variate normal distribution; Multiple testing.

1 Introduction

In recent years, matrix-valued data are becoming ubiquitous in a wide range of scientific and business applications, including bioinformatics (Yin and Li, 2012), brain imaging analysis (Reiss and Ogden, 2010; Aston and Kirch, 2012; Zhou and Li, 2014), finance (Leng and Tang, 2012), among many others. Accordingly, the matrix normal distribution is becoming increasingly popular in modeling the matrix-variate observations (Zhou, 2014). Our motivating example is an electroencephalography (EEG) data, which measures voltage value from electrodes placed at various brain locations over a period of time for a group of alcoholic subjects and normal controls. One scientific goal is to infer the connectivity patterns among those spatial locations. More generally, accurate and informative mapping of the human connectivity network is now at the center stage of neuroscience research. The objective is to infer brain connectivity network, which is commonly characterized as a graph consisting of nodes and links. Here nodes represent

neural elements, from micriscopic neurons to macroscopic brain regions, and links represent statistical dependencies between neural components (Johansen-Berg, 2013). Partial correlations, reported by a precision matrix, are frequently employed to describe such statistical dependencies (Fornito et al., 2013). This precision matrix, in turn, is to be derived from imaging modalities, such as EEG, magnetoencephalography, and functional magnetic resonance imaging. The data of those imaging modalities are in the common form of a two-dimensional matrix, with one spatial dimension and the other temporal dimension.

Adopting a matrix normal distribution framework, we formulate the brain connectivity network analysis as a precision matrix inference problem. Specifically, let $X \in \mathbb{R}^{p \times q}$ denote the spatial-temporal matrix data from an image modality, e.g., EEG. It is assumed to follow a matrix normal distribution with the Kronecker product covariance structure,

$$\text{cov}\{\text{vec}(X)\} = \Sigma_L \otimes \Sigma_T,$$

where the operator $\text{vec}(X)$ stacks the columns of the matrix X to a vector, \otimes is the Kronecker product, $\Sigma_L \in \mathbb{R}^{p \times p}$ denotes the covariance matrix of p spatial locations, $\Sigma_T \in \mathbb{R}^{q \times q}$ denotes the covariance matrix for q time points. Correspondingly,

$$\text{cov}^{-1}\{\text{vec}(X)\} = \Sigma_L^{-1} \otimes \Sigma_T^{-1} = \Omega_L \otimes \Omega_T,$$

where $\Omega_L \in \mathbb{R}^{p \times p}$ is the spatial precision matrix, $\Omega_T \in \mathbb{R}^{q \times q}$ is the temporal precision matrix. In brain connectivity analysis, our primary interest is to infer the the connectivity network characterized by the spatial precision matrix Ω_L . By contrast, the temporal precision matrix Ω_T is of little interest here and is to be treated as a nuisance parameter in our analysis. We also make some remarks regarding the assumptions of our adopted framework. First, the matrix normal assumption has been frequently adopted in various applications (Yin and Li, 2012; Leng and Tang, 2012), and is scientifically plausible in neuroimaging analysis. For instance, the majority standard neuroimaging processing software, such as SPM (Friston et al., 2007) and FSL (Smith et al., 2004), adopt a framework that assumes the data are normally distributed per voxel (location) with a noise factor and an autoregressive structure, which shares a similar spirit as the matrix normal formulation. Second, it is commonly assumed that the precision matrix is sparse, which

we adopt for our inferential procedure as well. Again, this sparsity assumption is scientifically justifiable, as it is known that brain region connections are energy consuming (Olshausen and Field, 2004), and biological units tend to minimize energy-consuming activities (Bullmore and Sporns, 2009).

In this article, we aim to address the following two hypothesis testing questions. The first is to test if all spatial locations are conditionally independent, namely, we test the null hypothesis

$$H_0 : \Omega_L \text{ is diagonal} \quad \text{versus} \quad H_1 : \Omega_L \text{ is not diagonal.} \quad (1)$$

The second is to identify those conditionally dependent pairs of locations with false discovery rate and false discovery proportion control; i.e., we simultaneously test

$$H_{0,i,j} : \omega_{L,i,j} = 0 \quad \text{versus} \quad H_{1,i,j} : \omega_{L,i,j} \neq 0, \quad \text{for } 1 \leq i < j \leq p, \quad (2)$$

where $\omega_{L,i,j}$ is the (i, j) th element of Ω_L .

In the literature, there have been a good number of methods proposed to estimate a sparse precision matrix under normal distribution (Meinshausen and Bühlmann, 2006; Yuan and Lin, 2007; Friedman et al., 2008; Yuan, 2010; Ravikumar et al., 2011; Cai et al., 2011). There are extensions of this line of work from a single precision matrix to multiple precision matrices (Danaher et al., 2014; Zhu et al., 2014), and from a Gaussian distribution to a more flexible class of nonparanormal distribution (Liu et al., 2012). Extension of sparse precision matrix estimation has also emerged for matrix-valued data under matrix normal distribution (Yin and Li, 2012; Leng and Tang, 2012; Zhou, 2014). However, all those methods tackle the *estimation* aspect of the problem and induce a connectivity network from an estimated precision matrix. Only recently, *hypothesis testing* procedures have been developed under Gaussian graphical model. In particular, Liu (2013) proposed a testing procedure to recover a network in the one-sample case, whereas Xia et al. (2015) proposed a method to test the equality of two precision matrices, so to infer the differential network structures in genomics. Both papers, however, worked with vector normal data instead of matrix normal data.

We aim at hypothesis testing for the spatial precision matrix, under the matrix normal framework, to induce the connectivity network of brain regions. We separate the spatial

and temporal dependence structures, then infer the precision matrix Ω_L through inverse regression models by relating its elements to the regression coefficients. Two procedures are considered. One is to assume the temporal covariance matrix is known, and we term the resulting method as an oracle procedure. The other is to use a data-driven approach to estimate and plug in the temporal covariance matrix, and accordingly we term it a data-driven procedure. We construct test statistics based on the covariances between the residuals from the inverse regression models. We first develop a global test for (1), and show it is particularly powerful against a sparse alternative, then develop a multiple testing procedure for simultaneously testing the hypotheses (2) with false discovery rate and false discovery proportion control. We study the theoretical properties of the two testing procedures, in both the oracle and data-driven scenarios. We show that the data-driven procedure performs asymptotically as well as the oracle procedure, and enjoys certain optimality under the regularity conditions. Our numerical analysis also supports such findings.

Our contributions are multi-fold. First, brain connectivity analysis is now becoming a central goal of neuroscience research (Fornito et al., 2013), and it constantly calls for statistical significance quantification of the inferred connection between neural elements. However, there is a paucity of systematic hypothesis testing solutions developed for this type of problems in the literature, and our proposal offers a timely response. Second, although various network sparse estimation methods have been successfully applied in neural connectivity analyses, network hypothesis testing is an utterly different problem than estimation. The key of estimation is to seek a bias-variance tradeoff, and many common sparse estimation solutions such as graphical lasso (Friedman et al., 2008) and constrained ℓ_1 -minimization for inverse matrix estimation (Cai et al., 2011, clime) are biased estimators. Such estimation methods do not produce a direct quantification of statistical significance for the network edges. By contrast, hypothesis testing starts with an nearly unbiased estimator, and produces an explicit significance quantification. Third, among the few network hypothesis testing solutions, Liu (2013) and Xia et al. (2015) focused on a vector-valued X following a normal distribution rather than a matrix-valued

data. Directly applying their methods to test the spatial conditional dependence, with no regard for the separable structure of Ω_L and Ω_T , is equivalent to assuming the columns of the matrix X are independent. This is clearly not true as the measurements at different time points can be highly correlated. Thus it is important to separate the spatial-temporal dependence structure before testing.

The following notations are used throughout this article. For any matrix normal data $\{X_k\}_{k=1}^n$, with $X_k \in \mathbb{R}^{p \times q}$, $X_{k,i,l}$ denotes the i -th spatial location at the l -th time point for the k -th sample, and $\bar{X}_{i,l} = \frac{1}{n} \sum_{k=1}^n X_{k,i,l}$; $X_{k,-i,l}$ denotes the column spatial vector with the i -th entry removed, and $\bar{X}_{\cdot,-i,l} = \frac{1}{n} \sum_{k=1}^n X_{k,-i,l} \in \mathbb{R}^{1 \times (p-1)}$. For a $p \times 1$ vector a , a_{-i} denotes the $(p-1) \times 1$ vector by removing the i th entry a_i from a . For an $nq \times p$ data matrix $A = (A_1, \dots, A_{nq})^T$, $A_{\cdot,-i}$ denotes an $nq \times (p-1)$ matrix $(A_{1,-i}^T, \dots, A_{nq,-i}^T)^T$, $\bar{A}_{\cdot,-i} = \frac{1}{nq} \sum_{k=1}^n A_{k,-i} \in \mathbb{R}^{1 \times (p-1)}$, $A_{(i)} = (A_{1,i}, \dots, A_{nq,i})^T \in \mathbb{R}^{nq \times 1}$, $\bar{A}_{(i)} = (\bar{A}_i, \dots, \bar{A}_i)^T \in \mathbb{R}^{nq \times 1}$, where $\bar{A}_i = \frac{1}{nq} \sum_{k=1}^n A_{k,i}$, and $\bar{A}_{(\cdot,-i)} = (\bar{A}_{\cdot,-i}^T, \dots, \bar{A}_{\cdot,-i}^T)^T \in \mathbb{R}^{nq \times (p-1)}$. Furthermore, for any $A \in \mathbb{R}^{p \times q}$, $A_{i,-j}$ denotes the i th row of A with its j th entry removed, and $A_{-i,j}$ denotes the j th column of A with its i th entry removed. $A_{-i,-j}$ denotes the submatrix of A with its i th row and j th column removed. If A is symmetric, then $\lambda_{\max}(A)$ and $\lambda_{\min}(A)$ denote the largest and smallest eigenvalues of A . We also use the following norms. For a vector $a \in \mathbb{R}^{p \times 1}$, define its ℓ_s norm as $|a|_s = (\sum_{i=1}^p |a_i|^s)^{1/s}$ for $1 \leq s \leq \infty$. For a matrix $A \in \mathbb{R}^{p \times p}$, the matrix 1-norm is the maximum absolute column sum, $\|A\|_{L_1} = \max_{1 \leq j \leq p} \sum_{i=1}^p |a_{i,j}|$. The matrix element-wise infinity norm is $\|A\|_{\infty} = \max_{1 \leq i,j \leq p} |a_{i,j}|$ and the element-wise ℓ_1 norm is $\|A\|_1 = \sum_{i=1}^p \sum_{j=1}^p |a_{i,j}|$. Finally, for two sequences of real numbers $\{a_n\}$ and $\{b_n\}$, $a_n = O(b_n)$ if there exists a constant c such that $|a_n| \leq c|b_n|$ holds for all n , $a_n = o(b_n)$ if $\lim_{n \rightarrow \infty} a_n/b_n = 0$, and $a_n \asymp b_n$ if there are positive constants c_0 and c_1 such that $c_0 \leq a_n/b_n \leq c_1$ for all n .

2 Methodology

In this section, we derive test statistics and testing procedures for the global hypothesis (1), and the entry-wise hypothesis (2) with false discovery rate control.

2.1 Separation of spatial-temporal dependency

Let $\{X_1, \dots, X_n\}$, each with dimension $p \times q$, denote n i.i.d. random samples from a matrix normal distribution. The mean, without loss of generality, is assumed to be zero, and the covariance is of the form $\Sigma = \Sigma_L \otimes \Sigma_T$. Our interest is to infer about $\Omega_L = \Sigma_L^{-1}$, while treating $\Omega_T = \Sigma_T^{-1}$ as a nuisance. To separate the spatial and temporal dependence structures, we build hypothesis testing procedures for (1) and (2) based upon the linear transformation of the original samples $\{X_k \Sigma_T^{-1/2}, k = 1, \dots, n\}$. Specifically, we consider two scenarios. We first treat Σ_T as known, and term the resulting testing procedure an oracle procedure. In practice, however, Σ_T is rarely known, and as such we plug in an estimator of Σ_T . When $\text{trace}(\Sigma_L) = p$ holds true, the sample covariance matrix $\hat{\Sigma}_T = \frac{1}{np} \sum_{k=1}^n X_k^T X_k$ is an unbiased estimator of Σ_T . However, when $\text{trace}(\Sigma_L) \neq p$, $\hat{\Sigma}_T$ is biased and we have $E(\hat{\Sigma}_T) = \{\text{trace}(\Sigma_L)/p\}\Sigma_T$. As we will show in Remark 2 in the next section, $\text{trace}(\Sigma_L)$ does not affect our proposed test statistics, and thus we can assume without loss of generality that $\text{trace}(\Sigma_L) = p$ to simplify the notations. Subsequently, we plug in the sample covariance matrix $\hat{\Sigma}_T$, develop the hypothesis testing based on the transformed samples, $\{X_k \hat{\Sigma}_T^{-1/2}, k = 1, \dots, n\}$, and term it a data-driven procedure. One may also use other estimators of Σ_T or Ω_T , and we will briefly discuss those alternatives in Section 6.

2.2 Test statistics

We first develop test statistics for the two hypotheses in the oracle case. The development of the data-driven statistics is very similar, so we omit the details and will remark clearly the difference between the oracle and data-driven cases. For simplicity, we also use the same set of notations for the two scenarios, and will only differentiate them when we

study their respective asymptotic properties in Section 3.

It is well established that, under the normal distribution, the precision matrix can be described in terms of regression models (Anderson, 2003, Sec 2.5). Specifically, letting $Y_k = X_k \Sigma_T^{-1/2}$, $k = 1, \dots, n$, denote the transformed samples, we have,

$$Y_{k,i,l} = Y_{k,-i,l}^\top \beta_i + \epsilon_{k,i,l}, \quad 1 \leq i \leq p, 1 \leq l \leq q, \quad (3)$$

where $\epsilon_{k,i,l} \sim N(0, \sigma_{L,i,i} - \Sigma_{L,i,-i} \Sigma_{L,-i,-i}^{-1} \Sigma_{L,-i,i})$ is independent of $Y_{k,-i,l}$. The regression coefficient vector β_i and the error term $\epsilon_{k,i}$ satisfy that

$$\beta_i = -\omega_{L,i,i}^{-1} \Omega_{L,-i,i}, \quad \text{and} \quad r_{i,j} = \text{cov}(\epsilon_{k,i,l}, \epsilon_{k,j,l}) = \frac{\omega_{L,i,j}}{\omega_{L,i,i} \omega_{L,j,j}}.$$

As such, the elements $\omega_{L,i,j}$ of Ω_L can be represented in terms of $r_{i,j}$. Next, we construct an estimator of $r_{i,j}$ and its bias-corrected version. We then build on this estimator to obtain an estimator of $\omega_{L,i,j}$, upon which we further develop our test statistics.

A natural estimator of $r_{i,j}$ is the sample covariance between the residuals $\hat{\epsilon}_{k,i,l} = Y_{k,i,l} - \bar{Y}_{i,l} - (Y_{k,-i,l} - \bar{Y}_{-i,l})^\top \hat{\beta}_i$,

$$\tilde{r}_{i,j} = \frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q \hat{\epsilon}_{k,i,l} \hat{\epsilon}_{k,j,l},$$

where $\hat{\beta}_i$, $i = 1, \dots, p$, are estimators of β_i that satisfy Condition (C1) in the oracle case and satisfies Condition (C1') in the data-driven case, and these estimators can be obtained via standard estimation methods such as the Lasso and Dantzig selector, as will be discussed in Section 3.1. When $i \neq j$, however, $\tilde{r}_{i,j}$ tends to be biased due to the correlation induced by the estimated parameters. We thus consider a bias-corrected estimator of $r_{i,j}$,

$$\hat{r}_{i,j} = -(\tilde{r}_{i,j} + \tilde{r}_{i,i} \hat{\beta}_{i,j} + \tilde{r}_{j,j} \hat{\beta}_{j-1,i}), \quad \text{for } 1 \leq i < j \leq p.$$

For $i = j$, we let $\hat{r}_{i,i} = \tilde{r}_{i,i}$, which is a nearly unbiased estimator of $r_{i,i}$. An estimator of the entry $\omega_{L,i,j}$ of the spatial precision matrix Ω_L can then be constructed as,

$$T_{i,j} = \frac{\hat{r}_{i,j}}{\hat{r}_{i,i} \cdot \hat{r}_{j,j}}, \quad 1 \leq i < j \leq p.$$

To further estimate the variance of $T_{i,j}$, note that

$$\theta_{i,j} = \text{var}\{\epsilon_{k,i} \epsilon_{k,j} / (r_{i,i} r_{j,j})\} / (nq) = (1 + \rho_{i,j}^2) / (nq r_{i,i} r_{j,j}), \quad (4)$$

where $\rho_{i,j}^2 = \beta_{i,j}^2 r_{i,i}/r_{j,j}$. Then $\theta_{i,j}$ can be estimated by

$$\hat{\theta}_{i,j} = (1 + \hat{\beta}_{i,j}^2 \hat{r}_{i,i}/\hat{r}_{j,j})/(nq\hat{r}_{i,i}\hat{r}_{j,j}).$$

Given $\{T_{i,j}, 1 \leq i < j \leq p\}$ are heteroscedastic and can possibly have a wide variability, we standardize $T_{i,j}$ by its standard error, which leads to the standardized statistics,

$$W_{i,j} = \frac{T_{i,j}}{\sqrt{\hat{\theta}_{i,j}}}, \quad 1 \leq i < j \leq p.$$

In the next section, we test hypotheses (1) and (2) based on $\{W_{i,j}\}_{i,j=1}^p$.

Remark 1 *Construction of the test statistics for the data-driven procedure is almost the same as that for the oracle procedure, except that the oracle procedure starts with transformed sample $Y_k = X_k \Sigma_T^{-1/2}$ in (3), whereas the data-driven one replaces it with $Y_k = X_k \hat{\Sigma}_T^{-1/2}$. Furthermore, the regression coefficients slightly vary at different time points in the data-driven scenario, and we shall replace (3) by $Y_{k,i,l} = Y_{k,-i,l}^\top \beta_{i,l} + \epsilon_{k,i,l}$, for $1 \leq i \leq p, 1 \leq l \leq q$.*

Remark 2 *When Σ_T is unknown, $E(\hat{\Sigma}_T) = \{\text{trace}(\Sigma_L)/p\}\Sigma_T$. If $\text{trace}(\Sigma_L) = cp$, with $c \neq 1$, an unbiased estimator of Σ_T becomes $\hat{\Sigma}_T/c$. Accordingly, we shall define the transformed data $Y_k = \sqrt{c}X_k \hat{\Sigma}_T^{-1/2}$, for $k = 1, \dots, n$. Then we have the bias-corrected estimator $\hat{r}_{i,j} = c\hat{r}_{i,j}$, which in turn leads to $T_{i,j} = T_{i,j}/c$, and $\hat{\theta}_{i,j} = \hat{\theta}_{i,j}/c^2$. Thus, the standardized statistic $W_{i,j}$ remains the same, as the constant c is cancelled. Therefore, c does not affect our final test statistics, and thus for notational simplicity, we set $c = 1$ from the beginning, without loss of any generality.*

2.3 Global testing procedure

We propose the following test statistic for testing the global null hypothesis $H_0 : \Omega_L$ is diagonal,

$$M_{nq} = \max_{1 \leq i < j \leq p} W_{i,j}^2.$$

Furthermore, we define the global test Ψ_α by

$$\Psi_\alpha = I(M_{nq} \geq q_\alpha + 4 \log p - \log \log p)$$

where q_α is the $1 - \alpha$ quantile of the type I extreme value distribution with the cumulative distribution function $\exp\{(8\pi)^{-1/2}e^{-t/2}\}$, i.e.,

$$q_\alpha = -\log(8\pi) - 2\log\log(1 - \alpha)^{-1}.$$

The hypothesis H_0 is rejected whenever $\Psi_\alpha = 1$.

The above test is developed based on the asymptotic properties of M_{nq} , which will be studied in detail in Section 3.2. Intuitively, $\{W_{i,j}\}_{i,j=1}^p$ are approximately standard normal variables under the null distribution, and are only weakly dependent under suitable conditions. Thus M_{nq} is the maximum of the squares of $p(p-1)/2$ such random variables, and its value should be close to $2\log\{p(p-1)/2\} \approx 4\log p$ under H_0 . We will later show that, under certain regularity conditions, $M_{nq} - 4\log p - \log\log p$ converges to a type I extreme value distribution under H_0 .

2.4 Multiple testing procedure

Next we develop a multiple testing procedure for $H_{0,i,j} : \omega_{L,i,j} = 0$, so to identify spatial locations that are conditionally dependent. The test statistic $W_{i,j}$ defined in Section 2.2 is employed. Since there are $(p^2 - p)/2$ simultaneous hypotheses to test, it is important to control the false discovery rate. Let t be the threshold level such that $H_{0,i,j}$ is rejected if $|W_{i,j}| \geq t$. Let $\mathcal{H}_0 = \{(i, j) : \Omega_{L,i,j} = 0, 1 \leq i < j \leq p\}$ be the set of true nulls. Denote by $R_0(t) = \sum_{(i,j) \in \mathcal{H}_0} I(|W_{i,j}| \geq t)$ the total number of false positives, and by $R(t) = \sum_{1 \leq i < j \leq p} I(|W_{i,j}| \geq t)$ the total number of rejections. The false discovery proportion and false discovery rate are then defined as

$$\text{FDP}(t) = \frac{R_0(t)}{R(t) \vee 1}, \quad \text{FDR}(t) = E\{\text{FDP}(t)\}.$$

An ideal choice of t would reject as many true positives as possible while controlling the false discovery rate and false discovery proportion at the pre-specified level α . That is, we select

$$t_0 = \inf \{0 \leq t \leq 2(\log p)^{1/2} : \text{FDP}(t) \leq \alpha\}.$$

We shall estimate $\sum_{(i,j) \in \mathcal{H}_0} I\{|W_{i,j}| \geq t\}$ by $2\{1 - \Phi(t)\}|\mathcal{H}_0|$, where $\Phi(t)$ is the standard normal cumulative distribution function. Note that $|\mathcal{H}_0|$ can be estimated by $(p^2 - p)/2$ due to the sparsity of Ω_L . This leads to the following multiple testing procedure.

Step 1. Calculate the test statistics $W_{i,j}$.

Step 2. For given $0 \leq \alpha \leq 1$, calculate

$$\hat{t} = \inf \left\{ 0 \leq t \leq 2(\log p)^{1/2} : \frac{2\{1 - \Phi(t)\}(p^2 - p)/2}{R(t) \vee 1} \leq \alpha \right\}.$$

If \hat{t} does not exist, set $\hat{t} = 2(\log p)^{1/2}$.

Step 3. For $1 \leq i < j \leq p$, reject $H_{0,i,j}$ if and only if $|W_{i,j}| \geq \hat{t}$.

3 Theory

In this section, we analyze the theoretical properties of the global and multiple testing procedures for both the oracle and data-driven scenarios. We show that the data-driven procedures perform asymptotically as well as the oracle procedures and enjoy certain optimality under the regularity conditions. For separate treatment of the oracle and data-driven procedures, we now distinguish the notations of the two, and add the superscript “ o ” to denote the statistics and tests of the oracle procedures, e.g., $\hat{\beta}_i^o, M_{nq}^o, \Psi_\alpha^o, \hat{t}^o$, and the superscript “ d ” to denote those of the data-driven procedures, e.g., $\hat{\beta}_i^d, M_{nq}^d, \Psi_\alpha^d$, and \hat{t}^d .

3.1 Regularity conditions

For the oracle procedure, we require the following set of regularity conditions.

(C1) Assume that $\max_{1 \leq i \leq p} |\hat{\beta}_i^o - \beta_i|_1 = o_p[\{\log \max(p, q, n)\}^{-1}]$, and $\max_{1 \leq i \leq p} |\hat{\beta}_i^o - \beta_i|_2 = o_p\{(nq \log p)^{-1/4}\}$.

(C2) Assume that $\log p = o\{(nq)^{1/5}\}$, and there are constants $c_0, c_1 > 0$ such that, $c_0^{-1} \leq \lambda_{\min}(\Omega_L) \leq \lambda_{\max}(\Omega_L) \leq c_0$, and $c_1^{-1} \leq \lambda_{\min}(\Omega_T) \leq \lambda_{\max}(\Omega_T) \leq c_1$.

(C3) Let D_L be the diagonal of Ω_L and let $R_L = D_L^{-1/2} \Omega_L D_L^{-1/2}$ with elements $\eta_{L,i,j}$, $1 \leq i, j \leq p$. Assume that $\max_{1 \leq i < j \leq p} |\eta_{L,i,j}| \leq \eta_L < 1$, for some constant $0 < \eta_L < 1$.

For the data-driven procedure, we replace the above condition (C1) with a slightly different one (C1'), then introduce a new condition (C4).

(C1') Assume that $\max_{1 \leq i \leq p, 1 \leq l \leq q} |\hat{\beta}_i^d - \beta_{i,l}|_1 = o_p[\{\log \max(p, q, n)\}^{-1}]$,
and $\max_{1 \leq i \leq p, 1 \leq l \leq q} |\hat{\beta}_i^d - \beta_{i,l}|_2 = o_p\{(nq \log p)^{-1/4}\}$.

(C4) Define $s_p = \max_{1 \leq l \leq q} \max_{1 \leq i \leq p} \sum_{j=1}^p \max\{I(\omega_{L,i,j} \neq 0), I(\omega_{l,i,j}^d \neq 0)\}$, where $(\omega_{l,i,j}^d)_{p \times p} = \Omega_l^d = \text{cov}^{-1}\{(X_k \hat{\Sigma}_T^{-1/2})_{\cdot, l}\}$. Assume that $\|\Omega_T\|_{L_1}^2 \|\Omega_L\|_{L_1}^2 = o\{\min(r_{1,n,p,q}, r_{2,n,p,q})\}$, where $r_{1,n,p,q} = [np/\{s_p^2 q^3 \log q \log^3 \max(p, q, n)\}]^{1/2}$, and $r_{2,n,p,q} = (np^2/[s_p^2 q^7 \{\log q \log \max(p, q, n)\}^2 \log p])^{1/4}$.

A few remarks are in order. The estimator $\hat{\beta}_i^o$ satisfying (C1) can be easily obtained via standard estimation methods such as Lasso and Dantzig selector. For instance, if one uses the Lasso estimator, then (C1) is satisfied under (C2) and the sparsity condition $\max_{1 \leq i \leq p} |\beta_i|_0 = o[(nq)^{1/2}/\{\log \max(p, q, n)\}^{3/2}]$. Similarly, $\hat{\beta}_i^d$ satisfying (C1') can be obtained by Lasso if (C4) holds and the data-driven regression coefficients $\beta_{i,l}$ satisfy the similar sparsity condition. Conditions (C2) and (C3) are regularity conditions that are commonly used in the high-dimensional hypothesis testing setting (Cai et al., 2013; Liu, 2013; Xia et al., 2015). (C4) is a mild technical condition. If Ω_T , Ω_L and Ω_l^d satisfy $\max_{1 \leq i \leq q} \sum_{j=1}^q I(\omega_{T,i,j} \neq 0) \leq s$ and $s_p \leq s$, for some constant $s > 0$, then the conditions on matrix 1-norms can be relaxed to the conditions only related to n, p and q , namely, $q^3 \log q \log^3 \max(p, q, n) = o(np)$ and $q^7 \{\log q \log \max(p, q, n)\}^2 \log p = o(np^2)$.

3.2 Oracle global testing procedure

We next analyze the limiting null distribution of the oracle global test statistic M_{nq}^o and the power of the corresponding test Ψ_α^o . We are particularly interested in the power of the test under the alternative when Ω_L is sparse, and show that the power is minimax rate optimal.

The following theorem states the asymptotic null distribution for M_{nq}^o , and indicates that, under H_0 , $M_{nq}^o - 4 \log p + \log \log p$ converges weakly to a Gumbel random variable with distribution function $\exp\{-(8\pi)^{-1/2} e^{-t/2}\}$.

Theorem 1 *Assume (C1), (C2) and (C3). Then under H_0 , for any $t \in \mathbb{R}$,*

$$\text{pr}(M_{nq}^o - 4 \log p + \log \log p \leq t) \rightarrow \exp\{-(8\pi)^{-1/2} \exp(-t/2)\}, \text{ as } nq, p \rightarrow \infty.$$

Under H_0 , the above convergence is uniform for all $\{X_k\}_{k=1}^n$ satisfying (C1)-(C3).

We next study the power of the corresponding test Ψ_α^o . We define the following class of precision matrices for spatial locations:

$$\mathcal{U}(c) = \left\{ \Omega_L : \max_{1 \leq i < j \leq p} \frac{|\omega_{L,i,j}|}{\theta_{i,j}^{1/2}} \geq c(\log p)^{1/2} \right\}. \quad (5)$$

This class of matrices include all precision matrices such that there exists one standardized off-diagonal entry having the magnitude exceeding $c(\log p)^{1/2}$. By the definition in (4), $\theta_{i,j}$ is of the order $1/(nq)$, and thus we only require one of the off-diagonal entries to have size larger than $C\{\log p/(nq)\}^{1/2}$ for some constant $C > 0$, where C is fully determined by c_0 and c_1 in Condition (C2). Then if we choose the constant $c = 4$, that is, if there exists one standardized off-diagonal entry having the magnitude larger or equal than $4(\log p)^{1/2}$, the next theorem shows that the null parameter set in which Ω_L is diagonal is asymptotically distinguishable from $\mathcal{U}(4)$ by the test Ψ_α^o . That is, H_0 is rejected by the test Ψ_α^o with overwhelming probability if $\Omega_L \in \mathcal{U}(4)$.

Theorem 2 *Assume (C1) and (C2). Then,*

$$\inf_{\Omega_L \in \mathcal{U}(4)} \text{pr}(\Psi_\alpha^o = 1) \rightarrow 1, \text{ as } nq, p \rightarrow \infty.$$

The next theorem further shows that this lower bound $4(\log p)^{1/2}$ is rate-optimal. Let \mathcal{T}_α be the set of all α -level tests, i.e., $\text{pr}(T_\alpha = 1) \leq \alpha$ under H_0 for all $T_\alpha \in \mathcal{T}_\alpha$.

Theorem 3 *Suppose that $\log p = o(nq)$. Let $\alpha, \beta > 0$ and $\alpha + \beta < 1$. Then there exists a constant $c_2 > 0$ such that for all sufficiently large nq and p ,*

$$\inf_{\Omega_L \in \mathcal{U}(c_2)} \sup_{T_\alpha \in \mathcal{T}_\alpha} \text{pr}(T_\alpha = 1) \leq 1 - \beta.$$

As Theorem 3 indicates, if c_2 is sufficiently small, then any α level test is unable to reject the null hypothesis correctly uniformly over $\Omega_L \in \mathcal{U}(c_2)$ with probability tending to one. So the order $(\log p)^{1/2}$ in the lower bound of $\max_{1 \leq i < j \leq p} \{|\omega_{L,i,j}| \theta_{i,j}^{-1/2}\}$ in (5) cannot be further improved.

3.3 Oracle multiple testing procedure

We next investigate the properties of the oracle multiple testing procedure. The following theorem shows that the oracle procedure controls the false discovery proportion and false discovery rate at the pre-specified level α asymptotically.

Theorem 4 *Assume (C1) and (C2), and let*

$$\mathcal{S}_\rho = \left\{ (i, j) : 1 \leq i < j \leq p, \frac{|\omega_{L,i,j}|}{\theta_{i,j}^{1/2}} \geq (\log p)^{1/2+\rho} \right\}.$$

Suppose for some $\rho, \delta > 0$, $|\mathcal{S}_\rho| \geq [1/\{(8\pi)^{1/2}\alpha\} + \delta](\log \log p)^{1/2}$. Suppose $l_0 = |\mathcal{H}_0| \geq c_0 p^2$ for some $c_0 > 0$, and $p \leq c(nq)^r$ for some $c, r > 0$. Letting $l = (p^2 - p)/2$, then,

$$\lim_{(nq,p) \rightarrow \infty} \frac{\text{FDR}(\hat{t}^o)}{\alpha l_0/l} = 1, \quad \frac{\text{FDP}(\hat{t}^o)}{\alpha l_0/l} \rightarrow 1$$

in probability, as $(nq, p) \rightarrow \infty$.

We comment that the condition $|\mathcal{S}_\rho| \geq [1/\{(8\pi)^{1/2}\alpha\} + \delta](\log \log p)^{1/2}$ in Theorem 4 is mild, because we have $(p^2 - p)/2$ hypotheses in total and this condition only requires a few entries of Ω_L having standardized magnitude exceeding $\{(\log p)^{1/2+\rho}/(nq)\}^{1/2}$ for some constant $\rho > 0$.

3.4 Data-driven procedures

We next turn to data-driven procedures for both the global testing and the multiple testing. We show that they perform as well as the oracle testing procedures asymptotically.

Theorem 5 *Assume (C1'), (C2)-(C4).*

(i) Under H_0 , for any $t \in \mathbb{R}$,

$$\text{pr}(M_{nq}^d - 4 \log p + \log \log p \leq t) \rightarrow \exp\{-(8\pi)^{-1/2} \exp(-t/2)\}, \text{ as } nq, p \rightarrow \infty.$$

Under H_0 , the above convergence is uniform for all $\{X_k\}_{k=1}^n$ satisfying (C1'), (C2)-(C4).

(ii) Furthermore, $\inf_{\Omega_L \in \mathcal{U}(4)} \text{pr}(\Psi_\alpha^d = 1) \rightarrow 1$, as $nq, p \rightarrow \infty$.

This theorem shows that M_{nq}^d has the same limiting null distribution as the oracle test statistics M_{nq}^o , and the power of the corresponding test Ψ_α^d performs as well as the oracle test and is thus minimax rate optimal. The same observation applies to Theorem 6 below, which shows that the data-driven multiple procedure also performs as well as the oracle case, and controls the false discovery proportion and false discovery rate at the pre-specified level α asymptotically.

Theorem 6 Assume (C1') and (C4). Then under the same conditions as in Theorem 4,

$$\lim_{(nq,p) \rightarrow \infty} \frac{\text{FDR}(\hat{t}^d)}{\alpha l_0/l} = 1, \quad \frac{\text{FDP}(\hat{t}^d)}{\alpha l_0/l} \rightarrow 1$$

in probability, as $(nq, p) \rightarrow \infty$.

4 Simulations

We study in this section the finite-sample performance of the proposed testing procedures. For the global testing of (1), we measure the size and power of the oracle test Ψ_α^o and the data-driven version Ψ_α^d , and for the multiple testing of (2), we measure the empirical FDR and power. We compare the oracle and data-driven testing procedures, as well as a simple alternative that was developed by Xia et al. (2015) under normal instead of matrix normal distribution, which ignores the separable spatial-temporal structure. The temporal covariance matrix Σ_T is constructed with elements $\sigma_{T,i,j} = 0.4^{|i-j|}$, $1 \leq i, j \leq p$. The sample size and the number of time points is set at $n = 20, q = 20$ and $n = 50, q = 30$, respectively, whereas the spatial dimension p varies among $\{50, 200, 400, 800\}$. We have

chosen this setting, since our primary interest is on inferring about spatial connectivity networks with different spatial dimensions. We keep the temporal dimension small, since it is a nuisance in our setup, and choose a relatively small sample size to reflect the fact that there is usually only a limited sample size in many neuroimaging studies.

For each generated dataset below, we use Lasso to estimate β_i as

$$\hat{\beta}_i = D_i^{-\frac{1}{2}} \arg \min_u \left\{ \frac{1}{2nq} \left| (Y_{\cdot, -i} - \bar{Y}_{(\cdot, -i)}) D_i^{-1/2} u - (Y_{(i)} - \bar{Y}_{(i)}) \right|_2^2 + \lambda_{n,i} |u|_1 \right\}, \quad (6)$$

where Y is the $nq \times p$ data matrix by stacking the transformed samples $\{(Y_{k,\cdot,l}, k = 1, \dots, n, l = 1, \dots, q)\}$, where $Y_k = X_k \Sigma_T^{-1/2}$ for the oracle procedure and $Y_k = X_k \hat{\Sigma}_T^{-1/2}$ for the data-driven procedure, $k = 1, \dots, n$, $D_i = \text{diag}(\hat{\Sigma}_{L, -i, -i})$, and $\hat{\Sigma}_L$ is the sample covariance matrix of Σ_L with nq transformed samples, and $\lambda_{n,i} = \kappa \{\hat{\Sigma}_{L,i,i} \log p / (nq)\}^{1/2}$.

4.1 Global testing simulation

For the global testing, the data $\{X_1, \dots, X_n\}$ are generated from a matrix normal distribution with mean zero and precision matrix $I \otimes \Omega_T$ under the null. To evaluate the power, let U be a matrix with eight random nonzero entries. The locations of four nonzero entries are selected randomly from the upper triangle of U , each with a magnitude generated randomly and uniformly from the set $[-4\{\log p / (nq)\}^{1/2}, -2\{\log p / (nq)\}^{1/2}] \cup [2\{\log p / (nq)\}^{1/2}, 4\{\log p / (nq)\}^{1/2}]$. The other four nonzero entries in the lower triangle are determined by symmetry. We set $\Omega_L = (I + U + \delta I) / (1 + \delta)$, with $\delta = |\lambda_{\min}(I + U)| + 0.05$, and choose the tuning parameter $\kappa = 2$ in (6).

The size and power, in percentage, of the global testing are reported in Table 1, based on 1000 data replications and the significance level $\alpha_1 = 0.05$. We see from Table 1 that the empirical sizes of the proposed oracle and data-driven procedures are well controlled under the significance level $\alpha_1 = 0.05$. However, for the vector normal based procedure that ignores the spatial-temporal dependence structure, there is a serious size distortion across all settings. The empirical sizes for the new procedures are slightly below the nominal level for high dimensions, due to the correlation among the variables. Similar phenomenon has also been observed and justified in Cai et al. (2013, Proposition 1).

We also see from the table that the new procedures are powerful in all settings, even though the two spatial precision matrices differ only in eight entries with the magnitude of difference of the order $\{\log p/(nq)\}^{1/2}$. For both the empirical sizes and powers, the data-driven procedure is seen to perform similarly as the oracle procedure.

4.2 Multiple testing simulation

For the multiple testing, the data $\{X_1, \dots, X_n\}$ are generated from a matrix normal distribution with mean zero and precision matrix $\Omega_L \otimes \Omega_T$. Three choices of Ω_L are considered:

Model 1: $\Omega_L^{(1)} = (\omega_{L,i,j}^{(1)})$ where $\omega_{L,i,i}^{(1)} = 1$, $\omega_{L,i,i+1}^{(1)} = \omega_{L,i+1,i}^{(1)} = 0.6$, $\omega_{L,i,i+2}^{(1)} = \omega_{L,i+2,i}^{(1)} = 0.3$ and $\omega_{L,i,j}^{(1)} = 0$ otherwise.

Model 2: $\Omega_L^{*(2)} = (\omega_{L,i,j}^{*(2)})$ where $\omega_{L,i,j}^{*(2)} = \omega_{L,j,i}^{*(2)} = 0.5$ for $i = 10(k-1) + 1$ and $10(k-1) + 2 \leq j \leq 10(k-1) + 10$, $1 \leq k \leq p/10$. $\omega_{L,i,j}^{*(2)} = 0$ otherwise. $\Omega_L^{(2)} = (\Omega_L^{*(2)} + \delta I)/(1 + \delta)$ with $\delta = |\lambda_{\min}(\Omega_L^{*(2)})| + 0.05$.

Model 3: $\Omega_L^{*(3)} = (\omega_{L,i,j}^{*(3)})$ where $\omega_{L,i,i}^{*(3)} = 1$, $\omega_{L,i,j}^{*(3)} = 0.8 \times \text{Bernoulli}(1, 2/p)$ for $i < j$ and $\omega_{L,j,i}^{*(3)} = \omega_{L,i,j}^{*(3)}$. $\Omega_L^{(3)} = (\Omega_L^{*(3)} + \delta I)/(1 + \delta)$ with $\delta = |\lambda_{\min}(\Omega_L^{*(3)})| + 0.05$.

We select the tuning parameters $\lambda_{n,i}$ in (6) in the Lasso estimation adaptively given the data, following the general principle of making $\sum_{(i,j) \in \mathcal{H}_0} I(|W_{i,j}| \geq t)$ and $\{2 - 2\Phi(t)\}(p^2 - p)/2$ close. The steps of parameter tuning are summarized as follows.

Step 1. Let $\lambda_{n,i} = b/20\sqrt{\hat{\Sigma}_{L,i,i} \log p/(nq)}$, for $b = 1, \dots, 40$. For each b , calculate $\hat{\beta}_i^{(b)}$, $i = 1, \dots, p$, and construct the corresponding standardized statistics $W_{i,j}^{(b)}$.

Step 2. Choose \hat{b} as the minimizer of

$$\sum_{s=1}^{10} \left(\frac{\sum_{(i,j) \in \mathcal{H}} I(|W_{i,j}|^{(b)} \geq \Phi^{-1}[1 - s\{1 - \Phi(\sqrt{\log p})\}/10])}{s\{1 - \Phi(\sqrt{\log p})\}/10 \cdot p(p-1)} - 1 \right)^2.$$

Step 3. The tuning parameters $\lambda_{n,i}$ are then set as,

$$\lambda_{n,i} = \hat{b}/20\sqrt{\hat{\Sigma}_{L,i,i} \log p/(nq)}.$$

	method	$p = 50$	$p = 200$	$p = 400$	$p = 800$
		Size			
$n = 30, q = 20$	oracle	3.6	3.5	2.8	2.9
	data-driven	3.8	3.8	2.9	2.9
	vector normal	36.4	56.7	64.7	75.3
		Size			
$n = 50, q = 30$	oracle	3.7	3.5	5.2	4.3
	data-driven	3.5	3.5	5.1	4.1
	vector normal	39.7	64.8	73.5	88.4
		Power			
$n = 30, q = 20$	oracle	77.9	85.6	87.7	90.9
	data-driven	83.1	88.1	87.8	90.7
	vector normal	86.2	94.1	95.0	99.1
$n = 50, q = 30$	oracle	62.4	74.3	68.2	75.2
	data-driven	66.1	74.7	68.3	75.5
	vector normal	76.2	92.1	93.1	90.1

Table 1: Empirical sizes and powers (%) for global testing. Three methods are compared: the proposed oracle and data-driven procedures based on the matrix normal distribution, and the simple alternative based on the vector normal distribution that ignores the spatial-temporal dependency. The results are based on 1000 data replications.

For comparison, we also carry out the alternative procedure that ignores the Kronecker product structure by using the stacked original data samples $\{X_{k,\cdot,l}, k = 1, \dots, n, l = 1, \dots, q\}$.

The empirical FDR and the empirical power of FDR control, in percentage, are summarized in Tables 2 and 3, based on 100 data replications and the FDR level set at $\alpha_2 = 0.1$ and $\alpha_3 = 0.01$, respectively. In particular, the power is calculated as

$$\frac{1}{100} \sum_{l=1}^{100} \frac{\sum_{(i,j) \in \mathcal{H}_1} I(|W_{i,j,l}| \geq \hat{t})}{|\mathcal{H}_1|},$$

where $W_{i,j,l}$ denotes standardized statistic for the l -th replication and \mathcal{H}_1 denotes the nonzero locations. We observe from Tables 2 and 3 a similar pattern as that from Table 1. That is, the empirical FDRs of the proposed oracle and data-driven procedures are both close to the significance levels across all settings, whereas the vector normal based procedure ignoring the spatial-temporal dependence structure yields empirical FDRs much larger than the significance levels. We also see from the table that the new procedures achieve a high power, and the data-driven procedure again performs similarly as the oracle procedure.

5 Real Data Analysis

We illustrate our testing method on an electroencephalography (EEG) data. The data was collected in a study examining EEG correlates of genetic predisposition to alcoholism and is available at <http://kdd.ics.uci.edu/datasets/eeg/eeg.data.html>. It consists of 77 alcoholic individuals and 45 controls, and each subject was fitted with a 61-lead electrode cap and was recorded 256 samples per second for one second. There were in addition a ground and two bipolar deviation electrodes, which are excluded from the analysis. The electrode positions were located at standard sites (Standard Electrode Position Nomenclature, American Electroencephalographic Association 1990), and were organized into frontal, central, parietal, occipital, left temporal and right temporal regions. Each subject performed 120 trials under three types of stimuli. More details of data collection can be

		Model 1				Model 2				Model 3			
α	p	50	200	400	800	50	200	400	800	50	200	400	800
		Empirical FDR (in %)											
0.1	oracle	7.4	6.7	6.4	6.0	8.9	8.5	8.2	7.8	8.9	8.4	8.0	7.8
	data-driven	8.0	6.9	6.5	6.0	11.4	9.9	9.0	8.2	11.4	9.3	8.4	7.9
	vector normal	18.9	23.1	22.7	20.7	24.5	34.1	36.5	36.5	26.1	32.0	33.5	33.8
0.01	oracle	0.6	0.5	0.4	0.7	0.9	0.8	0.6	0.5	0.8	0.7	0.6	0.7
	data-driven	0.6	0.5	0.4	0.7	1.2	0.9	0.7	0.5	1.2	0.9	0.7	0.7
	vector normal	2.5	3.2	2.6	5.6	4.6	5.6	5.4	2.2	4.4	5.3	0.5	5.3
		Empirical Power (in %)											
0.1	oracle	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	99.9	99.9	99.7
	data-driven	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	99.9	99.9	99.7
	vector normal	100.0	99.9	99.9	99.9	100.0	100.0	100.0	100.0	100.0	99.8	99.6	99.2
0.01	oracle	99.9	99.9	99.8	100.0	100.0	100.0	100.0	99.8	100.0	99.7	99.2	98.5
	data-driven	99.9	99.9	99.8	100.0	99.9	99.9	100.0	99.8	100.0	99.6	99.1	98.5
	vector normal	99.8	99.7	99.6	99.7	99.9	99.7	99.7	99.4	99.9	99.0	97.9	96.7

Table 2: Empirical FDRs and powers (%) for multiple testing with $n = 20$ and $q = 20$. Three methods are compared: the proposed oracle and data-driven procedures based on the matrix normal distribution, and the simple alternative based on the vector normal distribution that ignores the spatial-temporal dependency. The results are based on 100 data replications.

		Model 1				Model 2				Model 3			
α	p	50	200	400	800	50	200	400	800	50	200	400	800
		Empirical FDR (in %)											
0.1	oracle	8.1	7.7	8.0	7.7	8.9	9.1	8.8	8.6	8.9	8.7	8.4	8.2
	data-driven	8.4	7.9	8.1	7.8	10.3	10.0	9.2	8.9	11.7	9.3	8.8	8.2
	vector normal	23.3	29.8	33.6	36.1	28.2	39.8	44.4	49.2	29.0	37.5	42.2	45.8
0.01	oracle	0.7	0.6	0.6	0.6	1.1	0.8	0.7	0.8	0.7	0.8	0.8	0.7
	data-driven	0.7	0.6	0.6	0.6	1.2	0.9	0.8	0.8	0.9	0.9	0.8	0.7
	vector normal	3.8	4.8	5.5	5.6	5.9	7.1	7.3	9.1	5.1	6.5	7.3	8.3
		Empirical Power (in %)											
0.1	oracle	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
	data-driven	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
	vector normal	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
0.1	oracle	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
	data-driven	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
	vector normal	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Table 3: Empirical FDRs and powers (%) for multiple testing with $n = 50$ and $q = 30$. Three methods are compared: the proposed oracle and data-driven procedures based on the matrix normal distribution, and the simple alternative based on the vector normal distribution that ignores the spatial-temporal dependency. The results are based on 100 data replications.

found in Zhang et al. (1995). We preprocessed the data in the following ways. Similarly as Li et al. (2010), we focused only on the average of all trials under a single stimulus condition for each subject. We then performed an α band filtering on the signals following Hayden et al. (2006). Finally, we downsized the temporal dimension by averaging signals at eight consecutive times points. This is to facilitate estimation of the temporal covariance in our testing procedure since it is treated as a nuisance. More discussion of temporal covariance estimation is given in Section 6. The resulting data is a 61×32 matrix for each subject, and our goal is to infer the 61×61 connectivity network of the brain spatial locations.

We applied our testing procedures for the alcoholic and control groups separately. We first applied the global test and obtained the p -value 0 for the alcoholic group, and $1.89e^{-15}$ for the control, clearly indicating that some brain regions are connected in the two groups. We then applied the data-driven multiple testing procedure, with a pre-specified FDR significance level $\alpha = 0.01$. There were totally $61 \times 60/2 = 1,830$ pairs of spatial locations, among which we identified 208 significant pairs for the alcoholic group, and 155 pairs for the control group. For graphical illustration, we report the top 30 most significant pairs of spatial locations, ordered by their p -values, in Figure 1. Examining the connection patterns among those electrodes in the frontal region (denoted by symbols FP, AF, and F), we noted a slight decrease in connections and some asymmetry between the left and right frontal regions in the alcoholic group compared to the control. A similar phenomenon has been observed in Hayden et al. (2006). We observed similar connection decrease in the central region (FC and C) for the alcoholic group, but connection increase for the parietal region (CP and P). Such findings require additional scientific validation. We also repeated the multiple testing procedure for the downsized EEG data with temporal dimension equal to 16, and observed similar but less clear patterns. For the sake of space, we omit the plots here. In summary, our testing procedure produces a useful list of connections warranted for further examination.




Figure 1: Connectivity network inferred by the multiple testing procedure for the EEG data. The left panel is for the alcoholic group, and the right panel for the control. Top 30 significant links are shown in this graph.

6 Discussion

We have proposed in this article global and multiple testing procedures under the matrix normal distribution for detecting the conditional dependence between spatial locations. It has been shown that the procedures perform well both theoretically and numerically. In this section, we discuss the strength and some potential limitations of our methods. We also explore some alternatives and point out possible future extension of our work.

Our work is one of the few to tackle hypothesis testing aspect of connectivity analysis, and is a useful complement to the general literature of statistical estimation and inference regarding high-dimensional precision matrix. The tests and associated theory have been built upon some of the early works on precision matrix inference such as Liu (2013) and Xia et al. (2015). However, our work is the first that explicitly exploits the special covariance (precision) structure of a matrix-variate normal distribution, and our simulations suggest using such information would improve the test. A strength of our method is that

it works for an adequately large network; in our simulations, the number of nodes varied from 50 to 800, which encompasses the typical region/location-based connectivity analysis in neuroscience. A potential limitation is that our method treats the temporal component as nuisance, and the developed tests hinge on the accuracy of the temporal covariance estimation. In this article, we have simply used the usual sample covariance matrix $\hat{\Sigma}_T$ to estimate Σ_T . As such the method usually prefers a relatively small temporal dimension.

Then a potential extension of our proposal is to employ some alternative estimators of Σ_T . We note that, many such estimators can be used in conjunction with our proposed testing procedures, as long as the estimator $\tilde{\Sigma}_T$ satisfies the condition,

$$\|\tilde{\Sigma}_T - c\Sigma_T\|_\infty = O_p[\{\log q/(np)\}^{1/2}],$$

for arbitrary constant $c > 0$. As an example, if the temporal covariance matrix is sparse, in the sense that $\max_{1 \leq i \leq q} \sum_{j=1}^q I(\sigma_{T,i,j} \neq 0) \leq c'$ for some constant $c' > 0$, then the adaptive thresholding estimator proposed in Cai and Liu (2011) satisfies the above condition and can be used in our testing procedures. Another alternative is that one can directly estimate the precision matrix Ω_T , and base the testing procedures on $\{X_k \hat{\Omega}_T^{1/2}\}_{k=1}^n$, as long as the estimator $\tilde{\Omega}_T$ satisfies that,

$$\|\tilde{\Omega}_T - c\Omega_T\|_\infty = O_p\left[\|\Omega_T\|_{L_1}^2 \{\log q/(np)\}^{1/2}\right],$$

for any constant $c > 0$. For instance, if the precision matrix Ω_T belongs to the following uniformity class of matrices

$$\mathcal{U} = \left\{ \Omega_T : \max_{1 \leq i \leq q} \sum_{j=1}^q |\omega_{T,i,j}|^l \leq s_0(q) \right\},$$

for $0 \leq l < 1$, then the clime estimator of Cai et al. (2011) can be employed.

In this article, we have primarily focused on the one-sample inference scenario, where we assume the matrix-valued observations came from a single population. In the EEG data analysis, we simply carried out the tests for the two populations separately. Practically, it is of equal interest to study the two-sample scenario, where the aim is to

identify changes of the conditional dependence structures for spatial locations across two or more populations. Specifically, let $X, Y \in \mathbb{R}^{p \times q}$ follow two matrix normal distributions with the Kronecker product covariance structures, $\Sigma\{\text{vec}(X)\} = \Sigma_{L1} \otimes \Sigma_{T1}$, and $\Sigma\{\text{vec}(Y)\} = \Sigma_{L2} \otimes \Sigma_{T2}$, respectively. The goal becomes the global testing of

$$H_0 : \Omega_{L1} = \Omega_{L2} \text{ versus } H_1 : \Omega_{L1} \neq \Omega_{L2},$$

and simultaneous testing of

$$H_{0,i,j} : \omega_{L1,i,j} = \omega_{L2,i,j} \text{ versus } H_{1,i,j} : \omega_{L1,i,j} \neq \omega_{L2,i,j}, \quad 1 \leq i < j \leq p,$$

where $\omega_{L1,i,j}$ is the (i, j) th element of $\Sigma_{L1}^{-1} = \Omega_{L1} = (\omega_{L1,i,j})$, and $\omega_{L2,i,j}$ is the (i, j) th element of $\Sigma_{L2}^{-1} = \Omega_{L2} = (\omega_{L2,i,j})$. In the one-sample scenario, we have used the sample covariance matrix $\hat{\Sigma}_T$ to estimate temporal covariance matrix, because it can be turned into an unbiased estimator without affecting the test statistics, as shown in Remark 2. However, in the two-sample scenario, Remark 2 can no longer apply, and thus $\text{trace}(\Sigma_{L1})$ and $\text{trace}(\Sigma_{L2})$ need to be carefully estimated in order to get good estimators for temporal covariance matrices. Consequently, the two-sample comparison is technically much more challenging, and we leave it as our future research.

7 Appendix: Proofs

7.1 Technical Lemmas

We prove the main results in this section. We begin by collecting some technical lemmas.

Lemma 1 (Bonferroni inequality) *Let $B = \cup_{t=1}^p B_t$. For any $k < [p/2]$, we have*

$$\sum_{t=1}^{2k} (-1)^{t-1} F_t \leq \text{pr}(B) \leq \sum_{t=1}^{2k-1} (-1)^{t-1} F_t,$$

where $F_t = \sum_{1 \leq i_1 < \dots < i_t \leq p} \text{pr}(B_{i_1} \cap \dots \cap B_{i_t})$.

Define $U_{i,j} = \frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q \{\epsilon_{k,i,l} \epsilon_{k,j,l} / (r_{i,i} r_{j,j}) - E \epsilon_{k,i,l} \epsilon_{k,j,l} / (r_{i,i} r_{j,j})\}$. The follow lemma states the results in the oracle case.

Lemma 2 Suppose that (C1) and (C2) hold. Then we have

$$\max_{1 \leq i \leq p} |\hat{r}_{i,i} - r_{i,i}| = o_p[\{\log p/(nq)\}^{1/2}],$$

and

$$\tilde{r}_{i,j} = \tilde{R}_{i,j} - \tilde{r}_{i,i}(\hat{\beta}_{i,j} - \beta_{i,j}) - \tilde{r}_{j,j}(\hat{\beta}_{j-1,i} - \beta_{j-1,i}) + o_p\{(nq \log p)^{-1/2}\},$$

for $1 \leq i < j \leq p$, where $\tilde{R}_{i,j}$ is the empirical covariance between $\{\epsilon_{k,i,l}, k = 1, \dots, n, l = 1, \dots, q\}$ and $\{\epsilon_{k,j,l}, k = 1, \dots, n, l = 1, \dots, q\}$. Consequently, uniformly in $1 \leq i < j \leq p$,

$$\hat{r}_{i,j} - (\omega_{L,i,i}\hat{\sigma}_{i,i,\epsilon} + \omega_{L,j,j}\hat{\sigma}_{j,j,\epsilon} - 1)r_{i,j} = -U_{i,j} + o_p\{(nq \log p)^{-1/2}\},$$

where $(\hat{\sigma}_{i,j,\epsilon}) = \frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q (\epsilon_{k,,l} - \bar{\epsilon}_l)(\epsilon_{k,,l} - \bar{\epsilon}_l)'$, $\epsilon_{k,,l} = (\epsilon_{k,1,l}, \dots, \epsilon_{k,p,l})$ and $\bar{\epsilon}_l = \frac{1}{n} \sum_{k=1}^n \epsilon_{k,,l}$.

This lemma is essentially proved in Xia et al. (2015) with nq inverse regression models instead.

7.2 Proof of Theorem 1

Without loss of generality, throughout this section, we assume that $\omega_{L,i,i} = 1$ for $i = 1, \dots, p$. Let

$$V_{i,j} = -U_{i,j}/\{\text{var}(\epsilon_{k,i,1}\epsilon_{k,j,1})/nq\}^{1/2}.$$

By Lemma 2, we have

$$\max_{1 \leq i \leq p} |\hat{r}_{i,i} - r_{i,i}| = O_p[\{\log p/(nq)\}^{\frac{1}{2}}], \quad (7)$$

and

$$\max_{1 \leq i \leq p} |\hat{r}_{i,i} - \tilde{R}_{i,i}| = o_p\{(nq \log p)^{-1/2}\}.$$

Note that

$$\max_{1 \leq i < j \leq p} (\hat{\beta}_{i,j}^2 \hat{r}_{i,i} / \hat{r}_{j,j} - \rho_{i,j}^2) = o_p(1/\log p), \quad (8)$$

Then by Lemma 2, it is easy to see that, under conditions (C1) and (C2), we have under H_0

$$\max_{1 \leq i < j \leq p} ||W_{i,j}| - |V_{i,j}|| = o_p\{(\log p)^{-1/2}\}.$$

Thus it suffices to prove that

$$\text{pr}\left(\max_{1 \leq i < j \leq p} V_{i,j}^2 - 4 \log p + \log \log p \leq t\right) \rightarrow \exp\{-(8\pi)^{-1/2} \exp(-t/2)\}.$$

We arrange the indices $\{1 \leq i < j \leq p\}$ in any ordering and set them as $\{(i_m, j_m) : m = 1, \dots, s\}$ with $s = p(p-1)/2$. Let $\theta_m = \text{var}(\epsilon_{k,i_m,l} \epsilon_{k,j_m,l})$, and define $Z_{k,m,l} = \epsilon_{k,i_m,l} \epsilon_{k,j_m,l}$ for $1 \leq k \leq n$ and $1 \leq l \leq q$, $V_m = (nq\theta_m)^{-1/2} \sum_{k=1}^n \sum_{l=1}^q Z_{k,m,l}$, and $\hat{V}_m = (nq\theta_m)^{-1/2} \sum_{k=1}^n \sum_{l=1}^q \hat{Z}_{k,m,l}$, where $\hat{Z}_{k,m,l} = Z_{k,m,l} I(|Z_{k,m,l}| \leq \tau_n) - E\{Z_{k,m,l} I(|Z_{k,m,l}| \leq \tau_n)\}$, and $\tau_n = 32 \log(p + nq)$. Note that $\max_{1 \leq i < j \leq p} V_{i,j}^2 = \max_{1 \leq m \leq s} V_m^2$, and that

$$\begin{aligned} \max_{1 \leq m \leq s} (nq)^{-1/2} \sum_{k=1}^n \sum_{l=1}^q E[|Z_{k,m,l}| I\{|Z_{k,m,l}| \geq 32 \log(p + nq)\}] \\ \leq C(nq)^{1/2} \max_{1 \leq k \leq n} \max_{1 \leq l \leq q} \max_{1 \leq m \leq s} E[|Z_{k,m,l}| I\{|Z_{k,m,l}| \geq 32 \log(p + nq)\}] \\ \leq C(nq)^{1/2} (p + nq)^{-4} \max_{1 \leq k \leq n} \max_{1 \leq l \leq q} \max_{1 \leq m \leq s} E[|Z_{k,m,l}| \exp\{|Z_{k,m,l}|/8\}] \\ \leq C(nq)^{1/2} (p + nq)^{-4}. \end{aligned}$$

This yields to

$$\text{pr}\left\{\max_{1 \leq m \leq s} |V_m - \hat{V}_m| \geq (\log p)^{-1}\right\} \leq \text{pr}\left(\max_{1 \leq m \leq s} \max_{1 \leq k \leq n} \max_{1 \leq l \leq q} |Z_{k,m,l}| \geq \tau_n\right) = O(p^{-1}).$$

Note that

$$\left| \max_{1 \leq m \leq s} V_m^2 - \max_{1 \leq m \leq s} \hat{V}_m^2 \right| \leq 2 \max_{1 \leq m \leq s} |\hat{V}_m| \max_{1 \leq m \leq s} |V_m - \hat{V}_m| + \max_{1 \leq m \leq s} |V_m - \hat{V}_m|^2,$$

it suffices to prove that for any $t \in \mathbb{R}$, as $nq, p \rightarrow \infty$,

$$\text{pr}\left(\max_{1 \leq m \leq s} \hat{V}_m^2 - 4 \log p + \log \log p \leq t\right) \rightarrow \exp\{-(8\pi)^{-1/2} \exp(-t/2)\}. \quad (9)$$

By Lemma 1, for any integer l with $0 < h < s/2$,

$$\begin{aligned} \sum_{d=1}^{2h} (-1)^{d-1} \sum_{1 \leq m_1 < \dots < m_d \leq q} \text{pr}\left(\bigcap_{j=1}^d F_{m_j}\right) &\leq \text{pr}\left(\max_{1 \leq m \leq s} \hat{V}_m^2 \geq y_p\right) \\ &\leq \sum_{d=1}^{2h-1} (-1)^{d-1} \sum_{1 \leq m_1 < \dots < m_d \leq q} \text{pr}\left(\bigcap_{j=1}^d F_{m_j}\right), \quad (10) \end{aligned}$$

where $y_p = 4 \log p - \log \log p + t$ and $F_{m_j} = (\hat{V}_{m_j}^2 \geq y_p)$. Let $\tilde{Z}_{k,m,l} = \hat{Z}_{k,m,l}/(\theta_m)^{1/2}$ for $m = 1, \dots, s$ and $W_{k,l} = (\tilde{Z}_{k,m_1,l}, \dots, \tilde{Z}_{k,m_d,l})$, for $1 \leq k \leq n$ and $1 \leq l \leq q$. Define $|a|_{\min} = \min_{1 \leq i \leq d} |a_i|$ for any vector $a \in R^d$. Then we have

$$\Pr\left(\bigcap_{j=1}^d F_{m_j}\right) = \Pr\left(\left|(nq)^{-\frac{1}{2}} \sum_{k=1}^n \sum_{l=1}^q W_{k,l}\right|_{\min} \geq y_p^{\frac{1}{2}}\right).$$

Then it follows from Theorem 1 in Zaitsev (1987) that

$$\Pr\left(\left|(nq)^{-1/2} \sum_{k=1}^n \sum_{l=1}^q W_{k,l}\right|_{\min} \geq y_p^{1/2}\right) \leq \Pr\left\{|N_d|_{\min} \geq y_p^{1/2} - \epsilon_n(\log p)^{-1/2}\right\} + c_1 d^{\frac{5}{2}} \exp\left\{-\frac{(nq)^{1/2} \epsilon_n}{c_2 d^3 \tau_n (\log p)^{1/2}}\right\}, \quad (11)$$

where $c_1 > 0$ and $c_2 > 0$ are constants, $\epsilon_n \rightarrow 0$ which will be specified later and $N_d = (N_{m_1}, \dots, N_{m_d})$ is a normal random vector with $E(N_d) = 0$ and $\text{cov}(N_d) = \text{cov}(W_{1,1})$. Recall that d is a fixed integer which does not depend on n, p, q . Because $\log p = o((nq)^{1/5})$, we can let $\epsilon_n \rightarrow 0$ sufficiently slowly that, for any large $M > 0$

$$c_1 d^{5/2} \exp\left\{-\frac{(nq)^{1/2} \epsilon_n}{c_2 d^3 \tau_n (\log p)^{1/2}}\right\} = O(p^{-M}). \quad (12)$$

Combining (10), (11) and (12) we have

$$\Pr\left(\max_{1 \leq m \leq s} \hat{V}_m^2 \geq y_p\right) \leq \sum_{d=1}^{2h-1} (-1)^{d-1} \sum_{1 \leq m_1 < \dots < m_d \leq s} \Pr\left\{|N_d|_{\min} \geq y_p^{1/2} - \epsilon_n(\log p)^{-1/2}\right\} + o(1). \quad (13)$$

Similarly, using Theorem 1 in Zaitsev (1987) again, we can get

$$\Pr\left(\max_{1 \leq m \leq s} \hat{V}_m^2 \geq y_p\right) \geq \sum_{d=1}^{2h} (-1)^{d-1} \sum_{1 \leq m_1 < \dots < m_d \leq s} \Pr\left\{|N_d|_{\min} \geq y_p^{1/2} + \epsilon_n(\log p)^{-1/2}\right\} - o(1). \quad (14)$$

We recall the following lemma, which is shown in the supplementary material of Cai et al. (2013).

Lemma 3 For any fixed integer $d \geq 1$ and real number $t \in \mathbb{R}$,

$$\sum_{1 \leq m_1 < \dots < m_d \leq q} \Pr\left\{|N_d|_{\min} \geq y_p^{1/2} \pm \epsilon_n(\log p)^{-1/2}\right\} = \frac{1}{d!} \{(8\pi)^{-1/2} \exp(-t/2)\}^d \{1 + o(1)\}. \quad (15)$$

Then Lemma 3, (13) and (14) yield that

$$\begin{aligned}\limsup_{nq,p \rightarrow \infty} \Pr\left(\max_{1 \leq m \leq s} \hat{V}_m^2 \geq y_p\right) &\leq \sum_{d=1}^{2h} (-1)^{d-1} \frac{1}{d!} \{(8\pi)^{-1/2} \exp(-t/2)\}^d \\ \liminf_{nq,p \rightarrow \infty} \Pr\left(\max_{1 \leq m \leq s} \hat{V}_m^2 \geq y_p\right) &\geq \sum_{d=1}^{2h-1} (-1)^{d-1} \frac{1}{d!} \{(8\pi)^{-1/2} \exp(-t/2)\}^d\end{aligned}$$

for any positive integer h . By letting $h \rightarrow \infty$, we obtain (9) and Theorem 1 is proved. ■

7.3 Proof of Theorem 2

By Lemma 2, we have

$$\max_{1 \leq i < j \leq p} \left| \frac{T_{i,j} - \{1 + o(1)\} ET_{i,j}}{\hat{\theta}_{i,j}^{1/2}} - V_{i,j} \right| = o_p\{(\log p)^{-1/2}\}.$$

Let

$$M_{nq}^1 = \max_{1 \leq i < j \leq p} \frac{T_{i,j} - \{1 + o(1)\} ET_{i,j}}{\hat{\theta}_{i,j}^{1/2}}.$$

Then the proof of Theorem 1 yields

$$\Pr(M_{nq}^1 \leq 4 \log p - 2^{-1} \log \log p) \rightarrow 1,$$

as $nq, p \rightarrow \infty$. By (7), (8) and the fact that

$$\max_{1 \leq i < j \leq p} \omega_{L,i,j}^2 / \hat{\theta}_{i,j} = \max_{1 \leq i < j \leq p} [\{1 + o(1)\} ET_{i,j}]^2 / \hat{\theta}_{i,j} \leq 2M_{nq}^1 + 2M_{nq},$$

and the fact that

$$\max_{1 \leq i < j \leq p} |\omega_{L,i,j}| / \theta_{i,j}^{1/2} \geq 4(\log p)^{1/2},$$

we have

$$\Pr(M_{nq} \geq q_\alpha + 4 \log p - \log \log p) \rightarrow 1$$

as $nq, p \rightarrow \infty$. ■

7.4 Proof of Theorem 3

This theorem is essentially proved in Xia et al. (2015), we skip the proof here.

7.5 Proof of Theorem 4

By separation of spatial-temporal dependence structure, we have the following nq inverse regression models

$$(X_k \Sigma_T^{-1/2})_{i,l} = \alpha_{i,l} + (X_k \Sigma_T^{-1/2})_{-i,l} \beta_i + \epsilon_{k,i,l}, \quad 1 \leq k \leq n, 1 \leq l \leq q.$$

Then Theorem 4 is proved by applying Theorem 3.1 in Liu (2013) in nq regression models. ■

7.6 Proof of Theorems 5 and 6

We use the superscript “ d ” to denote the corresponding statistics we derived from the data-driven inverse regression models

$$Y_{k,i,l}^d = (Y_{k,-i,l}^d)^\top \beta_{i,l}^d + \epsilon_{k,i,l}^d, \quad 1 \leq i \leq p, 1 \leq l \leq q,$$

where $Y_k^d = X_k \hat{\Sigma}_T^{-1/2}$, for $k = 1, \dots, n$.

By the proofs of Theorem 1 and Theorem 4, it suffices to prove that uniformly in $1 \leq i \leq j \leq p$,

$$\hat{r}_{i,j}^d - (\omega_{L,i,i} \hat{\sigma}_{i,i,\epsilon} + \omega_{L,j,j} \hat{\sigma}_{j,j,\epsilon} - 1) r_{i,j} = -U_{i,j} + o_p((nq \log p)^{-1/2}), \quad (16)$$

where $\hat{r}_{i,j}^d = -(\tilde{r}_{i,j}^d + \tilde{r}_{i,i}^d \hat{\beta}_{i,j}^d + \tilde{r}_{j,j}^d \hat{\beta}_{j-1,i}^d)$ with $\tilde{r}_{i,j}^d = 1/(nq) \sum_{k=1}^n \sum_{l=1}^q \hat{\epsilon}_{k,i,l}^d \hat{\epsilon}_{k,j,l}^d$ and $\hat{\epsilon}_{k,i,l}^d = Y_k^d - (Y_{k,-i,l}^d)^\top \hat{\beta}_{i,l}^d$. Let $\tilde{\epsilon}_{k,i,l} = \epsilon_{k,i,l} - \bar{\epsilon}_{i,l}$. Then we have

$$\begin{aligned} \hat{\epsilon}_{k,i,l}^d \hat{\epsilon}_{k,j,l}^d &= \tilde{\epsilon}_{k,i,l} \tilde{\epsilon}_{k,j,l} - \tilde{\epsilon}_{k,i,l} (Y_{k,-j,l} - \bar{Y}_{-j,l})^\top (\hat{\beta}_j^d - \beta_j) - \tilde{\epsilon}_{k,j,l} (Y_{k,-i,l} - \bar{Y}_{-i,l})^\top (\hat{\beta}_i^d - \beta_i) \\ &\quad + (\hat{\beta}_i^d - \beta_i)^\top (Y_{k,-i,l} - \bar{Y}_{-i,l}) (Y_{k,-j,l} - \bar{Y}_{-j,l})^\top (\hat{\beta}_j^d - \beta_j) \\ &\quad + \left[\{Y_{k,i,l}^d - \bar{Y}_{i,l}^d - (Y_{k,-i,l}^d - \bar{Y}_{-i,l}^d)^\top \hat{\beta}_i^d\} \{Y_{k,j,l}^d - \bar{Y}_{j,l}^d - (Y_{k,-j,l}^d - \bar{Y}_{-j,l}^d)^\top \hat{\beta}_j^d\} - \right. \\ &\quad \left. \{Y_{k,i,l} - \bar{Y}_{i,l} - (Y_{k,-i,l} - \bar{Y}_{-i,l})^\top \hat{\beta}_i^d\} \{Y_{k,j,l} - \bar{Y}_{j,l} - (Y_{k,-j,l} - \bar{Y}_{-j,l})^\top \hat{\beta}_j^d\} \right]. \end{aligned}$$

Note that

$$\hat{\Sigma}_T = \frac{1}{np} \sum_{k=1}^n X_k^\top X_k = \frac{1}{np} \sum_{k=1}^n Z_k^\top \Sigma_L Z_k,$$

where $Z_k = \Sigma_L^{-1/2} X_k$. Let $\Sigma_L = U^\top \Lambda U$, to be the eigen-decomposition of Σ_L , then we have

$$\hat{\Sigma}_T = \frac{1}{np} \sum_{k=1}^n (U Z_k)^\top \Lambda (U Z_k),$$

where rows of $U Z_k$ are independent. Thus it is easy to show that

$$\|\hat{\Sigma}_T - \Sigma_T\|_\infty = O_p \left[\{\log q/(np)\}^{1/2} \right].$$

Thus, we have

$$\|\hat{\Sigma}_T^{-1} - \Sigma_T^{-1}\|_\infty = O_p \left[\|\Omega_T\|_{L_1}^2 \{\log q/(np)\}^{1/2} \right].$$

This yields that

$$\|\hat{\Sigma}_T^{-1/2} - \Sigma_T^{-1/2}\|_\infty = \|(\hat{\Sigma}_T^{-1/2} + \Sigma_T^{-1/2})^{-1}(\hat{\Sigma}_T^{-1} - \Sigma_T^{-1})\|_\infty = O_p[\{q \log q/(np)\}^{1/2} \|\Omega_T\|_{L_1}^2],$$

which implies, uniformly for $l = 1, \dots, q$, $k = 1, \dots, n$,

$$|(X_k \hat{\Sigma}_T^{-1/2})_{\cdot, l} - (X_k \Sigma_T^{-1/2})_{\cdot, l}|_\infty = O_p[\{q^3 \log q \log \max(p, q, n)/(np)\}^{1/2} \|\Omega_T\|_{L_1}^2].$$

We shall focus on the event $\{\|\hat{\Sigma}_T - \Sigma_T\|_\infty = O[\{\log q/(np)\}^{1/2}]\}$ in the following analysis. We have, uniformly for $l = 1, \dots, q$, $k = 1, \dots, n$,

$$\|\text{cov}^{-1}[(X_k \hat{\Sigma}_T^{-1/2})_{\cdot, l}] - \Omega_L\|_\infty = O[\{q^3 \log q \log \max(p, q, n)/(np)\}^{1/2} \|\Omega_T\|_{L_1}^2 \|\Omega_L\|_{L_1}^2].$$

Thus, by (C4), it is easy to show that

$$\max_{1 \leq i \leq p, 1 \leq l \leq q} |\beta_{i,l}^d - \beta_i|_1 = o[\{\log \max(p, q, n)\}^{-1}], \text{ and } \max_{1 \leq i \leq p, 1 \leq l \leq q} |\beta_{i,l}^d - \beta_i|_2 = o\{(nq \log p)^{-1/4}\}.$$

Thus we have

$$\max_{1 \leq i \leq p} |\hat{\beta}_i^d - \beta_i|_1 = o_p[\{\log \max(p, q, n)\}^{-1}], \text{ and } \max_{1 \leq i \leq p} |\hat{\beta}_i^d - \beta_i|_2 = o_p\{(nq \log p)^{-1/4}\}.$$

Hence, by the proof of Lemma 2, we have

$$\begin{aligned} & \frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q \left\{ \tilde{\epsilon}_{k,i,l} \tilde{\epsilon}_{k,j,l} - \tilde{\epsilon}_{k,i,l} (Y_{k,-j,l} - \bar{Y}_{-j,l})^\top (\hat{\beta}_j^d - \beta_j) - \tilde{\epsilon}_{k,j,l} (Y_{k,-i,l} - \bar{Y}_{-i,l})^\top (\hat{\beta}_i^d - \beta_i) \right. \\ & \quad \left. + (\hat{\beta}_i^d - \beta_i)^\top (Y_{k,-i,l} - \bar{Y}_{-i,l}) (Y_{k,-j,l} - \bar{Y}_{-j,l})^\top (\hat{\beta}_j^d - \beta_j) \right\} \end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q (\hat{\epsilon}_{k,i,l}^d)^2 \hat{\beta}_{i,j}^d - \frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q (\hat{\epsilon}_{k,j,l}^d)^2 \hat{\beta}_{j-1,i}^d - \\
&\quad (\omega_{L,i,i} \hat{\sigma}_{i,i,\epsilon} + \omega_{L,j,j} \hat{\sigma}_{j,j,\epsilon} - 1) r_{i,j} + U_{i,j} + o_p((nq \log p)^{-1/2}),
\end{aligned}$$

uniformly in $1 \leq i \leq j \leq p$. Note that

$$\begin{aligned}
&\frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q \left[\{Y_{k,i,l}^d - \bar{Y}_{i,l}^d - (Y_{k,-i,l}^d - \bar{Y}_{-i,l}^d)^\top \hat{\beta}_i^d\} \{Y_{k,j,l}^d - \bar{Y}_{j,l}^d - (Y_{k,-j,l}^d - \bar{Y}_{-j,l}^d)^\top \hat{\beta}_j^d\} - \right. \\
&\quad \left. \{Y_{k,i,l}^d - \bar{Y}_{i,l}^d - (Y_{k,-i,l}^d - \bar{Y}_{-i,l}^d)^\top \hat{\beta}_i^d\} \{Y_{k,j,l}^d - \bar{Y}_{j,l}^d - (Y_{k,-j,l}^d - \bar{Y}_{-j,l}^d)^\top \hat{\beta}_j^d\} \right] \\
&= \frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q \left(\left\{ Y_{k,i,l}^d - \bar{Y}_{i,l}^d - (Y_{k,-i,l}^d - \bar{Y}_{-i,l}^d)^\top \hat{\beta}_i^d \right\} \left[\{Y_{k,j,l}^d - \bar{Y}_{j,l}^d - (Y_{k,j,l}^d - \bar{Y}_{j,l}^d)\} \right. \right. \\
&\quad \left. \left. - \{Y_{k,-j,l}^d - \bar{Y}_{-j,l}^d - (Y_{k,-j,l}^d - \bar{Y}_{-j,l}^d)^\top \hat{\beta}_j^d\} \right] + \left\{ Y_{k,j,l}^d - \bar{Y}_{j,l}^d - (Y_{k,-j,l}^d - \bar{Y}_{-j,l}^d)^\top \hat{\beta}_j^d \right\} \right. \\
&\quad \left. \left[\{Y_{k,i,l}^d - \bar{Y}_{i,l}^d - (Y_{k,i,l}^d - \bar{Y}_{i,l}^d)\} - \{Y_{k,-i,l}^d - \bar{Y}_{-i,l}^d - (Y_{k,-i,l}^d - \bar{Y}_{-i,l}^d)^\top \hat{\beta}_i^d\} \right] \right)
\end{aligned}$$

It suffices to show that, uniformly in $1 \leq i \leq j \leq p$,

$$\begin{aligned}
s_{1,i,j} &= \frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q \left\{ Y_{k,i,l}^d - \bar{Y}_{i,l}^d - (Y_{k,-i,l}^d - \bar{Y}_{-i,l}^d)^\top \hat{\beta}_i^d \right\} \left[\{Y_{k,j,l}^d - \bar{Y}_{j,l}^d - (Y_{k,j,l}^d - \bar{Y}_{j,l}^d)\} \right. \\
&\quad \left. - \{Y_{k,-j,l}^d - \bar{Y}_{-j,l}^d - (Y_{k,-j,l}^d - \bar{Y}_{-j,l}^d)^\top \hat{\beta}_j^d\} \right] = o_p\{(nq \log p)^{-1/2}\}. \quad (17)
\end{aligned}$$

Recall that $\hat{\epsilon}_{k,i,l}^d = Y_{k,i,l}^d - \bar{Y}_{i,l}^d - (Y_{k,-i,l}^d - \bar{Y}_{-i,l}^d)^\top \hat{\beta}_i^d$, then we have

$$s_{1,i,j} = \frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q \left(\hat{\epsilon}_{k,i,l}^d \left[\{Y_{k,j,l}^d - \bar{Y}_{j,l}^d - (Y_{k,j,l}^d - \bar{Y}_{j,l}^d)\} - \sum_{h \neq j} \{Y_{k,h,l}^d - \bar{Y}_{h,l}^d - (Y_{k,h,l}^d - \bar{Y}_{h,l}^d)\} \hat{\beta}_{h,j}^d \right] \right).$$

Note that $\hat{\epsilon}_{k,i,l}^d = \epsilon_{k,i,l}^d + (Y_{k,-i,l}^d - \bar{Y}_{-i,l}^d)^\top (\beta_{i,l}^d - \hat{\beta}_i^d)$. Let event $A = (\max_{1 \leq i \leq p, 1 \leq l \leq q} |\beta_i^d - \hat{\beta}_i^d|_1 = o[\{\log \max(p, q, n)\}^{-1}])$. Then we have

$$\begin{aligned}
\max_{1 \leq i \leq p, 1 \leq l \leq q} |E(\hat{\epsilon}_{k,i,l}^d | A)| &\leq \max_{1 \leq i \leq p, 1 \leq l \leq q} |E\epsilon_{k,i,l}^d| + \max_{1 \leq i \leq p, 1 \leq l \leq q} |E\{(Y_{k,-i,l}^d - \bar{Y}_{-i,l}^d)^\top (\beta_{i,l}^d - \hat{\beta}_i^d) | A| \\
&= o[\{\log \max(p, q, n)\}^{-1/2}].
\end{aligned}$$

It is easy to check that, for arbitrarily small $\gamma > 0$,

$$\begin{aligned}
&\max_{1 \leq i \leq p, 1 \leq l \leq q, 1 \leq k \leq n} |\{Y_{k,i,l}^d - \bar{Y}_{i,l}^d - (Y_{k,i,l}^d - \bar{Y}_{i,l}^d)\}| \\
&= \max_{1 \leq i \leq p, 1 \leq l \leq q, 1 \leq k \leq n} \left| \left\{ \left(X_k - 1/n \sum_{k=1}^n X_k \right) \Sigma_T^{-1/2} \right\}_{i,l} - \left\{ \left(X_k - 1/n \sum_{k=1}^n X_k \right) \hat{\Sigma}_T^{-1/2} \right\}_{i,l} \right|
\end{aligned}$$

$$= o_p \left[\{q^3 \log q \log \max(p, q, n) \log^\gamma p / (np)\}^{1/2} \|\Omega_T\|_{L_1}^2 \right].$$

Let event

$$B = \left\{ \max_{1 \leq i \leq p, 1 \leq l \leq q, 1 \leq k \leq n} |\{Y_{k,i,l} - \bar{Y}_{i,l} - (Y_{k,i,l}^d - \bar{Y}_{i,l}^d)\}| \right. \\ \left. = o \left[\{q^3 \log q \log \max(p, q, n) \log^\gamma p / (np)\}^{1/2} \|\Omega_T\|_{L_1}^2 \right] \right\}.$$

Thus, by the fact that $\max_{1 \leq i \leq p, 1 \leq l \leq q, 1 \leq k \leq n} |\hat{\epsilon}_{k,i,l}^d - \epsilon_{k,i,l}| = o_p[\{\log \max(p, q, n)\}^{-1/2}]$, it can be shown that, for arbitrarily small $\gamma > 0$,

$$\begin{aligned} & \text{pr} \left(\max_{1 \leq i \leq p} \max_{1 \leq h \leq p} \left| \frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q \hat{\epsilon}_{k,i,l}^d \{Y_{k,h,l}^d - \bar{Y}_{h,l}^d - (Y_{k,h,l} - \bar{Y}_{h,l})\} \right| \right. \\ & \geq C \sqrt{\frac{q^2 \log q \log \max(p, q, n) \log^{1+\gamma} p}{n^2 p} \|\Omega_T\|_{L_1}^2} \Bigg) \\ & \leq \text{pr} \left(\max_{1 \leq i \leq p} \max_{1 \leq h \leq p} \left| \frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q \hat{\epsilon}_{k,i,l}^d \{Y_{k,h,l}^d - \bar{Y}_{h,l}^d - (Y_{k,h,l} - \bar{Y}_{h,l})\} \right| \right. \\ & \geq C \sqrt{\frac{q^2 \log q \log \max(p, q, n) \log^{1+\gamma} p}{n^2 p} \|\Omega_T\|_{L_1}^2}, A \cap B \Bigg) + \text{pr}(A^c) + \text{pr}(B^c) \\ & \leq \text{pr} \left(\max_{1 \leq i \leq p} \max_{1 \leq h \leq p} \left| \frac{1}{\sqrt{nq}} \sum_{k=1}^n \sum_{l=1}^q \hat{\epsilon}_{k,i,l}^d \frac{\{Y_{k,h,l}^d - \bar{Y}_{h,l}^d - (Y_{k,h,l} - \bar{Y}_{h,l})\}}{\sqrt{q^3 \log q \log \max(p, q, n) \log^\gamma p / np}} \right| \right. \\ & \geq C \sqrt{\log p}, A \cap B \Bigg) + o(1) = o(1). \end{aligned}$$

This, together with (C4), implies that

$$\begin{aligned} & \max_{1 \leq i \leq j \leq p} \left| \sum_{h \neq j} \frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q \hat{\epsilon}_{k,i,l}^d \{Y_{k,h,l}^d - \bar{Y}_{h,l}^d - (Y_{k,h,l} - \bar{Y}_{h,l})\} \hat{\beta}_{h,j}^d \right| \\ & = O_p \left(\sqrt{\frac{q^2 \log q \log \max(p, q, n) \log^{1+\gamma} p}{n^2 p}} \right) \|\Omega_T\|_{L_1}^2 \|\Omega_L\|_{L_1} = o_p\{(nq \log p)^{-1/2}\}. \end{aligned}$$

Thus (17) is proved. Thus we have

$$\begin{aligned} & \left| \frac{1}{nq} \sum_{k=1}^n \sum_{l=1}^q \left[\{Y_{k,i,l}^d - \bar{Y}_{i,l}^d - (Y_{k,-i,l}^d - \bar{Y}_{-i,l}^d)^\top \hat{\beta}_i^d\} \{Y_{k,j,l}^d - \bar{Y}_{j,l}^d - (Y_{k,-j,l}^d - \bar{Y}_{-j,l}^d)^\top \hat{\beta}_j^d\} - \right. \right. \\ & \quad \left. \left. \{Y_{k,i,l} - \bar{Y}_{i,l} - (Y_{k,-i,l} - \bar{Y}_{-i,l})^\top \hat{\beta}_i^d\} \{Y_{k,j,l} - \bar{Y}_{j,l} - (Y_{k,-j,l} - \bar{Y}_{-j,l})^\top \hat{\beta}_j^d\} \right] \right| \\ & = o_p\{(nq \log p)^{-1/2}\} \end{aligned}$$

uniformly in $1 \leq i \leq j \leq p$. Hence equation (16) is proved and Theorems 5 and 6 thus follow. ■

References

- Anderson, T. W. (2003). *An Introduction To Multivariate Statistical Analysis*. Wiley-Interscience, 3rd ed, New York.
- Aston, J. A. and Kirch, C. (2012). Estimation of the distribution of change-points with application to fmri data. *Annals of Applied Statistics*, 6:1906–1948.
- Bullmore, E. and Sporns, O. (2009). Complex brain networks: graph theoretical analysis of structural and functional systems. *Nature reviews. Neuroscience*, 10(3):186–198.
- Cai, T. T. and Liu, W. (2011). Adaptive thresholding for sparse covariance matrix estimation. *J. Am. Statist. Assoc.*, 106(494):672–684.
- Cai, T. T., Liu, W., and Luo, X. (2011). A constrained ℓ_1 minimization approach to sparse precision matrix estimation. *J. Amer. Statist. Assoc.*, 106(494):594–607.
- Cai, T. T., Liu, W., and Xia, Y. (2013). Two-sample covariance matrix testing and support recovery in high-dimensional and sparse settings. *J. Amer. Statist. Assoc.*, 108(501):265–277.
- Danaher, P., Wang, P., and Witten, D. M. (2014). The joint graphical lasso for inverse covariance estimation across multiple classes. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 76(2):373–397.
- Fornito, A., Zalesky, A., and Breakspear, M. (2013). Graph analysis of the human connectome: Promise, progress, and pitfalls. *NeuroImage*, 80:426–444.
- Friedman, J., Hastie, T., and Tibshirani, R. (2008). Sparse inverse covariance estimation with the graphical lasso. *Biostatistics*, 9(3):432–441.
- Friston, K., Ashburner, J., Kiebel, S., Nichols, T., and Penny, W., editors (2007). *Statistical Parametric Mapping: The Analysis of Functional Brain Images*. Academic Press.
- Hayden, E. P., Wiegand, R. E., Meyer, E. T., Bauer, L. O., O’Connor, S. J., Nurnberger, J. I., Chorlian, D. B., Porjesz, B., and Begleiter, H. (2006). Patterns of regional brain ac-

- tivity in alcohol-dependent subjects. *Alcoholism: Clinical and Experimental Research*, 30(12):1986–1991.
- Johansen-Berg, H. (2013). Human connectomics - what will the future demand? *NeuroImage*, 80:541–544.
- Leng, C. and Tang, C. Y. (2012). Sparse matrix graphical models. *J. Amer. Statist. Assoc.*, 107(499):1187–1200.
- Li, B., Kim, M. K., and Altman, N. (2010). On dimension folding of matrix- or array-valued statistical objects. *The Annals of Statistics*, 38(2):1094–1121.
- Liu, H., Han, F., Yuan, M., Lafferty, J., and Wasserman, L. (2012). High-dimensional semiparametric Gaussian copula graphical models. *Ann. Statist.*, 40(4):2293–2326.
- Liu, W. (2013). Gaussian graphical model estimation with false discovery rate control. *Ann. Statist.*, 41(6):2948–2978.
- Meinshausen, N. and Bühlmann, P. (2006). High-dimensional graphs and variable selection with the lasso. *The Annals of Statistics*, pages 1436–1462.
- Olshausen, B. and Field, D. (2004). Sparse coding of sensory inputs. *Current Opinion in Neurobiology*, 14(4):481–487.
- Ravikumar, P., Wainwright, M. J., Raskutti, G., and Yu, B. (2011). High-dimensional covariance estimation by minimizing ℓ_1 -penalized log-determinant divergence. *Electron. J. Stat.*, 5:935–980.
- Reiss, P. and Ogden, R. (2010). Functional generalized linear models with images as predictors. *Biometrics*, 66:61–69.
- Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E., Johansen-Berg, H., Bannister, P. R., Luca, M. D., Drobnjak, I., Flitney, D. E., Niazy, R. K., Saunders, J., Vickers, J., Zhang, Y., Stefano, N. D., Brady, J. M., and Matthews, P. M. (2004). Advances in functional and structural {MR} image analysis and implementation as {FSL}. *NeuroImage*, 23, Supplement 1:S208 – S219. Mathematics in Brain Imaging.

- Xia, Y., Cai, T., and Cai, T. T. (2015). Testing differential networks with applications to the detection of gene-gene interactions. *Biometrika*, 102:247–266.
- Yin, J. and Li, H. (2012). Model selection and estimation in the matrix normal graphical model. *Journal of Multivariate Analysis*, 107:119 – 140.
- Yuan, M. (2010). High dimensional inverse covariance matrix estimation via linear programming. *J. Mach. Learn. Res.*, 11:2261–2286.
- Yuan, M. and Lin, Y. (2007). Model selection and estimation in the gaussian graphical model. *Biometrika*, 94(1):19–35.
- Zaitsev, A. Y. (1987). On the Gaussian approximation of convolutions under multidimensional analogues of S.N. Bernstein’s inequality conditions. *Probab. Theory Rel.*, 74(4):535–566.
- Zhang, X., Begleiter, H., Porjesz, B., Wang, W., and Litke, A. (1995). Event related potentials during object recognition tasks. *Brain Research Bulletin*, 38(6):531–538.
- Zhou, H. and Li, L. (2014). Regularized matrix regression. *Journal of the Royal Statistical Society. Series B*, 76:463–483.
- Zhou, S. (2014). Gemini: graph estimation with matrix variate normal instances. *Ann. Statist.*, 42(2):532–562.
- Zhu, Y., Shen, X., and Pan, W. (2014). Structural Pursuit Over Multiple Undirected Graphs. *J. Amer. Statist. Assoc.*, 109(508):1683–1696.