ON REAL HYPERSURFACES IN NON-FLAT COMPLEX SPACE FORMS WITH A CONDITION ON THE STRUCTURE JACOBI OPERATOR

S.H. Kon, Tee-How Loo, Shiquan Ren Institute of Mathematical Sciences, University of Malaya, 50603 Kuala Lumpur, Malaysia

Abstract. In this paper we prove some classification theorems of real hypersurfaces in $M_n(c)$ satisfying certain conditions on the covariant derivative of the structure Jacobi operator. We also prove the non-existence of real hypersurfaces with Codazzi type structure Jacobi operator in $M_n(c)$.

Mathematics Subject Classification(2010): 53C15 53B25

Keywords: complex space form, real hypersurface, Jacobi operator, totally η -umbilical real hypersurface

1 Introduction

The study of real hypersurfaces of Kaehlerian manifolds has been an important subject in the geometry of submanifolds, especially when the ambient space is a complex space form. Complex space forms are the simplest Kaehlerian manifolds, and they are the complex case analogues of real space forms. The complex structure of the ambient space induced an almost contact structure on its real hypersurfaces. The interaction between the almost contact structure and shape operator results in interesting properties of real hypersurfaces in complex space forms (for instance, see [7, 11, 13, 14, 15, 19]).

Let $M_n(c)$ be an *n*-dimensional non-flat complex space form with constant holomorphic sectional curvature 4*c*. It is known that a complete and simply connected non-flat complex space form is either a complex projective space (c > 0), denoted by CP^n , or a complex hyperbolic space (c < 0), denoted by CH^n . Throughout this paper, all manifolds, vector fields, etc., will be considered of class C^{∞} unless otherwise stated and it is always assumed $c = \pm 1$ and n > 2.

We also assume that M is a connected real hypersurface in $M_n(c)$, without boundary. Since we only study local properties, we do not distinguish between M and an open subset of M. Let N be a locally defined unit normal vector field on M. Denote by ∇ the Levi-Civita connection on M induced from $M_n(c)$. Let \langle , \rangle denote the Riemannian metric of M induced from the Riemannian metric of $M_n(c)$ and A the shape operator of M in $M_n(c)$. We define a tensor field ϕ of type (1,1), a vector field ξ and a 1-form η by

$$JX = \phi X + \eta(X)N, \quad JN = -\xi,$$

where $X \in \Gamma(TM)$ and $\eta(X) = \langle X, \xi \rangle$. Let D, called the holomorphic distribution, denote the distribution on M given by all vectors orthogonal to ξ at each point of M. Let $\alpha = \langle A\xi, \xi \rangle$. A real hypersurface M in $M_n(c)$ is said to be Hopf if ξ is principal, i.e., $A\xi = \alpha\xi$; and M is said to be totally η -umbilical if there exists a function λ such that $AX = \lambda X$ for all $X \in \Gamma(D)$. It has been proved that λ must be a constant (cf. [2], [10]). The following celebrated theorems classify certain families of well-behaved Hopf hypersurfaces in $\mathbb{C}P^n$ and $\mathbb{C}H^n$ respectively.

Theorem 1. ([5]) Let M be a Hopf hypersurface with constant principal curvatures in \mathbb{CP}^n . Then M is locally congruent to a tube of radius r over one of the following Keahlerian manifolds:

 $\begin{array}{l} (A_1) \ hyperplane \ CP^{n-1}, \ where \ 0 < r < \pi/2; \\ (A_2) \ totally \ geodesic \ CP^p \ (0 < p < n-1), \ where \ 0 < r < \pi/2; \\ (B) \ complex \ quadric \ Q_{n-1}, \ where \ 0 < r < \pi/4; \\ (C) \ CP^1 \times CP^{(n-1)/2}, \ where \ 0 < r < \pi/4 \ and \ n > 4 \ is \ odd; \\ (D) \ complex \ Grassmann \ G_{2,5}, \ where \ 0 < r < \pi/4 \ and \ n = 9; \\ (E) \ Hermitian \ symmetric \ space \ SO(10)/U(5), \ where \ 0 < r < \pi/4 \ and \ n = 15. \end{array}$

Remark 2. In CP^n , a real hypersurface is a geodesic hypersphere of radius r if and only if it is a tube of radius $\frac{\pi}{2} - r$ over CP^{n-1} , where $0 < r < \pi/2$.

Theorem 3. ([1]) Let M be a Hopf hypersurface with constant principal curvatures in CH^n . Then M is locally congruent to one of the following:

- (A_0) a horosphere;
- (A_1) a geodesic hypersphere CH^0 or a tube over a hyperplane CH^{n-1} ;
- (A_2) a tube over a totally geodesic CH^p 0 ;

(B) a tube over a totally real hyperbolic space RH^n .

In the following, by a real hypersurface of type A, we mean of type A_1 , A_2 (resp. of A_0 , A_1 , A_2) for c > 0 (resp. c < 0). By using the above theorems, it is possible to obtain various classification theorems under conditions on certain symmetric operators, such as the shape operator, Ricci tensor, and the structure Jacobi operator.

The Jacobi operator R_X with respect to a tangent vector field X on an open subset of M, is defined by $R_X(Y) = R(Y, X)X$, for $Y \in \Gamma(TM)$, where R is the curvature tensor on M. In particular, R_{ξ} is called the structure Jacobi operator of M. Various conditions on the structure Jacobi operator have been considered for either characterizing certain classes of real hypersurfaces or non-existence problems. The study of real hypersurfaces with parallel R_{ξ} was initiated in [16]. In [8], a characterization of ruled real hypersurfaces in $M_n(c)$ under the condition $(\nabla_X R_{\xi})\xi = 0$ for all $X \in \Gamma(D)$ was given, and the non-existence of real hypersurfaces in $M_n(c)$ with D-parallel and D-recurrent R_{ξ} can be deduced due to this result. In [4], real hypersurfaces in $M_n(c)$ with cyclic-parallel R_{ξ} have been studied; and in [17], real hypersurfaces in CP^n with Lie ξ -parallel R_{ξ} were considered. In [6] and [9], the following conditions for the shape operator A and the Ricci tensor S

$$(\nabla_X A)Y = -c\{\langle \phi X, Y \rangle \xi + \eta(Y)\phi X\},\$$

$$(\nabla_X S)Y = k\{\langle \phi X, Y \rangle \xi + \eta(Y)\phi X\}$$

have been studied respectively for real hypersurfaces in CP^n . In [3], these two conditions were considered for real hypersurfaces in CH^n . The study of such conditions was motivated by the non-existence of real hypersurfaces in $M_n(c)$ with either parallel shape operator or parallel Ricci tensor (cf. page 243 and page 271 of [12]). On the other hand, there does not exist any real hypersurfaces in $M_n(c)$ with parallel R_{ξ} as proved in [16]. It is natural to study a similar condition on R_{ξ} in the following theorem.

Theorem 4. Let M be a real hypersurface in $M_n(c)$. Then M satisfies

$$(\nabla_X R_{\xi})Y = k(\langle \phi X, Y \rangle \xi + \eta(Y)\phi X), \tag{1}$$

for all $X, Y \in \Gamma(TM)$ if and only if M is totally η -umbilical, i.e., it is locally congruent to one of the following real hypersurfaces:

for c > 0, (a). geodesic hyperspheres in CP^n ; for c < 0, (a). geodesic hyperspheres in CH^n ; (b). tubes around complex hyperbolic hyperplane in CH^n ; (c). horospheres in CH^n . Furthermore, we have $k \neq 0$.

Let F be a tensor field of type (1, 1) on M. F is said to be of Codazzi type if for any $X, Y \in \Gamma(TM)$,

$$(\nabla_X F)Y = (\nabla_Y F)X. \tag{2}$$

The condition (2) is weaker than the parallelism of F, and it is natural since for a totally geodesic hypersurface in a Riemannian manifold (if it exists), the shape operator is of Codazzi type. A Riemannian manifold is said to have harmonic curvature if its Ricci tensor is of Codazzi type. This cannot happen for a Hopf hypersurface in $M_n(c)$ (cf. [12, page 279]). Moreover, in [18], the non-existence of real hypersurfaces in \mathbb{CP}^n with Codazzi type structure Jacobi operator has been obtained. We will generalize this statement in Theorem 5.

Theorem 5. There does not exist any real hypersurface M in $M_n(c)$ with its structure Jacobi operator of Codazzi type.

In order to prove Theorem 4 and Theorem 5 simultaneously, we consider a generalized condition in Theorem 6.

Theorem 6. A real hypersurface M in $M_n(c)$ satisfies

$$\langle (\nabla_X R_{\xi})Y - (\nabla_Y R_{\xi})X, W \rangle = k(2\eta(W)\langle \phi X, Y \rangle + \eta(Y)\langle \phi X, W \rangle - \eta(X)\langle \phi Y, W \rangle)$$
(3)

for all $X, Y, W \in \Gamma(TM)$, if and only if M is a totally η -umbilical real hypersurface, or an arbitrary Hopf hypersurface with $\alpha = k = 1$, c = -1. Furthermore, we have $k \neq 0$.

Finally, Theorem 4 and Theorem 6 give two equivalent characterizations for totally η -umbilical real hypersurfaces in \mathbb{CP}^n , as stated in the following corollary.

Corollary 7. For a real hypersurface M in CP^n , the following conditions are equivalent:

- (a). (1) holds for all $X, Y, W \in \Gamma(TM)$;
- (b). (3) holds for all $X, Y \in \Gamma(TM)$;
- (c). M is totally η -umbilical.

2 Preliminaries

Let M be a connected real hypersurface in $M_n(c)$ without boundary. The set of tensors $(\phi, \xi, \eta, \langle, \rangle)$ is an almost contact metric structure on M, i.e., they satisfy the following

$$\phi^2 X = -X + \eta(X)\xi, \quad \phi\xi = 0, \quad \eta(\phi X) = 0, \quad \eta(\xi) = 1.$$

From the parallelism of J, we get

$$(\nabla_X \phi)Y = \eta(Y)AX - \langle AX, Y \rangle \xi,$$

 $\nabla_X \xi = \phi AX.$

Let R be the curvature tensor of M. Then the Gauss and Codazzi equations are respectively given by

$$\begin{split} R(X,Y)Z &= c\{\langle Y,Z\rangle X - \langle X,Z\rangle \ Y + \langle \phi Y,Z\rangle \phi X - \langle \phi X,Z\rangle \phi Y \\ &- 2\langle \phi X,Y\rangle \phi Z\} + \langle AY,Z\rangle AX - \langle AX,Z\rangle AY, \end{split}$$

$$(\nabla_X A)Y - (\nabla_Y A)X = c\{\eta(X)\phi Y - \eta(Y)\phi X - 2\langle\phi X, Y\rangle\xi\}$$

From the Gauss equation, we have

$$R_{\xi}Y = c\{Y - \eta(Y)\xi\} + \alpha AY - \eta(AY)A\xi, \qquad (4)$$

$$(\nabla_X R_{\xi})Y = -c\langle Y, \phi AX \rangle \xi - c\eta(Y)\phi AX + (X\alpha)AY + \alpha(\nabla_X A)Y -\langle (\nabla_X A)Y, \xi \rangle A\xi - \langle Y, A\phi AX \rangle A\xi -\eta(AY)(\nabla_X A)\xi - \eta(AY)A\phi AX,$$
(5)

for any $X, Y \in \Gamma(TM)$.

For a Hopf hypersurface M in $M_n(c)$, it can be proved that α is a constant (cf. [7]). The following cited results will be used in the proof of our theorems.

Theorem 8. ([12, page 262-264]) Let M be a real hypersurface in $M_n(c)$. Then M is locally congruent to a Hopf hypersurface of type A if and only if

$$(\nabla_X A)Y = -c\{\langle \phi X, Y \rangle \xi + \eta(Y)\phi X\}$$

for any $X, Y \in \Gamma(TM)$.

Theorem 9. ([2], [10]) Let M be a real hypersurface in $M_n(c)$. Then M is totally η -umbilical if and only if it is one of the following:

for c > 0,

(a). geodesic hyperspheres in CP^n ;

for c < 0,

(a). geodesic hyperspheres in CH^n ;

(b). tubes around complex hyperbolic hyperplane in CH^n ;

(c). horospheres in CH^n .

The corresponding principal curvatures of the real hypersurfaces in Theorem 9 are in Table 1 (λ is the principal curvature such that $AX = \lambda X$ for $X \in \Gamma(D)$).

Table 1:				
	Case	Radius	α	λ
c > 0	(a)	r	$2\cot 2r$	$\cot r$
$\begin{array}{c} c < 0 \\ c < 0 \end{array}$	(a)	r	$2 \coth 2r$	$\coth r$
c < 0	(b)	r	$2 \coth 2r$	$\tanh r$
c < 0	(c)	-	2	1

Let $\beta = ||\phi A\xi||$. If M is a non-Hopf real hypersurface in $M_n(c)$ then $\beta > 0$. We can define a unit vector field U in $\Gamma(D)$ by $U = -\frac{1}{\beta}\phi^2 A\xi$ and a distribution D_U by

$$D_U = \{ X \in T_x M | X \perp \xi, U, \phi U \}, x \in M.$$

Lemma 10. ([8]) Let M be a non-Hopf real hypersurface in $M_n(c)$. Suppose M satisfies the following:

(a) $A\xi = \alpha \xi + \beta U$, $AU = \beta \xi + \gamma U$ and $A\phi U = \delta \phi U$ for some functions γ and δ on M;

(b) there exists a unit vector field $Z \in \Gamma(D_U)$ such that $AZ = \lambda Z$ and $A\phi Z = \lambda \phi Z$ for some function λ on M.

Then we have

$$\beta\lambda(\lambda-\delta) - (\lambda-\gamma)\phi U\lambda = 0.$$
(6)

Proof. Suppose M is such a real hypersurface satisfying (a) and (b). Taking inner product in the Codazzi equation

$$(\nabla_Z A)\xi - (\nabla_\xi A)Z = -c\phi Z$$

with ϕZ , we obtain

$$\beta \langle \nabla_Z U, \phi Z \rangle = \lambda^2 - \alpha \lambda - c.$$

Taking inner product in the Codazzi equation

$$(\nabla_Z A)\phi U - (\nabla_{\phi U} A)Z = 0$$

with Z, we obtain

$$(\delta - \lambda) \langle \nabla_Z \phi U, Z \rangle = \phi U \lambda.$$

By using

$$\nabla_Z \phi U = \phi \nabla_Z U,$$

we have

$$(\lambda - \delta)(\lambda^2 - \alpha \lambda - c) = \beta \phi U \lambda.$$
(7)

Taking inner product in the Codazzi equation $(\nabla_Z A)\phi Z - (\nabla_{\phi Z} A)Z = -2c\xi$ with ξ and U respectively, we obtain

$$\langle \nabla_{\phi Z} Z - \nabla_Z \phi Z, U \rangle = \frac{2(\lambda^2 - \alpha \lambda - c)}{\beta},$$
$$(\lambda - \gamma)(\langle \nabla_{\phi Z} Z, U \rangle - \langle \nabla_Z \phi Z, U \rangle) = 2\beta\lambda.$$

Combining these two equations, we obtain

$$(\lambda - \gamma)(\lambda^2 - \alpha\lambda - c) - \beta^2\lambda = 0.$$
(8)

From (7) and (8), we get (6).

Lemma 11. ([7]) Let M be a Hopf hypersurface in $M_n(c)$ with $A\xi = \alpha \xi$. Then

$$2A\phi A - \alpha(A\phi + \phi A) - 2c\phi = 0.$$

3 A Lemma

In this section, We shall obtain a lemma that is useful for the proof of the non-Hopf case of our theorems. The generalized condition in this lemma enables it to be applied in a number of problems.

Lemma 12. Let M be a non-Hopf real hypersurface in $M_n(c)$. Suppose M satisfies $A\xi = \epsilon c\xi + \beta U$, $AU = \beta \xi + \epsilon (\beta^2 - c)U$, $A\phi U = -\epsilon c\phi U$, where U is a unit vector field in D, β is a nonvanishing function defined on M and $\epsilon = \pm 1$. Then c > 0 and there exists a vector field $X \in \Gamma(D_U)$ such that $AX \neq -\epsilon X$. *Proof.* Suppose M is such a real hypersurface. Taking inner product in the Codazzi equation $(\nabla_U A)\phi U - (\nabla_{\phi U} A)U = -2c\xi$ with U and ξ respectively, we obtain

$$-\beta \langle \nabla_U \phi U, U \rangle + \beta^2 - 3c - 2\phi U\beta = 0, \qquad (9)$$

$$-\beta \langle \nabla_U \phi U, U \rangle + 3c\beta^2 - 4c^2 + 2c - \phi U\beta = 0.$$

From these two equations, we obtain

$$\beta^2 - 3c\beta^2 + 4c^2 - 5c - \phi U\beta = 0.$$
⁽¹⁰⁾

Taking inner product in the Codazzi equation

$$(\nabla_{\phi U}A)\xi - (\nabla_{\xi}A)\phi U = cU \tag{11}$$

with $U, \xi, \phi U$ respectively, we obtain

$$\epsilon\beta^2 \langle \nabla_{\xi} \phi U, U \rangle + 2c^2 - c\beta^2 - \beta^2 - c + \phi U\beta = 0, \qquad (12)$$

$$\langle \nabla_{\xi} \phi U, U \rangle - 4\epsilon c = 0, \tag{13}$$

$$\langle \nabla_{\phi U} U, \phi U \rangle = 0. \tag{14}$$

From (12) and (13), we obtain

$$3c\beta^2 + 2c^2 - \beta^2 - c + \phi U\beta = 0.$$
(15)

By summing up (10) and (15), we obtain

$$c(c-1) = 0,$$

which cannot happen when c = -1. Hence c = 1 and (10) becomes

$$\phi U\beta = -2\beta^2 - 1. \tag{16}$$

By substituting (16) into (9), we have

$$\beta \langle \nabla_U \phi U, U \rangle = 5\beta^2 - 1. \tag{17}$$

From (13), we have

$$\langle \nabla_{\xi} \phi U, U \rangle = 4\epsilon. \tag{18}$$

In order to prove the second assertion, we shall suppose to the contrary that for any $Z \in \Gamma(D_U)$, $AZ = -\epsilon Z$. Take inner product in the Codazzi equation $(\nabla_Z A)\xi - (\nabla_\xi A)Z = -\phi Z$ with $U, \phi U, \xi$ respectively,

$$Z\beta + \epsilon\beta^2 \langle \nabla_{\xi} Z, U \rangle = 0, \tag{19}$$

$$\langle \nabla_Z U, \phi U \rangle = 0, \tag{20}$$

$$\langle \nabla_{\xi} Z, U \rangle = 0. \tag{21}$$

(19) and (21) imply $Z\beta = 0$. Taking inner product in the Codazzi equation $(\nabla_Z A)U - (\nabla_U A)Z = 0$ with U and with the help of $Z\beta = 0$, we have

$$\langle \nabla_U Z, U \rangle = 0. \tag{22}$$

Taking inner product with Z in (11), we have

$$\langle \nabla_{\phi U} U, Z \rangle = 0. \tag{23}$$

From (14), (23) and $\langle \nabla_{\phi U} U, \xi \rangle = -\epsilon$, we obtain

$$\nabla_{\phi U} U = -\epsilon \xi. \tag{24}$$

Hence

$$\nabla_{\phi U}\phi U = (\nabla_{\phi U}\phi)U + \phi\nabla_{\phi U}U = 0.$$
(25)

From (17), (22) and $\langle \nabla_U U, \xi \rangle = 0$, we obtain

$$\nabla_U U = \frac{1 - 5\beta^2}{\beta} \phi U. \tag{26}$$

Hence

$$\nabla_U \phi U = \epsilon (1 - \beta^2) \xi + \frac{5\beta^2 - 1}{\beta} U.$$
(27)

From (18), (21) and $\langle \nabla_{\xi} U, \xi \rangle = 0$, we obtain

$$\nabla_{\xi} U = -4\epsilon \phi U. \tag{28}$$

Finally, we also have

$$\nabla_{\phi U}\xi = \phi A\phi U = \epsilon U. \tag{29}$$

Let X = U, $Y = \phi U$ and Z = U in the Gauss equation. Then we have

$$R(U,\phi U)U = (\beta^2 - 5)\phi U.$$
 (30)

On the other hand, it follows from (16), (24)–(29) and

$$R(U,\phi U)U = \nabla_U \nabla_{\phi U} U - \nabla_{\phi U} \nabla_U U - \nabla_{[U,\phi U]} U$$

that

$$R(U,\phi U)U = (10\beta^2 - 8)\phi U.$$
(31)

From (30) and (31), we see that β is a constant. This contradicts (16).

4 Auxiliary propositions

In this section we mainly focus on the Hopf case and prove some propositions about Hopf hypersurfaces in preparation for the proof of our theorems.

Proposition 13. Let M be a real hypersurface in $M_n(c)$. If M is totally η -umbilical then M satisfies (1). Furthermore, $k \neq 0$.

Proof. For a totally η -umbilical real hypersurface M, we have

$$AX = \lambda X + (\alpha - \lambda)\eta(X)\xi \tag{32}$$

for any $X \in \Gamma(TM)$, where λ and α are two constants. Applying (32) and Theorem 8 to (5), we have

$$(\nabla_X R_{\xi})Y = -\lambda(c + \alpha\lambda)\{\langle Y, \phi X \rangle \xi + \eta(Y)\phi X\}.$$
(33)

Hence *M* satisfies (1). From Lemma 11, we have $\lambda^2 = \alpha \lambda + c$, hence $\lambda \neq 0$. Then from the right-hand side of (33), we see that $k = -\lambda(c + \alpha \lambda) = -\lambda^3 \neq 0$.

Proposition 14. Let M be a Hopf hypersurface in CH^n with $\alpha = 1$. Then M satisfies (3) for k = 1 and does not satisfy (1) for k = 1.

Proof. We suppose k = 1. Then (5) reduces to

$$(\nabla_X R_{\xi})Y = \langle Y, \phi AX \rangle \xi + \eta(Y)\phi AX + (\nabla_X A)Y - \langle Y, (\nabla_X A)\xi \rangle \xi - \langle Y, A\phi AX \rangle \xi - \eta(Y)(\nabla_X A)\xi - \eta(Y)A\phi AX.$$
(34)

By a direct computation, we have $(\nabla_X A)\xi = \phi AX - A\phi AX$. Substituting this equation into (34), we obtain

$$(\nabla_X R_\xi) Y = (\nabla_X A) Y. \tag{35}$$

By using (35) and the Codazzi equation, we see that M satisfies (3) for k = 1.

Next, suppose there exists a Hopf hypersurface M satisfying (1) for k = 1. Then by (35), (1) becomes

$$(\nabla_X A)Y = \langle \phi X, Y \rangle \xi + \eta(Y)\phi X. \tag{36}$$

By Theorem 8, M is locally congruent to a Hopf hypersurface of type A with $\alpha = 1$. According to [12, page 254-257, 260], $\alpha \neq 1$ for real hypersurfaces of type A. This is a contradiction. Hence we conclude that M does not satisfy (1) for k = 1.

Let M be a real hypersurface in $M_n(c)$ satisfying (3). Then by applying (5) and the Codazzi equation, (3) becomes

$$0 = \langle (\alpha c + k)(\eta(X)\phi Y - \eta(Y)\phi X - 2\langle \phi X, Y\rangle \xi) + 2\langle AX, \phi AY\rangle A\xi + c(\langle X, (A\phi + \phi A)Y\rangle \xi + 2\langle \phi X, Y\rangle A\xi - \eta(Y)\phi AX + \eta(X)\phi AY) + (X\alpha)AY - (Y\alpha)AX + \eta(AX)(\nabla_Y A)\xi - \eta(AY)(\nabla_X A)\xi + \eta(AX)A\phi AY - \eta(AY)A\phi AX, W\rangle$$
(37)

for any $X, Y, W \in \Gamma(TM)$.

Proposition 15. Let M be a Hopf hypersurface in $M_n(c)$ satisfying (3). Then either

(1). M is a totally η-umbilical real hypersurface;
or
(2). α = k = 1, c = -1.

Proof. Recall that α is a constant when M is Hopf. Let $X = \xi$, $Y, W \in \Gamma(D)$ in (37), we have

$$(\alpha c + k)\langle \phi Y, W \rangle + (\alpha^2 + c)\langle \phi AY, W \rangle = 0.$$
(38)

Let $Y \in \Gamma(D)$ be a unit principal vector field with $AY = \lambda Y$ and let $W = \phi Y$ in (38). Then

$$(\alpha^2 + c)\lambda + (\alpha c + k) = 0.$$
(39)

We consider two cases: $\alpha^2 + c \neq 0$, $\alpha^2 + c = 0$.

Case-I. $\alpha^2 + c \neq 0$.

Since Y is an arbitrary unit principal vector field in $\Gamma(D)$, we have $AX = \lambda X$ for all $X \in \Gamma(D)$, where

$$\lambda = -\frac{\alpha c + k}{\alpha^2 + c}.$$

Therefore, M is totally η -umbilical.

Case-II. $\alpha^2 + c = 0.$

Then c = -1, and by replacing the unit normal vector field N with -N if necessary, we have $\alpha = 1$. By (39), we have $k = \alpha = 1$.

5 Proof of the theorems

In this section we first prove the non-existence of non-Hopf real hypersurface satisfying condition (1) or (3), then prove the theorems.

Let M be a real hypersurface in $M_n(c)$. Letting $W = \xi$ in (37), we have

$$2k\langle\phi X, Y\rangle + 2\alpha\langle A\phi AX, Y\rangle + c\langle(\phi A + A\phi)X, Y\rangle -\eta(AY)\eta(A\phi AX) + \eta(AX)\eta(A\phi AY) = 0$$
(40)

for any $X, Y \in \Gamma(TM)$. From (40), we have

$$2k\phi X + 2\alpha A\phi AX + c(\phi A + A\phi)X - \eta(A\phi AX)A\xi - \eta(AX)A\phi A\xi = 0$$
(41)

for any $X \in \Gamma(TM)$.

Proposition 16. There does not exist any non-Hopf real hypersurface M in $M_n(c)$ satisfying (3).

Proof. Suppose M is a non-Hopf real hypersurface satisfying (3) with $A\xi = \alpha\xi + \beta U$, β a non-vanishing function and $U \in \Gamma(D)$ a unit vector field.

Let $X = \xi$ in (41). Then we have $\alpha \neq 0$ and

$$A\phi U = -\frac{c}{\alpha}\phi U. \tag{42}$$

Let $X = \phi U$ in (41). Then with the help of (42) we obtain

$$AU = \beta\xi + \left(\frac{2k}{c} + \frac{\beta^2 - c}{\alpha}\right)U.$$
(43)

Hence D_U is A-invariant.

Let $X = \phi U, Y \in \Gamma(D_U), W \in \Gamma(TM)$ in (37). Then we have

$$(\phi U\alpha)AY + \frac{c}{\alpha}(Y\alpha)\phi U = 0.$$
(44)

Taking inner product with ϕU in (44), we have $Y\alpha = 0$.

Let $X = U, Y \in \Gamma(D_U), W \in \Gamma(TM)$ in (37). Then with the help of $Y\alpha = 0$, we have

$$(U\alpha)AY + \beta(\nabla_Y A)\xi + \beta A\phi AY = 0.$$
(45)

Let $X = \xi$, $Y \in \Gamma(D_U)$, $W \in \Gamma(TM)$ in (37). Then by the fact that $Y\alpha = 0$, we have

$$(\xi\alpha)AY + c\phi AY + (\alpha c + k)\phi Y + \alpha(\nabla_Y A)\xi + \alpha A\phi AY = 0.$$
(46)

From (45) and (46), we have

$$(\xi \alpha - \frac{\alpha}{\beta} U \alpha) AY + c \phi AY + (\alpha c + k) \phi Y = 0.$$
(47)

Let Y be unit vector field with $AY = \lambda Y$ and take inner product with ϕY in (47). Then we obtain

$$c\lambda + \alpha c + k = 0. \tag{48}$$

Therefore,

$$AX = \lambda X, (\lambda = -\frac{\alpha c + k}{c})$$
(49)

for any $X \in \Gamma(D_U)$. On the other hand, if we put X = Y in (41), then we have

$$\alpha \lambda^2 + c\lambda + k = 0. \tag{50}$$

From (48) and (50) we have

$$\lambda^2 = c. \tag{51}$$

Hence c = 1 and $\lambda = \pm 1$. It follows that $\phi U \lambda = 0$. So by Lemma 10, (42), (50) and (51), we obtain $\alpha = -\lambda (= \mp 1)$ and k = 0. By substituting all these quantities into (42), (43) and (49), we obtain a contradiction according to Lemma 12. This completes the proof.

Proof of Theorem 6. (\Leftarrow). It follows directly from Proposition 13 and Proposition 14.

(⇒). Suppose M satisfies (3). By Proposition 15 and Proposition 16, M is either totally η -umbilical or a Hopf hypersurface with $\alpha = k = 1, c = -1$. Furthermore, by Proposition 13, we can see that $k \neq 0$.

Proof of Theorem 4. (\Leftarrow). It has been proved in Proposition 13.

(⇒). Suppose M is a real hypersurface satisfying (1). Then M also satisfies (3). Hence from Theorem 6, M is either totally η -umbilical or a Hopf hypersurface with $\alpha = k = 1, c = -1$. However, by Proposition 14, the latter case cannot occur. We conclude that M is totally η -umbilical, so it is locally congruent to one of the real hypersurfaces listed in Theorem 9.

Proof of Theorem 5. Suppose the Jacobi operator is of Codazzi type. Then M satisfies (3) with k = 0. This contradicts the fact that $k \neq 0$ as stated in Theorem 6. The proof is completed.

References

- J. Berndt, Real hypersurfaces with constant principal curvatures in complex hyperbolic space, J. Reine Angew. Math. 395 (1989), 132-141.
- [2] T.E. Cecil and P.J. Ryan, Focal sets and real hypersurfaces in complex projective space, *Trans. Amer. Math. Soc.* 269 (1982), 481-499.
- [3] Y.W. Choe, Characterizations of certain real hypersurfaces of a complex space form, *Nihonkai Math. J.* 6 (1995), 97-114.
- [4] U.H. Ki and H. Kurihara, Real hypersurfaces with cyclic-parallel structure Jacobi operators in a non-flat complex space form, *Bull. Aust. Math. Soc.* 81 (2010), 260-273.
- [5] M. Kimura, Real hypersurfaces and complex submanifolds in complex projective space, *Trans. Amer. Math. Soc.* 296:1 (1986), 137-149.
- [6] M. Kimura and S. Maeda, Characterizations of geodesic hyperspheres in a complex projective space in terms of Ricci tensors, *Yokohama Math. J.* 40 (1992), 35-43.
- [7] M. Kon, Pseudo-Einstein real hypersurfaces in complex space forms, J. Differential Geom. 14 (1979), 339-354.
- [8] S.H. Kon, T.H. Loo and S. Ren, Real hypersurfaces in a complex space form with a condition on the structure Jacobi operator. *Math. Slovaca*, to appear.
- [9] Y. Maeda, On real hypersurfaces of a complex projective space, J. Math. Soc. Japan 28:3 (1976), 529-540.
- [10] S. Montiel, Real hypersurfaces of a complex hyperbolic space, J. Math. Soc. Japan 37:3 (1985), 515-535.
- [11] S. Montiel and A. Romero, On some real hypersurfaces of a complex hyperbolic space, *Geom. Dedicata* **20** (1986), 245-261.
- [12] R. Niebergall and P.J. Ryan, Real hypersurfaces in complex space forms, *Tight and taut submanifolds MSRI Publication* **32** (1997), 233-304.
- [13] M. Okumura, Certain almost contact hypersurfaces in Euclidean spaces, *Kodai Math. Sem. Rep.* 16 (1964), 44-54.
- [14] M. Okumura, Contact hypersurfaces in certain Kaehlerian manifolds, Tohoku Math. J. 18 (1966), 74-102.

- [15] M. Okumura, On some real hypersurfaces of a complex projective space, *Trans. Amer. Math. Soc.* **212** (1975), 355-364.
- [16] M. Ortega, J.D. Pérez and F.G. Santos, Non-existence of real hypersurfaces with parallel structure Jacobi operator in non-flat complex space forms. *Rocky Mountain J. Math.* 36 (2006), 1603-1613.
- [17] J.D. Pérez, F.G. Santos and Y.J. Suh, Real hypersurfaces in complex projective space whose structure Jacobi operator is Lie ξ -parallel, *Differential Geom. Appl.* **22** (2005), 181-188.
- [18] J.D. Pérez, F.G. Santos and Y.J. Suh, Real hypersurfaces in complex projective space whose structure Jacobi operator is of Codazzi type. *Canad. Math. Bull.* **50:3** (2007), 347-355.
- [19] M.H. Vernon, Contact hypersurfaces of a complex hyperbolic space, *Tohoku Math. J.* **39** (1987), 215-222.

Email:

S.H.Kon: shkon@um.edu.my, Tee-How Loo: looth@um.edu.my, Shiquan Ren: renshiquan@gmail.com