

GROWTH RATE OF ENDOMORPHISMS OF HOUGHTON'S GROUPS

JONG BUM LEE AND SANG RAE LEE

ABSTRACT. A Houghton's group $\mathcal{H}_n$ consists of translations at infinity of a n rays of discrete points on the plane. In this paper we study the growth rate of endomorphisms of Houghton's groups. We show that if the kernel of an endomorphism ϕ is not trivial then the growth rate $\text{GR}(\phi)$ equals either 1 or the spectral radius of the induced map on the abelianization. It turns out that every monomorphism ϕ of $\mathcal{H}_n$ determines a unique natural number ℓ such that $\phi(\mathcal{H}_n)$ is generated by translations with the same translation length ℓ . We use this to show that $\text{GR}(\phi)$ of a monomorphism ϕ of $\mathcal{H}_n$ is precisely ℓ for all $2 \leq n$.

1. INTRODUCTION

Let G be a finitely generated group with a generating set A . Let $\phi : G \rightarrow G$ be an endomorphism. For any $g \in G$, let $|g|$ denote the *word length* of g , that is, the minimum length of a word over $A \cup A^{-1}$ which represents g . Then the **growth rate** of ϕ is defined to be ([1])

$$\text{GR}(\phi) := \sup \left\{ \limsup_{k \rightarrow \infty} (|\phi^k(g)|)^{1/k} : g \in G \right\}.$$

The growth rate is well-defined, i.e., independent of the choice of a set of generators ([12, p.114]).

The problem of determining the growth rate of a group endomorphism was initiated by R. Bowen in ([1]). The growth rate of an endomorphism is related to algebraic entropy and topological entropy ([1], [12]). Algebraic entropy of ϕ is defined by $h_{\text{alg}}(\phi) = \log \text{GR}(\phi)$. Note that $h_{\text{alg}}(\phi)$ provides a lower bound for the topological entropy of a continuous self map f on a compact connected manifold M which induces the endomorphism ϕ of $\pi_1(M)$.

A group theoretic approach is discussed in [7] including the result that $\text{GR}(\phi)$ is finite and bounded by the maximum length of the image of a generator. In case ϕ is an automorphism of a nilpotent group it is shown in [13] that $\text{GR}(\phi)$ coincides with the growth rate of the induced automorphism on its abelianization. In [8], the first author extends this to all endomorphisms of nilpotent groups. In the same article it was proven that $\text{GR}(\phi)$ is an algebraic number if ϕ is an endomorphism of a torsion-free nilpotent or lattices of Sol.

For $n \in \mathbb{N}$, a Houghton's group $\mathcal{H}_n$ consists of *eventual translations* on a disjoint union of n copies of $\mathbb{N}$, each arranged along a ray emanating from the origin in the plane. The group $\mathcal{H}_2$ is finitely generated but not finitely presented. In general, by the work of K. Brown ([2]), $\mathcal{H}_n$ has finiteness type F_{n-1} but not F_n . For each n , $\mathcal{H}_n$ fits into the short exact sequence.

$$1 \longrightarrow \text{FSym}_n \longrightarrow \mathcal{H}_n \longrightarrow \mathbb{Z}^{n-1} \longrightarrow 1.$$

where FSym_n consists of permutations on the underlying set with finite supports. Note that every $\mathcal{H}_n$ contains all finite groups. Our main result is the following.

Theorem 1.1. *Suppose ϕ is an endomorphism of $\mathcal{H}_n$, $2 \leq n$. Every monomorphism ϕ of $\mathcal{H}_n$ determines $\ell \in \mathbb{N}$ such that $\text{GR}(\phi) = \ell$. If ϕ is not a monomorphism then $\text{GR}(\phi)$ equals either 1 or the spectral radius of the induced map on the abelianization.*

Date: February 14, 2022.

Key words and phrases. Houghton's groups, Growth rate of endomorphisms.

In Section 2 we define Houghton's groups and review basic facts including explicit presentations. In Section 3 we deal with automorphisms of $\mathcal{H}_n$. We calculate $\text{GR}(\phi)$ by using the structure of $\text{Aut}(\mathcal{H}_n)$ which is well understood. If ϕ is an endomorphism with non trivial kernel we can reduce the calculation of $\text{GR}(\phi)$ to the problem of growth rate of endomorphisms on finitely generated free abelian groups. In Section 4 we use the result of [8] to prove the second part of our main theorem above. The rest of paper is used to understand monomorphisms of $\mathcal{H}_n$. For $3 \leq n$, $\mathcal{H}_n$ is generated by $g_2, \dots, g_n$ where g_i translates points on the first ray toward i^{th} ray by 1. We use the relations of $\mathcal{H}_n$ together with the ray structure of the underlying set to characterize the behavior of monomorphisms in several steps carefully. It turns out that each monomorphism ϕ determines $\ell \in \mathbb{N}$ such that ϕ maps each generator g_i of $\mathcal{H}_n$ to a translation of length ℓ . This character of a monomorphism is essential for us to understand iterations ϕ^k applied to an element f with a finite support, which is the main obstruction in calculating $\text{GR}(\phi)$. Perhaps Houghton's groups are first examples of groups where relations were used extensively to calculate the growth rate of endomorphisms.

The following provides an effective calculation for $\text{GR}(\phi)$ ([7]), which will be used throughout the paper. For an endomorphism $\phi : G \rightarrow G$, let $|\phi^k|$ denote the maximum of $|\phi^k(a_i)|$ over a generating set $A = \{a_1, \dots, a_m\}$ of G , then

$$\text{GR}(\phi) = \lim_{k \rightarrow \infty} (|\phi^k|)^{1/k} = \inf_k \left\{ (|\phi^k|)^{1/k} \right\}.$$

2. HOUGHTON'S GROUPS $\mathcal{H}_n$

Let us use the following notational conventions. All bijections (or permutations) act on the right unless otherwise specified. Consequently gh means g followed by h . The conjugation by g is denoted by $\mu(g)$, $h^g = g^{-1}hg =: \mu(g)(h)$, and the commutator is defined by $[g, h] = ghg^{-1}h^{-1}$.

Our basic references are [9, 14] for Houghton's groups and [3] for their automorphism groups. Fix an integer $n \geq 1$. For each k with $1 \leq k \leq n$, let

$$R_k = \left\{ me^{i\theta} \in \mathbb{C} \mid m \in \mathbb{N}, \theta = \frac{\pi}{2} + (k-1)\frac{2\pi}{n} \right\}$$

and let $X_n = \bigcup_{k=1}^n R_k$ be the disjoint union of n copies of $\mathbb{N}$, each arranged along a ray emanating from the origin in the plane. We shall use the notation $\{1, \dots, n\} \times \mathbb{N}$ for X_n , letting (k, p) denote the point of R_k with distance p from the origin.

A bijection $g : X_n \rightarrow X_n$ is called an *eventual translation* if the following holds:

There exist an n -tuple $(m_1, \dots, m_n) \in \mathbb{Z}^n$ and a finite set $K_g \subset X_n$ such that

$$(2-1) \quad (k, p) \cdot g := (k, p + m_k) \quad \forall (k, p) \in X_n \setminus K_g.$$

An eventual translation acts as a translation on each ray outside a finite set. For each $n \in \mathbb{N}$ the *Houghton's group* $\mathcal{H}_n$ is defined to be the group of all eventual translations of X_n .

Let g_i be the translation on the ray of $R_1 \cup R_i$ by 1 for $2 \leq i \leq n$. Namely,

$$(j, p) \cdot g_i = \begin{cases} (1, p-1) & \text{if } j = 1 \text{ and } p \geq 2, \\ (i, 1) & \text{if } (j, p) = (1, 1), \\ (i, p+1) & \text{if } j = i, \\ (j, p) & \text{otherwise.} \end{cases}$$

Johnson provided a finite presentation for $\mathcal{H}_3$ in [11] and the second author gave a finite presentation for $\mathcal{H}_n$ with $n \geq 3$ in [14] as follows:

Theorem 2.1 ([14, Theorem C]). *For $n \geq 3$, $\mathcal{H}_n$ is generated by $g_2, \dots, g_n, \alpha$ with relations*

$$\alpha^2 = 1, (\alpha \alpha^{g_2})^3 = 1, [\alpha, \alpha^{g_2^2}] = 1, \alpha = [g_i, g_j], \alpha^{g_i^{-1}} = \alpha^{g_j^{-1}}$$

for $2 \leq i \neq j \leq n$.

From the definition of Houghton's groups, the assignment $g \in \mathcal{H}_n \mapsto (m_1, \dots, m_n) \in \mathbb{Z}^n$ defines a homomorphism $\pi = (\pi_1, \dots, \pi_n) : \mathcal{H}_n \rightarrow \mathbb{Z}^n$. Then we have:

Lemma 2.2 ([14, Lemma 2.3]). *For $n \geq 3$, we have $\ker \pi = [\mathcal{H}_n, \mathcal{H}_n]$.*

Note that $\pi(g_i) \in \mathbb{Z}^n$ has only two nonzero values -1 and 1 ,

$$\pi(g_i) = (-1, 0, \dots, 0, 1, 0, \dots, 0)$$

where 1 occurs in the i^{th} component. Since the image of $\mathcal{H}_n$ under π is generated by those elements, we have that

$$\pi(\mathcal{H}_n) = \left\{ (m_1, \dots, m_n) \in \mathbb{Z}^n \mid \sum_{i=1}^n m_i = 0 \right\},$$

which is isomorphic to the free Abelian group of rank $n - 1$. Consequently, $\mathcal{H}_n$ ($n \geq 3$) fits in the following short exact sequence

$$1 \longrightarrow \mathcal{H}'_n = [\mathcal{H}_n, \mathcal{H}_n] \longrightarrow \mathcal{H}_n \xrightarrow{\pi} \mathbb{Z}^{n-1} \longrightarrow 1.$$

The above abelianization, first observed by C. H. Houghton in [9], is the characteristic property of $\{\mathcal{H}_n\}$ for which he introduced those groups in the same paper. We may regard π as a homomorphism $\mathcal{H}_n \rightarrow \mathbb{Z}^n \rightarrow \mathbb{Z}^{n-1}$ given by

$$\pi : g_i \mapsto (-1, 0, \dots, 0, 1, 0, \dots, 0) \mapsto (0, \dots, 0, 1, 0, \dots, 0).$$

In particular, $\pi(g_2), \dots, \pi(g_n)$ form a set of free generators for $\mathbb{Z}^{n-1}$.

As definition, $\mathcal{H}_1$ is the symmetric group itself on X_1 with finite support, which is not finitely generated. Furthermore, $\mathcal{H}_2$ is

$$\mathcal{H}_2 = \langle g_2, \alpha \mid \alpha^2 = 1, (\alpha \alpha^{g_2})^3 = 1, [\alpha, \alpha^{g_2^k}] = 1 \text{ for all } |k| > 1 \rangle,$$

which is finitely generated, but not finitely presented. It is not difficult to see that $\mathcal{H}'_2 = \text{FAlt}_2$.

Notation.

- $\text{Sym}_n =$ the full symmetric group of X_n ,
- $\text{FSym}_n =$ the symmetric group of X_n with finite support,
- $\text{FAlt}_n =$ the alternating group of X_n with finite support.

Theorem 2.3 ([3, Theorem 2.2]). *For $n \geq 2$, we have*

$$\text{Aut}(\mathcal{H}_n) \cong \mathcal{H}_n \rtimes \Sigma_n$$

where Σ_n is the symmetric group that permutes n rays isometrically.

Here are some other known results for the Houghton's groups $\mathcal{H}_n$:

- K. S. Brown([2]) showed that $\mathcal{H}_n$ has type F_{n-1} but not F_n .
- By [14, Theorem 2.18], $\mathcal{H}_n$ is an amenable group.
- Röver [15] showed that for all $n \geq 1, r \geq 2, m \geq 1$, $\mathcal{H}_n$ embeds in Higman's groups $G_{r,m}$ (defined in [10]), and in particular all Houghton's groups are subgroups of Thompson's group V .

3. GROWTH RATES FOR AUTOMORPHISMS OF $\mathcal{H}_n$

In this section, we will study the growth rates for automorphisms of the Houghton's groups $\mathcal{H}_n$. We refer to [1, 8] for general information on growth rates for endomorphisms of finitely generated groups. Since $\mathcal{H}_1 = \text{FSym}_1$ is not finitely generated, we must assume $n \geq 2$.

For $n \geq 3$, we fix a generating set $\Gamma = \{g_2, \dots, g_n\}$ for $\mathcal{H}_n$. Let $|g|$ denote the word length $|g|_\Gamma$ of $g \in \mathcal{H}_n$ with respect to Γ . With $g_1 = \text{id}$, for any automorphism $\phi = \mu(\sigma\gamma)$ we have

$$\begin{aligned} g_i^\sigma &= g_{\sigma(1)}^{-1} g_{\sigma(i)} \text{ for } i = 2, \dots, n, \\ g_i^\gamma &= g_i^* = \text{an eventual } g_i. \end{aligned}$$

The second line follows from the fact that γ is a word of $g_j^{\pm 1}$'s and an observation that $g_j^{-1} g_i g_j$ is the translation on $R_1 \cup \{(1)\} \cup R_i$ by 1. Observe also that

- (1) $(g_i^*)^{-1} = (g_i^{-1})^*$, $(g_i^{\pm 1} g_j)^* = (g_i^*)^{\pm 1} g_j^*$,
- (2) $(g_i^*)^\sigma = (g_i^\sigma)^*$, $(g_i^*)^\gamma = (g_i^\gamma)^* = g_i^*$.

In particular, we have $\phi(g_i) = \phi(g_i^*) = \phi(g_i)^*$. Thus we have

$$\phi^m(g_i^*) = g_{\sigma^m(1)}^* g_{\sigma^m(i)}^*.$$

This shows that if σ is of order m , then $\phi^m(g_i) = \phi^m(g_i^*) = g_i^*$. Remark also that if $\gamma = 1$, then $\phi^m(g_i) = g_i$ for all $2 \leq i \leq n$, i.e., $\phi^m = \text{id}$. Hence $\text{GR}(\phi^m) = 1$ and so $\text{GR}(\phi) = 1$. In fact, we have:

Theorem 3.1. *For any automorphism ϕ of $\mathcal{H}_n$ ($n \geq 3$), we have $\text{GR}(\phi) = 1$.*

Proof. Let $|\gamma| = \ell$. Then we have $|\mu(\gamma)^k| \leq 1 + 2\ell k$. Indeed, since $\mu(\gamma)^k(g_i) = \gamma^{-k} g_i \gamma^k$, we have

$$\begin{aligned} |\mu(\gamma)^k(g_i)| &\leq |\gamma^{-k}| + |g_i| + |\gamma^k| \\ &= 1 + 2|\gamma^k| \leq 1 + 2k|\gamma| = 1 + 2k\ell. \end{aligned}$$

Thus we have $|\mu(\gamma)^k| \leq 1 + 2k\ell$. This implies that

$$\text{GR}(\mu(\gamma)) = \lim_{k \rightarrow \infty} |\mu(\gamma)^k|^{1/k} \leq \lim_{k \rightarrow \infty} (1 + 2k\ell)^{1/k} = 1.$$

Let $\phi = \mu(\sigma\gamma)$ with σ of order m . Then

$$\phi^m = \mu(\sigma\gamma)^m = \mu(\sigma^m \gamma^{\sigma^{m-1}} \dots \gamma^\sigma \gamma) = \mu(\gamma^{\sigma^{m-1}} \dots \gamma^\sigma \gamma).$$

Consequently, we have that

$$\text{GR}(\phi^m) = \text{GR}(\mu(\gamma^{\sigma^{m-1}} \dots \gamma^\sigma \gamma)) \leq 1.$$

Since $\text{GR}(\phi)^m = \text{GR}(\phi^m)$ (this follows from definition), we have $\text{GR}(\phi) \leq 1$. If $\text{GR}(\phi) < 1$, then by [8, Lemma 2.3], ϕ is an eventually trivial endomorphism. This is impossible because ϕ is an automorphism, and so we must have $\text{GR}(\phi) = 1$. $\square$

Next, we consider $\mathcal{H}_2$. By Theorem 2.3 again, every automorphism ϕ of $\mathcal{H}_2$ is induced from the conjugation by some element of $\sigma\gamma \in \mathcal{H}_2 \Sigma_2 \subset \text{Sym}_2$. We remark that the proof of the above theorem relies on only the fact that automorphisms are induced from conjugations. This makes possible to repeat the above proof verbatim for the set $\Gamma = \{g_2, \alpha\}$ of generators for $\mathcal{H}_2$. Therefore we have:

Theorem 3.2. *For any automorphism ϕ of $\mathcal{H}_2$, we have $\text{GR}(\phi) = 1$.*

4. ENDOMORPHISMS OF $\mathcal{H}_n$ WHICH ARE NOT MONIC

We recall the following results.

Theorem 4.1 ([6, Theorem 8.1A]). *Let Ω be any set with $|\Omega| > 4$. Then the nontrivial normal subgroups of $\text{Sym}(\Omega)$ are precisely: $\text{FAlt}(\Omega)$ and the subgroups of the form $\text{Sym}(\Omega, c)$ with $\aleph_0 \leq c \leq |\Omega|$. Here,*

$$\text{Sym}(\Omega, c) := \{x \in \text{Sym}(\Omega) \mid |\text{supp}(x)| < c\}.$$

In particular, $\text{Sym}(\Omega, \aleph_0) = \text{FSym}(\Omega)$, and so the nontrivial normal subgroups of Sym_n are precisely FAlt_n and FSym_n .

Theorem 4.2 ([16, 11.3.3]). *The nontrivial normal subgroups of $\text{Sym}(\Omega, B)$ are the groups $\text{Sym}(\Omega, D)$ with $\aleph_0 \leq D < B \leq 2^{|\Omega|}$ and $\text{FAlt}(\Omega)$. In particular, we have:*

- (1) *The only nontrivial normal subgroup of FSym_n is FAlt_n (when $B = \aleph_0$).*
- (2) *The only nontrivial normal subgroups of Sym_n are FSym_n and FAlt_n (when $B = 2^{\aleph_0}$).*

Corollary 4.3. *If N is a nontrivial normal subgroup of $\mathcal{H}_n$, then $N \cap \text{FSym}_n$ is either FAlt_n or FSym_n . In particular N contains FAlt_n .*

Proof. If N is a normal subgroup of $\mathcal{H}_n$, then $N \cap \text{FSym}_n$ is a normal subgroup of FSym_n . By Part (1) of Theorem 4.2, $N \cap \text{FSym}_n$ is either 1, FAlt_n or FSym_n .

If $N \cap \text{FSym}_n = \text{FAlt}_n$, then $\text{FAlt}_n \subset N$; if $N \cap \text{FSym}_n = \text{FSym}_n$, then $\text{FAlt}_n \subset \text{FSym}_n \subset N$.

We now consider the case $N \cap \text{FSym}_n = 1$. This implies that the surjection $\mathcal{H}_n \rightarrow \mathbb{Z}^{n-1}$ maps N isomorphically into $\mathbb{Z}^{n-1}$. If $N \neq 1$, then N contains an element $g \in \mathcal{H}_n$ of infinite order. There exists a point $x \in \text{supp}(g)$ whose orbit under g is infinite. Observe that for any transposition β exchanging two points in the orbit of x , $[\beta, g] = g^\beta g^{-1} \in N$ is a 3-cycle or a product of two commuting transpositions. In fact, if β exchanges two consecutive points x and $(x)g$ in $\text{supp}(g)$, then $[\beta, g] = ((x)g^{-1}, x, (x)g)$ is a 3-cycle; if β exchanges two non-consecutive points x and y in $\text{supp}(g)$, then we can show easily that $[\beta, g] = (x, y)((x)g^{-1}, (y)g^{-1})$ is a product of two commuting transpositions. In any case, N contains a non-trivial torsion element, and so $N \cap \text{FSym}_n \neq 1$. Consequently, $N = 1$ if $N \cap \text{FSym}_n = 1$. This completes the proof. $\square$

Corollary 4.4 ([5]). *The Houghton's group $\mathcal{H}_n$ is not residually finite.*

Proof. Suppose that $\mathcal{H}_n$ is residually finite. By definition, for any $x \in \mathcal{H}_n - \{1\}$ there exists a homomorphism ϕ from $\mathcal{H}_n$ to a finite group such that $\phi(x) \neq 1$. Equivalently, for any $x \in \mathcal{H}_n - \{1\}$ there exists a finite index normal subgroup N_x of $\mathcal{H}_n$ such that $x \notin N_x$. Since $N_x \neq 1$, Corollary 4.3 implies that $\text{FAlt}_n \subset N_x$ for all $x \in \mathcal{H}_n - \{1\}$; in particular, for $x \in \text{FAlt}_n - \{1\}$. This is a contradiction. $\square$

Remark 4.5. The Houghton's group $\mathcal{H}_n$ is not co-Hopfian for $n \geq 3$ ([3]). That is, there is an injection which is not an isomorphism. Indeed, $\phi : g_i \mapsto g_i^2$, $2 \leq i \leq n$, defines such an injection. It is direct to check that $\phi(r) = 1$ for all relators r of the presentation in Theorem 2.1. For the injectivity of ϕ one can realize $\phi(\mathcal{H}_n)$ as disjoint union of two subgroups each of which is isomorphic to $\mathcal{H}_n$. (See Example 5.6)

Remark 4.6. Recall that for a short exact sequence of groups $0 \rightarrow A \rightarrow G \rightarrow Q \rightarrow 1$ with A abelian, the following are equivalent:

- (1) Q acts trivially on A .
- (2) A lies in the center of G .

In this case, we say that the short exact sequence is *central*. Remark that if $A \cong \mathbb{Z}_2$ then the exact sequence is always central. In fact, for any $g \in G$ and $a \neq 1$ in A , we must have that $gag^{-1} = a$.

Corollary 4.7 ([5]). *The Houghton's group $\mathcal{H}_n$ is Hopfian.*

Proof. Let $\phi : \mathcal{H}_n \rightarrow \mathcal{H}_n$ be an epimorphism, but not an isomorphism. Put $K = \ker(\phi)$; then $K \neq 1$, $\mathcal{H}_n/K \cong \mathcal{H}_n$ and from the proof of Corollary 4.3, $K \cap \text{FSym}_n$ is either FAlt_n or FSym_n .

If $K \cap \text{FSym}_n = \text{FSym}_n$, then $\text{FSym}_n \subset K$ and thus we have

$$\mathcal{H}_n/\text{FSym}_n (\cong \mathbb{Z}^{n-1}) \longrightarrow \mathcal{H}_n/K \xrightarrow{\cong} \mathcal{H}_n.$$

This implies that $\mathcal{H}_n$ is abelian, which is absurd.

Consider next the case $K \cap \text{FSym}_n = \text{FAlt}_n$. Then we have a commuting diagram

$$\begin{array}{ccccccc} & & 1 & & 1 & & 1 \\ & & \uparrow & & \uparrow & & \uparrow \\ 1 & \longrightarrow & \text{FSym}_n/K \cap \text{FSym}_n & \longrightarrow & \mathcal{H}_n/K & \longrightarrow & \mathcal{H}_n/K \cdot \text{FSym}_n \longrightarrow 1 \\ & & \uparrow & & \uparrow & & \uparrow \\ 1 & \longrightarrow & \text{FSym}_n & \longrightarrow & \mathcal{H}_n & \longrightarrow & \mathbb{Z}^{n-1} \longrightarrow 1 \\ & & \uparrow & & \uparrow & & \uparrow \\ 1 & \longrightarrow & K \cap \text{FSym}_n & \longrightarrow & K & \longrightarrow & K/K \cap \text{FSym}_n \longrightarrow 1 \\ & & \uparrow & & \uparrow & & \uparrow \\ & & 1 & & 1 & & 1 \end{array}$$

Since $\text{FSym}_n/K \cap \text{FSym}_n = \text{FSym}_n/\text{FAlt}_n \cong \mathbb{Z}_2$, the top row exact sequence is a central extension by Remark 4.6. This implies that $\mathcal{H}_n/K \cong \mathcal{H}_n$ has a nontrivial center. But Lemma 4.8 below shows that $\mathcal{H}_n$ has trivial center. $\square$

Lemma 4.8. *For all n , the center of $\mathcal{H}_n$ is trivial.*

Proof. Assume that $\mathcal{Z}(\mathcal{H}_n)$ is nontrivial. By Corollary 4.3, $\text{FAlt}_n \subset \mathcal{Z}(\mathcal{H}_n)$. This implies that FAlt_n is abelian, a contradiction. $\square$

Let φ be an endomorphism of $\mathcal{H}_n$. In this section, we shall assume $K = \ker(\varphi) \neq 1$. [Since $\mathcal{H}_n$ is Hopfian by Corollary 4.7, φ cannot be epic.] By Corollary 4.3, $K \cap \text{FSym}_n$ is either FAlt_n or FSym_n .

Consider first the case $K \cap \text{FSym}_n = \text{FSym}_n$. Since $\text{FSym}_n \subset K$, φ induces the following diagram

$$\begin{array}{ccccc} \mathcal{H}_n & \longrightarrow & \mathcal{H}_n/\text{FSym}_n & \longrightarrow & \mathcal{H}_n/K \\ \downarrow \varphi & & \downarrow \bar{\varphi} & & \downarrow \hat{\varphi} \\ \mathcal{H}_n & \longrightarrow & \mathcal{H}_n/\text{FSym}_n & \longrightarrow & \mathcal{H}_n/K \end{array}$$

where the horizontal maps are canonical surjections. It is immediate from the definition of the growth rate that

$$\text{GR}(\hat{\varphi}) \leq \text{GR}(\bar{\varphi}) \leq \text{GR}(\varphi).$$

We claim now that $\text{GR}(\hat{\varphi}) = \text{GR}(\bar{\varphi}) = \text{GR}(\varphi)$. This follows from the fact that the endomorphism $\hat{\varphi} : \mathcal{H}_n/K \rightarrow \mathcal{H}_n/K$ induced by φ is simply the restriction of φ on the φ -invariant subgroup $\varphi(\mathcal{H}_n)$ of $\mathcal{H}_n$, $\varphi|_{\varphi(\mathcal{H}_n)} : \varphi(\mathcal{H}_n) \rightarrow \varphi(\mathcal{H}_n)$. It is known that $\text{GR}(\varphi) = \text{GR}(\varphi|_{\varphi(\mathcal{H}_n)}) = \text{GR}(\hat{\varphi})$, see for example [4, Sect. 4]. Therefore, we have

$$\text{GR}(\varphi) = \text{GR}(\bar{\varphi}).$$

Recall also that $\text{GR}(\bar{\varphi})$ is the spectral radius of the integer matrix determined by the endomorphism $\bar{\varphi}$ of $\mathcal{H}_n/\text{FSym}_n = \mathbb{Z}^{n-1}$.

Next we consider the case $K \cap \text{FSym}_n = \text{FAlt}_n$. Since $\text{FAlt}_n \subset K$, φ induces the following diagram

$$\begin{array}{ccccc} \mathcal{H}_n & \longrightarrow & \mathcal{H}_n/\text{FAlt}_n & \longrightarrow & \mathcal{H}_n/K \\ \downarrow \varphi & & \downarrow \tilde{\varphi} & & \downarrow \tilde{\varphi} \\ \mathcal{H}_n & \longrightarrow & \mathcal{H}_n/\text{FAlt}_n & \longrightarrow & \mathcal{H}_n/K \end{array}$$

where the horizontal maps are canonical surjections. Repeating the same argument as above, we have

$$\text{GR}(\varphi) = \text{GR}(\tilde{\varphi}).$$

Since FSym_n is the subgroup of $\mathcal{H}_n$ consisting of all elements that have finite order, FSym_n is ϕ -invariant. Consequently, ϕ induces the following diagram

$$\begin{array}{ccccc} \text{FSym}_n/\text{FAlt}_n & \longrightarrow & \mathcal{H}_n/\text{FAlt}_n & \longrightarrow & \mathcal{H}_n/\text{FSym}_n \\ \downarrow \phi' & & \downarrow \tilde{\phi} & & \downarrow \tilde{\phi} \\ \text{FSym}_n/\text{FAlt}_n & \longrightarrow & \mathcal{H}_n/\text{FAlt}_n & \longrightarrow & \mathcal{H}_n/\text{FSym}_n \end{array}$$

By [8, Lemma 2.5], we have

$$\text{GR}(\tilde{\phi}) = \max \{ \text{GR}(\phi'), \text{GR}(\tilde{\phi}) \}.$$

Remark also that $\text{FSym}_n/\text{FAlt}_n = \mathbb{Z}_2$, hence $\text{GR}(\phi') = 0$ or 1 depending on ϕ' is trivial or not trivial (then ϕ' is the identity). If $\text{GR}(\tilde{\phi}) = \text{sp}[\tilde{\phi}] < 1$ then $\tilde{\phi}$ is eventually trivial or $\text{GR}(\tilde{\phi}) = 0$. This implies that $\text{GR}(\tilde{\phi}) \geq 1$ if $\tilde{\phi}$ is not eventually trivial. So we can summarize our discussion in this section as follows.

Theorem 4.9. *Let ϕ be an endomorphism of $\mathcal{H}_n$ with nontrivial kernel. Then FSym_n is ϕ -invariant and so there results in the following commutative diagram*

$$\begin{array}{ccccc} \text{FSym}_n/\text{FAlt}_n & \longrightarrow & \mathcal{H}_n/\text{FAlt}_n & \longrightarrow & \mathcal{H}_n/\text{FSym}_n \\ \downarrow \phi' & & \downarrow \tilde{\phi} & & \downarrow \tilde{\phi} \\ \text{FSym}_n/\text{FAlt}_n & \longrightarrow & \mathcal{H}_n/\text{FAlt}_n & \longrightarrow & \mathcal{H}_n/\text{FSym}_n \end{array}$$

If ϕ' is nontrivial and $\tilde{\phi}$ is eventually trivial then $\text{GR}(\phi) = 1$, and otherwise we have

$$\text{GR}(\phi) = \text{sp}[\tilde{\phi}].$$

5. MONOMORPHISMS OF $\mathcal{H}_n$

Let $2 \leq n$ and let ϕ denote a monomorphism of $\mathcal{H}_n$ throughout this section unless otherwise stated. Let $\text{supp } g$ denote the support of $g \in \mathcal{H}_n$. Recall from Theorem 2.1 that if $n \geq 3$ then $\mathcal{H}_n$ has the following presentation

$$(5-1) \quad \alpha^2 = 1, (\alpha\alpha^{g_2})^3 = 1, [\alpha, \alpha^{g_2^2}] = 1, \alpha = [g_i, g_j], \alpha^{g_i^{-1}} = \alpha^{g_j^{-1}}$$

for $2 \leq i \neq j \leq n$.

For $i \neq j$ and $k \in \mathbb{N}$, the support of the transposition $\alpha^{g_i^{k+1}}$ belongs to R_i and does not intersect $\text{supp } g_j$. Consequently we have the following identities

$$(5-2) \quad [\alpha^{g_i^{k+1}}, g_j] = 1$$

for $k \in \mathbb{N}$ and $i \neq j$. By the same reason, we also have

$$(5-3) \quad [\alpha, \alpha^{g_i^{k+1}}] = 1$$

for all $k \in \mathbb{N}$. Note that the actions of g_i^{-1} and g_j^{-1} coincide on R_1 . So we have identities

$$(5-4) \quad \alpha^{g_i^{-k}} = \alpha^{g_j^{-k}}$$

for all $2 \leq i, j \leq n$ and $k \in \mathbb{N}$.

We say $P \in \text{supp } g$ is an *essential point* of g if its orbit under g is infinite. Let $\text{Esupp } g$ denote the set of essential points of g . For two elements $g, h \in \mathcal{H}_n$, we say g *intersects* h if $\text{supp } g \cap \text{supp } h \neq \emptyset$. Similarly we say g *intersects* $X \subset X_n$ if $\text{supp } g \cap X \neq \emptyset$.

For each pair $2 \leq i \neq j \leq n$, $[\phi(g_i), \phi(g_j)] = \phi(\alpha)$ is nontrivial and has order 2 for a monomorphism ϕ . So $\phi(\alpha)$ can be written

$$(5-5) \quad \phi(\alpha) = \tau_1 \cdots \tau_\ell$$

as a product of commuting transpositions. Let T_α denote the set of transpositions in (5-5), and let $S_\alpha = \text{supp } \phi(\alpha)$. Note that each $P \in S_\alpha$ determines a *unique* transposition in T_α which moves P . Let T_i be the collection of transpositions τ in (5-5) with $\text{supp } \tau \cap \text{Esupp } \phi(g_i) \neq \emptyset$, $2 \leq i \leq n$.

Lemma 5.1. *Suppose ϕ is a monomorphism of $\mathcal{H}_n$ with $n \geq 2$. For each i , $T_i \neq \emptyset$.*

Proof. We first consider a monomorphism when $n \geq 3$. Fix i . The conjugations $\alpha^{g_i^k}$ are all distinct for $k \in \mathbb{Z}$. A monomorphism ϕ induces a bijection between infinite sets $\{\alpha^{g_i^k} \mid k \in \mathbb{Z}\} \leftrightarrow \{\phi(\alpha^{g_i^k}) \mid k \in \mathbb{Z}\} = A$. Let us denote by β_k conjugation

$$\beta_k = \phi(\alpha)^{\phi(g_i)^k} = (\tau_1 \cdots \tau_\ell)^{\phi(g_i)^k}.$$

If all points of $\text{supp } \phi(\alpha)$ are not essential points of $\phi(g_i)$, then the set A must be finite since $\phi(g_i)^{k_0}$ conjugates $\phi(\alpha)$ to itself for some integer $k_0 \neq 0$. Suppose that all points of S_α are not essential points of $\phi(g_i)$. If $S_\alpha \cap \text{supp } \phi(g_i) = \emptyset$, then it is clear that $\phi(\alpha)^{\phi(g_i)} = \phi(\alpha)$. Next consider $S_\alpha \cap \text{supp } \phi(g_i) = \{P_1, \dots, P_s\} \neq \emptyset$. By the supposition, each P_j is not an essential point of $\phi(g_i)$, i.e., $(P_j)\phi(g_i)^{k_j} = P_j$ for some $k_j > 0$. Let $k_0 = \text{lcm}(k_1, \dots, k_s)$. Hence $(P_j)\phi(g_i)^{k_0} = P_j$ for all $j = 1, \dots, s$ and so $(P)\phi(g_i)^{k_0} = P$ for all $P \in S_\alpha$. This shows that $\beta_{k_0} = \phi(\alpha)$.

Let $P \in \text{supp } \phi(\alpha)$ be an essential point of $\phi(g_i)$. Let τ' denote the unique transposition in (5-5) which moves P to $Q = (P)\tau'$. We claim that Q is also an essential point of $\phi(g_i)$. Assume the contrary that

$$(Q)\phi(g_i)^{k_0} = Q$$

for some integer $k_0 \neq 0$ ($k_0 = 1$ when Q is fixed by $\phi(g_i)$, and $|k_0| \geq 2$ when $Q \in \text{supp } \phi(g_i)$). We need to examine β_k when k is a multiple of k_0 .

Since $\phi(g_i)^{mk_0}$ fixes Q for each nonzero $m \in \mathbb{Z}$, $\phi(g_i)^{mk_0}$ conjugates $\tau' = (P, Q)$ to the transposition, denoted by τ'_m , which exchanges $(P)\phi(g_i)^{mk_0}$ and Q . See Figure 1. Observe that

$$Q \in \text{supp } \tau' \cap \text{supp } \tau'_m \subset S_\alpha \cap \text{supp } \beta_{mk_0}$$

for each nonzero $m \in \mathbb{Z}$. To draw a contradiction, we use identities (5-3) which say in particular that $\phi(\alpha)$ commute with β_k for all $k = mk_0$ with $|mk_0| \geq 2$. We apply Lemma 5.2 to see that the intersection $I_m = S_\alpha \cap \text{supp } \beta_{mk_0}$ satisfies that

$$(5-6) \quad (I_m)\phi(\alpha) = I_m = (I_m)\beta_{mk_0}$$

for all m with $|mk_0| \geq 2$. In particular, S_α contains $(Q)\beta_{mk_0} = (Q)\tau'_m = (P)\phi(g_i)^{mk_0}$ for infinitely many $m \in \mathbb{Z}$. Since P is an essential point of $\phi(g_i)$, S_α must be an infinite set. This contradicts that an element $\phi(\alpha) \in \text{FSym}_n$ has a finite support. Therefore Q is also an essential point of $\phi(g_i)$, and hence the transposition τ' exchanges two essential points P and Q of $\phi(g_i)$.

For $n = 2$, recall that $\mathcal{H}_2$ has a presentation

$$(5-7) \quad \mathcal{H}_2 = \langle g_2, \alpha \mid \alpha^2 = 1, (\alpha\alpha^{g_2})^3 = 1, [\alpha, \alpha^{g_2^k}] = 1 \text{ for all } |k| > 1 \rangle$$

It follows from analogous arguments applied to g_2 with commutation relations above that $T_i \neq \emptyset$. $\square$




FIGURE 1. τ'_m moves Q to $(P)\phi(g_i)^{mk_0}$ for infinitely many $m \in \mathbb{Z}$.

Lemma 5.2. *Suppose two involutions $\beta, \gamma \in \text{FSym}_n$ commute. Then $I = (I)\beta = (I)\gamma$ where $I = \text{supp } \beta \cap \text{supp } \gamma$.*

Proof. We may assume the intersection $I = \text{supp } \beta \cap \text{supp } \gamma$ is non-empty. Since β is an involution, $B = I \cup (I)\beta$ is a β -invariant subset of $\text{supp } \beta$. Similarly $C = I \cup (I)\gamma$ satisfies $(C)\gamma = C$. Let β_B and γ_C denote the restrictions of β and γ on B and C respectively. We denote by β_{B^c} and γ_{C^c} the restrictions on the complements $B^c = \text{supp } \beta \setminus B$ and $C^c = \text{supp } \gamma \setminus C$ respectively. We can write

$$\beta = \beta_B \cdot \beta_{B^c} \text{ and } \gamma = \gamma_C \cdot \gamma_{C^c}.$$

Note that $B^c \cap \text{supp } \gamma = \emptyset$ and $C^c \cap \text{supp } \beta = \emptyset$.

Since $\beta^\gamma = \beta$ and

$$\beta^\gamma = (\beta_B \cdot \beta_{B^c})^\gamma = (\beta_B)^\gamma \cdot (\beta_{B^c})^\gamma = (\beta_B)^\gamma \cdot \beta_{B^c} = (\beta_B)^{\gamma_C} \cdot \beta_{B^c},$$

we have $(\beta_B)^{\gamma_C} \cdot \beta_{B^c} = \beta = \beta_B \cdot \beta_{B^c}$. So, γ_C conjugates β_B to itself, and B is an invariant set of γ_C . Similarly $\gamma^\beta = \gamma$ implies that $(\gamma_C)^{\beta_B} = \gamma_C$.

We first show $B \subset C$. If $P \in B \setminus C \subset (I)\beta \setminus (I)\gamma$, there exists $Q \in I$ such that $P = (Q)\beta_B$. Observe that P and Q determine a unique transposition in the cycle decomposition of β_B such that

$$P = (Q)\tau.$$

Since γ_C moves Q to $Q' \neq P$ while fixing $P \notin C$, γ_C conjugates τ to $\tau^{\gamma_C} \neq \tau$ which exchanges P and Q' . However this contradicts the uniqueness of the transposition τ of β_B . In other words γ_C fails to conjugate β_B to itself. Therefore $(I)\beta \subset (I)\gamma$ and so $B \subset C$.

For the converse $(I)\gamma \subset (I)\beta$ or $C \subset B$, one can use analogous arguments applied to $\gamma^\beta = \gamma$. In all,

$$(I)\beta = (I)\gamma, \quad B = I \cup (I)\beta = I \cup (I)\gamma = C.$$

Furthermore, if $P \in (I)\beta = (I)\gamma$ then $P = (Q)\beta = (Q')\gamma$ for some $Q, Q' \in I$, thus $P \in \text{supp } \beta \cap \text{supp } \gamma = I$. The reverse inclusion follows since β and γ are involutions, hence $(I)\beta = (I)\gamma = I$. $\square$

Remark 5.3. (Squares on I) An interesting consequence of Lemma 5.2 is that a point $P \in I$ with $(P)\beta \neq (P)\gamma$ determines a certain *square* (or *4-cycle of transpositions*) on I . Let τ_1 and τ'_1 be the transpositions of β and γ respectively which moves P to distinct points. The Lemma 5.2 says $Q = (P)\tau_1 \in I \subset \text{supp } \gamma$ and $R = (P)\tau'_1 \in I \subset \text{supp } \beta$. Let τ_2 and τ'_2 be the unique transpositions of β and γ respectively so that $R \in \text{supp } \tau_2$ and $Q \in \text{supp } \tau'_2$. The intersection $\text{supp } \tau_2 \cap \text{supp } \tau'_2$ contains a point S since $\beta\gamma = \gamma\beta$ applied to P implies

$$P \xrightarrow{\beta} (P)\beta = (P)\tau_1 = Q \xrightarrow{\gamma} (Q)\gamma = (Q)\tau'_2 = S,$$

$$P \xrightarrow{\gamma} (P)\gamma = (P)\tau'_1 = R \xrightarrow{\beta} (R)\beta = (R)\tau_2 = S.$$

Applying alternate compositions of β and γ , we have a square on the four points $P \xrightarrow{\tau'_1} R \xrightarrow{\tau_2} S \xrightarrow{\tau'_2} Q \xrightarrow{\tau_1} P$. Reading off four transpositions of β and γ in order we have a 4-cycle $\tau'_1 - \tau_2 - \tau'_2 - \tau_1$

of transpositions. (This should not be confused with a 4-cycle of FSym_n .) Figure 2 illustrates the square on four points and corresponding 4-cycle of transpositions.




FIGURE 2.

Next we consider a useful necessary condition when two involutions $\beta, \gamma \in \text{FSym}_n$ form $\beta\gamma$ of order 3. Suppose β and γ have cycle decompositions

$$\beta = \tau_1 \cdots \tau_k \text{ and } \gamma = \tau'_1 \cdots \tau'_\ell.$$

Let $\mathcal{S} = \{\tau_1, \dots, \tau_k\}$ and $\mathcal{T} = \{\tau'_1, \dots, \tau'_\ell\}$, and let $\mathfrak{P}(\mathcal{S})$ and $\mathfrak{P}(\mathcal{T})$ denote the power sets of $\mathcal{S}$ and $\mathcal{T}$ respectively. We define assignments

$$(5-8) \quad \xi : \mathcal{S} \longrightarrow \mathfrak{P}(\mathcal{T}) \text{ and } \chi : \mathcal{T} \longrightarrow \mathfrak{P}(\mathcal{S})$$

so that $\text{supp } \tau_i$ has non-trivial intersection with each support of $\tau'_j \in \xi(\tau_i)$. The assignment $\chi(\tau_i)$ is defined by the same manner after swapping $\mathcal{S}$ and $\mathcal{T}$. As we shall see below the cardinalities of $\xi(*)$ and $\chi(*)$ are either one or two. Let $\mathcal{S}_1 \subset \mathcal{S}$ and $\mathcal{T}_1 \subset \mathcal{T}$ be defined by

$$\mathcal{S}_1 = \{\tau_i \in \mathcal{S} : |\xi(\tau_i)| = 1\}, \quad \mathcal{T}_1 = \{\tau'_j \in \mathcal{T} : |\chi(\tau'_j)| = 1\}.$$

Lemma 5.4. *With the notations for two involutions β and γ such that $\beta\gamma$ has order 3 as above, $1 \leq |\xi(\tau_i)| \leq 2$ and $1 \leq |\chi(\tau'_j)| \leq 2$ for each $\tau_i \in \mathcal{S}$ and $\tau'_j \in \mathcal{T}$. There exists a bijection $\mathcal{S}_1 \leftrightarrow \mathcal{T}_1$ where correspondence is given by $\xi(\tau_i) = \{\tau'_j\} \Leftrightarrow \chi(\tau'_j) = \{\tau_i\}$.*

Proof. Fix a transposition $\tau_i \in \mathcal{S}$ which exchanges two points P and Q . It is obvious that $\mathcal{T}$ contains at most two transpositions which move P or Q . So the cardinality of $\xi(\tau_i)$ is at most 2. To show $|\xi(\tau_i)| \neq 0$, assume that every transposition of $\mathcal{T}$ fixes P and Q . For convenience we write $\beta = \tau_i \cdot \beta'$ where $\beta' = \tau_1 \cdots \widehat{\tau_i} \cdots \tau_k$ fixes P and Q . If γ fixes P and Q then the product $(\beta\gamma)^3$ can be written as

$$(\beta\gamma)^3 = (\tau_i \beta' \cdot \gamma)^3 = \tau_i^3 (\beta' \gamma)^3 = \tau_i (\beta' \gamma)^3$$

since τ_i commutes with β' and γ . The support of $\tau_i (\beta' \gamma)^3$ contains P and Q and so it is not empty. This means that $\beta\gamma$ is not an element of order 3. We have shown that $|\xi(\tau_i)|$ equals 1 or 2. For the cardinality of $\chi(*)$ one applies analogous arguments after swapping β and γ .

To establish the correspondence $\mathcal{S}_1 \leftrightarrow \mathcal{T}_1$, suppose $\xi(\tau_i) = \{\tau'_j\}$, i.e., $\tau'_j \in \mathcal{T}$ is the unique transposition which moves P or Q . If $\tau_i = \tau'_j$ it is obvious that $\xi(\tau_i) = \{\tau'_j\} \Leftrightarrow \chi(\tau'_j) = \{\tau_i\}$. We assume $\tau_i \neq \tau'_j$ and show that $\chi(\tau'_j) = \{\tau_i\}$ as follows. It suffices to show that $|\chi(\tau'_j)| = 1$ since $\tau_i \in \chi(\tau'_j)$. If $|\chi(\tau'_j)| = 2$ then $\mathcal{S}$ contains a transposition $\tau \neq \tau_i$, whose support intersects $\text{supp } \tau'_j$ nontrivially. Since τ'_j has to move P or Q , so we may assume τ'_j moves P . Let $P' = (P)\tau'_j$. Then τ exchanges P' and $R = (P')\tau$, and the point Q is fixed by γ . Applying the product $\beta\gamma$ repeatedly

to the point R we have $(R)(\beta\gamma)^3 = P' \neq R$ since

$$\begin{aligned} R &\xrightarrow{\beta} (R)\beta = (R)\tau = P' \xrightarrow{\gamma} (P')\gamma = (P')\tau'_j = P, \\ P &\xrightarrow{\beta} (P)\beta = (P)\tau_i = Q \xrightarrow{\gamma} (Q)\gamma = Q, \\ Q &\xrightarrow{\beta} (Q)\beta = (Q)\tau_i = P \xrightarrow{\gamma} (P)\gamma = (P)\tau'_j = P'. \end{aligned}$$

So $R \in \text{supp}(\beta\gamma)^3 \neq \emptyset$. This contradicts that the product $\beta\gamma$ has order 3. Figure 3 illustrates this situation.

So far we have shown that $\xi(\tau_i) = \{\tau'_j\}$ implies $\chi(\tau'_j) = \{\tau_i\}$. The converse follows because a symmetric argument shows that $\gamma\beta = (\beta\gamma)^{-1}$ fails to have order 3 if $\chi(\tau'_j) = \{\tau_i\}$ but $|\xi(\tau_i)| = 2$. Therefore $\xi(\tau_i) = \{\tau'_j\} \Leftrightarrow \chi(\tau'_j) = \{\tau_i\}$ determines the correspondence $\mathcal{S}_1 \leftrightarrow \mathcal{T}_1$. $\square$




FIGURE 3.

Remark 5.5. (Hexagons on I) To understand the relation between complements of $\mathcal{S}_1$ and $\mathcal{T}_1$ consider assignments ξ and χ on

$$\mathcal{S}_2 = \mathcal{S} \setminus \mathcal{S}_1 \text{ and } \mathcal{T}_2 = \mathcal{T} \setminus \mathcal{T}_1.$$

Lemma 5.4 implies that $|\xi(\tau_i)| = 2$ for all $\tau_i \in \mathcal{S}_2$, and conversely $|\chi(\tau'_j)| = 2$ for all $\tau'_j \in \mathcal{T}_2$. It is interesting to see that each element $\tau_i \in \mathcal{S}_2$ involves a certain *hexagon* (or 6-cycle of transpositions) on $I = \text{supp } \beta \cap \text{supp } \gamma$. Suppose $\tau_i \in \mathcal{S}_2$ exchanges P and Q . Let us describe the other 5 transpositions. Let $\xi(\tau_i) = \{\tau'_P, \tau'_Q\}$ where τ'_P exchanges P and $(P)\tau'_P = P_1$ and τ'_Q exchanges Q and $(Q)\tau'_Q = Q_1$. Since $|\chi(\tau'_P)| = 2$ and $\tau_i \in \chi(\tau'_P)$, $\chi(\tau'_P)$ contains $\tau_{P_1} \in \mathcal{S}_2$ which exchanges P_1 and $P_2 = (P_1)\tau_{P_1}$. Similarly $\chi(\tau'_Q)$ contains two transpositions τ_i and $\tau_{Q_1} \in \mathcal{S}_2$ which exchanges Q_1 and $Q_2 = (Q_1)\tau_{Q_1}$. One can check that $\tau_{P_1} \neq \tau_{Q_1}$. If $\tau_{P_1} = \tau_{Q_1}$ the four transpositions τ_i , τ'_Q , $\tau_{P_1} = \tau_{Q_1}$, and τ'_P form a square as described in Remark 5.3. It is direct to see that $\{P_1, Q_1\} \subset \text{supp}(\beta\gamma)^3$, which contradicts that $\beta\gamma$ has order 3. So far we have 5 transpositions τ_i , τ'_P , τ'_Q , τ_{P_1} , and τ_{Q_1} . Next we check that τ_{P_1} and τ_{Q_1} determine a unique transposition $\tau'_j \in \mathcal{T}_2$ so that $\tau'_j \in \xi(\tau_{P_1}) \cap \xi(\tau_{Q_1})$, namely τ'_j exchanges P_2 and Q_2 . Applying $\beta\gamma$ to the point Q_2 twice we have

$$\begin{aligned} Q_2 &\xrightarrow{\beta} (Q_2)\beta = (Q_2)\tau_{Q_1} = Q_1 \xrightarrow{\gamma} (Q_1)\gamma = (Q_1)\tau'_Q = Q \\ Q &\xrightarrow{\beta} (Q)\beta = (Q)\tau_i = P \xrightarrow{\gamma} (P)\gamma = (P)\tau'_P = P_1. \end{aligned}$$

Since $\beta\gamma$ has order 3 we must have $(Q_2)(\beta\gamma)^3 = (P_1)\beta\gamma = Q_2$. The later identity means that

$$(P_1)\tau_{P_1}\gamma = (P_2)\gamma = (P_2)\tau'_j = Q_2.$$

Thus the transposition τ'_j exchanges P_2 and Q_2 , and so it is the unique transposition of $\mathcal{T}_2$ such that $\tau'_j \in \xi(\tau_{P_1}) \cap \xi(\tau_{Q_1})$ as desired. Alternate composition β -and-then- γ applied to P determines

6 points

$$P \xrightarrow{\tau_i} Q \xrightarrow{\tau'_Q} Q_1 \xrightarrow{\tau_{Q_1}} Q_2 \xrightarrow{\tau'_j} P_2 \xrightarrow{\tau_{P_1}} P_1 \xrightarrow{\tau'_{P'}} P.$$

The six transpositions in order form a hexagon or 6-cycle of transpositions: $\tau_i - \tau'_Q - \tau_{Q_1} - \tau'_j - \tau_{P_1} - \tau'_{P'}$ (Again, one should distinguish this from a 6-cycle of $\mathcal{H}_n$.) Figure 4 illustrates the hexagon on four points and corresponding 6-cycle of transpositions.




FIGURE 4.

Partial translations. For $g \in \mathcal{H}_n$, it is useful to decompose g into *essential part* and *finitary part*. Let $E(g)$ and $F(g)$ denote the restrictions of g on $\text{Esupp } g$ and $\text{supp } g \setminus \text{Esupp } g$ respectively. As an element of Sym_n , the restriction of g on $\text{Esupp } g$ has a cycle decomposition. The set $\text{Esupp } g$ is partitioned into orbits of essential points of g . The element g restricts on each orbit to define an infinite cycle, which will be called a *partial translation* of g . Denoting partial translations of g by $\mathbf{p}_1, \dots, \mathbf{p}_\ell$, we can write g as

$$(5-9) \quad g = \mathbf{p}_1 \cdots \mathbf{p}_\ell \cdot f$$

where $\text{supp } f \subset \text{supp } g \setminus \text{Esupp } g$. Partial translations of g commute with each other. In case g has no essential points, (5-9) becomes $g = f$. Note that a partial translation is not an element of $\mathcal{H}_n$ in general. For example, g_2^2 has two partial translations, which are not eventual translations.

As an infinite cycle on X_n , each partial translation $\mathbf{p}$ can be realized as an embedding $\mathbb{Z} \hookrightarrow X_n$. Picking a base point $x_0 \in \text{supp } \mathbf{p}$ we identify $\text{supp } \mathbf{p} = \{(x_0)\mathbf{p}^k | k \in \mathbb{Z}\}$ with $\mathbb{Z}$. Let $[k]_{\mathbf{p}}$ denote the point of $\text{supp } \mathbf{p}$ corresponding to $k \in \mathbb{Z}$:

$$(5-10) \quad (x_0)\mathbf{p}^k \leftrightarrow [k]_{\mathbf{p}}.$$

Under the identification (5-10), a partial translation $\mathbf{p}$ of g translates points on $\text{supp } \mathbf{p}$ by $+1$ in the above identification, i.e.,

$$(5-11) \quad [k]_{\mathbf{p}} g^m = [k]_{\mathbf{p}} \mathbf{p}^m \leftrightarrow [k + m]_{\mathbf{p}}$$

for each $m \in \mathbb{Z}$.

By the definition (2-1) an eventual translation $g \in \mathcal{H}_n$ acts as a translation by m_i on R_i outside a finite set F_g where $\pi(g) = (m_1, \dots, m_n)$. We call a ray R_i a *source* of g if $m_i < 0$, and we call a ray R_i a *target* of g if $m_i > 0$. As a restriction of g on one of its orbits, a partial translation $\mathbf{p}$ of g has unique *source* and *target* so that $\mathbf{p}$ moves points *from the source towards the target*. More precisely there exist exactly two rays $R_{\mathbf{p}}^-$ and $R_{\mathbf{p}}^+$ in $\{R_1, \dots, R_n\}$, and $k_0 \in \mathbb{N}$ such that

$$(5-12) \quad [-k]_{\mathbf{p}} \in R_{\mathbf{p}}^- \text{ and } [k]_{\mathbf{p}} \in R_{\mathbf{p}}^+$$

for all $k \geq k_0$.

Note that the generators $g_2, \dots, g_n$ of $\mathcal{H}_n$ are all partial translations. Generators g_i 's share a unique source R_1 on which the actions of g_i^{-1} 's coincide as translations by $+1$. Moreover those generators have all distinct targets. Intuitively this behavior can be characterized as, for each pair $i \neq j$,

- (1) two actions of g_i and g_j are identical on $R_1 \setminus (1, 1)$,
- (2) once g_i and g_j ‘diverge’ from $(1, 1)$ then they never meet again,
- (3) the commutator $[g_i, g_j] = \alpha$ is the transposition which exchanges $(1, 1)$ and $(1, 2)$.

Note that $\text{supp } \alpha$ consists of two points which are the last two points before g_i and g_j diverge. It turns out that generators $\phi(g_2), \dots, \phi(g_n)$ of $\phi(\mathcal{H}_n)$ follow the same rule provided ϕ is a monomorphism. See Lemmas 5.7 and 5.8.

Example 5.6. Consider the monomorphism $\phi : \mathcal{H}_3 \rightarrow \mathcal{H}_3, g_i \mapsto g_i^2$ for $i = 2, 3$. The generators g_2^2 and g_3^2 of $\phi(\mathcal{H}_3)$ share a common source R_1 and their targets, R_2 and R_3 respectively, are distinct. Both g_2^2 and g_3^2 are products of two partial translations. Let $\mathbf{p}_i$ and $\mathbf{p}'_i$ denote the two partial transpositions of g_i^2 moving $(1, 1)$ and $(1, 2)$ respectively for $i = 2, 3$. Note that R_1 is the unique source for all four partial translations. The ray R_2 is the target of $\mathbf{p}_2$ and $\mathbf{p}'_2$, and R_3 is the target of $\mathbf{p}_3$ and $\mathbf{p}'_3$. As in (5–10), we label $\text{supp } \mathbf{p}_2$ and $\text{supp } \mathbf{p}_3$ by $\mathbb{Z}$ with a base point $(1, 3)$:

$$(1, 3) \leftrightarrow [0]_{\mathbf{p}_2} \text{ and } (1, 3) \leftrightarrow [0]_{\mathbf{p}_3}.$$

It is direct to check that $[k]_{\mathbf{p}_2} = [k]_{\mathbf{p}_3}$ for all $k \leq 0$ and so $\mathbf{p}_2$ and $\mathbf{p}_3$ agree on $\{[k]_{\mathbf{p}_2} | k \leq 1\}$. Note that $[\mathbf{p}_2, \mathbf{p}_3]$ is the transposition which exchanges $(1, 3) \leftrightarrow [0]_{\mathbf{p}_2}$ and $(1, 1) \leftrightarrow [1]_{\mathbf{p}_2}$, the last two consecutive points before they diverge. The partial translations $\mathbf{p}'_2$ and $\mathbf{p}'_3$ follow the same rule: they agree on the source R_1 except $(1, 2)$, and $[\mathbf{p}'_2, \mathbf{p}'_3]$ exchanges $(1, 2)$ and $(1, 4)$, the last two consecutive points before they diverge.

We also remark that two pairs of partial translations induce decomposition of the underlying set X_3 . Let X denote the orbit of $(1, 1)$ under $\phi(\mathcal{H}_3)$. In other words, X is the union of two supports of $\mathbf{p}_2$ and $\mathbf{p}_3$. Similarly $\mathbf{p}'_2$ and $\mathbf{p}'_3$ determine a subset X' which is the orbit of $(1, 2)$ under $\phi(\mathcal{H}_3)$. Obviously two invariant subsets provide a partition $X_3 = X \sqcup X'$. The corresponding decomposition of $\phi(\mathcal{H}_3)$ says that $\phi(\mathcal{H}_3)$ is the disjoint union of two identical subgroups H_3 and H'_3 where H_3 and H'_3 are generated by $\mathbf{p}_i$ and $\mathbf{p}'_i$ respectively, $i = 2, 3$. The isomorphism between H_3 and H'_3 can be described by the obvious bijection between X and X_3 . In particular, ϕ is injective.

Lemma 5.7. *A monomorphism $\phi : \mathcal{H}_n \rightarrow \mathcal{H}_n$ determines $\ell \in \mathbb{N}$ and a ray R_ℓ such that R_ℓ is a unique common source of $\phi(g_i)$, and $\phi(g_i)$ has translation length $-\ell$ on R_ℓ for all $2 \leq i \leq n$. Moreover targets of $\phi(g_i)$ and $\phi(g_j)$ do not share a ray if $2 \leq i \neq j \leq n$.*

Proof. The group $\mathcal{H}_2$ has only one generator g_2 of infinite order. So $\phi(g_2)$ satisfies the above automatically. To deal with a monomorphism ϕ of $\mathcal{H}_n$ with $3 \leq n$ recall notations: $S_\alpha = \text{supp } \phi(\alpha)$, $T_\alpha =$ the set of all transpositions in (5–5), $T_i \subset T_\alpha$ consisting of transpositions which intersect $\text{Esupp } (\phi(g_i))$. Being a monomorphism, ϕ implies that the order of every $\phi(g_i)$ is infinite. So $\phi(g_i)$ contains at least one partial translation when written as in (5–9). Let $\mathcal{P}_i$ denote the set of all partial translations of $\phi(g_i)$.

Fix i . Lemma 5.1 says that $T_i \neq \emptyset$, and so $\mathcal{P}_i$ contains a partial translation $\mathbf{p}$ such that

$$(5-13) \quad \text{supp } \mathbf{p} \cap S_\alpha \neq \emptyset.$$

Consider the source R_ℓ of $\mathbf{p}$ with the property (5–12). Since R_ℓ is also a source of $\phi(g_i)$, $\phi(g_i)$ has translation length $-\ell < 0$ on R_ℓ .

Step 1. In this step we show $\phi(g_j)$ has the same translation length $-\ell < 0$ on R_ℓ for all $j \neq i$. The identities (5–4) imply that two involutions $\phi(\alpha)^{\phi(g_i)^{-k}}$ and $\phi(\alpha)^{\phi(g_j)^{-k}}$ are identical for all $k \in \mathbb{N}$. So the two involutions must share their supports,

$$(5-14) \quad (S_\alpha)\phi(g_i)^{-k} = (S_\alpha)\phi(g_j)^{-k}$$

for all $k \in \mathbb{N}$. Since $\mathbf{p}$ intersects S_α , one can take $A \in \text{supp } \mathbf{p} \cap S_\alpha \neq \emptyset$. By (5–12), we have

$$(A)\mathbf{p}^{-k} \in \text{supp } \mathbf{p} \cap R_\ell$$

for all $k \geq k_0$ where $k_0 \in \mathbb{N}$. Hence $(A)\mathbf{p}^{-k} \in (S_\alpha)\mathbf{p}^{-k} \cap R_l$. Moreover, $(S_\alpha)\mathbf{p}^{-k} \subset (S_\alpha)\phi(g_i)^{-k}$ for each integer k because $\mathbf{p}$ is the restriction of $\phi(g_i)$ on $\text{supp } \mathbf{p}$. Therefore $(S_\alpha)\phi(g_i)^{-k} \cap R_l$ is non-empty for all but finitely many k . The identities (5-14) imply

$$(5-15) \quad (S_\alpha)\phi(g_i)^{-k} \cap R_l = (S_\alpha)\phi(g_j)^{-k} \cap R_l$$

for all $k \in \mathbb{N}$. The identities (5-15) force $\phi(g_j)$ to have the same translation length $-\ell < 0$ on R_l . Consider the two smallest balls on X_n (centered at the origin) which contain two sets in (5-15) respectively. If $\phi(g_j)$ moves points on R_l (up to a finite set) by $\ell' \neq -\ell$, the two balls have to have different sizes for infinitely many k . This means $\phi(g_i)$ and $\phi(g_j)$ fail to satisfy the identities (5-15). Therefore $\phi(g_i)$ and $\phi(g_j)$ have the same translation length $-\ell < 0$ on R_l . Since i and j were arbitrary the first assertion of the Lemma is verified. The uniqueness of the ray will be verified in the last step.

Step 2. Identify $\text{supp } \mathbf{p}$ with $\{[k]_{\mathbf{p}} : k \in \mathbb{Z}\}$ as described in (5-10). We will show that there exists $k_0 \in \mathbb{Z}$ such that for each $j \neq i$, $\mathbf{p}$ intersects only one partial translation $\mathbf{q} \in \mathcal{P}_j$ satisfying

$$(5-16) \quad \mathbf{p} = \mathbf{q} \text{ on } \{[k]_{\mathbf{p}} : k \leq k_0\}$$

and

$$(5-17) \quad \text{supp } \mathbf{p} \cap \text{supp } \mathbf{q} = \{[k]_{\mathbf{p}} : k \leq k_0 + 1\}.$$

From Step 1 we see that actions of $\phi(g_j)$ and $\phi(g_i)$ are identical on R_l as a translation by $-\ell < 0$ up to a finite set. In particular, there exists $k_1 \in \mathbb{Z}$ such that

$$(5-18) \quad (P)\phi(g_i) = (P)\phi(g_j)$$

for all $P \in \{[k]_{\mathbf{p}} : k \leq k_1\}$. The orbit of (any such point) P under $\phi(g_j)$ determines a partial translation $\mathbf{q}$ as $\mathbf{p}$ is the restriction of $\phi(g_i)$ on the orbit of P under $\phi(g_i)$. From the identity (5-18) it is immediate to check that $\mathbf{q}$ satisfies

$$(5-19) \quad \mathbf{q} = \mathbf{p} \text{ on } \{[k]_{\mathbf{p}} : k \leq k_1\}.$$

We first claim that $\mathbf{p} \neq \mathbf{q}$. If $\mathbf{p} = \mathbf{q}$ then $\phi(\alpha)$ restricted to $\text{supp } \mathbf{p}$ is the identity map since

$$(P)\phi(g_i)\phi(g_j) = (P)\mathbf{p}\mathbf{q} = (P)\mathbf{p}^2 = (P)\mathbf{q}\mathbf{p} = (P)\phi(g_j)\phi(g_i)$$

for all $P \in \text{supp } \mathbf{p}$. So $\mathbf{p} = \mathbf{q}$ implies that $\mathbf{p}$ does not intersect $\phi(\alpha)$. However this contradicts the condition (5-13).

From the condition (5-19) together with $\mathbf{p} \neq \mathbf{q}$ we can consider the largest $k_2 \geq 0$ such that

$$([k_1 + k_2]_{\mathbf{p}})\mathbf{p} = ([k_1 + k_2]_{\mathbf{p}})\mathbf{q},$$

or equivalently k_2 is the smallest number such that

$$([k_1 + k_2 + 1]_{\mathbf{p}})\mathbf{p} \neq ([k_1 + k_2 + 1]_{\mathbf{p}})\mathbf{q}.$$

It is obvious that $k_0 = k_1 + k_2$ satisfies (5-16) and $\text{supp } \mathbf{p} \cap \text{supp } \mathbf{q} \supset \{[k]_{\mathbf{p}} : k \leq k_0 + 1\}$. For the reverse inclusion we assume $[k]_{\mathbf{p}} \in \text{supp } \mathbf{q}$ with $k \geq k_0 + 2$ and then draw a contradiction using the identities (5-2). Observe that $[k_0]_{\mathbf{p}} \in S_\alpha$. By definition of k_0 , $[k_0]_{\mathbf{p}}$ satisfies

$$([k_0]_{\mathbf{p}})\phi(g_i)\phi(g_j) = ([k_0]_{\mathbf{p}})\mathbf{p}\phi(g_j) = ([k_0 + 1]_{\mathbf{p}})\phi(g_j) = ([k_0 + 1]_{\mathbf{p}})\mathbf{q}$$

which is distinct from

$$([k_0 + 1]_{\mathbf{p}})\mathbf{p} = ([k_0 + 1]_{\mathbf{p}})\phi(g_i) = ([k_0]_{\mathbf{p}})\mathbf{q}\phi(g_i) = ([k_0]_{\mathbf{p}})\phi(g_j)\phi(g_i).$$

So there exists a transposition $\tau \in T_\alpha$ which moves $[k_0]_{\mathbf{p}}$, and hence

$$(5-20) \quad [k_0]_{\mathbf{p}} \in S_\alpha.$$

Moreover the condition (5-16) implies that $[k]_{\mathbf{p}} \notin S_{\alpha}$ for all $k < k_0$, i.e., k_0 is the smallest number in

$$(5-21) \quad N_{\mathbf{p}} = \{k \in \mathbb{Z} : [k]_{\mathbf{p}} \in S_{\alpha}\}.$$

The identities in (5-2) imply that $[\beta_k, \phi(g_j)] = 1$ for all $k \geq k_0 + 2$ where β_k is the conjugation defined by

$$(5-22) \quad \beta_k := \phi(\alpha)^{\phi(g_i)^{k-k_0}}.$$

To complete the proof we show that $\phi(g_j)$ does not commute with β_k under the assumption $[k]_{\mathbf{p}} \in \text{supp } \mathbf{q}$ for some $k \geq k_0 + 2$. Our claim follows basically from that the involution $\beta_k \in \text{FSym}_n$ intersects $\mathbf{q}$. Since the cycle decomposition of β_k contains $\tau_k := \tau^{\phi(g_i)^{k-k_0}}$, β_k intersects $\mathbf{q}$ at the point

$$([k_0]_{\mathbf{p}})\phi(g_i)^{k-k_0} = ([k_0]_{\mathbf{p}})\mathbf{p}^{k-k_0} = [k_0 + (k - k_0)]_{\mathbf{p}} = [k]_{\mathbf{p}}$$

by the observation (5-11).

If $\text{supp } \tau_k \subset \text{supp } \mathbf{q}$ then we can apply a similar argument used in Corollary 4.3 which shows that an infinite order element g does not commute with a transposition β if $\text{supp } \beta \subset \text{Esupp } g$. Identify $\text{supp } \mathbf{q}$ with $\mathbb{Z}$ with an appropriate base point so that $[k]_{\mathbf{p}} = [k]_{\mathbf{q}}$. From $[k]_{\mathbf{q}} \in \text{supp } \tau_k$ and the minimality of $k_0 \in N_{\mathbf{p}}$ we need to consider two cases: $\tau_k = ([k]_{\mathbf{q}}, [k+1]_{\mathbf{q}})$ or $\tau_k = ([k]_{\mathbf{q}}, [k']_{\mathbf{q}})$ with $k' \geq k+2$. For the first case we check that $([k]_{\mathbf{q}})\beta_k\phi(g_j) \neq ([k]_{\mathbf{q}})\phi(g_j)\beta_k$ by

$$\begin{aligned} [k]_{\mathbf{q}} &\xrightarrow{\beta_k} ([k]_{\mathbf{q}})\tau_k = [k+1]_{\mathbf{q}} \xrightarrow{\phi(g_j)} ([k+1]_{\mathbf{q}})\mathbf{q} = [k+2]_{\mathbf{q}}, \\ [k]_{\mathbf{q}} &\xrightarrow{\phi(g_j)} ([k]_{\mathbf{q}})\mathbf{q} = [k+1]_{\mathbf{q}} \xrightarrow{\beta_k} ([k+1]_{\mathbf{q}})\tau_k = [k]_{\mathbf{q}}. \end{aligned}$$

So $\text{supp } [\beta_k, \phi(g_j)] \ni [k]_{\mathbf{q}}$ is nontrivial, and hence $[\beta_k, \phi(g_j)] \neq 1$. In case τ_k exchanges $[k]_{\mathbf{p}}$ and $[k']_{\mathbf{q}}$ with $k' \geq k+2$ we check $[\beta_k, \phi(g_j)]$ applied to $[k]_{\mathbf{q}}$:

$$[k]_{\mathbf{q}} \xrightarrow{\beta_k} ([k]_{\mathbf{q}})\tau_k = [k']_{\mathbf{q}} \xrightarrow{\phi(g_j)} ([k']_{\mathbf{q}})\mathbf{q} = [k'+1]_{\mathbf{p}} \xrightarrow{\beta_k^{-1}} ([k'+1]_{\mathbf{q}})\beta_k^{-1} \xrightarrow{\phi(g_j)^{-1}} ([k'+1]_{\mathbf{q}})\beta_k^{-1}\phi(g_j)^{-1}.$$

If β_k commutes with $\phi(g_j)$ then we have to have $([k'+1]_{\mathbf{q}})\beta_k^{-1}\phi(g_j)^{-1} = [k]_{\mathbf{q}}$ or equivalently

$$([k'+1]_{\mathbf{q}})\beta_k = ([k]_{\mathbf{q}})\phi(g_j) = [k+1]_{\mathbf{q}}.$$

In other words $[\beta_k, \phi(g_j)] = 1$ implies that β_k contains a transposition exchanging $[k+1]_{\mathbf{q}}$ and $[k'+1]_{\mathbf{q}}$. Inductively one applies the same argument to show that β_k contains infinitely many transpositions in its cycle decomposition. Consequently $\text{supp } \beta_k$ is an infinite set provided $[\beta_k, \phi(g_j)] = 1$. However every element of FSym_n must have a finite support.

We can apply an analogous argument when τ_k exchanges $[k]_{\mathbf{q}}$ and $P \notin \text{supp } \mathbf{q}$. Our claim is that β_k has an infinite support if it commutes with $\phi(g_j)$. From $\beta_k\phi(g_j) = \phi(g_j)\beta_k$ and

$$\begin{aligned} [k]_{\mathbf{q}} &\xrightarrow{\beta_k} ([k]_{\mathbf{q}})\tau_k = P \xrightarrow{\phi(g_j)} (P)\phi(g_j), \\ [k]_{\mathbf{p}} &\xrightarrow{\phi(g_j)} ([k+1]_{\mathbf{q}}) \xrightarrow{\beta_k} ([k+1]_{\mathbf{q}})\beta_k \end{aligned}$$

we see that $(P)\phi(g_j) = ([k+1]_{\mathbf{q}})\beta_k$. Since $\text{supp } \mathbf{q}$ is an $\phi(g_j)$ -invariant subset of $\text{supp } \phi(g_j)$, $(P)\phi(g_j) \notin \text{supp } \mathbf{q}$. In particular $[k+1]_{\mathbf{q}} \in \text{supp } \beta_k$. By induction argument as before, we can show that $\text{supp } \beta_k$ is an infinite set, which is a contradiction.

So far we have found a partial translation $\mathbf{q} \in \mathcal{P}_j$ and $k_0 \in \mathbb{Z}$ with the properties (5-16) and (5-17). For the uniqueness of $\mathbf{q}$ suppose $\mathbf{p}$ intersects $\mathbf{q}' \in \mathcal{P}_j$ with $\mathbf{q}' \neq \mathbf{q}$. Since $\mathbf{q}$ does not intersect $\mathbf{q}'$, if $[k]_{\mathbf{p}} \in \text{supp } \mathbf{q}'$ then $k \geq k_0 + 2$ by (5-16). One can apply the same argument as above to show that β_k defined in (5-22) does not commute with $\mathbf{q}'$, which contradicts (5-2).

We remark that the smallest number $k_0 \in N_{\mathbf{p}}$ does not depend on j . This follows from the relation $\alpha = [g_i, g_j]$ of (5-1) which implies that $\phi(\alpha) = \phi([g_i, g_{j'}]) = \phi([g_i, g_j])$ for pairwise distinct

i, j and j' in $\{2, \dots, n\}$. The minimality of k_0 in $N_{\mathbf{p}}$ implies that $\phi(g_{j'})$ determines a unique partial translation $\mathbf{q}_{j'} \in \mathcal{P}_{j'}$ which agrees with $\mathbf{p}$ on $\{[k]_{\mathbf{p}} : k \leq k_0\}$ and that $([k_0]_{\mathbf{p}})\mathbf{q}_{j'}^2 \neq ([k_0]_{\mathbf{p}})\mathbf{p}^2$. In other words k_0 satisfies identities (5–16) and (5–17) where $\mathbf{q}$ is replaced by $\mathbf{q}_{j'}$.

Step 3. In this step we verify the second part of the Lemma. For $2 \leq i \leq n$, let $\mathfrak{T}_i$ denote the set of all targets of $\phi(g_i)$, i.e., $\phi(g_i)$ has a positive translation length on a ray $R_k \in \mathfrak{T}_i$. If $\mathfrak{T}_i$ and $\mathfrak{T}_j$ share a ray R_k then $\phi(g_i)$ and $\phi(g_j)$ have positive translation lengths on R_k . Using the (positive) least common multiple of the two translation lengths one can find $\mathbf{p} \in \mathcal{P}_i$ and $\mathbf{q} \in \mathcal{P}_j$ such that $\text{supp } \mathbf{p} \cap \text{supp } \mathbf{q}$ contains infinitely many points on the target R_k . However this pair of partial translations fail to satisfy the conditions (5–16) and (5–17) of Step 2. Therefore $\phi(g_i)$ and $\phi(g_j)$ can not share any ray in their targets provided $i \neq j$.

Step 4. From Step 1, we know that there exists a ray R_l which is a source of $\phi(g_i)$ for all $2 \leq i \leq n$. So each target $\mathfrak{T}_i$ is a subset of $\{R_1, \dots, R_n\} \setminus \{R_l\}$. The only way for the targets $\mathfrak{T}_2, \dots, \mathfrak{T}_n$ to be pairwise disjoint is when they are all distinct singleton sets. A monomorphism ϕ defines a bijection $\gamma : \{2, \dots, n\} \rightarrow \{1, \dots, \hat{l}, \dots, n\}$ such that

$$(5-23) \quad \mathfrak{T}_i = \{R_{\gamma(i)}\}$$

for $2 \leq i \leq n$. Now it is obvious that R_l is only ray on which $\phi(g_i)$ has a negative translation length for all i . Consequently ϕ determines a unique ray R_l which is a common source of $\phi(g_i)$ for all i . $\square$

As we checked in Step 2 in the proof of Lemma 5.7, if two partial translations $\mathbf{p} \in \mathcal{P}_i$ and $\mathbf{q} \in \mathcal{P}_j$ ($i \neq j$) intersect then there exists a unique point $[k_0]_{\mathbf{p}}$ satisfying conditions (5–16) and (5–17). Intuitively those two conditions can be interpreted as (1) for each $\mathbf{p} \in \mathcal{P}_i$ there exists $\mathbf{q} \in \mathcal{P}_j$ such that $\mathbf{p}$ and $\mathbf{q}$ are identical all the way up to $[k_0]_{\mathbf{p}}$, (2) once $\mathbf{p}$ develops a different orbit from $[k_0 + 1]_{\mathbf{p}}$, $\mathbf{p}$ never intersects $\mathbf{q}$ again. Compare the above with the characterization (1) and (2) of the generators of $\mathcal{H}_n$ in page 12. With this intuition, let us call the point $[k_0 + 1]_{\mathbf{p}}$ the *diverging point* of $\mathbf{p}$ and denote it by $D_{\mathbf{p}}$. For all partial translations in $\mathcal{P} = \cup_{i=2}^n \mathcal{P}_i$ we label those supports with $\mathbb{Z}$ appropriately such that

$$(5-24) \quad D_{\mathbf{p}} = [0]_{\mathbf{p}}$$

for each $\mathbf{p} \in \mathcal{P}$. From now on let us use the above default labeling for $\text{supp } \mathbf{p}$ for all $\mathbf{p} \in \mathcal{P}$.

Lemma 5.8. *Let $i \in \{2, \dots, n\}$. Each partial translation $\mathbf{p}$ of $\phi(g_i)$ intersects exactly one transposition $\tau_{\mathbf{p}} \in T_i$ which exchanges $[-1]_{\mathbf{p}}$ and $[0]_{\mathbf{p}}$.*

Proof. Any transposition $\tau \in T_i$ is one of the three types:

- Type I: τ exchanges two consecutive points of one partial transposition of $\mathcal{P}_i$.
- Type II: τ exchanges two non-consecutive points of one partial transposition of $\mathcal{P}_i$.
- Type III: τ intersects two distinct partial transpositions of $\mathcal{P}_i$.

We show that each $\mathbf{p} \in \mathcal{P}_i$ can intersect only one transposition $\tau \in T_i$ of type I. From (5–20) we see that $k_0 = -1$ is indeed the smallest integer in $N_{\mathbf{p}}$ defined in (5–21). We first focus on the case $n \geq 3$.

Claim I: If $\mathbf{p} \in \mathcal{P}_i$ intersects τ of type I, then $\mathbf{p}$ does not intersect any other transposition of T_i . The first case to consider is when τ exchanges $[-1]_{\mathbf{p}}$ and $[0]_{\mathbf{p}}$ where -1 is the smallest number in $N_{\mathbf{p}}$. Suppose $\mathbf{p}$ intersects $\tau' \in T_{\alpha}$ of any type whose support contains $[q]_{\mathbf{p}}$. Since τ and τ' does not intersect, $q \geq 1$.

In case $q = 1$, we use the relation r_2 of (5–1) and the hexagon argument described in Remark 5.5. Two involutions $\phi(\alpha)$ and $\beta := \phi(\alpha)^{\phi(g_i)}$ satisfy

$$(\phi(\alpha)\beta)^3 = 1.$$

Observe that β contains a transposition $\tau^{\phi(g_i)} = \tau^p = ([0]_p, [1]_p)$ which intersects two transpositions τ and τ' of $\phi(\alpha)$. Therefore there exists a hexagon on 6 points

$$[-1]_p \xrightarrow{\tau} [0]_p \xrightarrow{\tau^p} [1]_p \xrightarrow{\tau'} Q \xrightarrow{\beta} (Q)\beta \xrightarrow{\phi(\alpha)} S \xrightarrow{\beta} [-1]_p$$

where $Q = ([1]_p)\tau'$ and $S = ([1]_p)\beta$. In particular, $\text{supp } \beta$ contains $[-1]_p$. This means that S_α contains $[-2]_p$, a contradiction to the minimality of $-1 \in N_p$.

Next, we may assume $q \geq 2$ and q is the smallest number in $N_p \setminus \{-1, 0\}$. Consider the conjugation defined by

$$\beta = \phi(\alpha)^{\phi(g_i)^q}.$$

Since $q \geq 2$ the identity (5-3) says that $\phi(\alpha)$ and β commute. Two involutions $\phi(\alpha)$ and β move $[q]_p$ to distinct points since

$$[q]_p\phi(\alpha) = ([q]_p)\tau' \neq [q-1]_p = ([q]_p)\beta = ([q]_p)\tau^{p^q} = [q-1]_p$$

where the inequality follows from the minimality of q in $N_p \setminus \{-1, 0\}$. We are in good position to apply the ‘square argument’ described in Remark 5.3. The square on $I = S_\alpha \cap \text{supp } \beta$ involving 4 points

$$[q-1]_p \xrightarrow{\beta} [q]_p \xrightarrow{\tau'} ([q]_p)\tau' \xrightarrow{\beta} ([q-1]_p)\phi(\alpha) \xrightarrow{\phi(\alpha)} [q-1]_p$$

implies that $[q-1]_p$ must belong to $N_p \setminus \{-1, 0\}$. Again, this contradicts to the minimality of q .

Claim II: If $p \in \mathcal{P}_i$ intersects $\tau \in T_i$ then τ can not be of type II. First we argue that $\text{supp } \tau$ does not contain $[-1]_p$ if τ is of type II. Assume $\tau = ([-1]_p, [q]_p)$ for some $q \geq 1$. The involution β defined by

$$\beta = \phi(\alpha)^{\phi(g_i)^{q+1}}$$

contains $\tau^{p^{q+1}}$ in its cycle decomposition. So β moves $[q]_p$ to $[q + (q+1)]_p$ but $\phi(\alpha)$ moves $[q]_p$ to $[q - (q+1)]_p = [-1]_p$. Therefore we can apply the square argument to see that $[-1]_p$ belongs to $\text{supp } \beta$, or equivalently $[-1 - (q+1)]_p \in N_p$ as in the proof of Claim I. However, the minimality of -1 in N_p says that this can not happen.

To complete the proof, we need to rule out the case when p intersects $\tau = ([q]_p, [r]_p)$ for some $q \geq 0$ and $r \geq q+2$. Recall that -1 is the smallest number in N_p , and so T_i contains a transposition τ' which moves $[-1]_p$. From the discussion so far we may assume that τ' is of type III. Consequently the point $[-1]_p$ gets mapped to distinct points by $\phi(\alpha)$ and β defined by

$$\beta = \phi(\alpha)^{\phi(g_i)^{-(r+1)}}$$

since $([-1]_p)\beta = [-1 - (r - q)]_p \in \text{supp } p$ but $([-1]_p)\phi(\alpha) = ([-1]_p)\tau' \notin \text{supp } p$. The square argument applied to two involutions $\phi(\alpha)$ and β implies that the point $[-1 - (r - q)]_p$ belongs to N_p , which can not happen.

Claim III: If p intersects $\tau \in T_i$ of type III then τ fixes the point $[-1]_p$ where -1 is the smallest number in N_p . Observe that Claim III together with Claim II implies that the transposition τ_p of T_i , which moves $[-1]_p$, must be of type I. Consequently p intersects only transposition τ_p which exchanges $[-1]_p$ and $[0]_p$, completing the proof for the Lemma.

To this end we draw a contradiction from the assumption that p intersects $\tau \in T_i$ of type III with $[-1]_p \in \text{supp } \tau$. Since τ is of type III, it relates p to a partial translation $p_1 \in \mathcal{P}_i$ that τ intersects. From Step II of Lemma 5.7 we know that there exists a smallest number in

$$N_{p_1} = \{k \in \mathbb{Z} : [k]_{p_1} \in S_\alpha\}.$$

By the convention 5-24 we can further say that the smallest number is -1 (after pre-composing an appropriate translation on $\mathbb{Z}$ to $\mathbb{Z} \hookrightarrow \text{supp } p_1$ if necessary). We can show that if $([-1]_p)\tau = [k_1]_{p_1}$ then $k_1 \geq 0$. If $k_1 = -1$, we have to have

$$([-1]_p)pq = ([-1]_p)\phi(g_i)\phi(g_j) = ([-1]_{p_1})\phi(g_j)\phi(g_i) = ([-1]_{p_1})q_1p_1$$

where $\mathbf{q}$ and $\mathbf{q}_1$ are unique partial translations of $\mathcal{P}_j$ which intersect $\mathbf{p}$ and $\mathbf{p}_1$ respectively. Consequently $\mathbf{q}$ intersects both of $\mathbf{p}$ and $\mathbf{p}_1$, a contradiction to the uniqueness of $\mathbf{q}$. Therefore τ fixes $[-1]_{\mathbf{p}_1}$ and so $k_1 \geq 0$.

The partial translation $\mathbf{p}_1$ intersects τ of type III and so Claim I implies that $\mathbf{p}_1$ can only intersect transpositions of type III. In particular the transposition $\tau_1 \in T_i$ with $[-1]_{\mathbf{p}_1} \in \text{supp } \tau_1$ must be of type III. Observe that τ_1 relates $\mathbf{p}_1$ to $\mathbf{p}_2 \in \mathcal{P}_i \setminus \{\mathbf{p}_1\}$ as τ relates $\mathbf{p}$ to $\mathbf{p}_1$. As argued above one can check that τ_1 intersects $\mathbf{p}_2$ at a point $[k_2]_{\mathbf{p}_2}$ with $k_2 > -1$ where -1 is the smallest number in $N_{\mathbf{p}_2} = \{k \in \mathbb{Z} : [k]_{\mathbf{p}_2} \in S_\alpha\}$. Inductively we can consider a cycle of partial translations

$$(5-25) \quad \mathbf{p}_0 = \mathbf{p} \xrightarrow{\tau_0=\tau} \mathbf{p}_1 \xrightarrow{\tau_1} \cdots \xrightarrow{\tau_{m-1}} \mathbf{p}_m \xrightarrow{\tau_m} \mathbf{p}_0.$$

Let $k_1, \dots, k_m$ be integers so that $\tau_r = ([-1]_{\mathbf{p}_r}, [k_{r+1}]_{\mathbf{p}_{r+1}})$ for $0 \leq r \leq m-1$ and $\tau_m = ([-1]_{\mathbf{p}_m}, [k_m]_{\mathbf{p}_0})$.

Next we show that $k_r = 0$ for $1 \leq r \leq m$. Assume the contrary that, for example, $k_1 \geq 1$. We can apply the square argument to two commuting involutions $\phi(\alpha)$ and β defined by

$$\beta = \phi(\alpha)^{\phi(g_i)^{k_1+1}}$$

since $([k_1]_{\mathbf{p}_1})\phi(\alpha) = [-1]_{\mathbf{p}_0} \neq [k_2 + k_1 + 1]_{\mathbf{p}_2} = ([k_1]_{\mathbf{p}_1})\beta$. Consequently $[-1]_{\mathbf{p}_0} \in \text{supp } \beta$ or equivalently $-1 - (k_1 + 1) \in N_{\mathbf{p}_0}$, contradicting to the minimality of $-1 \in N_{\mathbf{p}_0}$. A similar argument shows that $k_r = 0$ for all r .

So far we have shown that if $\mathbf{p}$ intersects $\tau \in T_i$ of type III then there exists a cycle of partial translations (5-25) which are related by transpositions $\tau_r = ([-1]_{\mathbf{p}_r}, [0]_{\mathbf{p}_{r+1}})$ for $0 \leq r \leq m-1$ and $\tau_m = ([-1]_{\mathbf{p}_m}, [0]_{\mathbf{p}_0})$. Next we show that $\phi(g_i)$ restricts on the set $C_1 = \{[1]_{\mathbf{q}_0}, \dots, [1]_{\mathbf{q}_m}\}$ to define a $(m+1)$ -cycle

$$\sigma_1 : [1]_{\mathbf{q}_m} \rightarrow [1]_{\mathbf{q}_{m-1}} \rightarrow \cdots \rightarrow [1]_{\mathbf{q}_0} \rightarrow [1]_{\mathbf{q}_m}.$$

For each $0 \leq r \leq m-1$, the transposition $\tau_r = ([-1]_{\mathbf{p}_r}, [0]_{\mathbf{p}_{r+1}})$ of $[\phi(g_i), \phi(g_j)]$ implies that

$$[-1]_{\mathbf{p}_r} \xrightarrow{\phi(g_i)} [0]_{\mathbf{p}_r} = [0]_{\mathbf{q}_r} \xrightarrow{\phi(g_j)} [1]_{\mathbf{q}_r} \xrightarrow{\phi(g_i)^{-1}} ([1]_{\mathbf{q}_r})\phi(g_i)^{-1} \xrightarrow{\phi(g_j)^{-1}} [0]_{\mathbf{q}_{r+1}}.$$

From the last arrow we have $([1]_{\mathbf{q}_r})\phi(g_i)^{-1} = [1]_{\mathbf{q}_{r+1}}$, or $([1]_{\mathbf{q}_{r+1}})\phi(g_i) = [1]_{\mathbf{q}_r}$ for $0 \leq r \leq m-1$. For $r = m$ it is direct to check that $[1]_{\mathbf{q}_0}\phi(g_i) = [1]_{\mathbf{q}_m}$. Therefore $\phi(g_i)$ restricted on the set C_1 is σ_1 .

Finally we show that $\phi(g_i)$ contains infinitely many copies of σ_1 in its cycle decomposition. Observe that the conjugation $\phi(\alpha)^{\phi(g_j)^2}$ contains a factor β_2 given by

$$(5-26) \quad \beta_2 = (\tau_0\tau_1 \cdots \tau_m)^{\phi(g_j)^2}.$$

The action of $\phi(g_j)^2$ on $\text{supp } \beta$ is the translation by $+2$ along partial translations $\mathbf{q}_0, \mathbf{q}_1, \dots, \mathbf{q}_m$. So β_2 can be written as a product of transpositions

$$([1]_{\mathbf{q}_0}, [2]_{\mathbf{q}_1}) \cdots ([1]_{\mathbf{q}_{m-1}}, [2]_{\mathbf{q}_m})([1]_{\mathbf{q}_m}, [2]_{\mathbf{q}_0})$$

Therefore the conjugation $(\sigma_1)^{\beta_2}$ is the $(m+1)$ -cycle

$$\sigma_2 : [2]_{\mathbf{q}_m} \rightarrow [2]_{\mathbf{q}_{m-1}} \rightarrow \cdots \rightarrow [2]_{\mathbf{q}_0} \rightarrow [2]_{\mathbf{q}_m}.$$

Intuitively speaking, taking the conjugation of σ_1 by β_2 results in translating the $(m+1)$ -cycle σ_1 globally by $+1$ along the partial translations $\mathbf{q}_0, \dots, \mathbf{q}_m$. Now the identity (5-4) says that $\phi(g_i)$ commutes with $\phi(\alpha)^{\phi(g_j)^2}$. In particular $\phi(g_i)^{\beta_2} = \phi(g_i)$. This means that $\phi(g_i)$ already contains σ_2 , a translated copy of σ_1 , in its cycle decomposition. Applying identity (5-4) repeatedly to $\phi(g_i)$ and $\phi(\alpha)^{\phi(g_j)^k}$, we see that $\phi(g_i)^{\beta_k} = \phi(g_i)$ for $k \geq 2$ where β_k is defined in an analogous way as in (5-26) with the power k instead of 2. Thus $\phi(g_i)$ contains infinitely many copies of σ_2 in its cycle decomposition, which is absurd. In all, a partial translation $\mathbf{p}$ which intersects $\tau \in T_i$ of type III forces that $\phi(g_i) \notin \mathcal{H}_n$. Claim III is verified and so we are done.

If $n = 2$, we can show the same result for a partial translation of $\phi(g_2)$ as a special case of the above discussion. The braid relation and commutation relation of (5–7) can be used for Claim 1 and Claim 2 respectively. One can verify Claim 3 immediately with braid relation again. $\square$

Corollary 5.9. *Let $3 \leq n$. Suppose that a partial translation $\mathbf{p}$ of $\phi(g_i)$ intersects a partial translation $\mathbf{q}$ of $\phi(g_j)$. Then $\mathbf{p}$ does not intersect any other cycles of $\phi(g_j)$ but $\mathbf{q}$.*

Proof. We first check that $[1]_{\mathbf{p}}$ is fixed under $\phi(g_j)$. Lemma 5.8 states that the unique transposition $\tau_{\mathbf{p}}$, which intersects $\mathbf{p}$, exchanges $[-1]_{\mathbf{p}}$ and $[0]_{\mathbf{p}}$. From the identity $([0]_{\mathbf{p}})[\phi(g_i), \phi(g_j)] = ([0]_{\mathbf{p}})\tau_{\mathbf{p}} = [-1]_{\mathbf{p}}$ we have $([0]_{\mathbf{p}})\phi(g_i)\phi(g_j) = ([-1]_{\mathbf{p}})\phi(g_i)\phi(g_j)$, and so

$$\begin{aligned} ([1]_{\mathbf{p}})\phi(g_j) &= ([0]_{\mathbf{p}})\mathbf{p}\phi(g_j) = ([0]_{\mathbf{p}})\phi(g_i)\phi(g_j) = ([-1]_{\mathbf{p}})\phi(g_j)\phi(g_i) \\ &= ([-1]_{\mathbf{p}})\mathbf{q}\phi(g_i) = ([0]_{\mathbf{p}})\phi(g_i) = ([0]_{\mathbf{p}})\mathbf{p} = [1]_{\mathbf{p}} \end{aligned}$$

since $([-1]_{\mathbf{p}})\mathbf{q} = ([-1]_{\mathbf{p}})\mathbf{p} = [0]_{\mathbf{p}}$ by (5–16). Suppose $\mathbf{p}$ intersects a component $f \neq \mathbf{q}$ in the cycle decomposition of $\phi(g_j)$. We may further assume that if k_1 is the smallest integer in $\{k \in \mathbb{Z} : [k]_{\mathbf{p}} \in \text{supp } f\}$ then $[k]_{\mathbf{p}}$ is fixed under $\phi(g_j)$ for all $1 \leq k \leq k_1 - 1$. To show $[k_1 - 1]_{\mathbf{p}} \in \text{supp } \phi(\alpha)$ we check that

$$[k_1 - 1]_{\mathbf{p}} \xrightarrow{\phi(g_i)} ([k_1 - 1]_{\mathbf{p}})\mathbf{p} = [k_1]_{\mathbf{p}} \xrightarrow{\phi(g_j)} ([k_1]_{\mathbf{p}})\phi(g_j) = ([k_1]_{\mathbf{p}})f$$

which is distinct from

$$[k_1 - 1]_{\mathbf{p}} \xrightarrow{\phi(g_j)} ([k_1 - 1]_{\mathbf{p}})\phi(g_j) = [k_1 - 1]_{\mathbf{p}} \xrightarrow{\phi(g_i)} ([k_1 - 1]_{\mathbf{p}})\mathbf{p} = [k_1]_{\mathbf{p}}.$$

Since $k_1 - 1 > 0$, $\phi(\alpha)$ contains a transposition $\beta \neq \tau_{\mathbf{p}}$ which intersects $\mathbf{p}$. However this contradicts Lemma 5.8. $\square$

The unique positive integer ℓ in Lemma 5.7 is called *the eventual length* of a monomorphism ϕ , and denoted by $\ell(\phi)$. Lemma 5.7 implies that $\mathcal{P}_i$ contains precisely $\ell = \ell(\phi)$ partial translations $\mathbf{p}_1, \dots, \mathbf{p}_{\ell}$ for each $2 \leq i \leq n$. Since they are only infinite cycles of $\phi(g_i)$ we have $\text{Esupp } \phi(g_i) = \bigcup_{j=1}^{\ell} \text{supp } \mathbf{p}_j$. Lemma 5.7 also implies that each ray of X_n is either a source or a target of $\phi(g_i)$ for some i . So the set $\bigcup_{i=2}^n \text{Esupp } \phi(g_i)$ contains all points of X_n but finitely many. Let $\text{ES}(\phi) \subset X_n$ be the set

$$\text{ES}(\phi) = \bigcup_{g \in \mathcal{H}_n} \text{Esupp } \phi(g).$$

Proposition 5.10. *With the notation defined above, $\text{ES}(\phi) = \bigcup_{i=2}^n \text{Esupp } \phi(g_i)$.*

Proof. One side inclusion is clear. For the reverse inclusion we claim that $F = X_n \setminus \bigcup_{i=2}^n \text{Esupp } \phi(g_i)$ is invariant under $\phi(g)$ for all $g \in \mathcal{H}_n$. Note that $P \in F$ if and only if P has a finite orbit under $\phi(g_i)$ for all $2 \leq i \leq n$. In other words each $\phi(g_i)$ contains a finite cycle which moves P . We argue by induction on the word length of g . If a finite cycle f of $\phi(g_i)$ moves P then obviously $(P)\phi(g_i) = (P)f$ has the same finite orbit under $\phi(g_i)$. If $(P)\phi(g_i)$ is fixed under $\phi(g_j)$ for some j then we have nothing to prove because the orbit under $\phi(g_j)$ is trivial. To show $(P)\phi(g_i) \in F$ for every i , we need to check that if $\phi(g_i)$ has a finite cycle f such that $(P)f \in \text{supp } \phi(g_j)$ for some $j \neq i$, then $\phi(g_j)$ also has a finite cycle which moves $(P)\phi(g_i)$. This follows from the last assertion of Corollary 5.9 which implies that if f intersects $\phi(g_j)$ then it can only intersect finite cycles of $\phi(g_j)$ for all $j \neq i$. The same argument shows that if $(P)\phi(g_i)^{-1}$ belongs to $\text{supp } \phi(g_j)$ then it has a finite orbit under $\phi(g_j)$ for every $j \neq i$. So $(P)\phi(g_i)^{\pm 1} \in F$. This establishes the base case.

Suppose $g \in \mathcal{H}_n$ with $|g| = k + 1$. Say the last letter of g is $g_i^{\pm 1}$, i.e., $g = g' \cdot g_i^{\pm 1}$ with $|g'| = k$. By induction assumption $(P)\phi(g') \in F$. So, for each i , if $(P)\phi(g') \in \text{supp } \phi(g_i)$ then there exists a finite cycle of $\phi(g_i)$ which moves the point $(P)\phi(g')$. A similar argument applied to $(P)\phi(g')\phi(g_i)^{\pm 1}$, instead of $(P)\phi(g_i)^{\pm 1}$ as in the base case, shows that the point $(P)\phi(g')\phi(g_i)^{\pm 1}$ has a finite orbit under $\phi(g_j)$ for all $2 \leq j \leq n$. Therefore $(P)\phi(g)\phi(g_i)^{\pm 1} \in F$, and hence F is an invariant set under

$\phi(g)$ for all $g \in \mathcal{H}_n$. Since F is finite $P \in F$ must have a finite orbit under $\phi(g)$. This means that $P \notin \text{ES}(\phi)$. $\square$

Consider the decomposition of $X_n = \text{ES}(\phi) \sqcup F$ where F is the finite set as in the above proof. Let $E(\phi(g))$ denote the restriction of $\phi(g)$ on the set $\text{ES}(\phi)$. From the decomposition we have $E(\phi(g))E(\phi(h)) = E(\phi(gh))$ for all $g, h \in \mathcal{H}_n$. One crucial observation is that, for each $2 \leq i \leq n$, $E(\phi(g_i))$ is nothing but the product of its commuting partial translations;

$$(5-27) \quad E(\phi(g_i)) = \prod_{\mathbf{p} \in \mathcal{P}_i} \mathbf{p}.$$

Consequently we have

$$(5-28) \quad \text{Esupp}(\phi(g_i)) = \bigcup_{\mathbf{p} \in \mathcal{P}_i} \text{supp } \mathbf{p}.$$

So Lemma 5.8 implies that T_i consists of ℓ transpositions each of which exchanges $[-1]_{\mathbf{p}}$ and $[0]_{\mathbf{p}}$ for some $\mathbf{p} \in \mathcal{P}_i$. Two conditions (5-16) and (5-17) show that $T_i = T_j$ for each pair $2 \leq i \neq j \leq n$. It follows from that $E(\phi(\alpha))$ is the product of ℓ transpositions in T_i for any i . Indeed we have the following.

Corollary 5.11. *Let D be the set of all diverging points of partial translations in $\mathcal{P} = \bigcup_{i=2}^n \mathcal{P}_i$. There exists bijections*

$$\mathcal{P}_i \leftrightarrow D \leftrightarrow T_i$$

for each $2 \leq i \leq n$.

Proof. Fix i and consider the sequence of maps

$$\mathcal{P}_i \xrightarrow{\iota_1} D \xrightarrow{\iota_2} T_i \xrightarrow{\iota_3} \mathcal{P}_i$$

defined by

$$\iota_1 : \mathbf{p} \mapsto [0]_{\mathbf{p}}, \quad \iota_2 : [0]_{\mathbf{p}} \mapsto \tau_{\mathbf{p}}, \quad \iota_3 : \tau_{\mathbf{p}} \mapsto \mathbf{p}.$$

We claim that all three maps are injective whose composition in a row yields the identity. Partial translations of $\mathcal{P}_i$ do not intersect each other, which explains ι_1 is injective. Every diverging point is given by $[0]_{\mathbf{q}}$ for some $\mathbf{q} \in \mathcal{P}_j$. If $j = i$, we have nothing to show that ι_2 is well-defined due to Lemma 5.8. If not, from Step 2 of Lemma 5.7 one can find a partial translation $\mathbf{p} \in \mathcal{P}_i$ with conditions (5-16) and (5-17). So we have $[0]_{\mathbf{q}} = [0]_{\mathbf{p}}$. The assignment $[0]_{\mathbf{p}} \mapsto \tau_{\mathbf{p}}$ is well-defined and injective due to Lemma 5.8. For ι_3 it suffices to check if $\tau \in T_i$ then τ intersects a unique partial translation in $\mathcal{P}_i$. By definition, if $\tau \in T_i$ then τ intersects $\text{Esupp } \phi(g_i)$. The union (5-28) forces τ to intersect at least one partial translation of $\mathcal{P}_i$. By Lemma 5.8 again, ι_3 is well-defined and injective. It follows immediately from definitions of three maps that the composition is the identity map. We have established bijections as desired. $\square$

Expanding map $\tilde{\phi}$. Recall that $\mathcal{H}_n$ acts on the underlying set $X_n = \{1, \dots, n\} \times \mathbb{N}$ transitively. So X_n can be considered as a single orbit of a base point $(1, 1)$. Let us to express points $(j, m) \in X_n$ ($1 \leq j \leq n, m \in \mathbb{N}$) as

$$(j, m) = \begin{cases} (1, 1)g_2^{-(m-1)} & \text{if } j = 1, \\ (1, 1)g_j^m & \text{if } 2 \leq j \leq n. \end{cases}$$

In case $(j, m) \in R_1$ we simply choose g_2^{-1} among inverses of $n-1$ generators whose actions coincide on R_1 as translations by $+1$. By taking g_2^{-1} as a default translation on R_1 we can make the

above expression unique. Take subscripts and exponents of the generators to obtain the following coordinate system for $(j, m) \in X_n$

$$(5-29) \quad (j, m) \leftrightarrow \begin{cases} [2, -(m-1)] & \text{if } j = 1, \\ [j, m] & \text{if } 2 \leq j \leq n. \end{cases}$$

Corollary 5.11 implies that a monomorphism ϕ determines $\ell = \ell(\phi)$ diverging points $D_1, \dots, D_\ell$. Let $\mathcal{Q}_l = \{\mathbf{p} \in \mathcal{P} : [0]_{\mathbf{p}} = D_l\}$ for $1 \leq l \leq \ell$. For each $2 \leq i \leq n$, $\mathcal{P}_i$ contains precisely one partial translation $\mathbf{p}$ such that $[0]_{\mathbf{p}} = D_l$. Let $\mathbf{p}_{l,i}$ denote such a unique partial translation $\mathbf{p} \in \mathcal{P}_i$, i.e., $\mathbf{p}_{l,i} = \iota_1^{-1}(D_l)$ where $\iota_1 : \mathcal{P}_i \rightarrow D$ is the bijection defined in proof of Corollary 5.11. We label partial translations of $\mathcal{Q}_l$ as

$$\mathcal{Q}_l = \{\mathbf{p}_{l,2}, \dots, \mathbf{p}_{l,n}\}$$

With this new labeling, $\mathcal{P}_i = \{\mathbf{p}_{l,i} : 1 \leq l \leq \ell\}$ for each $2 \leq i \leq n$, and the set of all partial translations $\mathcal{P}$ determined by ϕ has decompositions $\mathcal{P} = \sqcup_{i=2}^n \mathcal{P}_i = \sqcup_{l=1}^\ell \mathcal{Q}_l$. The last identity follows from the equivalence relation: partial translations $\mathbf{p}, \mathbf{q} \in \mathcal{P}$ belong to $\mathcal{Q}_l$ for some l if and only if $\mathbf{p}$ intersects $\mathbf{q}$. For better notation, let $[m]_{l,i}$ denote the point $[m]_{\mathbf{p}_{l,i}}$ from now on. Let $\mathcal{O}(D_l) \subset \text{ES}(\phi)$ denote the set

$$\mathcal{O}(D_l) = \bigcup_{2 \leq i \leq n} \text{supp } \mathbf{p}_{l,i}.$$

Since $D_l = [0]_{l,i}$ for all $2 \leq i \leq n$, every point in $\mathcal{O}(D_l)$ can be written as $[m]_{l,i} = (D_l)\mathbf{p}_{l,i}^m$ for some $\mathbf{p}_{l,i} \in \mathcal{Q}_l$ and $m \in \mathbb{Z}$. Note that this expression is not unique when $m \leq 0$. Each pair of partial translations of $\mathcal{Q}_l$ satisfies conditions (5-16) and (5-17). The first condition implies that the point $[m]_{l,i}$ with $m \leq 0$ and $3 \leq i \leq n$ is identified to $[m]_{l,2}$. The second condition implies that $[m]_{l,i} = [m]_{l,i'}$ if and only if $i = i'$ when $m \geq 1$. So every point $P \in \mathcal{O}(D_l)$ can be expressed uniquely as

$$(5-30) \quad P = \begin{cases} [m]_{l,2} & m \leq 0, \\ [m]_{l,i} & 2 \leq i \leq n, 1 \leq m. \end{cases}$$

Comparing (5-29) and (5-30) we obtain a canonical bijection $\tilde{\phi}_l : X_n \rightarrow \mathcal{O}(D_l)$, for each $l = 1, \dots, \ell$, defined by

$$(5-31) \quad ([i, m])\tilde{\phi}_l = [m]_{l,i}.$$

The expanding map $\tilde{\phi} : X_n \rightarrow (X_n)^\ell$ is defined by

$$\tilde{\phi} = \tilde{\phi}_1 \times \dots \times \tilde{\phi}_\ell.$$

Note that $\tilde{\phi}$ depends on the choice of diverging points. However components of $\tilde{\phi}$ are all distinct. This follows from that $\mathcal{O}(D_l)$ does not intersect $\mathcal{O}(D_{l'})$ if $l \neq l'$.

Remark 5.12. A monomorphism ϕ of $\mathcal{H}_n$ determines a group $G_l \leq \text{Sym}_n$ which consists of restrictions $g \in \phi(\mathcal{H}_n)$ on the orbit of D_l under $\phi(\mathcal{H}_n)$, $1 \leq l \leq \ell$. We remark that $G_l \cong \mathcal{H}_n$ for all l . Observe that each $\mathcal{O}(D_l)$ coincides with the orbit of D_l . This is because all partial translations in $\mathcal{P} \setminus \mathcal{Q}_l$ fix D_l . Indeed G_l is generated by $n-1$ partial translations of $\mathcal{Q}_l$. The map $\tilde{\phi}_l$ conjugates $\mathcal{H}_n$ to G_l in the ambient group Sym_n .

Proposition 5.13. Suppose $\tau = (P, Q)$ is the transposition exchanging P and Q . Then $E(\phi(\tau))$ coincides with a product of ℓ commuting transpositions $\prod_{j=1}^\ell ((P)\tilde{\phi}_j, (Q)\tilde{\phi}_j)$.

Proof. Assume $n \geq 3$. We consider the following four cases depending on which rays P and Q lie.
Case I. Suppose both P and Q lie on R_1 . Let $P = [2, m] = (1, -m + 1)$ and $Q = [2, m'] = (1, -m' + 1)$ with $m < m' \leq 0$. From the definition of $\tilde{\phi}$ we have

$$(P)\tilde{\phi}_l = ([2, m])\tilde{\phi}_l = [m]_{l,2}, \text{ and } (Q)\tilde{\phi}_l = ([2, m'])\tilde{\phi}_l = [m']_{l,2}$$

for each l . Since components of $\tilde{\phi}$ are all distinct, $\prod_{l=1}^{\ell} ((P)\tilde{\phi}_l, (Q)\tilde{\phi}_l) = \prod_{l=1}^{\ell} ([m]_{l,2}, [m']_{l,2})$ is a product of ℓ commuting transpositions.

On the other hand, one can check that τ can be written as $\tau = \alpha^h$ where

$$h = g_2 g_3^{m-m'+1} g_2^{m'-1}.$$

So $E(\phi(\tau)) = E(\phi(\alpha^h)) = E(\phi(\alpha))^{E(\phi(h))} = E(\phi(\alpha))^{\phi(h)}$. Recall that $\phi(g_2)$ contains partial translations $\mathbf{p}_{1,2}, \dots, \mathbf{p}_{\ell,2}$ with $D_l = [0]_{l,2}$ for $l = 1, \dots, \ell$. So the involution $E(\phi(\alpha))$ is the product of ℓ transpositions $\tau_1, \dots, \tau_{\ell}$ where τ_l exchanges $[0]_{l,2}$ and $[-1]_{l,2}$. We examine the effect of $E(\phi(h))$ on those points. Since $\phi(g_3)$ fixes $[m]_{l,2}$ for all $m \geq 1$, we have, for each l ,

$$[0]_{l,2} \xrightarrow{\phi(g_2)} ([0]_{l,2})\mathbf{p}_{l,2} = [1]_{l,2} \xrightarrow{\phi(g_3^{m-m'+1})} [1]_{l,2} \xrightarrow{\phi(g_2^{m'-1})} ([1]_{l,2})\mathbf{p}_{l,2}^{m'-1} = [m']_{l,2},$$

$$[-1]_{l,2} \xrightarrow{\phi(g_2)} [0]_{l,2} \xrightarrow{\phi(g_3^{m-m'+1})} ([0]_{l,2})\mathbf{q}_{l,3}^{m-m'+1} = [m-m'+1]_{l,2} \xrightarrow{\phi(g_2^{m'-1})} ([m-m'+1]_{l,2})\mathbf{p}_{l,2}^{m'-1} = [m]_{l,2}$$

where $\mathbf{q}_{l,3}$ is the partial translation of $\phi(g_3)$ such that $\mathbf{q}_{l,3} = \mathbf{p}_{l,2}$ on $\{[m]_{l,2} : m \leq -1\}$. Therefore $E(\phi(\tau))$ is the product of ℓ transpositions exchanging $[m']_{l,2}$ and $[m]_{l,2}$, $l = 1, \dots, \ell$, as expected.

To complete the proof, one can repeat similar calculation for $E(\phi(h))$ in the following cases:

Case II. $P = [2, m]$ and $Q = [i, m']$, with $m \leq 0$, $m' \geq 1$. Then $\tau = \alpha^h$ where

$$h = \begin{cases} g_2 g_3 g_2^{m'-1} g_3^{m-1} & i = 2 \\ g_2 g_i^{m'} g_2^{m-1} & 3 \leq i \leq n. \end{cases}$$

Case III. $P = [i, m]$ and $Q = [i, m']$ with $1 \leq m' < m$. Then $\tau = \alpha^h$ where

$$h = g_2 g_3^{-(m-m'-1)} g_2^{-1} g_i^m.$$

Case IV. $P = [i, m]$ and $Q = [j, m']$ with $i \neq j$. Then $\tau = \alpha^h$ where

$$h = g_i g_j g_i^{m-1} g_j^{m'-1}.$$

If $n = 2$ we can apply similar argument with $\alpha = ((1, 1), (1, 2))$. Suppose $P = [2, m]$ and $Q = [2, m']$ with $1 \leq m < m'$. The transposition $\tau = (P, Q)$ can be written as a conjugation $\tau = \alpha^h$ where

$$h = (g_2 \alpha)^{m'-m-1} g_2^{m+1}.$$

Since $\phi(\alpha)$ exchanges $[-1]_{l,2}$ and $[0]_{l,2}$, but fixes all other points for each l we have

$$\begin{aligned} [0]_{l,2} &\xrightarrow{\phi(g_2 \alpha)} [1]_{l,2} \xrightarrow{\phi(g_2 \alpha)} ([2]_{l,2}) \rightarrow \dots \rightarrow [m' - m - 1]_{l,2} \xrightarrow{\phi(g_2)^{m+1}} [m']_{l,2}, \\ [-1]_{l,2} &\xrightarrow{\phi(g_2 \alpha)} [-1]_{l,2} \rightarrow \dots \rightarrow [-1]_{l,2} \xrightarrow{\phi(g_2)^{m+1}} [m]_{l,2} \end{aligned}$$

Therefore $E(\phi(\tau)) = \prod_{l=1}^{\ell} ([m]_{l,2}, [m']_{l,2}) = \prod_{l=1}^{\ell} ((P)\tilde{\phi}_l, (Q)\tilde{\phi}_l)$. The other cases can be taken care of by similar calculation because a transposition of $\mathcal{H}_2$ can be expressed as a conjugation of α by $(g_2 \alpha)^{m_1} g_2^{m_2}$ for some m_1, m_2 . $\square$

Lemma 5.14. Suppose that $\phi(g_i) = g_i f_i$ for all $2 \leq i \leq n$ where $f_i \in \text{FSym}_n$. If τ is a transposition, there exists a constant A_2 , which do not depend on k , such that $|\phi^k(\tau)| \leq A_2 k$ for all $k \in \mathbb{N}$.

Proof. By the observation (5–27), each $\phi(g_i)$ is nothing but a partial translation $\mathbf{p}_i$. So $\mathbf{p}_i$ acts as a translation by -1 on R_1 and by 1 on R_i up to a finite set for $2 \leq i \leq n$. We first show that $|\phi^k(\tau)|$ for a transposition $\tau = (P, Q)$. By Proposition 5.13 we see that

$$\phi^k(\tau) = ((P)\tilde{\phi}^k, (Q)\tilde{\phi}^k)$$

where $\tilde{\phi} = \tilde{\phi}_1$ is the expanding map $X_n \rightarrow \text{ES}(\phi)$ defined in (5–31). We claim that both $(P)\tilde{\phi}^k$ and $(Q)\tilde{\phi}^k$ belong to $B_{n,r}$ with $r \leq m + ks$ for all $k \in \mathbb{N}$ for some constants m and s which are determined by ϕ . Let $[p]_i$ denote the point $[p]_{\mathbf{p}_i}$. For each $2 \leq i$ one can find the smallest integer $0 < k_i$ such that

$$[p]_i = (i, m_i + p - k_i)$$

for all $k_i \leq p$ where $[k_i]_i = (i, m_i)$. We take the largest number $k_1 \leq 0$ so that

$$[p]_2 = (1, m_1 + k_1 - p)$$

for all $p \leq k_1$ where $[k_1]_2 = (1, m_1)$. Note that k_1 is well defined since the actions of $\phi(g_2)^{-1}, \dots, \phi(g_n)^{-1}$ are identical on $\{[p]_2 : p \leq 0\}$. Let $B_{n,m}$ denote the ball of X_n with radius $m = \max\{m_1, \dots, m_n\}$. Since each $\phi(g_i)$ acts as a translation by ± 1 on each ray outside $B_{n,m}$, $P \in B_{n,s}$ implies that $(P)\tilde{\phi} \in B_{n,s+1}$. Observe that if $P = (i, p) \notin B_{n,m}$ then $(P)\tilde{\phi} \in B_{n,p+s}$ where $s = \max\{|m_2 - k_2|, \dots, |m_n - k_n|, |m_1 + k_1 - 1|\}$. One can check that if $k_i \leq p$ then, by the coordinate system (5–29),

$$([i, p])\tilde{\phi} = [p]_i = (i, m_i + p - k_i) = [i, p + (m_i - k_i)]$$

for each $2 \leq i$. Similarly if $p \leq k_1$ then

$$([2, p])\tilde{\phi} = [p]_2 = (1, m_1 + k_1 - p) = [2, p - (m_1 + k_1 - 1)]$$

by (5–29). Thus our claim is verified, and so the transposition $\phi^k(\tau) = ((P)\tilde{\phi}^k, (Q)\tilde{\phi}^k)$ has support in $B_{n,r}$ with $r \leq m + ks$ for all $k \in \mathbb{N}$. It is not difficult to show that a transposition τ_0 with $\text{supp } \tau_0 \subset B_{n,r}$ has length $< 10r$. Observe that τ_0 can be written as a conjugation $\tau_0 = \alpha^h$ where $h \in \mathcal{H}_n$ can be taken so that $2|h| + |\alpha| < 10r$ as in the proof of Proposition 5.13. Therefore we have

$$|\phi^k(\tau)| = |((P)\tilde{\phi}^k, (Q)\tilde{\phi}^k)| < 10(m + ks) < 10(m + s)k = A_2k$$

for all k . □

From now on we assume that a monomorphism ϕ with $2 \leq \ell(\phi)$ satisfies that $\phi(g_i)$ has a source R_1 and a target R_i for all $2 \leq i \leq n$. (In view of Proposition 5.24 this assumption is legitimate) We also assume that $P \in \text{ES}(\phi)$ unless otherwise stated.

Rooted trees induced by $\tilde{\phi}$. The iteration of $\tilde{\phi}$ applied to P determines a *labeled rooted ℓ -ary tree*, which we will denote by $\mathcal{T}_P$. The vertex set V_P of $\mathcal{T}_P$ is equipped with *label*, *level* and *ℓ -ary sequences*. The root of $\mathcal{T}_P$ is the unique vertex at level 0, labeled by P , which corresponds to the empty sequence. Inductively the label $L : V_P \rightarrow \text{ES}(\phi)$ and corresponding sequence $W : V_P \rightarrow \Omega_\ell$ are defined such that a vertex $v \in V_P$ at level $k \in \mathbb{N}$ is labeled by

$$L(v) = (P)\tilde{\phi}_{l_1}\tilde{\phi}_{l_2} \cdots \tilde{\phi}_{l_k}$$

and corresponds to a sequence

$$W(v) = l_1 l_2 \cdots l_k$$

where Ω_ℓ consists of ℓ -ary sequences on $\{1, \dots, \ell\}$ and $1 \leq l_j \leq \ell$ for all $1 \leq j \leq k$. Note that $W(v)$ can be realized as the unique edge path from the root to v with edge labels $\{1, \dots, \ell\}$. Consequently we have a bijective correspondence between V_P^k , the set of vertices at level k , and $\Omega_{\ell,k}$, the set of all ℓ -ary sequences of length k .

Example 5.15. Consider a monomorphism ϕ of $\mathcal{H}_3$ defined by $\phi(g_i) = (g_i)^2 \cdot f_i$ for $i = 2, 3$ where f_2 and f_3 are cycles given by

$$f_2 : (1, 1) \rightarrow (1, 2) \rightarrow (2, 1) \rightarrow (1, 1), \quad f_3 : (1, 1) \rightarrow (1, 2) \rightarrow (2, 1) \rightarrow (3, 2) \rightarrow (1, 1)$$

From direct computation we see that $\phi(\alpha) = E(\phi(\alpha))$ is the product of two transpositions τ_1 and τ_2 where

$$\tau_1 : ((2, 1), (1, 4)), \quad \tau_2 : ((1, 2), (1, 3)).$$

Set $D_1 = (2, 1)$ and $D_2 = (1, 2)$. Each of $\phi(g_2)$ and $\phi(g_3)$ has $\ell(\phi) = 2$ partial translations which are


$$p_{1,2} : \cdots \rightarrow (1, 2m+4) \rightarrow (1, 2m+2) \rightarrow \cdots \rightarrow (1, 4) \rightarrow (2, 1) \rightarrow \cdots \rightarrow (2, 2m-1) \rightarrow (2, 2m+1) \rightarrow \cdots$$

$$p_{2,2} : \cdots \rightarrow (1, 2m+3) \rightarrow (1, 2m+1) \rightarrow \cdots \rightarrow (1, 3) \rightarrow (1, 2) \rightarrow (1, 1) \rightarrow \cdots \rightarrow (2, 2m) \rightarrow (2, 2m+2) \rightarrow \cdots$$

$$p_{1,3} : \cdots \rightarrow (1, 2m+4) \rightarrow (1, 2m+2) \rightarrow \cdots \rightarrow (1, 4) \rightarrow (2, 1) \rightarrow \cdots \rightarrow (3, 2m) \rightarrow (3, 2m+2) \rightarrow \cdots$$

$$p_{2,3} : \cdots \rightarrow (1, 2m+3) \rightarrow (1, 2m+1) \rightarrow \cdots \rightarrow (1, 3) \rightarrow (1, 2) \rightarrow \cdots \rightarrow (3, 2m-1) \rightarrow (3, 2m+1) \rightarrow \cdots$$

where $m \in \mathbb{N}$. With $D_l = [0]_{l,2} = [0]_{l,3}$ for $l = 1, 2$, let us apply $\tilde{\phi}$ repeatedly to describe rooted trees of points in $\{(1, 1), (1, 2), (2, 1)\} = \text{supp } f_2$.



The above illustrates the tree $\mathcal{T}_{(1,1)}$ with labels up to level 3. Each vertex v has $\ell = 2$ children; the left child is $(v)\tilde{\phi}_1$ and the right child is $(v)\tilde{\phi}_2$. Accordingly all left edges are labelled by 1 ($\tilde{\phi}_1$) and all right edges are labeled by 2 ($\tilde{\phi}_2$). Each $v \in V_{(1,1)}^k$ corresponds to a unique sequence $W(v) \in \Omega_{2,k}$. For example the vertex $[2, 0]$ at level 2 corresponds to the sequence $12 \in \Omega_{2,2}$. One important observation is that $\mathcal{T}_{(1,1)}$ contains infinitely many copies of itself. The point $(1, 1)$ appears as a label for all vertex v with $W(v) = 1212 \cdots 12$ (as emphasized in blue color!) because the subtree spanned by v and its descendants is identical to $\mathcal{T}_{(1,1)}$.

The following figure illustrates the tree $\mathcal{T}_{(1,2)}$ up to level 3, which happens to be a subtree of $\mathcal{T}_{(1,1)}$. Observe that $\mathcal{T}_{(1,2)}$ never contains a copy of itself. This is because the vertices of the tree are all distinct. Yet another crucial fact is that each pair of children has labels which are *translations* of the labels $(1, 4)$ and $(1, 3)$ at level 1. For example, $\mathcal{T}_{(1,2)}$ has four pairs of labels at level 3, each of which is a translation of the pair $(1, 4)$ and $(1, 3)$ by some power of g_2 . Intuitively this is because $p_{1,2}$ and $p_{2,2}$ act as a translation by ± 2 on all points in their supports but finitely many. More precisely


$$([2, m])\tilde{\phi}_1 = [m]_{1,2} = \begin{cases} [2, 2m-1] = (1, -2m+2) & m \leq -1 \\ [2, 2m+1] = (2, 2m+1) & 1 \leq m \end{cases},$$

$$([2, m])\tilde{\phi}_2 = [m]_{2,2} = \begin{cases} [2, 2m] = (1, -2m + 1) & m \leq -1 \\ [2, 2m - 2] = (2, 2m - 2) & 2 \leq m \end{cases}$$


where the last equalities in each case follow from

$$[2, m] = \begin{cases} (1, -m + 1) & m \leq 0 \\ (2, m) & 1 \leq m \end{cases}$$

by (5–29). See Proposition 5.19.



The tree $\mathcal{T}_{(2,1)}$ below also contains infinitely many copies of itself. The subtrees determined by vertices with sequences $2121 \cdots 21$ are identical to $\mathcal{T}_{(2,1)}$. On the other hand, the subtree $\mathcal{T}_{(2,3)}$ illustrates the opposite behavior; it does not contain a copy of itself. Moreover any children at level $k \geq 2$ in $\mathcal{T}_{(2,3)}$ has labels which are translations of the labels $(2, 7)$ and $(2, 4)$.



For a vertex $v \in V_P$ with $\omega = W(v)$, let P_ω denote the label $L(v)$. From Proposition 5.13 we have the following.

Corollary 5.16. *Suppose $\tau = (P, Q)$ is the transposition on two points $P, Q \in \text{ES}(\phi)$. Then $E(\phi^k(\tau))$ coincides with a product of ℓ^k commuting transpositions $\prod_{\omega \in \Omega_{\ell,k}} (P_\omega, Q_\omega)$ for each $k \in \mathbb{N}$.*

Proof. The base case follows directly from Proposition 5.13 since $\Omega_{\ell,1} = \{1, \dots, \ell\}$. Suppose that $E(\phi^k(\tau)) = \prod_{\omega \in \Omega_{\ell,k}} \tau_\omega$ is a product of ℓ^k commuting transpositions for $k \geq 1$ where $\tau_\omega = (P_\omega, Q_\omega)$.

By Proposition 5.13 again, we have

$$\mathbb{E}(\phi(\tau_\omega)) = \prod_{j=1}^{\ell} \left((P_\omega) \tilde{\phi}_j, (Q_\omega) \tilde{\phi}_j \right)$$

for all $\omega \in \Omega_{\ell,k}$. So $\mathbb{E}(\phi(\tau_\omega)) = \prod_{\omega'} (P_{\omega'}, Q_{\omega'})$ where $\omega' = \omega l \in \Omega_{\ell,k+1}$ for $l = 1, \dots, \ell$. Therefore

$$\begin{aligned} \mathbb{E}(\phi^{k+1}(\tau)) &= \mathbb{E}(\phi \phi^k(\tau)) = \mathbb{E}\left(\phi \left(\prod_{\omega \in \Omega_{\ell,k}} \tau_\omega \right)\right) = \prod_{\omega \in \Omega_{\ell,k}} \left(\mathbb{E}(\phi(\tau_\omega)) \right) \\ &= \prod_{\omega \in \Omega_{\ell,k}} \prod_{j=1}^{\ell} \left((P_\omega) \tilde{\phi}_j, (Q_\omega) \tilde{\phi}_j \right) = \prod_{\omega \in \Omega_{\ell,k}} \prod_{\substack{\omega' = \omega l \\ 1 \leq l \leq \ell}} (P_{\omega'}, Q_{\omega'}) = \prod_{\omega \in \Omega_{\ell,k+1}} (P_\omega, Q_\omega). \end{aligned}$$

The commutativity of transpositions above follows from Proposition 5.17. $\square$

Recall the notation V_P^k which denotes the set of all vertices of $\mathcal{T}_P$ at level k .

Proposition 5.17. *The label map $L : V_P^k \rightarrow \text{ES}(\phi)$ is injective for each $k \in \mathbb{N}$.*

Proof. Induction on k . The decomposition $\text{ES}(\phi) = \sqcup_{l=1}^{\ell} \mathcal{O}(D_l)$ shows that if $l \neq l'$, $(X_n) \tilde{\phi}_l = \mathcal{O}(D_l)$ does not intersect $(X_n) \tilde{\phi}_{l'} = \mathcal{O}(D_{l'})$. This establishes the base case. Suppose $v \in V_P^{k+1}$. Observe that the last letter of $W(v)$ determines which $\mathcal{O}(D_l)$ the label $L(v)$ belongs to. More precisely, $L(v) \in \mathcal{O}(D_l)$ if and only if the last letter of $W(v)$ is l . If $L(v) = L(v') = Q \in \mathcal{O}(D_l)$ for vertices v and v' at level $k+1$, then $W(v)$ and $W(v')$ share the same last letter l , that is, $W(v) = \eta l$ and $W(v') = \eta' l$ for some $\eta, \eta' \in \Omega_{\ell,k}$. Since $\tilde{\phi}_l : X_n \rightarrow \mathcal{O}(D_l)$ is a bijection, the two vertices which correspond to η and η' respectively share the same label $(Q) \tilde{\phi}_l^{-1}$. However this contradicts induction assumption. $\square$

Stable points. In Example 5.15 we saw that the trees $\mathcal{T}_{(2,1)}$ and $\mathcal{T}_{(1,1)}$ contain copies of themselves. However the tree $\mathcal{T}_{(1,2)}$ illustrates the opposite behavior. Intuitively this is because each $\tilde{\phi}_l$ applied to (i, k) ‘doubles’ the second coordinate for all points of its support but finitely many. We need to formulate this rigorously.

Let $\ell \geq 2$. We want to define *stable points* for $\phi(g_i)$ with the assumption that $\phi(g_i)$ has a unique target R_i for $i = 2, \dots, n$. Fix i . Recall that $\phi(g_i)$ translates points of R_i by $+\ell$ up to a finite set. In other words, the action of $\phi(g_i)$ on R_i is *eventually stabilized* as a translation by $+\ell$. Being a restriction of $\phi(g_i)$, $\mathbf{p}_{l,i}$ is also eventually stabilized for all $1 \leq l \leq \ell$. From property (5–12) one can take an integer $0 \leq k_{l,i}$ for each l such that if $[k]_{l,i} = (i, m)$ then $[k+1]_{l,i} = (i, m+\ell)$ for all $k_{l,i} \leq k$. More precisely, for each l , there exists a smallest integer $k_{l,i}$ such that

$$(5-32) \quad [k]_{l,i} = (i, m_{l,i} + \ell(k - k_{l,i}))$$

for all $k_{l,i} \leq k$ where $[k_{l,i}]_{l,i} = (i, m_{l,i})$. The i^{th} *threshold* is the positive integer s_i defined by

$$(5-33) \quad s_i = \max_{1 \leq l \leq \ell} \left\{ \left\lceil \frac{\ell k_{l,i} - m_{l,i}}{\ell - 1} \right\rceil, k_{l,i} \right\}$$

where $\lceil * \rceil$ stands for the smallest integer function. A point (i, k) is called a *stable point* if $s_i < k$ where $2 \leq i \leq n$. Recall the coordinate system (5–29); $[i, k] = (i, k)$ for $2 \leq i \leq n$ and $k \in \mathbb{N}$. One crucial observation is that if $[i, k]$ is a stable point, $([i, k]) \tilde{\phi}_l$ is again a stable point for all l . A stable point $[i, k]$ satisfies $k_{l,i} < k$ for all l . By (5–32) we have

$$(5-34) \quad ([i, k]) \tilde{\phi}_l = [k]_{l,i} = (i, m_{l,i} + \ell(k - k_{l,i})).$$

for all $k > s_i$. Since

$$k < m_{l,i} + \ell(k - k_{l,i}) \Leftrightarrow \frac{\ell k_{l,i} - m_{l,i}}{\ell - 1} < k,$$

$s_i < k < m_{l,i} + \ell(k - k_{l,i})$ provided $s_i < k$. So $([i, k])\tilde{\phi}_l$ is a stable point if $[i, k]$ is a stable point for $2 \leq i \leq n$.

We can define stable points on R_1 in a similar manner because all the actions of $\phi(g_i)$'s on R_1 are also eventually stabilized as a translation by $-\ell$. However we need to change signs and take reverse inequalities accordingly. It suffices to consider one partial translation, say $\mathbf{p}_{l,2}$, of $\phi(g_2)$ since actions of $\mathbf{p}_{l,2}^{-1}$ and $\mathbf{p}_{l,i}^{-1}$ are identical on $\{[k]_{l,2} : k \leq 0\}$ for $3 \leq i \leq n$. There exists a largest integer $k_l \leq 0$ such that

$$(5-35) \quad [k]_{l,2} = (1, m_l + \ell(k_l - k))$$

for all $k \leq k_l$ where $[k_{l,2}]_l = (1, m_l)$. Now points $(1, k) \in R_1$ are called *stable points* for all $s_1 < k$ where s_1 is the positive integer defined by

$$(5-36) \quad s_1 = \max \left\{ \left\lceil \frac{\ell - \ell k_l - m_l}{\ell - 1} \right\rceil, -k_l + 1 \right\}$$

The above definition implies that the image of $(1, k)$ under $\tilde{\phi}_l$ is a stable point if $(i, k) = [2, -k + 1]$ (by the coordinate system(5-29)) is a stable point. Since $s_1 < k \Leftrightarrow -k + 1 < -s_1 + 1 < k_l$ we have, by (5-35),

$$([2, -k + 1])\tilde{\phi}_l = [-k + 1]_{l,2} = (1, m_l + \ell(k_l - (-k + 1))) = (1, m_l + \ell(k_l + k - 1))$$

for all stable points on R_1 . So we have $s_1 < k < m_l + \ell(k_l + k - 1)$ from

$$k < m_l + \ell(k_l + k - 1) \Leftrightarrow \frac{\ell - \ell k_l - m_l}{\ell - 1} < k.$$

Let $\mathcal{S}$ denote the set of all stable points. We summarize the discussion above as

Remark 5.18. If $P = (i, p)$ is a stable point then $(P)\tilde{\phi}_l = (i, q)$ with $p < q$ for each $1 \leq l \leq \ell$, $1 \leq i \leq n$. In particular, $\mathcal{S}$ is invariant under $\tilde{\phi}_l$ for each l .

Translations on a ray. Let $v \in V_P$ with $\omega = W(v)$. The *descendants* of v at a depth $m \in \mathbb{N}$ consists of vertices u such that $W(u) = \omega\omega_1$ for some $\omega_1 \in \Omega_{\ell,m}$. Let $D_{v,m}$ denote the set of descendants of v with depth m , and let $D_v = \cup_m D_{v,m}$. The *children* of v is the descendants of v at depth 1. Note that $V_P^k = \sqcup_{v \in V_P^{k-1}} D_{v,1}$ for all k . In case an ordered ℓ -tuple $L(D_{v,1}) = ((i, p_1), \dots, (i, p_\ell))$ we suppress the first coordinate and take a vector $\nu = [p_1, \dots, p_\ell] \in \mathbb{N}^\ell$ to express $L(D_{v,1})$. The *translation* of an ℓ -tuple ν by $t \in \mathbb{Z}$ is the ℓ -tuple

$$\nu + t = [p_1 + t, \dots, p_\ell + t].$$

We require that all points of ν and its translation $\nu + t$ stay in the same ray R_i . Let $B_{n,r}$ denote the ball of X_n with radius r ,

$$B_{n,r} = \{(i, p) | p \leq r\}.$$

Recall that $\phi(g_i)$ acts as a translation on $\{(i, p) | p > s_i\}$ and $\{(1, q) | q > s_1\}$ by ℓ and $-\ell$ respectively for all $2 \leq i \leq n$ where s_i is the i^{th} threshold defined in (5-33) and (5-36). Consequently $\phi(g_i)$ acts by the same manner on $R_i \setminus B_{n,s}$ and $R_1 \setminus B_{n,s}$ for

$$s = \max\{s_1, \dots, s_n\}.$$

Proposition 5.19. Suppose $P \in \mathcal{S} \cap R_i$, $1 \leq i \leq n$. The label map $L : V_P \rightarrow \text{ES}(\phi)$ is injective and $L(V_P) \subset \mathcal{S} \cap R_i$. For each $v \in V_P$, $L(D_{v,1})$ is a translation of $L(V_P^1)$. Moreover there exists a constant $A_0 = A_0(P)$ such that $L(V_P^k) \subset B_{n,r}$ with $r = A_0\ell^k + s$

Proof. First we consider the case when $P = (i, p)$ is a stable point for $2 \leq i$. By Remark 5.18, we have $L(v) = (i, q)$ with $p < q$ for all $v \in V_P$, and so $L(V_P) \subset \mathcal{S} \cap R_i$. Suppressing i in the first coordinate we can write $L(V_P^1) = [p_1, \dots, p_\ell]$ where

$$(i, p_l) = ([i, p])\tilde{\phi}_l = [p]_{l,i} = (i, m_{l,i} + \ell(p - k_{l,i}))$$

for $l = 1, \dots, \ell$ by (5-34). The label map L restricted on the root vertex and V_P^1 is injective since $p < p_l$ for all l and each p_l belongs to $\mathcal{O}(D_l)$ which are all disjoint for $l = 1, \dots, \ell$. With this base case assume that the map L is injective on the set of vertices up to level k . Since no vertex at level $k+1$ attains the label $(i, p) = P$ it suffices to check whether two vertices v and v' (other than the root vertex) at levels $\leq k+1$ share the same label $L(v) = L(v')$. We can apply similar argument as in Proposition 5.17 to draw a contradiction; $L(v) = L(v')$ implies that the ascendants of v and v' share the same label at levels $\leq k$.

For the second assertion suppose $v \in V_P$ with $L(v) = (i, q)$. Since $(1, q)$ is a stable point the identity (5-32) implies that $L(D_{v,1}) = [q_1, \dots, q_\ell]$ where

$$(i, q_l) = \left(i, m_{l,i} + \ell(q - k_{l,i})\right) = \left(i, m_{l,i} + \ell(p - k_{l,i}) + \ell(q - p)\right)$$

for $l = 1, \dots, \ell$. Since $p < q$

$$(5-37) \quad L(D_{v,1}) = L(V_P^1) + \ell(q - p)$$

is the translation of $L(V_P^1)$ by $\ell(q - p) > 0$.

Since $L(v) = (i, q)$ for all $v \in V_P^k$, we can consider a sequence of natural numbers $\{a_k\}$ so that a_k denotes the maximum of such q 's. We want to show $a_k \leq s\ell^k + s$ for all k . From the rewriting

$$(5-38) \quad m_{l,i} + \ell(p - k_{l,i}) = \ell\left(p - \frac{\ell k_{l,i} - m_{l,i}}{\ell - 1}\right) + \frac{\ell k_{l,i} - m_{l,i}}{\ell - 1}$$

we have $a_1 \leq A_0\ell + s$ where

$$A_0 = \max_{1 \leq l \leq \ell} \left\{ p - \frac{\ell k_{l,i} - m_{l,i}}{\ell - 1} \right\}.$$

(since $P = (i, p)$ is a stable point the above maximum is taken over positive numbers, and so $A_0 > 0$.) One can rewrite $a_{k+1} = m_{l,i} + \ell(a_k - k_{l,i})$, which follows from the second step above, as in (5-38) with p replaced by a_k to check that

$$(5-39) \quad a_{k+1} - s = \ell(a_k - s)$$

for all $k \in \mathbb{N}$. Therefore $L(V_P^k) \subset B_{n,a_k}$ and $a_k \leq A_0\ell^k + s$ for all $k \in \mathbb{N}$.

In case $P = (1, p)$ one can apply analogous arguments. Remark 5.18 together with ‘cancelling argument’ shows that $L(v) = (1, q)$ is a stable point for all $v \in V_P$. By $(1, q) = [2, -q + 1]$ and (5-35), we have that $((1, q))\tilde{\phi}_l$ becomes

$$([2, -q + 1])\tilde{\phi}_l = [-q + 1]_{l,2} = \left(1, m_l + \ell(k_l + q - 1)\right) = \left(1, m_l + \ell(k_l + p - 1) + \ell(q - p)\right)$$

for all l . Therefore $L(D_{v,1})$ is a translation of $L(V_P^1)$ by $\ell(q - p) > 0$. From the rewriting for the root $P = (1, p) = [2, -p + 1]$,

$$([2, -p + 1])\tilde{\phi}_l = m_l + \ell(k_l + p - 1) = \ell\left(p - \frac{\ell - m_l - \ell k_l}{\ell - 1}\right) + \frac{\ell - m_l - \ell k_l}{\ell - 1}$$

we find $a_1 = A_0\ell + s$ such that $L(V_P^1) \subset B_{n,a_1}$ where

$$A_0 = \max_{1 \leq l \leq \ell} \left\{ p - \frac{\ell - m_l - \ell k_l}{\ell - 1} \right\}.$$

Finally the identity $a_{k+1} = m_l + \ell(k_l + a_k - 1)$ implies that $L(V_P^k) \subset B_{n,a_k}$ where the sequence $\{a_k\}$ satisfies (5-39) for all $k \in \mathbb{N}$. $\square$

Remark 5.20. We remark that the radii of balls that contain $L(V_P^k)$ can be bound by a linear term of ℓ^k even when P is not a stable point. Intuitively this happens because the images of P under expanding map $\tilde{\phi}^k$ travel inside the ball $B_{n,s}$ for first finite steps. Only after does $L(V_P^k)$ contain stable points the radii of the balls contain V_P^k follow the growth as described in Proposition 5.19. More precisely, a ball B_{n,a_k} that contains $L(V_P^k)$ has radius

$$a_k \leq \begin{cases} s & k \leq k_0 \\ A_0 \ell^{k-k_0} + s & k_0 + 1 \leq k \end{cases}$$

for some constant A_0 . In all, every $P \in \text{ES}(\phi)$ determines a constant A_0 such that $L(V_P^k) \subset B_{n,r}$ with $r \leq A_0 \ell^k + s$.

Intervals of V_P^k . We want to decompose V_P^k into *intervals* which provide a coarser decomposition than $V_P^k = \sqcup_{v \in V_P^{k-1}} D_{v,1}$. Motivating examples come from stable points. Proposition 5.19 states that $L(D_{v,1})$ is a translation of $L(V_P^1)$ for all $v \in V_P^{k-1}$ if P is a stable point. The set $\{L(D_{v,1}) : v \in V_P^{k-1}\}$ can be viewed as a single orbit of V_P^1 under translation. In this case, we want to take V_P^k as a single interval.

Let $v \in V_P^{k_1}$ with $k_1 < k$. We say that $D_{v,k-k_1} \subset V_P^k$ is an *interval* of V_P^k if either $v \in V_P^{k-1}$ or $L(v)$ is a stable point. (The naming ‘interval’ comes from that if $D_{v,k-k_1}$ is an interval, $\Lambda(D_{v,k-k_1})$ forms an interval of $\mathbb{N}$ with size ℓ^{k-k_1} where $\Lambda : V_P^k \rightarrow \Omega_{\ell,k} \rightarrow \{1, \dots, \ell^k\}$ is the unique map such that $\Lambda(v) = j$ if $W(v)$ is the j^{th} sequence in the lexicographic order on $\Omega_{\ell,k}$.) Note that an interval may contain smaller intervals. For example, a descendant u of v determines an interval $D_u \cap V_P^k$ if $u \in V_P^{k_2}$ with $k_2 < k$. However we can take maximal intervals so that V_P^k consists of minimum number of intervals. For each $k \in \mathbb{N}$, let $N_P(k)$ denote the minimum number of intervals which cover V_P^k . We remark that V_P^k can be expressed as a disjoint union of $N_P(k)$ intervals.

In view of Proposition 5.19, we say a tree $\mathcal{T}_P$ is *stabilized* if $L : V_P \rightarrow \text{ES}(\phi)$ is injective. Let $\mathcal{U} \subset \text{ES}(\phi)$ denote the set of all points P such that $L : V_P \rightarrow \text{ES}(\phi)$ is not injective. Proposition 5.19 implies that the cardinality of $\mathcal{U}$ is finite since $\mathcal{U} \subset \text{ES}(\phi) \setminus \mathcal{S}$. (example)

Lemma 5.21. *There exists a constant $A_1 = A_1(\phi)$ such that $N_P(k) \leq (k-1)(\ell-1)A_1 + 1$ for all $P \in \text{ES}(\phi)$ and $k \in \mathbb{N}$.*

Proof. Proposition 5.19 implies that $N_P(k) = 1$ for all $P \in \mathcal{S}$. Indeed $N_P(k)$ is bounded by a constant for all $P \notin \mathcal{U}$. If $L : V_P \rightarrow \text{ES}(\phi)$ is injective, one can find $k_0 \in \mathbb{N}$ such that $L(V_P^k)$ consists of stable points for all $k \geq k_0$ since $|\text{ES}(\phi) \setminus \mathcal{S}|$ is finite. From the decomposition

$$V_P^k = \bigsqcup_{u \in V_P^{k_0}} D_{u,k-k_0},$$

where each component is an interval of V_P^k , we have $N_P(k) \leq |V_P^{k_0}| = \ell^{k_0}$ for all $k \geq k_0$. So $N_P(k) \leq \ell^{k_0}$ for all $k \in \mathbb{N}$ since the function $N_P(k)$ is monotone increasing on k . Taking maximum of upper bounds ℓ^{k_0} over all $P \in \text{ES}(\phi) \setminus \mathcal{S} \setminus \mathcal{U}$, which is a finite set, one obtains a constant $A_1 \geq 1$ such that $N_P(k) \leq A_1$ for all $P \in \text{ES}(\phi) \setminus \mathcal{U}$. To establish desired bound for points of $\mathcal{U}$ we need following steps together with induction on k .

Step 1. A vertex v of $\mathcal{T}_P$ and its descendants determine a unique subtree $\mathcal{T}_v$. Note that a pair of subtrees $\mathcal{T}_v$ and $\mathcal{T}_{v'}$ satisfies the following dichotomy: they do not intersect or one contains the other. In this step we show that if $L(v) = L(v')$ then either $\mathcal{T}_v \subset \mathcal{T}_{v'}$ or $\mathcal{T}_{v'} \subset \mathcal{T}_v$.

Suppose $L(v) = L(v')$ for $v \neq v'$. Then $W(v) = \omega l$ and $W(v') = \omega' l$ for some l . As in proof of Proposition 5.17, we can cancel the last same letters of two sequences. We have either ωl is a subword of $\omega' l$ or vice versa since $v \neq v'$. This precisely means that one subtree contains the other.

Step 2. We claim that $P \in \mathcal{U}$ if and only if $\mathcal{T}_P$ contains a vertex v other than the root with $L(v) = P$. We need to show that if the label map $L : \mathcal{T}_P \rightarrow \text{ES}(\phi)$ is not injective then V_P contains v at level $k \geq 1$ with $L(v) = P$. Suppose V_P contains v_1 and v_2 with $L(v_1) = L(v_2) = Q$. From Step 1 we know that one of subtrees contains the other. We may further assume that one of two trees, say $\mathcal{T}_{v_1}$, is maximal, which means that $\mathcal{T}_{v_1}$ contains all subtrees determined by v with $L(v) = Q$. Consequently $\omega_1 = W(v_1)$ is a subsequence of $\omega_2 = W(v_2)$, i.e., $\omega_2 = \omega_1\eta$ for some sequence η . The first case to consider is when $|\omega_1| \leq |\eta|$. Using the same canceling argument as in Step 1, one can show that $\eta = \eta_0\omega_1$ for some sequence η_0 . The sequence $\omega_1 = l_1 \cdots l_k$ determines a composition of injective maps such that $(P)\tilde{\phi}_{l_1} \cdots \tilde{\phi}_{l_k} = Q$. Observe that ω_1 also determines a unique path from the vertex u , which corresponds to $\omega_1\eta_0$, to v_2 . This path transforms into the same composition of maps such that $(L(u))\tilde{\phi}_{l_1} \cdots \tilde{\phi}_{l_k} = L(v_2) = Q$. So u is a vertex with $L(u) = P$. The next case to consider is when $|\omega_1| > |\eta|$. The cancelling argument implies that ω_1 is the concatenation $\omega_1 = \omega_0\eta$ for some sequence ω_0 . Since the sequence $\eta = l'_1 \cdots l'_m$ determines identity map $(Q)\tilde{\phi}_{l'_1} \cdots \tilde{\phi}_{l'_m} = (L(v_1))\tilde{\phi}_{l'_1} \cdots \tilde{\phi}_{l'_m} = L(v_2) = Q$ we have $(L(u))\tilde{\phi}_{l'_1} \cdots \tilde{\phi}_{l'_m} = L(v_1) = Q$ for the vertex u corresponding to ω_0 . So $L(u) = Q$. However this contradicts the maximality of the tree $\mathcal{T}_{v_1}$.

Step 3. Next we show that V_P^1 contains only one vertex v with $L(v) \in \mathcal{U}$. Suppose there exist two vertices v_1 and v_2 at level 1 such that $L(v_i) \in \mathcal{U}$, $i = 1, 2$. By Step 2, $\mathcal{T}_{v_i}$ contains a vertex u_i with $L(u_i) = L(v_i)$, $i = 1, 2$. Suppose $L(v_1) = (P)\tilde{\phi}_l$ for some l . Since $L(u_1) = L(v_1)$ the parent of u_1 has label P . So $\mathcal{T}_{v_1}$ contains a vertex w_1 such that $L(w_1) = P$. By the same reason $\mathcal{T}_{v_2}$ contains a vertex w_2 with $L(w_2) = P$. Since two subtrees $\mathcal{T}_{v_1}$ and $\mathcal{T}_{v_2}$ do not intersect, $\mathcal{T}_P$ contains two disjoint copies of itself which are determined by w_1 and w_2 . Obviously this can not occur by Step 1.

Step 4. We complete the proof by induction on k . The base case is obvious since $N_P^1 = 1$ for any $P \in \text{ES}(\phi)$. From Step 3, we know that $V_P^1 = \{v_1, \dots, v_\ell\}$ contains one vertex v_j with $L(v_j) = Q_j \in \mathcal{U}$. Since all other subtrees $\mathcal{T}_{v_l}$ are stabilized there exists a constant A_1 such that $N_{Q_l}(k-1) \leq A_1$ for all $Q_l = L(v_l)$ and $k \in \mathbb{N}$, $i \neq j$. We complete the proof by

$$N_P(k) \leq \sum_{1 \leq l \leq \ell} A_1 Q_l(k-1) \leq A_1 Q_j(k-1) + (\ell-1)A_1 \leq (k-1)(\ell-1)A_1 + 1.$$

□

Lemma 5.22. Suppose $\tau = (P, Q)$ is the transposition on two points $P, Q \in \text{ES}(\phi)$. There exists a polynomial ρ on $k \in \mathbb{N}$ whose degree does not depend on k such that $|\phi^k(\tau)| \leq \rho(k)\ell^k$ for all $k \in \mathbb{N}$.

Proof. First we show the existence of such a polynomial when P and Q are stable points, and then we extend the discussion to the general case. Suppose $P = (i, p)$ is a stable point. Observe that explicit expressions for $L(v)$ can be obtained for $v \in V_P^k$ whenever $W(v)$ is given. We want to find such a expressions using $L(V_P^1)$. For each $1 \leq l \leq \ell$, let $V_P^1 = \{v_1, \dots, v_\ell\}$ with $L(v_l) = (i, p_l) = (P)\tilde{\phi}_l$, and let $d_l = p_l - p$. We claim that if $W(v) = l_1 l_2 \cdots l_{k+1} \in \Omega_{\ell, k+1}$ then $L(v) = (i, p_v)$ where

$$(5-40) \quad p_v = d_{l_1}\ell^k + d_{l_2}\ell^{k-1} + \cdots + d_{l_k}\ell + p_{l_{k+1}}$$

for each $k \in \mathbb{N}$. The base case for $k = 1$ follows immediately from (5-37). Assume $u \in V_P^{k+2}$ with $W(u) = l_1 l_2 \cdots l_{k+2}$. The unique parent vertex v of u with $W(v) = l_1 l_2 \cdots l_{k+1}$ has label $L(v) = (i, p_v)$ which satisfies (5-40). By the identity (5-37), $L(u) = (i, p_u)$ satisfies

$$p_u = \ell(p_v - p) + p_{l_{k+2}} = \ell(d_{l_1}\ell^k + \cdots + d_{l_k}\ell + p_{l_{k+1}} - p) + p_{l_{k+2}} = d_{l_1}\ell^{k+1} + \cdots + d_{l_{k+1}}\ell + p_{l_{k+2}}$$

since u is the $(l_{k+2})^{\text{th}}$ child of v .

Next we want to describe the relationship between $L(D_{v_l, k})$ and $L(V_P^k)$ for $1 \leq l \leq \ell$ and $k \in \mathbb{N}$. A vertex $v_l \in V_P^1$ determines a set of vertices $D_{v_l, k} \subset V_P^{k+1}$ with $|D_{v_l, k}| = \ell^k = |V_P^k|$. Observe

that each vertex $u \in D_{v_l, k}$ corresponds to $W(u) = l\omega$ for some $\omega \in \Omega_{\ell, k}$. The concatenation $W(v) \mapsto lW(v)$ induces a bijection between V_P^k and $D_{v_l, k}$; $V_P^k \ni v \leftrightarrow u \in D_{v_l, k}$ if $W(u) = lW(v)$. We can further show that $L(D_{v_l, k})$ is a translation of $L(V_P^k)$ by $d_l \ell^k$ for each l and k . Using the expression (5–40) we compare $L(v)$ and $L(u)$ for a corresponding pair under the above bijection. If $v \in V_P^k$ with $W(v) = l_1 \cdots l_k$, the $L(v) = (i, p_v)$ satisfies

$$p_v = d_{l_1} \ell^{k-1} + \cdots + d_{l_{k-1}} \ell + p_{l_k}.$$

The corresponding vertex $u \in D_{v_l, k}$ with $W(u) = l l_1 \cdots l_k$ has label $L(u) = (i, p_u)$ where

$$p_u = d_l \ell^k + d_{l_1} \ell^{k-1} + \cdots + d_{l_{k-1}} \ell + p_{l_k}.$$

Then we have the desired difference $p_u - p_v = d_l \ell^k$. Therefore the above bijection between V_P^k and $D_{v_l, k}$ transforms into a translation of $L(V_P^k)$ by $d_l \ell^k$ for all l and k , i.e., the following diagram commute

$$\begin{array}{ccc} V_P^k & \longrightarrow & D_{v_l, k} \\ \downarrow L & & \downarrow L \\ L(V_P^k) \subset R_i & \xrightarrow{+d_l \ell^k} & L(D_{v_l, k}) \subset R_i \end{array}$$

We are ready to find a polynomial $\rho(k)$ such that $|\phi^k(\tau)| \leq \rho(k) \ell^k$ for all k when $P = (i, p)$ and $Q = (j, q)$ are stable points where $1 \leq i, j \leq n$. Note that we want to make sure that the degree of $\rho(k)$ does not depend on k . Let $V_Q^1 = \{u_1, \dots, u_\ell\}$ with $L(u_l) = (j, q_l) = (Q) \tilde{\phi}_l$, and let $d'_l = q_l - q$ for $1 \leq l \leq \ell$. To deal with the base case $k = 1$, we shall take $\rho(k)$ with $p(1) = |\phi(\tau)|$. Corollary 5.16 implies that

$$E(\phi^{k+1}(\tau)) = \prod_{\omega \in \Omega_{\ell, k+1}} (P_\omega, Q_\omega).$$

To establish desired bounds for $|\phi^{k+1}(\tau)|$ let us regroup those ℓ^{k+1} transpositions into ℓ subcollections as follows. The sets $V_P^1 = \{v_1, \dots, v_\ell\}$ and $V_Q^1 = \{u_1, \dots, u_\ell\}$ provide canonical decompositions

$$V_P^{k+1} = \bigsqcup_{1 \leq l \leq \ell} D_{v_l, k} \text{ and } V_Q^{k+1} = \bigsqcup_{1 \leq l \leq \ell} D_{u_l, k}.$$

Let σ_l be the product

$$\sigma_l = \prod_{\omega \in W(D_{v_l, k})} (P_\omega, Q_\omega) = \prod_{\substack{\omega = l\omega' \\ \omega' \in \Omega_{\ell, k}}} (P_\omega, Q_\omega)$$

for $1 \leq l \leq \ell$. Note that each σ_l is the restriction of $E(\phi^{k+1}(\tau))$ on a $\phi^{k+1}(\tau)$ -invariant subset $L(D_{v_l, k}) \cup L(D_{u_l, k})$. So $E(\phi^{k+1}(\tau)) = \sigma_1 \cdots \sigma_\ell$. Since $L(D_{v_l, k}) \subset R_i$ and $L(D_{u_l, k}) \subset R_j$ are translations of $L(V_P^k)$ and $L(V_Q^k)$ by $d_l \ell^k$ and $d'_l \ell^k$ respectively for each l , σ_l can be expressed as a conjugation of $E(\phi^k(\tau))$. We need to consider three cases depending on i and j .

Case I. $2 \leq i \neq j \leq n$. Define β_l by

$$\beta_l = g_i^{d_l \ell^k} g_j^{d'_l \ell^k}$$

for $1 \leq l \leq \ell$. Then σ_l is the conjugation of $\phi^k(\tau)$ by β_l . Setting $d = 1/4 \max\{d_1, \dots, d_\ell, d'_1, \dots, d'_\ell\}$ we can expect

$$|\sigma_l| = |E(\phi^k(\tau))| + 2|\beta_l| \leq \rho(k) \ell^k + 2(d_l + d'_l) \ell^k \leq (\rho(k) + d) \ell^k$$

for each l . Obviously there exists a polynomial $\rho(k)$ of degree d with the condition $p(1) = |\phi(\tau)|$ such that

$$(5-41) \quad |\mathbb{E}(\phi^{k+1}(\tau))| \leq \sum_{1 \leq \ell} |\sigma_l| = (\rho(k) + d)\ell^{k+1} < \rho(k+1)\ell^{k+1}.$$

For example one can take $\rho(k) = k^d + |\phi(\tau)|$.

Case II. $j = 1, 2 \leq i \leq n$. Take $i' \neq i$ to define β_l as

$$\beta_l = (g_{i'}^{-1} g_i)^{d_l \ell^k} g_{i'}^{-d'_{i'} \ell^k}$$

for $1 \leq l \leq \ell$. Since each σ_l is the conjugation of $\phi^k(\tau)$ by β_l for each l , we have

$$|\sigma_l| = |\mathbb{E}(\phi^k(\tau))| + 2|\beta_l| \leq \rho(k)\ell^k + 2(2d_l + d'_{i'})\ell^k \leq (\rho(k) + d)\ell^k$$

where $d = 1/6 \max\{d_1, \dots, d_\ell, d'_1, \dots, d'_\ell\}$. Therefore we can choose a polynomial $\rho(k)$ which satisfies the inequality (5-43) as well as the condition on $p(1)$.

Case III. $i = j$. In this case it may not be useful to express σ_l as a conjugation of $\mathbb{E}(\phi^k(\tau))$ by β_l . It seems that $|\beta_l|$ can be larger than we expected as β_l translates $L(D_{v_l, k})$ and $L(D_{u_l, k})$ independently by distinct amounts $d_l \ell^k$ and $d'_{i'} \ell^k$ on the same ray. Instead of induction on k , we can use a pattern that the transpositions of $\mathbb{E}(\phi^k(\tau))$ follow. Observe that if $P = (i, p)$ and $Q = (i, q)$ are stable points, then $L(V_P^k)$ is a translation of $L(V_Q^k)$ on the ray R_i for each $k \in \mathbb{N}$. Assume that $q < p$ and $2 \leq i$. For $k = 1$, it is each to check that

$$p_l - q_l = (m_{l,i} + \ell(p - k_{l,i})) - (m_{l,i} + \ell(q - k_{l,i})) = \ell(p - q)$$

for each l . So $L(V_P^1)$ is the translation of $L(V_Q^1)$ by $\ell(p - q)$. Since

$$d_l - d'_{i'} = (p_l - p) - (q_l - q) = (p_l - q_l) - (p - q) = (\ell - 1)(p - q)$$

for all l , we can use the expression (5-40) to compare $L(v) = (i, p_v)$ and $L(u) = (i, q_u)$ for each pair of vertices $v \in V_P^k$ and $u \in V_Q^k$ with $W(v) = W(u)$. If $W(v) = l_1 l_2 \dots l_k$, then we have

$$\begin{aligned} p_v - q_u &= (d_{l_1} \ell^{k-1} + \dots + d_{l_{k-1}} \ell + p_{l_k}) - (d'_{l_1} \ell^{k-1} + \dots + d'_{l_{k-1}} \ell + q_{l_k}) \\ &= (d_{l_1} - d'_{l_1}) \ell^{k-1} + \dots + (d_{l_{k-1}} - d'_{l_{k-1}}) \ell + (p_{l_k} - q_{l_k}) \\ &= (\ell - 1)(p - q)(\ell^{k-1} + \dots + \ell) + \ell(p - q), \end{aligned}$$

which does not depend on the choice of v and u . Therefore the whole set $L(V_P^k)$ is a translation of $L(V_Q^k)$ by $y = p_v - q_u$. In particular, all ℓ^k transpositions of $\phi^k(\tau)$ are translations of one transposition on R_i . So we can write $\phi^k(\tau)$ as

$$(5-42) \quad \mathbb{E}(\phi^k(\tau)) = \prod_{1 \leq m \leq \ell^k} (x_m, x_m + y)$$

where (x, x') stands for the transposition exchanging (i, x) and (i, x') . Since all transpositions in (5-42) commute with each other we can arrange them so that $x_m < x_{m'}$ for all $m < m'$. Set $\tau_1 = (x_1, x_1 + y)$ to express $(x_m, x_m + y)$ as a conjugation of τ_1 by $g_i^{x_m - x_1}$ for all m . Cancelling subwords $g_i g_i^{-1}$ we obtain

$$\begin{aligned} |\mathbb{E}(\phi^k(\tau))| &= |\tau_1 \tau_1^{g_i^{x_2 - x_1}} \dots \tau_1^{g_i^{x_m - x_1}}| \\ &= |\tau_1 g_i^{-(x_2 - x_1)} \tau_1 g_i^{x_2 - x_1} g_i^{-(x_3 - x_1)} \dots g_i^{-(x_m - x_1)} \tau_1 g_i^{x_m - x_1}| \\ &\leq \ell |\tau_1| + 2(x_m - x_1) \end{aligned}$$

To bound $|\tau_1|$ we can use the fact that a transposition τ_0 with $\text{supp } \tau_0 \subset B_{n,r}$ has length at most $5r$. Proposition 5.19 implies that $\text{supp } \tau_1 \subset B_{n,r}$ with $r \leq A_0 \ell^k + s$ for some constant A_0 which does not depend on k . Thus $|\tau_1| \leq 5(A_0 \ell^k + s)$. Proposition 5.19 also implies that $x_m + y \leq A_0 \ell^k + s$ for all m . So

$$|E(\phi^k(\tau))| \leq 5\ell(A_0 \ell^k + s) + 2(A_0 \ell^k + s)$$

for all k . Therefore there exists a polynomial $\rho(k) = 5A_0 \ell + 2A_0 + 5s\ell + 2s$ such that $|E(\phi^k(\tau))| \leq \rho(k)\ell^k$ for all $k \in \mathbb{N}$.

In case P and Q are stable points of R_1 one can apply similar argument and calculation after expressing transpositions of (5-42) as conjugations of τ_1 by negative powers of g_i . So far we have shown that if τ exchanges two stable points there exists a polynomial ρ such that

$$(5-43) \quad |E(\phi^k(\tau))| \leq \rho(k)\ell^k$$

for all k where the degree of $\rho(k)$ does not depend on k .

Suppose $P, Q \in \text{ES}(\phi)$. For each $k \in \mathbb{N}$, V_P^k and V_Q^k can be expressed as disjoint unions of $N_P(k)$ and $N_Q(k)$ intervals respectively. The two collections of intervals determine two partitions on $\{1, \dots, \ell^k\}$ via the map $A : V_P^k \rightarrow \{1, \dots, \ell^k\}$. The common refinement of two partitions determine a set of common intervals for both V_P^k and V_Q^k . Let $\mathcal{I}_k = \{I_1, \dots, I_N\}$ and $\mathcal{I}'_k = \{I'_1, \dots, I'_N\}$ denote the common intervals for V_P^k and V_Q^k respectively, i.e., $W(I_j) = W(I'_j)$ for all j . Lemma 5.21 guarantees that $N \leq 2A_1 k \ell$ where A_1 is a constant determined by ϕ . Using those common intervals we can write $E(\phi(\tau)^k) = \prod_{1 \leq j \leq N} \sigma_j$ where

$$\sigma_j = \prod_{\omega \in W(I_j)} (P_\omega, Q_\omega)$$

for all k . Note that an interval of has size ℓ^m for some $m \in \mathbb{N}$ by definition. If an interval I_j has size ℓ then σ_j is a product of ℓ transpositions on $L(I_j) \cup L(I'_j)$ whose supports belong to a ball of radius $A_0 \ell^k + s$. So we have a bound

$$(5-44) \quad |\sigma_j| \leq 5(A_0 \ell^k + s) \leq (5A_0 + s)\ell^k = \rho_j(k)\ell^k.$$

If an interval I_j has size ℓ^m with $2 \leq m \leq k$ then I_j and I'_j were common subintervals of original intervals for V_P^k and V_Q^k that we started with. Thus there exist $u \in V_P$ and $v \in V_Q$ such that $I_j = D_{u,m} \cap V_P^k$, and $I'_j = D_{v,m} \cap V_Q^k$. Since both $L(u)$ and $L(v)$ are stable points, we have a polynomial as in (5-43) to bound $|\sigma_j|$ by

$$(5-45) \quad |\sigma_j| \leq \rho_j(m)\ell^m.$$

So far we have found upper bounds for individual σ_j depending on the size of the interval I_j . To obtain a universal bound consider the set of all polynomials $\{\rho_1, \dots, \rho_N\}$ that we need for (5-44) and (5-45). Let ρ denote the polynomial in the above set with the largest degree (or fastest growth). Assuming that ρ is increasing on $\mathbb{N}$, we have

$$(5-46) \quad |\sigma_j| \leq \rho(k)\ell^k$$

for all j . Thus we have found a polynomial $\ell\rho(k)$ such that

$$|E(\phi(\tau)^k)| = \sum_{1 \leq j \leq N} |\sigma_j| \leq \ell\rho(k)\ell^k$$

for all $k \in \mathbb{N}$. □

We remark that the expression (5-40) can be used to show that $V_P^k \subset B_{n,r}$ with r bounded by a linear term of ℓ^k as in Proposition 5.19.

Theorem 5.23. *Let ϕ be a monomorphism of $\mathcal{H}_n$ with $2 \leq n$. Then $\text{GR}(\phi) = \ell$ where $\ell = \ell(\phi)$.*

Proof. If $\ell(\phi) = 1$ it suffices to establish an upper bound since $\text{GR}(\phi) < 1$ means that ϕ is an eventually trivial map. In view of Proposition 5.24 let us assume $\phi(g_i) = g_i f_i$ for each i with $f_i \in \text{FSym}_n$. Each f_i is a product of transpositions; $f_i = \tau_1, \dots, \tau_{F_i}$. Lemma 5.14 implies that there exists a constant A_2 such that

$$|\phi^k(f_i)| = |\phi^k(\tau_1) \cdots \phi^k(\tau_{F_i})| \leq A_2 F_i k$$

for all $2 \leq i \leq n$ and $k \in \mathbb{N}$. Let F denote the maximum of F_i 's. From $\phi^{k+1}(g_i) = \phi^k(g_i)\phi^k(f_i)$, we have

$$|\phi^{k+1}(g_i)| \leq |\phi^k(g_i)| + |\phi^k(f_i)| \leq |\phi^k(g_i)| + A_2 F k$$

for all i and k . Take a polynomial ρ such that

$$\rho(k) + A_2 F k \leq \rho(k+1) \text{ and } G \leq \rho(1)$$

where $G = \max\{|\phi(g_i)| : 2 \leq i \leq n\}$. By induction on k , we check that

$$|\phi^{k+1}(g_i)| \leq \rho(k) + A_2 F k \leq \rho(k+1)$$

for all i . Thus we have a desired upper bound

$$\text{GR}(\phi) \leq \lim_{k \rightarrow \infty} \rho(k)^{1/k} = 1.$$

Therefore every monomorphism ϕ with $\ell(\phi) = 1$ satisfies $\text{GR}(\phi) \leq 1$ since $\text{GR}(\phi)^d = \text{GR}(\phi^d) \leq 1$ by Proposition 5.24.

Let us assume that $\ell(\phi) \geq 2$ and that ϕ satisfies $\phi(g_i) = g_i^\ell f_i$ for all $2 \leq i \leq n$ with $f_i \in \text{FSym}_n$. We first establish an upper bound for $\text{GR}(\phi)$. Each f_i can be written as a product of F_i transpositions, $f_i = \tau_1, \dots, \tau_{F_i}$. By Lemma 5.22 we can find polynomials $\rho_1, \dots, \rho_{F_i}$ such that

$$|\phi^k(f_i)| = |\phi^k(\tau_1) \cdots \phi^k(\tau_{F_i})| \leq (\rho_1(k) + \cdots + \rho_{F_i}(k)) \ell^k$$

for all $2 \leq i \leq n$ and $k \in \mathbb{N}$. Taking ρ to be the polynomial with largest degree over all $F_2 + \cdots + F_n$ polynomials, we have

$$|\phi^k(f_i)| \leq F \rho(k) \ell^k$$

for all i and k where F denotes the maximum over all F_i 's. Since $\phi^{k+1}(g_i) = (\phi^k(g_i))^\ell \phi^k(f_i)$, we have

$$(5-47) \quad |\phi^{k+1}(g_i)| \leq \ell |\phi^k(g_i)| + |\phi^k(f_i)| \leq \ell |\phi^k(g_i)| + F \rho(k) \ell^k$$

for all k . We seek a polynomial $\bar{\rho}_i$ such that $|\phi^k(g_i)| \leq \bar{\rho}_i(k) \ell^k$ for all k and i . There is no obstruction to take a polynomial $\bar{\rho}_i$ such that

$$\bar{\rho}_i(k) + F \rho(k) \leq \bar{\rho}_i(k+1) \text{ and } |\phi(g_i)| \leq \bar{\rho}_i(1)$$

for each i . Take a polynomial $\bar{\rho}$ such that $\bar{\rho}_i(k) \leq \bar{\rho}(k)$ for all i and k . Now (5-47) becomes

$$|\phi^{k+1}(g_i)| \leq \bar{\rho}(k) \ell^{k+1} + F \rho(k) \ell^k \leq (\bar{\rho}(k) + F \rho(k)) \ell^{k+1} \leq \bar{\rho}(k+1) \ell^{k+1}$$

for all i . Therefore we have a universal bound $|\phi^k(g_i)| \leq \bar{\rho}(k) \ell^k$ for all i and k where $\bar{\rho}$ is a polynomial on k . We have a desired upper bound

$$\text{GR}(\phi) \leq \lim_{k \rightarrow \infty} (\bar{\rho}(k) \ell^k)^{1/k} = \ell$$

The induced homomorphism $\bar{\phi}$ on the abelianization of $\mathcal{H}_n$ maps g_i to g_i^ℓ for all i . So $\text{GR}(\phi) \geq \text{GR}(\bar{\phi}) = \ell$. So far we have shown that $\text{GR}(\phi) = \ell(\phi)$ if ϕ satisfies the above assumption. Now Proposition 5.24 completes the proof because $\text{GR}(\phi)^d = \text{GR}(\phi^d) = \ell^d$. $\square$

Recall that R_1 is the common source of all generators $g_2, \dots, g_{n-1}$ and that each of them has a unique target such that targets are pairwise distinct. Lemma 5.7 tells us that the behavior of $\phi(g_i)$'s is similar to this up to an element of Σ_n , the group of outer automorphisms of $\mathcal{H}_n$ described in Theorem 2.3. Observe that a monomorphism ϕ of $\mathcal{H}_n$ defines a permutation δ_ϕ on $\{1, \dots, n\}$. For each i , $\phi(g_i)$ has a unique target, which defines the bijection γ as in (5–23); the target of $\phi(g_i)$ is $R_{\gamma(i)}$. We extend $\gamma : \{2, \dots, n\} \hookrightarrow \{1, \dots, n\}$ to get a bijection δ_ϕ on $\{1, \dots, n\}$ by setting $R_{\delta_\phi(1)}$ to be the unique source of $\phi(g_i)$'s. Note that δ_ϕ determines an arrangement of n rays

$$(5-48) \quad R_1 \mapsto R_{\delta_\phi(1)} \text{ and } R_i \mapsto R_{\gamma(i)} = R_{\delta_\phi(i)}$$

for $2 \leq i \leq n$. Lemma 5.7 guarantees that δ_ϕ is a bijection.

Being an element of the symmetric group Σ_n , $\delta = \delta_\phi$ determines a permutation matrix of $A_\delta \in \text{GL}(n, \mathbb{Z})$, which has exactly one entry of 1 in each row and each column and 0's elsewhere, with respect to the standard basis $\{e_1, \dots, e_n\}$ of $\mathbb{Z}^n$. The matrix A_δ has the form

$$\delta(i) \begin{pmatrix} & i \\ & \vdots \\ \cdots & 1 & \cdots \\ & \vdots \end{pmatrix}.$$

Under the correspondence $R_i \leftrightarrow e_i$ A_δ realizes δ in (5–48) as $e_{\delta(i)} = A_\delta(e_i)$ for each i . Recall that $\pi : \mathcal{H}_n \rightarrow \mathbb{Z}^n$ measures translation lengths on the rays; $\pi(g_i) = e_i - e_1$ for all i . The image $\pi(\mathcal{H}_n) = \{(m_1, \dots, m_n) \in \mathbb{Z}^n \mid \sum m_i = 0\}$ is freely generated by $\bar{g}_2, \dots, \bar{g}_n$ where $\bar{g} = \pi(g)$. The map A_δ restricts on the A_δ -invariant subgroup $\pi(\mathcal{H}_n)$ to define a matrix $\bar{A}_\delta \in \text{GL}(n-1, \mathbb{Z})$ with respect to the ordered basis $\{\bar{g}_2, \dots, \bar{g}_n\}$.

On the other hand, the induced map $\bar{\phi}$ in the following commutative diagram can be expressed as a matrix $A_{\bar{\phi}}$ with respect to the ordered basis $\{\bar{g}_2, \dots, \bar{g}_n\}$ of $\mathbb{Z}^{n-1}$.

$$\begin{array}{ccccccc} 1 & \longrightarrow & [\mathcal{H}_n, \mathcal{H}_n] & \longrightarrow & \mathcal{H}_n & \xrightarrow{\pi} & \mathbb{Z}^{n-1} \longrightarrow 1 \\ & & \downarrow \phi' & & \downarrow \phi & & \downarrow \bar{\phi} \\ 1 & \longrightarrow & [\mathcal{H}_n, \mathcal{H}_n] & \longrightarrow & \mathcal{H}_n & \xrightarrow{\pi} & \mathbb{Z}^{n-1} \longrightarrow 1 \end{array}$$

Since $\phi(g_i)$ has a unique target $R_{\delta(i)}$ and a unique source $R_{\delta(1)}$, where $\phi(g_i)$ acts as a translation by ℓ and $-\ell$ respectively, we have

$$\bar{\phi}(\bar{g}_i) = \overline{\phi(g_i)} = \ell(e_{\delta(i)} - e_{\delta(1)}) = \ell A_\phi(e_i - e_1) = \ell \bar{A}_\delta(\bar{g}_i)$$

for each i where $\ell = \ell(\phi)$. Thus we have

$$A_{\bar{\phi}} = \ell \bar{A}_\phi.$$

We also have

$$A_{\bar{\phi}^k} = A_{(\bar{\phi})^k} = (A_{\bar{\phi}})^k = \ell^k (\bar{A}_\phi)^k$$

for all $k \in \mathbb{N}$. In particular, $A_{\bar{\phi}^d} = (\bar{A}_\phi)^d = \ell^d I$ where $d = d(\delta_\phi)$ denotes the order of δ_ϕ and $I \in \text{GL}(n-1, \mathbb{Z})$ is the identity matrix. This proves Proposition 5.24.

Proposition 5.24. *A monomorphism ϕ of $\mathcal{H}_n$ satisfies that for each i , $2 \leq i \leq n$,*

$$\phi^d(g_i) \sim (g_i)^{\ell^d}$$

where $d = d(\delta_\phi)$ and $\ell = \ell(\phi)$.

REFERENCES

- [1] R. Bowen, Entropy and the fundamental group, *The structure of attractors in dynamical systems* (Proc. Conf., North Dakota State Univ., Fargo, N.D., 1977), pp. 21–29, Lecture Notes in Math., Vol. 668, Springer, Berlin, 1978.
- [2] K. Brown, Finiteness properties of groups, Proceedings of the Northwestern conference on cohomology of groups (Evanston, Ill., 1985), J. Pure Appl. Algebra Vol. 44, 1987
- [3] J. Burillo, S. Cleary, A. Martino and C. E. Röver, Commensurations and metric properties of Houghton’s groups, arXiv:1403.0026.
- [4] J. S. Cánovas, Commutativity formulas for fundamental group entropy, Int. J. Math. Game Theory Algebra, **18** (2009), 489–509.
- [5] Y. Cornulier, L. Guyot and W. Pitsch’ On the isolated points in the space of groups, J. Algebra, 307(1), (2007), 254277.
- [6] J. D. Dixon and B. Mortimer, Permutation groups, Springer-Verlag, New York, 1996.
- [7] K. J. Falconer, B. Fine and D. Kahrobaei, Growth rate of an endomorphism of a group, Groups. Complexity. Cryptology, (2011), 285–300.
- [8] A. Fel’shtyn, J. H. Jo and J. B. Lee, Growth rate for endomorphisms of finitely generated nilpotent groups and solvable groups, arXiv:1411.6360.
- [9] C. H. Houghton, The first cohomology of a group with permutation module coefficients, Arch. Math. (Basel), **31** (1978/79), 254–258.
- [10] G. Higman, Finitely presented infinite simple groups, Notes on Pure Mathematics, No. 8 (1974), Department of Pure Mathematics, Department of Mathematics, I.A.S. Australian National University, Canberra, 1974.
- [11] D. L. Johnson, Embedding some recursively presented groups, Groups St. Andrews 1997 in Bath, II, 410–416, London Math. Soc. Lecture Note Ser., **261**, Cambridge Univ. Press, Cambridge, 1999.
- [12] A. Katok and B. Hasselblatt, Introduction to the modern theory of dynamical systems, Encyclopedia of Mathematics and its Applications, 54, Cambridge University Press, Cambridge, 1995.
- [13] T. Koberda, Entropy of automorphisms, homology and the intrinsic polynomial structure of nilpotent groups, in the tradition of Ahlfors-Bers VI, 8799, Contemp. Math., **590**, Amer. Math. Soc.,
- [14] S. R. Lee, Geometry of Houghton’s groups, arXiv:1212.0257.
- [15] C. E. Röver, Subgroups of finitely presented simple groups, Ph.D. Thesis, University of Oxford, 1999.
- [16] W. R. Scott, Group Theory, Dover, 1987.

DEPARTMENT OF MATHEMATICS, SOGANG UNIVERSITY, SEOUL 121-742, KOREA
E-mail address: jlee@sogang.ac.kr

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TEXAS 77843, USA
E-mail address: srlee@math.tamu.edu