

Harmonic oscillator model for the helium atom

Martin Carlsen

*Department of Applied Mathematics and Computer Science,
Technical University of Denmark, Building 303B, DK-2800 Kongens Lyngby, Denmark**

(Dated: December 22, 2021)

A harmonic oscillator model in four dimensions is presented for the helium atom to estimate the distance to the inner and outer electron from the nucleus, the angle between electrons and the energy levels. The method is algebraic and is not based on the choice of correct trial wave function. Three harmonic oscillators and thus three quantum numbers are sufficient to describe the two-electron system. We derive a simple formula for the energy in the general case and in the special case of the Wannier Ridge. For a set of quantum numbers the distance to the electrons and the angle between the electrons are uniquely determined as the intersection between three surfaces. We show that the excited states converge either towards ionization thresholds or towards extreme parallel or antiparallel states and provide an estimate of the ground state energy.

I. INTRODUCTION

The Bohr-Sommerfeld quantization method did not succeed to predict the ground state of helium and the three-body Coulomb potential has since then been one of the major challenges in quantum physics. The Bohr-Sommerfeld quantization failed since they assumed ad-hoc rules that later turned out not to be valid. They for instance assumed that the electron distance to the core is the same for the two electrons and that the quantum numbers are integers. The spectrum of helium can be divided into 1) bound states that consist of the ground state and discrete singly excited states (singlet and triplet states) 2) doubly excited states that often are autoionizing and 3) the continuum above the three-particle breakup threshold[1]. The Hartree-Fock self-consistent-field method has shown to be particularly good at calculating bound states but it fails for doubly excited states where the electron-electron correlation is important (discovered by [2] and discussed in [3]). Numerical methods such as the complex rotation method based on a Hylleraas-type basis set or a Sturmian basis set using perimetric coordinates have been developed to study doubly excited states[4–12]. They have in addition been used to estimate the ground state energy with high accuracy[13]. An important question is why the single-particle picture of the Hartree-Fock method cannot be applied for doubly excited states. The molecular adiabatic approximation answers this question by considering the two-electron system as a H_2^+ molecule using the adiabatic distance as the interelectronic separation and the relative motion of the center of mass[14–16]. In this model the spatial wave function is separated into rotational, vibrational and molecular orbital wave functions, a more coherent picture than the single-particle picture. This model provides accurate estimations of the energy levels for doubly excited states. Classification methods to determine approximate quantum numbers of the two-electron system are primarily based on algebraic methods[17–20]. Finally, several semiclassical methods have been developed that consider the dynamics of a classical three-body potential[21–24]. These are used to study chaos.

The methods described above to determine the energy levels of a two-electron system have in common that the choice of wave functions are crucial to the success of these methods (for the molecular adiabatic method for instance it leads to quasiseparability of the three-body Coulomb Schrödinger equation). The algebraic classification methods are used to determine the structure of the energy levels but do not by themselves predict the energy levels. This means that no successful method to determine the bound and doubly excited energy levels of helium exists today that does not depend on the choice of spatial wave function.

In this work, the energy levels of helium are determined using an algebraic method based on the harmonic oscillator in four dimensions[25]. This leads to a system that consists of three oscillators and thus three quantum numbers N_1 , N_2 and N_3 . These three quantum numbers uniquely determine the energy E , the distance between the electrons and the nucleus r_1 and r_2 and the angle between the two electrons θ . The ground state is found at $N_1 = 1$, $N_2 = N_3 = 3/2$ with $E = -2.8827\text{a.u.}$, $\tan \alpha = \frac{r_1}{r_2} = 1.2635$, $r_2 = 1.0481\text{\AA}$ and $\cos \theta = -0.22725$. We analyze data for 1000 different combinations of the quantum numbers and derive a simple formula for the total energy. Finally, we consider the special case of the Wannier Ridge where $\tan \alpha = 1$.

* macar@dtu.dk

II. METHODS

A. The harmonic oscillator in 4 dimensions

In this section, we derive the formulas to calculate $r = r_2$, $\cos \theta$, $\tan \alpha$ and E . The derivation is based on a second order expansion of the helium potential and the algebraic method for a simple harmonic oscillator generalized to 4 dimensions. We derive the second order expansion in the next section.

We consider the Hamiltonian

$$H = \frac{1}{2m} \mathbf{p}^T \mathbf{p} + \frac{\hbar^2}{amr_{12}^3} \left(\frac{1}{2} \mathbf{x}^T \mathbf{C}^2 \mathbf{x} - \mathbf{c}^T \mathbf{x} \right) + \text{constant}, \quad (1)$$

where \mathbf{x} , \mathbf{p} , \mathbf{c} and \mathbf{C}^2 are defined as

$$\mathbf{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_4 \end{pmatrix}, \quad \mathbf{p} = \begin{pmatrix} p_1 \\ \vdots \\ p_4 \end{pmatrix}, \quad \mathbf{c} = \begin{pmatrix} c_1 \\ \vdots \\ c_4 \end{pmatrix}, \quad \mathbf{C}^2 = \begin{pmatrix} c_{11} & \dots & c_{14} \\ \vdots & \ddots & \vdots \\ c_{41} & \dots & c_{44} \end{pmatrix}, \quad (2)$$

and m is the mass of each electron, \hbar is Planck's constant, r_{12} is the distance between the electrons and a is the Bohr radius defined by

$$a = \frac{4\pi\epsilon_0\hbar^2}{me^2}. \quad (3)$$

In concordance with the standard algebraic method for simple harmonic oscillators we define the annihilation operator and creation operator as

$$\begin{aligned} \boldsymbol{\mu} &= \left(\frac{1}{4ar_{12}^3} \right)^{1/4} \left(\mathbf{C}\mathbf{x} + i\sqrt{\frac{ar^3}{\hbar^2}} \mathbf{p} \right) \\ \boldsymbol{\mu}^\dagger &= \left(\frac{1}{4ar_{12}^3} \right)^{1/4} \left(\mathbf{C}\mathbf{x} - i\sqrt{\frac{ar^3}{\hbar^2}} \mathbf{p} \right). \end{aligned} \quad (4)$$

Let \mathbf{e}_i and λ_i be the eigenvectors and eigenvalues of \mathbf{C}^2 . We assume that λ_i is nonnegative. The projections onto \mathbf{e}_i are given by

$$\begin{aligned} x_i &= \mathbf{e}_i^T \mathbf{x} \\ p_i &= \mathbf{e}_i^T \mathbf{p} \\ \mu_i &= \mathbf{e}_i^T \boldsymbol{\mu} \\ \mu_i^\dagger &= \mathbf{e}_i^T \boldsymbol{\mu}^\dagger \\ c_i &= \mathbf{e}_i^T \mathbf{c}. \end{aligned} \quad (5)$$

Clearly, x_i and p_i have the canonical commutation relation:

$$[x_i, p_i] = i\hbar. \quad (6)$$

When we add the creation and annihilation operator we find

$$\mathbf{C}\mathbf{x} = \frac{(ar_{12}^3)^{1/4}}{\sqrt{2}} (\boldsymbol{\mu} + \boldsymbol{\mu}^\dagger), \quad (7)$$

so x_i can be written as

$$x_i = \frac{(ar_{12}^3)^{1/4}}{\sqrt{2}\sqrt{\lambda_i}} (\mu_i + \mu_i^\dagger). \quad (8)$$

Using the canonical commutation relation, we find

$$\begin{aligned} [\mu_i, \mu_i^\dagger] &= \frac{-i\sqrt{\lambda_i}}{\hbar} [x_i, p_i] \\ &= \sqrt{\lambda_i}. \end{aligned} \quad (9)$$

As a result, we have

$$\begin{aligned} \mu_i \mu_i^\dagger &= \sqrt{\lambda_i} N_i + \frac{\sqrt{\lambda_i}}{2} \\ \mu_i^\dagger \mu_i &= \sqrt{\lambda_i} N_i - \frac{\sqrt{\lambda_i}}{2} \\ \mu_i \mu_i^\dagger + \mu_i^\dagger \mu_i &= 2\sqrt{\lambda_i} N_i. \end{aligned} \quad (10)$$

It is easy to see that

$$\mu_i^\dagger \mu_i = \left(\frac{r_{12}}{a}\right)^{3/2} \frac{1}{2E_0} H_i - \frac{\sqrt{\lambda_i}}{2}, \quad (11)$$

where

$$E_0 = \frac{\hbar^2}{2ma^2}, \quad (12)$$

so the total energy is given by

$$\begin{aligned} E &= \sum_i E_i \\ &= \sum_i 2E_0 \left(\frac{a}{r_{12}}\right)^{3/2} \sqrt{\lambda_i} N_i. \end{aligned} \quad (13)$$

We made the assumption that λ_i is nonnegative. This cannot be assumed in general. If λ_i is nonpositive then $\sqrt{\lambda_i}$ should be used instead of $\sqrt{\lambda_i}$ which ensures that $\sqrt{\lambda_i} N_i$ is negative. For simplicity, we will assume that N_i in this case is imaginary. It is also clear that N_i is an integer. However, N_i equal to $1/2, 3/2, 5/2$ is also a valid choice. We will thus assume that N_i can be either half-integers or integers.

To calculate r , $\cos \theta$ and $\tan \alpha$, we use

$$(\mathbf{c}^T \mathbf{e}_i)^2 = \frac{1}{2} (\mathbf{x}^T \mathbf{C}^2 \mathbf{e}_i)^2, \quad (14)$$

or

$$c_i^2 = \frac{1}{2} \lambda_i^2 x_i^2, \quad (15)$$

which can be reduced to

$$\begin{aligned} c_i^2 &= \frac{1}{2} \lambda_i^2 x_i^2 \\ &= \frac{\lambda_i}{4} \sqrt{ar_{12}^3} (\mu_i + \mu_i^\dagger)^2 \\ &= \frac{\lambda_i^{3/2}}{2} \sqrt{ar_{12}^3} N_i. \end{aligned} \quad (16)$$

We thus see that

$$r_{12}^3 = \frac{4c_i^4}{N_i^2 \lambda_i^3 a}. \quad (17)$$

This equation is very important. As c_i is proportional to r shown below this equation gives a formula for r as a function of θ and α for each of the harmonic oscillators. Since the equation has to be satisfied for all of the oscillators it can be used to give an estimate of r , $\cos \theta$ and $\tan \alpha$ and thus the energy.

When using equation (17) the energy can be written as

$$E = \sum_i E_0 \left(\frac{a\lambda_i N_i}{c_i}\right)^2 = 2E_0 \frac{2a}{r_{12}^3} \sum_i \frac{c_i^2}{\lambda_i}. \quad (18)$$

Finally, we remark that the method presented here easily can be generalized to any dimension.

B. Second order expansion of the helium potential

Let r_1 and r_2 be the distance between the electrons and the nucleus and let r_{12} be the distance between the electrons. The helium potential is given by

$$V = \frac{\hbar^2}{am} \left(-\frac{2}{r_1} - \frac{2}{r_2} + \frac{1}{r_{12}} \right). \quad (19)$$

We define \mathbf{x}_1 and \mathbf{x}_2 as the coordinates of the electrons and \mathbf{x}_0 as the coordinates of the nucleus. The unit vectors \mathbf{u}_1 , \mathbf{v}_1 and \mathbf{w}_1 are defined as

$$\mathbf{u}_1 = \frac{1}{r_1} (\mathbf{x}_1 - \mathbf{x}_0), \quad \mathbf{v}_1 = \frac{1}{r_2} (\mathbf{x}_2 - \mathbf{x}_0), \quad \mathbf{w}_1 = \frac{r_2}{r_{12}} \mathbf{v}_1 - \frac{r_1}{r_{12}} \mathbf{u}_1. \quad (20)$$

Let $\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$, $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ and $\{\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3\}$ be three orthonormal bases. Furthermore, let

$$\tan \alpha = \frac{r_1}{r_2}, \quad r = r_2, \quad (21)$$

and

$$\mathbf{v}_1 = \cos \theta \mathbf{u}_1 + \sin \theta \mathbf{u}_2. \quad (22)$$

It follows that

$$r_{12}^2 = r_1^2 + r_2^2 - 2r_1 r_2 \cos \theta = r^2 (1 + \tan^2 \alpha - 2 \tan \alpha \cos \theta). \quad (23)$$

In the three bases we define the operators:

$$\begin{aligned} \mathbf{P}_1 &= \mathbf{u}_1 \mathbf{u}_1^T, \quad \mathbf{P}_\perp = I - \mathbf{P}_1 \\ \mathbf{Q}_1 &= \mathbf{v}_1 \mathbf{v}_1^T, \quad \mathbf{Q}_\perp = I - \mathbf{Q}_1 \\ \mathbf{R}_1 &= \mathbf{w}_1 \mathbf{w}_1^T = \frac{1}{r_{12}^2} (r_1^2 \mathbf{u}_1 \mathbf{u}_1^T + r_2^2 \mathbf{v}_1 \mathbf{v}_1^T - r_1 r_2 \mathbf{u}_1 \mathbf{v}_1^T - r_1 r_2 \mathbf{v}_1 \mathbf{u}_1^T) \\ &= \frac{1}{r_{12}^2} \left((r_1 - r_2 \cos \theta)^2 \mathbf{u}_1 \mathbf{u}_1^T + r_2^2 \sin^2 \theta \mathbf{u}_2 \mathbf{u}_2^T - \sin \theta (r_1 r_2 - r_2^2 \cos \theta) (\mathbf{u}_1 \mathbf{u}_2^T + \mathbf{u}_2 \mathbf{u}_1^T) \right) \\ &= \frac{r^2}{r_{12}^2} \left((\tan \alpha - \cos \theta)^2 \mathbf{u}_1 \mathbf{u}_1^T + \sin^2 \theta \mathbf{u}_2 \mathbf{u}_2^T - \sin \theta (\tan \alpha - \cos \theta) (\mathbf{u}_1 \mathbf{u}_2^T + \mathbf{u}_2 \mathbf{u}_1^T) \right) \\ \mathbf{R}_\perp &= I - \mathbf{R}_1, \end{aligned} \quad (24)$$

such that \mathbf{P}_1 and \mathbf{P}_\perp in the basis $\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$ have the matrix representations

$$\mathbf{P}_1 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad \mathbf{P}_\perp = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad (25)$$

and similarly for \mathbf{Q}_1 and \mathbf{Q}_\perp in the basis $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ and \mathbf{R}_1 and \mathbf{R}_\perp in the basis $\{\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3\}$.

The second order perturbation of V is given by [26]

$$\begin{aligned} &\frac{\hbar^2}{am} \left(\frac{2}{r_1^2} \mathbf{x}^T \mathbf{u}_1 \frac{2}{r_2^2} \mathbf{y}^T \mathbf{v}_1 - \frac{1}{r_{12}^2} (\mathbf{x} - \mathbf{y})^T \mathbf{w}_1 \right. \\ &\left. - \frac{1}{r_1^3} \mathbf{x}^T (2\mathbf{P}_1 - \mathbf{P}_\perp) \mathbf{x} - \frac{1}{r_2^3} \mathbf{y}^T (2\mathbf{Q}_1 - \mathbf{Q}_\perp) \mathbf{y} + \frac{1}{2r_{12}^3} (\mathbf{x} - \mathbf{y})^T (2\mathbf{R}_1 - \mathbf{R}_\perp) (\mathbf{x} - \mathbf{y}) \right), \end{aligned} \quad (26)$$

where I have subtracted the zeroth order terms. This motivates us to consider the following potential

$$\frac{\hbar^2}{am} \left(\frac{2}{r_1^2} \mathbf{x}^T \mathbf{u}_1 - \frac{2}{r_2^2} \mathbf{y}^T \mathbf{v}_1 - \frac{1}{r_{12}^2} (\mathbf{x} - \mathbf{y})^T \mathbf{w}_1 - \frac{1}{r_1^3} \mathbf{x}^T 2\mathbf{P}_1 \mathbf{x} - \frac{1}{r_2^3} \mathbf{y}^T 2\mathbf{Q}_1 \mathbf{y} + \frac{1}{2r_{12}^3} (\mathbf{x} - \mathbf{y})^T 2\mathbf{R}_1 (\mathbf{x} - \mathbf{y}) \right). \quad (27)$$

There are two important changes: 1) We only consider terms involving \mathbf{u}_1 , \mathbf{v}_1 and \mathbf{w}_1 and 2) We consider $\mathbf{x} + \mathbf{y}$ instead of $\mathbf{x} - \mathbf{y}$ and let $\mathbf{y} \rightarrow -\mathbf{y}$.

This is an atomic system in four dimensions. Define $\mathbf{x} = x_1 \mathbf{u}_1 + x_2 \mathbf{u}_2$ and $\mathbf{y} = y_1 \mathbf{u}_1 + y_2 \mathbf{u}_2$. The 4×4 matrix \mathbf{C}^2 in equation (1) is given by

$$\mathbf{C}^2 = -4r_{12}^3 \begin{pmatrix} \frac{1}{r_1^3} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{r_2^3} \cos^2 \theta & \frac{1}{r_2^3} \cos \theta \sin \theta \\ 0 & 0 & \frac{1}{r_2^3} \cos \theta \sin \theta & \frac{1}{r_2^3} \sin^2 \theta \end{pmatrix} - \frac{2}{r_{12}^2} \begin{pmatrix} -(r_1 - r_2 \cos \theta)^2 & \sin \theta (r_1 r_2 - r_2^2 \cos \theta) & (r_1 - r_2 \cos \theta)^2 & -\sin \theta (r_1 r_2 - r_2^2 \cos \theta) \\ \sin \theta (r_1 r_2 - r_2^2 \cos \theta) & -r_2^2 \sin^2 \theta & -\sin \theta (r_1 r_2 - r_2^2 \cos \theta) & r_2^2 \sin^2 \theta \\ (r_1 - r_2 \cos \theta)^2 & -\sin \theta (r_1 r_2 - r_2^2 \cos \theta) & -(r_1 - r_2 \cos \theta)^2 & \sin \theta (r_1 r_2 - r_2^2 \cos \theta) \\ -\sin \theta (r_1 r_2 - r_2^2 \cos \theta) & r_2^2 \sin^2 \theta & \sin \theta (r_1 r_2 - r_2^2 \cos \theta) & -r_2^2 \sin^2 \theta \end{pmatrix}, \quad (28)$$

and the vector \mathbf{c} is given by

$$\mathbf{c} = \begin{pmatrix} \frac{2r_{12}^3}{r_1^2} + r_1 - r_2 \cos \theta \\ -r_2 \sin \theta \\ -\frac{2r_{12}^3}{r_2^2} \cos \theta - r_1 + r_2 \cos \theta \\ -\frac{2r_{12}^3}{r_2^2} \sin \theta + r_2 \sin \theta \end{pmatrix}. \quad (29)$$

They can be written as

$$\mathbf{C}^2 = -4\frac{r_{12}^3}{r^3} \begin{pmatrix} \tan^{-3} \alpha & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \cos^2 \theta & \cos \theta \sin \theta \\ 0 & 0 & \cos \theta \sin \theta & \sin^2 \theta \end{pmatrix} - \frac{2r^2}{r_{12}^2} \begin{pmatrix} -(\tan \alpha - \cos \theta)^2 & \sin \theta (\tan \alpha - \cos \theta) & (\tan \alpha - \cos \theta)^2 & -\sin \theta (\tan \alpha - \cos \theta) \\ \sin \theta (\tan \alpha - \cos \theta) & -\sin^2 \theta & -\sin \theta (\tan \alpha - \cos \theta) & \sin^2 \theta \\ (\tan \alpha - \cos \theta)^2 & -\sin \theta (\tan \alpha - \cos \theta) & -(\tan \alpha - \cos \theta)^2 & \sin \theta (\tan \alpha - \cos \theta) \\ -\sin \theta (\tan \alpha - \cos \theta) & \sin^2 \theta & \sin \theta (\tan \alpha - \cos \theta) & -\sin^2 \theta \end{pmatrix}, \quad (30)$$

and

$$\mathbf{c} = r \begin{pmatrix} 2\frac{r_{12}^3}{r^3} \tan^{-2} \alpha + \tan \alpha - \cos \theta \\ -\sin \theta \\ -2\frac{r_{12}^3}{r^3} \cos \theta - \tan \alpha + \cos \theta \\ -2\frac{r_{12}^3}{r^3} \sin \theta + \sin \theta \end{pmatrix}. \quad (31)$$

The eigenvectors and eigenvalues of \mathbf{C}^2 can be found numerically so the expressions for \mathbf{C}^2 and \mathbf{c} as a function of θ and α are enough to use the formulas derived in the preceding section. It is, however, useful to derive the analytical expression for \mathbf{e}_i and λ_i as it can be used to prove a formula for the energy.

C. Derivation of \mathbf{e}_i , λ_i and c_i

The eigenvalues and eigenvectors of \mathbf{C}^2 can be calculated using the equations

$$\begin{aligned} -\frac{r_{12}^2}{2r^2} \lambda a &= \left(\frac{2r_{12}^5}{r^5} \tan^{-3} \alpha - (\tan \alpha - \cos \theta)^2 \right) a + \sin \theta (\tan \alpha - \cos \theta) b + (\tan \alpha - \cos \theta)^2 c - \sin \theta (\tan \alpha - \cos \theta) d \\ -\frac{r_{12}^2}{2r^2} \lambda b &= \sin \theta (\tan \alpha - \cos \theta) a - \sin^2 \theta b - \sin \theta (\tan \alpha - \cos \theta) c + \sin^2 \theta d \\ -\frac{r_{12}^2}{2r^2} \lambda c &= (\tan \alpha - \cos \theta)^2 a - \sin \theta (\tan \alpha - \cos \theta) b + \left(\frac{2r_{12}^5}{r^5} \cos^2 \theta - (\tan \alpha - \cos \theta) \right) c + \left(\frac{2r_{12}^5}{r^5} \cos \theta \sin \theta + \sin \theta (\tan \alpha - \cos \theta) \right) d \\ -\frac{r_{12}^2}{2r^2} \lambda d &= -\sin \theta (\tan \alpha - \cos \theta) a + \sin^2 \theta b + \left(\frac{2r_{12}^5}{r^5} \cos \theta \sin \theta + \sin \theta (\tan \alpha - \cos \theta) \right) c + \left(\frac{2r_{12}^5}{r^5} \sin^2 \theta - \sin^2 \theta \right) d. \end{aligned} \quad (32)$$

From these four equations, we deduce

$$\begin{aligned}
-\frac{r_{12}^2}{2r^2}\lambda(\sin\theta a + (\tan\alpha - \cos\theta)b) &= \frac{2r_{12}^5}{r^5} \tan^{-3}\alpha \sin\theta a \\
-\frac{r_{12}^2}{2r^2}\lambda(\sin\theta c - (\tan\alpha - \cos\theta)b) &= \frac{2r_{12}^5}{r^5} (\cos^2\theta c + \cos\theta \sin\theta d) \sin\theta \\
-\frac{r_{12}^2}{2r^2}\lambda(\sin\theta a - (\tan\alpha - \cos\theta)d) &= \frac{2r_{12}^5}{r^5} (a \sin\theta \tan^{-3}\alpha - \cos\theta \sin\theta (\tan\alpha - \cos\theta)c - \sin^2\theta (\tan\alpha - \cos\theta)d) \\
-\frac{r_{12}^2}{2r^2}\lambda(a + c) &= \frac{2r_{12}^5}{r^5} (a \tan^{-3}\alpha + \cos^2\theta c + \cos\theta \sin\theta d) \\
-\frac{r_{12}^2}{2r^2}\lambda(b + d) &= \frac{2r_{12}^5}{r^5} (\cos\theta \sin\theta c + \sin^2\theta d) \\
-\frac{r_{12}^2}{2r^2}\lambda(\sin\theta(a + c) - \cos\theta(d + b)) &= \frac{2r_{12}^5}{r^5} \tan^{-3}\alpha \sin\theta a.
\end{aligned} \tag{33}$$

First we find that $\lambda = 0$ is an eigenvalue with $a = 0$, $b = 1$, $c = -\frac{\sin\theta}{\tan\alpha}$ and $d = \frac{-\cos\theta}{\sin\theta}c = \frac{\cos\theta}{\tan\alpha}$. Next, we can without loss of generality assume that $a = 1$. When combining the equations we find the relation

$$\sin\theta c - \cos\theta d = b \tan\alpha. \tag{34}$$

Furthermore, we obtain

$$\frac{1}{\tan\alpha \sin\theta + (\tan\alpha - \cos\theta)(\sin\theta c - \cos\theta d)} = \frac{(\cos\theta c + \sin\theta d) \tan^3\alpha}{\sin\theta c + (\tan\alpha - \cos\theta)d}. \tag{35}$$

This equation can be reduced to the quadratic equation

$$\begin{aligned}
(1 - \tan^4\alpha \cos\theta) \sin\theta c + ((\tan\alpha - \cos\theta) - \tan^4\alpha \sin^2\theta) d + (\tan\alpha - \cos\theta) \tan^3\alpha \cos\theta \sin\theta (d^2 - c^2) \\
+ (\tan\alpha - \cos\theta) \tan^3\alpha (\cos^2\theta - \sin^2\theta) cd = 0,
\end{aligned} \tag{36}$$

which can be written as

$$\begin{aligned}
\sin\theta c - \cos\theta d - \tan^4\alpha \sin\theta (\cos\theta c + \sin\theta d) + \tan\alpha d - (\tan\alpha - \cos\theta) \tan^3\alpha (\sin\theta c - \cos\theta d) (\cos\theta c + \sin\theta d) \\
= \tan\alpha \sin\theta (1 - \tan^3\alpha) \gamma_1 + (1 - \tan\alpha \cos\theta) \gamma_2 - (\tan\alpha - \cos\theta) \tan^3\alpha \gamma_1 \gamma_2 = 0,
\end{aligned} \tag{37}$$

or

$$-\left(\gamma_1 - \frac{1 - \tan\alpha \cos\theta}{(\tan\alpha - \cos\theta) \tan^3\alpha}\right)\left(\gamma_2 - \frac{\tan\alpha \sin\theta (1 - \tan^3\alpha)}{(\tan\alpha - \cos\theta) \tan^3\alpha}\right) + \frac{(1 - \tan\alpha \cos\theta)(1 - \tan^3\alpha) \sin\theta}{(\tan\alpha - \cos\theta)^2 \tan^5\alpha} = 0, \tag{38}$$

where γ_1 and γ_2 are defined as

$$\begin{aligned}
\gamma_1 &= \cos\theta c + \sin\theta d \\
\gamma_2 &= \sin\theta c - \cos\theta d,
\end{aligned} \tag{39}$$

and c and d are given by

$$\begin{aligned}
c &= \cos\theta \gamma_1 + \sin\theta \gamma_2 \\
d &= \sin\theta \gamma_1 - \cos\theta \gamma_2.
\end{aligned} \tag{40}$$

For γ_1 we have

$$\gamma_1 = \frac{(1 - \tan^3\alpha)(1 - \tan\alpha \cos\theta) \sin\theta}{(\tan\alpha - \cos\theta)^2 \tan^5\alpha \gamma_2 - \tan^3\alpha \sin\theta (1 - \tan^3\alpha) (\tan\alpha - \cos\theta)} + \frac{1 - \tan\alpha \cos\theta}{(\tan\alpha - \cos\theta) \tan^3\alpha}. \tag{41}$$

The following equality also holds:

$$\sin\theta (\tan\alpha - \cos\theta) - \sin^2\theta \frac{\gamma_2}{\tan\alpha} - \sin\theta \tan\alpha (\cos\theta \gamma_1 + \sin\theta \gamma_2) + \sin\theta \gamma_1 = \frac{2 \frac{r_{12}^5}{r^5} \tan^{-3}\alpha \sin\theta \frac{\gamma_2}{\tan\alpha}}{\sin\theta + (\tan\alpha - \cos\theta) \frac{\gamma_2}{\tan\alpha}}, \tag{42}$$

and it leads to the quadratic equation

$$\begin{aligned} \sin \theta (\tan \alpha - \cos \theta) + \sin \theta (1 - \tan \alpha \cos \theta) \gamma_1 + \left(-\frac{2r_{12}^5}{r^5} \tan^{-4} \alpha + \frac{(\tan \alpha - \cos \theta)^2}{\tan \alpha} - \frac{\sin^2 \theta}{\tan \alpha} - \sin^2 \theta \tan \alpha \right) \gamma_2 \\ + (1 - \tan \alpha \cos \theta) \frac{(\tan \alpha - \cos \theta)}{\tan \alpha} \gamma_1 \gamma_2 - \sin \theta \left(\frac{1}{\tan^2 \alpha} + 1 \right) (\tan \alpha - \cos \theta) \gamma_2^2 = 0. \end{aligned} \quad (43)$$

Using the equation above for $\gamma_1 \gamma_2$ we get

$$\begin{aligned} \sin \theta (\tan \alpha - \cos \theta) + \sin \theta \tan^{-3} \alpha (1 - \tan \alpha \cos \theta) \gamma_1 \\ + \left(-\frac{2r_{12}^5}{r^5} \tan^{-4} \alpha + \frac{(\tan \alpha - \cos \theta)^2}{\tan \alpha} - \frac{\sin^2 \theta}{\tan \alpha} - \sin^2 \theta \tan \alpha + \tan^{-4} \alpha (1 - \tan \alpha \cos \theta)^2 \right) \gamma_2 \\ - \sin \theta \left(\frac{1}{\tan^2 \alpha} + 1 \right) (\tan \alpha - \cos \theta) \gamma_2^2 = 0, \end{aligned} \quad (44)$$

and inserting the equation for γ_1 we finally obtain the cubic equation

$$\begin{aligned} -\sin^2 \theta (1 - \tan^3 \alpha) \tan^3 \alpha (\tan \alpha - \cos \theta)^2 \\ + (\sin \theta \tan^{-1} \alpha (1 - \tan \alpha \cos \theta)^2 + \sin \theta \tan^5 \alpha (\tan \alpha - \cos \theta)^2) (\tan \alpha - \cos \theta) \gamma_2 \\ - \tan^3 \alpha \sin \theta (1 - \tan^3 \alpha) \left(-\frac{2r_{12}^5}{r^5} \tan^{-4} \alpha + \frac{(\tan \alpha - \cos \theta)^2}{\tan \alpha} - \frac{\sin^2 \theta}{\tan \alpha} - \sin^2 \theta \tan \alpha + \tan^{-4} \alpha (1 - \tan \alpha \cos \theta)^2 \right) (\tan \alpha - \cos \theta) \gamma_2 \\ + \tan^3 \alpha \sin^2 \theta (1 - \tan^3 \alpha) \left(\frac{1}{\tan^2 \alpha} + 1 \right) (\tan \alpha - \cos \theta)^2 \gamma_2^2 \\ + \tan^5 \alpha \left(-\frac{2r_{12}^5}{r^5} \tan^{-4} \alpha + \frac{(\tan \alpha - \cos \theta)^2}{\tan \alpha} - \frac{\sin^2 \theta}{\tan \alpha} - \sin^2 \theta \tan \alpha + \tan^{-4} \alpha (1 - \tan \alpha \cos \theta)^2 \right) (\tan \alpha - \cos \theta)^2 \gamma_2^2 \\ - \tan^5 \alpha \sin \theta \left(\frac{1}{\tan^2 \alpha} + 1 \right) (\tan \alpha - \cos \theta)^3 \gamma_2^3 = 0. \end{aligned} \quad (45)$$

We denote the solutions to this equation $\gamma_{2,1}$, $\gamma_{2,2}$ and $\gamma_{2,3}$. $\gamma_{1,1}$, $\gamma_{1,2}$ and $\gamma_{1,3}$ can be found by calculating γ_1 . The eigenvectors of \mathbf{C}^2 are thus

$$\begin{aligned} \mathbf{e}_1 &= \frac{1}{\sqrt{1 + \gamma_{2,1}^2 (1 + \tan^{-2} \alpha) + \gamma_{1,1}^2}} \begin{pmatrix} 1 \\ \tan^{-1} \alpha \gamma_{2,1} \\ \cos \theta \gamma_{1,1} + \sin \theta \gamma_{2,1} \\ \sin \theta \gamma_{1,1} - \cos \theta \gamma_{2,1} \end{pmatrix}, \\ \mathbf{e}_2 &= \frac{1}{\sqrt{1 + \gamma_{2,2}^2 (1 + \tan^{-2} \alpha) + \gamma_{1,2}^2}} \begin{pmatrix} 1 \\ \tan^{-1} \alpha \gamma_{2,2} \\ \cos \theta \gamma_{1,2} + \sin \theta \gamma_{2,2} \\ \sin \theta \gamma_{1,2} - \cos \theta \gamma_{2,2} \end{pmatrix}, \\ \mathbf{e}_3 &= \frac{1}{\sqrt{1 + \gamma_{2,3}^2 (1 + \tan^{-2} \alpha) + \gamma_{1,3}^2}} \begin{pmatrix} 1 \\ \tan^{-1} \alpha \gamma_{2,3} \\ \cos \theta \gamma_{1,3} + \sin \theta \gamma_{2,3} \\ \sin \theta \gamma_{1,3} - \cos \theta \gamma_{2,3} \end{pmatrix}, \\ \mathbf{e}_4 &= \frac{1}{\sqrt{1 + \tan^{-2} \alpha}} \begin{pmatrix} 0 \\ 1 \\ -\sin \theta \tan^{-1} \alpha \\ \cos \theta \tan^{-1} \alpha \end{pmatrix}. \end{aligned} \quad (46)$$

The eigenvalues of \mathbf{C}^2 are

$$\begin{aligned}\lambda_1 &= -\frac{2r^2}{r_{12}^2 \gamma_{2,1}} \sin \theta \tan \alpha \xi_1 \\ \lambda_2 &= -\frac{2r^2}{r_{12}^2 \gamma_{2,2}} \sin \theta \tan \alpha \xi_2 \\ \lambda_3 &= -\frac{2r^2}{r_{12}^2 \gamma_{2,3}} \sin \theta \tan \alpha \xi_3 \\ \lambda_4 &= 0,\end{aligned}\tag{47}$$

where ξ_i are defined as

$$\xi_i = \tan \alpha - \cos \theta + (1 - \tan \alpha \cos \theta) \gamma_{1,i} - \sin \theta \tan^{-1} \alpha (1 + \tan^2 \alpha) \gamma_{2,i}.\tag{48}$$

Using the expressions for \mathbf{e}_i we find that $c_i = \mathbf{e}_i^T \mathbf{c}_i$ are given by

$$\begin{aligned}c_1 &= r \frac{\frac{2r_{12}^3}{r^3} (\tan^{-2} \alpha - \gamma_{1,1}) + \xi_1}{\sqrt{1 + \gamma_{2,1}^2 (1 + \tan^{-2} \alpha) + \gamma_{1,1}^2}} \\ c_2 &= r \frac{\frac{2r_{12}^3}{r^3} (\tan^{-2} \alpha - \gamma_{1,2}) + \xi_2}{\sqrt{1 + \gamma_{2,2}^2 (1 + \tan^{-2} \alpha) + \gamma_{1,2}^2}} \\ c_3 &= r \frac{\frac{2r_{12}^3}{r^3} (\tan^{-2} \alpha - \gamma_{1,3}) + \xi_3}{\sqrt{1 + \gamma_{2,3}^2 (1 + \tan^{-2} \alpha) + \gamma_{1,3}^2}} \\ c_4 &= 0.\end{aligned}\tag{49}$$

D. Formula for E

In this section we prove that the energy is given by the expression:

$$E = 2E_0 \frac{2a}{r} \left(-1 - \tan^{-1} \alpha + \frac{r}{2r_{12}} \right).\tag{50}$$

First, the orthogonality of the eigenvectors implies

$$\begin{aligned}1 + \gamma_{1,1} \gamma_{1,2} + (1 + \tan^{-2} \alpha) \gamma_{2,1} \gamma_{2,2} &= 0 \\ 1 + \gamma_{1,1} \gamma_{1,3} + (1 + \tan^{-2} \alpha) \gamma_{2,1} \gamma_{2,3} &= 0 \\ 1 + \gamma_{1,2} \gamma_{1,3} + (1 + \tan^{-2} \alpha) \gamma_{2,2} \gamma_{2,3} &= 0.\end{aligned}\tag{51}$$

Using the three equations we deduce the following useful relations

$$\begin{aligned}\sum_i \frac{1}{1 + \gamma_{1,i}^2 + (1 + \tan^2 \alpha) \gamma_{2,i}^2} &= 1, \quad \sum_i \frac{\gamma_{1,i}}{1 + \gamma_{1,i}^2 + (1 + \tan^2 \alpha) \gamma_{2,i}^2} = 0, \quad \sum_i \frac{\gamma_{2,i}}{1 + \gamma_{1,i}^2 + (1 + \tan^2 \alpha) \gamma_{2,i}^2} = 0 \\ \sum_i \frac{\gamma_{1,i}^2}{1 + \gamma_{1,i}^2 + (1 + \tan^2 \alpha) \gamma_{2,i}^2} &= 1, \quad \sum_i \frac{\gamma_{2,i}^2}{1 + \gamma_{1,i}^2 + (1 + \tan^2 \alpha) \gamma_{2,i}^2} = \frac{\tan^2 \alpha}{1 + \tan^2 \alpha}, \quad \sum_i \frac{\gamma_{1,i} \gamma_{2,i}}{1 + \gamma_{1,i}^2 + (1 + \tan^2 \alpha) \gamma_{2,i}^2} = 0 \\ \sum_i \frac{1}{\xi_i} \frac{\gamma_{1,i} \gamma_{2,i}}{1 + \gamma_{1,i}^2 + (1 + \tan^2 \alpha) \gamma_{2,i}^2} &= 0, \quad \sum_i \frac{1}{\xi_i} \frac{\gamma_{2,i}}{1 + \gamma_{1,i}^2 + (1 + \tan^2 \alpha) \gamma_{2,i}^2} = \frac{\sin \theta \tan^4 \alpha r^5}{2r_{12}^5}, \\ \sum_i \frac{1}{\xi_i} \frac{\gamma_{1,i}^2 \gamma_{2,i}}{1 + \gamma_{1,i}^2 + (1 + \tan^2 \alpha) \gamma_{2,i}^2} &= \frac{\sin \theta \tan \alpha r^5}{2r_{12}^5}.\end{aligned}\tag{52}$$




FIG. 1. Showing the three surfaces for $N_1 = 1$, $N_2 = N_3 = 3/2$. Equation (17) is satisfied for all of the three oscillators when the three surfaces intersect.

The first relation is proven in the Appendix A. Given the above relations we can readily prove the formula for the energy. It is given by

$$\begin{aligned}
 E &= 2E_0 \frac{2a}{r_{12}^3} \sum_i \frac{c_i^2}{\lambda_i} \\
 &= -2E_0 \frac{a}{r} \frac{r}{r_{12} \sin \theta \tan \alpha} \sum_i \frac{\gamma_{2,i} \left(2 \frac{r_{12}^3}{r^3} (\tan^{-2} \alpha - \gamma_{1,i}) + \xi_i \right)^2}{\xi_i (1 + \gamma_{1,i}^2 + \gamma_{2,i}^2 (1 + \tan^{-2} \alpha))} \\
 &= -2E_0 \frac{a}{r} \frac{r}{r_{12} \sin \theta \tan \alpha} \sum_i \gamma_{2,i} \frac{\frac{4 \frac{r_{12}^6}{r^6} (\tan^{-2} \alpha - \gamma_{1,i})^2}{\xi_i} + 4 \frac{r_{12}^3}{r^3} (\tan^{-2} \alpha - \gamma_{1,i}) + \xi_i}{1 + \gamma_{1,i}^2 + \gamma_{2,i}^2 (1 + \tan^{-2} \alpha)}.
 \end{aligned} \tag{53}$$

Several terms vanish due to the relations mentioned above. The non-vanishing terms are

$$E = -2E_0 \frac{a}{r} \frac{r}{r_{12} \sin \theta \tan \alpha} \sum_i \frac{\frac{4 \frac{r_{12}^6}{r^6} (\tan^{-4} \alpha \gamma_{2,i} + \gamma_{1,i}^2 \gamma_{2,i})}{\xi_i} - \sin \theta \tan^{-1} \alpha (1 + \tan^2 \alpha) \gamma_{2,i}^2}{1 + \gamma_{1,i}^2 + \gamma_{2,i}^2 (1 + \tan^{-2} \alpha)}, \tag{54}$$

so we finally obtain the expression for the energy of the helium atom

$$E = 2E_0 \frac{2a}{r} \left(-1 - \tan^{-1} \alpha + \frac{r}{2r_{12}} \right). \tag{55}$$

The derivation is not valid when $\tan \alpha = 1$. The formula for the energy in this special case is proven in Appendix B.

E. Numerical estimation of r , $\tan \alpha$, $\cos \theta$ and E

Equation (17) viewed as the function $r(\theta, \alpha, N_i)$ determines r for each oscillator. To evaluate r requires a calculation of e_i , λ_i and c_i that were found by calculating the three analytical solutions to equation (45). For a given N_i , r is a surface as a function θ and α . It is clear that r , θ and α must be uniquely determined. Hence, r , θ and α can be found by calculating the intersection points between the three surfaces. Observations suggest that only one intersection point exists although no proof has been given in this work. Given N_1 , N_2 and N_3 , r , θ and α can thus be estimated by calculating the unique intersection point where equation (17) is satisfied for all of the three oscillators. This is




FIG. 2. For $N_1 = 1, 2, 3, 4$ and 5 we plot the energy (left) and the distance to the inner electron (right) for different values of N_2 and N_3 . The surface plot of $N_1 = 1$ is lowest. The surface plot of $N_1 = 2$ is between $N_1 = 1$ and $N_1 = 3$ and so forth. The data can also be found in the supporting material.

illustrated in Figure 1. Let s_1, s_2 and s_3 be r for each of the three oscillators. The intersection point is estimated by calculating

$$\text{residual} = \min_{\theta, \alpha} (|s_1 - s_2| + |s_1 - s_3| + |s_2 - s_3|) \quad (56)$$

using a fine grid of 1000×1000 where the range of $\cos \theta$ is from -0.99999 to 0.99999 and the range of $\tan \alpha$ is from 1.00001 to 10 . The method is then iterated by decreasing the range of $\cos \theta$ and $\tan \alpha$ to be close to the estimated intersection point. For the estimated values of r, θ and α the energy can be calculated either by use of the analytical formula derived above or by equation (18).

III. RESULTS

The observables $r, \tan \alpha, \cos \theta$ and E were calculated using the intersection method described in the preceding section for different values of N_1, N_2 and N_3 under the assumption that the quantum numbers are either half-integers or integers. The estimations of $r, \tan \alpha, \cos \theta$ and E and the residuals given by equation (56) for 1000 different combinations of N_1, N_2 and N_3 with a range from 0.5 to 5 were tabulated in the supporting material.

To investigate how the quantum numbers determine the energy five three dimensional surface plots are shown in Figure 2(a) of the energy for different integer numbers of N_1 (1,2,3,4 and 5) and all combinations of N_2 and N_3 . Here, the surface plot of $N_1 = 1$ is lowest in the plot, over this plot is the surface plot of $N_1 = 2$ and so forth. We see that the energy is increased by a substantial amount when N_1 increases. Furthermore, the data suggests that the energy converges to the ionization thresholds for fixed N_1 and $N_2 = N_3 \rightarrow \infty$ and the three-particle breakup threshold for $N_1 = N_2 = N_3 \rightarrow \infty$. In Figure 2(b) the corresponding surface plots of r are shown. As for the energy the surface plot of $N_1 = 1$ is lowest in the plot, over this plot is the surface plot of $N_1 = 2$ and so forth. We see that r increases substantially when N_1 increases and when N_3 increases in particular for high values of N_1 . The equivalent surface plots of $\cos \theta$ and $\tan \alpha$ are not shown but the data can be found in the supporting material. The data shows that $\cos \theta$ has a little dependence on N_1 but is mainly determined by N_2 and N_3 while $\tan \alpha$ converges towards 1 when $N_1 \rightarrow \infty$.

Next, we investigate $\cos \theta, \tan \alpha, r$, and E for N_1 set to 1. Four surface plots are shown in Figure 3. The data clearly shows that an increase in N_2 leads to a decrease in $\cos \theta$ while an increase in N_3 leads to an increase in $\cos \theta$. The results of an increase in N_2 (N_3) is thus that the motion of the two electrons becomes more parallel (antiparallel). The results suggest that an increase in $\tan \alpha$ requires an increase in both N_2 and N_3 . Furthermore, r is almost




FIG. 3. For $N_1 = 1$ a surface plot of $\cos \theta$ (top left), $\tan \alpha$ (top right), r (bottom left) and E (bottom right) are shown for different values of N_2 and N_3 . The data can also be found in the supporting material.

constant for large values of N_2 and N_3 ($r_2 \approx a$) while the energy as described above converges to the first ionization threshold at -2 a.u.. When the system is excited by raising N_2 and N_3 simultaneously it thus follows that the angle between the electrons and the distance to the inner electron more or less are unchanged while the distance to the outer electron is increased substantially.

Finally, the data suggests (see the supporting material for the exact values) that 1 is the lowest state for N_1 since the energies of $N_1 = 0.5$ are too low. Furthermore, it is interesting that $N_1 = 1$, $N_2 = N_3 = 3/2$ gives the energy -78.44eV , which only is 0.57eV from the experimental value of -79.01eV .

IV. CONCLUSION

The object of this paper was to introduce a harmonic oscillator model to calculate the distance to the inner and outer electron, the angle between the electrons and the energy levels of helium. The method is based on a second order expansion of a two-electron Coulomb potential and the algebraic method for a harmonic oscillator. We derived

a simple formula for the energy of helium in the general case and in the specific case of the Wannier ridge where $\tan \alpha = 1$. A diagonalization of \mathbf{C}^2 led to three harmonic oscillators and the system can thus be described by three quantum numbers. The three harmonic oscillators are related through equation (17) which gives an equation for r as a function of θ and α . We found that the intersection point between the three surfaces uniquely determines r , $\cos \theta$ and $\tan \alpha$ and thus E for a choice of N_1 , N_2 and N_3 and calculated the values for 1000 different combinations of the three quantum numbers using an iterative method.

Our analysis of the dependence of r , $\cos \theta$, $\tan \alpha$ and E on the three quantum numbers indicates that the outer electron is excited towards the ionization threshold when both N_2 and N_3 is increased. The system becomes highly parallel or antiparallel if only one of the two quantum numbers is increased. Finally, the ground state is at $N_1 = 1$, $N_2 = N_3 = 3/2$ with energy -78.44eV .

Appendix A: Proof of the first relation in equation (52)

In this section we prove the following relation:

$$\sum_i \frac{1}{1 + \gamma_{1,i}^2 + (1 + \tan^2 \alpha) \gamma_{2,i}^2} = 1 \quad (\text{A1})$$

It is easy to see that we have to prove

$$\begin{aligned} & (1 + \gamma_{1,1}^2 + (1 + \tan^2 \alpha) \gamma_{2,1}^2) (1 + \gamma_{1,2}^2 + (1 + \tan^2 \alpha) \gamma_{2,2}^2) (1 + \gamma_{1,3}^2 + (1 + \tan^2 \alpha) \gamma_{2,3}^2) \\ &= (1 + \gamma_{1,1}^2 + (1 + \tan^2 \alpha) \gamma_{2,1}^2) (1 + \gamma_{1,2}^2 + (1 + \tan^2 \alpha) \gamma_{2,2}^2) \\ &+ (1 + \gamma_{1,1}^2 + (1 + \tan^2 \alpha) \gamma_{2,1}^2) (1 + \gamma_{1,3}^2 + (1 + \tan^2 \alpha) \gamma_{2,3}^2) \\ &+ (1 + \gamma_{1,2}^2 + (1 + \tan^2 \alpha) \gamma_{2,2}^2) (1 + \gamma_{1,3}^2 + (1 + \tan^2 \alpha) \gamma_{2,3}^2). \end{aligned} \quad (\text{A2})$$

This equation can be written as

$$\begin{aligned} & (\gamma_{1,1}^2 + (1 + \tan^2 \alpha) \gamma_{2,1}^2) (\gamma_{1,2}^2 + (1 + \tan^2 \alpha) \gamma_{2,2}^2) (\gamma_{1,3}^2 + (1 + \tan^2 \alpha) \gamma_{2,3}^2) \\ &= \gamma_{1,1}^2 \gamma_{1,2}^2 \gamma_{1,3}^2 + \gamma_{2,1}^2 \gamma_{2,2}^2 \gamma_{2,3}^2 (1 + \tan^{-2} \alpha)^3 + (\gamma_{2,1}^2 \gamma_{1,2}^2 \gamma_{1,3}^2 + \gamma_{2,2}^2 \gamma_{1,1}^2 \gamma_{1,3}^2 + \gamma_{2,3}^2 \gamma_{1,1}^2 \gamma_{1,2}^2) (1 + \tan^{-2} \alpha) \\ &+ (\gamma_{2,1}^2 \gamma_{2,2}^2 \gamma_{1,3}^2 + \gamma_{2,1}^2 \gamma_{2,3}^2 \gamma_{1,2}^2 + \gamma_{2,2}^2 \gamma_{2,3}^2 \gamma_{1,1}^2) (1 + \tan^{-2} \alpha)^2 \\ &= -1 + \sum_i (1 + \gamma_{1,i}^2 + (1 + \tan^2 \alpha) \gamma_{2,i}^2). \end{aligned} \quad (\text{A3})$$

Now, using equation (51) we obtain the equations:

$$\begin{aligned} & \gamma_{2,1}^2 \gamma_{2,2}^2 \gamma_{2,3}^2 (1 + \tan^{-2} \alpha)^3 = -1 - \gamma_{1,1} \gamma_{1,2} - \gamma_{1,1} \gamma_{1,3} - \gamma_{1,2} \gamma_{1,3} - \gamma_{1,1}^2 \gamma_{1,2} \gamma_{1,3} - \gamma_{1,2}^2 \gamma_{1,1} \gamma_{1,3} - \gamma_{1,3}^2 \gamma_{1,1} \gamma_{1,2} \\ & (\gamma_{2,1}^2 \gamma_{1,2}^2 \gamma_{1,3}^2 + \gamma_{2,2}^2 \gamma_{1,1}^2 \gamma_{1,3}^2 + \gamma_{2,3}^2 \gamma_{1,1}^2 \gamma_{1,2}^2) (1 + \tan^{-2} \alpha) = \gamma_{2,1}^2 \gamma_{1,2} \gamma_{1,3} (-1 - \gamma_{2,2} \gamma_{2,3} (1 + \tan^{-2} \alpha)) (1 + \tan^{-2} \alpha) \\ & + \gamma_{2,2}^2 \gamma_{1,1} \gamma_{1,3} (-1 - \gamma_{2,1} \gamma_{2,3} (1 + \tan^{-2} \alpha)) (1 + \tan^{-2} \alpha) + \gamma_{2,3}^2 \gamma_{1,1} \gamma_{1,2} (-1 - \gamma_{2,1} \gamma_{2,2} (1 + \tan^{-2} \alpha)) (1 + \tan^{-2} \alpha) \\ & (\gamma_{2,1}^2 \gamma_{2,2}^2 \gamma_{1,3}^2 + \gamma_{2,1}^2 \gamma_{2,3}^2 \gamma_{1,2}^2 + \gamma_{2,2}^2 \gamma_{2,3}^2 \gamma_{1,1}^2) (1 + \tan^{-2} \alpha)^2 = \gamma_{1,1}^2 \gamma_{2,2} \gamma_{2,3} (-1 - \gamma_{1,2} \gamma_{1,3}) (1 + \tan^{-2} \alpha) \\ & + \gamma_{1,2}^2 \gamma_{2,1} \gamma_{2,3} (-1 - \gamma_{1,1} \gamma_{1,3}) (1 + \tan^{-2} \alpha) + \gamma_{1,3}^2 \gamma_{2,1} \gamma_{2,2} (-1 - \gamma_{1,1} \gamma_{1,2}) (1 + \tan^{-2} \alpha) \\ & \gamma_{2,1}^2 \gamma_{1,2} \gamma_{1,3} + \gamma_{2,2}^2 \gamma_{1,1} \gamma_{1,3} + \gamma_{2,3}^2 \gamma_{1,1} \gamma_{1,2} = \gamma_{2,1}^2 + \gamma_{2,2}^2 + \gamma_{2,3}^2 + 2 (\gamma_{2,1}^2 \gamma_{2,2} \gamma_{2,3} + \gamma_{2,2}^2 \gamma_{2,1} \gamma_{2,3} + \gamma_{2,3}^2 \gamma_{2,1} \gamma_{2,2}) (1 + \tan^{-2} \alpha) \\ & + 3 \gamma_{2,1}^2 \gamma_{2,2}^2 \gamma_{2,3}^2 (1 + \tan^{-2} \alpha)^2, \end{aligned} \quad (\text{A4})$$

which when combined lead to the result.

Appendix B: The special case of the Wannier ridge

The eigenvalues and eigenvectors for the special case of $\alpha = \pi/4$ and $r_1 = r_2 = r$, the so called Wannier ridge[27], have to be derived separately. This important exception is treated below.

Clearly, $\lambda = 0$ is still an eigenvalue with the same eigenvector. Using equation (38) for $\tan \alpha = 1$ we find that $\gamma_1 = 1$ and $\gamma_2 = 0$ are two solutions to the equation. When $\gamma_2 = 0$ we have $\gamma_1 = -1$. On the other hand when $\gamma_1 = 1$ the cubic equation reduces to the quadratic equation:

$$-1 + b^2 - \beta b = 0, \quad (\text{B1})$$

with

$$\beta = \frac{2^4 \sin^3 \frac{\theta}{2} + 2 \cos \theta}{-\sin \theta}, \quad (\text{B2})$$

and

$$b = \gamma_{\pm}. \quad (\text{B3})$$

The solution to the quadratic equation is

$$\gamma_{\pm} = \frac{\beta}{2} \left(1 \pm \sqrt{1 + \frac{4}{\beta^2}} \right). \quad (\text{B4})$$

The solution has the following properties

$$\begin{aligned} \gamma_+ \gamma_- &= -1 \\ \beta = \gamma_+ + \gamma_- &= \gamma_+ - \frac{1}{\gamma_+} = \frac{\gamma_+^2 - 1}{\gamma_+} = \frac{\gamma_-^2 - 1}{\gamma_-} \\ \gamma_+ - \gamma_- &= \frac{\gamma_+^2 + 1}{\gamma_+} = -\frac{\gamma_-^2 + 1}{\gamma_-}. \end{aligned} \quad (\text{B5})$$

To conclude, the eigenvalues of \mathbf{C}^2 when $\tan \alpha = 1$ are given by

$$\begin{aligned} \lambda_1 &= -2^5 \sin^3 \frac{\theta}{2} = -4 \frac{r_{12}^3}{r^3} \\ \lambda_2 &= -4 \frac{r^2}{r_{12}^2} \sin \theta \frac{1 - \cos \theta - \sin \theta \gamma_+}{\gamma_+} \\ \lambda_3 &= -4 \frac{r^2}{r_{12}^2} \sin \theta \frac{1 - \cos \theta - \sin \theta \gamma_-}{\gamma_-} \\ \lambda_4 &= 0. \end{aligned} \quad (\text{B6})$$

and the eigenvectors are given by

$$\mathbf{e}_1 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ -\cos \theta \\ -\sin \theta \end{pmatrix}, \quad \mathbf{e}_2 = \frac{1}{\sqrt{2}} \frac{1}{\sqrt{\gamma_+^2 + 1}} \begin{pmatrix} 1 \\ \gamma_+ \\ \cos \theta + \sin \theta \gamma_+ \\ \sin \theta - \cos \theta \gamma_+ \end{pmatrix}, \quad \mathbf{e}_3 = \frac{1}{\sqrt{2}} \frac{1}{\sqrt{\gamma_-^2 + 1}} \begin{pmatrix} 1 \\ \gamma_- \\ \cos \theta + \sin \theta \gamma_- \\ \sin \theta - \cos \theta \gamma_- \end{pmatrix}, \quad \mathbf{e}_4 = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ -\sin \theta \\ \cos \theta \end{pmatrix}. \quad (\text{B7})$$

The reader can readily confirm that the eigenvectors are orthogonal.

Next, we calculate $c_i = \mathbf{e}_i^T \mathbf{c}_i$:

$$\begin{aligned} c_1 &= -\sqrt{2} 2^4 \sin^3 \frac{\theta}{2} r = -\frac{2\sqrt{2} r_{12}^3}{r^2} \\ c_2 &= -\sqrt{2} r \frac{1 - \cos \theta - \sin \theta \gamma_+}{\sqrt{\gamma_+^2 + 1}} \\ c_3 &= -\sqrt{2} r \frac{1 - \cos \theta - \sin \theta \gamma_-}{\sqrt{\gamma_-^2 + 1}} \\ c_4 &= 0. \end{aligned} \quad (\text{B8})$$

The formulas for the eigenvalues and the eigenvalues and the c_i 's are so simple that we readily can derive a formula for r and for the different energy terms. Using equation (17) for c_1 and λ_1 we obtain

$$r_{12}^3 = 4 \frac{\left(-2\sqrt{2} \frac{r_{12}^3}{r^2}\right)^4}{N_1^2 \left(-4 \frac{r_{12}^3}{r^3}\right)^3 a}. \quad (\text{B9})$$

Hence, we find that r is given by

$$r = -N_1^2 a = n_1^2 a, \quad (\text{B10})$$

where $N_1 = in_1$. To use equation (18), we calculate

$$\begin{aligned} \frac{c_1^2}{\lambda_1} &= -2 \frac{r_{12}^3}{r} \\ \frac{c_2^2}{\lambda_2} &= -\frac{r_{12}^2}{2} \frac{1}{\sin \theta} \frac{\gamma_+}{\gamma_+^2 + 1} (1 - \cos \theta - \sin \theta \gamma_+) \\ \frac{c_3^2}{\lambda_3} &= -\frac{r_{12}^2}{2} \frac{1}{\sin \theta} \frac{\gamma_-}{\gamma_-^2 + 1} (1 - \cos \theta - \sin \theta \gamma_-) \end{aligned} \quad (\text{B11})$$

Finally using $\frac{\gamma_-}{\gamma_-^2 + 1} = -\frac{1}{\gamma_+ - \gamma_-}$ and $\frac{\gamma_+}{\gamma_+^2 + 1} = \frac{1}{\gamma_+ - \gamma_-}$ we obtain the formula for the total energy of the helium atom at the Wannier ridge:

$$\begin{aligned} E &= 2E_0 \frac{2a}{r_{12}^3} \sum_i \frac{c_i^2}{\lambda_i} \\ &= 2E_0 \frac{2a}{r} \left(-2 + \frac{r}{2r_{12}} \right). \end{aligned} \quad (\text{B12})$$

- [1] G. Tanner, K. Richter, and J.-M. Rost, Rev. Mod. Phys. **72**, 497 (2000)
- [2] R. Madden and K. Codling, Phys. Rev. Lett. **10**, 516 (1963)
- [3] U. Fano, Rep. Prog. Phys. **46**, 97 (1983)
- [4] E. Balslev and J.-M. Combes, Commun. Math. Phys. **22**, 280 (1971)
- [5] Y. Ho, Phys. Rev. A **23**, 2137 (1981)
- [6] B. Junker, Adv. At. Mol. Phys. **18**, 207 (1982)
- [7] Y. K. Ho, Phys. Rep. **99**, 1 (1983)
- [8] Y. Ho and J. Callaway, J. Phys. B **18**, 3481 (1985)
- [9] Y. Ho, Phys. Rev. A **34**, 4402 (1986)
- [10] Y. Ho and A. Bhatia, Phys. Rev. A **44**, 2895 (1991)
- [11] Y. Ho, Phys. Rev. A **48**, 3598 (1993)
- [12] E. Lindroth, Phys. Rev. A **49**, 4473 (1994)
- [13] S. Salomonson and P. Öster, Phys. Rev. A **40**, 5559 (1989)
- [14] J. Macek, J. Phys. B **1**, 831 (1968)
- [15] J. M. Feagin and J. S. Briggs, Phys. Rev. Lett. **57**, 984 (1986)
- [16] J. M. Feagin and J. S. Briggs, Phys. Rev. A **37**, 4599 (1988)
- [17] C. Wulfman, Phys. Lett. A **26**, 397 (1968)
- [18] D. R. Herrick and O. Sinanolu, Phys. Rev. A **11**, 97 (1975)
- [19] D. R. Herrick, Adv. Chem. Phys., 1(1983)
- [20] C. Lin, Phys. Rev. A **29**, 1019 (1984)
- [21] M. C. Gutzwiller, J. Math. Phys. **8**, 1979 (1967)
- [22] M. C. Gutzwiller, J. Math. Phys. **12**, 343 (1971)
- [23] G. Ezra, K. Richter, G. Tanner, and D. Wintgen, J. Phys. B **24**, L413 (1991)
- [24] D. Wintgen, K. Richter, and G. Tanner, Chaos **2**, 19 (1992)
- [25] M. Moshinsky, *The harmonic oscillator in modern physics* (Gordon and Breach, 1969)
- [26] M. Carlsen, J. Comput. Chem **35**, 1149 (2014)
- [27] K. Richter and D. Wintgen, J. Phys. B **23**, L197 (1990)

Appendix C: Supporting material

Numerical data for r , $\cos \theta$, $\tan \alpha$, E and the residuals. The value of N_1 is in the first row and first column. The value of N_2 is in first row from the second to the eleventh column. The value of N_3 is in the first column from the second to the eleventh row. The data is tabulated from the second to the eleventh row and from the second to the eleventh column. The data was calculated using the iterative intersection method described in main document.

1. The distance to the inner electron - r in Angstrom

0.5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	0.197	0.20277	0.20418	0.20468	0.20492	0.20505	0.20512	0.20517	0.20521	0.20523
1	0.26474	0.29101	0.29164	0.29152	0.29139	0.29131	0.29125	0.29122	0.29119	0.29117
1.5	0.16824	0.26868	0.28316	0.28448	0.28467	0.28467	0.28464	0.28461	0.28458	0.28456
2	0.16388	0.14321	0.25997	0.26861	0.27117	0.27235	0.27298	0.27337	0.27363	0.2738
2.5	0.18006	0.10637	0.18006	0.25609	0.2608	0.26304	0.26436	0.26521	0.2658	0.26621
3	0.20569	0.093861	0.13046	0.20526	0.25403	0.25669	0.25834	0.25948	0.26029	0.26089
3.5	0.23935	0.087936	0.11058	0.15766	0.21666	0.25278	0.25438	0.25555	0.25643	0.25712
4	0.28259	0.084672	0.10009	0.13427	0.17753	0.24858	0.25209	0.253	0.25381	0.25449
4.5	0.33989	0.082701	0.093741	0.12068	0.15471	0.19034	0.22247	0.24153	0.25276	0.25293
5	0.42506	0.08143	0.089566	0.11197	0.14005	0.17007	0.19848	0.22276	0.24213	0.24262

1	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	0.48825	0.49568	0.50073	0.50266	0.50358	0.50408	0.50438	0.50458	0.50471	0.50481
1	0.7923	0.788	0.8037	0.81107	0.8147	0.81671	0.81793	0.81873	0.81928	0.81967
1.5	1.0127	1.0259	1.0481	1.0595	1.0652	1.0683	1.0702	1.0714	1.0722	1.0728
2	1.1968	1.059	1.1521	1.164	1.1663	1.1666	1.1663	1.1661	1.1658	1.1656
2.5	1.4868	0.79027	1.0932	1.1566	1.1658	1.1668	1.1662	1.1653	1.1646	1.1639
3	1.9304	0.67297	0.75021	1.0747	1.1212	1.1327	1.1365	1.1379	1.1385	1.1387
3.5	2.5843	0.64693	0.54307	0.82873	1.0539	1.0876	1.0997	1.1055	1.1087	1.1106
4	3.5147	0.65551	0.47032	0.57284	0.90557	1.0399	1.0636	1.0744	1.0807	1.0847
4.5	4.7524	0.68204	0.43573	0.47563	0.64594	0.95537	1.0307	1.0472	1.0562	1.0621
5	6.2552	0.72026	0.41726	0.42547	0.52912	0.72024	0.98624	1.0244	1.036	1.0432

1.5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	0.91144	0.89901	0.90584	0.91073	0.91313	0.91445	0.91526	0.91578	0.91614	0.9164
1	1.3913	1.3142	1.3342	1.3483	1.3558	1.36	1.3626	1.3643	1.3655	1.3663
1.5	1.9248	1.7549	1.773	1.7992	1.8153	1.8249	1.8309	1.8349	1.8376	1.8396
2	2.4821	2.1271	2.1592	2.192	2.2153	2.2299	2.2392	2.2454	2.2497	2.2528
2.5	3.1408	2.2522	2.4099	2.4608	2.4849	2.4985	2.5065	2.5116	2.5149	2.5172
3	3.9755	2.2162	2.3827	2.5657	2.6065	2.6191	2.6233	2.6246	2.6247	2.6245
3.5	5.0122	2.2451	1.9694	2.4563	2.5954	2.6287	2.6375	2.639	2.6382	2.6368
4	6.2466	2.3653	1.65	1.971	2.4506	2.5586	2.5903	2.601	2.6046	2.6054
4.5	7.6598	2.5636	1.5142	1.4703	2.0606	2.4181	2.5038	2.5348	2.5485	2.5551
5		2.835	1.4633	1.2551	1.4896	2.1396	2.3853	2.4529	2.4821	2.4973

2	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	1.4807	1.4114	1.4244	1.4339	1.439	1.4418	1.4435	1.4447	1.4455	1.446
1	2.1216	1.953	1.9619	1.9827	1.9954	2.0029	2.0076	2.0107	2.0128	2.0143
1.5	2.8685	2.5565	2.5381	2.5693	2.5937	2.6092	2.6192	2.6258	2.6305	2.6338
2	3.7031	3.1692	3.1193	3.152	3.1887	3.2148	3.2323	3.2443	3.2527	3.2588
2.5	4.6352	3.6838	3.6487	3.6827	3.7263	3.7611	3.7858	3.803	3.8153	3.8242
3	5.7013	4.0509	4.0112	4.1035	4.1547	4.1924	4.2193	4.238	4.2512	4.2608
3.5	6.9224	4.3843	4.0464	4.3326	4.4329	4.4772	4.5024	4.5179	4.5278	4.5345
4	8.2999	4.7873	3.8361	4.2359	4.5215	4.6086	4.6418	4.6561	4.6625	4.6652
4.5	9.8276	5.3013	3.6672	3.7095	4.3511	4.5827	4.6568	4.6839	4.6936	4.6963
5	11.4984	5.9471	3.6151	3.1611	3.7639	4.3727	4.5601	4.6266	4.6529	4.6633

2.5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	2.1863	2.0183	2.0538	2.0722	2.0811	2.0864	2.0897	2.0918	2.0933	2.0944
1	2.9726	2.7268	2.7018	2.7257	2.7447	2.7566	2.7643	2.7694	2.773	2.7755
1.5	3.8802	3.4876	3.4045	3.4294	3.4609	3.4832	3.498	3.5081	3.5152	3.5203
2	4.9003	4.3029	4.1466	4.1575	4.1993	4.2345	4.2595	4.277	4.2894	4.2985
2.5	6.0292	5.1152	4.8901	4.8799	4.925	4.9721	5.0087	5.0354	5.0548	5.0692
3	7.2752	5.8734	5.5557	5.5544	5.5989	5.6531	5.6991	5.7343	5.7606	5.7803
3.5	8.6519	6.5991	6.0218	6.1102	6.1763	6.2352	6.2858	6.3254	6.3554	6.3781
4	10.1662	7.3719	6.2432	6.4298	6.6002	6.6815	6.736	6.7756	6.8046	6.826
4.5	11.8187	8.2589	6.3556	6.3784	6.7888	6.9563	7.032	7.0746	7.1012	7.1186
5	13.6072	9.3022	6.4989	6.0194	6.6186	7.0142	7.1626	7.2272	7.2588	7.2752

3	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	3.0162	2.8367	2.8404	2.7572	2.8392	2.8479	2.8534	2.8572	2.8597	2.8616
1	3.9374	3.6458	3.5616	3.596	3.6059	3.6234	3.635	3.6429	3.6485	3.6525
1.5	4.9841	4.5636	4.3942	4.3971	4.4318	4.4611	4.4818	4.4963	4.5065	4.514
2	6.1515	5.5653	5.2888	5.2568	5.2948	5.3369	5.3696	5.3932	5.4104	5.4231
2.5	7.4366	6.6214	6.2191	6.141	6.1715	6.2239	6.2698	6.305	6.3312	6.3509
3	8.8392	7.6992	7.1307	7.0197	7.0355	7.092	7.1495	7.1968	7.2333	7.2613
3.5	10.3637	8.7897	7.9425	7.839	7.8523	7.9084	7.9731	8.0301	8.0761	8.1121
4	12.0158	9.9284	8.593	8.5082	8.5718	8.6366	8.7053	8.7678	8.8197	8.8611
4.5	13.7984	11.171	9.1145	8.9111	9.1178	9.2329	9.3134	9.3794	9.433	9.4756
5	15.713	12.5631	9.6048	9.0087	9.3793	9.6398	9.7654	9.8432	9.8985	9.9395

3.5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	3.9665	3.6993	3.6817	3.7561	3.8052	3.8043	3.7344	3.7403	3.7444	3.7474
1	5.0179	4.7125	4.5476	4.5373	4.5802	4.6036	4.6202	4.6317	4.6399	4.6459
1.5	6.1958	5.7852	5.5194	5.4819	5.5128	5.5479	5.5751	5.5948	5.609	5.6195
2	7.4972	6.9618	6.5658	6.4708	6.4933	6.5387	6.5787	6.6091	6.6318	6.6487
2.5	8.9205	8.2261	7.6721	7.5064	7.5065	7.5565	7.6092	7.6526	7.686	7.7114
3	10.4644	9.5596	8.801	8.5658	8.5338	8.5801	8.6428	8.6994	8.7451	8.7809
3.5	12.13	10.9496	9.9001	9.6081	9.5496	9.5856	9.653	9.7208	9.7788	9.8256
4	13.9184	12.4043	10.9203	10.5658	10.514	10.5439	10.6116	10.6866	10.7545	10.8113
4.5	15.8337	13.9543	11.8517	11.3503	11.3641	11.4163	11.4877	11.5666	11.6405	11.7043
5	17.8765	15.6346	12.7397	11.9013	12.0074	12.1483	12.2456	12.3318	12.4093	12.4761

4	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	5.0377	4.8204	4.7053	4.5539	4.6494	4.7124	4.7342	4.741	4.7474	4.7517
1	6.2172	5.9228	5.6874	5.6455	5.6775	5.6977	5.7198	5.7357	5.7473	5.7558
1.5	7.5231	7.1465	6.7877	6.6917	6.7079	6.7459	6.7795	6.8051	6.8242	6.8384
2	8.9537	8.4862	7.987	7.812	7.805	7.8476	7.8934	7.9309	7.9597	7.9817
2.5	10.5079	9.9323	9.2645	8.9962	8.9497	8.9881	9.0441	9.0948	9.1356	9.1674
3	12.1852	11.4739	10.5947	10.226	10.1288	10.1523	10.2138	10.277	10.3311	10.3748
3.5	13.9852	13.1012	11.9446	11.4695	11.3226	11.3229	11.3829	11.4559	11.5231	11.5795
4	15.9018	14.8125	13.2796	12.6769	12.4992	12.4771	12.5299	12.608	12.6861	12.7548
4.5	17.9537	16.6189	14.5795	13.7839	13.6082	13.5832	13.6295	13.7086	13.7938	13.8725
5	20.1266	18.5408	15.8569	14.7351	14.5762	14.5948	14.65	14.7308	14.82	14.9051

4.5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	6.2311	6.0361	5.768	5.7412	5.7827	5.8762	5.8497	5.8582	5.8774	5.8897
1	7.5373	7.2694	6.9525	6.9004	6.9032	6.987	6.9338	6.9548	6.9704	6.9821
1.5	8.9705	8.6387	8.2029	8.0333	8.0233	8.0911	8.0963	8.1281	8.1526	8.1712
2	10.5284	10.1308	9.556	9.2888	9.2374	9.2694	9.3174	9.3615	9.3969	9.4245
2.5	12.2103	11.7389	11.0013	10.6221	10.5134	10.5292	10.5832	10.6396	10.6877	10.7264
3	14.0169	13.4565	12.522	12.0178	11.8401	11.8277	11.8801	11.9467	12.0081	12.0595
3.5	15.9462	15.2769	14.0973	13.4508	13.2014	13.1513	13.1936	13.266	13.3398	13.4047
4		17.1962	15.7039	14.8845	14.5714	14.4822	14.5072	14.5798	14.6631	14.7408
4.5		19.2164	17.3232	16.2732	15.9098	15.7942	15.8007	15.8684	15.957	16.0453
5		21.3465	18.9527	17.5728	17.1593	17.0488	17.0475	17.1092	17.1994	17.2949

5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	7.5478	7.3746	7.185	6.7936	6.9716	6.9912	7.0834	7.0723	7.0996	7.1142
1	8.9803	8.7451	8.3754	8.0733	8.1195	8.2153	8.2626	8.2888	8.3091	8.3245
1.5	10.5392	10.2545	9.7666	9.498	9.5475	9.4872	9.5298	9.5642	9.5945	9.6182
2	12.2241	11.8903	11.2722	10.9073	10.7972	10.807	10.8537	10.9028	10.9449	10.9787
2.5	14.0321	13.6472	12.8819	12.391	12.2062	12.1873	12.2326	12.2916	12.3463	12.392
3		15.5206	14.5838	13.9505	13.6796	13.6182	13.6526	13.7178	13.7847	13.8435
3.5		17.5066	16.3646	15.5662	15.2036	15.0894	15.1031	15.1684	15.2455	15.3176
4		19.6012	18.2089	17.2115	16.7565	16.5865	16.5714	16.6298	16.7131	16.7973
4.5		21.8039	20.1022	18.854	18.3059	18.0876	18.041	18.0864	18.1709	18.2644
5		24.1117	22.0373	20.4606	19.8076	19.5605	19.4896	19.5197	19.6009	19.7

2. The angle between the electrons - $\cos \theta$

0.5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	-0.082998	-0.70383	-0.86231	-0.92135	-0.94931	-0.96466	-0.97398	-0.98005	-0.98422	-0.98721
1	0.58678	-0.26771	-0.62124	-0.7752	-0.85253	-0.89621	-0.92313	-0.94084	-0.95309	-0.96191
1.5	0.90507	0.25099	-0.31316	-0.57422	-0.71378	-0.79564	-0.84727	-0.88174	-0.90583	-0.92329
2	0.94921	0.87727	0.049786	-0.37733	-0.57815	-0.69533	-0.77013	-0.82072	-0.85646	-0.88259
2.5	0.96571	0.94411	0.83001	-0.11693	-0.43767	-0.59679	-0.69487	-0.76085	-0.8076	-0.84196
3	0.97431	0.9652	0.92722	0.80271	-0.24125	-0.48586	-0.61628	-0.70004	-0.75842	-0.80113
3.5	0.97954	0.97564	0.95508	0.9128	0.80579	-0.32931	-0.52279	-0.63305	-0.70627	-0.75865
4	0.98299	0.9818	0.96852	0.94526	0.90586	0.35851	-0.40233	-0.55106	-0.64667	-0.71199
4.5	0.98538	0.98582	0.9764	0.96117	0.93865	0.90564	0.85016	0.76034	-0.60718	-0.6669
5	0.98704	0.98861	0.98151	0.97058	0.95553	0.93562	0.90958	0.87149	0.78525	0.80564

1	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	0.36351	-0.43969	-0.72568	-0.84054	-0.8964	-0.92747	-0.94646	-0.95888	-0.96744	-0.97358
1	0.69917	-0.082998	-0.50667	-0.70383	-0.80481	-0.86231	-0.89789	-0.92135	-0.9376	-0.94931
1.5	0.86433	0.26324	-0.22725	-0.50837	-0.667	-0.76161	-0.82166	-0.86185	-0.88998	-0.91037
2	0.92295	0.58678	0.071609	-0.26771	-0.48326	-0.62124	-0.71251	-0.7752	-0.8198	-0.85253
2.5	0.9489	0.83294	0.38182	-0.022741	-0.28759	-0.46596	-0.5885	-0.67489	-0.73749	-0.78402
3	0.96401	0.90507	0.77714	0.25099	-0.091993	-0.31316	-0.46553	-0.57422	-0.65392	-0.71378
3.5	0.97436	0.9337	0.8954	0.69564	0.14577	-0.1548	-0.34447	-0.47641	-0.5727	-0.64517
4	0.98202	0.94921	0.93142	0.87727	0.59763	0.049786	-0.2131	-0.37733	-0.49262	-0.57815
4.5	0.98767	0.95898	0.94952	0.92268	0.8534	0.48828	-0.038512	-0.26582	-0.40887	-0.51045
5	0.99158	0.96571	0.96054	0.94411	0.91041	0.83001	0.34179	-0.11693	-0.31225	-0.43767

1.5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	0.6531	-0.14382	-0.54953	-0.73123	-0.82332	-0.87553	-0.90776	-0.92898	-0.94367	-0.95425
1	0.83032	0.15769	-0.32365	-0.58062	-0.71954	-0.80062	-0.85146	-0.88525	-0.90878	-0.92579
1.5	0.91134	0.41103	-0.082998	-0.40034	-0.58883	-0.70383	-0.77764	-0.82738	-0.86231	-0.88772
2	0.94812	0.61809	0.15188	-0.19679	-0.42909	-0.58041	-0.68117	-0.75057	-0.80001	-0.83632
2.5	0.96694	0.77384	0.37119	0.014617	-0.24681	-0.43086	-0.55982	-0.6516	-0.71841	-0.7682
3	0.97807	0.8582	0.58678	0.22154	-0.058529	-0.26771	-0.42202	-0.53603	-0.62124	-0.68596
3.5	0.98517	0.89933	0.77857	0.43852	0.12821	-0.10425	-0.28075	-0.41503	-0.51778	-0.59722
4	0.98982	0.92239	0.8667	0.70217	0.33318	0.060852	-0.14101	-0.29526	-0.41464	-0.50802
4.5	0.99287	0.93706	0.90507	0.84906	0.61588	0.25099	0.0044592	-0.17565	-0.31316	-0.42055
5		0.94725	0.92627	0.89934	0.82659	0.52945	0.17947	-0.046979	-0.20977	-0.33366

2	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	0.81882	0.1203	-0.35437	-0.60151	-0.73396	-0.81102	-0.85926	-0.8913	-0.9136	-0.92972
1	0.90659	0.36351	-0.13239	-0.43969	-0.61805	-0.72568	-0.79439	-0.84054	-0.8729	-0.8964
1.5	0.94766	0.5537	0.079939	-0.26405	-0.48457	-0.62444	-0.71612	-0.77866	-0.82294	-0.85532
2	0.96837	0.69917	0.27188	-0.082998	-0.3357	-0.50667	-0.62285	-0.70383	-0.76194	-0.80481
2.5	0.97969	0.80156	0.44227	0.094581	-0.17605	-0.37295	-0.51323	-0.61393	-0.68758	-0.74263
3	0.9864	0.86433	0.59528	0.26324	-0.012872	-0.22725	-0.38833	-0.50837	-0.59846	-0.667
3.5	0.99061	0.90061	0.72659	0.4246	0.1473	-0.077052	-0.25371	-0.39069	-0.49661	-0.579
4	0.99336	0.92295	0.81608	0.58678	0.30468	0.071609	-0.11684	-0.26771	-0.38774	-0.48326
4.5	0.99519	0.93799	0.86651	0.73993	0.47078	0.22019	0.018649	-0.14489	-0.27755	-0.38508
5	0.99644	0.9489	0.89597	0.83294	0.66205	0.38182	0.15661	-0.022741	-0.16846	-0.28759

2.5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	0.90456	0.32805	-0.15897	-0.45872	-0.63167	-0.73566	-0.80194	-0.84643	-0.8776	-0.90025
1	0.94729	0.52678	0.047679	-0.2923	-0.50647	-0.64115	-0.72905	-0.78888	-0.83119	-0.86209
1.5	0.96866	0.66921	0.23113	-0.12372	-0.37067	-0.53494	-0.64549	-0.7221	-0.77688	-0.81721
2	0.98023	0.77236	0.39015	0.03975	-0.22798	-0.41812	-0.55114	-0.64548	-0.71392	-0.7648
2.5	0.9869	0.84228	0.52712	0.19285	-0.082998	-0.29263	-0.44624	-0.55843	-0.64138	-0.70383
3	0.99098	0.88735	0.64456	0.33389	0.05969	-0.16146	-0.33186	-0.46079	-0.55845	-0.63319
3.5	0.9936	0.91602	0.74071	0.46437	0.19692	-0.028592	-0.21056	-0.35366	-0.46521	-0.55236
4	0.99535	0.9349	0.8107	0.58667	0.3283	0.10238	-0.086283	-0.24003	-0.36357	-0.4624
4.5	0.99654	0.9481	0.85638	0.6979	0.45669	0.23014	0.0373	-0.12401	-0.25718	-0.36629
5	0.99738	0.9579	0.88588	0.78416	0.58678	0.35736	0.15904	-0.0087104	-0.1498	-0.26771

3	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	0.94707	0.5096	0.033159	-0.32338	-0.52005	-0.65133	-0.73684	-0.795	-0.8361	-0.86612
1	0.96873	0.6531	0.20757	-0.14382	-0.38925	-0.54953	-0.65695	-0.73123	-0.78429	-0.82332
1.5	0.98037	0.75777	0.36351	0.011252	-0.25383	-0.43969	-0.5687	-0.65977	-0.72568	-0.77459
2	0.98707	0.83032	0.49557	0.15769	-0.11768	-0.32365	-0.47269	-0.58062	-0.66	-0.71954
2.5	0.99114	0.87895	0.60695	0.29086	0.015331	-0.20384	-0.36995	-0.49399	-0.58704	-0.65777
3	0.99373	0.91134	0.69917	0.41103	0.14228	-0.082998	-0.26197	-0.40034	-0.50667	-0.58883
3.5	0.99543	0.93311	0.77221	0.51975	0.26178	0.036231	-0.15074	-0.3007	-0.41914	-0.5125
4	0.9966	0.94812	0.82638	0.61809	0.3741	0.15188	-0.038653	-0.19679	-0.32539	-0.42909
4.5	0.99742	0.95889	0.86433	0.70442	0.48069	0.26324	0.072133	-0.090931	-0.22725	-0.33981
5	0.99801	0.96694	0.89056	0.77384	0.58292	0.37119	0.18043	0.014617	-0.12714	-0.24681

3.5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	0.96875	0.63407	0.18951	-0.15363	-0.39091	-0.55202	-0.66516	-0.73773	-0.78953	-0.82763
1	0.98043	0.74863	0.34497	-0.009685	-0.27039	-0.45317	-0.57949	-0.66847	-0.73279	-0.78047
1.5	0.98713	0.82349	0.4768	0.13618	-0.13838	-0.34162	-0.48765	-0.59297	-0.67023	-0.7281
2	0.9912	0.87418	0.58632	0.26606	-0.01019	-0.2275	-0.39059	-0.51154	-0.60188	-0.67036
2.5	0.99378	0.90837	0.67681	0.38208	0.11151	-0.11322	-0.28969	-0.42481	-0.5279	-0.60717
3	0.99548	0.93171	0.7496	0.48539	0.22516	-0.001036	-0.18658	-0.33362	-0.44859	-0.53851
3.5	0.99663	0.94794	0.80605	0.57728	0.33048	0.10724	-0.082998	-0.23907	-0.36446	-0.46448
4	0.99744	0.95947	0.84831	0.65829	0.42809	0.21059	0.019383	-0.14256	-0.27634	-0.38544
4.5	0.99802	0.96788	0.87915	0.72761	0.519	0.30886	0.11927	-0.045612	-0.18545	-0.30213
5	0.99845	0.9742	0.90151	0.7837	0.60372	0.4027	0.21602	0.050386	-0.093273	-0.21578

4	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	0.98046	0.74081	0.33296	-0.040662	-0.29022	-0.46481	-0.58802	-0.67541	-0.73842	-0.78513
1	0.98716	0.81882	0.46342	0.1203	-0.15253	-0.35437	-0.49811	-0.60151	-0.67727	-0.73396
1.5	0.99122	0.87123	0.57267	0.24913	-0.027455	-0.24315	-0.40397	-0.52276	-0.61128	-0.67828
2	0.9938	0.90659	0.66299	0.36351	0.09161	-0.13239	-0.30688	-0.43969	-0.54063	-0.61805
2.5	0.9955	0.93079	0.73599	0.4646	0.20231	-0.024133	-0.2083	-0.35312	-0.4657	-0.55341
3	0.99665	0.94766	0.79338	0.5537	0.3042	0.079939	-0.10978	-0.26405	-0.38704	-0.48457
3.5	0.99746	0.95967	0.83742	0.63178	0.39761	0.17875	-0.012757	-0.17362	-0.3053	-0.41184
4	0.99803	0.96837	0.87067	0.69917	0.48329	0.27188	0.081575	-0.082998	-0.22134	-0.3357
4.5	0.99846	0.97481	0.89559	0.75576	0.56193	0.35953	0.17246	0.0067006	-0.13613	-0.25682
5	0.99877	0.97969	0.91427	0.80156	0.63374	0.44227	0.25964	0.094581	-0.050699	-0.17605

4.5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	0.98718	0.81502	0.44491	0.10032	-0.16945	-0.36159	-0.50701	-0.60895	-0.68285	-0.73835
1	0.99123	0.86909	0.56173	0.23876	-0.038588	-0.24877	-0.41422	-0.53128	-0.61836	-0.68421
1.5	0.99381	0.90543	0.6531	0.34983	0.077021	-0.14382	-0.31917	-0.45019	-0.54953	-0.6256
2	0.9955	0.9302	0.72682	0.45018	0.18623	-0.040184	-0.22317	-0.36626	-0.4771	-0.56323
2.5	0.99666	0.94744	0.78513	0.53819	0.28643	0.061467	-0.12762	-0.28041	-0.40159	-0.49735
3	0.99746	0.9597	0.83032	0.615	0.37778	0.15769	-0.033839	-0.19365	-0.32365	-0.42832
3.5	0.99804	0.96858	0.86489	0.68133	0.46089	0.24797	0.057062	-0.10704	-0.24403	-0.35657
4		0.97513	0.89123	0.73758	0.53647	0.33229	0.14431	-0.021558	-0.16354	-0.28266
4.5		0.98004	0.91134	0.7842	0.605	0.41103	0.22752	0.061979	-0.082998	-0.20721
5		0.9838	0.92676	0.82192	0.66655	0.48468	0.30669	0.143	-0.0031901	-0.13094

5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	0.99124	0.86734	0.55698	0.20183	-0.059451	-0.27179	-0.42331	-0.5403	-0.62512	-0.68964
1	0.99381	0.90456	0.64501	0.32805	0.058236	-0.15897	-0.32922	-0.45872	-0.55671	-0.63167
1.5	0.99551	0.92979	0.71987	0.43811	0.1793	-0.052614	-0.2344	-0.37624	-0.48567	-0.57058
2	0.99666	0.94729	0.77923	0.52678	0.27315	0.047679	-0.14077	-0.2923	-0.41206	-0.50647
2.5	0.99747	0.95969	0.82543	0.60322	0.36351	0.14247	-0.048944	-0.20781	-0.33646	-0.43969
3		0.96866	0.86097	0.66921	0.44538	0.23113	0.039846	-0.12372	-0.25955	-0.37067
3.5		0.97528	0.88822	0.72538	0.51942	0.3136	0.12485	-0.040942	-0.18209	-0.29991
4		0.98023	0.90919	0.77236	0.58622	0.39015	0.20564	0.03975	-0.10481	-0.22798
4.5		0.984	0.92541	0.81098	0.64611	0.46118	0.2821	0.11779	-0.028406	-0.15547
5		0.9869	0.93807	0.84228	0.69917	0.52712	0.35437	0.19285	0.046535	-0.082998

3. The ratio between the inner and outer electron - $\tan \alpha$

	0.5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	1.1308	1.1356	1.1364	1.1367	1.1368	1.1369	1.1369	1.1369	1.137	1.137	1.137
1	1.4276	1.498	1.5451	1.5668	1.5777	1.584	1.5878	1.5903	1.5921	1.5933	
1.5	1.6387	2.1492	2.2456	2.2888	2.312	2.3257	2.3342	2.34	2.344	2.3469	
2	1.6425	2.186	3.2773	3.2658	3.2246	3.1996	3.1842	3.1743	3.1676	3.1628	
2.5	1.6264	2.1042	2.7489	4.7326	4.5053	4.3344	4.2261	4.1549	4.106	4.071	
3	1.6057	2.071	2.4545	3.4008	6.4124	5.9399	5.6162	5.4016	5.2541	5.1488	
3.5	1.5829	2.0519	2.3616	2.8151	3.9887	8.2731	7.5608	7.0716	6.7343	6.4942	
4	1.5584	2.0395	2.3142	2.6404	3.1691	10.1026	10.1111	9.3671	8.7044	8.2319	
4.5	1.5311	2.0309	2.2853	2.5532	2.9142	3.4869	4.6429	7.2826	10.1111	10.1111	
5	1.4977	2.0246	2.266	2.5007	2.7846	3.172	3.7527	4.7846	7.7297	8.1248	

	1	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	1.057	1.0599	1.0594	1.0591	1.059	1.0589	1.0588	1.0588	1.0588	1.0587	
1	1.1071	1.1308	1.1344	1.1356	1.1361	1.1364	1.1366	1.1367	1.1368	1.1368	
1.5	1.1719	1.243	1.2635	1.2772	1.2852	1.29	1.2931	1.2952	1.2966	1.2977	
2	1.2069	1.4276	1.46	1.498	1.5262	1.5451	1.5579	1.5668	1.5731	1.5777	
2.5	1.2048	1.5982	1.7347	1.7865	1.8322	1.8657	1.8894	1.9062	1.9184	1.9275	
3	1.1837	1.6387	1.9431	2.1492	2.2093	2.2456	2.2708	2.2888	2.3021	2.312	
3.5	1.1531	1.6455	1.9348	2.3209	2.6657	2.7138	2.726	2.7313	2.7346	2.7371	
4	1.1196	1.6425	1.9218	2.186	2.8158	3.2773	3.2894	3.2658	3.2429	3.2246	
4.5	1.0888	1.6354	1.9112	2.1334	2.4514	3.4861	3.9714	3.9273	3.8601	3.8031	
5	1.0647	1.6264	1.9024	2.1042	2.3363	2.7489	4.4053	4.7326	4.6211	4.5053	

	1.5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	1.0341	1.0416	1.0413	1.041	1.0409	1.0408	1.0407	1.0407	1.0406	1.0406	
1	1.048	1.0785	1.0803	1.0803	1.0801	1.0799	1.0798	1.0798	1.0797	1.0797	
1.5	1.0603	1.1226	1.1308	1.1336	1.1349	1.1356	1.136	1.1363	1.1364	1.1365	
2	1.0717	1.1826	1.1994	1.2094	1.2162	1.2207	1.2237	1.2257	1.2271	1.2282	
2.5	1.0729	1.2621	1.2948	1.3124	1.3294	1.3428	1.3526	1.3598	1.3651	1.369	
3	1.0654	1.3252	1.4276	1.4472	1.4733	1.498	1.5179	1.5334	1.5451	1.5542	
3.5	1.0545	1.3558	1.5617	1.6205	1.6497	1.6822	1.7111	1.7346	1.7531	1.7676	
4	1.0437	1.3663	1.6192	1.7946	1.8611	1.9	1.9338	1.962	1.9846	2.0027	
4.5	1.0345	1.3656	1.6387	1.8413	2.0478	2.1492	2.1932	2.2231	2.2456	2.2634	
5		1.3581	1.6449	1.8458	2.0453	2.3438	2.4826	2.5255	2.5461	2.559	

	2	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	1.0197	1.0321	1.0325	1.0323	1.0322	1.0321	1.032	1.0319	1.0319	1.0319	
1	1.0231	1.057	1.0599	1.0599	1.0596	1.0594	1.0592	1.0591	1.059	1.059	
1.5	1.0248	1.081	1.09	1.0916	1.0919	1.0919	1.0919	1.0918	1.0918	1.0917	
2	1.0272	1.1071	1.1258	1.1308	1.1331	1.1344	1.1351	1.1356	1.1359	1.1361	
2.5	1.0285	1.1389	1.1709	1.1801	1.1864	1.1908	1.1939	1.196	1.1976	1.1988	
3	1.0271	1.1719	1.2302	1.243	1.2538	1.2635	1.2712	1.2772	1.2817	1.2852	
3.5	1.024	1.1954	1.3038	1.3243	1.3378	1.353	1.367	1.3788	1.3883	1.3959	
4	1.0205	1.2069	1.3693	1.4276	1.4414	1.46	1.4798	1.498	1.5134	1.5262	
4.5	1.0172	1.2091	1.4098	1.5349	1.5679	1.5869	1.6101	1.6333	1.654	1.6717	
5	1.0143	1.2048	1.4312	1.5982	1.7022	1.7347	1.7604	1.7865	1.8108	1.8322	

	2.5	0.5	1	1.5	2	2.5	3	3.5	4	4.5	5
0.5	1.0106	1.0259	1.0268	1.0269	1.0268	1.0267	1.0266	1.0266	1.0266	1.0265	
1	1.0115	1.0439	1.0485	1.0488	1.0486	1.0484	1.0482	1.0481	1.048	1.0479	
1.5	1.0115	1.058	1.0698	1.0716	1.0718	1.0717	1.0715	1.0714	1.0713	1.0712	
2	1.0118	1.0704	1.0927	1.0973	1.0987	1.0992	1.0994	1.0995	1.0995	1.0995	
2.5	1.0123	1.0838	1.1186	1.1272	1.1308	1.1328	1.134	1.1347	1.1353	1.1356	
3	1.0123	1.099	1.1497	1.163	1.1693	1.1737	1.177	1.1793	1.1811	1.1824	
3.5	1.0117	1.1123	1.1874	1.2073	1.216	1.2233	1.2297	1.2348	1.2389	1.2421	
4	1.0106	1.1204	1.2273	1.2628	1.273	1.2828	1.2926	1.3016	1.3092	1.3155	
4.5	1.0094	1.1228	1.2601	1.3283	1.3432	1.3537	1.3664	1.3793	1.3911	1.4013	
5	1.0082	1.1206	1.2824	1.39	1.4276	1.4383	1.4521	1.468	1.4836	1.498	

4. Energy in a.u.

5. Residuals

