On concentration for (regularized) empirical risk minimization

Sara van de Geer Seminar for Statistics, ETH Zürich

and

Martin J. Wainwright Department of Statistics and Department of EECS University of California, Berkeley

Abstract. Rates of convergence for empirical risk minimizers have been well studied in the literature. In this paper, we aim to provide a complementary set of results, in particular by showing that after normalization, the risk of the empirical minimizer concentrates on a single point. Such results have been established by Chatterjee [2014] for constrained estimators in the normal sequence model. We first generalize and sharpen this result to regularized least squares with convex penalties, making use of a "direct" argument based on Borell's theorem. We then study generalizations to other loss functions, including the negative log-likelihood for exponential families combined with a strictly convex regularization penalty. The results in this general setting are based on more "indirect" arguments as well as on concentration inequalities for maxima of empirical processes.

1 Introduction

Empirical risk minimization (ERM) is an important methodology in statistics and machine learning, widely used for estimating high-dimensional and/or nonparametric parameters of interest. The idea is to express the parameter as minimizer of an expected loss, the so-called population or theoretical risk. Given that the distribution of data is not known or difficult to assess, one replaces the theoretical expectation by an empirical counterpart defined by samples. The technique of ERM is known under various names, including M-estimation and minimum contrast estimation.

By the law of large numbers, empirical averages of various types of random variables, with the i.i.d. setting being the canonical case, are close to their expectation. This elementary fact is the motivation for ERM and the starting point for studying its theoretical properties. There is much literature developing the theory for a broad spectrum of estimation problems. The more recent literature takes a non-asymptotic point of view, in which context concentration inequalities play a major role. Concentration inequalities describe the amount of concentration of certain (complex) quantities around their mean. We refer to Talagrand [1995] as a key paper in the area, and to the important monographs Ledoux [2001] and more recently Boucheron et al. [2013]. The key point is that the deviation from the mean is generally of much smaller order than the mean itself. Moreover, at least in a certain sense, the deviation does not depend on the

complexity of the original object. In statistics, the usefulness of concentration inequalities has been excellently outlined and studied in Massart [2000]. We also refer the reader to Koltchinskii [2011] for an in-depth treatment in the context of high-dimensional problems.

Some statistical papers address concentration for the parameter of interest itself; for instance, see Boucheron and Massart [2011] and Saumard [2012]. The present paper is along the lines of Chatterjee [2014]. The latter examines the concentration properties of constrained estimators for the normal sequence model, or alternatively phrased in the regression setting, for the least-squares problem with fixed design and Gaussian errors. The author shows that the statistical error of the least squares estimator satisfies a concentration inequality where the amount of concentration still depends on the complexity of the problem, but is in the nonparametric case of smaller order than the statistical error itself. In Muro and van de Geer [2015], the situation is studied where a regularization penalty based on a squared norm is added to the least squares loss function. In Section 2, we provide a "direct" argument for concentration of the regularized least squares in the normal sequence setting. Our argument here is elementary, using standard facts from convex analysis [Rockafellar, 1970], and concentration for Lipschitz functions of Gaussian vectors [Borell, 1975]. Our next goal is to extend such results to more general problems. The main obstacle is that the direct concentration for Lipschitz functions holds only for the Gaussian case. Accordingly, we make use of more general one-sided concentration results for maxima of empirical processes, as given by Klein [2002] and Klein and Rio [2005].

Our theory allows us to treat a number of new examples in which concentration holds. However, as (asymptotically) exact values for the expectation of maxima of the empirical process are generally not available, we cannot always provide explicit expressions for the point of concentration in terms of the parameters of the model.

Set-up and notation

Consider independent observations X_1, \ldots, X_n taking values in a space \mathcal{X} , a given class \mathcal{F} of real-valued functions on \mathcal{X} and a non-negative regularization penalty pen : $\mathcal{F} \to [0, \infty)$. The empirical measure P_n of a function $f : \mathcal{X} \to \mathbb{R}$ is defined by the average $P_n f := \frac{1}{n} \sum_{i=1}^n f(X_i)$, whereas the theoretical or population measure is given by $Pf := \mathbb{E}P_n f$.

We let \mathcal{F} denote a class of loss functions, say indexed by a parameter g in a parameter space \mathcal{G} . As a concrete example, in the case of least-squares regression, the observations consist of covariates along with real-valued responses of the form $\{(X_i, Y_i)\}_{i=1}^n$. In the least-squares case, the loss class takes the form $\mathcal{F} = \{f_g(x, y) = (y - g(x))^2 : g \in \mathcal{G}\}$, where \mathcal{G} is some underlying collection of regression functions.

With this set-up, the regularized empirical risk estimator is defined¹ as

$$\hat{f} = \arg\min_{f \in \mathcal{F}} \bigg\{ P_n f + \operatorname{pen}(f) \bigg\}.$$
(1)

We define the associated target function $f^0 := \arg \min_{f \in \mathcal{F}} Pf$, corresponding to the population minimizer, and we let

$$\tau^{2}(f) := P(f - f^{0}) + \operatorname{pen}(f)$$
(2)

the penalized excess risk.

In order to simplify the exposition, we give asymptotic statements at places, using the classical scaling in which the sample size n tends to infinity. For a sequence of positive numbers $\{z_n\}_{n=1}^{\infty}$, we write

$$z_n = \mathcal{O}(1)$$
 if $\limsup z_n < \infty$, and $z_n = o(1)$ if $z_n \to 0$,

as well as $z_n \approx 1$ if both $z_n = \mathcal{O}(1)$ and $1/z_n = \mathcal{O}(1)$. For two positive sequences $\{y_n\}_{n=1}^{\infty}$ and $\{z_n\}_{n=1}^{\infty}$, we write $z_n = \mathcal{O}(y_n)$ if $z_n/y_n = \mathcal{O}(1)$, along with analogous definitions for the *o*- and \approx -notation. We furthermore use the stochastic order symbols are $\mathcal{O}_{\mathbf{P}}$ and $o_{\mathbf{P}}$. In all our uses of these forms of order notation, the arguments depend on *n*, but we often omit this dependence to simplify notation.

With this set-up, the main results of this paper involve showing that under certain conditions, we have

$$\left|\tau(\hat{f}) - s_0\right| = o_{\mathbf{P}}(s_0)$$

Here s_0 is a deterministic quantity defined by the problem under consideration; see equation (8) for its precise definition. In our context, the result requires the complexity of the problem to be in the nonparametric regime in which $\sqrt{\log n/n} = o(s_0)$. When a certain concavity condition is met, the log *n*-term can be removed. This concavity condition holds in the normal sequence model, as well as in all the examples given in Section 6. In Section 2, there is no log *n*term as well, but the concentration result there is for $\sqrt{P(\hat{f} - f^0)}$ as opposed to $\tau(\hat{f})$.

Organization

The remainder of this paper is organized as follows. In Section 2, we provide a concentration result for the normal sequence model and least squares with convex penalty, based on a "direct" argument. We then consider more general models and loss functions, using the more indirect route originally taken

¹In order to avoid digressions, we assume throughout this paper that "argmin's" and "argmax's" exist and are unique in a suitable sense.

by Chatterjee [2014]. In Section 3, we discuss the deterministic counterpart of empirical risk minimization, corresponding the population-level optimization problem. Our theory requires a certain amount of curvature of the objective function around its minimum, a requirement that we term a second order margin condition. In Section 4, we present a concentration result (Theorem 4.1) for a general loss function. Section 5 is devoted to a more careful analysis of quadratic second-order margin conditions. Section 6 is devoted to the detailed analysis of two examples in which the empirical process is linear in its parameter—projection estimators for densities and linearized least squares whereas Section 7 provides results for nonparametric estimation involving exponential families. In Section 8, we present the concentration inequalities that underlie the proof of our indirect approach. In Section 9, we provide a similar result as in Section 4 but now for a shifted version of $\tau^2(\hat{f})$. Finally, all proofs are in provided in Section 10.

2 Direct approach to normal sequence model

In this section, we analyze the concentration properties of regularized leastsquares estimators in the normal sequence setting. The main contribution of this section is to provide a direct argument that generalizes and sharpens the previous result of Chatterjee [2014].

Let $Y_i \in \mathbb{R}$ be a response variable and X_i be a fixed co-variable in some space $\mathcal{X}, i = 1, \ldots, n$. The normal sequence model is given by

$$Y_i = g^0(X_i) + \epsilon_i \qquad \text{for } i = 1, \dots, n, \tag{3}$$

where $\epsilon_1, \ldots, \epsilon_n$ are i.i.d. mean-zero Gaussians with variance σ^2 , and the regression vector $g^0 := (g^0(X_1), \ldots, g^0(X_n))^T$ is unknown. Let us write the vector of responses as $Y = (Y_1, \ldots, Y_n)^T$, and the noise vector as $\epsilon := (\epsilon_1, \ldots, \epsilon_n)$.

Let pen : $\mathbb{R}^n \to \mathbb{R} \cup \{+\infty\}$ be a complexity penalty, assumed to be convex. The regularized least squares estimator is given by

$$\hat{g} := \arg\min_{g \in \mathbb{R}^n} \bigg\{ \|Y - g\|_n^2 + \operatorname{pen}(g) \bigg\},\tag{4}$$

where for any vector $v \in \mathbb{R}^n$, we use the standard notation $||v||_n^2 := \frac{v^T v}{2} = \frac{||v||_2^2}{n}$. In past work, Chatterjee [2014] analyzed the concentration of the constrained variant of this estimator, given by

$$\hat{g} := \arg\min_{g \in \mathcal{G}} \left\{ \|Y - g\|_n^2 \right\},\tag{5}$$

where $\mathcal{G} \subseteq \mathbb{R}^n$ is a closed, convex set. Note that this constrained estimator (5) is a special case of the regularized estimator (4), in which the penalty function

takes the form

$$pen(g) := \begin{cases} 0 & \text{if } g \in \mathcal{G} \\ +\infty & \text{otherwise.} \end{cases}$$
(6)

The following result guarantees that with for any convex penalty, the estimation error $\|\hat{g} - g^0\|_n$ of the regularized estimator (4) is sharply concentrated around its expectation $m_0 := \mathbb{E}\|\hat{g} - g^0\|_n$.

Theorem 2.1 For any convex penalty pen : $\mathbb{R}^n \to \mathbb{R} \cup \{+\infty\}$, the error in the regularized estimator (4) satisfies

$$\mathbb{P}\left(\left|\|\hat{g} - g^0\|_n - m_0\right| \ge \sigma \sqrt{2t/n}\right) \le \exp[-t] \quad \text{for all } t > 0.$$
(7)

See Section 10.1 for the proof of this claim. The argument is direct, using some basic facts from convex analysis, and the concentration of Lipschitz functions of Gaussian vectors (see Borell [1975]).

Remarks: In terms of asymptotic behaviour, it follows that when $\sigma = \mathcal{O}(1)$ and $1/\sqrt{n} = o(m_0)$ —the latter condition corresponding to the non-parametric regime—it holds that

$$\left| \|\hat{g} - g^0\|_n - m_0 \right| = o_{\mathbf{P}}(m_0).$$

Moreover, it follows from its proof that Theorem 2.1 remains true if the population minimizer g^0 is replaced by any other vector $g \in \mathbb{R}^n$. Thus, for instance, we can take g as the minimizer of the penalized noiseless problem

$$g^* := \arg\min_{g \in \mathbb{R}^n} \left\{ \|g - g^0\|_n^2 + \operatorname{pen}(g) \right\}$$

With this choice, we also have concentration of $\|\hat{g} - g^*\|_n$ around its expectation $\mathbb{E}\|\hat{g} - g^*\|_n$.

With the choice (6), the result also applies to the constrained least squares estimate (5), for any closed convex set $\mathcal{G} \subseteq \mathbb{R}^n$. In this context, Theorem 2.1 sharpens the previous result of Chatterjee [2014].

3 Theoretical version of minimization problem

For the remainder of the paper, we study the general empirical risk minimizer. Let $\tau(\hat{f})$ be the excess risk (2) associated with the empirical minimizer \hat{f} from equation (1). We define the minimum possible excess risk

$$\tau_{\min}^2 := \min_{f \in \mathcal{F}} \tau^2(f) = \min_{f \in \mathcal{F}} \bigg\{ P(f - f^0) + \operatorname{pen}(f) \bigg\}.$$

For all $s \ge \tau_{\min}$, we define the set $\mathcal{F}_s := \{f \in \mathcal{F} \mid \tau(f) \le s\}$, and the functions

$$\hat{\mathbf{E}}_n(s) = \max_{f \in \mathcal{F}_s} (P_n - P)(f^0 - f), \text{ and } \mathbf{E}(s) := \mathbb{E}(\hat{\mathbf{E}}_n(s)).$$

In addition, we define the minimizers

$$\hat{s} := \arg\min_{s \ge \tau_{\min}} \{ s^2 - \hat{\mathbf{E}}_n(s) \}, \text{ and } s_0 := \arg\min_{s \ge \tau_{\min}} \{ s^2 - \mathbf{E}(s) \}.$$
 (8)

From hereon, we refer to minimizing the function $s \mapsto s^2 - \mathbf{E}(s)$ as the theoretical problem, and to minimizing $s \mapsto s^2 - \hat{\mathbf{E}}_n(s)$ as the empirical problem. That the latter indeed yields the risk associated with the original ERM estimate (1) is guaranteed by the following lemma.

Lemma 3.1 We have $\tau(\hat{f}) = \hat{s}$.

See Section 10.2 for the proof of this claim.

In order to prove concentration results, we need a certain amount of curvature of the function $s \mapsto s^2 - \mathbf{E}(s)$ around its minimum. The following condition is known as a second-order margin condition; we also introduce a first-order margin condition in Definition 4.1.

Definition 3.1 Let G be a strictly convex increasing function with G(0) = 0and let $\underline{\delta} \geq 0$ and $\tau_{\max} \geq s_0 + \underline{\delta}$ be constants. If $s_0 > \tau_{\min}$ we say that the second order margin condition holds in the range $[\tau_{\min}, s_0) \cup (s_0 + \underline{\delta}, \tau_{\max}]$ with margin function G if for all $s \in [\tau_{\min}, s_0) \cup (s_0 + \underline{\delta}, \tau_{\max}]$

$$[s^{2} - \mathbf{E}(s)] - [s_{0}^{2} - \mathbf{E}(s_{0})] \ge G(|s - s_{0}|).$$
(9)

If inequality (9) is only true for all $s \in (s_0 + \underline{\delta}, \tau_{\max}]$, then we say that the right-sided second order margin condition holds in the range $(s_0 + \underline{\delta}, \tau_{\max}]$ with margin function G. If no reference is made to any range, it means that the condition holds $\underline{\delta} = 0$ and $\tau_{\max} = \infty$.

In the two-sided case in Definition 3.1, we do not allow for a gap for values of s to the left of s_0 . This choice is merely for simplicity and corresponds to our examples.

An important special case is quadratic margin behaviour, as formalized in the following:

Definition 3.2 If $s_0 > \tau_{\min}$ we say that the second order quadratic margin condition is met in the range $[\tau_{\min}, s_0) \cup (s_0 + \underline{\delta}, \tau_{\max}]$ with margin constant c > 0 if for all $s \in [\tau_{\min}, s_0) \cup (s_0 + \underline{\delta}, \tau_{\max}]$, it holds that

$$\left[s^{2} - \mathbf{E}(s)\right] - \left[s_{0}^{2} - \mathbf{E}(s_{0})\right] \ge (s - s_{0})^{2}/c^{2}.$$

If this is only true for $s \in (s_0 + \underline{\delta}, \tau_{\max}]$, then we say that the right-sided second order quadratic margin condition holds in the range $(s_0 + \underline{\delta}, \tau_{\max}]$ with margin constant c > 0.

 $^{^{2}}$ Recall that throughout this paper, we tacitly assume that the minimizers are unique.

When the two-sided condition in Definition 3.2 holds with $\underline{\delta} = 0$, then it corresponds to a form of *strong convexity* of the function $s \mapsto s^2 - \mathbf{E}(s)$ at s_0 .

Clearly, if $s \mapsto \mathbf{E}(s)$ is concave then $s \mapsto s^2 - \mathbf{E}(s)$ is strictly convex. In fact, then the second order quadratic margin condition holds with margin constant cat most equal to 1. This type of condition holds in the normal sequence setting, as exploited by Chatterjee [2014]. In the latter paper, the map $s \mapsto \hat{\mathbf{E}}_n(s)$ is concave and hence then also $s \mapsto \mathbf{E}(s)$ is concave. Moreover, the empirical function $s \mapsto s^2 - \hat{\mathbf{E}}_n(s)$ is then convex, which allows one to remove the log *n*factor. We will consider conditions for (right-sided) second order quadratic margin behaviour in Section 5.

4 Concentration of ERM

We now turn to the statement of our main result on concentration of ERM in the general setting. We begin by specifying some conditions that underlie the result. First, we require a uniform boundedness condition:

Condition 4.1 The function class \mathcal{F} is uniformly bounded, meaning that

$$K := \max_{f \in \mathcal{F}} \|f - f^0\|_{\infty} < \infty.$$

We note that this condition can be removed if one first shows that, for a suitable constant K, the minimizer \hat{f} satisfies the bound $\|\hat{f} - f^0\|_{\infty} \leq K$ with high probability.

When Condition 4.1 holds, one may take

$$\tau_{\max}^2 := 2K + \operatorname{pen}(f^0).$$

However, in order to obtain a sharper result, one may first want to prove that $\tau^2(\hat{f})$ is much smaller than $2K + \text{pen}(f^0)$ with high probability. In fact, there is a substantial literature on techniques for showing that $\tau(\hat{f}) = \mathcal{O}_{\mathbf{P}}(s_0)$ for various problems. As we discuss in Section 9, similar results exist for the shifted version, in particular showing that $\tau^2(\hat{f}) - \tau_*^2 = \mathcal{O}_{\mathbf{P}}(s_0^2 - \tau_*^2)$, where τ_*^2 is a suitably chosen constant.

We now come to our "first order" margin condition, which quantifies the curvature of $f \mapsto Pf$ around its minimum. To avoid confusion with the "second order" margin condition we call it a "curvature condition".

For $f \in \mathcal{F}$, define the variance

$$\sigma^{2}(f) := \frac{1}{n} \sum_{i=1}^{n} \left[\mathbb{E}f^{2}(X_{i}) - (\mathbb{E}f(X_{i}))^{2} \right].$$

Definition 4.1 A quadratic curvature condition with constant C > 0 is said to hold if

$$P(f - f^0) \ge \frac{\sigma^2(f - f^0)}{C^2} \quad for \ all \ f \in \mathcal{F}.$$
 (10)

We take the quadratic curvature condition as basis for our results. An extension to more general curvature is omitted here to avoid digressions.

Let \mathcal{J} be a strictly increasing function defined on $[\tau_{\min}, \infty)$ and such that $\mathcal{J}(\tau_{\min}) = 0$. We then define a new function

$$\Phi_{\mathcal{J}}(u) := [\mathcal{J}^{-1}(u)]^2 \quad \text{for all } u > 0.$$
(11)

Our choice of the square here is linked to the quadratic curvature condition (10).

Condition 4.2 There is a constant m_n , a strictly increasing function \mathcal{J} such that the function $\Phi_{\mathcal{J}}$ is strictly convex, and such that the bound

$$\mathbf{E}(s) \le \frac{\mathcal{J}(s)}{m_n} \tag{12}$$

holds for all $s \geq \tau_{\min}$.

We also define the convex conjugate function $\Phi^*_{\mathcal{J}}(v) := \sup_{u>0} \Big\{ vu - \Phi_{\mathcal{J}}(u) \Big\}$, and make use of the Fenchel-Young inequality

$$uv \leq \Phi_{\mathcal{J}}(u) + \Phi^*_{\mathcal{J}}(v),$$
 valid for all pairs $u, v > 0.$

Finally, in terms of the previously defined quantities, we define

$$r_0^2 := 2C^2 \Phi_{\mathcal{J}}^* (4K/(m_n C^2))$$

With this notation, the following theorem is our main result:

Theorem 4.1 Suppose that:

- Conditions 4.1 and 4.2, as well as the quadratic curvature condition with constant C hold.
- The right-sided second order margin condition holds in the range $(s_0 + \underline{\delta}, \tau_{\max}]$ with margin function G, with associated convex conjugate G^* .

Then there is a constant $c_0 = c_0(C, K)$, and a function $\delta : (0, \infty) \to (0, \infty)$ such that

$$G(\delta(t)) \le G^* \left(c_0 \sqrt{[t + \log(1 + \sqrt{n\tau_{\max}^2})]/n}) \right) + c_0 \left((s_0 + r_0) \sqrt{[t + \log(1 + \sqrt{n\tau_{\max}^2})]/n} + [t + \log(1 + \sqrt{n\tau_{\max}^2})]/n \right),$$

and such that the following deviation inequality holds:

$$\mathbb{P}\left(\tau_{\max} \ge \tau(\hat{f}) > s_0 + \max\{\underline{\delta}, \delta(t)\}\right) \le \exp[-t] \quad \text{for all } t > 0.$$

If, in fact, the two-sided version of the second order margin condition holds over $[\tau_{\min}, s_0) \cup (s_0 + \underline{\delta}, \tau_{\max}]$, then we have

$$\mathbb{P}\left(|\tau(\hat{f}) - s_0| > \max\{\underline{\delta}, \delta(t)\}, \ \tau(\hat{f}) \le \tau_{\max}\right) \le 2\exp[-t] \quad \text{for all } t > 0.$$

Asymptotics: If the second order margin condition holds with the quadratic function $G(u) = \frac{u^2}{2c^2}$, then its convex conjugate $G^*(v) = \frac{c^2v^2}{2}$ is also quadratic. Thus, under the scalings $C = \mathcal{O}(1)$, $K = \mathcal{O}(1)$ and $r_0 \approx s_0$, we then find that

$$\delta(t) = \mathcal{O}\left((\log n/n)^{1/2} + (s_0^2 \log n/n)^{1/4} \right)$$

for each fixed t. Hence, whenever $\sqrt{\frac{\log n}{n}} = o_{\mathbf{P}}(s_0)$, then we are guaranteed that $\delta(t) = o_{\mathbf{P}}(s_0)$.

5 Second order quadratic margin behaviour

In this section, we investigate conditions under which the (right-sided) second order quadratic margin condition holds over an appropriate range. In particular, we extend the setting of Chatterjee [2014] to the case where one has a strictly convex penalty in Lemma 5.1, and to approximate forms of concavity in Lemmas 5.2 and 5.3. We note that it is possible to formulate different results with other combinations of conditions, but we omit this here.

Lemma 5.1 Let $\mathcal{F} := \{f_g \mid g \in \mathcal{G}\}$ be a class of loss functions indexed by a parameter g in a parameter space \mathcal{G} . Assume \mathcal{G} is a convex subset of a linear vector space, the mapping $g \mapsto f_g - Pf_g$ is linear, and that for some q > 1, the mapping $g \mapsto \tau^{2/q}(f_g)$ is convex. For some constant M > 0, define $\tau_{\max} := (M+1)s_0$. Then the right-sided second order quadratic margin condition holds in the range $(s_0, \tau_{\max}]$ with constant

$$c = \sqrt{2q^{-1}(q-1)(M+1)^{-\frac{2(2-q)}{q}}}$$

Moreover, when q = 2 and $s_0 > \tau_{\min}$, then the (two-sided) second order quadratic margin condition holds with c = 1.

We note that the latter two-sided second order quadratic margin condition corresponds to the favourable setting of the normal sequence model, as studied by Chatterjee [2014].

Asymptotics: The idea in the above lemma is that one first proves by separate means that $\tau(\hat{f}) = \mathcal{O}_{\mathbf{P}}(s_0)$. There is a large literature on bounds of this form; for example, see Koltchinskii [2011] and references therein. One can then take $M = \mathcal{O}(1)$.

We sometimes write $\hat{\mathbf{E}}_n(\cdot) =: \hat{\mathbf{E}}_n^{\tau}(\cdot)$ and $\mathbf{E}(\cdot) =: \mathbf{E}^{\tau}(\cdot)$ so as to highlight their dependence on τ . For $f \in \mathcal{F}$, define the functionals

$$\varsigma^2(f) := c^2 \sigma^2(f - f^0) + pen(f), \text{ and } \varsigma_{\min} := \min_{f \in \mathcal{F}} \varsigma(f),$$

where c > 0 is some constant. Moreover, for $s \ge \varsigma_{\min}$, let us define

$$\hat{\mathbf{E}}_{n}^{\varsigma}(s) := \max_{f \in \mathcal{F}: \ \varsigma(f) \le s} (P_{n} - P)(f^{0} - f), \text{ and } \mathbf{E}^{\varsigma}(s) := \mathbb{E}\hat{\mathbf{E}}_{n}^{\varsigma}(s).$$

Lemma 5.2 Suppose that the function $s \mapsto \mathbf{E}^{\varsigma}(s)$ is concave, and that

$$\frac{\varsigma^2(f)}{A} \leq \tau^2(f) \leq A\varsigma^2(f), \quad \text{for all } f \in \mathcal{F}, \text{ and } \tau_f \leq \tau_{\max},$$

where $A^2 = 1 + \epsilon$ for some $\epsilon > 0$ satisfying $\sqrt{\epsilon}(1+\epsilon) < 1/2$. Let $\tau_{\max} := (M+1)s_0$ for some M > 0 and

$$\underline{\delta} := 2 \left[\sqrt{\epsilon} (2\sqrt{\epsilon}M + 1) \right]^{1/2} s_0.$$

Then when $s_0 > \tau_{\min}$, the quadratic second order margin condition holds in the range $[\tau_{\min}, s_0) \cup (s_0 + \underline{\delta}, \tau_{\max}]$ with constant c = 4.

Asymptotics As in Lemma 5.1 one may first prove by separate means that $\tau(\hat{f}) = \mathcal{O}_{\mathbf{P}}(s_0)$ and then take $M = \mathcal{O}(1)$.

Lemma 5.2 requires the function \mathbf{E}^{ς} to be concave. We now present conditions under which this is indeed the case.

Lemma 5.3 Let $\mathcal{F} := \{f_g \mid g \in \mathcal{G}\}$ be a class of loss functions indexed by the parameter g in a parameter space \mathcal{G} . Assume \mathcal{G} is a convex subset of a linear vector space, and that $g \mapsto \sqrt{\operatorname{pen}(f_g)}$ is convex and $g \mapsto f_g - Pf_g$ is linear. Then the function $s \mapsto \hat{\mathbf{E}}_n^{\varsigma}(s)$ is concave.

In fact, we show concavity of the empirical version $\hat{\mathbf{E}}_{n}^{\varsigma}$, which then implies concavity of $\mathbf{E}_{n}^{\varsigma}$. The reasoning is along the lines of Chatterjee [2014], and of the corresponding part of the proof of Lemma 5.1.

6 Some "pure" cases

In this section, we examine a number of problems that are "pure" in the sense that the empirical process enters in a linear manner. The simplest example of such a pure case is the normal sequence model studied in Section 2, and we examine some other examples here.

More precisely, consider a class of the form $\mathcal{F} := \{f_g \mid g \in \mathcal{G}\}$, where \mathcal{G} is a convex subset of a normed linear vector space $(\bar{\mathcal{G}}, \|\cdot\|)$. The pure case corresponds to problems in which the mapping $g \mapsto f_g - Pf_g$ is linear, and moreover, we have $P(f_g - f^0) = \|g - g^0\|^2$, where $g_0 = \arg\min_{g \in \mathcal{G}} Pf_g$, which ensures the equivalence $f^0 = f_{g^0}$.

6.1 Density estimation using projection

Let X_1, \ldots, X_n be i.i.d. random variables with distribution P taking values in a space \mathcal{X} . For a sigma-finite measure ν on \mathcal{X} , let $\|\cdot\|$ denote the $L^2(\nu)$ -norm. Let \mathcal{G} be a convex subset of a linear vector space $\overline{G} \subset L^2(\nu)$, and suppose that density $g^0 := dP/d\nu$ is a member of the model class \mathcal{G} . With this set-up, we consider the estimator

$$\hat{g} := \arg\min\left\{-P_n g + \frac{\|g\|^2}{2} + \lambda^2 I^q(g)\right\},\$$

where I denotes some pseudo-norm on \overline{G} , the exponent $q \in (1, 2]$, and $\lambda \ge 0$ is a regularization parameter.

In order to analyze the concentration properties of this estimator using our general theory, we begin by casting it within our framework. For each $g \in \mathcal{G}$, define $f_g := -g + \frac{1}{2} ||g||^2$, as well as the associated function class $\mathcal{F} := \{f_g \mid g \in \mathcal{G}\}$. With these choices, for all $g \in \mathcal{G}$, we have $(P_n - P)f_g = -(P_n - P)g$, and moreover

$$P(f_g - f_{g_0}) = -P(g - g^0) + \frac{\|g\|^2}{2} - \frac{\|g^0\|^2}{2} = \|g - g^0\|^2.$$

We split our analysis into several cases, depending on the nature of the penalty I.

6.1.1 Case 1: No penalty

In this case, we assume: \circ Condition 4.1 holds with $K = \mathcal{O}(1)$, $\circ \mathbf{E}(s) \leq \mathcal{J}(s)/\sqrt{n}$, $\circ \mathcal{J}(s) = As^{1-\alpha}, \exists A = \mathcal{O}(1), \exists 0 < \alpha < 1$ not depending on n, $\circ s_0 \simeq n^{-\frac{1}{2(1+\alpha)}}$.

It then follows from Theorem 4.1 combined with Lemma 5.1 that

$$\left|\frac{\|\hat{g} - g^0\| - s_0}{s_0}\right| = \mathcal{O}_{\mathbf{P}}\left(\sqrt{\frac{\log n}{n^{\frac{\alpha}{1+\alpha}}}}\right) = o_{\mathbf{P}}(1).$$

In fact, in this case, the log *n*-term can be removed because Lemma 5.3 ensures that the map $s \mapsto \hat{\mathbf{E}}_n(s)$ is concave.

6.1.2 Case 2: Quadratic penalty

In this case, we assume: $\circ q = 2,$ \circ Condition 4.1 with $K = \mathcal{O}(1),$ $\circ \mathbf{E}(s) \leq \mathcal{J}(s)/m_n,$ $\circ \mathcal{J}(s) = As, m_n = \sqrt{n}\lambda^{\alpha}, \exists A = \mathcal{O}(1), \exists 0 < \alpha < 1 \text{ not depending on } n,$ $\circ s_0 \approx 1/(\sqrt{n}\lambda^{\alpha}),$ $\circ \lambda^{2\alpha} \log n = o(1).$

Then Theorem 4.1 combined with Lemma 5.1 implies that

$$\left|\frac{\tau(\hat{f}) - s_0}{s_0}\right| = \mathcal{O}_{\mathbf{P}}\left(\sqrt{\lambda^{2\alpha}\log n}\right) = o_{\mathbf{P}}(1)$$

As before, the $\log n$ -factor can be removed.

One sees from the condition $\lambda^{2\alpha} \log n = o(1)$ that the regularization parameter λ must be sufficiently small. Moreover, hiding in our conditions is the fact that the penalty $I(f^0)$ is not too large. Indeed, we have $s_0^2 \simeq \frac{1}{n\lambda^{2\alpha}} \geq \tau_{\min}^2$. Consequently, when $\tau_{\min}^2 \simeq \operatorname{pen}(f^0) = \lambda^2 I^2(f^0)$, we must have

$$I^{2}(f^{0}) = \mathcal{O}\left(\left(n\lambda^{2(1+\alpha)}\right)^{-1}\right).$$
(13)

As a special case, suppose that we take $\lambda = n^{-\frac{1}{2(1+\alpha)}}$. Then we find $s_0 \sim n^{-\frac{1}{2(1+\alpha)}}$ as in the previous section. Then the bound (13) yields $I^2(f^0) = \mathcal{O}(1)$. Otherwise, we see that $\tau(\hat{f})$ concentrates on the boundary τ_{\min} , and that in this example, we have

$$\tau_{\min}^{2} = \min_{g \in \mathcal{G}} \bigg\{ \|g - g^{0}\|^{2} + \lambda^{2} I^{2}(g) \bigg\}.$$

6.1.3 Case 3: Strictly convex penalty

In this case, we assume: $\circ 1 < q \leq 2 \text{ not depending on } n,$ $\circ \text{ Condition 4.1 with } K = \mathcal{O}(1),$ $\circ \mathbf{E}(s) \leq \mathcal{J}(s)/m_n,$ $\circ \mathcal{J}(s) = As^{1+(2/q-1)\alpha}, m_n = \sqrt{n}\lambda^{2\alpha/q}, \exists A = \mathcal{O}(1), \exists 0 < \alpha < 1 \text{ not depending on } n,$ $\circ s_0 \asymp (\sqrt{n}\lambda^{2\alpha/q})^{-\frac{q}{q-(2-q)\alpha}},$ $\circ \log n/(n^{2-q}\lambda^4)^{\frac{\alpha}{q-(2-q)\alpha}} = o(1).$

Under this condition, Theorem 4.1 combined with Lemma 5.1 implies the deviation result

$$\frac{\tau(\hat{f}) - s_0}{s_0} \le z_n, \text{ where } z_n = \mathcal{O}_{\mathbf{P}}\left(\frac{\log n}{ns_0^2}\right) + o_{\mathbf{P}}(1) = o_{\mathbf{P}}(1).$$

6.2 Linearized least squares regression

Let $\{(X_i, Y_i)\}_{i=1}^n$ be i.i.d. samples taking values in $\mathbb{R}^p \times \mathbb{R}$. We assume the model

$$Y_i = g^0(X_i) + \epsilon_i, \quad \text{for } i = 1, \dots, n,$$

where $\epsilon_i \sim \mathcal{N}(0,1)$ is independent of X_i , and the function g^0 belongs to a convex model class \mathcal{G} . Assume \mathcal{G} is a convex subset of a linear vector space $\overline{\mathcal{G}}$, and moreover that

$$K_X := \max_{g \in \mathcal{G}} \|g - g^0\|_{\infty} < \infty.$$

We moreover assume $K_0 := ||g^0||_{\infty} < \infty$. Let *I* be some pseudo-norm on $\overline{\mathcal{G}}$, we consider the estimator

$$\hat{g} := \arg\min_{g \in \mathcal{G}} \left\{ -\frac{1}{n} \sum_{i=1}^{n} Y_i g(X_i) + \frac{Pg^2}{2} + \lambda^2 I^2(g) \right\}.$$
(14)

Note that implementation of this estimator requires that Pg^2 is known or can be computed for all $g \in \mathcal{G}$.

Given the form of the estimator (14), we have

$$\mathcal{F} = \left\{ f_g(x, y) = -yg(x) + Pg^2/2 \mid g \in \mathcal{G} \right\}$$

This class has an envelope function F satisfying the conditions of Lemma 8.1. To see this, note the bounds

$$\max_{\substack{g \in \mathcal{G} \\ g \in \mathcal{G}}} |\epsilon_i(g(X_i) - g^0(X_i))| \le |\epsilon_i| K_X,$$

$$\max_{g \in \mathcal{G}} |g^0(X_i)(g(X_i) - g^0(X_i))| \le K_X K_0, \quad \text{and}$$

$$\max_{g \in \mathcal{G}} |Y_i(g(X_i) - g^0(X_0))| \le (|\epsilon_i| + K_0) K_X.$$

Lemma 8.1 can now be used as concentration tool. Still, as the class \mathcal{F} is not uniformly bounded in this case one cannot apply Lemma 8.2. The strategy may then be to first prove that $\tau(\hat{f}) = \mathcal{O}_{\mathbf{P}}(s_0)$ and then that $\mathbf{E}(s) \simeq s_0$ for $s \simeq s_0$. The results are then as in the previous subsection albeit that we are facing an additional log *n* factor.

Linearized least-squares regression refers to the special case of a linear model. More concretely, define the design matrix $X \in \mathbb{R}^{n \times p}$ rows X_i^T for $i = 1, \ldots, n$, as well as the sample covariance matrix $\hat{\Sigma} := X^T X/n$. The linear model consists of the function class

$$\mathcal{G} = \{ g_{\beta}(x) = x^T \beta : \beta \in \mathcal{B} \},\$$

along with the true function $g^0(x) = x^T \beta^0$. Our conditions then require that the population covariance matrix $\Sigma_0 := \mathbb{E}\hat{\Sigma}$ is known, and moreover that \mathcal{B} is a convex subset of \mathbb{R}^p and satisfying, for some constants K_0 and K_X , $|X_1^T\beta^0| \leq K_0$ and $|X_1^T(\beta - \beta^0)| \leq K_X$ for all $\beta \in \mathcal{B}$. The latter is for example true when $||X_1|| \leq K_X$ and $||\beta - \beta^0||_1 \leq 1$ (say) for all $\beta \in \mathcal{B}$.

7 Exponential families with squared norm penalty

We now turn to some examples involving exponential families. Throughout this section, we specialize to the case of squared norm penalties, noting that more general penalties can be studied as in the previous section.

7.1 Density estimation

Suppose that X_1, \ldots, X_n are i.i.d. random variables with distribution P taking values in \mathcal{X} . Given a sigma-finite measure ν on \mathcal{X} , let us define the function class

$$\bar{\mathcal{G}} := \bigg\{g: \ \mathcal{X} \to \mathbb{R}: \ \int \exp[g(x)] d\nu(x) < \infty \bigg\},$$

and note that it is convex. We define a functional on $\bar{\mathcal{G}}$ via

$$d(g) := \log\left(\int \exp[g(x)]d\nu(x)\right).$$

Let $\mathcal{G} \subset \overline{\mathcal{G}}$ be a convex subset and define the function $f_g = -g + d(g)$, for each $g \in \mathcal{G}$, along with the associated function class $\mathcal{F} := \{f_g \mid g \in \mathcal{G}\}$. Letting I be a pseudo-norm on \overline{G} along with with a non-negative regularization weight λ , we consider the estimator

$$\hat{g} := \arg\min_{g \in \mathcal{G}} \bigg\{ -P_n g + \mathbf{d}(g) + \lambda^2 I^2(g) \bigg\},\$$

and define $\hat{f} := f_{\hat{g}}$.

For identification purposes, we take the functions in \mathcal{G} to be centered— that is, such that $\int g d\nu = 0$ for each $g \in \mathcal{G}$. For simplicity and without loss of generality, we take $g^0 \equiv 0$ so that $f^0 \equiv 1$ and $\nu = P$. Since P is unknown, the centering of the functions (in actual practice) is done with respect to some other measure. This difference does not alter the theory but should be kept in mind when examining the assumptions.

The following lemma relates the function d to the second moment:

Lemma 7.1 Suppose that $K := \max_{g \in \mathcal{G}} \|g\|_{\infty} < \infty$. Then we have

$$\max_{g \in \mathcal{G}} \left| \frac{\mathrm{d}(tg)}{t^2 P g^2} - \frac{1}{2} \right| = \mathcal{O}(t), \qquad \text{valid as } t \downarrow 0.$$

As a useful corollary, it gives us an asymptotic expression for d(g) as the ℓ_{∞} -norm of g shrinks. In particular, we let define $\mathcal{G}_{\infty}(\eta) := \{g : \|g\|_{\infty} \leq \eta\}.$

Corollary 7.1 For each $g \in \mathcal{G}_{\infty}(\eta)$, we have

$$d(g) = \frac{1}{2} P g^2 (1 + \mathcal{O}(\eta)), \qquad valid \ as \ \eta \downarrow 0.$$

We are now equipped to state a result. Suppose that:

 $\begin{aligned} &\circ \ r_0 = \mathcal{O}(s_0), \\ &\circ \ \tau(\hat{f}) = \mathcal{O}_{\mathbf{P}}(s_0), \\ &\circ \ \|\hat{g}\|_{\infty} = o_{\mathbf{P}}(1), \\ &\circ \ \sqrt{\log n/n} = o_{\mathbf{P}}(s_0). \end{aligned}$

Combining Theorem 4.1 with Lemmas 5.2 and 5.3, along with Corollary 7.1, guarantees that $\tau(\hat{f})$ concentrates on s_0 , and in particular, we have

$$|\tau(f) - s_0| = o_{\mathbf{P}}(s_0).$$

7.2 Regression with fixed design

Let $\{(X_i, Y_i)\}_{i=1}^n$ be independent observations taking values in the Cartesian product space $\mathcal{X} \times \mathbb{R}$. We assume the design $\{X_i\}_{i=1}^n$ is fixed, and that for some given sigma-finite measure ν , the log-density of Y_i given X_i takes the log-linear form

$$-Y_i g^0(X_i) + \mathrm{d}(g^0(X_i)),$$

where the function $d(\xi) := \log \left(\int \exp[y\xi] d\nu \right)$ has domain

$$\Xi := \{\xi \in \mathbb{R} \mid \int \exp[y\xi] d\nu < \infty\}.$$

We define $g := (g(X_1), \ldots, g(X_n))^T \in \mathbb{R}^n$, let \mathcal{G} be a convex subset of \mathbb{R}^n and use, for $v \in \mathbb{R}^n$, the notation $||v||_n^2 := v^T v/n$. Letting I be a pseudo-norm on \mathbb{R}^n , we consider the estimator

$$\hat{g} := \arg \max_{g \in \mathcal{G}} \left\{ -Y^T g/n + \sum_{i=1}^n \mathrm{d}(g(X_i))/n + \lambda^2 I^2(g) \right\}.$$

In this case, the effective function class takes the form

$$\mathcal{F} = \{ f_g(x, y) = -yg(x) + d(g(x)) : g \in \mathcal{G} \},\$$

and we have $\hat{f} = f_{\hat{g}}$.

Let us assume that: $\circ \sup_{\tau^{2}(f_{g}) \leq \eta} \sum_{i=1}^{n} d(g(X_{i})) - d(g^{0}(X_{i}))) / (n ||g - g^{0}||_{n}^{2}) = c^{2}(1 + o(1)), \ \eta \downarrow 0,$ $\circ ||\hat{g} - g^{0}||_{n} = o_{\mathbf{P}}(1),$ $\circ r_{0} = \mathcal{O}(s_{0}),$ $\circ \tau(\hat{f}) = \mathcal{O}_{\mathbf{P}}(s_{0}),$ $\circ \sqrt{\log n/n} = o_{\mathbf{P}}(s_{0}).$

Then Theorem 4.1 in conjunction with Lemmas 5.2 and 5.3 guarantees that $\tau(\hat{f})$ concentrates on s_0 , and moreover that

$$|\tau(\hat{f}) - s_0| = o_{\mathbf{P}}(s_0).$$

8 Concentration for maxima of empirical processes

The following result of Klein [2002] (see also Klein and Rio [2005]) is our main tool.

Theorem 8.1 Define $K := \max_{f \in \mathcal{F}_s} ||f - f^0||_{\infty}$ and $\sigma_s := \max_{f \in \mathcal{F}_s} \sigma(f - f^0)$. Then for all $t \ge 0$, we have

$$\hat{\mathbf{E}}_n(s) \ge \mathbf{E}(s) - \sqrt{8K\mathbf{E}(s) + 2\sigma_s^2}\sqrt{t/n} - \frac{Kt}{n}, \quad and$$
$$\hat{\mathbf{E}}_n(s) \le \mathbf{E}(s) + \sqrt{8K\mathbf{E}(s) + 2\sigma_s^2}\sqrt{t/n} + \frac{2Kt}{3n},$$

where each bound holds with probability at least $1 - \exp[-t]$,

We next present a consequence for the case where the functions in \mathcal{F}_s are not uniformly bounded, but have a (sub-Gaussian) envelope function. This result is invoked in the analysis of Section 6.2.

Lemma 8.1 Assume that for some constants $c_F \ge 1$ and $C_F \ge 1$, the envelope function $F(\cdot) := \max_{f \in \mathcal{F}_s} |f(\cdot) - f^0(\cdot)|$ satisfies the bounds

$$PF^{2}{}^{1}{F > t} \le c_{F}^{2} \exp[-t^{2}/C_{F}^{2}], \ t > 0.$$
(15)

Then for all t > 0 with probability at least $1 - \exp[-t] - 1/t^2$

$$\hat{\mathbf{E}}_n(s) \ge \mathbf{E}(s) - \sqrt{8C_F \sqrt{\log n}} [\mathbf{E}(s) + 2c_F (c_F + t)/n] + 2\sigma_s^2 \sqrt{t/n} - C_F t \sqrt{\log n}/n - c_F (4c_F + t)/n,$$

and with probability at least $1 - \exp[-t] - 1/t^2$

$$\hat{\mathbf{E}}_{n}(s) \leq \mathbf{E}(s) - \sqrt{8C_{F}\sqrt{\log n}}[\mathbf{E}(s) + 2c_{F}^{2}/n] + 2\sigma_{s}^{2}\sqrt{t/n} - 2C_{F}t\sqrt{\log n}/(3n) - c_{F}(4c_{F}+t)/n.$$

In the next lemma, we replace the quantity $\mathbf{E}(s)$ appearing in the square-root of Theorem 8.1 by a suitable upper bound.

Lemma 8.2 Under Conditions 4.1 and 4.2, we have

$$\hat{\mathbf{E}}_n(s) \ge \mathbf{E}(s) - 2Cs\sqrt{t/n} + r_0\sqrt{t/n} - Kt/n, \quad and \\ \hat{\mathbf{E}}_n(s) \le \mathbf{E}(s) + 2Cs\sqrt{t/n} + r_0\sqrt{t/n} + 2Kt/(3n),$$

where each bound holds with probability at least $1 - \exp[-t]$.

9 The shifted version

For a scalar $\tau_*^2 \geq \tau_{\min}^2$ to be chosen, we study in this section the "shifted" function

$$\mathbf{F}(s) := \max_{\tau^2(f) \le \tau_*^2 + s^2} (P_n - P)(f^0 - f), \quad \text{defined for } s^2 \ge \tau_*^2 - \tau_{\min}^2.$$

This shifted version may be of interest when $\tau^2(\hat{f})$ is of larger order than $P(\hat{f} - f^0)$. The idea is then to replace g^0 in the previous sections by the function $g^* := \arg \min_{g \in \mathcal{G}} \tau^2(g)$. One then needs curvature conditions on $R(g) - R(g^*)$ instead of $R(g) - R(g^0)$. This we handle here by the notion of an "oracle potential", as defined in Definition 9.1 below.

Lemma 9.1 shows that curvature conditions on the function $\mathcal{Q}(s; \mathbf{E}) := s^2 - \mathbf{E}(s)$ are weaker than those on the function $\mathcal{Q}(s; \mathbf{F}) := s^2 - \mathbf{F}(s)$. Using the shorthand $s_*^2 = s_0^2 - \tau_*^2$, the following lemma summarizes this fact:

Lemma 9.1 For any $s \ge \tau_{\min}$ and $\tilde{s}^2 = s^2 - \tau_*^2$, we have

$$\mathcal{Q}(s; \mathbf{E}) - \mathcal{Q}(s_0; \mathbf{E}) = \mathcal{Q}(\tilde{s}; \mathbf{F}) - \mathcal{Q}(s_*; \mathbf{F})$$

and $|\tilde{s} - s_*| \ge |s - s_0|$.

We define for $s^2 \ge \tau_*^2 - \tau_{\min}^2$

$$\kappa_s^2 := \max\left\{ P(f - f^0) : f \in \mathcal{F}, \ P(f - f^0) + \operatorname{pen}(f) \le \tau_*^2 + s^2 \right\}.$$

Definition 9.1 We say that the oracle potential holds if

$$\Gamma := \sup_{s>0} \kappa_s / s < \infty. \tag{16}$$

For the shifted version, the counterpart of Condition 4.2 replaces $\mathbf{E}(\cdot)$ by $\mathbf{F}(\cdot)$.

Condition 9.1 There is a constant m_n , a strictly increasing function \mathcal{J} such that the function $\Phi_{\mathcal{J}}$ is strictly convex, and such that the bound

$$\mathbf{F}(s) \le \frac{\mathcal{J}(s)}{m_n} \tag{17}$$

holds for all $s \geq 0$.

When Conditions 4.1 and 9.1 hold, we define

$$r_*^2 := 2C^2 \Phi_{\mathcal{J}}^* (4K/(m_n C^2))$$

where $\Phi_{\mathcal{J}}^*$ is the convex conjugate of $\Phi_{\mathcal{J}}$.

Theorem 9.1 Suppose that:

- Conditions 4.1 and 9.1, as well as the quadratic curvature condition with constant C hold.
- The shifted mapping Q(·; F) satisfies the right-sided second order margin condition over the interval (s_{*} + δ, √τ²_{max} τ²_{*}] with margin function G.
- The oracle potential condition (16) holds.

Then there is a constant c_0 depending on C, K and Γ , such that for all t > 0and for a constant $\delta(t)$ such that $G(\delta(t))$ is not larger than

$$G^* \left(c_0 \sqrt{[t + \log(1 + \sqrt{n(\tau_{\max}^2 - \tau_*^2)})]/n}) \right) + c_0 \left((s_* + r_*) \sqrt{[t + \log(1 + \sqrt{n(\tau_{\max}^2 - \tau_*^2)})]/n} + [t + \log(1 + \sqrt{n(\tau_{\max}^2 - \tau_*^2)})]/n) \right)$$

one has the deviation inequality

$$\mathbb{P}\left(\sqrt{\tau_{\max}^2 - \tau_*^2} \ge \sqrt{\tau^2(\hat{f}) - \tau_*^2} > s_* + \max\{\delta(t), \underline{\delta}\}\right) \le \exp[-t].$$
(18)

Moreover, if $s_*^2 > \tau_{\min}^2 - \tau_*^2$ and in fact the two-sided second order margin condition holds for $\mathcal{Q}(\cdot; \mathbf{F})$ in the range $[\sqrt{\tau_{\min}^2 - \tau_*^2}, s_*) \cup (s_* + \underline{\delta}, \sqrt{\tau_{\max}^2 - \tau_*^2}]$ with margin function G, then one has the concentration inequality

$$\mathbb{P}\left(|\sqrt{\tau^2(\hat{f}) - \tau_*^2} - s_*| > \max\{\delta(t), \underline{\delta}\}, \ \tau(\hat{f}) \le \tau_{\max}\right) \le 2\exp[-t].$$

10 Proofs

This section is devoted to the proofs of all our results.

10.1 Proof of Theorem 2.1

After some simple algebra, we may write

$$\hat{g} = \arg\min_{g \in \mathbb{R}^n} \left\{ \frac{1}{2} \tau^2(g) - \epsilon^T g/n \right\},$$

where $\tau^2(g) := \|g - g^0\|_n^2 + \text{pen}(g)$. The function τ^2 is convex, so that it has a sub-differential, denoted by $\partial \tau^2(g)$. With this notation, the minimizing argument \hat{g} must satisfy the relation

$$\epsilon/n \in \partial \tau^2(\hat{g})/2. \tag{19}$$

Due to the strong convexity and coercivity of $g \mapsto \tau^2(g)$, the inclusion (19) always has a unique solution \hat{g} .

We now use a classical fact from convex analysis (Rockafellar [1970]): since the function $g \mapsto \tau^2(g)/2$ is (1/n)-strongly convex, the sub-differential mapping $g \mapsto \partial \tau^2(g)/2$ is (1/n)-strongly monotone, which means that for any pair of vectors $u, u' \in \mathbb{R}^n$

$$||u - u'||_n^2 \le (v - v')^T (u - u'), \tag{20}$$

where v, v' denote any members of $\partial \tau^2(u)/2$ and $\partial \tau^2(u')/2$ respectively.

By Borell's theorem (1975) on the concentration of Lipschitz functions of Gaussian vectors, it suffices to show that the mapping $\epsilon \mapsto \hat{m} := \|\hat{g} - g^0\|_n$ is Lipschitz with parameter $1/\sqrt{n}$. Let ϵ and ϵ' be two realizations of the noise vector, with corresponding solutions \hat{g} and \hat{g}' , along with their associated errors $\hat{m} = \|\hat{g} - g^0\|_n$ and $\hat{m}' = \|\hat{g}' - g^0\|_n$. By the triangle inequality, we have

$$|\hat{m} - \hat{m}'| = \left| \|\hat{g} - g^0\|_n - \|\hat{g}' - g^0\|_n \right| \le \|\hat{g} - \hat{g}'\|_n,$$

so that it suffices to prove that

$$\|\hat{g} - \hat{g}'\|_n \le \|\epsilon - \epsilon'\|_n = \frac{1}{\sqrt{n}} \|\epsilon - \epsilon'\|_2.$$

Now consider the pair $u = \hat{g}$ and $u' = \hat{g}'$, along with the corresponding elements $v = \epsilon/n$ and $v' = \epsilon'/n$. Applying the monotone property (20) to these pairs yields the inequality

$$\|\hat{g} - \hat{g}'\|_n^2 \le (\epsilon - \epsilon')^T (\hat{g} - \hat{g}')/n \le \|\epsilon - \epsilon'\|_n \|\hat{g} - \hat{g}'\|_n,$$

where the final step follows from the Cauchy-Schwarz inequality. Cancelling terms completes the proof. $\hfill \Box$

10.2 Proofs for Section 3

In this section, we collect the proofs of all results stated in Section 3.

Proof of Lemma 3.1: For any scalar s and $f \in \mathcal{F}$ such that $\tau(f) \leq s$, we have

$$P_n(f - f^0) + pen(f) = \tau^2(f) - (P_n - P)(f^0 - f) \le s^2 - (P_n - P)(f^0 - f).$$

Consequently, we have

$$\begin{aligned} \tau^{2}(\hat{f}) - (P_{n} - P)(f^{0} - \hat{f}) &= \min_{f \in \mathcal{F}} \left\{ \tau^{2}(f) - (P_{n} - P)(f^{0} - f) \right\} \\ &= \min_{s \ge \tau_{\min}} \min_{\tau(f) \le s} \left\{ \tau^{2}(f) - (P_{n} - P)(f^{0} - f) \right\} \\ &\leq \min_{s \ge \tau_{\min}} \min_{\tau(f) \le s} \left\{ s^{2} - (P_{n} - P)(f^{0} - f) \right\} \\ &= \min_{s \ge \tau_{\min}} \left\{ s^{2} - \max_{\tau(f) \le s} (P_{n} - P)(f^{0} - f) \right\} \\ &= \min_{s \ge \tau_{\min}} \left\{ s^{2} - \hat{\mathbf{E}}_{n}(s) \right\}. \end{aligned}$$

On the other hand, for any $f \in \mathcal{F}$, we have the lower bound

$$\tau^{2}(f) - (P_{n} - P)(f^{0} - f) \ge \tau^{2}(f) - \max_{\tilde{f} \in \mathcal{F}_{\tau(f)}} (P_{n} - P)(f^{0} - \tilde{f}) = \tau^{2}(f) - \hat{\mathbf{E}}_{n}(\tau(f)),$$

which implies that

$$\tau^{2}(\hat{f}) - (P_{n} - P)(f^{0} - \hat{f}) \ge \tau^{2}(\hat{f}) - \hat{\mathbf{E}}_{n}(\tau(\hat{f})) \ge \min_{s \ge \tau_{\min}} \{s^{2} - \hat{\mathbf{E}}_{n}(s)\}.$$

Since the minimizing argument $\hat{s} = \arg \min_{s \ge \tau_{\min}} \{s^2 - \hat{\mathbf{E}}_n(s)\}$ is unique by assumption, we conclude that $\tau(\hat{f}) = \hat{s}$, as claimed. \Box

10.3 Proofs for Section 4

We first state and prove an auxiliary lemma that serves as a tool in the proof of Theorem 4.1.

Lemma 10.1 Let G be a real-valued function with convex conjugate G^* . Then all for positive scalars a, b and c such that $G(a) \ge G^*(2b) + 2c$, we have $G(a) - ab - c \ge 0$.

Proof of Lemma 10.1. By the Fenchel-Young inequality, we have

$$ab = a\frac{2b}{2} \le \frac{G(a)}{2} + \frac{G^*(2b)}{2},$$

and consequently,

$$G(a) - ab - c \ge G(a) - \frac{G(a)}{2} - \frac{G^*(2b)}{2} - c = \frac{G(a)}{2} - \frac{G^*(2b)}{2} - c \ge 0.$$

We now turn the proof of the main theorem.

Proof of Theorem 4.1. Assume first that the right-sided second order margin condition holds with $\underline{\delta} = 0$. Let t > 0 be arbitrary, and define

$$\bar{z}(t) := 2Cs_0\sqrt{\frac{t}{n}} + r_0\sqrt{\frac{t}{n}} + \frac{2Kt}{3n}, \text{ and } \underline{z}(t) := 2Cs_0\sqrt{\frac{t}{n}} + r_0\sqrt{\frac{t}{n}} + \frac{Kt}{n}.$$

Our strategy is to apply a "peeling argument" so as to transition from the fixed s-result of Theorem 8.1 to a result that holds uniformly in s. For a parameter $\epsilon > 0$ to be chosen later, define the intervals

$$I_j := ((j-1)\epsilon + \delta + s_0, s_0 + \delta + j\epsilon], \quad \text{for } j = 1, \dots, J,$$

where $J := \lceil \frac{\tau_{\max}}{\epsilon} \rceil$, as well as the associated probabilities

$$\mathbb{P}_j = \mathbb{P}\left(\exists \ s \in I_j \text{ such that } s^2 - \hat{\mathbf{E}}_n(s) \le s_0^2 - \mathbf{E}(s_0) + \bar{z}(t)\right)$$

Then for a parameter $\delta > 0$ to be chosen later (and leading to the $\delta(t)$ in the theorem statement), we have

$$\mathbb{P}\left(s_0 + \delta < \hat{s} \le \tau_{\max}, \ \hat{\mathbf{E}}_n(s_0) > \mathbf{E}(s_0) + \bar{z}(t)\right)$$

$$\leq \mathbb{P}\left(\exists s \in (s_0 + \delta, \tau_{\max}] \text{ such that } s^2 - \hat{\mathbf{E}}_n(s) \le s_0^2 - \mathbf{E}(s_0) + \bar{z}(t)\right) \le \sum_{j=1}^J \mathbb{P}_j.$$

For each index j and for all $s \in I_j$, we have

$$s^{2} - \hat{\mathbf{E}}_{n}(s) \ge ((j-1)\epsilon + \delta + s_{0})^{2} - \hat{\mathbf{E}}_{n}(s_{0} + \delta + j\epsilon).$$

Moreover, for all u > 0, we have by Theorem 8.1

$$\hat{\mathbf{E}}_n(s_0+\delta+j\epsilon) \le \mathbf{E}(s_0+\delta+j\epsilon) - 2C(\delta+j\epsilon)\sqrt{u/n} - \underline{z}(u)$$

with probability at least $1 - \exp[-u]$. Furthermore, by the one-sided form of second order margin condition, we have the lower bound

$$(s_0 + \delta + (j-1)\epsilon)^2 - \mathbf{E}(s_0 + \delta + j\epsilon) \ge s_0^2 - \mathbf{E}(s_0) + G(\delta + j\epsilon) + (s_0 + \delta(j-1)\epsilon)^2 - (s_0 + \delta + j\epsilon)^2 = s_0^2 - \mathbf{E}(s_0) + G(\delta + j\epsilon) - 2\epsilon(s_0 + \delta + j\epsilon) + \epsilon^2.$$

Putting together the pieces, for all $s \in I_j$, we have

$$s^2 - \hat{\mathbf{E}}_n(s) \ge s_0^2 - \mathbf{E}(s_0) + G(\delta + j\epsilon) - 2\left(C\sqrt{\frac{u}{n}} + \epsilon\right)(\delta + j\epsilon) - 2\epsilon s_0 + \epsilon^2 - \underline{z}(u),$$

with probability at least $1 - \exp[-u]$. We now apply Lemma 10.1 with the choices $a := \delta + j\epsilon$, $b := 2(C\sqrt{u/n} + \epsilon)$, and $c := 2\epsilon s_0 - \epsilon^2 + \underline{z}(u) + \overline{z}(t)$. In order to be able to do so, we require that

$$G(\delta) \ge G^* \left(4(C\sqrt{u/n} + \epsilon) \right) + 2\left(2\epsilon s_0 - \epsilon^2 + \underline{z}(u) + \overline{z}(t) \right).$$
(21)

We now settle the choice of ϵ and u. Taking $\epsilon = 1/\sqrt{n}$, we are then guaranteed that

$$J \le 1 + \frac{\tau_{\max}}{\epsilon} = 1 + \sqrt{n\tau_{\max}^2}.$$

Moreover, recalling the arbitrary t > 0 introduced at the start of the proof, we set $u = t + \log(1 + \sqrt{n\tau_{\max}^2})$, and then the condition (21) on $\delta := \delta(t)$ becomes

$$G(\delta(t)) := G^* \left(4(C\sqrt{[t + \log(1 + \sqrt{n\tau_{\max}^2})]/n} + 1/\sqrt{n}) \right) + 2\left(2s_0/\sqrt{n} - 1/n + \underline{z}([t + \log(1 + \sqrt{n\tau_{\max}^2})]) + \overline{z}(t) \right).$$

We are then guaranteed that for each $j \in \{1, \ldots, J\}$, with probability at least $1 - \exp[-(t + \log(1 + \sqrt{n\tau_{\max}^2})])$, for all $s \in I_j$ it holds that

$$s^2 - \hat{\mathbf{E}}_n(s) \ge s_0^2 - \mathbf{E}(s_0) + \bar{z}(t).$$

It follows that $\mathbb{P}_j \leq \exp\left\{-\left[t + \log(1 + \sqrt{n\tau_{\max}^2})\right]\right\}$, and hence

$$\sum_{j=1}^{J} \mathbb{P}_j \le J \exp\left[-[t + \log(1 + \sqrt{n\tau_{\max}^2})]\right] \le \exp[-t].$$

One easily verifies that for some constant c_0 depending on C and K, we have

$$G(\delta(t)) \le G^* \left(c_0 \sqrt{[t + \log(1 + \sqrt{n\tau_{\max}^2})]/n}) \right) + c_0 \left((s_0 + r_0) \sqrt{[t + \log(1 + \sqrt{n\tau_{\max}^2})]/n} + [t + \log(1 + \sqrt{n\tau_{\max}^2})]/n} \right).$$

Overall, we have established that $\mathbb{P}(\hat{s} > s_0 + \delta(t)) \leq 2 \exp[-t]$.

In stating the bound in the theorem, we removed the pre-factor of 2 for cosmetic reasons. This can be done by replacing t by $t + \log 2$. In order to prove the lower bound, one may follow the same argument, instead using the left-sided version of the second order margin condition.

In our argument thus far, we assumed $\underline{\delta} = 0$. If the second order margin condition only holds at distance $\underline{\delta} > 0$, it is clear that one simply can take the maximum of $\delta(t)$ and $\underline{\delta}$ in the bounds.

10.4 Proofs for Section 5

Proof of Lemma 5.1. Let us introduce the shorthand notation $\tilde{s} := s^{2/q}$ and $\tilde{s}_0 = s_0^{2/q}$, let $\tilde{s}_1 \ge \tau_{\min}^{2/q}$ and $\tilde{s}_2 \ge \tau_{\min}^{2/q}$ be arbitrary, and define

$$\hat{f}_1 := f_{\hat{g}_1} := \arg \max_{\tau^{2/q}(f) \le \tilde{s}_1} (P_n - P)(f^0 - f), \text{ as well as}$$
$$\hat{f}_2 := f_{\hat{g}_2} := \arg \max_{\tau^{2/q}(f) \le \tilde{s}_2} (P_n - P)(f^0 - f).$$

With these choices, we have

$$\tau^{2/q}(f_{t\hat{g}_1+(1-t)\hat{g}_2}) \le t\tau^{2/q}(\hat{f}_1) + (1-t)\tau^{2/q}(\hat{f}_2) \le t\tilde{s}_1 + (1-t)\tilde{s}_2 \quad \text{for all } t \in [0,1]$$

Moreover, we have the lower bound

$$\hat{\mathbf{E}}_{n}((t\tilde{s}_{1}+(1-t)\tilde{s}_{2})^{q/2}) \geq (P_{n}-P)(f^{0}-f_{t\hat{g}_{1}+(1-t)\hat{g}_{2}})$$

= $t(P_{n}-P)(f^{0}-\hat{f}_{1}) + (1-t)(P_{n}-P)(f^{0}-\hat{f}_{2})$
= $t\hat{\mathbf{E}}_{n}(\tilde{s}_{1}^{q/2}) + (1-t)\hat{\mathbf{E}}_{n}(\tilde{s}_{2}^{q/2}).$

Taking expectations yields the lower bound

$$\mathbf{E}((t\tilde{s}_1 + (1-t)\tilde{s}_2)^{q/2}) \ge t\mathbf{E}(\tilde{s}_1^{q/2}) + (1-t)\mathbf{E}(\tilde{s}_2^{q/2})$$

Using the fact that

$$[(t\tilde{s}_1 + (1-t)\tilde{s}_2)^q - \mathbf{E}((t\tilde{s}_1 + (1-t)\tilde{s}_2)^{q/2})] - [\tilde{s}_0^q - \mathbf{E}(\tilde{s}_0^{q/2})] \ge 0,$$

we have

$$[\tilde{s}^{q} - \mathbf{E}(\tilde{s}^{q/2})] - [\tilde{s}_{0}^{q} - \mathbf{E}(\tilde{s}_{0}^{q/2})] \ge \tilde{s}^{q} - \tilde{s}_{0}^{q} + \frac{1}{t}\tilde{s}_{0}^{q} - (t\tilde{s} + (1-t)\tilde{s}_{0})^{q}.$$

Taking $t \downarrow 0$ then gives

$$[\tilde{s}^{q} - \mathbf{E}(\tilde{s}^{q/2})] - [\tilde{s}_{0}^{q} - \mathbf{E}(\tilde{s}_{0}^{q/2})] \ge \tilde{s}^{q} - \tilde{s}_{0}^{q} - q\tilde{s}_{0}^{q-1}(\tilde{s} - \tilde{s}_{0}) \ge q(q-1)(M+1)^{-\frac{2(2-q)}{q}} \tilde{s}_{0}^{-(2-q)}(\tilde{s} - \tilde{s}_{0})^{2}/2,$$

valid when $q \in (1,2]$ and $\tilde{s} > (M+1)^{2/q} \tilde{s}_0$. Furthermore, for $1 < q \le 2$ we get for some $s_0 < \bar{s} \le s$

$$\tilde{s} - \tilde{s}_0 = s^{2/q} - s_0^{2/q} = 2\bar{s}^{\frac{2-q}{q}}(s-s_0)/q \ge 2s_0^{\frac{2-q}{q}}(s-s_0)/q.$$

Consequently, for all $q \in (1, 2]$, we have

$$[\tilde{s}^{q} - \mathbf{E}(\tilde{s}^{q/2})] - [\tilde{s}_{0}^{q} - \mathbf{E}(\tilde{s}_{0}^{q/2})] \ge 2(q-1)(M+1)^{-\frac{2(2-q)}{q}}(s-s_{0})^{2}/q,$$

as claimed.

Proof of Lemma 5.2. For all $s \ge \tau_{\min}$, we have

$$\mathbf{E}^{\tau}(As) \ge \mathbf{E}^{\varsigma}(A^{1/2}s) \ge \mathbf{E}^{\tau}(s).$$

For $ts + (1-t)s_0 \ge \tau_{\min}$, the concavity of \mathbf{E}^{ς} yields

$$\mathbf{E}^{\tau} (A(ts + (1-t)s_0)) \geq \mathbf{E}^{\varsigma} (A^{1/2}(ts + (1-t)s_0))$$

$$\geq t \mathbf{E}^{\varsigma} (A^{1/2}s) + (1-t) \mathbf{E}^{\varsigma} (A^{1/2}s_0)$$

$$\geq t \mathbf{E}^{\tau}(s) + (1-t) \mathbf{E}^{\tau}(s_0)$$

$$= t(\mathbf{E}(s) - \mathbf{E}(s_0)) + \mathbf{E}(s_0).$$

Since s_0 is the minimizer of $s^2 - \mathbf{E}(s)$, we have

$$A^{2}(ts + (1-t)s_{0})^{2} - \mathbf{E}(A^{2}(ts + (1-t)s_{0})) \ge s_{0}^{2} - \mathbf{E}(s_{0}),$$

and consequently

$$\mathbf{E}(A(ts+(1-t)s_0)) \le A^2(ts+(1-t)s_0)^2 - s_0^2 + \mathbf{E}(s_0).$$

It follows that

$$t(\mathbf{E}(s) - \mathbf{E}(s_0)) \le A^2 (ts + (1-t)s_0)^2 - s_0^2$$

= $A^2 (t^2 s^2 + s_0^2 - 2ts_0^2 + t^2 s_0^2 + 2tss_0 - 2t^2 ss_0) - s_0^2$
= $A^2 (t^2 (s - s_0)^2 + s_0^2 - 2ts_0^2 + 2tss_0) - s_0^2$
= $t^2 (s - s_0)^2 - 2ts_0^2 + 2tss_0 + \epsilon \left(t^2 (s - s_0)^2 - 2ts_0^2 + 2tss_0\right) + \epsilon s_0^2.$

Putting together the pieces, we have

$$[s^{2} - \mathbf{E}(s)] - [s_{0}^{2} - \mathbf{E}(s_{0})] = s^{2} - s_{0}^{2} - \frac{t(\mathbf{E}(s) - \mathbf{E}(s_{0}))}{t}$$

$$\geq s^{2} - s_{0}^{2} - t(s - s_{0})^{2} + 2s_{0}^{2} - 2ss_{0} - \epsilon \left(t(s - s_{0})^{2} - 2s_{0}^{2} + 2ss_{0}\right) - \epsilon s_{0}^{2}/t$$

$$= (s - s_{0})^{2} - t(s - s_{0})^{2} - \epsilon \left(t(s - s_{0})^{2} + 2s_{0}(s - s_{0})\right) - \epsilon s_{0}^{2}/t.$$

We now choose $t = \sqrt{\epsilon}$ to find that

$$[s^{2} - \mathbf{E}(s)] - [s_{0}^{2} - \mathbf{E}(s_{0})] \ge (s - s_{0})^{2} - \sqrt{\epsilon}(s - s_{0})^{2} - \epsilon \left(\sqrt{\epsilon}(s - s_{0})^{2} + 2s_{0}(s - s_{0})\right) - \sqrt{\epsilon}s_{0}^{2}.$$

Next we use our assumption that $s \leq \tau_{\max} = (M+1)s_0$. We then get

$$[s^{2} - \mathbf{E}(s)] - [s_{0}^{2} - \mathbf{E}(s_{0})] \ge \left(1 - \sqrt{\epsilon}(1+\epsilon)\right)(s-s_{0})^{2} - \epsilon\left(2s_{0}^{2}M\right) - \sqrt{\epsilon}s_{0}^{2}$$
$$= (1 - \sqrt{\epsilon}(1+\epsilon))(s-s_{0})^{2} - \sqrt{\epsilon}(2\sqrt{\epsilon}M+1)s_{0}^{2}$$
$$\ge (s-s_{0})^{2}/2 - \sqrt{\epsilon}(2\sqrt{\epsilon}M+1)s_{0}^{2}.$$

Now we take $|s - s_0| \ge 2 \left[\sqrt{\epsilon}(2\sqrt{\epsilon}M + 1)\right]^{1/2} s_0$, and conclude that

$$[s^{2} - \mathbf{E}(s)] - [s_{0}^{2} - \mathbf{E}(s_{0})] \ge (s - s_{0})^{2}/4.$$

Proof of Lemma 5.3. Let $s_1 \ge \varsigma_{\min}$ and $s_2 \ge \varsigma_{\min}$ be arbitrary, and define $\hat{f}_1 := f_{\hat{g}_1} := \arg \max_{\varsigma(f) \le s_1} (P_n - P)(f^0 - f), \text{ and } \hat{f}_2 := f_{\hat{g}_2} := \arg \max_{\varsigma(f) \le s_2} (P_n - P)(f^0 - f).$

For all $t \in [0, 1]$, we have

$$\varsigma(f_{t\hat{g}_1+(1-t)\hat{g}_2}) \le t\varsigma(\hat{f}_1) + (1-t)\varsigma(\hat{f}_2) \le ts_1 + (1-t)s_2.$$

In addition, we have

$$\begin{aligned} \hat{\mathbf{E}}_{n}^{\varsigma}(ts_{1}+(1-t)s_{2}) &\geq (P_{n}-P)(f^{0}-f_{t\hat{g}_{1}+(1-t)\hat{g}_{2}}) \\ &= t(P_{n}-P)(f^{0}-\hat{f}_{1})+(1-t)(P_{n}-P)(f^{0}-\hat{f}_{2}) \\ &= t\hat{\mathbf{E}}_{n}^{\varsigma}(s_{1})+(1-t)\hat{\mathbf{E}}_{n}^{\varsigma}(s_{2}), \end{aligned}$$

which completes the proof.

10.5 Proofs for Section 7

Proof of Lemma 7.1. Throughout this proof, we let $0 \leq \tilde{t} \leq t$ be some intermediate point, not the same at each appearance. The function h(t) = d(tg) is infinitely differentiable with h(0) = 0 and h'(0) = 0, so a second-order Taylor series expansion yields $d(tg) = \frac{1}{2}t^2h''(t)$. Consequently, it suffices to show that

$$h''(t) = Pg^2(1 + \mathcal{O}(t)).$$
(22)

Computing derivatives, we have

$$h'(t) = [P \exp[tg]]^{-1} P(\exp[tg]g), \text{ and} h''(t) = [P \exp[tg]]^{-1} P(\exp[tg]g^2) - \{[P \exp[tg]]^{-1} P(\exp[tg]g)\}^2.$$

By a Taylor series expansion of the exponential, we have

$$P \exp[tg] = 1 + tPg + t^2 P(\exp[\tilde{t}g]g^2)/2 = 1 + \mathcal{O}(t^2)Pg^2 = 1 + \mathcal{O}(t^2), \text{ and}$$
$$P \exp[tg]g = tPg^2 + \frac{t^2}{2}P(\exp[\tilde{t}g]g^3) = tPg^2 + \mathcal{O}(t^2)Pg^2 = tPg^2(1 + \mathcal{O}(t)).$$

Combining the pieces, we find that

$$\left[P\exp[tg]\right]^{-1}P(\exp[tg]g) = \left[1 + \mathcal{O}(t^2)\right]^{-1}\left[tPg^2(1 + \mathcal{O}(t))\right] = tPg^2(1 + \mathcal{O}(t)).$$

It follows that

$$\left\{ \left[P \exp[tg] \right]^{-1} P(\exp[tg]g) \right\}^2 = t^2 (Pg^2)^2 (1 + \mathcal{O}(t)) = \mathcal{O}(t^2 Pg^2).$$

But $P(\exp[tg]g^2) = Pg^2 + tP(\exp[\tilde{t}g]g^3) = Pg^2(1+\mathcal{O}(t))$, and hence the bound (22) follows, which completes the proof.

10.6 Proofs for Section 8

Proof of Lemma 8.1.

For each t > 0, we have

$$(P_n - P)(f^0 - f) \le (P_n - P)(f^0 - f) \{F \le t\} + (P_n - P)F \{F > t\} + 2PF \{F > t\},\$$

and also

$$(P_n - P)(f^0 - f) \ge (P_n - P)(f^0 - f) \{F \le t\} - (P_n - P)F \{F > t\} - 2PF \{F > t\}$$

Taking t here equal to $t_0 := C_F \sqrt{\log n}$ (and assuming $t_0 > 1$) we see that

$$2PFl\{F > t_0\} \le 2PF^2l\{F > t_0\} \le 2c_F^2/n.$$

Moreover, for all t > 0, with probability at least $1 - 1/t^2$

$$|(P_n - P)Fl\{F > t_0\}| \le t(P(F^2l\{F > t_0\})^{1/2}/\sqrt{n} \le tc_F/n.$$

Write the truncated versions as

$$\hat{\mathbf{E}}_{n}^{\mathrm{trunc}}(s) := \max_{f \in \mathcal{F}_{s}} (P_{n} - P)(f^{0} - f) \mathbb{I}\{F \leq t\}, \text{ and}$$
$$\mathbf{E}^{\mathrm{trunc}}(s) := \mathbb{E}\left(\max_{f \in \mathcal{F}_{s}} (P_{n} - P)(f^{0} - f) \mathbb{I}\{F \leq t\}\right).$$

Then $|\mathbf{E}(s) - \mathbf{E}^{\text{trunc}}(s)| \le 2c_F^2/n$, and moreover, with probability at least $1 - 1/t^2$, we have

$$\left| \hat{\mathbf{E}}_n(s) - \hat{\mathbf{E}}_n^{\text{trunc}}(s) \right| \le c_F 2(c_F + t)/n.$$

Now Theorem 8.1 ensures that, for all $t \ge 0$,

$$\hat{\mathbf{E}}_{n}^{\text{trunc}}(s) \geq \mathbf{E}^{\text{trunc}}(s) - \sqrt{8C_F\sqrt{\log n}\mathbf{E}^{\text{trunc}}(s) + 2\sigma_s^2}\sqrt{t/n} - C_Ft\sqrt{\log n}/n, \text{ and } \\ \hat{\mathbf{E}}_{n}^{\text{trunc}}(s) \leq \mathbf{E}^{\text{trunc}}(s) + \sqrt{8C_F\sqrt{\log n}\mathbf{E}^{\text{trunc}}(s) + 2\sigma_s^2}\sqrt{t/n} + 2C_Ft\sqrt{\log n}/(3n), \\ \text{where each bound holds with probability at least } 1 - \exp[-t], \square$$

Proof of Lemma 8.2. By Theorem 8.1 of Klein and Rio, for all $t \ge 0$, with probability at least $1 - \exp[-t]$,

$$\hat{\mathbf{E}}_n(s) \ge \mathbf{E}(s) - \sqrt{8K\mathbf{E}(s) + 2\sigma_s^2}\sqrt{t/n} - Kt/n$$

But

$$8K\mathbf{E}(s) \le 8K\mathcal{J}(s)/m_n \le 2C^2s^2 + 2C^2\Phi^*(4K/(m_nC^2)) = 2C^2s^2 + r_0^2.$$

Moreover, we have $\sigma_s^2 \leq C^2 s^2$ by the quadratic curvature condition, and hence

$$\sqrt{8K\mathbf{E}(s) + 2\sigma_s^2}\sqrt{t/n} \le \sqrt{4C^2s^2 + r_0^2}\sqrt{t/n} \le 2Cs\sqrt{t/n} + r_0\sqrt{t/n}.$$

10.7 Proofs for Section 9

Proof of Lemma 9.1. Since $\mathbf{F}(\tilde{s}) = \mathbf{E}(s)$, it follows that

$$\tilde{s}^2 - \mathbf{F}(\tilde{s}) = s^2 - \mathbf{E}(s) - \tau_*^2.$$

We also have

$$|s - s_0| = \left|\sqrt{\tilde{s}^2 + \tau_*^2} - \sqrt{s_*^2 + \tau_*^2}\right| = |\tilde{s} - s_*| \frac{\tilde{s} + s_*}{\sqrt{\tilde{s}^2 + \tau_*^2} + \sqrt{\tilde{s}_*^2 + \tau_*^2}} \le |\tilde{s} - s_*|$$

Proof of Theorem 9.1. By the oracle potential (see Definition 9.1), we know that for all scalars s such that $s^2 \ge \tau_*^2 - \tau_{\min}^2$, we have

$$\sup_{f \in \mathcal{F}: \tau^2(f) \le \tau_*^2 + s^2} P(f - f_0) \le \Gamma^2 s^2,$$

and hence $\sup_{f \in \mathcal{F}, : \tau^2(f) \le \tau_*^2 + s^2} \sigma^2(f - f^0) \le \Gamma^2 C^2 s^2$. We can thus proceed along

the same lines as in the proof of Theorem 4.1, replacing C by ΓC .

References

- C. Borell. The Brunn-Minkowski inequality in Gauss space. Inventiones Mathematicae, 30(2):207–216, 1975.
- S. Boucheron and P. Massart. A high-dimensional Wilks phenomenon. Probability Theory and Related Fields, 150(3-4):405–433, 2011.
- S. Boucheron, G. Lugosi, and P. Massart. *Concentration inequalities: A* nonasymptotic theory of independence. OUP Oxford, 2013.
- S. Chatterjee. A new perspective on least squares under convex constraint. The Annals of Statistics, 42(6):2340–2381, 2014.
- T. Klein. Une inégalité de concentration à gauche pour les processus empiriques. Comptes Rendus Mathematique, 334(6):501–504, 2002.
- T. Klein and E. Rio. Concentration around the mean for maxima of empirical processes. *The Annals of Probability*, 33(3):1060–1077, 2005.
- V. Koltchinskii. Oracle Inequalities in Empirical Risk Minimization and Sparse Recovery Problems: Ecole dEté de Probabilités de Saint-Flour XXXVIII-2008, volume 38. Springer Science & Business Media, 2011.
- M. Ledoux. *The concentration of measure phenomenon*, volume 89. American Mathematical Society, 2001.
- P. Massart. Some applications of concentration inequalities to statistics. In Annales de la Faculté des sciences de Toulouse: Mathématiques, volume 9, pages 245–303, 2000.
- A. Muro and S. van de Geer. Concentration behavior of the penalized least squares estimator, 2015. arXiv:1511.08698.
- R.T. Rockafellar. Convex Analysis. Princeton University Press, 1970.
- A. Saumard. Optimal upper and lower bounds for the true and empirical excess risks in heteroscedastic least-squares regression. *Electronic Journal of Statistics*, 6:579–655, 2012.

M. Talagrand. Concentration of measure and isoperimetric inequalities in product spaces. *Publications Mathématiques de l'IHES*, 81:73–205, 1995.