

Classifying RRATs and FRBs

E.F. Keane¹

¹ *SKA Organisation, Jodrell Bank Observatory, Lower Withington, Macclesfield, Cheshire, SK11 9DL, UK*

6 March 2022

ABSTRACT

In this paper we consider the fact that the simple criterion used to label fast radio transient events as either fast radio bursts (FRBs, thought to be extragalactic with as yet unknown progenitors) or rotating radio transients (RRATs, thought to be Galactic neutron stars) is uncertain. We identify single pulse events reported in the literature which have never been seen to repeat, and which have been labelled as RRATs, but are potentially mis-labelled FRBs. We examine the probability that such ‘grey area’ events are within the Milky Way. The uncertainty in the RRAT/FRB labelling criterion, as well as Galactic-latitude dependent reporting bias may be contributing to the observed latitude dependence of the FRB rate, in addition to effects such as Eddington bias due to scintillation.

Key words: surveys — intergalactic medium — methods: data analysis

1 INTRODUCTION

Fast radio burst (FRB) is the name given to a handful of sub-second radio transient signals detected over the last decade (Keane & Petroff 2015). Because of their many potential uses as tools for studying extreme physical environments and the Universe up to redshifts of $\sim 2 - 3$ (Macquart et al. 2015), there is a large effort underway to search for these signals, both in dedicated ongoing surveys and in archival data (Thornton et al. 2013). FRBs manifest themselves as millisecond-duration events with dispersion measures of several hundred $\text{cm}^{-3} \text{pc}$. No FRB has yet been observed to emit more than one pulse (Petroff et al. 2015b).

Rotating radio transient (RRATs) is the name given to a group of sporadically pulsing sources (Keane & McLaughlin 2011). The majority of RRATs have been seen to repeatedly emit detectable pulses, and it seems clear, from the results of applying pulsar timing techniques (McLaughlin et al. 2006, 2009; Keane et al. 2011), that these sources are Galactic neutron stars with intermittent and/or highly variable pulsar emission.

Single-instance RRAT pulses are no different in appearance to FRB pulses. The criterion for deciding whether such pulses should be classified as FRB or RRAT is comparison of the dispersion measure (DM) with the predicted Galactic maximum of this parameter. In this paper, in § 2, we examine the uncertainty in this classification. Then, in § 3, we estimate the probability that various transients are Galactic or extragalactic, before presenting our concluding thoughts in § 4.

2 NE2001 DISTANCES

The parameter used in deciding whether a dispersed radio burst is an FRB, i.e. extragalactic, is DM_{MW} , the maximum Galactic DM contribution for the particular line of sight. If the burst’s DM exceeds DM_{MW} then it is extragalactic, an FRB, with an inferred $\sim \text{Gpc}$ -scale distance. If the burst’s DM is less than DM_{MW} then the source is Galactic, a RRAT, with an inferred $\sim \text{kpc}$ -scale distance. In practice the model used has thusfar always been NE2001 (Cordes & Lazio 2002); more specifically “NE2001b” as per the labelling of Schnitzeler et al. (2012). Here we investigate the uncertainty of this model, and thus the uncertainty in the RRAT/FRB classification of dispersed radio bursts which have not been seen to repeat.

There are a number of studies in the literature which have addressed the question of the accuracy of the NE2001 model for predicting distances. Deller et al. (2009) used distances to 41 pulsars, determined from VLBI and pulsar timing measurements of parallax, to estimate the distribution of distance errors of the model. More recently, Verbiest et al. (2010, 2012) identified and corrected the Lutz-Kelker bias in the by-then enlarged sample of pulsar distance measurements derived from parallax, supernova remnant associations and neutral hydrogen absorption. Taking the 120 sources in Verbiest et al. (2012) with DM measurements (i.e. those detected at radio wavelengths) we produce a Lutz-Kelker corrected version of the distance error distribution of Deller et al. (2009), shown in Figure 1.

An overestimate (underestimate) of the distance corresponds to an underestimate (overestimate) in the DM inferred *at the correct distance*, and correspondingly an over-

arXiv:1512.02513v2 [astro-ph.HE] 22 Apr 2016




Figure 1. The distribution of the errors in the NE2001 distance estimate, based on the 120 radio-detected sources in Verbiest et al. (2012). This distribution also represents (see text) the error in $x = DM/DM_{MW}$. The error distribution is expressed in decibels, such that a negative (positive) value corresponds to an underestimate (overestimate). The three distributions which are overlapped are those based on the entire Verbiest et al. (2012) sample and the two sub-samples of sources which are below and above a Galactic latitude of 10 degrees.

estimate (underestimate) in $x = DM/DM_{MW}$. Thus the distribution of the NE2001 distance errors maps directly to NE2001 x errors, i.e. RRAT/FRB classification. We can see that the distribution is skewed, with overestimates more likely with a ratio of 70 : 50. However, there is evidence of latitude dependence in the distribution. Considering only those pulsars within 10 degrees of the Galactic plane the ratio of underestimated:overestimated distances is 31 : 55, but is 19 : 15 for sources that are more than 10 degrees from the plane. Assuming that finer resolution in Galactic latitude would see an insufficiently large sample for further resolution, below we consider only these three distributions in estimating the accuracy of RRAT/FRB labelling.

3 FRB OR RRAT?

A search of the literature (see Table 1 for references) shows that there are 12 sources labelled as RRATs, which have never been seen to show a second radio pulse. In Table 1 these are considered in addition to the 16 published FRBs. The probability that each of these events is extragalactic is tabulated, for each of the three distributions of NE2001 errors discussed above: derived from (i) the entire Verbiest et al. (2012) sample; (ii) the $|b| < 10$ degrees sub-sample; and (iii) the $|b| > 10$ degrees sub-sample. Based solely on this simple assessment of NE2001 uncertainties, FRB classifications seem to be mostly sound, with perhaps one source being of uncertain progeny: FRB 010621 has a $\sim 70\%$ probability of being extragalactic. Of the ‘RRAT’ sources, J1354+24 appears to have the highest likelihood of being a mis-labelled FRB, with a probability of $\sim 20-40\%$ of being extragalactic.

Additional outside information can be used to build upon these estimates, where it is available. For example

while one might argue the lack of scattering in FRB 010621 favours an extragalactic origin (Lorimer et al. 2013), identifying a Galactic explanation for FRB 010621 has been the subject of investigation by Keane et al. (2012) and Bannister & Madsen (2014), the latter suggesting that ionised gas unaccounted for in NE2001 can explain much or all of the excess dispersion. Another key piece of additional information is the limit on repeatability. For FRB 121102, Spitler et al. (2014) suggest the observed spectrum, which is dramatically positive, is not intrinsic and a result of a far side-lobe detection. Kulkarni et al. (2015) suggest the observed spectrum is an effect of free-free absorption in a nebula surrounding an extragalactic magnetar. In both scenarios the ~ 4.5 hours of followup may be insufficient to rule out underlying repetition. Obtaining key additional information like this will be essential to elucidate the origins of sources like J1354+24.

4 DISCUSSION & CONCLUSIONS

We have considered the uncertainty in the DM excess parameter $x = DM/DM_{MW}$, crucial in RRAT/FRB classifications in high time resolution radio surveys of the sky. By considering a simple probability-based estimator, we have determined that the FRB identifications made so far are reasonably secure with perhaps one exception (FRB 010621). On the other side of the classification, we find the ‘RRAT’ J1354+24 has a $\sim 20-40\%$ chance of being a mis-labelled FRB; this suggest it merits a deep observational study to address the question of whether it repeats. Similarly, the cumulative probability of at least one RRAT being mis-labelled is at the $\sim 30-80\%$ level; a dedicated campaign to search for a second pulse from each of these sources is merited.

A further concern is a potential bias in the reported single pulse events. In pulsar and transient surveys it is standard practice to consider a source to be a *bona fide* astrophysical source only when it has been observed on at least two epochs (Lorimer & Kramer 2005). This rule, coupled with finite available observing time for re-observing pulsar and transient candidates, continues to contribute to the difficulty in identifying those intermittent pulsars and RRATs with the longest repetition timescales (Kramer et al. 2006). However, this rule is not applied to FRBs: they are simply reported when detected. Given this, and that much more observing time is spent closer to the Galactic plane (Burke-Spolaor & Bannister 2014) we suspect a Galactic-latitude dependent bias in the reported events. To enable correction for this effect survey teams should report single pulse events which appear to be of astrophysical origin, regardless of whether they are seen on a second epoch or have $x < 1$. Probability-based classifiers could then be used for determining the Galactic/extragalactic nature.

The observed FRB rate at Parkes (Petroff et al. 2014a) is higher at higher latitudes. With a perfect electron density model and no reporting bias, such a dependence can still arise, as a result of Eddington bias of the population due to diffractive scintillation at high latitudes (Macquart & Johnston 2015). In this scenario the true rate is that at low latitudes, and the ‘boost factor’ depends directly on the $\log N - \log S$ distribution — the steeper the distribution the larger the boost. But it is clear that the difference between the observed and intrinsic rates, and hence in our ability to

Table 1. Here we list the 12 RRAT sources in the literature which have not yet been seen to show repeat pulses, as well as the 16 published FRBs. The references in the table are given in the second column and are: [1] Thornton et al. (2013); [2] Champion et al. (2016); [3] Petroff et al. (2015a); [4] Masui et al. (2015); [5] Ravi et al. (2015); [6] Lorimer et al. (2007); [7]; Burke-Spolaor & Bannister (2014); [8] Spitler et al. (2014); [9] Keane et al. (2011); [10] Karako-Argaman et al. (2015); [11] Burke-Spolaor & Bailes (2010); [12] Burke-Spolaor et al. (2011). For each signal we then give the excess DM parameter x in the third column and (in dB) in the fourth column. The fifth, sixth and seventh columns each give the probability that the signal is an FRB, for the 3 distributions considered in the text. In each case the exact number of sources is given in parentheses next to the probability; the error on the probabilities is simply the Poisson 95% confidence level (Gehrels 1986). Based on each source’s Galactic latitude the more relevant distribution out of P_{ii} and P_{iii} has been emboldened. The 10 FRBs referred in the first row, are: FRBs 110703, 090625, 130729, 110220, 121002, 120127, 140514, 110626, 110523 and 130626. Similarly, the 8 RRATs referred to in the final row are: J0441–04, J0923–31, J1311–59, J0513–04, J1709–43, J1649–46, J1541–42 and J0845–36.

Source	Reference	x	x (dB)	P_i	P_{ii}	P_{iii}
10 further FRBs	[1,2,3,4]	7.2 to 34.1	11.5 to 15.3	> 0.99 (120/120)	> 0.99 (86/86)	> 0.97 (34/34)
FRB 131104	[5]	10.9	10.4	$0.99^{+0.01}_{-0.14}$ (119/120)	$0.99^{+0.01}_{-0.18}$ (85/86)	> 0.97 (34/34)
FRB 130628	[2]	9.0	9.5	$0.98^{+0.02}_{-0.15}$ (117/120)	$0.97^{+0.03}_{-0.17}$ (83/86)	> 0.97 (34/34)
FRB 010724	[6]	8.4	9.2	$0.97^{+0.03}_{-0.14}$ (116/120)	$0.97^{+0.03}_{-0.17}$ (83/86)	0.97 $^{+0.03}_{-0.26}$ (33/34)
FRB 010125	[7]	7.2	8.6	$0.97^{+0.03}_{-0.14}$ (116/120)	$0.97^{+0.03}_{-0.17}$ (83/86)	0.91 $^{+0.09}_{-0.36}$ (31/34)
FRB 121102	[8]	3.0	4.8	$0.83^{+0.17}_{-0.08}$ (106/120)	0.90 $^{+0.10}_{-0.17}$ (77/86)	$0.85^{+0.15}_{-0.23}$ (29/34)
FRB 010621	[9]	1.4	1.5	$0.74^{+0.14}_{-0.12}$ (89/120)	0.73 $^{+0.27}_{-0.14}$ (63/86)	$0.77^{+0.23}_{-0.24}$ (26/34)
J1354+24	[10]	0.9	−0.4	$0.25^{+0.09}_{-0.07}$ (30/120)	$0.20^{+0.10}_{-0.07}$ (17/86)	0.38 $^{+0.24}_{-0.14}$ (13/34)
J1610−17	[11]	0.7	−1.7	$0.07^{+0.04}_{-0.04}$ (8/120)	$0.02^{+0.05}_{-0.01}$ (2/86)	0.18 $^{+0.17}_{-0.09}$ (6/34)
J1135−49	[12]	0.7	−1.8	$0.07^{+0.04}_{-0.04}$ (8/120)	$0.02^{+0.05}_{-0.01}$ (2/86)	0.18 $^{+0.17}_{-0.09}$ (6/34)
J1059−01	[10]	0.5	−3.0	$0.03^{+0.05}_{-0.02}$ (4/120)	$0.01^{+0.04}_{-0.01}$ (1/86)	0.09 $^{+0.15}_{-0.06}$ (3/34)
8 further RRATs	[9,10,11,12]	−0.4 to 0.1	−4.1 to −10.7	< 0.02 (0/120)	< 0.03 (0/86)	< 0.09 (0/34)

uncover the true $\log N - \log S$ and luminosity distributions, is a combination of a number of effects. In this paper we have tried to tackle the effect of the uncertainty of NE2001. With this, and a clear picture of single epoch pulse events, including those with $x < 1$ (most, but perhaps not all, will be within our Galaxy), we will be able to determine trustworthy metrics for determining the cosmic history of FRBs, whether or not they are standard candles, and their general utility as tools for precision cosmology.

ACKNOWLEDGEMENTS

The author would like to thank the referee, J.-P. Macquart, for valuable input that improved the quality of this paper.

REFERENCES

Bannister K. W., Madsen G. J., 2014, MNRAS, 440, 353
 Burke-Spolaor, S., Bailes M., 2010, MNRAS, 402, 855
 Burke-Spolaor S., et al., 2011, MNRAS, 416, 2465
 Burke-Spolaor, S., Bannister K. W., 2014, ApJ, 792, 19
 Champion D. J., et al., 2016, MNRAS, in press, (astro-ph/1511.07746)
 Cordes J. M., Lazio T. J. W., 2002 (astro-ph/0207156)
 Deller A. T., Tingay, S. J., Bailes, M., Reynolds, J. E., 2009, ApJ, 701, 1243
 Gehrels, N., 1986, ApJ, 303, 336
 Karakao-Argaman C., et al., 2015, ApJ, 809, 67
 Keane E. F., Kramer M., Lyne A. G., Stappers B. W., McLaughlin M. A., 2011, MNRAS, 415, 3065

Keane E. F., McLaughlin M. A., 2011, Bulletin of the Astronomical Society of India, 39, 333
 Keane E. F., Stappers B. W., Kramer M., Lyne A. G., 2012, MNRAS, 425, L71
 Keane E. F., Petroff E., 2015, MNRAS, 447, 2852
 Kramer M., Lyne A. G., O’Brien J. T., Jordan C. A., Lorimer D. R., 2006, Science, 312, 549
 Kulkarni S. R., Ofek E. O., Neill J. D., 2015, preprint (astro-ph/1511.09137).
 Lorimer D. R., Bailes M., McLaughlin M. A., Narkevic D. J., Crawford F., 2007, Science, 318, 777
 Lorimer D. R., Karastergiou A., McLaughlin M. A., Johnston S., 2013, MNRAS, 436, L5
 Lorimer D. R., Kramer M., 2005, Cambridge University Press
 Macquart J.-P., Keane, E. F., Grainge, K., et al. 2015, “Fast transients at cosmological distances”, in proc. *Advancing Astrophysics with the Square Kilometre Array*, PoS(AASKA14)055
 Macquart J.-P., Johnston S., 2015, MNRAS, 451, 3278
 Masui K., et al., 2015, Nature, 528, 523
 McLaughlin M. A. et al., 2006, Nature, 493, 817
 McLaughlin M. A. et al., 2009, MNRAS, 400, 1431
 Petroff E. et al., 2014, ApJ, 789, L26
 Petroff E. et al., 2015a, MNRAS, 447, 246
 Petroff E. et al., 2015b, MNRAS, 451, 3922
 Ravi V., Shannon, M. R., Jameson, A., 2015, ApJ, 799, L5
 Schnitzeler D. et al., 2012, MNRAS, 427, 664
 Spitler L. G. et al., 2014, ApJ, 790, 101
 Thornton D. et al., 2013, Science, 341, 53
 Verbiest J. et al., 2010, MNRAS, 405, 564
 Verbiest J. et al., 2012, ApJ, 755, 39