

Matching of gauge invariant dimension-six operators for $b \rightarrow s$ and $b \rightarrow c$ transitions

Jason Aebischer^a Andreas Crivellin^b Matteo Fael^a Christoph Greub^a

^a*Albert Einstein Center for Fundamental Physics, Institute for Theoretical Physics,
University of Bern, CH-3012 Bern, Switzerland.*

^b*CERN Theory Division, CH-1211 Geneva 23, Switzerland*

E-mail: aebischer@itp.unibe.ch, andreas.crivellin@cern.ch,
fael@itp.unibe.ch, greub@itp.unibe.ch

ABSTRACT: New physics realized above the electroweak scale can be encoded in a model independent way in the Wilson coefficients of higher dimensional operators which are invariant under the Standard Model gauge group. In this article, we study the matching of the $SU(3)_C \times SU(2)_L \times U(1)_Y$ gauge invariant dimension-six operators on the standard B physics Hamiltonian relevant for $b \rightarrow s$ and $b \rightarrow c$ transitions. The matching is performed at the electroweak scale (after spontaneous symmetry breaking) by integrating out the top quark, W , Z and the Higgs particle. We first carry out the matching of the dimension-six operators that give a contribution at tree level to the low energy Hamiltonian. In a second step, we identify those gauge invariant operators that do not enter $b \rightarrow s$ transitions already at tree level, but can give relevant one-loop matching effects.

KEYWORDS: Beyond Standard Model, B-Physics, Rare Decays

Contents

1	Introduction	1
2	Gauge invariant operators	5
2.1	Operator formalism	5
2.2	EW symmetry breaking	6
2.3	$Q_{d\varphi}$ and $Q_{u\varphi}$	8
3	Tree level matching	9
3.1	$\Delta B = \Delta S = 2$	9
3.2	$\Delta B = \Delta C = 1$	10
3.3	$\Delta B = \Delta S = 1$	11
4	One-loop matching corrections	16
4.1	Contribution of 4-fermion operators to 4-fermion operators ($\Delta B = \Delta S = 1$)	17
4.2	Contribution of 4-fermion operators to 4-fermion operators ($\Delta B = \Delta S = 2$)	20
4.3	Contributions of 4-fermion operators to O_7 and O_8	20
4.4	Contributions of right-handed Z couplings to O_9 , O_{10} and O_{3-6}^q	20
4.5	Contributions of right-handed W couplings to O_7 and O_8	21
4.6	Contributions of magnetic operators to O_7, O_8, O_9, O_{10} and O_4^q	21
5	Phenomenological example	24
6	Conclusions	25
A	Dimension-six operators in the mass basis	26

1 Introduction

The Standard Model (SM) of particle physics as the gauge theory of strong and electroweak (EW) interactions has been tested and confirmed to a high precision since many years [1].

Furthermore, the observation of a Higgs boson at the LHC [2, 3] and the first measurements of its production and decay channels are consistent with the SM Higgs mechanism of EW symmetry breaking.

Nevertheless, the SM is expected to constitute only an effective theory valid up to a new physics (NP) scale Λ where additional dynamic degrees of freedom enter. A renormalizable quantum field theory of NP, realized at a scale higher than the EW one, satisfies in general the following requirements:

- (i) Its gauge group must contain the SM gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y$ as a subgroup.
- (ii) All SM degrees of freedom should be contained, either as fundamental or as composite fields.
- (iii) At low-energies the SM should be reproduced, provided no undiscovered weakly coupled *light* particles exist (like axions or sterile neutrinos).

In most theories of physics beyond the SM that have been considered, the SM is recovered via the decoupling of heavy particles, with masses $\Lambda \gg M_Z$, guaranteed, at the perturbative level, by the Appelquist-Carazzone decoupling theorem [4]. Therefore, NP can be encoded in higher-dimensional operators which are suppressed by powers of the NP scale Λ :

$$\mathcal{L}_{SM} = \mathcal{L}_{SM}^{(4)} + \frac{1}{\Lambda} C_{\nu\nu}^{(5)} Q_{\nu\nu}^{(5)} + \frac{1}{\Lambda^2} \sum_k C_k^{(6)} Q_k^{(6)} + \mathcal{O}\left(\frac{1}{\Lambda^3}\right). \quad (1.1)$$

Here $\mathcal{L}_{SM}^{(4)}$ is the usual renormalizable SM Lagrangian which contains only dimension-two and dimension-four operators, $Q_{\nu\nu}^{(5)}$ is the Weinberg operator giving rise to neutrino masses [5], $Q_k^{(6)}$ and $C_k^{(6)}$ denote the dimension-six operators and their corresponding Wilson coefficients, respectively [6, 7].

Even if the ultimate theory of NP was not a quantum field theory, at low energies it would be described by an effective non-renormalizable Lagrangian [8] and it would be possible to parametrize its effects at the EW scale in terms of the Wilson coefficients associated to these operators. Thus, one can search for NP in a model independent way by studying the SM extended with higher-dimensional gauge-invariant operators. Once a specific NP model is chosen, the Wilson coefficients can be expressed in terms of the NP parameters by matching the beyond the SM theory under consideration on the SM enlarged with such higher dimensional operators.

Flavor observables, especially flavor changing neutral current processes, are excellent probes of physics beyond the SM: since in the SM they are suppressed by the Fermi constant G_F as well as by small CKM elements and loop factors they are very sensitive to even small NP contributions. Therefore, on one hand flavor processes can stringently constrain the Wilson coefficients of the dimension-six operators induced by NP. On the other hand, if deviations from the SM were uncovered, flavor physics could be used as a guideline towards




Figure 1: Mass scale hierarchy: the matching of the NP model onto the gauge invariant dimension-six operators is performed at the mass scale Λ . After the EW symmetry breaking, the matching of dimension-six operators on the effective Hamiltonian governing B physics is performed at the mass scale μ_W . The Wilson coefficients at different mass scales are connected via RGE evolution.

the construction of a theory of physics beyond the SM. The second point is especially interesting nowadays in light of the discrepancies between the SM predictions and the measurements of $b \rightarrow s\mu^+\mu^-$ and $b \rightarrow c\tau\nu$ processes: the combination of $B \rightarrow D^*\tau\nu$ and $B \rightarrow D\tau\nu$ branching fractions disagrees with the SM prediction [9] at the level of 3.9 standard deviations (σ) [10]. Furthermore, $b \rightarrow s\ell^+\ell^-$ global fits even show deviations between 4σ and 5σ [11–13]. These deviations have been extensively studied recently. Many NP models have been proposed to explain the anomalies, (see for example [14–35] for $b \rightarrow s\mu^+\mu^-$ and [34–47] for $b \rightarrow c\tau\nu$). Therefore, at the moment, B physics is probably our best guideline towards NP.

The effective field theory approach is an essential ingredient of all B physics calculations within and beyond the SM. However, the Hamiltonian governing $b \rightarrow s$ and $b \rightarrow c$ transitions is not invariant under the full SM gauge group, but only under $SU(3)_C \times U(1)_{\text{EM}}$ since it is defined below the EW scale where $SU(2)_L \times U(1)_Y$ is broken (see for example [48, 49] for a review of the use of effective Hamiltonians in B physics). As a consequence, the SM extended with gauge invariant dimension-six operators must be matched onto the low energy effective Hamiltonian governing B physics (see figure 1). In the flavor sector only partial analyses exist: the matching effects in the lepton sector were calculated in refs. [50–52],¹ while in the quark sector $b \rightarrow s\mu^+\mu^-$ transitions and their correlations with $B \rightarrow K^{(*)}\nu\nu$ and $B \rightarrow D^{(*)}\tau\nu$ were studied in refs. [54–58]. However a systematic and complete phe-

¹See ref. [53] for an analysis of non-gauge invariant effective operators for tau decays.

nomenological study of the gauge invariant dimension-six operators in B physics is still missing. Such analysis proceeds, in a bottom-up approach, in the following three steps.

- (i) The matching at the EW scale μ_W , of the order of M_W , of the gauge invariant operators onto the low-energy B physics Hamiltonian by integrating out the heavy degrees of freedom represented by the top quark, the Higgs and the Z and W bosons. It is the aim of this article to perform such systematic matching of the gauge invariant operators.
- (ii) The evolution of the effective Hamiltonian's Wilson coefficients from the scale μ_W down to the B meson scale μ_b , where μ_b is of the order of m_b . This is obtained by solving the appropriate renormalization group equation (RGE). We note that after the matching procedure the set of operators in the B physics Hamiltonian is larger than the SM one since new Lorentz structures must be taken into account, therefore the anomalous dimension matrices get also bigger compared to the SM.²
- (iii) The assessment of the constraints on the dimension-six operators' Wilson coefficients (defined at the EW scale μ_W) stemming from the available flavor observables. An example of such analysis can be found in the section 5, while the complete numerical analysis will be given in a subsequent publication.

The purpose of the outlined study is to depict the general pattern of deviations observed in B physics employing dimension-six operators. It is worth noting however that in the framework of higher dimensional operators, in order to correctly interpret any deviations of the SM in terms of a specific NP model, it is necessary to map the pattern of deviations observed at the EW scale back to the scale Λ where the BSM physics was supposedly integrated out (see figure 1). Indeed due to operator mixing, the pattern of deviations at the EW scale differs from the pattern of Wilson coefficients at the matching scale Λ . The connection between these two mass scales is given by the RGE evolution of dimension-six operators [63–65].

The outline of this article is as follows: in section 2 we list the operators relevant for our analysis and discuss the EW symmetry breaking. Then, in section 3, we establish our conventions for the B physics Hamiltonian and we perform the complete matching of the dimension-six operators that give contributions to $b \rightarrow s$ or $b \rightarrow c$ transitions at tree level. In section 4 we identify and calculate the leading one-loop EW matching corrections for $b \rightarrow s$ processes for those operators which do not enter $b \rightarrow s$ transitions already at tree-level. A phenomenological application of the computed matching conditions will be given in section 5. Finally we conclude.

²For the anomalous dimension matrices beyond the SM for $\Delta F = 2$ processes see for example refs. [59, 60], for 4-fermion operators ref. [60] and for $b \rightarrow s\gamma$ refs. [61, 62].

2 Gauge invariant operators

In this section we list the gauge invariant operators, following the conventions of ref. [7], that contribute to $b \rightarrow s$ or $b \rightarrow c$ transitions at tree-level. Here we only consider operators involving quark fields. The importance of flavor physics in constraining operators which modify triple gauge couplings has been studied in ref. [66]. Recall that the gauge invariant dimension-six operators are defined before EW symmetry breaking, implying that they are given in the interaction basis (as the mass basis it not yet defined). After the EW symmetry breaking, the fermions acquire their masses and the necessary diagonalizations of their mass matrices affect the Wilson coefficients. As we will see, all these rotations can be absorbed by a redefinition of the Wilson coefficients, except for the misalignment between the left-handed up-quark and down-quark rotations, i.e. the Cabibbo-Kobayashi-Maskawa matrix (CKM) which relates charged and neutral currents.

2.1 Operator formalism

In table 1 we list the operators contributing to $b \rightarrow s$ at the tree level (and possibly also to $b \rightarrow c$ transitions), while table 2 gives the operators generating at tree level $b \rightarrow c$ but not $b \rightarrow s$. For the SM Lagrangian we adopt the standard definition

$$\mathcal{L}_{SM}^{(4)} = -\frac{1}{4}G_{\mu\nu}^A G^{A\mu\nu} - \frac{1}{4}W_{\mu\nu}^I W^{I\mu\nu} - \frac{1}{4}B_{\mu\nu}B^{\mu\nu} + (D_\mu\varphi)^\dagger(D^\mu\varphi) + m^2\varphi^\dagger\varphi - \frac{1}{2}\lambda\left(\varphi^\dagger\varphi\right)^2 + i\left(\bar{\ell}\not{D}\ell + \bar{e}\not{D}e + \bar{q}\not{D}q + \bar{u}\not{D}u + \bar{d}\not{D}d\right) - \left(\bar{\ell}Y_e e\varphi + \bar{q}Y_u u\tilde{\varphi} + \bar{q}Y_d d\varphi + \text{h.c.}\right), \quad (2.1)$$

where ℓ , q and φ stand for the lepton, quark and Higgs $SU(2)_L$ doublets, respectively, while the right-handed isospin singlets are denoted by e , u and d . Here $\tilde{\varphi}^i = \varepsilon_{ij}(\varphi^j)^*$, where ε_{ij} is the totally antisymmetric tensor with $\varepsilon_{12} = +1$. Flavor indices $i, j, k, l = 1, 2, 3$ are implicitly assigned to each fermion field appearing in (2.1), and the Yukawa couplings $Y_{e,u,d}$ are matrices in the generation space. Therefore, in table 1 the operator names in the left column of each block should be supplemented with generation indices of the fermion fields whenever necessary. Covariant derivatives are defined with the plus sign, i.e. for example

$$D_\mu q = (\partial_\mu + ig_s T^A G_\mu^A + ig \frac{\tau^I}{2} W_\mu^I + ig' Y B_\mu) q, \quad (2.2)$$

where Y is the hypercharge and $T^A = \frac{1}{2}\lambda^A$; λ^A and τ^I are the Gell-Mann and Pauli matrices, respectively. With the above definition for the covariant derivative, the gauge field strength tensors read

$$\begin{aligned} G_{\mu\nu}^A &= \partial_\mu G_\nu^A - \partial_\nu G_\mu^A - g_s f^{ABC} G_\mu^B G_\nu^C, \\ W_{\mu\nu}^I &= \partial_\mu W_\nu^I - \partial_\nu W_\mu^I - g \varepsilon^{IJK} W_\mu^J W_\nu^K, \\ B_{\mu\nu} &= \partial_\mu B_\nu - \partial_\nu B_\mu. \end{aligned} \quad (2.3)$$

Moreover the Hermitian derivative terms are defined as

$$\varphi^\dagger i\overleftrightarrow{D}_\mu \varphi = i\varphi^\dagger(D_\mu - \overleftarrow{D}_\mu)\varphi, \quad \varphi^\dagger i\overleftrightarrow{D}_\mu^I \varphi = i\varphi^\dagger(\tau^I D_\mu - \overleftarrow{D}_\mu \tau^I)\varphi, \quad (2.4)$$

$(\overline{LR})(\overline{RL})$ or $(\overline{LR})(\overline{LR})$		$(\overline{LL})(\overline{LL})$		$\psi^2 X \varphi$	
$Q_{\ell edq}$	$(\bar{\ell}_i^a e_j)(\bar{d}_k q_l^a)$	$Q_{qq}^{(1)}$	$(\bar{q}_i \gamma_\mu q_j)(\bar{q}_k \gamma^\mu q_l)$	Q_{dW}	$(\bar{q}_i \sigma^{\mu\nu} d_j) \tau^I \varphi W_{\mu\nu}^I$
$Q_{quqd}^{(1)}$	$(\bar{q}_i^a u_j) \varepsilon_{ab} (\bar{q}_k^b d_l)$	$Q_{\ell q}^{(1)}$	$(\bar{\ell}_i \gamma_\mu \ell_j)(\bar{q}_k \gamma^\mu q_l)$	Q_{dB}	$(\bar{q}_i \sigma^{\mu\nu} d_j) \varphi B_{\mu\nu}$
$Q_{quqd}^{(8)}$	$(\bar{q}_i^a T^A u_j) \varepsilon_{ab} (\bar{q}_k^b T^A d_l)$	$Q_{qq}^{(3)}$	$(\bar{q}_i \gamma_\mu \tau^I q_j)(\bar{q}_k \gamma^\mu \tau^I q_l)$	Q_{dG}	$(\bar{q}_i \sigma^{\mu\nu} T^A d_j) \varphi G_{\mu\nu}^A$
$(\overline{LL})(\overline{RR})$		$Q_{\ell q}^{(3)}$	$(\bar{\ell}_i \gamma_\mu \tau^I \ell_j)(\bar{q}_k \gamma^\mu \tau^I q_l)$	$\psi^2 \varphi^3$	
$Q_{\ell d}$	$(\bar{\ell}_i \gamma_\mu \ell_j)(\bar{d}_k \gamma^\mu d_l)$	$(\overline{RR})(\overline{RR})$		$Q_{d\varphi}$	$(\varphi^\dagger \varphi)(\bar{q}_i d_j \varphi)$
Q_{qe}	$(\bar{q}_i \gamma_\mu q_j)(\bar{e}_k \gamma^\mu e_l)$	$\psi^2 \varphi^2 D$			
$Q_{qu}^{(1)}$	$(\bar{q}_i \gamma_\mu q_j)(\bar{u}_k \gamma^\mu u_l)$	Q_{dd}	$(\bar{d}_i \gamma_\mu d_j)(\bar{d}_k \gamma^\mu d_l)$	$Q_{\varphi q}^{(1)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{q}_i \gamma^\mu q_j)$
$Q_{qd}^{(1)}$	$(\bar{q}_i \gamma_\mu q_j)(\bar{d}_k \gamma^\mu d_l)$	Q_{ed}	$(\bar{e}_i \gamma_\mu e_j)(\bar{d}_k \gamma^\mu d_l)$	$Q_{\varphi q}^{(3)}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu^I \varphi)(\bar{q}_i \tau^I \gamma^\mu q_j)$
$Q_{qu}^{(8)}$	$(\bar{q}_i \gamma_\mu T^A q_j)(\bar{u}_k \gamma^\mu T^A u_l)$	$Q_{ud}^{(1)}$	$(\bar{u}_i \gamma_\mu u_j)(\bar{d}_k \gamma^\mu d_l)$	$Q_{\varphi d}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{d}_i \gamma^\mu d_j)$
$Q_{qd}^{(8)}$	$(\bar{q}_i \gamma_\mu T^A q_j)(\bar{d}_k \gamma^\mu T^A d_l)$	$Q_{ud}^{(8)}$	$(\bar{u}_i \gamma_\mu T^A u_j)(\bar{d}_k \gamma^\mu T^A d_l)$	$Q_{\varphi ud}$	$i(\tilde{\varphi}^\dagger D_\mu \varphi)(\bar{u}_i \gamma^\mu d_j)$

Table 1: Complete list of the dimension-six operators that contribute to $b \rightarrow s$ (and possibly also to $b \rightarrow c$) transitions at tree level.

$(\overline{LR})(\overline{LR})$	
$Q_{\ell equ}^{(1)}$	$(\bar{\ell}_i^a e_j) \varepsilon_{ab} (\bar{q}_k^b u_l)$
$Q_{\ell equ}^{(3)}$	$(\bar{\ell}_i^a \sigma^{\mu\nu} e_j) \varepsilon_{ab} (\bar{q}_k^b \sigma_{\mu\nu} u_l)$

Table 2: The two dimension-six operators that contribute to $b \rightarrow c$ but not to $b \rightarrow s$ transitions at tree level.

where $\varphi^\dagger \overleftrightarrow{D}_\mu \varphi = (D_\mu \varphi)^\dagger \varphi$. For further details concerning conventions and notations, we refer the reader to ref. [7].

For the operators in the classes $(\overline{LL})(\overline{LL})$, $(\overline{LL})(\overline{RR})$, $(\overline{RR})(\overline{RR})$ and $\psi^2 \varphi^2 D$ (except for $Q_{\varphi ud}$), hermitian conjugation is equivalent to the transposition of generation indices in each of the fermion currents. Moreover, the operators $Q_{qq}^{(1)}$, $Q_{qq}^{(3)}$, Q_{uu} and Q_{dd} are symmetric under exchange of the flavor indices $ij \leftrightarrow kl$. Therefore, we will restrict ourselves to the operators satisfying $[ij] < [kl]$, where $[ij]$ denotes the two digit number $[ij] = 10i + j$.

2.2 EW symmetry breaking

Although the set of gauge invariant dimension-six operators we have just introduced is written in term of the flavor basis, actual calculations that confront theory with experiment are performed using the mass eigenbasis which is defined after the EW symmetry breaking. In the broken phase, flavor and mass eigenstates are not identical and the $SU(2)_L$ doublet components are distinguishable. Therefore, we need to rotate the weak eigenstates into mass eigenstates via the following transformations:

$$u_L^i \rightarrow S_{L\,ij}^u u_L^j, \quad u_R^i \rightarrow S_{R\,ij}^u u_R^j, \quad (2.5)$$

$$d_L^i \rightarrow S_{L\,ij}^d d_L^j, \quad d_R^i \rightarrow S_{R\,ij}^d d_R^j, \quad (2.6)$$

where S_L^d, S_R^d, S_L^u and S_R^u are the 3×3 unitary matrices in flavor space that diagonalize the mass matrix as

$$S_{L\,ii'}^{q\dagger} m_q^{i'j'} S_{R\,j'j}^q = m_{q_i} \delta_{ij}. \quad (2.7)$$

With these definitions, the CKM matrix V is given by

$$V = (S_L^u)^\dagger S_L^d. \quad (2.8)$$

After these necessary field redefinitions, there are no flavor changing neutral currents at tree-level in the SM, due to the unitarity of the transformation matrices, and mixing between generations only occurs in the charged quark current. When dimension-six operators are included in the Lagrangian, the effect on them by the matrices $S_{L,R}^q$ cannot be eliminated by unitarity. However, these rotations can be absorbed into the Wilson coefficients. As a first example, we consider the operator $Q_{\varphi d}$ which takes the form:

$$C_{\varphi d}^{mn} Q_{\varphi d}^{mn} = C_{\varphi d}^{mn} \left(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) \left(\bar{d}_R^m \gamma^\mu d_R^n \right) \rightarrow C_{\varphi d}^{mn} \left(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) \left(\bar{d}_R^i S_{R\,im}^{d\dagger} \gamma^\mu S_{R\,nj}^d d_R^j \right). \quad (2.9)$$

Redefining

$$\tilde{C}_{\varphi d}^{ij} = C_{\varphi d}^{mn} S_{R\,im}^{d\dagger} S_{R\,nj}^d, \quad (2.10)$$

we can indeed absorb $S_{L,R}^q$ into the overall coefficient:

$$C_{\varphi d}^{mn} Q_{\varphi d}^{mn} = \tilde{C}_{\varphi d}^{ij} \left(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) \left(\bar{d}_R^i \gamma^\mu d_R^j \right). \quad (2.11)$$

In contrast to the SM, it is not possible anymore to avoid the appearance of flavor changing neutral currents for all operators. Moreover, the redefinitions of the Wilson coefficients are not unique, in general. Let us consider as a second example the operator $Q_{\varphi q}^{(1)}$:

$$C_{\varphi q}^{(1)mn} Q_{\varphi q}^{(1)mn} = C_{\varphi q}^{(1)mn} \left(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) \left(\bar{u}_L^m \gamma^\mu u_L^n + \bar{d}_L^m \gamma^\mu d_L^n \right) \quad (2.12)$$

$$\rightarrow C_{\varphi q}^{(1)mn} \left(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) \left(\bar{u}_L^i S_{L\,im}^{u\dagger} \gamma^\mu S_{L\,nj}^u u_L^j + \bar{d}_L^i S_{L\,im}^{d\dagger} \gamma^\mu S_{L\,nj}^d d_L^j \right). \quad (2.13)$$

In this case we cannot absorb at the same time the rotation for the up quarks and for the down quarks, so that we can choose to define

$$\tilde{C}_{\varphi q}^{(1)ij} = C_{\varphi q}^{(1)mn} S_{L\,im}^{d\dagger} S_{L\,nj}^d, \quad (2.14)$$

or

$$\check{C}_{\varphi q}^{(1)ij} = C_{\varphi q}^{(1)mn} S_{L\,im}^{u\dagger} S_{L\,nj}^u, \quad (2.15)$$

obtaining the two equivalent expressions

$$C_{\varphi q}^{(1)mn} Q_{\varphi q}^{(1)mn} = \tilde{C}_{\varphi q}^{(1)ij} \left(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) \left(V_{ki} V_{lj}^* \bar{u}_L^k \gamma^\mu u_L^l + \bar{d}_L^i \gamma^\mu d_L^j \right) \quad (2.16)$$

$$= \check{C}_{\varphi q}^{(1)ij} \left(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) \left(\bar{u}_L^i \gamma^\mu u_L^j + V_{ik}^* V_{jl} \bar{d}_L^k \gamma^\mu d_L^l \right). \quad (2.17)$$

For both definitions, the mass diagonalization leads to flavor changing neutral currents either in the up sector, for the coefficient denoted with the tilde (\sim), or in the down sector for that one with the check (\checkmark). The two notations are related through the identity

$$\check{C}^{ij} = V_{ik} V_{jl}^* \tilde{C}^{kl}. \quad (2.18)$$

All operators reported in table 1 must be analogously expressed in the mass basis. We report in appendix A the explicit expressions for the Wilson coefficients \tilde{C} .

2.3 $Q_{d\varphi}$ and $Q_{u\varphi}$

The operators $Q_{d\varphi}$ and $Q_{u\varphi}$ play a special role as they contribute to the quark mass matrices after the EW symmetry breaking. For example, the down-quark mass matrix receives two contributions, one from the SM Yukawa interactions and one from the operator $Q_{d\varphi}$:

$$m_d^{ij} = \frac{v}{\sqrt{2}} \left(Y_d^{ij} - \frac{1}{2} \frac{v^2}{\Lambda^2} C_{d\varphi}^{ij} \right), \quad (2.19)$$

where Y_d is the Yukawa matrix of the SM and $v = 246$ GeV is the vacuum expectation value of the SM Higgs field. For the coupling of the Higgs with the down-type quarks, defined by the Lagrangian term $\mathcal{L}_H = -h \bar{d}_L \Gamma^h d_R + \text{h.c.}$, the extra contribution is enhanced by a combinatorial factor of three compared to the contribution to the mass term:

$$\Gamma_{d_i d_j}^h = \frac{1}{\sqrt{2}} \left(Y_d^{ij} - \frac{3}{2} \frac{v^2}{\Lambda^2} C_{d\varphi}^{ij} \right) = \frac{m_d^{ij}}{v} - \frac{1}{\sqrt{2}} \frac{v^2}{\Lambda^2} C_{d\varphi}^{ij}. \quad (2.20)$$

Unlike in the pure dimension-four SM, the mass matrix and the quark-Higgs coupling cannot be diagonalized simultaneously: a flavor changing interaction between the SM Higgs and the quarks appears [51, 67, 68]. Indeed the first term in eq. (2.20) is rendered diagonal by a field redefinition as in (2.6),

$$U_{L ii'}^{d\dagger} m_d^{i' j'} U_{R j' j}^d = m_{d_i} \delta_{ij}, \quad (2.21)$$

where the new $U_{L,R}^d$ matrices, necessary to diagonalize the mass in the presence of the $Q_{d\varphi}$ operator, differ from $S_{L,R}^d$ by terms of order $1/\Lambda^2$. The quark-Higgs coupling matrix is now given by

$$\Gamma_{d_i d_j}^h = \frac{m_{d_i}}{v} \delta_{ij} - \frac{1}{\sqrt{2}} \frac{v^2}{\Lambda^2} \tilde{C}_{d\varphi}^{ij}, \quad (2.22)$$

where we have defined

$$\tilde{C}_{d\varphi}^{ij} = \left(U_L^{d\dagger} C_{d\varphi} U_R^d \right)_{ij} = \left(S_L^{d\dagger} C_{d\varphi} S_R^d \right)_{ij} + O\left(\frac{1}{\Lambda^2}\right). \quad (2.23)$$

Note that in this approximation all Wilson coefficients of the operators discussed above remain unchanged since the extra rotation induced by the $Q_{d\varphi}$ operator would lead to a $1/\Lambda^4$ effect. Similar considerations apply to the operator $Q_{u\varphi}$.

3 Tree level matching

In this section we perform the tree-level matching of the gauge invariant dimension-six operators relevant for $b \rightarrow s$ and $b \rightarrow c$ transitions. This matching is performed at the EW scale on the effective Hamiltonian governing B physics, which is defined below the EW scale. Therefore, the effective B physics Hamiltonian contains the SM fields without W , Z , the Higgs and the top quark, while these are dynamical fields of the gauge invariant dimension-six operator basis. As we will see, the B -physics Hamiltonian contains operators with additional Lorentz structures compared to the ones relevant in the SM.

3.1 $\Delta B = \Delta S = 2$

In this section we consider B_s - \overline{B}_s mixing. Here, following the conventions of refs. [59, 69], the effective Hamiltonian is given by

$$\mathcal{H}_{\text{eff}}^{\Delta B = \Delta S = 2} = \sum_{j=1}^5 C_j O_j + \sum_{j=1}^3 C'_j O'_j + \text{h.c.}, \quad (3.1)$$

with the operators defined as

$$\begin{aligned} O_1 &= (\overline{s} \gamma_\mu P_L b) (\overline{s} \gamma^\mu P_L b), & O_2 &= (\overline{s} P_L b) (\overline{s} P_L b), \\ O_3 &= (\overline{s}_\alpha P_L b_\beta) (\overline{s}_\beta P_L b_\alpha), & O_4 &= (\overline{s} P_L b) (\overline{s} P_R b), \\ O_5 &= (\overline{s}_\alpha P_L b_\beta) (\overline{s}_\beta P_R b_\alpha), \end{aligned} \quad (3.2)$$

where α and β are color indices. The primed operators $O'_{1,2,3}$ are obtained from $O_{1,2,3}$ by interchanging P_L with P_R .

The contributions from the four-fermion operators to the Hamiltonian in eq. (3.1) read:

$$C_1 = -\frac{1}{\Lambda^2} \left[\tilde{C}_{qq}^{(1)2323} + \tilde{C}_{qq}^{(3)2323} \right], \quad (3.3)$$

$$C'_1 = -\frac{1}{\Lambda^2} \tilde{C}_{dd}^{2323}, \quad (3.4)$$

$$C_4 = \frac{1}{\Lambda^2} \tilde{C}_{qd}^{(8)2323}, \quad (3.5)$$

$$C_5 = \frac{1}{\Lambda^2} \left[2\tilde{C}_{qd}^{(1)2323} - \frac{1}{N_c} \tilde{C}_{qd}^{(8)2323} \right], \quad (3.6)$$

where N_c denotes the number of colors. In addition, we include for completeness the effects of $Q_{d\varphi}$ even though they are formally suppressed by $1/\Lambda^4$ because the $1/\Lambda^2$ effect in the B -physics Hamiltonian is suppressed due to the m_f/v coupling of the Higgs to the light

fermions.³ Here we get

$$C_2 = -\frac{1}{2m_h^2} \left(\Gamma_{bs}^{h*} \right)^2, \quad (3.7)$$

$$C'_2 = -\frac{1}{2m_h^2} \left(\Gamma_{sb}^h \right)^2, \quad (3.8)$$

$$C_4 = -\frac{1}{m_h^2} \Gamma_{sb}^h \Gamma_{bs}^{h*}, \quad (3.9)$$

where $\Gamma_{d_i d_j}^h$ is defined in eq. (2.20). Note that we do not include the analogous contributions from a modified Z coupling since in this case the coupling to light fermions are not suppressed and especially $b \rightarrow s\mu^+\mu^-$ processes will give relevant tree-level constraints at the $1/\Lambda^2$ level.

3.2 $\Delta B = \Delta C = 1$

For the charged current process $b \rightarrow c\ell_i\nu_j$ we write the effective Hamiltonian as

$$\mathcal{H}_{\text{eff}}^{\Delta B=\Delta C=1} = -\frac{4G_F}{\sqrt{2}} \left[C_T O_T + \sum_{i=S,V} C_i O_i + C'_i O'_i \right], \quad (3.10)$$

where the operators are

$$O_V = (\bar{c} \gamma^\mu P_L b) (\bar{\ell} \gamma_\mu P_L \nu), \quad O_T = (\bar{c} \sigma^{\mu\nu} P_L b) (\bar{\ell} \sigma_{\mu\nu} P_L \nu), \quad O_S = (\bar{c} P_L b) (\bar{\ell} P_L \nu), \quad (3.11)$$

and the prime operators are obtained by interchanging $P_L \leftrightarrow P_R$ in the quark current.⁴

The four-fermion operators lead to the following contribution to the effective Hamiltonian:

$$C_V = \frac{v^2}{\Lambda^2} V_{ci} \tilde{C}_{\ell q}^{(3)lli3}, \quad C'_S = \frac{v^2}{2\Lambda^2} V_{ci} \tilde{C}_{\ell eq}^{*ll3i}, \quad (3.12)$$

$$C_S = \frac{v^2}{2\Lambda^2} V_{ib} \tilde{C}_{\ell equ}^{*(1)lli2}, \quad C_T = \frac{v^2}{2\Lambda^2} V_{ib} \tilde{C}_{\ell equ}^{*(3)lli2}, \quad (3.13)$$

where the summation over $i = 1, 2, 3$ is understood. The operators $Q_{\varphi ud}$ and $Q_{\varphi q}^{(3)}$ induce an anomalous u - d - W coupling. Their contribution to the $b \rightarrow c\ell\nu$ transition reads:

$$C'_V = -\frac{v^2}{2\Lambda^2} \tilde{C}_{\varphi ud}^{23}, \quad (3.14)$$

$$C_V = -\frac{v^2}{\Lambda^2} V_{ci} \tilde{C}_{\varphi q}^{(3)i3}. \quad (3.15)$$

³Note that this counting argument already suggest, that the EFT approach to flavor changing Higgs decays has quite limited applicability.

⁴The operator O'_T is identically zero due to Fierz transformations.

The effect of such modified W couplings to quarks on the determination of V_{cb} (and analogously on V_{ub}) has been discussed in refs. [70–80].

In principle, also momentum dependent modifications of the W - c - b coupling can lead to effects in $b \rightarrow c\ell\nu$ transitions as examined in refs. [73, 78] at the level of non-gauge invariant operators. However, these effects scale like $m_b v/(m_W^2 \Lambda^2)$. Furthermore, also corrections to Z - b - b couplings can appear which are stringently constrained, making the possible contributions tiny [79]. Therefore we do not include these effects here.

3.3 $\Delta B = \Delta S = 1$

We describe the $b \rightarrow s\ell^-\ell'^+$ and $b \rightarrow s\gamma$ transition via the effective Hamiltonian

$$\mathcal{H}_{\text{eff}}^{\Delta B=\Delta S=1} = -\frac{4G_F}{\sqrt{2}} \left(\sum_i C_i O_i + C'_i O'_i + \sum_i \sum_q C_i^q O_i^q + C_i^{'q} O_i^{'q} \right), \quad (3.16)$$

where the index q runs over all light quarks $q = u, d, c, s, b$. The operators contributing in the first part are:

$$\begin{aligned} O_1 &= (\bar{s} T^A \gamma_\mu P_L c) (\bar{c} T^A \gamma^\mu P_L b), & O_2 &= (\bar{s} \gamma_\mu P_L c) (\bar{c} \gamma^\mu P_L b), \\ O_7 &= \frac{e}{16\pi^2} m_b (\bar{s} \sigma_{\mu\nu} P_R b) F^{\mu\nu}, & O_8 &= \frac{g_s}{16\pi^2} m_b (\bar{s} T^A \sigma_{\mu\nu} P_R b) G^{\mu\nu A}, \\ O_9^{\ell\ell'} &= \frac{e^2}{16\pi^2} (\bar{s} \gamma_\mu P_L b) (\bar{\ell} \gamma^\mu \ell'), & O_{10}^{\ell\ell'} &= \frac{e^2}{16\pi^2} (\bar{s} \gamma_\mu P_L b) (\bar{\ell} \gamma^\mu \gamma_5 \ell'), \\ O_S^{\ell\ell'} &= (\bar{s} P_R b) (\bar{\ell} \ell'), & O_P^{\ell\ell'} &= (\bar{s} P_R b) (\bar{\ell} \gamma_5 \ell'), \\ O_T^{\ell\ell'} &= (\bar{s} \sigma^{\mu\nu} b) (\bar{\ell} \sigma_{\mu\nu} \ell'), & O_{T5}^{\ell\ell'} &= (\bar{s} \sigma^{\mu\nu} b) (\bar{\ell} \sigma_{\mu\nu} \gamma_5 \ell'). \end{aligned} \quad (3.17)$$

While in the second part of the Hamiltonian we have four-quark operators with vectorial Lorentz structures,

$$\begin{aligned} O_3^q &= (\bar{s} \gamma_\mu P_L b) (\bar{q} \gamma^\mu q), & O_4^q &= (\bar{s} T^A \gamma_\mu P_L b) (\bar{q} T^A \gamma^\mu q), \\ O_5^q &= (\bar{s} \gamma_\mu \gamma_\nu \gamma_\rho P_L b) (\bar{q} \gamma^\mu \gamma^\nu \gamma^\rho q), & O_6^q &= (\bar{s} T^A \gamma_\mu \gamma_\nu \gamma_\rho P_L b) (\bar{q} T^A \gamma^\mu \gamma^\nu \gamma^\rho q), \end{aligned} \quad (3.18)$$

and four-quark operators with scalar and tensor Lorentz structure (with the notation of [62]),

$$\begin{aligned} O_{15}^q &= (\bar{s} P_R b) (\bar{q} P_R q), & O_{16}^q &= (\bar{s}_\alpha P_R b_\beta) (\bar{q}_\beta P_R q_\alpha), \\ O_{17}^q &= (\bar{s} P_R b) (\bar{q} P_L q), & O_{18}^q &= (\bar{s}_\alpha P_R b_\beta) (\bar{q}_\beta P_L q_\alpha), \\ O_{19}^q &= (\bar{s} \sigma^{\mu\nu} P_R b) (\bar{q} \sigma_{\mu\nu} P_R q), & O_{20}^q &= (\bar{s}_\alpha \sigma^{\mu\nu} P_R b_\beta) (\bar{q}_\beta \sigma_{\mu\nu} P_R q_\alpha). \end{aligned} \quad (3.19)$$

The primed operators are obtained by interchanging everywhere $P_L \leftrightarrow P_R$. We recall that in the SM only the vector operators receive contributions, while for the scalar/tensor operator the matching contribution is zero. However, NP is expected to contribute to the Hamiltonian also via scalar/tensor operators. We also note that the operators in (3.16) are redundant since O_1 and O_2 can be obtained from O_{3-6}^q , when $q = c$, via Fierz rearrangements. We will include all NP contributions into the definition of C_{3-6}^q even though for

$q = c$ they could be absorbed in C_1 and C_2 as well. Interestingly, at the leading-logarithmic order only the operators O_{15-20}^q mix into the magnetic and chromomagnetic operators O_7 and O_8 . The vector operators on the other hand mix neither into the magnetic and chromomagnetic nor into the scalar-tensor four-quark operators. The scalar-tensor operators however mix into the vector ones [62].

Four fermion operators that involve two right handed currents ($Q_{dd}, Q_{ud}^{(1)}$, and $Q_{ud}^{(8)}$), give the following contribution to the effective Hamiltonian:

$$C_3'^{q=d,s,b} = -\frac{v^2}{6\Lambda^2} \tilde{C}_{dd}^{1123, 2223, 2333}, \quad C_5'^{q=d,s,b} = \frac{v^2}{24\Lambda^2} \tilde{C}_{dd}^{1123, 2223, 2333}. \quad (3.20)$$

Through a Fierz rearrangement also the operator \tilde{Q}_{dd}^{1321} contributes to

$$C_3'^d = -\frac{v^2}{6\Lambda^2} \frac{1}{N_c} \tilde{C}_{dd}^{1321}, \quad C_5'^d = \frac{v^2}{24\Lambda^2} \frac{1}{N_c} \tilde{C}_{dd}^{1321}, \quad (3.21)$$

$$C_4'^d = -\frac{v^2}{3\Lambda^2} \tilde{C}_{dd}^{1321}, \quad C_6'^d = \frac{v^2}{12\Lambda^2} \tilde{C}_{dd}^{1321}. \quad (3.22)$$

Operators with up-type quarks give:

$$C_3'^{q=u,c} = -\frac{v^2}{6\Lambda^2} \tilde{C}_{ud}^{(1) 1123, 2223}, \quad C_5'^{q=u,c} = \frac{v^2}{24\Lambda^2} \tilde{C}_{ud}^{(1) 1123, 2223}, \quad (3.23)$$

$$C_4'^{q=u,c} = -\frac{v^2}{6\Lambda^2} \tilde{C}_{ud}^{(8) 1123, 2223}, \quad C_6'^{q=u,c} = \frac{v^2}{24\Lambda^2} \tilde{C}_{ud}^{(8) 1123, 2223}. \quad (3.24)$$

In the set $(\bar{L}L)(\bar{R}R)$ in table 1, the operators with right-handed up-type quarks give the following contributions:

$$C_3^{q=u,c} = \frac{2v^2}{3\Lambda^2} \tilde{C}_{qu}^{(1) 2311, 2322}, \quad C_5^{q=u,c} = -\frac{v^2}{24\Lambda^2} \tilde{C}_{qu}^{(1) 2311, 2322}, \quad (3.25)$$

$$C_4^{q=u,c} = \frac{2v^2}{3\Lambda^2} \tilde{C}_{qu}^{(8) 2311, 2322}, \quad C_6^{q=u,c} = -\frac{v^2}{24\Lambda^2} \tilde{C}_{qu}^{(8) 2311, 2322}. \quad (3.26)$$

For the same operator set, but with left-handed up-type quarks, we obtain

$$C_3'^{q=u,c} = \frac{2v^2}{3\Lambda^2} \tilde{C}_{qd}^{(1) 1123, 2223}, \quad C_5'^{q=u,c} = -\frac{v^2}{24\Lambda^2} \tilde{C}_{qd}^{(1) 1123, 2223}, \quad (3.27)$$

$$C_4'^{q=u,c} = \frac{2v^2}{3\Lambda^2} \tilde{C}_{qd}^{(8) 1123, 2223}, \quad C_6'^{q=u,c} = -\frac{v^2}{24\Lambda^2} \tilde{C}_{qd}^{(8) 1123, 2223}, \quad (3.28)$$

where $\tilde{C}_{qd}^{(1,8)ijkl} = V_{im} V_{jn}^* \tilde{C}_{qd}^{(1,8)m n k l}$, as defined in section 2. The operators with four down-type quarks give

$$C_3'^{q=d,s,b} = \frac{2v^2}{3\Lambda^2} \tilde{C}_{qd}^{(1) 1123, 2223, 3323}, \quad C_5'^{q=d,s,b} = -\frac{v^2}{24\Lambda^2} \tilde{C}_{qd}^{(1) 1123, 2223, 3323}, \quad (3.29)$$

$$C_3^{q=d,s,b} = \frac{2v^2}{3\Lambda^2} \tilde{C}_{qd}^{(1) 2311, 2322, 2333}, \quad C_5^{q=d,s,b} = -\frac{v^2}{24\Lambda^2} \tilde{C}_{qd}^{(1) 2311, 2322, 2333}, \quad (3.30)$$

$$C_4'^{q=d,s,b} = \frac{2v^2}{3\Lambda^2} \tilde{C}_{qd}^{(8) 1123, 2223, 3323}, \quad C_6'^{q=d,s,b} = -\frac{v^2}{24\Lambda^2} \tilde{C}_{qd}^{(8) 1123, 2223, 3323}, \quad (3.31)$$

$$C_4^{q=d,s,b} = \frac{2v^2}{3\Lambda^2} \tilde{C}_{qd}^{(8) 2311, 2322, 2333}, \quad C_6^{q=d,s,b} = -\frac{v^2}{24\Lambda^2} \tilde{C}_{qd}^{(8) 2311, 2322, 2333}. \quad (3.32)$$

Let us now investigate the set of four-fermion operators with the Dirac structure $(\bar{L}L)(\bar{L}L)$. Recalling that for this class of operators we consider only those that fulfill $[ij] \leq [kl]$. We obtain the following matching contribution from the vertices involving four left-handed down-type quarks:

$$C_3^{q=s,b} = -\frac{v^2}{6\Lambda^2} \left[\tilde{C}_{qq}^{(1)2223,2333} + \tilde{C}_{qq}^{(3)2223,2333} \right], \quad (3.33)$$

$$C_5^{q=s,b} = +\frac{v^2}{24\Lambda^2} \left[\tilde{C}_{qq}^{(1)2223,2333} + \tilde{C}_{qq}^{(3)2223,2333} \right], \quad (3.34)$$

$$C_3^d = -\frac{v^2}{6\Lambda^2} \left[\tilde{C}_{qq}^{(1)1123} + \tilde{C}_{qq}^{(3)1123} + \frac{1}{N_c} \left(\tilde{C}_{qq}^{(1)1321} + \tilde{C}_{qq}^{(3)1321} \right) \right], \quad (3.35)$$

$$C_5^d = +\frac{v^2}{24\Lambda^2} \left[\tilde{C}_{qq}^{(1)1123} + \tilde{C}_{qq}^{(3)1123} + \frac{1}{N_c} \left(\tilde{C}_{qq}^{(1)1321} + \tilde{C}_{qq}^{(3)1321} \right) \right], \quad (3.36)$$

$$C_4^d = -\frac{v^2}{3\Lambda^2} \left(\tilde{C}_{qq}^{(1)1321} + \tilde{C}_{qq}^{(3)1321} \right), \quad (3.37)$$

$$C_6^d = +\frac{v^2}{12\Lambda^2} \left(\tilde{C}_{qq}^{(1)1321} + \tilde{C}_{qq}^{(3)1321} \right). \quad (3.38)$$

From the operators with two left-handed up-type quarks we obtain

$$C_3^{q=u,c} = -\frac{v^2}{6\Lambda^2} \left(\chi_{u,c}^{(1)} - \chi_{u,c}^{(3)} + \frac{2}{N_c} \Xi_{u,c}^{(3)} \right), \quad (3.39)$$

$$C_5^{q=u,c} = +\frac{v^2}{24\Lambda^2} \left(\chi_{u,c}^{(1)} - \chi_{u,c}^{(3)} + \frac{2}{N_c} \Xi_{u,c}^{(3)} \right), \quad (3.40)$$

$$C_4^{q=u,c} = -\frac{2v^2}{3\Lambda^2} \Xi_{u,c}^{(3)}, \quad (3.41)$$

$$C_6^{q=u,c} = +\frac{v^2}{6\Lambda^2} \Xi_{u,c}^{(3)}, \quad (3.42)$$

where the symbols χ_q and Ξ_q stand for

$$\chi_q^{(1)} = \sum_{[kl]<[23]} \tilde{C}_{qq}^{(1)kl23} V_{qk} V_{ql}^* + \sum_{[kl]>[23]} \tilde{C}_{qq}^{(1)23kl} V_{qk} V_{ql}^* + 2\tilde{C}_{qq}^{(1)2323} V_{qs} V_{qb}^*, \quad (3.43)$$

$$\chi_q^{(3)} = \sum_{[kl]<[23]} \tilde{C}_{qq}^{(3)kl23} V_{qk} V_{ql}^* + \sum_{[kl]>[23]} \tilde{C}_{qq}^{(3)23kl} V_{qk} V_{ql}^* + 2\tilde{C}_{qq}^{(3)2323} V_{qs} V_{qb}^*, \quad (3.44)$$

$$\Xi_q^{(3)} = \sum_{[2j]<[k3]} \tilde{C}_{qq}^{(3)2jk3} V_{qj}^* V_{qk} + \sum_{[j3]<[3k]} \tilde{C}_{qq}^{(3)j32k} V_{qk}^* V_{qj} + 2\tilde{C}_{qq}^{(3)2323} V_{qb}^* V_{qs}. \quad (3.45)$$

Dim-6 operators involving scalar currents generate the following matching contribution for

the operators O_{15-20} in eq. (3.16) involving u or c quarks:

$$C_{15}^{i=u,c} = \frac{v^2}{2\Lambda^2} \left(\tilde{C}_{quqd}^{(1)ii23} - \frac{1}{2N_c} \tilde{C}_{quqd}^{(8)ii23} + \frac{1}{4} V_{ms}^* V_{in} \tilde{C}_{quqd}^{(8)min3} \right), \quad (3.46)$$

$$C'_{15}{}^{i=u,c} = \frac{v^2}{2\Lambda^2} \left(\tilde{C}_{quqd}^{*(1)ii32} - \frac{1}{2N_c} \tilde{C}_{quqd}^{*(8)ii32} + \frac{1}{4} V_{in}^* V_{mb} \tilde{C}_{quqd}^{*(8)min2} \right), \quad (3.47)$$

$$C_{16}^{i=u,c} = \frac{v^2}{4\Lambda^2} \left[\tilde{C}_{quqd}^{(8)ii23} + V_{ms}^* V_{in} \left(\tilde{C}_{quqd}^{(1)min3} - \frac{1}{2N_c} \tilde{C}_{quqd}^{(8)min3} \right) \right], \quad (3.48)$$

$$C'_{16}{}^{i=u,c} = \frac{v^2}{4\Lambda^2} \left[\tilde{C}_{quqd}^{*(8)ii32} + V_{in}^* V_{mb} \left(\tilde{C}_{quqd}^{*(1)min2} - \frac{1}{2N_c} \tilde{C}_{quqd}^{*(8)min2} \right) \right], \quad (3.49)$$

$$C_{19}^{i=u,c} = \frac{v^2}{32\Lambda^2} V_{ms}^* V_{in} \tilde{C}_{quqd}^{(8)min3}, \quad (3.50)$$

$$C'_{19}{}^{i=u,c} = \frac{v^2}{32\Lambda^2} V_{in}^* V_{mb} \tilde{C}_{quqd}^{*(8)min2}, \quad (3.51)$$

$$C_{20}^{i=u,c} = \frac{v^2}{16\Lambda^2} V_{ms}^* V_{in} \left(\tilde{C}_{quqd}^{(1)min3} - \frac{1}{2N_c} \tilde{C}_{quqd}^{(8)min3} \right), \quad (3.52)$$

$$C'_{20}{}^{i=u,c} = \frac{v^2}{16\Lambda^2} V_{in}^* V_{mb} \left(\tilde{C}_{quqd}^{*(1)min2} - \frac{1}{2N_c} \tilde{C}_{quqd}^{*(8)min2} \right), \quad (3.53)$$

The operators $Q_{\varphi q}^{(1)}, Q_{\varphi q}^{(3)}, Q_{\varphi ud}$ and $Q_{\varphi d}$, involving a Z and W coupling with right-handed fermions, contribute to the four-quark operators in eq. (3.10) in the following way:

$$C_3^i = \frac{v^2}{\Lambda^2} \left[\frac{1}{3N_c} \left(T_3^i + \frac{1}{2} \right) \Sigma_{\varphi q}^i - \left(\frac{T_3^i}{3} + Q_i \sin^2 \theta_w \right) \left(\tilde{C}_{\varphi q}^{(1)23} + \tilde{C}_{\varphi q}^{(3)23} \right) \right], \quad (3.54)$$

$$C'_3{}^i = \frac{v^2}{\Lambda^2} \left(\frac{4}{3} T_3^i - Q_i \sin^2 \theta_w \right) \tilde{C}_{\varphi d}^{23}, \quad (3.55)$$

$$C_4^i = \frac{2v^2}{3\Lambda^2} \left(T_3^i + \frac{1}{2} \right) \Sigma_{\varphi q}^i, \quad (3.56)$$

$$C_5^i = \frac{v^2}{\Lambda^2} \left[\frac{T_3^i}{12} \left(\tilde{C}_{\varphi q}^{(1)23} + \tilde{C}_{\varphi q}^{(3)23} \right) - \frac{1}{12N_c} \left(T_3^i + \frac{1}{2} \right) \Sigma_{\varphi q}^i \right], \quad (3.57)$$

$$C'_5{}^i = -\frac{v^2}{\Lambda^2} \frac{T_3^i}{12} \tilde{C}_{\varphi d}^{23}, \quad (3.58)$$

$$C_6^i = -\frac{v^2}{6\Lambda^2} \left(T_3^i + \frac{1}{2} \right) \Sigma_{\varphi q}^i, \quad (3.59)$$

$$C_{18}^i = -\frac{v^2}{\Lambda^2} \left(T_3^i + \frac{1}{2} \right) V_{is}^* \tilde{C}_{\varphi ud}^{i3}, \quad (3.60)$$

$$C'_{18}{}^i = -\frac{v^2}{\Lambda^2} \left(T_3^i + \frac{1}{2} \right) V_{ib} \tilde{C}_{\varphi ud}^{*i2}, \quad (3.61)$$

where $i = u, d, c, s, b$ and Q_i and T_3^i denote its charge and third isospin component, respectively. Moreover we introduced the short notation $\Sigma_{\varphi q}^i = \tilde{C}_{\varphi q}^{(3)j3} V_{ij} V_{is}^* + \tilde{C}_{\varphi q}^{(3)2j} V_{ib} V_{ij}^*$.

The operators involving a vector-current with left-handed quarks directly appear at

tree level in the coefficients for O_9, O_{10} in eq. (3.17):

$$C_9^{ij} = \frac{\pi}{\alpha} \frac{v^2}{\Lambda^2} \left[\tilde{C}_{\ell q}^{(1)ij23} + \tilde{C}_{\ell q}^{(3)ij23} + \tilde{C}_{qe}^{23ij} \right], \quad (3.62)$$

$$C_{10}^{ij} = \frac{\pi}{\alpha} \frac{v^2}{\Lambda^2} \left[\tilde{C}_{qe}^{23ij} - \tilde{C}_{\ell q}^{(1)ij23} - \tilde{C}_{\ell q}^{(3)ij23} \right], \quad (3.63)$$

where the indices $i, j = 1, 2, 3$, corresponding to e, μ and τ . Similar contributions appear for the operators O'_9, O'_{10} from vector-currents involving right-handed quarks:

$$C_9'^{ij} = \frac{\pi}{\alpha} \frac{v^2}{\Lambda^2} \left[\tilde{C}_{\ell d}^{ij23} + \tilde{C}_{ed}^{ij23} \right], \quad (3.64)$$

$$C_{10}'^{ij} = \frac{\pi}{\alpha} \frac{v^2}{\Lambda^2} \left[\tilde{C}_{ed}^{ij23} - \tilde{C}_{\ell d}^{ij23} \right]. \quad (3.65)$$

Scalar operators contribute to the coefficients of O'_P, O'_S :

$$C_S'^{ij} = C_P'^{ij} = \frac{v^2}{4\Lambda^2} \tilde{C}_{\ell edq}^{ij23}. \quad (3.66)$$

Also, for the operators O_P, O_S we have

$$C_S^{ij} = -C_P^{ij} = \frac{v^2}{4\Lambda^2} \tilde{C}_{\ell edq}^{*ij32}, \quad (3.67)$$

where the hermitian conjugate of the operator $Q_{\ell edq}^{ijmn}$ is defined as $\tilde{C}_{\ell edq}^{*ijmn} \left(\bar{e}_R^j \ell_L^i \right) \left(\bar{q}_L^n d_R^m \right)$. These results agree with those in [57] in the case of lepton flavor conservation. Also the operators Q_{dB} and Q_{dW} appear already at tree-level in the effective Hamiltonian through O_7 and O_7' :

$$C_7 = 2\sqrt{2} \sin \theta_w \frac{\pi}{\alpha} \frac{M_W}{m_b} \frac{v^2}{\Lambda^2} \left(\cos \theta_w \tilde{C}_{dB}^{23} - \sin \theta_w \tilde{C}_{dW}^{23} \right), \quad (3.68)$$

$$C_7' = 2\sqrt{2} \sin \theta_w \frac{\pi}{\alpha} \frac{M_W}{m_b} \frac{v^2}{\Lambda^2} \left(\cos \theta_w \tilde{C}_{dB}^{*32} - \sin \theta_w \tilde{C}_{dW}^{*32} \right). \quad (3.69)$$

The operators O_9 and O_{10} , and similarly O'_9 and O'_{10} , receive the following lepton flavor conserving tree-level contribution through the effective \bar{s} - b - Z coupling appearing in the operators $Q_{\varphi d}, Q_{\varphi q}^{(1)}$ and $Q_{\varphi q}^{(3)}$:

$$C_9^{ii} = \frac{\pi}{\alpha} \frac{v^2}{\Lambda^2} \left(\tilde{C}_{\varphi q}^{(1)23} + \tilde{C}_{\varphi q}^{(3)23} \right) (-1 + 4 \sin^2 \theta_w), \quad (3.70)$$

$$C_{10}^{ii} = \frac{\pi}{\alpha} \frac{v^2}{\Lambda^2} \left(\tilde{C}_{\varphi q}^{(1)23} + \tilde{C}_{\varphi q}^{(3)23} \right), \quad (3.71)$$

$$C_9'^{ii} = \frac{\pi}{\alpha} \frac{v^2}{\Lambda^2} \tilde{C}_{\varphi d}^{23} (-1 + 4 \sin^2 \theta_w), \quad (3.72)$$

$$C_{10}'^{ii} = \frac{\pi}{\alpha} \frac{v^2}{\Lambda^2} \tilde{C}_{\varphi d}^{23}. \quad (3.73)$$

The operator Q_{dG} contributes to the Wilson coefficients of \mathcal{O}_8 and \mathcal{O}'_8 in the following way:

$$C_8 = \sqrt{2} \frac{8\pi^2}{g g_s} \frac{M_W}{m_b} \frac{v^2}{\Lambda^2} \tilde{C}_{dG}^{23}, \quad (3.74)$$

$$C'_8 = \sqrt{2} \frac{8\pi^2}{g g_s} \frac{M_W}{m_b} \frac{v^2}{\Lambda^2} \tilde{C}_{dG}^{*32}, \quad (3.75)$$

where g and g_s are the $SU(2)_L$ and $SU(3)_C$ coupling constants, respectively. Interestingly, as already noted in ref. [57], there is no matching contribution to tensor operators at the dimension-six level.

The tree level contribution to the four-quark scalar operators stemming from the operator $Q_{d\varphi}$ is given by

$$C_{15}^b = C_{17}^b = -\frac{M_W m_b}{m_h^2} \frac{\sin \theta_W}{\sqrt{2}e} \frac{v^2}{\Lambda^2} \tilde{C}_{d\varphi}^{23}, \quad (3.76)$$

$$C_{15}'^b = C_{17}'^b = -\frac{M_W m_b}{m_h^2} \frac{\sin \theta_W}{\sqrt{2}e} \frac{v^2}{\Lambda^2} \tilde{C}_{d\varphi}^{*32}. \quad (3.77)$$

4 One-loop matching corrections

In this section we analyze the leading one-loop matching corrections to the $b \rightarrow s$ transitions arising from the dimension-six operators in (1.1). Let us define what we mean by “leading” one-loop matching corrections. First of all, if one of the gauge invariant operators can contribute already at tree-level to $b \rightarrow s$ transitions, a calculation of loop effects is not necessary, since the corresponding Wilson coefficient would already be stringently constrained. Therefore, the loop contribution would only be a subleading effect. With this argument, one can already eliminate all operators that do not contain right-handed up-type quarks: left-handed up quarks always come with their $SU(2)_L$ down quark partner that then contributes to the Hamiltonian at the tree level. Note that it might be possible that an operator containing quark doublets is flavor-violating for up-type quarks but flavor conserving concerning down-type quarks (i.e. not contributing $b \rightarrow s$ transitions due to an alignment in flavor space). However, we do not consider this possibility here and focus on operators with up-quark $SU(2)_L$ singlets. Therefore, we are left with the operators given in table 3.

In the following, we will identify six different classes of matching effects which can be numerically relevant and discuss each of them in a separate subsection. We have the following contributions of gauge invariant dimension-six operators to the ones of the B physics Hamiltonian:

1. 4-fermion operators to 4-fermion operators ($\Delta B = \Delta S = 1$).
2. 4-fermion operators to 4-fermion operators ($\Delta B = \Delta S = 2$).
3. 4-fermion operators to O_7 and O_8 .

$\psi^2 X \varphi$		$(\bar{R}R)(\bar{R}R)$		$(\bar{L}L)(\bar{R}R)$	
Q_{uW}	$(\bar{q}_i \sigma^{\mu\nu} u_j) \tau^I \tilde{\varphi} W_{\mu\nu}^I$	Q_{eu}	$(\bar{e}_i \gamma_\mu e_j)(\bar{u}_k \gamma^\mu u_l)$	$Q_{\ell u}$	$(\bar{\ell}_i \gamma_\mu \ell_j)(\bar{u}_k \gamma^\mu u_l)$
Q_{uB}	$(\bar{q}_i \sigma^{\mu\nu} u_j) \tilde{\varphi} B_{\mu\nu}$	Q_{uu}	$(\bar{u}_i \gamma_\mu u_j)(\bar{u}_k \gamma^\mu u_l)$	$Q_{qu}^{(1)}$	$(\bar{q}_i \gamma_\mu q_j)(\bar{u}_k \gamma^\mu u_l)$
Q_{uG}	$(\bar{q}_i \sigma^{\mu\nu} T^A u_j) \tilde{\varphi} G_{\mu\nu}^A$	$Q_{ud}^{(1)}$	$(\bar{u}_i \gamma_\mu u_j)(\bar{d}_k \gamma^\mu d_l)$	$Q_{qu}^{(8)}$	$(\bar{q}_i \gamma_\mu T^A q_j)(\bar{u}_k \gamma^\mu T^A u_l)$
$\psi^2 \varphi^2 D$		$Q_{ud}^{(8)}$	$(\bar{u}_i \gamma_\mu T^A u_j)(\bar{d}_k \gamma^\mu T^A d_l)$	$(\bar{L}R)(\bar{L}R)$	
$Q_{\varphi ud}$	$(\tilde{\varphi}^\dagger i D_\mu \varphi)(\bar{u}_i \gamma^\mu d_j)$			$Q_{quqd}^{(1)}$	$(\bar{q}_i^a u_j) \varepsilon_{ab} (\bar{q}_k^b d_l)$
$Q_{\varphi u}$	$(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi)(\bar{u}_i \gamma^\mu u_j)$			$Q_{quqd}^{(8)}$	$(\bar{q}_i^a T^A u_j) \varepsilon_{ab} (\bar{q}_k^b T^A d_l)$

Table 3: Dim-6 operators that contribute to $b \rightarrow s$ transitions at the one-loop level.

4. Right-handed Z couplings to O_9 , O_{10} and O_{3-6}^q .
5. Right-handed W couplings to O_7 and O_8 .
6. Magnetic operators to O_7 , O_8 , O_9 , O_{10} and O_4^q .

We perform the matching of the operators in table 3 by integrating out the heavy degrees of freedom represented by the Higgs and the top quark, together with the W and Z bosons. The amplitudes are evaluated at vanishing external momenta, setting all lepton and quark masses to zero except for the top quark mass. To calculate the contribution to the magnetic operators O_7 and O_8 , as well as the photon and gluon penguins, we expanded the amplitudes up to the second order in external momenta and small quark-masses. In order to check our result we performed the calculation in a general R_ξ gauge, and we explicitly verified the cancellation of the ξ dependent part in the final results.

In several cases, the amplitudes have ultraviolet (UV) divergences. Such divergences signal the running and/or the mixing of different gauge invariant operators between the NP scale Λ and the EW scale. The divergences can be (and are) removed via renormalization for which we choose the $\overline{\text{MS}}$ scheme. The residual finite terms constitute in these cases the matching result. To indicate the exact origin of the logarithms, we used the notation $\log(m_t^2/\mu_W^2)$ for the one-loop contributions where only the top quark appears in the loop internal legs, while $\log(M_W^2/\mu_W^2)$ signals the presence of at least one W -boson in the loop.

4.1 Contribution of 4-fermion operators to 4-fermion operators ($\Delta B = \Delta S = 1$)

We start by reporting the matching contribution to the semi-leptonic operators O_9 and O_{10} from four-fermion operators that couple up-type quarks and charged leptons: $Q_{\ell u}$ and Q_{eu} . Obviously, only a charged particle (i.e. the W and the charged Goldstone) can give a contribution to a bs operator which is only possible via a genuine vertex correction. Moreover, the result turns out to be proportional to $m_{u_j}^2$. Therefore, we include only the top-quark contribution while u or c quark effects are vanishing in the massless limit.




Figure 2: One-loop diagrams in unitary gauge contributing to the low energy theory generated by the four-fermion operators in table 3.

Calculating the diagram in figure 2a (and the analogous Goldstone contribution unless one is working in unitary gauge) gives the following matching contributions:

$$C_9^{ij} = \frac{\lambda_t}{\sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left[\tilde{C}_{\ell u}^{ij33} + \tilde{C}_{eu}^{ij33} \right] I(x_t), \quad (4.1)$$

$$C_{10}^{ij} = \frac{\lambda_t}{\sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left[\tilde{C}_{eu}^{ij33} - \tilde{C}_{\ell u}^{ij33} \right] I(x_t), \quad (4.2)$$

where $x_t = m_t^2/M_W^2$ and

$$I(x_t) = \frac{x_t}{16} \left[-\ln \left(\frac{M_W^2}{\mu_W^2} \right) + \frac{x_t - 7}{2(1 - x_t)} - \frac{x_t^2 - 2x_t + 4}{(x_t - 1)^2} \ln(x_t) \right]. \quad (4.3)$$

The four-fermion operators involving only quark fields can also contribute to $C_9^{(\prime)}$ and $C_{10}^{(\prime)}$ through a closed top loop (figure 2c) to which an off-shell Z or photon is attached. In this case the contribution is evidently lepton flavor conserving:

$$C_9^{ii} = \tilde{C}_{qu}^{(1)2333} \frac{v^2}{\Lambda^2} \left(\frac{3x_t}{8\sin^2 \theta_W} - \frac{3x_t}{2} - \frac{2}{3} \right) \ln \left(\frac{m_t^2}{\mu_W^2} \right), \quad (4.4)$$

$$C_9^{\prime ii} = \tilde{C}_{ud}^{(1)3323} \frac{v^2}{\Lambda^2} \left(\frac{3x_t}{8\sin^2 \theta_W} - \frac{3x_t}{2} - \frac{2}{3} \right) \ln \left(\frac{m_t^2}{\mu_W^2} \right), \quad (4.5)$$

$$C_{10}^{ii} = -\tilde{C}_{qu}^{(1)2333} \frac{v^2}{\Lambda^2} \frac{3x_t}{8\sin^2 \theta_W} \ln \left(\frac{m_t^2}{\mu_W^2} \right), \quad (4.6)$$

$$C_{10}^{\prime ii} = -\tilde{C}_{ud}^{(1)3323} \frac{v^2}{\Lambda^2} \frac{3x_t}{8\sin^2 \theta_W} \ln \left(\frac{m_t^2}{\mu_W^2} \right). \quad (4.7)$$

Furthermore, through a W-boson exchange (figure 2b) the operators under discussion give a one-loop matching contribution to $\Delta B = \Delta S = 1$ four-quark operators of the form:

$$C_3^i = \tilde{C}_{qu}^{(1)2333} \frac{\alpha}{4\pi} \frac{v^2}{\Lambda^2} \left\{ \ln \left(\frac{m_t^2}{\mu_W^2} \right) \left[Q_i \left(\frac{3x_t}{2} + \frac{2}{3} \right) + T_3^i \frac{x_t}{2\sin^2 \theta_W} \right] + \frac{2}{3} \left(T_3^i - \frac{1}{2} \right) \frac{|V_{ti}|^2 I(x_t)}{\sin^2 \theta_W} \right\}, \quad (4.8)$$

$$C_3^{\prime i} = \tilde{C}_{ud}^{(1)3323} \frac{\alpha}{4\pi} \frac{v^2}{\Lambda^2} \left\{ \ln \left(\frac{m_t^2}{\mu_W^2} \right) \left[Q_i \left(\frac{3x_t}{2} + \frac{2}{3} \right) - T_3^i \frac{2x_t}{\sin^2 \theta_W} \right] - \frac{8}{3} \left(T_3^i - \frac{1}{2} \right) \frac{|V_{ti}|^2 I(x_t)}{\sin^2 \theta_W} \right\}, \quad (4.9)$$

$$C_4^i = \tilde{C}_{qu}^{(8)2333} \frac{v^2}{\Lambda^2} \left\{ \frac{\alpha_s}{24\pi} \ln \left(\frac{m_t^2}{\mu_W^2} \right) + \frac{\alpha}{6\pi} \left(T_3^i - \frac{1}{2} \right) \frac{|V_{ti}|^2 I(x_t)}{\sin^2 \theta_W} \right\}, \quad (4.10)$$

$$C_4'^i = \tilde{C}_{ud}^{(8)3323} \frac{v^2}{\Lambda^2} \left\{ \frac{\alpha_s}{24\pi} \ln \left(\frac{m_t^2}{\mu_W^2} \right) - \frac{2\alpha}{3\pi} \left(T_3^i - \frac{1}{2} \right) \frac{|V_{ti}|^2 I(x_t)}{\sin^2 \theta_W} \right\}, \quad (4.11)$$

$$C_5^i = -\tilde{C}_{qu}^{(1)2333} \frac{\alpha}{32\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left[T_3^i x_t \ln \left(\frac{m_t^2}{\mu_W^2} \right) + \frac{4}{3} \left(T_3^i - \frac{1}{2} \right) |V_{ti}|^2 I(x_t) \right], \quad (4.12)$$

$$C_5'^i = +\tilde{C}_{ud}^{(1)3323} \frac{\alpha}{32\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left[T_3^i x_t \ln \left(\frac{m_t^2}{\mu_W^2} \right) + \frac{4}{3} \left(T_3^i - \frac{1}{2} \right) |V_{ti}|^2 I(x_t) \right], \quad (4.13)$$

$$C_6^i = \tilde{C}_{qu}^{(8)2333} \frac{\alpha}{24\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left(\frac{1}{2} - T_3^i \right) |V_{ti}|^2 I(x_t), \quad (4.14)$$

$$C_6'^i = \tilde{C}_{ud}^{(8)3323} \frac{\alpha}{24\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left(T_3^i - \frac{1}{2} \right) |V_{ti}|^2 I(x_t), \quad (4.15)$$

where Q_i is the charge of the quark, $T_3^i = 1/2$ for $q = u, c$ and $T_3^i = -1/2$ for $q = d, s, b$. Four-fermion operators not containing the flavor violating current $\bar{s}b$ contribute to the four-quarks operators in (3.16) in the following way:

$$C_3^{i=d,s,b} = \lambda_t \frac{\alpha}{6\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left[4 \tilde{C}_{ud}^{(1)33ii} - \tilde{C}_{qu}^{(1)ii33} \right] I(x_t), \quad (4.16)$$

$$C_3^{i=u,c} = \lambda_t \frac{\alpha}{6\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left[4 \left(\tilde{C}_{uu}^{ii33} - \frac{1}{N_c} \tilde{C}_{uu}^{i33i} \right) - \tilde{C}_{qu}^{(1)ii33} \right] I(x_t), \quad (4.17)$$

$$C_4^{i=d,s,b} = \lambda_t \frac{\alpha}{6\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left[4 \tilde{C}_{ud}^{(8)33ii} - \tilde{C}_{qu}^{(8)ii33} \right] I(x_t), \quad (4.18)$$

$$C_4^{i=u,c} = \lambda_t \frac{\alpha}{6\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left[-8 \tilde{C}_{uu}^{i33i} - \tilde{C}_{qu}^{(8)ii33} \right] I(x_t), \quad (4.19)$$

$$C_5^{i=d,s,b} = \lambda_t \frac{\alpha}{24\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left[\tilde{C}_{qu}^{(1)ii33} - \tilde{C}_{ud}^{(1)33ii} \right] I(x_t), \quad (4.20)$$

$$C_5^{i=u,c} = \lambda_t \frac{\alpha}{24\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left[\tilde{C}_{qu}^{(1)ii33} - \tilde{C}_{uu}^{ii33} + \frac{1}{N_c} \tilde{C}_{uu}^{i33i} \right] I(x_t), \quad (4.21)$$

$$C_6^{i=d,s,b} = \lambda_t \frac{\alpha}{24\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left[\tilde{C}_{qu}^{(8)ii33} - \tilde{C}_{ud}^{(8)33ii} \right] I(x_t), \quad (4.22)$$

$$C_6^{i=u,c} = \lambda_t \frac{\alpha}{24\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} \left[\tilde{C}_{qu}^{(8)ii33} + 2 \tilde{C}_{uu}^{i33i} \right] I(x_t), \quad (4.23)$$

where here we used also the notation introduced in section 2: $\tilde{C}_{qu}^{(1,8)ijkl} = V_{im} V_{jn}^* \tilde{C}_{qu}^{(1,8)mnlk}$.

4.2 Contribution of 4-fermion operators to 4-fermion operators ($\Delta B = \Delta S = 2$)

The Hamiltonian for B_s - \bar{B}_s mixing in eq. (3.1) gets a one-loop matching contribution through the graph in figure 2b:

$$C_1 = \lambda_t \frac{\alpha}{2\pi \sin^2 \theta_W} \frac{1}{\Lambda^2} I(x_t) \left[\left(-1 + \frac{1}{N_c} \right) \tilde{C}_{qu}^{(8) 2333} - 2\tilde{C}_{qu}^{(1) 2333} \right], \quad (4.24)$$

$$C_4 = \lambda_t \frac{\alpha}{\pi \sin^2 \theta_W} \frac{1}{\Lambda^2} I(x_t) \tilde{C}_{ud}^{(8) 3323}, \quad (4.25)$$

$$C_5 = \lambda_t \frac{\alpha}{\pi \sin^2 \theta_W} \frac{1}{\Lambda^2} I(x_t) \left[2\tilde{C}_{ud}^{(1) 3323} - \frac{1}{N_c} \tilde{C}_{ud}^{(8) 3323} \right]. \quad (4.26)$$

4.3 Contributions of 4-fermion operators to O_7 and O_8

Four-fermion operators with scalar currents contribute to the low energy Hamiltonian (3.16) through the diagram in figure 2d:

$$C_7 = -\frac{1}{6} \frac{m_t}{m_b} \frac{v^2}{\Lambda^2} \ln \left(\frac{m_t^2}{\mu_W^2} \right) \left[\tilde{C}_{quqd}^{(1) 2333} + C_F \tilde{C}_{quqd}^{(8) 2333} \right], \quad (4.27)$$

$$C_7' = -\frac{1}{6} \frac{m_t}{m_b} \frac{v^2}{\Lambda^2} \ln \left(\frac{m_t^2}{\mu_W^2} \right) \left[\tilde{C}_{quqd}^{*(1) 3332} + C_F \tilde{C}_{quqd}^{*(8) 3332} \right], \quad (4.28)$$

$$C_8 = -\frac{1}{4} \frac{m_t}{m_b} \frac{v^2}{\Lambda^2} \ln \left(\frac{m_t^2}{\mu_W^2} \right) \left[\tilde{C}_{quqd}^{(1) 2333} - \frac{1}{2N_c} \tilde{C}_{quqd}^{(8) 2333} \right], \quad (4.29)$$

$$C_8' = -\frac{1}{4} \frac{m_t}{m_b} \frac{v^2}{\Lambda^2} \ln \left(\frac{m_t^2}{\mu_W^2} \right) \left[\tilde{C}_{quqd}^{*(1) 3332} - \frac{1}{2N_c} \tilde{C}_{quqd}^{*(8) 3332} \right], \quad (4.30)$$

where $C_F = (N_c^2 - 1)/(2N_c)$. Note that the contribution to C_7 or C_8 from 4-fermion operators involving vector currents vanishes (excluding QCD corrections).

4.4 Contributions of right-handed Z couplings to O_9 , O_{10} and O_{3-6}^q

The operator $Q_{\varphi u}$, involving only right-handed up-type quarks, gives through a Z -penguin (figure 3f) a matching contribution to the $\Delta B = \Delta S = 1$ Hamiltonian in eq. (3.16) of the form:

$$C_3^i = -\lambda_t \frac{\alpha}{\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} I(x_t) \tilde{C}_{\varphi u}^{33} \left(Q_i \sin^2 \theta_W + \frac{1}{3} T_3^i \right), \quad (4.31)$$

$$C_5^i = \lambda_t \frac{\alpha}{12\pi \sin^2 \theta_W} \frac{v^2}{\Lambda^2} I(x_t) \tilde{C}_{\varphi u}^{33} T_3^i, \quad (4.32)$$

$$C_9^{ii} = \frac{\lambda_t}{\sin^2 \theta_W} \frac{v^2}{\Lambda^2} \tilde{C}_{\varphi u}^{33} I(x_t) (-1 + 4 \sin^2 \theta_W), \quad (4.33)$$

$$C_{10}^{ii} = \frac{\lambda_t}{\sin^2 \theta_W} \frac{v^2}{\Lambda^2} \tilde{C}_{\varphi u}^{33} I(x_t), \quad (4.34)$$

where $I(x_t)$ has been defined in eq. (4.3). The possibility to probe the anomalous couplings of the Z boson to top quark with rare meson decays were also studied in [81].




Figure 3: One-loop diagrams in the unitary gauge for $b \rightarrow sV$ transitions (with $V = Z, \gamma, g$) originating from the operators Q_{uB}, Q_{uW} and Q_{uG} . The red dots represent an operator insertion. For each of these diagrams a symmetric one must also be considered, with the effective operator in the W - t - b vertex. Box diagrams and self energies on the external legs (not depicted here) must also be included.

4.5 Contributions of right-handed W couplings to O_7 and O_8

The operator $Q_{\varphi ud}$ couples the W boson to right-handed quarks, which induces a non-zero contribution only to the magnetic terms O_7, O_8 :

$$C_7 = \frac{m_t}{m_b} \frac{v^2}{\Lambda^2} E_{\varphi ud}^7(x_t) \tilde{C}_{\varphi ud}^{33} V_{ts}^*, \quad (4.35)$$

$$C_7' = \frac{m_t}{m_b} \frac{v^2}{\Lambda^2} E_{\varphi ud}^7(x_t) \tilde{C}_{\varphi ud}^{*32} V_{tb}, \quad (4.36)$$

$$C_8 = \frac{m_t}{m_b} \frac{v^2}{\Lambda^2} E_{\varphi ud}^8(x_t) \tilde{C}_{\varphi ud}^{33} V_{ts}^*, \quad (4.37)$$

$$C_8' = \frac{m_t}{m_b} \frac{v^2}{\Lambda^2} E_{\varphi ud}^8(x_t) \tilde{C}_{\varphi ud}^{*32} V_{tb}, \quad (4.38)$$

where the x_t -functions, in agreement with [82, 83], are

$$E_{\varphi ud}^7(x_t) = \frac{-5x_t^2 + 31x_t - 20}{24(x_t - 1)^2} + \frac{x_t(2 - 3x_t)}{4(x_t - 1)^3} \ln(x_t), \quad (4.39)$$

$$E_{\varphi ud}^8(x_t) = -\frac{x_t^2 + x_t + 4}{8(x_t - 1)^2} + \frac{3x_t}{4(x_t - 1)^3} \ln(x_t). \quad (4.40)$$

4.6 Contributions of magnetic operators to O_7, O_8, O_9, O_{10} and O_4^q

In this subsection we summarize the matching contributions arising from the magnetic operators in table 3. The operators Q_{uB} and Q_{uW} contribute to the effective Hamiltonian for $b \rightarrow s\gamma$ and $b \rightarrow s\bar{\ell}\ell$ transitions via the one-loop diagrams in figure 3.

For simplicity, let us first consider the operators \tilde{C}_{uW}^{33} and \tilde{C}_{uB}^{33} that generate an extra term for the top anomalous magnetic moment resulting in a chirality flipping vertex with the W boson. We will later analyse the case when the vertices with the photon and the Z are flavor violating. Here we include only the contributions to four-quark operators arising from gluon-penguin diagrams, which are of $O(\alpha_s)$, and we neglect the subleading EW penguin diagrams, of $O(\alpha)$. We obtained the following contributions to the effective Hamiltonian in eq. (3.16):

$$C_4^i = \lambda_t \frac{\alpha_s}{\pi} \frac{m_t}{M_W} \frac{\sqrt{2} v^2}{\Lambda^2} A_{uW}(x_t) \text{Re}(\tilde{C}_{uW}^{33}), \quad (4.41)$$

$$C_7 = \lambda_t \frac{m_t}{M_W} \frac{\sqrt{2} v^2}{\Lambda^2} \left\{ \tilde{C}_{uW}^{33} E_{uW}^7(x_t) + \tilde{C}_{uW}^{*33} F_{uW}^7(x_t) + \frac{\cos \theta_W}{\sin \theta_W} \left[\tilde{C}_{uB}^{33} E_{uB}^7(x_t) + \tilde{C}_{uB}^{*33} F_{uB}^7(x_t) \right] \right\}, \quad (4.42)$$

$$C_8 = \lambda_t \frac{m_t}{M_W} \frac{\sqrt{2} v^2}{\Lambda^2} \left[\tilde{C}_{uW}^{33} E_{uW}^8(x_t) + \tilde{C}_{uW}^{*33} F_{uW}^8(x_t) \right], \quad (4.43)$$

$$C_9^{ii} = \lambda_t \frac{m_t}{M_W} \frac{\sqrt{2} v^2}{\Lambda^2} \left[\text{Re}(\tilde{C}_{uW}^{33}) \left(\frac{Y_{uW}(x_t)}{\sin^2 \theta_W} - Z_{uW}(x_t) \right) - \frac{\cos \theta_W}{\sin \theta_W} \text{Re}(\tilde{C}_{uB}^{33}) Z_{uB}(x_t) \right], \quad (4.44)$$

$$C_{10}^{ii} = -\lambda_t \frac{m_t}{M_W} \frac{\sqrt{2} v^2}{\Lambda^2} \frac{Y_{uW}(x_t)}{\sin^2 \theta_W} \text{Re}(\tilde{C}_{uW}^{33}), \quad (4.45)$$

where the explicit expressions for the x_t -dependent functions are

$$E_{uW}^7(x_t) = \frac{1}{8} \ln \left(\frac{M_W^2}{\mu_W^2} \right) + \frac{-9x_t^3 + 63x_t^2 - 61x_t + 19}{48(x_t - 1)^3} + \frac{3x_t^4 - 12x_t^3 - 9x_t^2 + 20x_t - 8}{24(x_t - 1)^4} \ln(x_t), \quad (4.46)$$

$$F_{uW}^7(x_t) = -\frac{3x_t^3 - 17x_t^2 + 4x_t + 4}{24(x_t - 1)^3} + \frac{x_t(2 - 3x_t)}{4(x_t - 1)^4} \ln(x_t), \quad (4.47)$$

$$E_{uB}^7(x_t) = -\frac{1}{8} \ln \left(\frac{M_W^2}{\mu_W^2} \right) - \frac{(x_t + 1)^2}{16(x_t - 1)^2} - \frac{x_t^2(x_t - 3)}{8(x_t - 1)^3} \ln(x_t), \quad (4.48)$$

$$F_{uB}^7(x_t) = -\frac{1}{8}, \quad (4.49)$$

$$E_{uW}^8(x_t) = \frac{3x_t^2 - 13x_t + 4}{8(x_t - 1)^3} + \frac{5x_t - 2}{4(x_t - 1)^4} \ln(x_t), \quad (4.50)$$

$$F_{uW}^8(x_t) = \frac{x_t^2 - 5x_t - 2}{8(x_t - 1)^3} + \frac{3x_t}{4(x_t - 1)^4} \ln(x_t), \quad (4.51)$$

$$A_{uW}(x_t) = \frac{5x_t^2 - 19x_t + 20}{24(x_t - 1)^3} + \frac{x_t - 2}{4(x_t - 1)^4} \ln(x_t), \quad (4.52)$$

$$Y_{uW}(x_t) = \frac{3x_t}{4(x_t - 1)} - \frac{3x_t}{4(x_t - 1)^2} \ln(x_t), \quad (4.53)$$

$$Z_{uW}(x_t) = \frac{99x_t^3 - 136x_t^2 - 25x_t + 50}{36(x_t - 1)^3} - \frac{24x_t^3 - 45x_t^2 + 17x_t + 2}{6(x_t - 1)^4} \ln(x_t), \quad (4.54)$$

$$Z_{uB}(x_t) = -\frac{x_t^2 + 3x_t - 2}{4(x_t - 1)^2} + \frac{3x_t - 2}{2(x_t - 1)^3} \ln(x_t). \quad (4.55)$$

We found that the expressions for the functions $E_{uW}^i, F_{uW}^i, Y_{uW}$ and Z_{uW} are in agreement with the results reported in [82, 83], while A_{uW}, Z_{uB}, E_{uB}^7 and F_{uB}^7 are new to the best of our knowledge. Note that the effect on the magnetic operators O_7 and O_8 is divergent while it is finite for the four-fermion operators. Moreover, all these effects scale like $1/\Lambda^2$ and do not possess an additional suppression by $1/M_W^2$.

Now we turn our attention to the operators Q_{uW}^{i3} and Q_{uB}^{i3} , where $i = 1, 2$.⁵ These operators lead to an anomalous W - t - d^i coupling, plus two flavor-violating neutral currents $(Z/\gamma)tc$ and $(Z/\gamma)tu$, so then in the diagram 3b one top quark propagator becomes $q = u, c$. However, we recall that this amplitude is non-zero only for the γ penguin, or the transition $b \rightarrow s\gamma$ — the effective coupling is proportional to $\sigma^{\mu\nu}q_\nu$, where q is the momentum of the boson. Only the functions arising from a γ penguin will be modified in this case, i.e. the functions Z, E^7, F^7 . Repeating the calculations performed for \tilde{C}_{uB}^{33} and \tilde{C}_{uW}^{33} we obtain the following results for the matching:

$$C_4^i = \frac{\alpha_s}{\pi} \frac{m_t}{M_W} \frac{\sqrt{2}v^2}{\Lambda^2} A_{uW}(x_t) \Sigma_{uW}, \quad (4.56)$$

$$C_7 = \frac{m_t}{M_W} \frac{\sqrt{2}v^2}{\Lambda^2} \left\{ \tilde{C}_{uW}^{i3} V_{is}^* V_{tb} E_{uW}'^7(x_t) + \tilde{C}_{uW}^{*i3} V_{ib} V_{ts}^* F_{uW}^7(x_t) \right. \\ \left. + \frac{\cos \theta_W}{\sin \theta_W} \left[\tilde{C}_{uB}^{i3} V_{is}^* V_{tb} E_{uB}'^7(x_t) + \tilde{C}_{uB}^{*i3} V_{ib} V_{ts}^* F_{uB}^7(x_t) \right] \right\}, \quad (4.57)$$

$$C_8 = \frac{m_t}{M_W} \frac{\sqrt{2}v^2}{\Lambda^2} \left[\tilde{C}_{uW}^{i3} V_{is}^* V_{tb} E_{uW}^8(x_t) + \tilde{C}_{uW}^{*i3} V_{ib} V_{ts}^* F_{uW}^8(x_t) \right], \quad (4.58)$$

$$C_9^{ii} = \frac{m_t}{M_W} \frac{\sqrt{2}v^2}{\Lambda^2} \left[\Sigma_{uW} \left(\frac{Y_{uW}(x_t)}{\sin^2 \theta_W} - Z'_{uW}(x_t) \right) - \frac{\cos \theta_W}{\sin \theta_W} \Sigma_{uB} Z'_{uB}(x_t) \right], \quad (4.59)$$

$$C_{10}^{ii} = -\frac{m_t}{M_W} \frac{\sqrt{2}v^2}{\Lambda^2} \frac{Y_{uW}(x_t)}{\sin^2 \theta_W} \Sigma_{uW}, \quad (4.60)$$

where $\Sigma_{uW} = (\tilde{C}_{uW}^{i3} V_{is}^* V_{tb} + \tilde{C}_{uW}^{*i3} V_{ib} V_{ts}^*)/2$ and $\Sigma_{uB} = (\tilde{C}_{uB}^{i3} V_{is}^* V_{tb} + \tilde{C}_{uB}^{*i3} V_{ib} V_{ts}^*)/2$ (the summation over $i = 1, 2$ is implied). The new functions introduced above are:

$$Z'_{uW}(x_t) = \frac{54x_t^3 - 59x_t^2 - 35x_t + 34}{18(x_t - 1)^3} - \frac{15x_t^3 - 27x_t^2 + 10x_t + 1}{3(x_t - 1)^4} \ln(x_t), \quad (4.61)$$

$$Z'_{uB}(x_t) = \frac{1}{1 - x_t} \ln(x_t), \quad (4.62)$$

$$E_{uW}'^7(x_t) = \frac{1}{8} \ln \left(\frac{M_W^2}{\mu_W^2} \right) + \frac{-3x_t^3 + 63x_t^2 - 67x_t + 19}{48(x_t - 1)^3} + \frac{3x_t^4 - 18x_t^3 - 3x_t^2 + 20x_t - 8}{24(x_t - 1)^4} \ln(x_t), \quad (4.63)$$

$$E_{uB}'^7(x_t) = -\frac{1}{8} \ln \left(\frac{M_W^2}{\mu_W^2} \right) + \frac{x_t + 1}{16(x_t - 1)} - \frac{x_t^2}{8(x_t - 1)^2} \ln(x_t). \quad (4.64)$$

The operator Q_{uG}^{33} gives a chromo-magnetic coupling with the top quark, that contributes at one-loop to O_8 and O_4 through the gluon-penguin diagrams in figure 3b,3f.

⁵The effect of a right-handed W - t - d coupling on $b \rightarrow d\gamma$ was studied in ref. [84].

The explicit matching contributions are

$$C_4^i = \lambda_t \frac{gg_s}{16\pi^2} \frac{m_t}{M_W} \frac{\sqrt{2}v^2}{\Lambda^2} \text{Re}(\tilde{C}_{uG}^{33}) A_{uG}(x_t), \quad (4.65)$$

$$C_8 = \lambda_t \frac{g}{g_s} \frac{m_t}{M_W} \frac{\sqrt{2}v^2}{\Lambda^2} \left[\tilde{C}_{uG}^{33} E_{uG}^8(x_t) + \tilde{C}_{uG}^{*33} F_{uG}^8(x_t) \right], \quad (4.66)$$

where $A_{uG} = Z_{uB}$, $E_{uG}^8 = E_{uB}^7$ and $F_{uG}^8 = F_{uB}^7$. Moreover, the operators Q_{uG}^{i3} lead to a flavor violating neutral current involving a gluon and up-type quarks, whose effects in the effective Hamiltonian are

$$C_4^i = \frac{gg_s}{16\pi^2} \frac{m_t}{M_W} \frac{\sqrt{2}v^2}{\Lambda^2} A'_{uG}(x_t) \frac{\tilde{C}_{uG}^{i3} V_{tb} V_{is}^* + \tilde{C}_{uG}^{*i3} V_{ib} V_{ts}^*}{2}, \quad (4.67)$$

$$C_8 = \frac{g}{g_s} \frac{m_t}{M_W} \frac{\sqrt{2}v^2}{\Lambda^2} \left[\tilde{C}_{uG}^{i3} V_{tb} V_{is}^* E'_{uG}(x_t) + \tilde{C}_{uG}^{*i3} V_{ib} V_{ts}^* F_{uG}^8(x_t) \right], \quad (4.68)$$

where $A'_{uG} = Z'_{uB}$ and $E'_{uG}^8 = E'_{uB}^7$.

5 Phenomenological example

As an example of phenomenological applications of the matching conditions reported in sections 3 and 4, we will consider the operator $\tilde{Q}_{\varphi ud}^{33}$ that gives rise to a one-loop contribution to C_7 and C_8 (see eqs.(4.35) and (4.37)). We can employ the inclusive $\bar{B} \rightarrow X_s \gamma$ branching ratio to constrain the the Wilson coefficient $\tilde{C}_{\varphi ud}^{33}$. Let us denote the Wilson coefficients in (3.16) as $C_i(\mu) = C_i^{\text{SM}}(\mu) + \Delta C_i(\mu)$, where $\Delta C_i(\mu)$ are possible non-SM terms. The calculation of the contribution to the decay $\bar{B} \rightarrow X_s \gamma$ proceeds precisely as in the SM case:

- The evolution of the Wilson coefficients in (3.16), from the mass scale $\mu = \mu_W$ down to $\mu = \mu_b$, where μ_b is of the order of m_b , by solving the appropriate RGE.
- The evaluation of the corrections to the matrix elements $\langle s\gamma | O_i(\mu) | b \rangle$ at the scale $\mu = \mu_b$, and the subsequent shift induced in the branching ratio $\mathcal{B}(\bar{B} \rightarrow X_s \gamma)$.

The beyond-SM effect on $\mathcal{B}(\bar{B} \rightarrow X_s \gamma)$, driven by the new additive contribution involving $\Delta C_{7,8}$, can be compactly written as

$$\mathcal{B}^{\text{th}}(\bar{B} \rightarrow X_s \gamma) \times 10^4 = (3.36 \pm 0.23) - \frac{8.22 \Delta C_7 + 1.99 \Delta C_8}{V_{tb} V_{ts}^*}, \quad (5.1)$$

where $\Delta C_{7,8}$ are defined at the mass scale $\mu_W = 160$ GeV [85]. The theoretical prediction (5.1) incorporates NNLO QCD corrections as well as nonperturbative effects. The input parameters and their uncertainties can be found in Appendix D of ref. [86]. Moreover it is assumed that the quadratic terms in $\Delta C_{7,8}$ are negligible.

For the purpose of this example we assume $\tilde{C}_{\varphi ud}^{33}$ to be real and we neglect the imaginary part of V_{tb} and V_{ts} . Identifying the non-SM terms $\Delta C_{7,8}$ with the results in eqs.(4.35) and (4.37), and taking into account the current world average [10]

$$\mathcal{B}^{\text{exp}}(\bar{B} \rightarrow X_s \gamma) = (3.43 \pm 0.21 \pm 0.07) \times 10^{-4}, \quad (5.2)$$

we can find the current 95% C.L. bounds

$$-3.3 \times 10^{-3} \leq \tilde{C}_{\varphi ud}^{33} [\mu_W = 160 \text{ GeV}] \frac{v^2}{\Lambda^2} \leq 2.7 \times 10^{-3}. \quad (5.3)$$

This quite strong bound takes place due to a relative enhancement m_t/m_b compared to the SM case: the SM chiral suppression factor m_b/M_W is replaced by the factor m_t/M_W [87]. It can be interesting to compare (5.3) with the Wtb vertex structure searches at the LHC. The 8 TeV data on the single top quark production cross section and the measurements of the W -boson helicity fractions allowed the authors of [88] to set a bound on $\tilde{C}_{\varphi ud}^{33} (v^2/\Lambda^2)$ at the level of 10^{-1} . Also, ATLAS searches for anomalous couplings in the Wtb vertex from the measurement of double differential angular decay rates of single top quarks produced in the t -channel show similar sensitivities [89].

6 Conclusions

In this article, we calculated (at the EW scale) the matching of the gauge invariant dimension-six operators on the B physics Hamiltonian (including lepton flavor violating operators) integrating out the top, W , Z and the Higgs. After performing the EW symmetry breaking and diagonalizing the mass matrices, we first presented the complete tree-level matching coefficients for $b \rightarrow s$ and $b \rightarrow c$ transitions. Operators involving top quarks do not contribute to $b \rightarrow s$ processes at the tree level, as the top is not a dynamical degree of freedom of the B physics Hamiltonian. Therefore, we identified all operators involving right-handed top quarks which can give numerically important contributions at the one loop-level:

1. 4-fermion operators to 4-fermion operators ($\Delta B = \Delta S = 1$).
2. 4-fermion operators to 4-fermion operators ($\Delta B = \Delta S = 2$).
3. 4-fermion operators to O_7 and O_8 .
4. Right-handed Z couplings to O_9 , O_{10} and O_{3-6}^q .
5. Right-handed W couplings to O_7 and O_8 .
6. Magnetic operators to O_7 , O_8 , O_9 , O_{10} and O_4^q .

Once the necessary running between the EW scale and the B meson scale is performed, our results can be used systematically to test the sensitivity of B physics observables on the dimension six operators.

Acknowledgments

We thank M. González-Alonso, U. Haisch and D. Straub for useful correspondence. We also thank K. Teppe for confirming missing factors of 2 in eqs. (4.24-4.26) in earlier versions and the journal version. J.A., M.F. and C.G. acknowledge the support from the Swiss National Science Foundation. A.C. is supported by a Marie Curie Intra-European Fellowship of the European Community's 7th Framework Programme under contract number PIEF-GA-2012-326948.

A Dimension-six operators in the mass basis

Here we explicitly relate the Wilson coefficients of the gauge invariant operators in the interaction basis to the mass basis. This translation is necessary, if the results obtained in this article have to be related to a UV complete model, where the interaction basis is specified. For the notation, we refer the reader to the original paper in ref. [7].

Table 4: Operators with quarks, gauge and/or Higgs bosons

Operator	Definition in the mass basis
Q_{dB}	$\tilde{C}_{dB}^{ij} \left[V_{ki} \bar{u}_L^k \sigma^{\mu\nu} d_R^j \varphi^+ + \bar{d}_L^i \sigma^{\mu\nu} d_R^j \left(\frac{v+h+i\varphi^0}{\sqrt{2}} \right) \right] B_{\mu\nu}$ $\tilde{C}_{dB}^{ij} = C_{dB}^{mn} S_{Lim}^{d\dagger} S_{Rnj}^d$
Q_{dW}	$\tilde{C}_{dW}^{ij} \left[V_{ki} \bar{u}_L^k \sigma^{\mu\nu} d_R^j \varphi^+ - \bar{d}_L^i \sigma^{\mu\nu} d_R^j \left(\frac{v+h+i\varphi^0}{\sqrt{2}} \right) \right] W_{\mu\nu}^3 + \dots$ $\tilde{C}_{dW}^{ij} = C_{dW}^{mn} S_{Lim}^{d\dagger} S_{Rnj}^d$
Q_{uB}	$\tilde{C}_{uB}^{ij} \left[\bar{u}_L^i \sigma^{\mu\nu} u_R^j \left(\frac{v+h-i\varphi^0}{\sqrt{2}} \right) - V_{ik}^* \bar{d}_L^k \sigma^{\mu\nu} u_R^j \varphi^- \right] B_{\mu\nu}$ $\tilde{C}_{uB}^{ij} = C_{uB}^{mn} S_{Lim}^{u\dagger} S_{Rnj}^u$
Q_{uW}	$\tilde{C}_{uW}^{ij} \left[\bar{u}_L^i \sigma^{\mu\nu} u_R^j \left(\frac{v+h-i\varphi^0}{\sqrt{2}} \right) + V_{ik}^* \bar{d}_L^k \sigma^{\mu\nu} u_R^j \varphi^- \right] W_{\mu\nu}^3 + \dots$ $\tilde{C}_{uW}^{ij} = C_{uW}^{mn} S_{Lim}^{u\dagger} S_{Rnj}^u$
Q_{dG}	$\tilde{C}_{dG}^{ij} \left[V_{ki} \bar{u}_L^k \sigma^{\mu\nu} T^A d_R^j \varphi^+ + \bar{d}_L^i \sigma^{\mu\nu} T^A d_R^j \left(\frac{v+h+i\varphi^0}{\sqrt{2}} \right) \right] G_{\mu\nu}^A$ $\tilde{C}_{dG}^{ij} = C_{dG}^{mn} S_{Lim}^{d\dagger} S_{Rnj}^d$
Q_{uG}	$\tilde{C}_{uG}^{ij} \left[\bar{u}_L^i \sigma^{\mu\nu} T^A u_R^j \left(\frac{v+h-i\varphi^0}{\sqrt{2}} \right) - V_{ik}^* \bar{d}_L^k \sigma^{\mu\nu} T^A u_R^j \varphi^- \right] G_{\mu\nu}^A$ $\tilde{C}_{uG}^{ij} = C_{uG}^{mn} S_{Lim}^{u\dagger} S_{Rnj}^u$
$Q_{\varphi q}^{(1)}$	$\tilde{C}_{\varphi q}^{(1)ij} \left(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) \left(V_{mi} V_{nj}^* \bar{u}_L^m \gamma^\mu u_L^n + \bar{d}_L^i \gamma^\mu d_L^j \right)$ $\tilde{C}_{\varphi q}^{(1)ij} = C_{\varphi q}^{(1)mn} S_{Lim}^{d\dagger} S_{Lnj}^d$
$Q_{\varphi q}^{(3)}$	$\tilde{C}_{\varphi q}^{(3)ij} \left(\varphi^\dagger i \overleftrightarrow{D}_\mu^1 \varphi \right) \left(V_{mi} \bar{u}_L^m \gamma^\mu d_L^j + V_{nj}^* \bar{d}_L^i \gamma^\mu u_L^n \right) + \dots$ $\tilde{C}_{\varphi q}^{(3)ij} = C_{\varphi q}^{(3)mn} S_{Lim}^{d\dagger} S_{Lnj}^d$
$Q_{\varphi d}$	$\tilde{C}_{\varphi d}^{ij} \left(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) \left(\bar{d}_R^i \gamma^\mu d_R^j \right)$ $\tilde{C}_{\varphi d}^{ij} = C_{\varphi d}^{mn} S_{Rim}^{d\dagger} S_{Rnj}^d$

Operator	Definition in the mass basis
$Q_{\varphi u}$	$\tilde{C}_{\varphi u}^{ij} \left(\varphi^\dagger i \overleftrightarrow{D}_\mu \varphi \right) \left(\bar{u}_R^i \gamma^\mu u_R^j \right)$ $\tilde{C}_{\varphi u}^{ij} = C_{\varphi u}^{mn} S_{Rim}^{u\dagger} S_{Rnj}^u$
$Q_{\varphi ud}$	$\tilde{C}_{\varphi ud}^{ij} \left(\varphi^\dagger i D_\mu \varphi \right) \left(\bar{u}_R^i \gamma^\mu d_R^j \right)$ $\tilde{C}_{\varphi ud}^{ij} = C_{\varphi ud}^{mn} S_{Rim}^{u\dagger} S_{Rnj}^d$
$Q_{d\varphi}$	$\tilde{C}_{d\varphi}^{ij} \left(\varphi^\dagger \varphi \right) \left[V_{mi} \bar{u}_L^m d_R^j \varphi^+ + \bar{d}_L^i d_R^j \left(\frac{v+h+i\varphi^0}{\sqrt{2}} \right) \right]$ $\tilde{C}_{d\varphi}^{ij} = C_{d\varphi}^{mn} S_{Lim}^{d\dagger} S_{Rnj}^d$

Table 5: Four-fermion operators with four quarks

Operator	Definition in the mass basis
$Q_{qq}^{(1)}$	$\tilde{C}_{qq}^{(1)ijkl} \left(V_{mi} V_{nj}^* \bar{u}_L^m \gamma^\mu u_L^n + \bar{d}_L^i \gamma^\mu d_L^j \right) \left(V_{mk} V_{nl}^* \bar{u}_L^m \gamma_\mu u_L^n + \bar{d}_L^k \gamma_\mu d_L^l \right)$ $\tilde{C}_{qq}^{(1)ijkl} = C_{qq}^{(1)pqrs} S_{Lip}^{d\dagger} S_{Lqj}^d S_{Lkr}^{d\dagger} S_{Lsl}^d$
$Q_{qq}^{(3)}$	$\tilde{C}_{qq}^{(3)ijkl} \left(V_{mi} V_{nj}^* \bar{u}_L^m \gamma^\mu u_L^n - \bar{d}_L^i \gamma^\mu d_L^j \right) \left(V_{mk} V_{nl}^* \bar{u}_L^m \gamma_\mu u_L^n - \bar{d}_L^k \gamma_\mu d_L^l \right) + \dots$ $\tilde{C}_{qq}^{(3)ijkl} = C_{qq}^{(3)pqrs} S_{Lip}^{d\dagger} S_{Lqj}^d S_{Lkr}^{d\dagger} S_{Lsl}^d$
$Q_{qd}^{(1)}$	$\tilde{C}_{qd}^{(1)ijkl} \left(V_{mi} V_{nj}^* \bar{u}_L^m \gamma^\mu u_L^n + \bar{d}_L^i \gamma^\mu d_L^j \right) \left(\bar{d}_R^k \gamma_\mu d_R^l \right)$ $\tilde{C}_{qd}^{(1)ijkl} = C_{qd}^{(1)pqrs} S_{Lip}^{d\dagger} S_{Lqj}^d S_{Rkr}^{d\dagger} S_{Rsl}^d$
$Q_{qd}^{(8)}$	$\tilde{C}_{qd}^{(8)ijkl} \left(V_{mi} V_{nj}^* \bar{u}_L^m \gamma^\mu T^A u_L^n + \bar{d}_L^i \gamma^\mu T^A d_L^j \right) \left(\bar{d}_R^k \gamma_\mu T^A d_R^l \right)$ $\tilde{C}_{qd}^{(8)ijkl} = C_{qd}^{(8)pqrs} S_{Lip}^{d\dagger} S_{Lqj}^d S_{Rkr}^{d\dagger} S_{Rsl}^d$
$Q_{qu}^{(1)}$	$\tilde{C}_{qu}^{(1)ijkl} \left(V_{mi} V_{nj}^* \bar{u}_L^m \gamma^\mu u_L^n + \bar{d}_L^i \gamma^\mu d_L^j \right) \left(\bar{u}_R^k \gamma_\mu u_R^l \right)$ $\tilde{C}_{qu}^{(1)ijkl} = C_{qu}^{(1)pqrs} S_{Lip}^{d\dagger} S_{Lqj}^d S_{Rkr}^{u\dagger} S_{Rsl}^u$
$Q_{qu}^{(8)}$	$\tilde{C}_{qu}^{(8)ijkl} \left(V_{mi} V_{nj}^* \bar{u}_L^m \gamma^\mu T^A u_L^n + \bar{d}_L^i \gamma^\mu T^A d_L^j \right) \left(\bar{u}_R^k \gamma_\mu T^A u_R^l \right)$ $\tilde{C}_{qu}^{(8)ijkl} = C_{qu}^{(8)pqrs} S_{Lip}^{d\dagger} S_{Lqj}^d S_{Rkr}^{u\dagger} S_{Rsl}^u$
$Q_{ud}^{(1)}$	$\tilde{C}_{ud}^{(1)ijkl} \left(\bar{u}_R^i \gamma^\mu u_R^j \right) \left(\bar{d}_R^k \gamma_\mu d_R^l \right)$ $\tilde{C}_{ud}^{(1)ijkl} = C_{ud}^{(1)pqrs} S_{Rip}^{u\dagger} S_{Rqj}^u S_{Rkr}^{d\dagger} S_{Rsl}^d$
$Q_{ud}^{(8)}$	$\tilde{C}_{ud}^{(8)ijkl} \left(\bar{u}_R^i T^A \gamma^\mu u_R^j \right) \left(\bar{d}_R^k T^A \gamma_\mu d_R^l \right)$ $\tilde{C}_{ud}^{(8)ijkl} = C_{ud}^{(8)pqrs} S_{Rip}^{u\dagger} S_{Rqj}^u S_{Rkr}^{d\dagger} S_{Rsl}^d$
Q_{dd}	$\tilde{C}_{dd}^{ijkl} \left(\bar{d}_R^i \gamma^\mu d_R^j \right) \left(\bar{d}_R^k \gamma_\mu d_R^l \right)$ $\tilde{C}_{dd}^{ijkl} = C_{dd}^{pqrs} S_{Rip}^{d\dagger} S_{Rqj}^d S_{Rkr}^{d\dagger} S_{Rsl}^d$
Q_{uu}	$\tilde{C}_{uu}^{ijkl} \left(\bar{u}_R^i \gamma^\mu u_R^j \right) \left(\bar{u}_R^k \gamma_\mu u_R^l \right)$ $\tilde{C}_{uu}^{ijkl} = C_{uu}^{pqrs} S_{Rip}^{u\dagger} S_{Rqj}^u S_{Rkr}^{u\dagger} S_{Rsl}^u$
$Q_{quqd}^{(1)}$	$\tilde{C}_{quqd}^{(1)ijkl} \left[\left(\bar{u}_L^i u_R^j \right) \left(\bar{d}_L^k d_R^l \right) - V_{im}^* V_{nk} \left(\bar{d}_L^m u_R^j \right) \left(\bar{u}_L^n d_R^l \right) \right]$ $\tilde{C}_{quqd}^{(1)ijkl} = C_{quqd}^{(1)pqrs} S_{Lip}^{u\dagger} S_{Rqj}^u S_{Lkr}^{d\dagger} S_{Rsl}^d$
$Q_{quqd}^{(8)}$	$\tilde{C}_{quqd}^{(8)ijkl} \left[\left(\bar{u}_L^i T^A u_R^j \right) \left(\bar{d}_L^k T^A d_R^l \right) - V_{im}^* V_{nk} \left(\bar{d}_L^m T^A u_R^j \right) \left(\bar{u}_L^n T^A d_R^l \right) \right]$ $\tilde{C}_{quqd}^{(8)ijkl} = C_{quqd}^{(8)pqrs} S_{Lip}^{u\dagger} S_{Rqj}^u S_{Lkr}^{d\dagger} S_{Rsl}^d$

Table 6: Four-fermion operators with two quarks and two leptons

Operator	Definition in the mass basis
$Q_{\ell q}^{(1)}$	$\tilde{C}_{\ell q}^{(1)ijkl} \left(\bar{\nu}_L^i \gamma^\mu \nu_L^j + \bar{e}_L^i \gamma^\mu e_L^j \right) (V_{mk} V_{nl}^* \bar{u}_L^m \gamma_\mu u_L^n + \bar{d}_L^k \gamma_\mu d_L^l)$ $\tilde{C}_{\ell q}^{(1)ijkl} = C_{\ell q}^{(1)ijmn} S_{Lkm}^{d\dagger} S_{Lnl}^d$
$Q_{\ell q}^{(3)}$	$\tilde{C}_{\ell q}^{(3)ijkl} \left(\bar{\nu}_L^i \gamma^\mu \nu_L^j - \bar{e}_L^i \gamma^\mu e_L^j \right) (V_{mk} V_{nl}^* \bar{u}_L^m \gamma_\mu u_L^n - \bar{d}_L^k \gamma_\mu d_L^l) \dots$ $\tilde{C}_{\ell q}^{(3)ijkl} = C_{\ell q}^{(3)ijmn} S_{Lkm}^{d\dagger} S_{Lnl}^d$
Q_{eu}	$\tilde{C}_{eu}^{ijkl} \left(\bar{e}_R^i \gamma^\mu e_R^j \right) (\bar{u}_R^k \gamma_\mu u_R^l)$ $\tilde{C}_{eu}^{ijkl} = C_{eu}^{ijmn} S_{Rkm}^{u\dagger} S_{Rnl}^u$
Q_{ed}	$\tilde{C}_{ed}^{ijkl} \left(\bar{e}_R^i \gamma^\mu e_R^j \right) (\bar{d}_R^k \gamma_\mu d_R^l)$ $\tilde{C}_{ed}^{ijkl} = C_{ed}^{ijmn} S_{Rkm}^{d\dagger} S_{Rnl}^d$
$Q_{\ell u}$	$\tilde{C}_{\ell u}^{ijkl} \left(\bar{\nu}_L^i \gamma^\mu \nu_L^j + \bar{e}_L^i \gamma^\mu e_L^j \right) (\bar{u}_R^k \gamma_\mu u_R^l)$ $\tilde{C}_{\ell u}^{ijkl} = C_{\ell u}^{ijmn} S_{Rkm}^{u\dagger} S_{Rnl}^u$
$Q_{\ell d}$	$\tilde{C}_{\ell d}^{ijkl} \left(\bar{\nu}_L^i \gamma^\mu \nu_L^j + \bar{e}_L^i \gamma^\mu e_L^j \right) (\bar{d}_R^k \gamma_\mu d_R^l)$ $\tilde{C}_{\ell d}^{ijkl} = C_{\ell d}^{ijmn} S_{Rkm}^{d\dagger} S_{Rnl}^d$
Q_{qe}	$\tilde{C}_{qe}^{ijkl} \left(V_{mi} V_{nj}^* \bar{u}_L^m \gamma_\mu u_L^n + \bar{d}_L^i \gamma_\mu d_L^j \right) (\bar{e}_R^k \gamma^\mu e_R^l)$ $\tilde{C}_{qe}^{ijkl} = C_{qe}^{mnkl} S_{Lim}^{d\dagger} S_{Lnj}^d$
Q_{ledq}	$\tilde{C}_{ledq}^{ijkl} \left[V_{nl}^* \left(\bar{\nu}_L^i e_R^j \right) (\bar{d}_R^k u_L^n) + \left(\bar{e}_L^i e_R^j \right) (\bar{d}_R^k d_L^l) \right]$ $\tilde{C}_{ledq}^{ijkl} = C_{ledq}^{ijmn} S_{Rkm}^{d\dagger} S_{Lnl}^d$
$Q_{lequ}^{(1)}$	$\tilde{C}_{lequ}^{(1)ijkl} \left[V_{km}^* \left(\bar{\nu}_L^i e_R^j \right) (\bar{d}_L^m u_R^l) - \left(\bar{e}_L^i e_R^j \right) (\bar{u}_L^k u_R^l) \right]$ $\tilde{C}_{lequ}^{(1)ijkl} = C_{lequ}^{(1)ijmn} S_{Lkm}^{u\dagger} S_{Rnl}^u$
$Q_{lequ}^{(3)}$	$\tilde{C}_{lequ}^{(3)ijkl} \left[V_{km}^* \left(\bar{\nu}_L^i \sigma^{\mu\nu} e_R^j \right) (\bar{d}_L^m \sigma_{\mu\nu} u_R^l) - \left(\bar{e}_L^i \sigma^{\mu\nu} e_R^j \right) (\bar{u}_L^k \sigma_{\mu\nu} u_R^l) \right]$ $\tilde{C}_{lequ}^{(3)ijkl} = C_{lequ}^{(3)ijmn} S_{Lkm}^{u\dagger} S_{Rnl}^u$

References

- [1] **Particle Data Group** Collaboration, J. Beringer et al., *Review of Particle Physics (RPP)*, *Phys.Rev.* **D86** (2012) 010001.
- [2] **ATLAS** Collaboration, G. Aad et al., *Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC*, *Phys. Lett.* **B716** (2013) 1–29, [[arXiv:1207.7214](#)].
- [3] **CMS** Collaboration, S. Chatrchyan et al., *Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC*, *Phys. Lett.* **B716** (2013) 30–61, [[arXiv:1207.7235](#)].
- [4] T. Appelquist and J. Carazzone, *Infrared Singularities and Massive Fields*, *Phys.Rev.* **D11** (1975) 2856.
- [5] S. Weinberg, *Baryon and Lepton Nonconserving Processes*, *Phys. Rev. Lett.* **43** (1979) 1566–1570.

- [6] W. Buchmuller and D. Wyler, *Effective Lagrangian Analysis of New Interactions and Flavor Conservation*, *Nucl.Phys.* **B268** (1986) 621.
- [7] B. Grzadkowski, M. Iskrzynski, M. Misiak, and J. Rosiek, *Dimension-Six Terms in the Standard Model Lagrangian*, *JHEP* **1010** (2010) 085, [[arXiv:1008.4884](#)].
- [8] S. Weinberg, *The Quantum theory of fields. Vol. 1: Foundations*, .
- [9] S. Fajfer, J. F. Kamenik, and I. Nisandzic, *On the $B \rightarrow D^* \tau \bar{\nu}_\tau$ Sensitivity to New Physics*, *Phys. Rev.* **D85** (2012) 094025, [[arXiv:1203.2654](#)].
- [10] **Heavy Flavor Averaging Group (HFAG)** Collaboration, Y. Amhis et al., *Averages of b -hadron, c -hadron, and τ -lepton properties as of summer 2014*, online update at <http://www.slac.stanford.edu/xorg/hfag/>, [arXiv:1412.7515](#).
- [11] T. Hurth, F. Mahmoudi, and S. Neshatpour, *Global fits to $b \rightarrow s \ell \ell$ data and signs for lepton non-universality*, *JHEP* **12** (2014) 053, [[arXiv:1410.4545](#)].
- [12] W. Altmannshofer and D. M. Straub, *New physics in $b \rightarrow s$ transitions after LHC run 1*, *Eur. Phys. J.* **C75** (2015), no. 8 382, [[arXiv:1411.3161](#)].
- [13] S. Descotes-Genon, L. Hofer, J. Matias, and J. Virto, *Global analysis of $b \rightarrow s \ell \ell$ anomalies*, [arXiv:1510.04239](#).
- [14] S. Descotes-Genon, J. Matias, and J. Virto, *Understanding the $B \rightarrow K^* \mu^+ \mu^-$ Anomaly*, *Phys. Rev.* **D88** (2013) 074002, [[arXiv:1307.5683](#)].
- [15] R. Gauld, F. Goertz, and U. Haisch, *On minimal Z' explanations of the $B \rightarrow K^* \mu^+ \mu^-$ anomaly*, *Phys. Rev.* **D89** (2014) 015005, [[arXiv:1308.1959](#)].
- [16] A. J. Buras and J. Girrbach, *Left-handed Z' and Z FCNC quark couplings facing new $b \rightarrow s \mu^+ \mu^-$ data*, *JHEP* **12** (2013) 009, [[arXiv:1309.2466](#)].
- [17] R. Gauld, F. Goertz, and U. Haisch, *An explicit Z' -boson explanation of the $B \rightarrow K^* \mu^+ \mu^-$ anomaly*, *JHEP* **01** (2014) 069, [[arXiv:1310.1082](#)].
- [18] A. J. Buras, F. De Fazio, and J. Girrbach, *331 models facing new $b \rightarrow s \mu^+ \mu^-$ data*, *JHEP* **02** (2014) 112, [[arXiv:1311.6729](#)].
- [19] W. Altmannshofer, S. Gori, M. Pospelov, and I. Yavin, *Quark flavor transitions in $L_\mu - L_\tau$ models*, *Phys. Rev.* **D89** (2014) 095033, [[arXiv:1403.1269](#)].
- [20] B. Gripaios, M. Nardecchia, and S. A. Renner, *Composite leptoquarks and anomalies in B -meson decays*, *JHEP* **05** (2015) 006, [[arXiv:1412.1791](#)].
- [21] A. Crivellin, G. D'Ambrosio, and J. Heeck, *Explaining $h \rightarrow \mu^\pm \tau^\mp$, $B \rightarrow K^* \mu^+ \mu^-$ and $B \rightarrow K \mu^+ \mu^- / B \rightarrow K e^+ e^-$ in a two-Higgs-doublet model with gauged $L_\mu - L_\tau$* , *Phys. Rev. Lett.* **114** (2015) 151801, [[arXiv:1501.00993](#)].
- [22] A. Crivellin, G. D'Ambrosio, and J. Heeck, *Addressing the LHC flavor anomalies with horizontal gauge symmetries*, *Phys. Rev.* **D91** (2015), no. 7 075006, [[arXiv:1503.03477](#)].
- [23] D. Bećirević, S. Fajfer, and N. Košnik, *Lepton flavor nonuniversality in $b \rightarrow s \ell^+ \ell^-$ processes*, *Phys. Rev.* **D92** (2015), no. 1 014016, [[arXiv:1503.09024](#)].
- [24] C. Niehoff, P. Stangl, and D. M. Straub, *Violation of lepton flavour universality in composite Higgs models*, *Phys. Lett.* **B747** (2015) 182–186, [[arXiv:1503.03865](#)].
- [25] I. de Medeiros Varzielas and G. Hiller, *Clues for flavor from rare lepton and quark decays*, *JHEP* **06** (2015) 072, [[arXiv:1503.01084](#)].

- [26] D. Aristizabal Sierra, F. Staub, and A. Vicente, *Shedding light on the $b \rightarrow s$ anomalies with a dark sector*, *Phys. Rev.* **D92** (2015), no. 1 015001, [[arXiv:1503.06077](#)].
- [27] A. Celis, J. Fuentes-Martin, M. Jung, and H. Serodio, *Family nonuniversal Z models with protected flavor-changing interactions*, *Phys. Rev.* **D92** (2015), no. 1 015007, [[arXiv:1505.03079](#)].
- [28] S. Sahoo and R. Mohanta, *Leptoquark effects on $b \rightarrow s\nu\bar{\nu}$ and $B \rightarrow Kl^+l^-$ decay processes*, [arXiv:1509.06248](#).
- [29] G. Bélanger, C. Delaunay, and S. Westhoff, *A Dark Matter Relic From Muon Anomalies*, *Phys. Rev.* **D92** (2015) 055021, [[arXiv:1507.06660](#)].
- [30] A. Falkowski, M. Nardecchia, and R. Ziegler, *Lepton Flavor Non-Universality in B-meson Decays from a $U(2)$ Flavor Model*, [arXiv:1509.01249](#).
- [31] B. Grinstein and J. M. Camalich, *Weak decays of unstable b-mesons*, [arXiv:1509.05049](#).
- [32] B. Gripaios, M. Nardecchia, and S. A. Renner, *Linear flavour violation and anomalies in B physics*, [arXiv:1509.05020](#).
- [33] A. Carmona and F. Goertz, *Lepton Flavor and Non-Universality from Minimal Composite Higgs Setups*, [arXiv:1510.07658](#).
- [34] M. Bauer and M. Neubert, *One Leptoquark to Rule Them All: A Minimal Explanation for $R_{D^{(*)}}$, R_K and $(g-2)_\mu$* , [arXiv:1511.01900](#).
- [35] S. Fajfer and N. Konik, *Vector leptoquark resolution of R_K and $R_{D^{(*)}}$ puzzles*, [arXiv:1511.06024](#).
- [36] A. Crivellin, C. Greub, and A. Kokulu, *Explaining $B \rightarrow D\tau\nu$, $B \rightarrow D^*\tau\nu$ and $B \rightarrow \tau\nu$ in a 2HDM of type III*, *Phys. Rev.* **D86** (2012) 054014, [[arXiv:1206.2634](#)].
- [37] A. Datta, M. Duraisamy, and D. Ghosh, *Diagnosing New Physics in $b \rightarrow c\tau\nu_\tau$ decays in the light of the recent BaBar result*, *Phys. Rev.* **D86** (2012) 034027, [[arXiv:1206.3760](#)].
- [38] A. Celis, M. Jung, X.-Q. Li, and A. Pich, *Sensitivity to charged scalars in $B \rightarrow D^{(*)}\tau\nu_\tau$ and $B \rightarrow \tau\nu_\tau$ decays*, *JHEP* **01** (2013) 054, [[arXiv:1210.8443](#)].
- [39] A. Crivellin, A. Kokulu, and C. Greub, *Flavor-phenomenology of two-Higgs-doublet models with generic Yukawa structure*, *Phys. Rev.* **D87** (2013), no. 9 094031, [[arXiv:1303.5877](#)].
- [40] X.-Q. Li, Y.-D. Yang, and X.-B. Yuan, *Exclusive radiative B-meson decays within minimal flavor-violating two-Higgs-doublet models*, *Phys. Rev.* **D89** (2014), no. 5 054024, [[arXiv:1311.2786](#)].
- [41] G. Faisel, *Charged Higgs contribution to $\bar{B}_s \rightarrow \phi\pi^0$ and $\bar{B}_s \rightarrow \phi\rho^0$* , *Phys. Lett.* **B731** (2014) 279–286, [[arXiv:1311.0740](#)].
- [42] M. Atoui, V. Morénas, D. Bečirevic, and F. Sanfilippo, *$B_s \rightarrow D_s\ell\nu_\ell$ near zero recoil in and beyond the Standard Model*, *Eur. Phys. J.* **C74** (2014), no. 5 2861, [[arXiv:1310.5238](#)].
- [43] Y. Sakaki, M. Tanaka, A. Tayduganov, and R. Watanabe, *Testing leptoquark models in $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$* , *Phys. Rev.* **D88** (2013), no. 9 094012, [[arXiv:1309.0301](#)].
- [44] I. Doršner, S. Fajfer, N. Košnik, and I. Nišandžić, *Minimally flavored colored scalar in $\bar{B} \rightarrow D^{(*)}\tau\bar{\nu}$ and the mass matrices constraints*, *JHEP* **11** (2013) 084, [[arXiv:1306.6493](#)].
- [45] P. Biancofiore, P. Colangelo, and F. De Fazio, *Rare semileptonic $B \rightarrow K^*\ell^+\ell^-$ decays in RS_c model*, *Phys. Rev.* **D89** (2014), no. 9 095018, [[arXiv:1403.2944](#)].

- [46] A. Crivellin, J. Heeck, and P. Stoffer, *A perturbed lepton-specific two-Higgs-doublet model facing experimental hints for physics beyond the Standard Model*, [arXiv:1507.07567](#).
- [47] M. Freytsis, Z. Ligeti, and J. T. Ruderman, *Flavor models for $\overline{B} \rightarrow D^{(*)}\tau\overline{\nu}$* , *Phys. Rev.* **D92** (2015), no. 5 054018, [[arXiv:1506.08896](#)].
- [48] G. Buchalla, A. J. Buras, and M. E. Lautenbacher, *Weak decays beyond leading logarithms*, *Rev. Mod. Phys.* **68** (1996) 1125–1144, [[hep-ph/9512380](#)].
- [49] A. J. Buras, *Weak Hamiltonian, CP violation and rare decays*, in *Probing the standard model of particle interactions. Proceedings, Summer School in Theoretical Physics, NATO Advanced Study Institute, 68th session, Les Houches, France, July 28-September 5, 1997. Pt. 1, 2*, pp. 281–539, 1998. [hep-ph/9806471](#).
- [50] A. Crivellin, S. Najjari, and J. Rosiek, *Lepton Flavor Violation in the Standard Model with general Dimension-Six Operators*, *JHEP* **1404** (2014) 167, [[arXiv:1312.0634](#)].
- [51] A. Crivellin, M. Hoferichter, and M. Procura, *Improved predictions for $\mu \rightarrow e$ conversion in nuclei and Higgs-induced lepton flavor violation*, *Phys.Rev.* **D89** (2014) 093024, [[arXiv:1404.7134](#)].
- [52] G. M. Pruna and A. Signer, *The $\mu \rightarrow e\gamma$ decay in a systematic effective field theory approach with dimension 6 operators*, *JHEP* **1410** (2014) 14, [[arXiv:1408.3565](#)].
- [53] B. Dassinger, T. Feldmann, T. Mannel, and S. Turczyk, *Model-independent analysis of lepton flavour violating tau decays*, *JHEP* **0710** (2007) 039, [[arXiv:0707.0988](#)].
- [54] B. Bhattacharya, A. Datta, D. London, and S. Shivashankara, *Simultaneous Explanation of the R_K and $R(D^{(*)})$ Puzzles*, *Phys.Lett.* **B742** (2015) 370–374, [[arXiv:1412.7164](#)].
- [55] R. Alonso, B. Grinstein, and J. M. Camalich, *Lepton universality violation and lepton flavor conservation in B -meson decays*, [arXiv:1505.05164](#).
- [56] L. Calibbi, A. Crivellin, and T. Ota, *Effective field theory approach to $b \rightarrow s\ell\ell^{(\prime)}$, $B \rightarrow K^{(*)}\nu\overline{\nu}$ and $B \rightarrow D^{(*)}\tau\nu$ with third generation couplings*, [arXiv:1506.02661](#).
- [57] R. Alonso, B. Grinstein, and J. Martin Camalich, *$SU(2) \times U(1)$ gauge invariance and the shape of new physics in rare B decays*, *Phys.Rev.Lett.* **113** (2014) 241802, [[arXiv:1407.7044](#)].
- [58] A. J. Buras, J. Girrbach-Noe, C. Niehoff, and D. M. Straub, *$B \rightarrow K^{(*)}\nu\overline{\nu}$ decays in the Standard Model and beyond*, *JHEP* **1502** (2015) 184, [[arXiv:1409.4557](#)].
- [59] M. Ciuchini, E. Franco, V. Lubicz, G. Martinelli, I. Scimemi, and L. Silvestrini, *Next-to-leading order QCD corrections to $\Delta F = 2$ effective Hamiltonians*, *Nucl. Phys.* **B523** (1998) 501–525, [[hep-ph/9711402](#)].
- [60] A. J. Buras, M. Misiak, and J. Urban, *Two loop QCD anomalous dimensions of flavor changing four quark operators within and beyond the standard model*, *Nucl.Phys.* **B586** (2000) 397–426, [[hep-ph/0005183](#)].
- [61] F. Borzumati and C. Greub, *2HDMs predictions for $\overline{B} \rightarrow X_s\gamma$ in NLO QCD*, *Phys. Rev.* **D58** (1998) 074004, [[hep-ph/9802391](#)].
- [62] F. Borzumati, C. Greub, T. Hurth, and D. Wyler, *Gluino contribution to radiative B decays: Organization of QCD corrections and leading order results*, *Phys. Rev.* **D62** (2000) 075005, [[hep-ph/9911245](#)].
- [63] E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the*

- Standard Model Dimension Six Operators I: Formalism and lambda Dependence*, *JHEP* **1310** (2013) 087, [[arXiv:1308.2627](#)].
- [64] E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the Standard Model Dimension Six Operators II: Yukawa Dependence*, *JHEP* **01** (2014) 035, [[arXiv:1310.4838](#)].
 - [65] R. Alonso, E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the Standard Model Dimension Six Operators III: Gauge Coupling Dependence and Phenomenology*, *JHEP* **04** (2014) 159, [[arXiv:1312.2014](#)].
 - [66] C. Bobeth and U. Haisch, *Anomalous triple gauge couplings from B-meson and kaon observables*, *JHEP* **09** (2015) 018, [[arXiv:1503.04829](#)].
 - [67] G. Blankenburg, J. Ellis, and G. Isidori, *Flavour-Changing Decays of a 125 GeV Higgs-like Particle*, *Phys.Lett.* **B712** (2012) 386–390, [[arXiv:1202.5704](#)].
 - [68] S. Davidson and P. Verdier, *LHC sensitivity to the decay $h \rightarrow \tau^\pm \mu^\mp$* , *Phys. Rev.* **D86** (2012) 111701, [[arXiv:1211.1248](#)].
 - [69] D. Becirevic, M. Ciuchini, E. Franco, V. Gimenez, G. Martinelli, et al., *$B_d - \bar{B}_d$ mixing and the $B_d \rightarrow J/\psi K_s$ asymmetry in general SUSY models*, *Nucl.Phys.* **B634** (2002) 105–119, [[hep-ph/0112303](#)].
 - [70] M. B. Voloshin, *Bound on $V + A$ admixture in the $b \rightarrow c$ current from inclusive versus exclusive semileptonic decays of B mesons*, *Mod. Phys. Lett.* **A12** (1997) 1823–1827, [[hep-ph/9704278](#)].
 - [71] B. M. Dassinger, R. Feger, and T. Mannel, *Testing the left-handedness of the $b \rightarrow c$ transition*, *Phys. Rev.* **D75** (2007) 095007, [[hep-ph/0701054](#)].
 - [72] C.-H. Chen and S.-h. Nam, *Left-right mixing on leptonic and semileptonic $b \rightarrow u$ decays*, *Phys. Lett.* **B666** (2008) 462–466, [[arXiv:0807.0896](#)].
 - [73] B. Dassinger, R. Feger, and T. Mannel, *Complete Michel Parameter Analysis of inclusive semileptonic $b \rightarrow c$ transition*, *Phys. Rev.* **D79** (2009) 075015, [[arXiv:0803.3561](#)].
 - [74] A. Crivellin, *Effects of right-handed charged currents on the determinations of $|V_{ub}|$ and $|V_{cb}|$* , *Phys. Rev.* **D81** (2010) 031301, [[arXiv:0907.2461](#)].
 - [75] V. Cirigliano, J. Jenkins, and M. Gonzalez-Alonso, *Semileptonic decays of light quarks beyond the Standard Model*, *Nucl. Phys.* **B830** (2010) 95–115, [[arXiv:0908.1754](#)].
 - [76] X.-G. He, J. Tandean, and G. Valencia, *Probing New Physics in Charm Couplings with FCNC*, *Phys. Rev.* **D80** (2009) 035021, [[arXiv:0904.2301](#)].
 - [77] A. J. Buras, K. Gemmler, and G. Isidori, *Quark flavour mixing with right-handed currents: an effective theory approach*, *Nucl. Phys.* **B843** (2011) 107–142, [[arXiv:1007.1993](#)].
 - [78] R. Feger, T. Mannel, V. Klose, H. Lacker, and T. Luck, *Limit on a Right-Handed Admixture to the Weak $b \rightarrow c$ Current from Semileptonic Decays*, *Phys. Rev.* **D82** (2010) 073002, [[arXiv:1003.4022](#)].
 - [79] A. Crivellin and S. Pokorski, *Can the differences in the determinations of V_{ub} and V_{cb} be explained by New Physics?*, *Phys. Rev. Lett.* **114** (2015), no. 1 011802, [[arXiv:1407.1320](#)].
 - [80] F. U. Bernlochner, Z. Ligeti, and S. Turczyk, *New ways to search for right-handed current in B decay*, *Phys. Rev.* **D90** (2014), no. 9 094003, [[arXiv:1408.2516](#)].

- [81] J. Brod, A. Greljo, E. Stamou, and P. Uttayarat, *Probing anomalous $t\bar{t}Z$ interactions with rare meson decays*, *JHEP* **02** (2015) 141, [[arXiv:1408.0792](#)].
- [82] B. Grzadkowski and M. Misiak, *Anomalous Wtb coupling effects in the weak radiative B -meson decay*, *Phys. Rev.* **D78** (2008) 077501, [[arXiv:0802.1413](#)]. [Erratum: *Phys. Rev.* **D84**, 059903(2011)].
- [83] J. Drobnak, S. Fajfer, and J. F. Kamenik, *Probing anomalous tWb interactions with rare B decays*, *Nucl. Phys.* **B855** (2012) 82–99, [[arXiv:1109.2357](#)].
- [84] A. Crivellin and L. Mercolli, *$B \rightarrow X_d \gamma$ and constraints on new physics*, *Phys. Rev.* **D84** (2011) 114005, [[arXiv:1106.5499](#)].
- [85] M. Misiak et al., *Updated NNLO QCD predictions for the weak radiative B -meson decays*, *Phys. Rev. Lett.* **114** (2015), no. 22 221801, [[arXiv:1503.01789](#)].
- [86] M. Czakon, P. Fiedler, T. Huber, M. Misiak, T. Schutzmeier, and M. Steinhauser, *The $(Q_7, Q_{1,2})$ contribution to $\bar{B} \rightarrow X_s \gamma$ at $\mathcal{O}(\alpha_s^2)$* , *JHEP* **04** (2015) 168, [[arXiv:1503.01791](#)].
- [87] P. L. Cho and M. Misiak, *$b \rightarrow s \gamma$ decay in $SU(2)_L \times SU(2)_R \times U(1)$ extensions of the Standard Model*, *Phys. Rev.* **D49** (1994) 5894–5903, [[hep-ph/9310332](#)].
- [88] C. Bernardo, N. F. Castro, M. C. N. Fiolhais, H. Gonalves, A. G. C. Guerra, M. Oliveira, and A. Onofre, *Studying the Wtb vertex structure using recent LHC results*, *Phys. Rev.* **D90** (2014), no. 11 113007, [[arXiv:1408.7063](#)].
- [89] **ATLAS** Collaboration, G. Aad et al., *Search for anomalous couplings in the Wtb vertex from the measurement of double differential angular decay rates of single top quarks produced in the t -channel with the ATLAS detector*, [[arXiv:1510.03764](#)].