

A review of μ - τ flavor symmetry in neutrino physics

Zhi-zhong Xing¹ and Zhen-hua Zhao^{1,2}

¹Institute of High Energy Physics and School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

²Department of Physics, Liaoning Normal University, Dalian 116029, China

E-mail: xingzz@ihep.ac.cn and zhaozhenhua@ihep.ac.cn

First version: December 2015; Modified version: March 2016

Abstract. Behind the observed pattern of lepton flavor mixing is a partial or approximate μ - τ flavor symmetry — a milestone on our road to the true origin of neutrino masses and flavor structures. In this review article we first describe the features of μ - τ permutation and reflection symmetries, and then explore their various consequences on model building and neutrino phenomenology. We pay particular attention to soft μ - τ symmetry breaking, which is crucial for our deeper understanding of the fine effects of flavor mixing and CP violation.

Keywords: flavor mixing, μ - τ symmetry, neutrino mass and oscillation, particle physics

Contents

1	Introduction	2
1.1	A brief history of the neutrino families	2
1.2	The μ - τ flavor symmetry stands out	5
2	Behind the lepton flavor mixing pattern	7
2.1	Lepton flavor mixing and neutrino oscillations	7
2.2	Current neutrino oscillation experiments	10
2.3	The observed pattern of the PMNS matrix	13
3	An overview of the μ-τ flavor symmetry	16
3.1	The μ - τ permutation symmetry	20
3.2	The μ - τ reflection symmetry	23
3.3	Breaking of the μ - τ permutation symmetry	26
3.4	Breaking of the μ - τ reflection symmetry	35
3.5	RGE-induced μ - τ symmetry breaking effects	38
3.6	Flavor mixing from the charged-lepton sector	45

4	Larger flavor symmetry groups	49
4.1	Neutrino mixing and flavor symmetries	49
4.2	Model building with discrete flavor symmetries	52
4.3	Generalized CP and spontaneous CP violation	57
5	Realization of the μ-τ flavor symmetry	61
5.1	Models with the μ - τ permutation symmetry	62
5.2	Models with the μ - τ reflection symmetry	65
5.3	On the TM1 and TM2 neutrino mixing patterns	68
5.4	When the sterile neutrinos are concerned	71
6	Some consequences of the μ-τ symmetry	75
6.1	Neutrino oscillations in matter	75
6.2	Flavor distributions of UHE cosmic neutrinos	78
6.3	Matter-antimatter asymmetry via leptogenesis	82
6.4	Fermion mass matrices with the Z_2 symmetry	85
7	Summary and outlook	88
8	Appendix	90

1. Introduction

1.1. A brief history of the neutrino families

Soon after the French physicist Henri Becquerel first discovered the radioactivity of uranium in 1896 [1], some nuclear physicists began to focus their attention on the beta decays $(A, Z) \rightarrow (A, Z + 1) + e^-$, where the energy spectrum of the outgoing electrons was expected to be *discrete* as constrained by the energy and momentum conservations. Surprisingly, the British physicist James Chadwick observed a *continuous* electron energy spectrum of the beta decay in 1914 [2], and such a result was firmly established in the 1920s [3]. At that time there were two typical points of view towards resolving this discrepancy between the *observed* and *expected* energy spectra of electrons: (a) the Danish theorist Niels Bohr intended to give up the energy conservation law, and later his idea turned out to be wrong; (b) the Austrian theorist Wolfgang Pauli preferred to add in a new particle, which marked the birth of a new science. In 1930 Pauli pointed out that a light, spin-1/2 and neutral particle — known as the electron antineutrino today — appeared in the beta decay and carried away some energy and momentum in an invisible way, and thus the energy spectrum of electrons in the process $(A, Z) \rightarrow (A, Z + 1) + e^- + \bar{\nu}_e$ was continuous. Three years later the Italian theorist Enrico Fermi took this hypothesis seriously and developed an effective field theory of the beta decay [4], which made it possible to calculate the reaction rates of nucleons and electrons or positrons interacting with neutrinos or antineutrinos.

In 1936 the German physicist Hans Bethe pointed out that an inverse beta decay mode of the form $\bar{\nu}_e + (A, Z) \rightarrow (A, Z - 1) + e^+$ should be a feasible way to verify the existence of electron antineutrinos produced from fission bombs or reactors [5]. This bright idea was elaborated by the Italian theorist Bruno Pontecorvo in 1946 [6], and it became more realistic with the development of the liquid scintillation counting techniques in the 1950s. For example, the invisible incident $\bar{\nu}_e$ triggers the reaction $\bar{\nu}_e + p \rightarrow n + e^+$, in which the emitted positron annihilates with an electron and the daughter nucleus is captured in the detector. Both events can be observed since they emit gamma rays, and the corresponding flashes in the liquid scintillator are separated by some microseconds. The American experimentalists Frederick Reines and Clyde Cowan did the first reactor antineutrino experiment and confirmed Pauli's hypothesis in 1956 [7]. Such a discovery motivated Pontecorvo to speculate on the possibility of lepton number violation and neutrino-antineutrino transitions in 1957 [8]. His viewpoint was based on a striking conjecture made by Fermi's doctoral student Ettore Majorana in 1937: a massive neutrino could be its own antiparticle [9]. Whether Majorana's hypothesis is true or not remains an open question in particle physics.

In 1962 the muon neutrino — a puzzling sister of the electron neutrino — was first observed by the American experimentalists Leon Lederman, Melvin Schwartz and Jack Steinberger in a new accelerator-based experiment [10]. Their discovery more or less motivated the Japanese theorists Ziro Maki, Masami Nakagawa and Shoichi Sakata to think about lepton flavor mixing and $\nu_e \leftrightarrow \nu_\mu$ transitions [11]. The tau neutrino, another sister of the electron neutrino, was finally observed at the Fermilab in 2001 [12]. Today we are left with three lepton families consisting of the charged members (e, μ, τ) and the neutral members (ν_e, ν_μ, ν_τ), together with their antiparticles. Table 1.1 shows the total lepton number (L) and individual flavor numbers (L_e, L_μ, L_τ) assigned to all the known leptons and antileptons in the standard theory of electroweak interactions. So far the nonconservation of L_e, L_μ and L_τ has been observed in a number of neutrino oscillation experiments.

The standard electroweak theory about charged leptons and neutrinos was first formulated by the American theorist Steven Weinberg in 1967 [13]. In this seminal paper the neutrinos were assumed to be massless, and hence there should be no lepton flavor conversion. Just one year later, a preliminary experimental evidence for finite neutrino masses and lepton flavor mixing appeared thanks to the first observation of solar neutrinos and their deficit as compared with the prediction of the standard solar model [14]. The point was that the observed deficit of solar electron neutrinos could easily be attributed to $\nu_e \rightarrow \nu_\mu$ and $\nu_e \rightarrow \nu_\tau$ oscillations [15] — a pure quantum phenomenon which would not take place if every neutrino were massless and the lepton flavor were conserved. Hitherto the flavor oscillations of solar, atmospheric, accelerator and reactor neutrinos (or antineutrinos) have all been established [16], and thus a nontrivial extension of the standard theory of electroweak interactions is unavoidable in order to explain the origin of nonzero but tiny neutrino masses and the dynamics of significant lepton flavor mixing.

Table 1.1. The total lepton number (L) and individual flavor numbers (L_e, L_μ, L_τ) of three families of leptons and antileptons in the standard theory of electroweak interactions.

	1st family		2nd family		3rd family	
	e^-	ν_e	μ^-	ν_μ	τ^-	ν_τ
L	+1	+1	+1	+1	+1	+1
L_e	+1	+1	0	0	0	0
L_μ	0	0	+1	+1	0	0
L_τ	0	0	0	0	+1	+1
	e^+	$\bar{\nu}_e$	μ^+	$\bar{\nu}_\mu$	τ^+	$\bar{\nu}_\tau$
L	-1	-1	-1	-1	-1	-1
L_e	-1	-1	0	0	0	0
L_μ	0	0	-1	-1	0	0
L_τ	0	0	0	0	-1	-1

At present the most popular and intriguing idea for neutrino mass generation is the *seesaw* mechanism, which attributes the tiny masses of three neutrinos to the existence of a few heavy degrees of freedom and lepton number violation. There are three typical seesaw mechanisms on the market:

- Type-I seesaw — two or three heavy right-handed neutrinos are added into the standard theory and the lepton number is violated by their Majorana mass term [17, 18, 19, 20, 21];
- Type-II seesaw — one heavy Higgs triplet is added into the standard theory and the lepton number is violated by its interactions with both the lepton doublet and the Higgs doublet [22, 23, 24, 25, 26];
- Type-III seesaw — three heavy triplet fermions are introduced into the standard theory and the lepton number is violated by their Majorana mass term [27, 28].

After the heavy degrees of freedom are integrated out, all the three seesaw mechanisms are convergent to a unique effective Majorana neutrino mass operator [29]. Although a given seesaw scenario can qualitatively explain why the neutrinos may have tiny masses, it is not powerful enough to determine the flavor structures of charged leptons and neutrinos. Hence a combination of the seesaw idea and possible flavor symmetries is desirable so as to achieve some testable quantitative predictions in the lepton sector.

In comparison with three lepton families, there exist three quark families consisting of the up-type quarks (u, c, t) and the down-type quarks (d, s, b), together with their antiparticles. All these leptons and quarks constitute the flavor part of particle physics, and their mass spectra, flavor mixing properties and CP violation are the central issues of flavor dynamics. In this review article we shall focus on the lepton flavors, especially the μ - τ flavor symmetry in the neutrino sector and its striking impacts on the phenomenology of neutrino physics.

1.2. The μ - τ flavor symmetry stands out

Since Noether's theorem was first published in 1918 [30], symmetries have been playing a very crucial role in understanding the fundamental laws of Nature. In fact, symmetries are so powerful that they can help simplify the complicated problems, classify the intricate systems, pin down the conservation laws and even determine the dynamics of interactions. In elementary particle physics there are many successful examples of this kind, such as the continuous space-time translation symmetries, the $SU(3)_q$ quark flavor symmetry and the $SU(3)_c \times SU(2)_L \times U(1)_Y$ gauge symmetries. Historically these examples led us to the momentum and energy conservation laws, the quark model and the standard model (SM) of electroweak and strong interactions, respectively [16]. The original symmetries in a given theory may either keep exact or be broken, so as to make our description of the relevant phenomena consistent with the experimental observations. For instance, the electromagnetic $U(1)_{em}$ gauge symmetry, the strong $SU(3)_c$ gauge symmetry and the continuous space-time translation symmetries are all exact; but the $SU(3)_q$ quark flavor symmetry, the electroweak $SU(2)_L \times U(1)_Y$ gauge symmetry and the P and CP symmetries in weak interactions must be broken. That is why exploring new symmetries and studying possible symmetry-breaking effects have been one of the main streams in particle physics, normally from lower energies to higher energies.

Although the SM has proved to be very successful in describing the phenomena of strong, weak and electromagnetic interactions, it is by no means a complete theory. The flavor part of this theory is particularly unsatisfactory, because it involves many free parameters but still cannot provide any solution to a number of burning problems, such as the origin of neutrino masses and lepton flavor mixing, the baryon number asymmetry of the Universe and the nature of cold or warm dark matter. To go beyond the SM in this connection, flavor symmetries are expected to serve for a powerful guideline.

Flavor symmetries can be either Abelian or non-Abelian, either local or global, either continuous or discrete, and either spontaneously broken or explicitly broken. All these possibilities have been extensively explored in the past few decades, so as to explain the observed lepton and quark mass spectra and the observed flavor mixing patterns [31, 32, 33, 34]. Given the peculiar pattern of lepton flavor mixing which is suggestive of a constant unitary matrix with some special entities (e.g., $1/\sqrt{2}$, $1/\sqrt{3}$ or $1/\sqrt{6}$), a lot of attention has been paid to the global and discrete flavor symmetry groups in the model building exercises. The advantages of such a choice are obvious at least in the following aspects: (a) the model does not involve any Goldstone bosons or additional gauge bosons which may mediate harmful flavor-changing-neutral-current processes; (b) the discrete group may come from some string compactifications or be embedded in a continuous symmetry group; (c) the model contains no family-dependent D-terms contributing to the sfermion masses if it is built in a supersymmetric framework. Although many discrete flavor symmetry groups have been taken into account in building viable neutrino mass models, it remains unclear which one can finally stand out as the unique basis of

the true flavor dynamics of leptons and quarks.

But it turns out to be clear that any promising discrete flavor symmetry group in the neutrino sector has to accommodate the simplest μ - τ flavor symmetry — a sort of Z_2 transformation symmetry with respect to the ν_μ and ν_τ neutrinos, because the latter has convincingly revealed itself through the currently available neutrino oscillation data (as one can see in the next section). In other words, a partial or approximate μ - τ flavor symmetry must be behind the observed pattern of the 3×3 Pontecorvo-Maki-Nakagawa-Sakata (PMNS) lepton flavor mixing matrix U [35], and thus it may serve as an especially useful low-energy window to look into the underlying structures of lepton flavors at either the electroweak scale or superhigh-energy scales. It is just this observation that motivates us to review what we have learnt from

- the μ - τ *permutation* symmetry — the neutrino mass term is unchanged under the transformations

$$\nu_e \rightarrow \nu_e, \quad \nu_\mu \rightarrow \nu_\tau, \quad \nu_\tau \rightarrow \nu_\mu; \quad (1.1)$$

- the μ - τ *reflection* symmetry — the neutrino mass term keeps unchanged under the transformations

$$\nu_e \rightarrow \nu_e^c, \quad \nu_\mu \rightarrow \nu_\tau^c, \quad \nu_\tau \rightarrow \nu_\mu^c, \quad (1.2)$$

where the superscript “ c ” denotes the charge conjugation of the relevant neutrino field, and to explore their interesting implications and consequences on various aspects of neutrino phenomenology.

In particular, we stress that slight or soft breaking of the μ - τ flavor symmetry is expected to help resolve the octant of the largest neutrino mixing angle θ_{23} and even the quadrant of the CP-violating phase δ in the standard parametrization of the PMNS matrix U [16], which consists of three rotation angles $(\theta_{12}, \theta_{13}, \theta_{23})$ and one (δ in the Dirac case) or three (δ, ρ, σ in the Majorana case) CP-violating phases. It may also offer a straightforward link between the neutrino mass spectrum and the lepton flavor mixing pattern. All these issues are very important on the theoretical side and highly concerned in the ongoing and upcoming neutrino experiments.

The remaining parts of this review paper are organized in the following way. In section 2 we begin with a brief introduction to the phenomenology of lepton flavor mixing and neutrino oscillations, followed by a short description of the main outcomes of various solar, atmospheric, reactor and accelerator neutrino oscillation experiments. In the three-flavor scheme a global analysis of current experimental data leads us to a preliminary pattern of lepton flavor mixing, which exhibits an approximate μ - τ flavor symmetry. Section 3 is devoted to an overview of the μ - τ flavor symmetry of the Majorana neutrino mass matrix and its connection with the flavor mixing parameters. Two kinds of symmetries, the μ - τ *permutation* symmetry and the μ - τ *reflection* symmetry, will be classified and discussed. The typical and instructive ways to slightly break the μ - τ flavor symmetry are introduced. In particular, the effects of μ - τ symmetry breaking induced by radiative corrections are described in some

detail. The contribution of the charged-lepton sector to lepton flavor mixing is also discussed. In section 4 we turn to some larger discrete flavor symmetry groups to illustrate how the μ - τ symmetry can naturally arise as a residual symmetry. Both the bottom-up approach and the top-down approach will be described in this connection. In section 5 we concentrate on the strategies of model building with the help of the μ - τ permutation or reflection symmetry. A combination of the seesaw mechanism and the μ - τ flavor symmetry is taken into account, and the relationship between the Friedberg-Lee symmetry and the μ - τ flavor symmetry is explored. We also comment on the model-building exercises associated with the light sterile neutrinos. Section 6 is devoted to some phenomenological consequences of the μ - τ flavor symmetry on some interesting topics in neutrino physics, including neutrino oscillations in terrestrial matter, flavor distributions of the Ultrahigh-energy (UHE) cosmic neutrinos at neutrino telescopes, a possible connection between the leptogenesis and low-energy CP violation, and a likely unified flavor texture of leptons and quarks. The concluding remarks and an outlook are finally made in section 7.

2. Behind the lepton flavor mixing pattern

To see why an approximate μ - τ flavor symmetry is behind the observed pattern of the PMNS lepton flavor mixing matrix U , let us first introduce some basics about neutrino mixing and flavor oscillations and then discuss current experimental constraints on the structure of U .

2.1. Lepton flavor mixing and neutrino oscillations

Just similar to quark flavor mixing, lepton flavor mixing can also take place provided leptonic weak charged-current interactions and Yukawa interactions coexist in a simple extension of the SM. The standard form of weak charged-current interactions of the charged leptons and neutrinos reads

$$-\mathcal{L}_{\text{cc}} = \frac{g}{\sqrt{2}} \sum_{\alpha} [\overline{\alpha'_L} \gamma^{\mu} \nu_{\alpha L} W_{\mu}^{-} + \text{h.c.}] , \quad (2.1)$$

where α runs over e , μ and τ , and the superscript “ \prime ” denotes the flavor eigenstate of a charged lepton. The leptonic Yukawa interactions are expected to be responsible for the mass generation of both charged leptons and neutrinos after spontaneous electroweak symmetry breaking, although the origin of neutrino masses is very likely to involve some new degrees of freedom and lepton number violation [36]. Without going into details of a specific neutrino mass model, here we assume massive neutrinos to be the Majorana particles and write out the effective lepton mass terms as follows:

$$-\mathcal{L}_{\text{m}} = \frac{1}{2} \overline{\nu_{\alpha L}} (M_{\nu})_{\alpha\beta} \nu_{\beta R}^c + \overline{\alpha'_L} (M_l)_{\alpha\beta} \beta'_{\text{R}} + \text{h.c.} \quad (2.2)$$

with M_ν being symmetric and M_l being arbitrary. Given the unitary matrices O_l and O_ν , M_ν and $M_l M_l^\dagger$ can be diagonalized through the transformations

$$\begin{aligned} O_\nu^\dagger M_\nu O_\nu^* &= D_\nu \equiv \begin{pmatrix} m_1 & 0 & 0 \\ 0 & m_2 & 0 \\ 0 & 0 & m_3 \end{pmatrix}, \\ O_l^\dagger M_l M_l^\dagger O_l &= D_l^2 \equiv \begin{pmatrix} m_e^2 & 0 & 0 \\ 0 & m_\mu^2 & 0 \\ 0 & 0 & m_\tau^2 \end{pmatrix}. \end{aligned} \quad (2.3)$$

Then it is possible to reexpress \mathcal{L}_m in terms of the mass eigenstates of charged leptons and neutrinos:

$$-\mathcal{L}_m = \frac{1}{2} \overline{\nu_{iL}} (D_\nu)_{ij} \nu_{jR}^c + \overline{\alpha_L} (D_l)_{\alpha\beta} \beta_R + \text{h.c.} \quad (2.4)$$

In doing so, one must consistently reexpress \mathcal{L}_{cc} in terms of the relevant mass eigenstates:

$$-\mathcal{L}_{cc} = \frac{g}{\sqrt{2}} \overline{(e \ \mu \ \tau)_L} \gamma^\mu U \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}_L W_\mu^- + \text{h.c.}, \quad (2.5)$$

where $U = O_l^\dagger O_\nu$ is just the unitary PMNS matrix which describes the strength of lepton flavor mixing in weak interactions [†].

In a commonly chosen basis where the flavor eigenstates of three charged leptons are identified with their mass eigenstates, the flavor eigenstates of three neutrinos can be expressed as

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu1} & U_{\mu2} & U_{\mu3} \\ U_{\tau1} & U_{\tau2} & U_{\tau3} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}. \quad (2.6)$$

The nine elements of U can be parameterized in terms of three rotation angles and three CP-violating phases. For example, $U = V P_\nu$ with $V = O_{23} O_\delta O_{13} O_\delta^\dagger O_{12}$ and $P_\nu = \text{Diag} \{e^{i\rho}, e^{i\sigma}, 1\}$, where

$$\begin{aligned} O_{12} &= \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \\ O_{13} &= \begin{pmatrix} c_{13} & 0 & s_{13} \\ 0 & 1 & 0 \\ -s_{13} & 0 & c_{13} \end{pmatrix}, \\ O_{23} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix}, \end{aligned} \quad (2.7)$$

[†] Whether U is really unitary or not actually depends on the mechanism of neutrino mass generation. In the canonical seesaw mechanism [17, 18, 19, 20, 21], for instance, the mixing between light and heavy Majorana neutrinos leads to tiny unitarity-violating effects for U itself. However, the unitarity of U has been tested at the percent level [37, 38], and thus it makes sense to assume U to be exactly unitary for the time being.

and $O_\delta = \text{Diag}\{1, 1, e^{i\delta}\}$ with $c_{ij} \equiv \cos \theta_{ij}$ and $s_{ij} \equiv \sin \theta_{ij}$ (for $ij = 12, 13, 23$). To be more explicit,

$$V = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ V_{\mu 1} & V_{\mu 2} & c_{13}s_{23} \\ V_{\tau 1} & V_{\tau 2} & c_{13}c_{23} \end{pmatrix}, \quad (2.8)$$

in which

$$\begin{aligned} V_{\mu 1} &= -s_{12}c_{23} - c_{12}s_{13}s_{23}e^{i\delta}, \\ V_{\mu 2} &= c_{12}c_{23} - s_{12}s_{13}s_{23}e^{i\delta}, \\ V_{\tau 1} &= s_{12}s_{23} - c_{12}s_{13}c_{23}e^{i\delta}, \\ V_{\tau 2} &= -c_{12}s_{23} - s_{12}s_{13}c_{23}e^{i\delta}. \end{aligned} \quad (2.9)$$

Without loss of generality, the three mixing angles are all arranged to lie in the first quadrant, while δ may vary from 0 to 2π . The fact that the elements of V in its first row and third column are very simple functions of the flavor mixing angles makes the latter have straightforward relations with the amplitudes of solar (θ_{12}), reactor (θ_{13}) and atmospheric (θ_{23}) neutrino oscillations, respectively [39]. In this parametrization δ is usually referred to as the “Dirac” phase, while ρ and σ are the Majorana phases which have nothing to do with neutrino oscillations. If massive neutrinos were the Dirac particles, one would simply forget the Majorana phase matrix P_ν . Throughout this review we mainly concentrate on the Majorana neutrinos, because they are well motivated from a theoretical point of view. Then the symmetric Majorana neutrino mass matrix can be reconstructed in terms of the neutrino masses m_i (for $i = 1, 2, 3$) and the PMNS matrix U in the chosen flavor basis:

$$M_\nu = \begin{pmatrix} M_{ee} & M_{e\mu} & M_{e\tau} \\ M_{e\mu} & M_{\mu\mu} & M_{\mu\tau} \\ M_{e\tau} & M_{\mu\tau} & M_{\tau\tau} \end{pmatrix} = U D_\nu U^T. \quad (2.10)$$

In a specific neutrino mass model which is able to fix the texture of M_ν , one may reversely obtain some testable predictions for the neutrino masses and flavor mixing parameters.

\mathcal{L}_{cc} in Eq. (2.5) tells us that a ν_α neutrino flavor can be produced from the $W^+ + \alpha^- \rightarrow \nu_\alpha$ interaction, and a ν_β neutrino flavor can be detected through the $\nu_\beta + W^- \rightarrow \beta^-$ interaction (for $\alpha, \beta = e, \mu, \tau$). The effective Hamiltonian responsible for the propagation of ν_i in vacuum is expressed as

$$\mathcal{H}_{\text{eff}} = \frac{1}{2E} M_\nu M_\nu^\dagger = \frac{1}{2E} V D_\nu^2 V^\dagger, \quad (2.11)$$

where $E \gg m_i$ is the neutrino beam energy, and the Majorana phase matrix P_ν has been cancelled. Thanks to a quintessentially quantum-mechanical effect, the $\nu_\alpha \rightarrow \nu_\beta$ oscillation happens if the ν_i beam travels a proper distance L . The probability of such a flavor oscillation is given by [40]

$$P(\nu_\alpha \rightarrow \nu_\beta) = \delta_{\alpha\beta} - 4 \sum_{i < j} \text{Re} \Diamond_{\alpha\beta}^{ij} \sin^2 \Delta_{ji}$$

$$+8\text{Im}\diamond_{\alpha\beta}^{ij}\prod_{i<j}\sin\Delta_{ji}, \quad (2.12)$$

where $\Delta_{ji} \equiv \Delta m_{ji}^2 L / (4E)$ and $\diamond_{\alpha\beta}^{ij} \equiv U_{\alpha i} U_{\beta j} U_{\alpha j}^* U_{\beta i}^*$ (for $i, j = 1, 2, 3$ and $\alpha, \beta = e, \mu, \tau$). The unitarity of the PMNS matrix U leads us to

$$\text{Im}\diamond_{\alpha\beta}^{ij} = \mathcal{J} \sum_{\gamma} \epsilon_{\alpha\beta\gamma} \sum_k \epsilon_{ijk}, \quad (2.13)$$

with $\mathcal{J} = c_{12}s_{12}c_{13}^2s_{13}c_{23}s_{23}\sin\delta$ being the so-called Jarlskog invariant [41], which is a universal measure of leptonic CP and T violation in neutrino oscillations. The probability of $\bar{\nu}_\alpha \rightarrow \bar{\nu}_\beta$ oscillations can be directly read off from Eq. (2.12) by making the replacement $U \rightarrow U^*$. There are in general two categories of neutrino oscillation experiments: *appearance* ($\alpha \neq \beta$) and *disappearance* ($\alpha = \beta$). Both the solar neutrino oscillations ($\nu_e \rightarrow \nu_e$) and the reactor antineutrino oscillations ($\bar{\nu}_e \rightarrow \bar{\nu}_e$) are of the disappearance type. In comparison, the atmospheric muon-neutrino (or muon-antineutrino) oscillations are essentially of the disappearance type, and the accelerator neutrino (or antineutrino) oscillations can be of either type.

Given Eq. (2.5), the reactions $\nu_e + e^- \rightarrow \nu_e + e^-$ and $\bar{\nu}_e + e^- \rightarrow \bar{\nu}_e + e^-$ can take place via the charged-current interactions. That is why the behavior of neutrino (or antineutrino) flavor conversion in a dense medium may be modified by the coherent forward $\nu_e e^-$ or $\bar{\nu}_e e^-$ scattering. This matter effect is also referred to as the Mikheyev-Smirnov-Wolfenstein (MSW) effect [42, 43]. In this case Eqs. (2.11)–(2.13) have to be replaced by their counterparts in matter [44].

2.2. Current neutrino oscillation experiments

The fact that neutrinos are massive and lepton flavors are mixed has been firmly established in the past two decades, thanks to a number of solar, atmospheric, reactor and accelerator neutrino (or antineutrino) oscillation experiments [16]. Let us briefly go over some of them in the following, before we discuss what is behind the observed pattern of lepton flavor mixing.

A. Solar neutrino oscillations

The solar ^8B neutrinos were first observed in the Homestake experiment in 1968, but the measured flux was only about one third of the value predicted by the standard solar model (SSM) [14, 45]. This anomaly was later confirmed by other experiments, such as GALLEX/GNO [46, 47], SAGE [48], Super-Kamiokande [49] and SNO [50]. The SNO experiment was particularly crucial because it provided the first model-independent evidence for the flavor conversion of solar ν_e neutrinos into ν_μ and ν_τ neutrinos.

The target material of the SNO detector is heavy water, which allows the solar ^8B neutrinos to be observed via the charged-current (CC) reaction $\nu_e + \text{D} \rightarrow e^- + p + p$, the neutral-current (NC) reaction $\nu_\alpha + \text{D} \rightarrow \nu_\alpha + p + n$ and the elastic-scattering process $\nu_\alpha + e^- \rightarrow \nu_\alpha + e^-$ (for $\alpha = e, \mu, \tau$) [50]. The neutrino fluxes extracted from these three channels can be expressed as $\phi_{\text{CC}} = \phi_e$, $\phi_{\text{NC}} = \phi_e + \phi_{\mu\tau}$ and




Figure 2.1. The allowed region of $\sin^2 2\theta_{13}$ changing with the Dirac phase δ as constrained by the present T2K neutrino oscillation data (curves) and the reactor antineutrino oscillation data (green band). A preliminary hint for $\delta \sim -\pi/2$ can therefore be observed.

$\phi_{\text{ES}} = \phi_e + 0.155\phi_{\mu\tau}$, where $\phi_{\mu\tau}$ denotes a sum of the fluxes of ν_μ and ν_τ neutrinos. If there were no flavor conversion, $\phi_{\mu\tau} = 0$ and $\phi_{\text{CC}} = \phi_{\text{NC}} = \phi_{\text{ES}}$ would hold. The SNO data yielded $\phi_{\text{CC}} = 1.68^{+0.06}_{-0.06}(\text{stat})^{+0.08}_{-0.09}(\text{syst})$, $\phi_{\text{ES}} = 2.35^{+0.22}_{-0.22}(\text{stat})^{+0.15}_{-0.15}(\text{syst}) > \phi_{\text{CC}}$ and $\phi_{\text{NC}} = 4.94^{+0.21}_{-0.21}(\text{stat})^{+0.38}_{-0.34}(\text{syst}) > \phi_{\text{ES}}$ [51], demonstrating the existence of flavor conversion (i.e., $\phi_{\mu\tau} \neq 0$) and supporting the SSM prediction for ϕ_{NC} in a convincing way. In fact, the observed deficit of solar ^8B neutrinos is attributed to $\nu_e \rightarrow \nu_\mu$ and $\nu_e \rightarrow \nu_\tau$ transitions modified by significant matter effects in the core of the Sun. The survival probability of ^8B neutrinos may roughly approximate to $P(\nu_e \rightarrow \nu_e) \simeq \sin^2 \theta_{12} \simeq 0.32$ [52], leading us to $\theta_{12} \simeq 34^\circ$.

Note that the recent Borexino experiment has done a real-time measurement of the mono-energetic solar ^7Be neutrinos and observed a remarkable deficit corresponding to $P(\nu_e \rightarrow \nu_e) = 0.56 \pm 0.1$ [53]. This result can approximately be interpreted as a vacuum oscillation effect, since the low-energy ^7Be neutrino oscillation is not very sensitive to matter effects in the Sun [52]. So one is left with the averaged survival probability $P(\nu_e \rightarrow \nu_e) \simeq 1 - 0.5 \sin^2 2\theta_{12} \simeq 0.56$ for solar ^7Be neutrinos, from which $\theta_{12} \simeq 35^\circ$ can be obtained. Such a result is apparently consistent with the aforementioned value of θ_{12} extracted from the data of solar ^8B neutrinos.

B. Atmospheric neutrino oscillations

It is well known that the atmospheric ν_μ , $\bar{\nu}_\mu$, ν_e and $\bar{\nu}_e$ events are produced in the Earth's atmosphere by cosmic rays, mainly through the decay modes $\pi^+ \rightarrow \mu^+ + \nu_\mu$ with $\mu^+ \rightarrow e^+ + \nu_e + \bar{\nu}_\mu$ and $\pi^- \rightarrow \mu^- + \bar{\nu}_\mu$ with $\mu^- \rightarrow e^- + \bar{\nu}_e + \nu_\mu$. If there were nothing wrong with the atmospheric neutrinos that enter and excite an underground detector, they would have an almost perfect spherical symmetry (i.e., the downward- and upward-going neutrino fluxes should be equal, $\Phi_e(\theta_z) = \Phi_e(\pi - \theta_z)$ and $\Phi_\mu(\theta_z) = \Phi_\mu(\pi - \theta_z)$ with respect to the zenith angle θ_z). In 1998 the Super-Kamiokande Collaboration

observed an approximate up-down flux symmetry for atmospheric ν_e and $\bar{\nu}_e$ events and a significant up-down flux asymmetry for atmospheric ν_μ and $\bar{\nu}_\mu$ events [54]. Such a striking anomaly could naturally be attributed to $\nu_\mu \rightarrow \nu_\tau$ and $\bar{\nu}_\mu \rightarrow \bar{\nu}_\tau$ oscillations for those upward-going ν_μ and $\bar{\nu}_\mu$ events, since the detector itself was insensitive to ν_τ and $\bar{\nu}_\tau$ events. This observation was actually the first *model-independent* discovery of neutrino flavor oscillations, and it marked an important turning point in experimental neutrino physics.

In 2004 the Super-Kamiokande Collaboration did a careful analysis of the disappearance probability of ν_μ and $\bar{\nu}_\mu$ events as a function of the neutrino flight length L over the neutrino energy E , and observed a clear dip in the L/E distribution as the first *direct* evidence for atmospheric neutrino oscillations [55]. Such a dip is consistent with the sinusoidal probability of neutrino flavor oscillations but incompatible with exotic new physics, such as the neutrino decay and neutrino decoherence scenarios.

It is a great challenge to directly observe the atmospheric $\nu_\mu \rightarrow \nu_\tau$ oscillation because this requires the neutrino beam energy greater than a threshold of 3.5 GeV, such that a tau lepton can be produced via the charged-current interaction of incident ν_τ with the target nuclei in the detector. The Super-Kamiokande data are found to be best described by neutrino oscillations that include the ν_τ appearance in addition to the overwhelming signature of the ν_μ disappearance. In particular, a neural network analysis of the zenith-angle distribution of multi-GeV contained events has recently demonstrated the ν_τ appearance effect at the 3.8σ level [56].

C. Accelerator neutrino oscillations

If the atmospheric ν_μ and $\bar{\nu}_\mu$ flavors oscillate, a fraction of the accelerator-produced ν_μ and $\bar{\nu}_\mu$ events may also disappear on their way to a remote detector. This expectation has been confirmed by two long-baseline neutrino oscillation experiments: K2K [57] and MINOS [58]. Both of them observed a reduction of the ν_μ flux and a distortion of the ν_μ energy spectrum, implying $\nu_\mu \rightarrow \nu_\mu$ oscillations. The most amazing result obtained from the atmospheric and accelerator neutrino oscillation experiments is $\sin^2 2\theta_{23} \simeq 1$ or equivalently $\theta_{23} \simeq 45^\circ$, which hints at a likely μ - τ flavor symmetry in the lepton sector.

Today the most important accelerator neutrino oscillation experiment is the T2K experiment, which has discovered the $\nu_\mu \rightarrow \nu_e$ appearance oscillations and carried out a precision measurement of the $\nu_\mu \rightarrow \nu_\mu$ disappearance oscillations. Since its preliminary data were first released in 2011, the T2K experiment has proved to be very successful in establishing the ν_e appearance out of a ν_μ beam at the 7.3σ level and constraining the neutrino mixing parameters θ_{13} , θ_{23} and δ [59, 60, 61]. The point is that the leading term of $P(\nu_\mu \rightarrow \nu_e)$ is sensitive to $\sin^2 2\theta_{13} \sin^2 \theta_{23}$, and its sub-leading term is sensitive to δ and terrestrial matter effects [62]. Fig. 2.1 is an illustration of the allowed region of $\sin^2 2\theta_{13}$ as a function of the CP-violating phase δ , as constrained by the present T2K data [61]. One can observe an unsuppressed value of θ_{13} in this plot, together with a preliminary hint of δ around $-\pi/2$. The latter is also suggestive of a possible μ - τ reflection symmetry, as one will see in section 3.

In comparison with K2K, MINOS and T2K, the accelerator-based OPERA experiment was designed to search for the ν_τ appearance in a ν_μ beam. After several years of data taking, the OPERA Collaboration reported five ν_τ events in 2015. These events have marked a discovery of the $\nu_\mu \rightarrow \nu_\tau$ appearance oscillations with the 5.1σ significance [63].

D. Reactor antineutrino oscillations

Since the first observation of the $\bar{\nu}_e$ events emitted from the Savannah River reactor in 1956 [7], nuclear reactors have been playing a special role in neutrino physics. In particular, θ_{12} and θ_{13} have been measured in the KamLAND [64] and Daya Bay [65, 66] reactor antineutrino experiments, respectively.

Given the average baseline length $L = 180$ km, the KamLAND experiment was sensitive to the Δm_{21}^2 -driven $\bar{\nu}_e \rightarrow \bar{\nu}_e$ oscillations and could accomplish a terrestrial test of the large-mixing-angle MSW solution to the long-standing solar neutrino problem. In fact, it succeeded in doing so in 2003 [64], with an impressive determination of $\theta_{12} \simeq 34^\circ$. A very striking sinusoidal behavior of $P(\bar{\nu}_e \rightarrow \bar{\nu}_e)$ against L/E was also observed by the KamLAND Collaboration later on [67].

The Daya Bay experiment was designed to probe the smallest lepton flavor mixing angle θ_{13} with an unprecedented sensitivity $\sin^2 2\theta_{13} \sim 1\%$ by measuring the Δm_{31}^2 -driven $\bar{\nu}_e \rightarrow \bar{\nu}_e$ oscillation with a baseline length $L \simeq 2$ km. In 2012 the Daya Bay Collaboration announced a 5.2σ discovery of $\theta_{13} \neq 0$ and obtained $\sin^2 2\theta_{13} = 0.092 \pm 0.016(\text{stat}) \pm 0.005(\text{syst})$ [65]. Some similar but less significant results were also achieved in the RENO [68] and Double Chooz [69] reactor antineutrino oscillation experiments.

The Daya Bay experiment has also measured the energy dependence of the $\bar{\nu}_e$ disappearance and seen a nearly full oscillation cycle against L/E [70]. The updated result $\sin^2 2\theta_{13} = 0.090^{+0.008}_{-0.009}$ is obtained in the three-flavor framework. A combination of the Daya Bay measurement of θ_{13} and the T2K measurement of a relatively strong $\nu_\mu \rightarrow \nu_e$ appearance signal [61] drives a slight but intriguing preference for $\delta \sim -\pi/2$, as shown in Fig. 2.1. In addition, the relatively large θ_{13} is so encouraging that the next-generation precision experiments should be able to determine the neutrino mass ordering and the CP-violating phase δ in the foreseeable future [71].

2.3. The observed pattern of the PMNS matrix

In the three-flavor scheme there are six independent parameters which govern the behaviors of neutrino oscillations: two neutrino mass-squared differences Δm_{21}^2 and Δm_{31}^2 , three flavor mixing angles θ_{12} , θ_{13} and θ_{23} , and the Dirac CP-violating phase δ . Those successful atmospheric, solar, accelerator and reactor neutrino oscillation experiments discussed above allow us to determine Δm_{21}^2 , $|\Delta m_{31}^2|$, θ_{12} , θ_{13} and θ_{23} to a good degree of accuracy. The ongoing and future neutrino oscillation experiments are expected to fix the sign of Δm_{31}^2 and pin down the value of δ .

Table 2.1. The best-fit values, together with the 1σ and 3σ intervals, for the six three-flavor neutrino oscillation parameters from a global analysis of current experimental data [72].

Parameter	Best fit	1σ range	3σ range
Normal mass ordering ($m_1 < m_2 < m_3$)			
$\Delta m_{21}^2/10^{-5} \text{ eV}^2$	7.54	7.32 — 7.80	6.99 — 8.18
$\Delta m_{31}^2/10^{-3} \text{ eV}^2$	2.47	2.41 — 2.53	2.26 — 2.65
$\sin^2 \theta_{12}/10^{-1}$	3.08	2.91 — 3.25	2.59 — 3.59
$\sin^2 \theta_{13}/10^{-2}$	2.34	2.15 — 2.54	1.76 — 2.95
$\sin^2 \theta_{23}/10^{-1}$	4.37	4.14 — 4.70	3.74 — 6.26
δ/π	1.39	1.12 — 1.77	0.00 — 2.00
Inverted mass ordering ($m_3 < m_1 < m_2$)			
$\Delta m_{21}^2/10^{-5} \text{ eV}^2$	7.54	7.32 — 7.80	6.99 — 8.18
$\Delta m_{13}^2/10^{-3} \text{ eV}^2$	2.42	2.36 — 2.48	2.22 — 2.60
$\sin^2 \theta_{12}/10^{-1}$	3.08	2.91 — 3.25	2.59 — 3.59
$\sin^2 \theta_{13}/10^{-2}$	2.40	2.18 — 2.59	1.78 — 2.98
$\sin^2 \theta_{23}/10^{-1}$	4.55	4.24 — 5.94	3.80 — 6.41
δ/π	1.31	0.98 — 1.60	0.00 — 2.00




Figure 2.2. A schematic illustration of the fermion mass spectrum of the SM at the electroweak scale, where the three neutrino masses are assumed to have a normal ordering.

A global analysis of the available data on solar (SNO, Super-Kamiokande, Borexino), atmospheric (Super-Kamiokande), accelerator (MINOS, T2K) and reactor (KamLAND, Daya Bay, RENO) neutrino (or antineutrino) oscillations has been done by several groups [72, 73, 74]. Here we quote the main results obtained by Capozzi *et al* [72] in Table 2.1 ‡. Some immediate comments are in order.

‡ The notations $\delta m^2 \equiv m_2^2 - m_1^2$ and $\Delta m^2 \equiv m_3^2 - (m_1^2 + m_2^2)/2$ have been used in Ref. [72]. Their

- The unfixed sign of Δm_{31}^2 implies two possible neutrino mass orderings: normal ($m_1 < m_2 < m_3$) or inverted ($m_3 < m_1 < m_2$). Here “normal” means that the mass ordering of the neutrinos is parallel to that of the charged leptons or the quarks of the same charge (i.e., $m_e \ll m_\mu \ll m_\tau$, $m_u \ll m_c \ll m_t$ and $m_d \ll m_s \ll m_b$, as shown in Fig. 2.2 [75]). A good theoretical reason for $\Delta m_{31}^2 > 0$ or $\Delta m_{13}^2 > 0$ has been lacking §.
- The output values of θ_{13} , θ_{23} and δ in such a global fit are more or less sensitive to the sign of Δm_{31}^2 . That is why it is crucial to determine the neutrino mass hierarchy in the ongoing and upcoming reactor (JUNO [77]), atmospheric (PINGU [78]) and accelerator (NO ν A [79] and LBNE [80]) neutrino oscillation experiments.
- The hint $\delta \neq 0$ (or π) at the 1σ level is preliminary but encouraging, simply because it implies a potential effect of leptonic CP violation which is likely to show up in some long-baseline neutrino oscillation experiments in the foreseeable future. In particular, the best-fit value of δ is quite close to $-\pi/2$, implying an approximate μ - τ reflection symmetry as one can see later on.
- The possibility of $\theta_{23} = \pi/4$ cannot be excluded at the 1σ or 2σ level, and hence a more precise determination of θ_{23} is desirable so as to resolve its octant. Since $\theta_{23} = \pi/4$ is a natural consequence of the μ - τ flavor symmetry in the neutrino sector, the positive or negative deviation of θ_{23} from $\pi/4$ may have profound implications on the structure of the PMNS matrix U and the building of viable neutrino mass models.

In short, the sign of Δm_{31}^2 , the octant of θ_{23} and the value of δ remain unknown. Whether these three open issues are potentially correlated with one another is an intriguing question.

Combining Eqs. (2.8) and (2.9) with Table 2.1, we obtain the remarkable result

$$|U_{\mu 1}| \simeq |U_{\tau 1}|, \quad |U_{\mu 2}| \simeq |U_{\tau 2}|, \quad |U_{\mu 3}| \simeq |U_{\tau 3}| \quad (2.14)$$

to a reasonably good degree of accuracy. This result becomes more transparent when the allowed ranges of the nine PMNS matrix elements are explicitly given at the 3σ level:

$$|U| \simeq \begin{pmatrix} 0.79 - 0.85 & 0.50 - 0.59 & 0.13 - 0.17 \\ 0.19 - 0.56 & 0.41 - 0.74 & 0.60 - 0.78 \\ 0.19 - 0.56 & 0.41 - 0.74 & 0.60 - 0.78 \end{pmatrix} \quad (2.15)$$

in the $\Delta m_{31}^2 > 0$ case; or

$$|U| \simeq \begin{pmatrix} 0.89 - 0.85 & 0.50 - 0.59 & 0.13 - 0.17 \\ 0.19 - 0.56 & 0.40 - 0.73 & 0.61 - 0.79 \\ 0.20 - 0.56 & 0.41 - 0.74 & 0.59 - 0.78 \end{pmatrix} \quad (2.16)$$

relations with Δm_{21}^2 and Δm_{31}^2 are $\Delta m_{21}^2 = \delta m^2$ and $\Delta m_{31}^2 = \Delta m^2 + \delta m^2/2$.

§ If the neutrino mass ordering is finally found to be inverted, one may always reorder it to $m'_1 < m'_2 < m'_3$ by setting $m'_1 = m_3$, $m'_2 = m_1$ and $m'_3 = m_2$, equivalent to a transformation $(\nu_1, \nu_2, \nu_3) \rightarrow (\nu'_2, \nu'_3, \nu'_1)$. In this case the elements of U must be reordered in a self-consistent way: $U \rightarrow U'$, in which $U'_{\alpha 1} = U_{\alpha 3}$, $U'_{\alpha 2} = U_{\alpha 1}$ and $U'_{\alpha 3} = U_{\alpha 2}$ (for $\alpha = e, \mu, \tau$) [76]. Of course, such a reordering treatment does not change any physical content of massive neutrinos.

in the $\Delta m_{31}^2 < 0$ case. In either case the pattern of U is significantly different from that of quark flavor mixing described by the Cabibbo-Kobayashi-Maskawa (CKM) matrix V_{CKM} [81, 82]. The latter is close to the identity matrix because its maximal flavor mixing angle is the Cabibbo angle $\theta_C \simeq 13^\circ$ [16].

In fact, the equality $|U_{\mu i}| = |U_{\tau i}|$ (for $i = 1, 2, 3$) exactly holds if either of the following two sets of conditions can be satisfied [83]:

$$|U_{\mu i}| = |U_{\tau i}| \iff \begin{cases} \theta_{23} = \frac{\pi}{4}, & \theta_{13} = 0; \\ \text{or} \\ \theta_{23} = \frac{\pi}{4}, & \delta = \pm \frac{\pi}{2}. \end{cases} \quad (2.17)$$

The possibility of $\theta_{13} = 0$ has been ruled out, but $\theta_{23} = \pi/4$ and $\delta = -\pi/2$ are both allowed at the 1σ or 2σ level (and $\delta = \pi/2$ is allowed at the 3σ level) as shown in Table 2.1. That is why we claim that there must be an approximate μ - τ flavor symmetry behind the observed pattern of U . In this case the μ - τ symmetry is expected to be a good starting point for model building, no matter what larger flavor groups it belongs to. On the experimental side, it is imperative to measure θ_{23} and δ as accurately as possible, so as to fix the strength of μ - τ symmetry breaking.

At this point it is also worth mentioning that three of the six unitarity triangles of U (the so-called Majorana triangles) in the complex plane, defined by the three orthogonality relations

$$\begin{aligned} \Delta_1 : & U_{e2}U_{e3}^* + U_{\mu 2}U_{\mu 3}^* + U_{\tau 2}U_{\tau 3}^* = 0, \\ \Delta_2 : & U_{e3}U_{e1}^* + U_{\mu 3}U_{\mu 1}^* + U_{\tau 3}U_{\tau 1}^* = 0, \\ \Delta_3 : & U_{e1}U_{e2}^* + U_{\mu 1}U_{\mu 2}^* + U_{\tau 1}U_{\tau 2}^* = 0, \end{aligned} \quad (2.18)$$

will become the *isosceles* triangles provided $\theta_{23} = \pi/4$ and $\delta = \pm\pi/2$ are satisfied. This interesting property, which can be intuitively seen in Fig. 2.3 [84], is another reflection of the μ - τ flavor symmetry as an instructive guideline for the study of leptonic CP violation. We shall go into details of this flavor symmetry and its breaking mechanisms in the subsequent sections.

3. An overview of the μ - τ flavor symmetry

The approximate μ - τ flavor symmetry displayed in the PMNS matrix U is also expected to show up in the mass matrix of either charged leptons or neutrinos. From a model-building point of view, one needs to know what forms of M_l and M_ν can give rise to the observed pattern of lepton flavor mixing as well as the correct mass spectra of (e, μ, τ) and (ν_1, ν_2, ν_3) . Since $U = O_l^\dagger O_\nu$ is a measure of the mismatch between O_l and O_ν , which have been used to diagonalize $M_l M_l^\dagger$ and M_ν in Eq. (2.3), we prefer to work in the basis where O_l equals the identity matrix (i.e., M_l itself is simply a diagonal matrix). In this basis the neutrino mixing effects are completely determined by the structure of M_ν . The latter can be reconstructed in terms of U and the neutrino masses as shown




Figure 2.3. The real shapes and orientations of three Majorana unitarity triangles in the complex plane, plotted by assuming $(\rho, \sigma) = (0, \pi/4)$ and inputting the best-fit values of θ_{12} , θ_{13} , θ_{23} and δ [74] in the normal mass ordering (NMO: left panel) or inverted mass ordering (IMO: right panel) case. The dashed triangles correspond to $(\rho, \sigma) = (0, 0)$ for comparison.

in Eq. (2.10). The six independent elements of M_ν turn out to be

$$M_{\alpha\beta} \equiv \langle m \rangle_{\alpha\beta} = \sum_{i=1}^3 m_i U_{\alpha i} U_{\beta i} , \quad (3.1)$$

where α and β run over e, μ, τ . Of course, M_ν totally consists of nine independent physical parameters.

With the help of the standard parametrization of U in Eqs. (2.7)—(2.9), we express the six effective neutrino mass terms as follows:

$$\begin{aligned} \langle m \rangle_{ee} &= \overline{m}_1 c_{12}^2 c_{13}^2 + \overline{m}_2 s_{12}^2 c_{13}^2 + m_3 \tilde{s}_{13}^{*2} , \\ \langle m \rangle_{\mu\mu} &= \overline{m}_1 (s_{12} c_{23} + c_{12} \tilde{s}_{13} s_{23})^2 \\ &\quad + \overline{m}_2 (c_{12} c_{23} - s_{12} \tilde{s}_{13} s_{23})^2 + m_3 c_{13}^2 s_{23}^2 , \\ \langle m \rangle_{\tau\tau} &= \overline{m}_1 (s_{12} s_{23} - c_{12} \tilde{s}_{13} c_{23})^2 \\ &\quad + \overline{m}_2 (c_{12} s_{23} + s_{12} \tilde{s}_{13} c_{23})^2 + m_3 c_{13}^2 c_{23}^2 , \\ \langle m \rangle_{e\mu} &= -\overline{m}_1 c_{12} c_{13} (s_{12} c_{23} + c_{12} \tilde{s}_{13} s_{23}) \\ &\quad + \overline{m}_2 s_{12} c_{13} (c_{12} c_{23} - s_{12} \tilde{s}_{13} s_{23}) \\ &\quad + m_3 c_{13} \tilde{s}_{13}^* s_{23} , \\ \langle m \rangle_{e\tau} &= \overline{m}_1 c_{12} c_{13} (s_{12} s_{23} - c_{12} \tilde{s}_{13} c_{23}) \\ &\quad - \overline{m}_2 s_{12} c_{13} (c_{12} s_{23} + s_{12} \tilde{s}_{13} c_{23}) \\ &\quad + m_3 c_{13} \tilde{s}_{13}^* c_{23} , \\ \langle m \rangle_{\mu\tau} &= -\overline{m}_1 (s_{12} s_{23} - c_{12} \tilde{s}_{13} c_{23}) (s_{12} c_{23} + c_{12} \tilde{s}_{13} s_{23}) \\ &\quad - \overline{m}_2 (c_{12} s_{23} + s_{12} \tilde{s}_{13} c_{23}) (c_{12} c_{23} - s_{12} \tilde{s}_{13} s_{23}) \\ &\quad + m_3 c_{13}^2 c_{23} s_{23} , \end{aligned} \quad (3.2)$$

where $\overline{m}_1 \equiv m_1 e^{2i\rho}$, $\overline{m}_2 \equiv m_2 e^{2i\sigma}$ and $\tilde{s}_{13} \equiv s_{13} e^{i\delta}$ are defined for the sake of simplicity. The numerical profiles of $|\langle m \rangle_{\alpha\beta}|$ versus the lightest neutrino mass (i.e., m_1 in the normal hierarchy (NH) case or m_3 in the inverted hierarchy (IH) case) are illustrated in Fig. 3.1 [85], where the best-fit values of two neutrino mass-squared differences and three flavor mixing angles presented in Table 2.1 are input while the Dirac and Majorana phases are allowed to vary in the intervals $[0, 2\pi)$ and $[0, \pi)$, respectively. It is obvious that the relations $|\langle m \rangle_{\mu\mu}| \simeq |\langle m \rangle_{\tau\tau}|$ and $|\langle m \rangle_{e\mu}| \simeq |\langle m \rangle_{e\tau}|$ hold in most of the parameter space. Instead of regarding this observation as the reflection of a kind of flavor “anarchy” [86], we treat it seriously as an indication of the μ - τ symmetry which can always be embedded in a much larger flavor symmetry group.

The μ - τ flavor symmetry itself is so powerful that it constrains the texture of M_ν by establishing some equalities or linear relations among its six independent elements. In other words, the μ - τ symmetry has been identified as the *minimal* (and thus most convincing) flavor symmetry in the neutrino sector to give rise to $|\langle m \rangle_{\mu\mu}| = |\langle m \rangle_{\tau\tau}|$ and $|\langle m \rangle_{e\mu}| = |\langle m \rangle_{e\tau}|$. It is therefore expected to be a successful bridge between neutrino phenomenology and model building. In this section we shall mainly describe the salient features of μ - τ permutation and reflection symmetries and outline some typical ways to




Figure 3.1. The profiles of $|\langle m \rangle_{\alpha\beta}|$ versus m_1 (normal hierarchy or NH: red region) or m_3 (inverted hierarchy or IH: green region), where $\delta \in [0, 2\pi)$, $\rho \in [0, \pi)$, $\sigma \in [0, \pi)$, and the best-fit values of Δm_{21}^2 , Δm_{31}^2 , θ_{12} , θ_{13} and θ_{23} listed in Table 2.1 are input.

softly break them. The larger flavor symmetry groups and model-building exercises will be discussed in sections 4 and 5, respectively.

3.1. The μ - τ permutation symmetry

Let us begin with the μ - τ permutation symmetry which has been defined in Eq. (1.1) [87, 88, 89, 90]. Historically, the discussion about this simple flavor symmetry was motivated by the experimental facts $\theta_{23} \sim \pi/4$ and $\sin^2 2\theta_{13} < 0.18$ [91], which point to $\theta_{23} = \pi/4$ and $\theta_{13} = 0$ as an ideal possibility. In this case the PMNS matrix U is greatly simplified and may take the following form in a chosen phase convention:

$$U = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2} c_{12} & \sqrt{2} s_{12} & 0 \\ -s_{12} & c_{12} & -\kappa \\ -\kappa s_{12} & \kappa c_{12} & 1 \end{pmatrix}, \quad (3.3)$$

where $\kappa = \pm 1$ will manifest itself in the corresponding neutrino mass matrix M_ν . The latter can be easily reconstructed as follows:

$$M_\nu = \frac{1}{2} \begin{pmatrix} 2m_{11} & -\sqrt{2} m_{12} & -\kappa\sqrt{2} m_{12} \\ \cdots & m_{22} + m_3 & \kappa(m_{22} - m_3) \\ \cdots & \cdots & m_{22} + m_3 \end{pmatrix}, \quad (3.4)$$

in which

$$\begin{aligned} m_{11} &= \overline{m}_1 c_{12}^2 + \overline{m}_2 s_{12}^2, \\ m_{12} &= (\overline{m}_1 - \overline{m}_2) c_{12} s_{12}, \\ m_{22} &= \overline{m}_1 s_{12}^2 + \overline{m}_2 c_{12}^2. \end{aligned} \quad (3.5)$$

It is clear that the elements of M_ν satisfy the relations $M_{e\mu} = \kappa M_{e\tau}$ and $M_{\mu\mu} = M_{\tau\tau}$. So M_ν is invariant under the μ - τ permutation operation $\nu_\mu \leftrightarrow \kappa \nu_\tau$, which is represented by the transformation matrix

$$S^\pm = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & \kappa \\ 0 & \kappa & 0 \end{pmatrix}, \quad (3.6)$$

where S^\pm corresponds to $\kappa = \pm 1$. So the most general neutrino mass matrix respecting the μ - τ permutation symmetry can be parameterized as

$$M_\nu = \begin{pmatrix} A & B & \kappa B \\ B & C & D \\ \kappa B & D & C \end{pmatrix}, \quad (3.7)$$

where A, B, C and D are in general complex. For simplicity, here these four parameters are all taken to be real [32]. Both $\kappa = \pm 1$ lead to $\theta_{23} = \pi/4$, as the phases or signs of the elements of U can always be rearranged to assure its mixing angles to lie in the first quadrant. This kind of rephasing is implemented by redefining the phases of relevant lepton fields. In the following we shall focus on the $\kappa = +1$ case. Note that the μ - τ permutation symmetry has no definite prediction for θ_{12} . In fact,

$$\tan 2\theta_{12} = \frac{2\sqrt{2} B}{C + D - A}, \quad (3.8)$$

and the three neutrino mass eigenvalues of M_ν are given by

$$m_1 = \frac{1}{2} \left[A + C + D - \sqrt{(A - C - D)^2 + 8B^2} \right],$$

$$m_2 = \frac{1}{2} \left[A + C + D + \sqrt{(A - C - D)^2 + 8B^2} \right],$$

$$m_3 = C - D. \quad (3.9)$$

Depending on the specific values of the relevant free parameters, the neutrino mass spectrum can be either normal ($\Delta m_{31}^2 > 0$) or inverted ($\Delta m_{31}^2 < 0$).

The neutrino mass matrix M_ν in Eq. (3.7) will become more predictive if its free parameters are further constrained. Let us consider two simple but instructive cases for illustration:

- If $C + D - A = B$ is assumed, then Eq. (3.8) leads us to the prediction $\sin \theta_{12} = 1/\sqrt{3}$ or equivalently $\theta_{12} \simeq 35.3^\circ$. In this case we are left with the well-known tri-bimaximal (TB) mixing pattern of massive neutrinos [92, 93]:

$$U_{\text{TB}} = \frac{1}{\sqrt{6}} \begin{pmatrix} 2 & \sqrt{2} & 0 \\ -1 & \sqrt{2} & -\sqrt{3} \\ -1 & \sqrt{2} & \sqrt{3} \end{pmatrix}, \quad (3.10)$$

together with the mass eigenvalues $m_1 = C + D - 2B$, $m_2 = C + D + B$ and $m_3 = C - D$.

- If $C + D - A = 0$ is taken, then Eq. (3.8) yields $\theta_{12} = \theta_{23} = \pi/4$. In this case one arrives at another special pattern of the neutrino mixing matrix — the so-called bi-maximal (BM) flavor mixing pattern [94, 95]:

$$U_{\text{BM}} = \frac{1}{2} \begin{pmatrix} \sqrt{2} & \sqrt{2} & 0 \\ -1 & 1 & -\sqrt{2} \\ -1 & 1 & \sqrt{2} \end{pmatrix}, \quad (3.11)$$

as well as the mass eigenvalues $m_1 = C + D - \sqrt{2}B$, $m_2 = C + D + \sqrt{2}B$ and $m_3 = C - D$.

Considering other constraints on the free parameters of M_ν in Eq. (3.7), one may similarly derive other neutrino mixing patterns which maintain $\theta_{23} = \pi/4$ and $\theta_{13} = 0$ but predict different values of θ_{12} (e.g., the golden-ratio [96] and hexagonal [97, 98] patterns). In such examples all the three flavor mixing angles are constants, which have nothing to do with the neutrino masses. It is possible to link θ_{12} with the ratio of m_1 to m_2 in the following way [99, 100]:

$$\tan \theta_{12} = \sqrt{\frac{m_1}{m_2}}, \quad (3.12)$$

if $A = 0$ (i.e., $\langle m \rangle_{ee} = 0$) holds. On the other hand, taking $C = D$ will lead to $m_3 = 0$, as one can see from Eq. (3.9). Such a special neutrino mass spectrum is not in conflict with the present experimental data, and it may have some interesting consequences in neutrino phenomenology [84].

At this point let us point out that the canonical μ - τ permutation symmetry can be generalized in a way described by the transformation matrix [101]

$$S(\theta, \phi) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos 2\theta & \sin 2\theta e^{-2i\phi} \\ 0 & \sin 2\theta e^{2i\phi} & -\cos 2\theta \end{pmatrix}. \quad (3.13)$$

Obviously, this symmetry will be reduced to the canonical one if $\theta = \pi/4$ and $\phi = 0$ are taken. The requirement of M_ν being invariant under $S(\theta, \phi)$ imposes the following constraints on its elements:

$$\begin{aligned} M_{e\tau} - e^{2i\phi} M_{e\mu} \tan \theta &= 0, \\ (e^{-2i\phi} M_{\tau\tau} - e^{2i\phi} M_{\mu\mu}) + 2M_{\mu\tau} \cos 2\theta &= 0. \end{aligned} \quad (3.14)$$

Such an M_ν will result in a flavor mixing matrix U consisting of $\theta_{13} = 0$ and $\theta_{23} = \theta$. The point is that $S(\theta, \phi)$ has an eigenvector

$$u = \begin{pmatrix} 0 \\ e^{-i\phi} \sin \theta \\ -e^{i\phi} \cos \theta \end{pmatrix} \quad (3.15)$$

with the eigenvalue -1 . Hence the relation $S(\theta, \phi)u = -u$ together with $[S(\theta, \phi)]^\dagger M_\nu [S(\theta, \phi)]^* = M_\nu$ will lead us to $S(\theta, \phi)(M_\nu u^*) = -(M_\nu u^*)$, indicating that $M_\nu u^*$ is proportional to u . One may therefore draw the conclusion that u constitutes one column of U . On the other hand, it is straightforward to show that the reconstructed M_ν via Eq. (2.10) in terms of U featuring $\theta_{13} = 0$ satisfies Eq. (3.14). In other words, M_ν always assumes an $S(\theta, \phi)$ symmetry of the form given in Eq. (3.13) for the case of $\theta_{13} = 0$. But in what follows we shall focus on the canonical μ - τ permutation symmetry, simply because it is much simpler and favored by the experimental result $\theta_{23} \simeq \pi/4$.

Finally, it is worth pointing out that there is a remarkable variant of the μ - τ permutation symmetry — the so-called μ - τ permutation *antisymmetry* [102, 103]. When the latter is concerned, the neutrino mass matrix satisfies the transformation

$$S^+ M_\nu^{\text{AS}} S^+ = -M_\nu^{\text{AS}}, \quad (3.16)$$

and thus it has a special form

$$M_\nu^{\text{AS}} = \begin{pmatrix} 0 & B & -B \\ B & C & 0 \\ -B & 0 & -C \end{pmatrix}. \quad (3.17)$$

The unitary matrix used to diagonalize this special mass matrix is given by

$$U_{\text{AS}} = \frac{1}{2N} \begin{pmatrix} 2B^* & 2B^* & -2C \\ C^* + iN & C^* - iN & 2B \\ C^* - iN & C^* + iN & 2B \end{pmatrix}, \quad (3.18)$$

where $N = \sqrt{2|B|^2 + |C|^2}$. In this case θ_{12} and θ_{23} are both equal to $\pi/4$, and a finite value of θ_{13} is allowed. The three mass eigenvalues of M_ν^{AS} are iN , $-iN$ and 0 , respectively. To fit current neutrino oscillation data, one has to introduce proper perturbations to M_ν^{AS} so as to break its μ - τ permutation antisymmetry and arrive at an acceptable neutrino mass spectrum [103]. Note that an arbitrary Majorana neutrino mass matrix can always be decomposed into two parts: the first part respects the μ - τ permutation symmetry, and the second part possesses the μ - τ permutation antisymmetry. In such a treatment the second part may serve as a perturbation term to characterize the μ - τ symmetry breaking effects. This point will become clearer in section 3.3.

To summarize, the μ - τ permutation symmetry was motivated by the early experimental data of neutrino oscillations and has played an important role in understanding the lepton flavor structures. The discovery of a relatively large value of θ_{13} [66] requires one to go beyond the original model-building scope in this respect, either by taking into account much larger μ - τ symmetry breaking effects [104, 105] or by paying particular attention to the μ - τ *reflection* symmetry. The latter approach is more attractive because it can both produce a nonzero θ_{13} and predict $\delta = \pm\pi/2$ in the first place (i.e., in the symmetry limit).

3.2. The μ - τ reflection symmetry

The μ - τ reflection symmetry was originally put forward by Harrison and Scott [106], who were inspired by the approximate μ - τ universality in the neutrino oscillation data. They started from the possibility of $|U_{\mu i}| = |U_{\tau i}|$ (for $i = 1, 2, 3$), switched off two Majorana phases and arrived at the following parametrization of U in a proper phase convention:

$$U = \begin{pmatrix} u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \\ v_1^* & v_2^* & v_3^* \end{pmatrix}, \quad (3.19)$$

where the elements u_i have been arranged to be real. Thanks to its unitarity, U contains four independent real parameters which can be chosen as u_1 , u_2 , $|v_1|$ and $\gamma_1 \equiv \arg(v_1)$. For instance, the phase difference $\gamma_{21} \equiv \gamma_2 - \gamma_1$ is determined via

$$\cos \gamma_{21} = \frac{-u_2 u_1}{2|v_2 v_1|} = \frac{-u_2 u_1}{\sqrt{(1-u_2^2)(1-u_1^2)}}. \quad (3.20)$$

The Jarlskog invariant [41] of this neutrino mixing pattern turns out to be

$$\begin{aligned} |\mathcal{J}| &= |\text{Im}(u_1 u_2^* v_1^* v_2)| = u_1 u_2 |v_1 v_2 \sin \gamma_{21}| \\ &= \frac{1}{2} u_1 u_2 \sqrt{1-u_1^2-u_2^2} = \frac{1}{2} |U_{e1} U_{e2} U_{e3}|. \end{aligned} \quad (3.21)$$

On the other hand, \mathcal{J} is given by

$$\begin{aligned} |\mathcal{J}| &= c_{12} s_{12} c_{13}^2 s_{13} c_{23} s_{23} |\sin \delta| \\ &= \frac{1}{2} |U_{e1} U_{e2} U_{e3} \sin \delta| \sin 2\theta_{23} \end{aligned} \quad (3.22)$$

in the standard parametrization of U , as pointed out below Eq. (2.13). If $\theta_{13} \neq 0$ or equivalently $U_{e3} \neq 0$, one will immediately obtain $\sin 2\theta_{23} |\sin \delta| = 1$ from Eqs. (3.21) and (3.22). This interesting result implies that the condition $|U_{\mu i}| = |U_{\tau i}|$ must yield $\theta_{23} = \pi/4$ and $\delta = \pm\pi/2$ — the same conclusion has been drawn in Eq. (2.17). Current neutrino oscillation data have given $\theta_{13} \simeq 9^\circ$ [66] and provided a preliminary but noteworthy hint $\delta \sim -\pi/2$ [61]. That is why the special mixing pattern in Eq. (3.19), which is invariant under the μ - τ reflection symmetry transformation (namely, $(S^+ U)^* = U$), is phenomenologically appealing.

Now let us turn to the neutrino mass matrix. It is easy to show that an M_ν obeying the condition

$$(S^+ M_\nu S^+)^* = M_\nu \quad (3.23)$$

can lead us to the neutrino mixing matrix in Eq. (3.19). In other words, M_ν is required to be invariant with respect to the μ - τ reflection transformations defined by Eq. (1.2)[†]. To be explicit, one may parametrize this kind of M_ν as follows:

$$M_\nu = \begin{pmatrix} A & B & B^* \\ B & C & D \\ B^* & D & C^* \end{pmatrix}, \quad (3.24)$$

where A and D are two real parameters.

Note that it is possible to obtain a neutrino mass matrix of the above form without invoking the μ - τ reflection symmetry. In Refs. [109, 110] the authors started from a particular neutrino mass matrix at a seesaw scale,

$$M_\nu^0 = m \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}, \quad (3.25)$$

and allowed it to run down to a much lower scale via the one-loop renormalization-group equations (RGEs) in the supersymmetry framework [111, 112]. Then the resulting neutrino mass matrix at the electroweak scale reads as

$$\begin{aligned} M_\nu &= (I + \Delta^\dagger) M_\nu^0 (I + \Delta^*) \\ &= m \begin{pmatrix} 1 + 2\delta_{ee} & \delta_{e\mu}^* + \delta_{e\tau} & \delta_{e\mu} + \delta_{e\tau}^* \\ \cdots & 2\delta_{\mu\tau} & 1 + \delta_{\mu\mu} + \delta_{\tau\tau} \\ \cdots & \cdots & 2\delta_{\mu\tau}^* \end{pmatrix} \end{aligned} \quad (3.26)$$

with I being the identity matrix and Δ representing the Hermitian radiative correction matrix,

$$\Delta = \begin{pmatrix} \delta_{ee} & \delta_{e\mu} & \delta_{e\tau} \\ \delta_{e\mu}^* & \delta_{\mu\mu} & \delta_{\mu\tau} \\ \delta_{e\tau}^* & \delta_{\mu\tau}^* & \delta_{\tau\tau} \end{pmatrix}. \quad (3.27)$$

It is obvious that the texture of M_ν in Eq. (3.26) is the same as that in Eq. (3.24), but the former as a model-building example crucially depends on the special form of M_ν^0 and the relevant RGEs. We shall subsequently concentrate on the pattern of M_ν in Eq. (3.24) and assume it to result from the μ - τ reflection symmetry.

The correspondence between the flavor mixing matrix in Eq. (3.19) and the neutrino mass matrix in Eq. (3.24) (i.e., the former as a consequence of the latter) was first noticed in Ref. [113]. It can be verified by taking $M_\nu = U D_\nu U^T$ in Eq. (3.23),

$$S^+ U^* D_\nu U^\dagger S^+ = U D_\nu U^T, \quad (3.28)$$

[†] This is actually a kind of generalized CP symmetry [107, 108] — a framework which enables us to combine the flavor symmetry with the CP symmetry in a consistent way. Such a scenario will be discussed in section 4.3.

where D_ν has been defined in Eq. (2.3). Hence S^+U^* is identical to U up to a diagonal phase matrix X — namely, $S^+U^* = UX$, in which X_{ii} is either an arbitrary phase factor for $m_i = 0$ or ± 1 for $m_i \neq 0$. We are therefore led to a particular neutrino mixing matrix of the form

$$U = \begin{pmatrix} \sqrt{X_{11}}u_1 & \sqrt{X_{22}}u_2 & \sqrt{X_{33}}u_3 \\ v_1 & v_2 & v_3 \\ X_{11}v_1^* & X_{22}v_2^* & X_{33}v_3^* \end{pmatrix}. \quad (3.29)$$

It is clear that the relation $|U_{\mu i}| = |U_{\tau i}|$ holds, and this pattern of U will have the same form as the one given in Eq. (3.19) if $X_{ii} = 1$ (for $i = 1, 2, 3$) is taken. Provided X_{ii} happens to be -1 , one may redefine the corresponding neutrino mass eigenstate ν_i as $\nu'_i = i\nu_i$ to absorb the imaginary factor in the first row of U and assure the second and third rows of U to satisfy $U_{\mu i} = U_{\tau i}^*$. In the meantime the mass eigenvalue m_i should be replaced by $m'_i = -m_i$, implying that ν'_i has a Majorana phase equal to $\pi/2$.

In short, as for the Majorana neutrinos, the flavor mixing matrix resulting from the mass matrix M_ν in Eq. (3.24) can also be parameterized as that given by Eq. (3.19), but the relevant Majorana phases are fixed to be 0 or $\pi/2$. One may see this interesting conclusion from another angle. Let us convert M_ν in Eq. (3.24) into a real matrix by a unitary transformation in the (2,3) plane [114]:

$$U_{23}^\dagger M_\nu U_{23}^* = \begin{pmatrix} A & \sqrt{2} \operatorname{Im} B & \sqrt{2} \operatorname{Re} B \\ \cdots & D - \operatorname{Re} C & \operatorname{Im} C \\ \cdots & \cdots & D + \operatorname{Re} C \end{pmatrix} \quad (3.30)$$

with

$$U_{23} = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2} & 0 & 0 \\ 0 & i & 1 \\ 0 & -i & 1 \end{pmatrix}. \quad (3.31)$$

The mass matrix in Eq. (3.30) can subsequently be diagonalized by a real orthogonal matrix O to get real mass eigenvalues, and the resulting neutrino mixing matrix $U = U_{23}O$ takes the form given by Eq. (3.19) with special Majorana phases. To conclude, the μ - τ reflection symmetry can not only predict $\theta_{23} = \pi/4$ and $\delta = \pm\pi/2$ but also constrain the corresponding Majorana phases to be 0 or $\pi/2$.

It is worth stressing that Eq. (3.24) is not the only form of M_ν which is able to generate both $\theta_{23} = \pi/4$ and $\delta = \pm\pi/2$. In fact, the elements of a neutrino mass matrix just need to satisfy the following relation to achieve this goal [102, 115, 116, 117, 118]:

$$\sum_{\alpha=e}^{\tau} M_{e\alpha}^* M_{\alpha\tau} = \sum_{\alpha=e}^{\tau} M_{e\alpha} M_{\alpha\mu}^*. \quad (3.32)$$

There are some other solutions to this equation, besides that given by Eq. (3.24). For example,

$$M_{ee} = M_{\mu\tau}, \quad M_{e\mu} = M_{e\tau}^*, \quad M_{\mu\mu} = M_{\tau\tau}^*, \quad (3.33)$$

leading to a special neutrino mass matrix of the form

$$M_\nu = \begin{pmatrix} A & B & B^* \\ B & C & A \\ B^* & A & C^* \end{pmatrix}. \quad (3.34)$$




Figure 3.2. The profiles of $|\epsilon_1|$ and $|\epsilon_2|$ versus the lightest neutrino mass m_1 (normal hierarchy: red region) or m_3 (inverted hierarchy: green region), where the 1σ ranges of relevant neutrino oscillation parameters listed in Table 2.1 have been input.

Note that this matrix is not equivalent to that given by Eq. (3.24) even if $D = A$ is taken, because here A is a complex parameter instead of a real one. Note also that the present matrix does not possess a transparent flavor symmetry (e.g., the μ - τ reflection symmetry), and thus it is difficult to be derived from an underlying flavor model. That is why we pay particular attention to the pattern of M_ν in Eq. (3.24), which is much more favored from the model-building point of view.

3.3. Breaking of the μ - τ permutation symmetry

Now that the μ - τ permutation symmetry gives rise to $\theta_{13} = 0$ and $\theta_{23} = \pi/4$, the observed result of θ_{13} ($\simeq 9^\circ$) and a possible deviation of θ_{23} from $\pi/4$ definitely signify the breaking of this interesting flavor symmetry. One nevertheless should discuss the symmetry-breaking effects at the mass matrix level, in light of the fact that the symmetry is realized for the mass matrix rather than for the flavor mixing matrix. To quantify the strength of μ - τ symmetry breaking, it is necessary to introduce a characteristic measure of such effects. In fact, the most general perturbation to a Majorana neutrino mass matrix $M_\nu^{(0)}$ with the μ - τ permutation symmetry can be decomposed into a symmetry-conserving part and a symmetry-violating part:

$$\begin{aligned}
 M_\nu^{(1)} = & \frac{1}{2} \begin{pmatrix} 2\delta_{ee} & \delta_{e\mu} + \delta_{e\tau} & \delta_{e\mu} + \delta_{e\tau} \\ \delta_{e\mu} + \delta_{e\tau} & \delta_{\mu\mu} + \delta_{\tau\tau} & 2\delta_{\mu\tau} \\ \delta_{e\mu} + \delta_{e\tau} & 2\delta_{\mu\tau} & \delta_{\mu\mu} + \delta_{\tau\tau} \end{pmatrix} \\
 & + \frac{1}{2} \begin{pmatrix} 0 & \delta_{e\mu} - \delta_{e\tau} & \delta_{e\tau} - \delta_{e\mu} \\ \delta_{e\mu} - \delta_{e\tau} & \delta_{\mu\mu} - \delta_{\tau\tau} & 0 \\ \delta_{e\tau} - \delta_{e\mu} & 0 & \delta_{\tau\tau} - \delta_{\mu\mu} \end{pmatrix}
 \end{aligned} \tag{3.35}$$

whose parameters are small in magnitude. Because the symmetry-conserving part can be absorbed via a redefinition of the original elements of $M_\nu^{(0)}$, we are left with the full

neutrino mass matrix $M_\nu = M_\nu^{(0)} + M_\nu^{(1)}$ in the following form [101]:

$$M_\nu = \begin{pmatrix} A' & B'(1+\epsilon_1) & B'(1-\epsilon_1) \\ B'(1+\epsilon_1) & C'(1+\epsilon_2) & D' \\ B'(1-\epsilon_1) & D' & C'(1-\epsilon_2) \end{pmatrix} \quad (3.36)$$

with

$$\begin{aligned} A' &= A + \delta_{ee}, & B' &= B + \frac{\delta_{e\mu} + \delta_{e\tau}}{2}, \\ D' &= D + \delta_{\mu\tau}, & C' &= C + \frac{\delta_{\mu\mu} + \delta_{\tau\tau}}{2}, \\ \epsilon_1 &= \frac{\delta_{e\mu} - \delta_{e\tau}}{2B'}, & \epsilon_2 &= \frac{\delta_{\mu\mu} - \delta_{\tau\tau}}{2C'}. \end{aligned} \quad (3.37)$$

So it is convenient to use the dimensionless parameters

$$\epsilon_1 = \frac{M_{e\mu} - M_{e\tau}}{M_{e\mu} + M_{e\tau}}, \quad \epsilon_2 = \frac{M_{\mu\mu} - M_{\tau\tau}}{M_{\mu\mu} + M_{\tau\tau}} \quad (3.38)$$

to measure the effects of μ - τ symmetry breaking. If $|\epsilon_1|$ and $|\epsilon_2|$ are both small enough (e.g., $\lesssim 0.2$), then one may argue that the neutrino mass matrix M_ν possesses an approximate μ - τ permutation symmetry.

The small parameters ϵ_1 and ϵ_2 can be linked to the observable quantities through a reconstruction of $M_{\alpha\beta}$ in terms of the neutrino mass and flavor mixing parameters. Note that $|\epsilon_{1,2}|$ are not rephasing-invariant — namely, they are sensitive to the phase transformations

$$\nu_e \rightarrow e^{i\phi_1}\nu_e, \quad \nu_\mu \rightarrow e^{i\phi_2}\nu_\mu, \quad \nu_\tau \rightarrow e^{i\phi_3}\nu_\tau. \quad (3.39)$$

In this case the values of $|\epsilon_{1,2}|$ will accordingly change:

$$\begin{aligned} |\epsilon_1| &\rightarrow \frac{|M_{e\mu} - M_{e\tau}e^{i\phi_{32}}|}{|M_{e\mu} + M_{e\tau}e^{i\phi_{32}}|}, \\ |\epsilon_2| &\rightarrow \frac{|M_{\mu\mu} - M_{\tau\tau}e^{2i\phi_{32}}|}{|M_{\mu\mu} + M_{\tau\tau}e^{2i\phi_{32}}|}, \end{aligned} \quad (3.40)$$

where $\phi_{32} \equiv \phi_3 - \phi_2$. Hence the values of $|\epsilon_{1,2}|$ are not fully physical. Nevertheless, ϕ_{32} is supposed to be strongly suppressed in the presence of an approximate μ - τ permutation symmetry. So one expects that this small phase difference should not change the main features of $|\epsilon_{1,2}|$. Based on this reasonable argument, we neglect ϕ_{32} for the time being and shall take account of it when necessary. Fig. 3.2 illustrates the possible values of $|\epsilon_{1,2}|$ against the lightest neutrino mass, where the relevant neutrino oscillation parameters used to reconstruct $M_{\alpha\beta}$ take values in their 1σ ranges given by Table 2.1. We see that $|\epsilon_1|$ and $|\epsilon_2|$ can be simultaneously small when the neutrino mass spectrum has an inverted hierarchy, and $|\epsilon_1|$ is unacceptably large when m_1 is very small (i.e., when the neutrino mass hierarchy is normal). The numerical results shown in Fig. 3.2 can be understood by using the approximate expressions of $|\epsilon_{1,2}|$ to be given below. Thanks to the smallness of $\theta_{13} \sim 0.15$ and $\Delta\theta_{23} \equiv \theta_{23} - \pi/4 \in [-0.09, +0.09]$, it is

possible to obtain

$$\begin{aligned}\epsilon_1 &\sim -\Delta\theta_{23} - \frac{m_{11}\tilde{\theta}_{13} - m_3\tilde{\theta}_{13}^*}{m_{12}}, \\ \epsilon_2 &\sim 2\Delta\theta_{23} - \frac{4m_{22}\Delta\theta_{23} + 2m_{12}\tilde{\theta}_{13}}{m_{22} + m_3},\end{aligned}\quad (3.41)$$

where the notations introduced in Eqs. (3.2) and (3.5) have been used (in particular, here $\tilde{\theta}_{13} \equiv \theta_{13}e^{i\delta}$). For the sake of simplicity, we consider the properties of $|\epsilon_{1,2}|$ in three typical cases [119].

- $m_1 \ll m_2 \ll m_3$. Given this highly hierarchical neutrino mass spectrum, we obtain

$$|\epsilon_1| \sim \frac{m_3\theta_{13}}{m_2c_{12}s_{12}} \simeq 1.9, \quad |\epsilon_2| \sim 2|\Delta\theta_{23}|, \quad (3.42)$$

implying that M_ν itself does not really exhibit an approximate μ - τ permutation symmetry.

- $m_1 \simeq m_2 \gg m_3$. Because of the near equality of m_1 and m_2 in this case, the Majorana phases ρ and σ become relevant:

$$\begin{aligned}\epsilon_1 &\sim -\Delta\theta_{23} - \frac{c_{12}^2 + s_{12}^2 e^{2i(\sigma-\rho)}}{[1 - e^{2i(\sigma-\rho)}] c_{12}s_{12}} \tilde{\theta}_{13}, \\ \epsilon_2 &\sim -2\Delta\theta_{23} - 2 \frac{[1 - e^{2i(\sigma-\rho)}] c_{12}s_{12}}{c_{12}^2 + s_{12}^2 e^{2i(\sigma-\rho)}} \tilde{\theta}_{13}.\end{aligned}\quad (3.43)$$

Note that the coefficient of $\tilde{\theta}_{13}$ in ϵ_1 is inversely proportional to that in ϵ_2 . So $|\epsilon_1|$ will be much larger than 1 for $\sigma - \rho \sim 0$, and $|\epsilon_2|$ will have a too large value for $\sigma - \rho \sim \pi/2$. A careful analysis indicates that $|\epsilon_1|$ and $|\epsilon_2|$ have no chance to be small enough at the same time.

- $m_1 \simeq m_2 \simeq m_3$. In this case let us consider four sets of special but typical values of ρ and σ . Given $(\rho, \sigma) = (0, 0)$ or $(\pi/2, \pi/2)$, for example, $|\epsilon_1|$ will be much larger than 1 owing to a nearly complete cancellation between the two components of m_{12} as one can see in Eq. (3.5). If $(\rho, \sigma) = (0, \pi/2)$ or $(\pi/2, 0)$, then $\epsilon_{1,2}$ can approximate to

$$\begin{aligned}\epsilon_1 &\sim -\Delta\theta_{23} + \frac{e^{-i\delta} \mp (c_{12}^2 - s_{12}^2) e^{i\delta}}{2c_{12}s_{12}} \theta_{13}, \\ \epsilon_2 &\sim \frac{2(1 \mp c_{12}^2 \pm s_{12}^2) \Delta\theta_{23} \mp 4c_{12}s_{12}\tilde{\theta}_{13}}{1 \pm c_{12}^2 \mp s_{12}^2},\end{aligned}\quad (3.44)$$

which are allowed to be simultaneously small.

We conclude that a quasi-degenerate neutrino mass spectrum allows M_ν to show an approximate μ - τ permutation symmetry. This conclusion can also be put in another way: given $m_1 \simeq m_2 \simeq m_3$, the Majorana neutrino mass matrix with an approximate μ - τ permutation symmetry is able to generate a phenomenologically viable pattern of neutrino mixing — a pattern compatible with current experimental data, as one will

see later on. A similar approach has been discussed in Ref. [120] to explore the gap between U_{TB} and the experimentally-favored pattern as well as what such a gap implies on the underlying texture of M_ν . Because U_{TB} can be obtained by invoking the condition $M_{ee} + M_{e\mu} = M_{\mu\mu} + M_{\mu\tau}$ for M_ν on the basis of the μ - τ permutation symmetry, one needs ϵ_1 , ϵ_2 and an extra small parameter to describe the actual departure of M_ν from $M_\nu^{\text{TB}} \equiv U_{\text{TB}} D_\nu U_{\text{TB}}^T$. Hence a similar conclusion has been reached [120]: only in the $m_1 \simeq m_2 \simeq m_3$ case can M_ν possess an approximate μ - τ symmetry and give rise to a viable neutrino mixing pattern which is very close to the form of U_{TB} . In this sense U_{TB} itself is accidental, at least from a point of view of model building, although its predictions for θ_{12} and θ_{23} are quite close to the experimental values.

Instead of reconstructing M_ν in terms of neutrino masses and flavor mixing parameters to constrain the sizes of $\epsilon_{1,2}$, now we follow a more straightforward way to study the texture of M_ν with an approximate μ - τ permutation symmetry (i.e., both ϵ_1 and ϵ_2 are assumed to be small from the beginning) and explore the direct dependence of $\tilde{\theta}_{13}$ and $\Delta\theta_{23}$ on $\epsilon_{1,2}$. There are two good reasons for doing so: (a) it is important to identify what kind of symmetry breaking can give rise to the viable phenomenological consequences in a model-building exercise; (b) some specific symmetry-breaking patterns are able to predict a few interesting correlations among the physical parameters and thus may bridge the gap between the unknown and known parameters. Let us first derive the expressions of $\tilde{\theta}_{13}$ and $\Delta\theta_{23}$ arising from the most general symmetry breaking pattern. In the standard parametrization the unitary matrix used for diagonalizing a Majorana neutrino mass matrix takes the form $U = P_\phi V P_\nu$, where V and P_ν have been given in Eqs. (2.6)–(2.9), and $P_\phi = \text{Diag}\{e^{i\phi_1}, e^{i\phi_2}, e^{i\phi_3}\}$. Although ϕ_1 , ϕ_2 and ϕ_3 have no physical meaning, they are necessary in the diagonalization process and thus should not be ignored from here. When the μ - τ permutation symmetry is exact, ϕ_2 is automatically equal to ϕ_3 . In the presence of small symmetry-breaking effects, ϕ_{32} becomes a finite but small quantity. By taking $M_\nu = U D_\nu U^T$ in Eq. (3.36) and doing perturbation expansions for those small quantities, one can obtain the following relations which connect ϕ_{32} , $\Delta\theta_{23}$ and $\tilde{\theta}_{13}$ with the perturbation parameters $\epsilon_{1,2}$:

$$\begin{aligned} m_{12}\Delta\theta_{23} + m_{11}\tilde{\theta}_{13} - m_3\tilde{\theta}_{13}^* + \frac{m_{12}}{2}(2\epsilon_1 + i\phi_{32}) &= 0, \\ m_{22-3}\Delta\theta_{23} + m_{12}\tilde{\theta}_{13} + \frac{m_{22+3}}{2}(\epsilon_2 + i\phi_{32}) &= 0, \end{aligned} \quad (3.45)$$

in which $m_{22\pm 3} \equiv m_{22} \pm m_3$ are defined. After solving these equations in a straightforward way, we obtain $\tilde{\theta}_{13}^*$ and $\Delta\theta_{23}$ as functions of $\epsilon_{1,2}$ [101]:

$$\begin{aligned} \tilde{\theta}_{13}^* &= (2\Delta m_{31}^2)^{-1} (2m_3 m_{12} c_{12}^2 \epsilon_1 + 2\bar{m}_1 m_{12}^* c_{12}^2 \epsilon_1^* \\ &\quad + m_3 m_{22+3} c_{12} s_{12} \epsilon_2 + \bar{m}_1 m_{22+3}^* c_{12} s_{12} \epsilon_2^*) \\ &\quad + (2\Delta m_{32}^2)^{-1} (2m_3 m_{12} s_{12}^2 \epsilon_1 + 2\bar{m}_2 m_{12}^* s_{12}^2 \epsilon_1^* \\ &\quad - m_3 m_{22+3} c_{12} s_{12} \epsilon_2 - \bar{m}_2 m_{22+3}^* c_{12} s_{12} \epsilon_2^*) , \\ \Delta\theta_{23} &= \text{Re} \left\{ (2\Delta m_{31}^2)^{-1} [2m_{12} c_{12} s_{12} (\bar{m}_1^* \epsilon_1 + m_3 \epsilon_1^*) \right. \\ &\quad \left. + m_{22+3} s_{12}^2 (\bar{m}_1^* \epsilon_2 + m_3 \epsilon_2^*)] \right\} \end{aligned}$$

$$\begin{aligned}
& - (2\Delta m_{32}^2)^{-1} [2m_{12}c_{12}s_{12} (\overline{m}_2^*\epsilon_1 + m_3\epsilon_1^*) \\
& - m_{22+3}c_{12}^2 (\overline{m}_2^*\epsilon_2 + m_3\epsilon_2^*)] \} .
\end{aligned} \tag{3.46}$$

With the help of these results, one may get a ball-park feeling of the dependence of $\tilde{\theta}_{13}$ and $\Delta\theta_{23}$ on $\epsilon_{1,2}$ in some particular cases to be discussed below. One will see that the values of θ_{13} and $\Delta\theta_{23}$ are strongly correlated with the neutrino mass spectrum when the strength of μ - τ permutation symmetry breaking (i.e., the size of $\epsilon_{1,2}$) is specified.

In the assumption of CP conservation, Eq. (3.46) is simplified to

$$\begin{aligned}
\theta_{13} &= \frac{2m_{12}c_{12}^2\epsilon_1 + m_{22+3}c_{12}s_{12}\epsilon_2}{2(m_3 \mp m_1)} \\
&+ \frac{2m_{12}s_{12}^2\epsilon_1 - m_{22+3}c_{12}s_{12}\epsilon_2}{2(m_3 \mp m_2)} , \\
\Delta\theta_{23} &= \frac{2m_{12}c_{12}s_{12}\epsilon_1 + m_{22+3}s_{12}^2\epsilon_2}{2(m_3 \mp m_1)} \\
&- \frac{2m_{12}c_{12}s_{12}\epsilon_1 - m_{22+3}c_{12}^2\epsilon_2}{2(m_3 \mp m_2)} ,
\end{aligned} \tag{3.47}$$

in which the signs “ \mp ” correspond to $\overline{m}_{1,2} = \pm m_{1,2}$ for $\rho, \sigma = 0$ or $\pi/2$. Some more specific discussions and comments are in order.

(1) As for the neutrino mass spectrum with a vanishingly small m_1 , the expression of θ_{13} can be further simplified to

$$\theta_{13} \sim \frac{1}{2}\sqrt{r} c_{12}s_{12} (2\epsilon_1 - \epsilon_2) \simeq 0.04 (2\epsilon_1 - \epsilon_2) , \tag{3.48}$$

where $r \equiv \Delta m_{21}^2/\Delta m_{31}^2 \simeq 0.03$. Given $|\epsilon_{1,2}| \lesssim 0.2$, it is impossible to get the observed value of θ_{13} from the above expression.

(2) In the $m_1 \simeq m_2 \gg m_3$ case, θ_{13} and $\Delta\theta_{23}$ are sensitive to the combination of ρ and σ ‡. If ρ and σ are equal, θ_{13} will be highly suppressed:

$$\theta_{13} \sim \frac{1}{4}rc_{12}s_{12} (2\epsilon_1 - \epsilon_2) \simeq 0.004 (2\epsilon_1 - \epsilon_2) . \tag{3.49}$$

Otherwise, θ_{13} turns out to be

$$\theta_{13} \sim \frac{1}{2} \cos 2\theta_{12} \sin 2\theta_{12} (2\epsilon_1 - \epsilon_2) , \tag{3.50}$$

which is still unable to fit its experimental value.

(3) When the neutrino mass spectrum is nearly degenerate, one may consider the following four special but typical cases.

- $(\rho, \sigma) = (0, 0)$. In this case θ_{13} approximates to

$$\theta_{13} \sim \frac{2m_0^2}{\Delta m_{31}^2} rc_{12}s_{12}\epsilon_2 , \tag{3.51}$$

where m_0 denotes the overall neutrino mass scale. The factor $m_0^2/\Delta m_{31}^2$ can enhance the value of θ_{13} as m_0 goes up, but $m_0 \leq 0.1$ eV is expected to hold in light of

‡ In the CP-conserving case under consideration, ρ and σ can only take a value of 0 or $\pi/2$.

the present cosmological upper bound on the sum of three neutrino masses (e.g., $m_1 + m_2 + m_3 \simeq 3m_0 < 0.23$ eV at the 95% confidence level [121]). So θ_{13} is at most 0.03. Such a result is certainly unacceptable.

- $(\rho, \sigma) = (0, \pi/2)$. In this case we obtain

$$\begin{aligned}\theta_{13} &\sim \frac{2m_0^2}{\Delta m_{31}^2} c_{12} s_{12} (2c_{12}^2 \epsilon_1 + s_{12}^2 \epsilon_2) , \\ \Delta\theta_{23} &\sim \frac{2m_0^2}{\Delta m_{31}^2} s_{12}^2 (2c_{12}^2 \epsilon_1 + s_{12}^2 \epsilon_2) .\end{aligned}\quad (3.52)$$

Thanks to the enhancement factor $m_0^2/\Delta m_{31}^2$, θ_{13} is easy to reach the measured value. Moreover, there is a correlation between $|\Delta\theta_{23}|$ and θ_{13} , i.e., $|\Delta\theta_{23}| \sim \theta_{13} s_{12}/c_{12} \simeq 6^\circ$. This relatively large $|\Delta\theta_{23}|$ will be tested in the near future.

- $(\rho, \sigma) = (\pi/2, 0)$. In this case the results are

$$\begin{aligned}\theta_{13} &\sim \frac{2m_0^2}{\Delta m_{31}^2} c_{12} s_{12} (2s_{12}^2 \epsilon_1 + c_{12}^2 \epsilon_2) , \\ \Delta\theta_{23} &\sim \frac{2m_0^2}{\Delta m_{31}^2} c_{12}^2 (2s_{12}^2 \epsilon_1 + c_{12}^2 \epsilon_2) .\end{aligned}\quad (3.53)$$

The correlation between $|\Delta\theta_{23}|$ and θ_{13} predicts $|\Delta\theta_{23}| \sim \theta_{13} c_{12}/s_{12} \simeq 13^\circ$, which is too large to be acceptable.

- $(\rho, \sigma) = (\pi/2, \pi/2)$. This special case is disfavored because θ_{13} is extremely suppressed by the factor $\Delta m_{21}^2/m_0^2$, as one can see from

$$\theta_{13} \sim \frac{\Delta m_{21}^2}{4m_0^2} c_{12} s_{12} \epsilon_1 . \quad (3.54)$$

To summarize, in the CP-conserving case considered above only the example of a quasi-degenerate neutrino mass spectrum with $(\rho, \sigma) = (0, \pi/2)$ is still allowed by current experimental data.

When CP violation is taken into account, one has to deal with some more free parameters. But this possibility is certainly more realistic and more interesting, because it is related to the asymmetry between matter and antimatter in a variety of weak interaction processes including neutrino oscillations. Here let us take two typical examples for illustration.

(A) In the first example we assume the Majorana phases to take trivial values (i.e., 0 or $\pi/2$) and the symmetry-breaking parameters to be purely imaginary (i.e., $\epsilon_{1,2} = i|\epsilon_{1,2}|$). Then Eq. (3.46) leads us to the results $\Delta\theta_{23} = 0$ and

$$\begin{aligned}\tilde{\theta}_{13}^* &= i \frac{2m_{12}c_{12}^2|\epsilon_1| + m_{22+3}c_{12}s_{12}|\epsilon_2|}{2(m_3 \pm m_1)} \\ &\quad + i \frac{2m_{12}s_{12}^2|\epsilon_1| - m_{22+3}c_{12}s_{12}|\epsilon_2|}{2(m_3 \pm m_2)} ,\end{aligned}\quad (3.55)$$

which implies $\delta = \pm\pi/2$. These results are actually the same as those predicted by the μ - τ reflection symmetry, simply because the perturbation under consideration is

so special that the overall neutrino mass matrix M_ν as given in Eq. (3.36) respects this flavor symmetry. Note that θ_{13} has the same expressions as those shown in Eqs. (3.48)—(3.50) when m_1 or m_3 is vanishingly small. If the quasi-degenerate neutrino mass spectrum is concerned, then

$$\begin{aligned}
(\rho, \sigma) = (0, 0) : \quad \theta_{13} &\sim \frac{\Delta m_{21}^2}{8m_0^2} c_{12} s_{12} (2|\epsilon_1| - |\epsilon_2|) , \\
(\rho, \sigma) = (0, \pi/2) : \quad \theta_{13} &\sim \frac{2m_0^2}{\Delta m_{31}^2} c_{12} s_{12}^3 (2|\epsilon_1| - |\epsilon_2|) , \\
(\rho, \sigma) = (\pi/2, 0) : \quad \theta_{13} &\sim \frac{2m_0^2}{\Delta m_{31}^2} c_{12}^3 s_{12} (2|\epsilon_1| - |\epsilon_2|) , \\
(\rho, \sigma) = (\pi/2, \pi/2) : \quad \theta_{13} &\sim \frac{r}{2} c_{12} s_{12} (2|\epsilon_1| - |\epsilon_2|)
\end{aligned} \tag{3.56}$$

can be obtained. Similar to the results achieved in the CP-conserving case, an acceptable value of θ_{13} is only obtainable from Eq. (3.56) with $(\rho, \sigma) = (0, \pi/2)$ or $(\pi/2, 0)$. It is worth mentioning that θ_{13} is always proportional to $2|\epsilon_1| - |\epsilon_2|$ in this case, because it is the combination $2\epsilon_1 - \epsilon_2$ that is invariant under the phase transformations in Eq. (3.39) when $\epsilon_{1,2}$ are imaginary.

(B) In the second example we relax ρ and σ to see whether θ_{13} is possible to fit current experimental data in the $m_1 \simeq m_2 \gg m_3$ case, where

$$\begin{aligned}
\theta_{13} &= \sqrt{\cos^2 2\theta_{12} [1 - \cos 2(\sigma - \rho)]^2 + \sin^2 2(\sigma - \rho)} \\
&\quad \times \frac{1}{2} c_{12} s_{12} |2\epsilon_1 - \epsilon_2| .
\end{aligned} \tag{3.57}$$

The maximal value of θ_{13} turns out to be $\theta_{13}^{\max} \simeq 0.25 |2\epsilon_1 - \epsilon_2|$ at $\sigma - \rho = \pi/4$. It is therefore hard to make θ_{13} compatible with the data unless $\epsilon_{1,2}$ have a relative phase of π and take their upper limit ~ 0.2 (for a valid perturbation expansion). In this extreme case $\Delta\theta_{23} \simeq -0.03 \cos[\arg(\epsilon_1)] \lesssim 2^\circ$. Of course, the chance for this extreme case to happen is rather small. So it is fair to say that the $m_3 \rightarrow 0$ limit seems not to be favored by M_ν in Eq. (3.36) even when CP violation is taken into account. Finally, it is worth emphasizing that δ may take any value in the interval $[0, 2\pi)$ regardless of the strength of μ - τ symmetry breaking. This point can be interpreted as follows. The vanishing of θ_{13} and $\Delta\theta_{23}$ is protected by the μ - τ permutation symmetry, so their realistic magnitudes are subject to the small effects of symmetry breaking. In contrast, δ is not well defined when the μ - τ permutation symmetry is exact, and it turns out to have physical meaning only after this symmetry is broken and the resulting neutrino mass matrix $M_\nu = M_\nu^{(0)} + M_\nu^{(1)}$ contains the nontrivial phases. Hence δ is sensitive to the phases of $\epsilon_{1,2}$. As shown in Eq. (3.55), $\delta = \pm\pi/2$ can emerge if $\epsilon_{1,2}$ are purely imaginary. Note that a finite value of δ may arise even when $\epsilon_{1,2}$ are real, if $M_\nu^{(0)}$ itself involves the nontrivial Majorana phases [101]. Here let us take $\epsilon_2 = 2\epsilon_1$ as an example which resembles the symmetry breaking induced by the RGE running effects, as one can see later. In this special case δ is given by

$$\tan \delta = \frac{m_2 \sin 2\sigma - m_1 \sin 2\rho}{m_1 \cos 2\rho - m_2 \cos 2\sigma - m_3 r} , \tag{3.58}$$

which is not directly associated with $\epsilon_{1,2}$ and may be large no matter whether the neutrino mass spectrum exhibits a normal or inverted hierarchy.

The above discussions are subject to the smallness of μ - τ symmetry breaking — namely, the symmetry-breaking terms are relatively small as compared with the entries of $M_\nu^{(0)}$. In a given neutrino mass model the elements of $M_\nu^{(0)}$ may not be at the same order of magnitude, and some of them are even possible to vanish at the tree level. Provided the higher-order contributions break the flavor symmetry, then they may play a dominant role in those originally vanishing entries to make the predictions of the resulting neutrino mass matrix M_ν compatible with current neutrino oscillation data. Of course, there is nothing wrong with this situation, but it motivates us to reconsider the physical implications of an approximate μ - τ permutation symmetry. Though a concrete model may more or less correlate the symmetry-breaking terms appearing in different elements of M_ν , we just treat them as independent parameters in our subsequent analysis. A good example of this kind is a hierarchical neutrino mass matrix which can lead us to the mass spectrum $m_1 < m_2 \ll m_3$. In this case the $\mu\mu$, $\mu\tau$ and $\tau\tau$ entries of M_ν are expected to be much larger than the others in magnitude by a factor $\sim \sqrt{1/r} = \sqrt{\Delta m_{31}^2/\Delta m_{21}^2}$. If these large entries arise from a simple flavor symmetry and those smaller ones come from the symmetry-breaking effects, a typical form of M_ν may be [122]

$$M_\nu = m_0 \begin{pmatrix} d\epsilon & c\epsilon & b\epsilon \\ c\epsilon & 1 + a\epsilon & -1 \\ b\epsilon & -1 & 1 + \epsilon \end{pmatrix}, \quad (3.59)$$

where ϵ denotes a small perturbation (or symmetry-breaking) parameter, and a , b , c and d are all the real coefficients of $\mathcal{O}(1)$. A straightforward calculation yields the flavor mixing angles and neutrino mass eigenvalues as follows:

$$\begin{aligned} \theta_{13} &\simeq \frac{1}{2\sqrt{2}} (b - c) \epsilon, \\ \Delta\theta_{23} &\simeq \frac{1}{4} (a - 1) \epsilon, \\ \tan 2\theta_{12} &\simeq \frac{2\sqrt{2} (b + c)}{a + 1 - 2d}; \end{aligned} \quad (3.60)$$

and

$$\begin{aligned} m_1 &\simeq \frac{1}{4} \epsilon m_0 (2d + a + 1 - \Delta), \\ m_2 &\simeq \frac{1}{4} \epsilon m_0 (2d + a + 1 + \Delta), \\ m_3 &\simeq 2m_0, \end{aligned} \quad (3.61)$$

in which $\Delta = \sqrt{(2d - a - 1)^2 + 8(b + c)^2}$. The small perturbation parameter ϵ is found to be

$$\epsilon \simeq \frac{4\sqrt{r}}{\sqrt{\Delta} (2d + a + 1)}. \quad (3.62)$$

We see that θ_{13} and $\Delta\theta_{23}$ are proportional to $b - c$ and $a - 1$, respectively. Moreover, θ_{13} is about $\sqrt{r} = \sqrt{\Delta m_{21}^2 / \Delta m_{31}^2}$ up to an $\mathcal{O}(1)$ factor, in agreement with the experimental result. So the smallness of θ_{13} is connected to the hierarchy between $\sqrt{\Delta m_{21}^2}$ and $\sqrt{\Delta m_{31}^2}$ in this scenario.

If $m_1 \simeq m_2 \gg m_3$ holds, M_ν is expected to exhibit a different hierarchical structure, in which the ee , $\mu\mu$, $\mu\tau$ and $\tau\tau$ entries should be much larger than the others. To illustrate, a simple but instructive example of this kind is

$$M_\nu = m_0 \begin{pmatrix} 2 + d\epsilon & c\epsilon & b\epsilon \\ c\epsilon & 1 + a\epsilon & 1 \\ b\epsilon & 1 & 1 + \epsilon \end{pmatrix}. \quad (3.63)$$

The results for three flavor mixing angles are similar to those given in Eq. (3.60), but the neutrino mass eigenvalues turn out to be

$$\begin{aligned} m_1 &\simeq 2m_0 + \frac{1}{4}\epsilon m_0 (2d + a + 1 - \Delta), \\ m_2 &\simeq 2m_0 + \frac{1}{4}\epsilon m_0 (2d + a + 1 + \Delta), \\ m_3 &\simeq \frac{1}{2}\epsilon m_0 (a + 1). \end{aligned} \quad (3.64)$$

In this case $\epsilon \simeq 2r/\Delta$, leading to a high suppression of θ_{13} and $\Delta\theta_{23}$. The point is that m_1 is equal to m_2 at the leading order, and their degeneracy is lifted at the next-to-leading order. So ϵ should be at the order of $(m_2 - m_1)/(m_2 + m_1) \sim 0.01$. To get around this problem, one may assume $a = -1$, $b = -c$ and $d = 0$, such that $r = \Delta = 0$ holds and ϵ may not be strongly suppressed. In this situation it is possible to obtain an appreciable value of θ_{13} , but lifting the degeneracy of m_1 and m_2 requires a higher-order perturbation.

Last but not least, there is a well-known form of M_ν which yields an inverted neutrino mass spectrum with $m_1 = -m_2$ at the tree level:

$$M_\nu = m_0 \begin{pmatrix} e\epsilon & 1 + d\epsilon & 1 + c\epsilon \\ 1 + d\epsilon & b\epsilon & \epsilon \\ 1 + c\epsilon & \epsilon & a\epsilon \end{pmatrix}, \quad (3.65)$$

in which the most general next-to-leading-order terms have been included. This particular texture of M_ν can be attributed to an $L_e - L_\mu - L_\tau$ flavor symmetry [123] — namely, the leading-order entries have a vanishing $L_e - L_\mu - L_\tau$ charge but the others do not. After a straightforward calculation, the phenomenological consequences of this neutrino mass matrix can be summarized as follows:

$$\begin{aligned} \theta_{13} &\simeq \frac{(a - b)\epsilon}{2\sqrt{2}}, \\ \Delta\theta_{23} &\simeq \frac{(c - d)\epsilon}{2}, \\ \tan 2\theta_{12} &\simeq \frac{4\sqrt{2}}{(2 + a + b - 2e)\epsilon}; \end{aligned} \quad (3.66)$$

and

$$\begin{aligned} m_1 &\simeq \frac{1}{4} \left[(2e + 2 + a + b) \epsilon - 4\sqrt{2} \right] m_0 , \\ m_2 &\simeq \frac{1}{4} \left[(2e + 2 + a + b) \epsilon + 4\sqrt{2} \right] m_0 , \\ m_3 &\simeq \frac{1}{2} (a + b - 2) \epsilon m_0 . \end{aligned} \quad (3.67)$$

Similar to the previous case, the symmetry-breaking parameter ϵ has to be exceedingly small unless $a = -b$ and $e = -1$ are assumed. But such an assumption will lead to $\tan 2\theta_{12} \simeq \sqrt{2}/\epsilon$, implying an unacceptably large value of θ_{12} for $\epsilon \lesssim 0.2$. Namely, the pattern of M_ν in Eq. (3.65) is not favored by current experimental data. We have certainly assumed CP conservation in the above examples. When CP violation is taken into account, some of the model parameters become complex [124] and the procedure for diagonalizing a complex and hierarchical neutrino mass matrix can be found in Ref. [125].

3.4. Breaking of the μ - τ reflection symmetry

Let us proceed to discuss possible ways of breaking the μ - τ reflection symmetry. Without involving any model details, the most general correction to a neutrino mass matrix with the μ - τ reflection symmetry is in the form as given by Eq. (3.35). It can be divided into two parts: the part respecting the μ - τ reflection symmetry and the part violating this interesting symmetry. Namely,

$$\begin{aligned} M_\nu^{(1)} &= \frac{1}{2} \begin{pmatrix} 2\text{Re}\delta_{ee} & \delta_{e\mu} + \delta_{e\tau}^* & \delta_{e\mu}^* + \delta_{e\tau} \\ \delta_{e\mu} + \delta_{e\tau}^* & \delta_{\mu\mu} + \delta_{\tau\tau}^* & 2\text{Re}\delta_{\mu\tau} \\ \delta_{e\mu}^* + \delta_{e\tau} & 2\text{Re}\delta_{\mu\tau} & \delta_{\mu\mu}^* + \delta_{\tau\tau} \end{pmatrix} \\ &+ \frac{1}{2} \begin{pmatrix} 2i\text{Im}\delta_{ee} & \delta_{e\mu} - \delta_{e\tau}^* & \delta_{e\tau} - \delta_{e\mu}^* \\ \delta_{e\mu} - \delta_{e\tau}^* & \delta_{\mu\mu} - \delta_{\tau\tau}^* & 2i\text{Im}\delta_{\mu\tau} \\ \delta_{e\tau} - \delta_{e\mu}^* & 2i\text{Im}\delta_{\mu\tau} & \delta_{\tau\tau} - \delta_{\mu\mu}^* \end{pmatrix} \end{aligned} \quad (3.68)$$

in a way similar to Eq. (3.35). As the first part of $M_\nu^{(1)}$ can be absorbed into the original neutrino mass matrix given in Eq. (3.24), which is now defined as $M_\nu^{(0)}$ in the μ - τ reflection symmetry limit, the full neutrino mass matrix $M_\nu = M_\nu^{(0)} + M_\nu^{(1)}$ can be parametrized as

$$M_\nu = \begin{pmatrix} \hat{A}(1 + i\epsilon_3) & \hat{B}(1 + \epsilon_1) & \hat{B}^*(1 - \epsilon_1^*) \\ \cdots & \hat{C}(1 + \epsilon_2) & \hat{D}(1 + i\epsilon_4) \\ \cdots & \cdots & \hat{C}^*(1 - \epsilon_2^*) \end{pmatrix} \quad (3.69)$$

with

$$\begin{aligned} \hat{A} &= A + \text{Re}\delta_{ee} , \quad \hat{B} = B + \frac{\delta_{e\mu} + \delta_{e\tau}^*}{2} , \\ \hat{D} &= D + \text{Re}\delta_{\mu\tau} , \quad \hat{C} = C + \frac{\delta_{\mu\mu} + \delta_{\tau\tau}^*}{2} ; \end{aligned} \quad (3.70)$$

and

$$\begin{aligned}
\epsilon_1 &= \frac{M_{e\mu} - M_{e\tau}^*}{M_{e\mu} + M_{e\tau}^*} = \frac{\delta_{e\mu} - \delta_{e\tau}^*}{2\widehat{B}} , \\
\epsilon_2 &= \frac{M_{\mu\mu} - M_{\tau\tau}^*}{M_{\mu\mu} + M_{\tau\tau}^*} = \frac{\delta_{\mu\mu} - \delta_{\tau\tau}^*}{2\widehat{C}} , \\
\epsilon_3 &= \frac{\text{Im}M_{ee}}{\text{Re}M_{ee}} = \frac{\text{Im}\delta_{ee}}{\widehat{A}} , \\
\epsilon_4 &= \frac{\text{Im}M_{\mu\tau}}{\text{Re}M_{\mu\tau}} = \frac{\text{Im}\delta_{\mu\tau}}{\widehat{D}} .
\end{aligned} \tag{3.71}$$

Hence these dimensionless quantities characterize the strength of μ - τ reflection symmetry breaking. They should be small (e.g., $\lesssim 0.2$) so that M_ν in Eq. (3.69) may possess an approximate μ - τ reflection symmetry. Note that rephasing the neutrino fields actually allows us to remove $\epsilon_{3,4}$, so we shall omit $\epsilon_{3,4}$ without loss of generality in the following discussions. As pointed out in section 3.2, the μ - τ reflection symmetry of a neutrino mass matrix $M_\nu^{(0)}$ requires the corresponding neutrino mixing matrix $U^{(0)} = P_\phi^{(0)} V^{(0)} P_\nu^{(0)}$ to have a special pattern in which $\theta_{23}^{(0)} = \pi/4$, $\delta^{(0)} = \pm\pi/2$, $\rho^{(0)} = 0$ or $\pi/2$ and $\sigma^{(0)} = 0$ or $\pi/2$ hold. Moreover, the three phases of $P_\phi^{(0)}$ have to satisfy $\phi_1^{(0)} = \phi_2^{(0)} + \phi_3^{(0)} = 0$ although they have no physical meaning. When this flavor symmetry is slightly broken, the resulting flavor mixing parameters will depart from the above values. Let us define the relevant deviations as $\Delta\phi_1 \equiv \phi_1 - 0$, $\Delta\phi \equiv (\phi_2 + \phi_3)/2 - 0$, $\Delta\theta_{23} \equiv \theta_{23} - \pi/4$, $\Delta\delta \equiv \delta - \delta^{(0)}$, $\Delta\rho \equiv \rho - \rho^{(0)}$ and $\Delta\sigma \equiv \sigma - \sigma^{(0)}$, and suppose that all of them are small quantities governed by $\epsilon_{1,2}$. The explicit relations of these quantities with $\epsilon_{1,2}$ can be established by doing a perturbation expansion for $M_\nu = U D_\nu U^T$. After some calculations, one arrives at

$$\begin{aligned}
&\overline{m}_1 c_{12}^2 \Delta\rho + \overline{m}_2 s_{12}^2 \Delta\sigma = -m_{11} \Delta\phi_1 , \\
&\overline{m}_1 s_{12}^2 \Delta\rho + \overline{m}_2 c_{12}^2 \Delta\sigma - 2m_{12} \tilde{s}_{13} \Delta\theta_{23} = -m_{22-3} \Delta\phi , \\
&\overline{m}_1 c_{12} (i s_{12} + c_{12} \tilde{s}_{13}) \Delta\rho - \overline{m}_2 s_{12} (i c_{12} - s_{12} \tilde{s}_{13}) \Delta\sigma \\
&\quad - \frac{1}{2} (m_{12} + i m_{11+3} \tilde{s}_{13}) \Delta\theta_{23} + \frac{1}{2} m_{11-3} \tilde{s}_{13} \Delta\delta \\
&\quad = \frac{1}{2} (m_{12} - i m_{11+3} \tilde{s}_{13}) (\epsilon_1 - i \Delta\phi_1 - i \Delta\phi) , \\
&\overline{m}_1 s_{12} (i s_{12} + 2 c_{12} \tilde{s}_{13}) \Delta\rho + \overline{m}_2 c_{12} (i c_{12} - 2 s_{12} \tilde{s}_{13}) \Delta\sigma \\
&\quad - m_{22-3} \Delta\theta_{23} + m_{12} \tilde{s}_{13} \Delta\delta \\
&\quad = \frac{1}{2} (m_{22+3} - 2 i m_{12} \tilde{s}_{13}) (\epsilon_2 - 2 i \Delta\phi) ,
\end{aligned} \tag{3.72}$$

where $m_{11\pm 3}$ and $m_{22\pm 3}$ stand for $m_{11} \pm m_3$ and $m_{22} \pm m_3$, respectively. Moreover, \overline{m}_1 (or \overline{m}_2) and \tilde{s}_{13} are equal to $\pm m_1$ (or $\pm m_2$) and $\pm s_{13}$, respectively, for $\rho^{(0)} = 0$ or $\pi/2$ (or $\sigma^{(0)} = 0$ or $\pi/2$) and $\delta^{(0)} = \pm\pi/2$. Note that the above four equations actually correspond to the conditions of μ - τ reflection symmetry breaking defined in Eq. (3.71). After solving these equations, one will see the clear dependence of $\Delta\theta_{23}$, $\Delta\delta$, $\Delta\rho$ and $\Delta\sigma$ on the perturbation parameters $R_{1,2} \equiv \text{Re}(\epsilon_{1,2})$ and $I_{1,2} \equiv \text{Im}(\epsilon_{1,2})$. For clarity, we

express their results in the following parametrizations:

$$\begin{aligned}
\Delta\theta_{23} &= c_{r1}^\theta R_1 + c_{i1}^\theta I_1 + c_{r2}^\theta R_2 + c_{i2}^\theta I_2 , \\
\Delta\delta &= c_{r1}^\delta R_1 + c_{i1}^\delta I_1 + c_{r2}^\delta R_2 + c_{i2}^\delta I_2 , \\
\Delta\rho &= c_{r1}^\rho R_1 + c_{i1}^\rho I_1 + c_{r2}^\rho R_2 + c_{i2}^\rho I_2 , \\
\Delta\sigma &= c_{r1}^\sigma R_1 + c_{i1}^\sigma I_1 + c_{r2}^\sigma R_2 + c_{i2}^\sigma I_2 .
\end{aligned} \tag{3.73}$$

The lengthy expressions of these 16 coefficients are listed in the Appendix for reference.

Since the Majorana phases cannot be pinned down in a foreseeable future, we only discuss the properties of $\Delta\theta_{23}$ and $\Delta\delta$ in several typical schemes regarding the neutrino mass spectrum.

(1) As for $m_1 \ll m_2 \ll m_3$, the expressions of $\Delta\theta_{23}$ and $\Delta\delta$ can approximate to

$$\begin{aligned}
\Delta\theta_{23} &\simeq (2m_3^2)^{-1} (-2m_2^2 c_{12}^2 s_{12}^2 R_1 - 4m_2^2 c_{12} s_{12}^3 \tilde{s}_{13} I_1 \\
&\quad + m_3^2 R_2 - \bar{m}_2 m_3 c_{12} s_{12} \tilde{s}_{13} I_2) , \\
\Delta\delta &\simeq (2\bar{m}_2 m_3 c_{12} s_{12} \tilde{s}_{13})^{-1} [m_2^2 c_{12}^2 s_{12}^2 (2R_1 + R_2) \\
&\quad + m_3^2 (c_{12}^2 - s_{12}^2) s_{13}^2 (2R_1 - R_2)] \\
&\quad - 2s_{12}^2 I_1 - (2\bar{m}_2)^{-1} m_3 I_2 .
\end{aligned} \tag{3.74}$$

In this case the largest coefficient for $\Delta\theta_{23}$ is $|c_{r2}^\theta| \simeq 0.5$. On the other hand, $\Delta\theta_{23}$ is almost insensitive to R_1 and $I_{1,2}$. In contrast, δ is more sensitive to the symmetry breaking because of an enhancement factor $1/s_{13}$. Numerically, the coefficients $|c_{r1,i1,i2}^\delta|$ (or $|c_{r2}^\delta|$) take values of $\mathcal{O}(1)$ (or $\mathcal{O}(0.1)$) around $m_1 \sim 0.001$ eV.

(2) The results in the $m_1 \simeq m_2 \gg m_3$ case strongly depend on the values of $\rho^{(0)}$ and $\sigma^{(0)}$. When these two phases are equal, one will have

$$\begin{aligned}
\Delta\theta_{23} &\simeq -s_{13}^2 R_1 - r c_{12} s_{12} \tilde{s}_{13} \left(I_1 - \frac{3}{4} I_2 \right) - \frac{1}{2} R_2 , \\
\Delta\delta &\simeq (4c_{12} s_{12} \tilde{s}_{13})^{-1} [(c_{12}^2 - s_{12}^2) s_{13}^2 (2R_1 + R_2) \\
&\quad - r c_{12}^2 s_{12}^2 (2R_1 - R_2)] + I_1 - \frac{3}{4} I_2 ,
\end{aligned} \tag{3.75}$$

where $r \equiv \Delta m_{21}^2 / \Delta m_{31}^2 \simeq 0.03$. Among the relevant coefficients, $|c_{r2}^\theta|$ and $|c_{i1,i2}^\delta|$ are of $\mathcal{O}(1)$, and the others are much smaller. If $\rho^{(0)}$ and $\sigma^{(0)}$ take different values, such as $\rho^{(0)} = 0$ and $\sigma^{(0)} = \pi/2$, the expressions of $\Delta\theta_{23}$ and $\Delta\delta$ turn out to be

$$\begin{aligned}
\Delta\theta_{23} &\simeq -4c_{12}^2 s_{12}^2 R_1 - (c_{12}^2 - s_{12}^2) c_{12} s_{12} \tilde{s}_{13} (4I_1 - I_2) \\
&\quad - \frac{1}{2} (c_{12}^2 - s_{12}^2)^2 R_2 , \\
\Delta\delta &\simeq (r c_{12} s_{12})^{-1} [c_{12} s_{12} (c_{12}^2 - s_{12}^2) (4I_1 - I_2) \\
&\quad - \tilde{s}_{13} (2R_1 + R_2)] .
\end{aligned} \tag{3.76}$$

In this case $|c_{r1}^\theta| \sim 1$ is the largest coefficient for $\Delta\theta_{23}$ while the other three are smaller by at least one order of magnitude. δ is very unstable against the symmetry breaking as its coefficients can easily exceed 10 due to the enhancement factor $1/r$.

(3) When the mass spectrum $m_1 \simeq m_2 \simeq m_3$ is concerned, the effects of μ - τ reflection symmetry breaking in M_ν may be significantly magnified, as one can see from the corresponding results of $\Delta\theta_{23}$ and $\Delta\delta$. In the $\rho^{(0)} = \sigma^{(0)} = 0$ case one obtains

$$\begin{aligned}\Delta\theta_{23} &\simeq 2\frac{m_1^2}{\Delta m_{31}^2} [2s_{13}^2 R_1 - rc_{12}s_{12}\tilde{s}_{13} (2I_1 - I_2) + R_2], \\ \Delta\delta &\simeq (c_{12}s_{12}\tilde{s}_{13})^{-1} \left\{ [rc_{12}^2s_{12}^2 + (c_{12}^2 - s_{12}^2)s_{13}^2] R_1 \right. \\ &\quad + 2\frac{m_1^2}{\Delta m_{31}^2} [rc_{12}^2s_{12}^2 - (c_{12}^2 - s_{12}^2)s_{13}^2] R_2 \\ &\quad \left. - \frac{m_1^2}{\Delta m_{31}^2} (c_{12}^2 - s_{12}^2) (2I_1 - I_2) \right\}. \end{aligned} \quad (3.77)$$

Given $m_1 \sim 0.1$ eV, for example, $|c_{r2}^\theta|$ can reach 6 while $|c_{r1,i1,i2}^\theta|$ are much smaller. $\Delta\delta$ is highly sensitive to $I_{1,2}$ whose relevant coefficients are of $\mathcal{O}(10)$, but it is insensitive to $R_{1,2}$ whose relevant coefficients are of $\mathcal{O}(0.1)$. On the other hand, $(\rho^{(0)}, \sigma^{(0)}) = (0, \pi/2)$ will lead $\Delta\theta_{23}$ and $\Delta\delta$ to

$$\begin{aligned}\Delta\theta_{23} &\simeq 2\frac{m_1^2}{\Delta m_{31}^2} [2c_{12}^2s_{12}^2R_1 + 2(c_{12}^2 - s_{12}^2)c_{12}s_{12}\tilde{s}_{13}I_1 \\ &\quad + s_{12}^4R_2 + c_{12}s_{12}^3\tilde{s}_{13}I_2], \\ \Delta\delta &\simeq \frac{m_1^2}{\Delta m_{21}^2} \left[8\frac{m_1^2}{\Delta m_{31}^2} c_{12}s_{12}\tilde{s}_{13} (2c_{12}^2R_1 - s_{12}^2R_2) \right. \\ &\quad \left. + 4(c_{12}^2 - s_{12}^2)I_1 + 2c_{12}^2I_2 \right]. \end{aligned} \quad (3.78)$$

In this case $|c_{r1,r2}^\theta|$ and $|c_{i1,i2}^\theta|$ are respectively of $\mathcal{O}(1)$ and $\mathcal{O}(0.1)$, and all the coefficients associated with $\Delta\delta$ may be enhanced to 100.

In short, $\Delta\delta$ and $\Delta\theta_{23}$ tend to be enlarged when the degeneracy of the neutrino masses grows and $\rho^{(0)}$ takes a different value from $\sigma^{(0)}$. Furthermore, δ is in general more unstable than θ_{23} with respect to the effects of μ - τ reflection symmetry breaking. In particular, the coefficients associated with $\Delta\delta$ can reach a value about 100 provided $m_1 \simeq m_2 \simeq m_3$ and $\rho^{(0)} \neq \sigma^{(0)}$. Furthermore, $\Delta\theta_{23}$ is more sensitive to $R_{1,2}$ while $\Delta\delta$ is more sensitive to $I_{1,2}$. With the help of such observations, one may break the μ - τ reflection symmetry in a proper way whenever necessary in a model-building exercise.

3.5. RGE-induced μ - τ symmetry breaking effects

If a certain flavor symmetry is associated with the neutrino mass generation and lepton flavor mixing at a superhigh energy scale Λ_{FS} , it will be broken due to the RGE evolution of M_ν from Λ_{FS} down to the electroweak scale $\Lambda_{\text{EW}} \sim 10^2$ GeV. In this case the significant difference between m_μ and m_τ is just a source of symmetry breaking which can be transmitted to M_ν via the RGE running effects. From a model-building point of view, it is in general natural to introduce a kind of flavor symmetry at an energy scale much higher than Λ_{EW} (e.g., the seesaw scale). So the RGE-triggered symmetry breaking effects should be taken into account when such a model is confronted with the available experimental data at low energies [126]. It is therefore worthwhile to explore how the

μ - τ flavor symmetry is broken via the RGEs. At the one-loop level the RGE running behavior of M_ν is described by [127, 128, 129, 130]

$$\frac{dM_\nu}{dt} = C \left(Y_l^\dagger Y_l \right)^T M_\nu + C M_\nu \left(Y_l^\dagger Y_l \right) + \alpha M_\nu \quad (3.79)$$

with $t \equiv (1/16\pi^2) \ln(\mu/\Lambda_{\text{FS}})$, in which μ denotes an energy scale between Λ_{EW} and Λ_{FS} , $C = 1$ and

$$\alpha \simeq -\frac{6}{5}g_1^2 - 6g_2^2 + 6y_t^2 \quad (3.80)$$

within the minimal supersymmetry standard model (MSSM). In Eq. (3.79) the α -term just provides an overall rescaling factor which will be referred to as I_α , while the other two terms may change the structure of M_ν . In the basis chosen for Eq. (3.79), the Yukawa coupling matrix of three charged leptons is diagonal: $Y_l = \text{Diag}\{y_e, y_\mu, y_\tau\}$. Because of $y_e \ll y_\mu \ll y_\tau$, it is reasonable to neglect the contributions of both y_e and y_μ in the subsequent discussions. So we are left with $y_\tau^2 = (1 + \tan^2 \beta) m_\tau^2/v^2$ in the MSSM with $v \simeq 246$ GeV being the vacuum expectation value of the Higgs fields. There are normally two ways to proceed: a) one may follow the method described in Refs. [131, 132, 133] to derive the differential RGEs of neutrino masses and flavor mixing parameters from Eq. (3.79); b) one may first integrate Eq. (3.79) to arrive at the RGE-corrected neutrino mass matrix at Λ_{EW} and then diagonalize the latter to obtain neutrino masses and flavor mixing parameters. Here we follow the second approach so as to see how M_ν is modified by the RGE running effects in a transparent way. After integrating Eq. (3.79), we get [134, 135]

$$M'_\nu = I_\alpha I_\tau^\dagger M_\nu I_\tau^* \quad (3.81)$$

at Λ_{EW} , where $I_\tau \simeq \text{Diag}\{1, 1, 1 - \Delta_\tau\}$ and

$$I_\alpha = \exp \left(\int_{\Lambda_{\text{FS}}}^{\Lambda_{\text{EW}}} \alpha dt \right), \quad \Delta_\tau = \int_{\Lambda_{\text{EW}}}^{\Lambda_{\text{FS}}} y_\tau^2 dt. \quad (3.82)$$

Given $\Lambda_{\text{FS}} \sim 10^{14}$ GeV, for example, I_α changes from 0.9 to 0.8 and Δ_τ changes from 0.002 to 0.044 when $\tan \beta$ varies from 10 to 50.

Now let us consider a specific neutrino mass matrix respecting the μ - τ permutation symmetry at Λ_{FS} , such as the one given in Eq. (3.4) with $\kappa = +1$. With the help of Eq. (3.81), one may calculate the RGE corrections to this matrix at Λ_{EW} and express the result in a way parallel to Eq. (3.4):

$$M'_\nu \simeq I_\alpha \left[M_\nu - \Delta_\tau \begin{pmatrix} 0 & 0 & M_{e\tau} \\ 0 & 0 & M_{\mu\tau} \\ M_{e\tau} & M_{\mu\tau} & 2M_{\tau\tau} \end{pmatrix} \right]. \quad (3.83)$$

It becomes obvious that the term proportional to Δ_τ measures the strength of μ - τ symmetry breaking. One may diagonalize M'_ν via the unitary transformation $U' = P'_\phi V' P'_\nu$, where V' is an analogue of V shown in Eq. (2.8), and P'_ϕ (or P'_ν) is an analogue of P_ϕ (or P_ν) as given above Eq. (3.45). Note again that P'_ϕ should not be ignored in

this treatment so as to keep everything consistent, although its phases have no physical meaning. After a lengthy but straightforward calculation, we obtain the neutrino masses

$$\begin{aligned} m'_1 &\simeq I_\alpha (1 - s_{12}^2 \Delta_\tau) m_1 , \\ m'_2 &\simeq I_\alpha (1 - c_{12}^2 \Delta_\tau) m_2 , \\ m'_3 &\simeq I_\alpha (1 - \Delta_\tau) m_3 \end{aligned} \quad (3.84)$$

at Λ_{EW} ; and the flavor mixing angles [136]

$$\begin{aligned} \theta'_{12} &\simeq \theta_{12} + \frac{1}{2} c_{12} s_{12} \frac{|\overline{m}_1 + \overline{m}_2|^2}{\Delta m_{21}^2} \Delta_\tau , \\ \theta'_{13} &\simeq c_{12} s_{12} \frac{m_3 |\overline{m}_1 - \overline{m}_2|}{\Delta m_{31}^2} \Delta_\tau , \\ \theta'_{23} &\simeq \frac{\pi}{4} + \frac{|\overline{m}_1 + m_3|^2 s_{12}^2 + |\overline{m}_2 + m_3|^2 c_{12}^2}{2 \Delta m_{31}^2} \Delta_\tau \end{aligned} \quad (3.85)$$

at Λ_{EW} . In addition, the two Majorana phases at the electroweak scale are given by

$$\begin{aligned} \rho' &\simeq \rho - \frac{m_1 m_2 \sin 2(\rho - \sigma) c_{12}^2}{\Delta m_{21}^2} \Delta_\tau , \\ \sigma' &\simeq \sigma - \frac{m_1 m_2 \sin 2(\rho - \sigma) s_{12}^2}{\Delta m_{21}^2} \Delta_\tau . \end{aligned} \quad (3.86)$$

Note that the validity of these interesting analytical results depends on the prerequisite that none of the physical parameters evolves violently from Λ_{FS} down to Λ_{EW} to make the perturbation expansions invalid. In this sense it is more general to discuss the running effects of relevant flavor parameters by deriving their differential RGEs from Eq. (3.79). But Eqs. (3.84)–(3.86) remain very useful for us to see some salient features of radiative corrections to the neutrino masses and flavor mixing parameters. Some more comments and discussions are in order.

- If the neutrino mass ordering is inverted or quasi-degenerate, the radiative correction to θ_{12} tends to be appreciable because it may be enhanced by a factor of $\mathcal{O}(100)$ in either of these two cases, although y_τ itself is strongly suppressed even in the MSSM. Nevertheless, a proper choice of $\rho - \sigma \sim \pm\pi/2$ can give rise to a significant cancellation in $|\overline{m}_1 + \overline{m}_2|$ and thus suppress the aforementioned enhancement and stabilize the running behavior of θ_{12} [137, 138]. If the neutrino mass ordering is normal, $\theta'_{12} - \theta_{12} \sim c_{12} s_{12} \Delta_\tau / 2$ is expected to be small. The higher the scale Λ_{FS} is, the smaller the corresponding value of θ_{12} will be as compared with its value at Λ_{EW} , because it always increases when running down in the MSSM scheme [139].
- As for θ_{13} , the RGE running effects are small no matter whether the neutrino mass ordering is normal or inverted. To generate $\theta_{13} \sim 9^\circ$ at Λ_{EW} from $\theta_{13} = 0$ at Λ_{FS} , the value of $\tan\beta$ has to be larger than 50 unless the absolute neutrino mass scale is unreasonably higher than 0.1 eV [140]. In this case the bottom-quark Yukawa coupling eigenvalue y_b would lie in the non-perturbation regime at a superhigh

energy scale — a result which is definitely unacceptable [141]. Namely, the RGE-induced μ - τ permutation symmetry breaking is hard to fit the observed value of θ_{13} , and thus one has to introduce a nonzero θ_{13} at Λ_{FS} . In the latter case the analytical approximations obtained in Eqs. (3.84)—(3.86) remain valid §, simply because θ_{13} itself does not significantly affect other parameters [133].

- An appreciable deviation of θ'_{23} from $\pi/4$ can be produced to fit the present experimental data when the neutrino masses are nearly degenerate. In particular, such a deviation will be positive in the MSSM scheme if $\Delta m_{31}^2 > 0$ holds, as one can see from Eq. (3.85). This interesting observation offers a potential correlation between the octant of θ_{23} and the neutrino mass hierarchy, and two numerical examples will be presented below [142].

Finally it is worth pointing out that a finite and meaningful value of δ will be generated along with a nonzero value of θ_{13} after the exact μ - τ permutation symmetry at Λ_{FS} is broken via the RGE running effects. Of course, the real seeds of such a radiative generation of δ must be the nontrivial Majorana phases ρ and σ at Λ_{FS} [143, 144].

The above approach also applies to the RGE-triggered μ - τ reflection symmetry breaking. Consider that a neutrino mass matrix M_ν exhibiting the μ - τ reflection symmetry is generated at a high scale Λ_{FS} . Then the corresponding neutrino mass matrix M'_ν at the electroweak scale Λ_{EW} can be deduced through Eq. (3.81) when the RGE running effects are taken into account. It is possible to reparametrize M'_ν in the form given by Eq. (3.69) with $\epsilon_2 = 2\epsilon_1 = \Delta_\tau$ as well as $\epsilon_{3,4} = 0$ [114]. Therefore, we can derive the RGE-induced $\Delta\theta_{23}$, $\Delta\delta$, $\Delta\rho$ and $\Delta\sigma$ from Eq. (3.73) by taking $R_2 = 2R_1 = \Delta_\tau$ and $I_{1,2} = 0$. While the result of $\Delta\theta_{23}$ is the same as that shown in Eq. (3.85), the result of $\Delta\delta$ appears as

$$\begin{aligned} \frac{\Delta\delta}{\Delta_\tau} = & \frac{m_3(\bar{m}_2 - \bar{m}_1)c_{12}s_{12}}{(m_3^2 - m_1^2)\tilde{s}_{13}} + \left(\frac{\bar{m}_1 s_{12}^2}{\bar{m}_1 - m_3} + \frac{\bar{m}_2 c_{12}^2}{\bar{m}_2 - m_3} \right) \\ & \cdot \frac{(\bar{m}_1 + m_3)c_{12}^2 - (\bar{m}_2 + m_3)s_{12}^2}{\bar{m}_1 + \bar{m}_2} \cdot \frac{\tilde{s}_{13}}{c_{12}s_{12}}. \end{aligned} \quad (3.87)$$

For illustration, we show the possible values of $\Delta\theta_{23}$, $\Delta\delta$, $\Delta\rho$ and $\Delta\sigma$ against the lightest neutrino mass in four typical cases in Fig. 3.3. Case (a): $(\rho^{(0)}, \sigma^{(0)}) = (0, 0)$ and $\tan\beta = 50$ with a normal neutrino mass ordering; Case (b): $(\rho^{(0)}, \sigma^{(0)}) = (0, 0)$ and $\tan\beta = 30$ with an inverted mass ordering; Case (c): $(\rho^{(0)}, \sigma^{(0)}) = (0, \pi/2)$ and $\tan\beta = 10$ with a normal neutrino mass ordering; and Case (d): $(\rho^{(0)}, \sigma^{(0)}) = (0, \pi/2)$ and $\tan\beta = 10$ with an inverted mass ordering. We have taken $\delta^{(0)} = -\pi/2$ in our numerical calculations. As one can see, the RGE running effects may be significant in the $m_1 \simeq m_2 \gg m_3$ case, and especially in the $m_1 \simeq m_2 \simeq m_3$ case with $\sigma^{(0)} \neq \rho^{(0)}$. Moreover, $\Delta\theta_{23}$ tends to be larger than $\Delta\delta$ when $\rho^{(0)}$ and $\sigma^{(0)}$ take the same values, or vice versa.

§ In this case the expression of θ'_{13} in Eq. (3.85) should be modified as $\theta'_{13} \simeq \theta_{13} + c_{12}s_{12}m_3|\bar{m}_1 - \bar{m}_2|\Delta_\tau/\Delta m_{31}^2$.




Figure 3.3. The possible values of $\Delta\theta_{23}$, $\Delta\delta$, $\Delta\rho$ and $\Delta\sigma$ against the lightest neutrino mass in four typical cases. We have taken $\Lambda_{\text{FS}} \sim 10^{14}$ GeV for the flavor symmetry scale. The sign “+−” or “++” corresponds to $(\rho^{(0)}, \sigma^{(0)}) = (0, 0)$ or $(\rho^{(0)}, \sigma^{(0)}) = (0, \pi/2)$.

We have discussed the RGE-induced corrections to a Majorana neutrino mass matrix which initially possesses the μ - τ permutation or reflection symmetry. Now we turn to radiative corrections to $|U_{\mu i}| = |U_{\tau i}|$, since these equalities are a straightforward consequence of the μ - τ symmetry. In other words, the observed effects of μ - τ symmetry breaking in the PMNS matrix U can be attributed to the RGE-induced corrections to $|U_{\mu i}| = |U_{\tau i}|$ when the neutrino masses and flavor mixing parameters run from Λ_{FS} down to Λ_{EW} . In this case it is possible to get the correct octant of θ_{23} and even the correct quadrant of δ [142]. Let us illustrate this striking point with the help of the one-loop RGEs in the MSSM framework. Note that the framework of the SM for the RGE running is less interesting in this connection for two simple reasons: (a) it is difficult to make the deviation of θ_{23} from 45° appreciable even if the neutrinos have a nearly degenerate mass spectrum; (b) the SM itself largely suffers from the vacuum-stability problem for the measured value of the Higgs mass ($\simeq 125$ GeV) as the flavor symmetry scale Λ_{FS} is above 10^{10} GeV [145]. For simplicity, here we start from $\theta_{23} = \pi/4$ and $\delta = -\pi/2$ at $\Lambda_{\text{FS}} \sim 10^{14}$ GeV to fit the observed pattern of the PMNS matrix U at $\Lambda_{\text{EW}} \sim 10^2$ GeV by taking account of the RGE evolution. Hence we are not subject to the special values of the Majorana phases as constrained by the μ - τ reflection symmetry imposed on the neutrino mass matrix M_ν . Namely, we mainly concentrate on the radiative breaking of

the equalities $|U_{\mu i}| = |U_{\tau i}|$ without going into details of the model-building issues.

We first define three μ - τ “asymmetries” of the PMNS matrix U and then figure out their expressions in the standard parametrization as follows:

$$\begin{aligned}
\Delta_1 &\equiv |U_{\tau 1}|^2 - |U_{\mu 1}|^2 \\
&= (\cos^2 \theta_{12} \sin^2 \theta_{13} - \sin^2 \theta_{12}) \cos 2\theta_{23} \\
&\quad - \sin 2\theta_{12} \sin \theta_{13} \sin 2\theta_{23} \cos \delta , \\
\Delta_2 &\equiv |U_{\tau 2}|^2 - |U_{\mu 2}|^2 \\
&= (\sin^2 \theta_{12} \sin^2 \theta_{13} - \cos^2 \theta_{12}) \cos 2\theta_{23} \\
&\quad + \sin 2\theta_{12} \sin \theta_{13} \sin 2\theta_{23} \cos \delta , \\
\Delta_3 &\equiv |U_{\tau 3}|^2 - |U_{\mu 3}|^2 = \cos^2 \theta_{13} \cos 2\theta_{23} .
\end{aligned} \tag{3.88}$$

It is clear that the three asymmetries satisfy the sum rule $\Delta_1 + \Delta_2 + \Delta_3 = 0$, and they vanish when the exact μ - τ flavor symmetry holds (i.e., when $\theta_{13} = 0$ and $\theta_{23} = \pi/4$, or when $\delta = \pm\pi/2$ and $\theta_{23} = \pi/4$). The present best-fit results of neutrino mixing parameters listed in Table 2.1 indicate that the possibility of $\delta = -\pi/2$ and $\theta_{23} = \pi/4$ is slightly more favored than either the possibility of $\delta = \pi/2$ and $\theta_{23} = \pi/4$ or that of $\theta_{13} = 0$ and $\theta_{23} = \pi/4$. Hence we infer that the natural condition for all the three Δ_i to vanish should be $\delta = -\pi/2$ and $\theta_{23} = \pi/4$ at $\Lambda_{\text{FS}} \sim 10^{14}$ GeV, and the observed pattern of lepton flavor mixing at $\Lambda_{\text{EW}} \sim 10^2$ GeV should be a consequence of the most “economical” μ - τ symmetry breaking triggered by the RGE running effects from Λ_{FS} to Λ_{EW} [142]. Explicitly, the one-loop RGEs of Δ_i are given by

$$\begin{aligned}
\frac{d\Delta_1}{dt} &= -y_\tau^2 [\xi_{21} (|U_{\tau 1}|^2 \Delta_2 + |U_{\tau 2}|^2 \Delta_1 + |U_{e3}|^2) \\
&\quad + \xi_{31} (|U_{\tau 1}|^2 \Delta_3 + |U_{\tau 3}|^2 \Delta_1 + |U_{e2}|^2) \\
&\quad + \zeta_{21} (|U_{\tau 1}|^2 \Delta_2 + |U_{\tau 2}|^2 \Delta_1 + |U_{e3}|^2) \cos \Phi_{12} \\
&\quad + \zeta_{31} (|U_{\tau 1}|^2 \Delta_3 + |U_{\tau 3}|^2 \Delta_1 + |U_{e2}|^2) \cos \Phi_{13} \\
&\quad + \mathcal{J} (\zeta_{21} \sin \Phi_{12} - \zeta_{31} \sin \Phi_{13})] , \\
\frac{d\Delta_2}{dt} &= y_\tau^2 [\xi_{21} (|U_{\tau 1}|^2 \Delta_2 + |U_{\tau 2}|^2 \Delta_1 + |U_{e3}|^2) \\
&\quad - \xi_{32} (|U_{\tau 2}|^2 \Delta_3 + |U_{\tau 3}|^2 \Delta_2 + |U_{e1}|^2) \\
&\quad + \zeta_{21} (|U_{\tau 1}|^2 \Delta_2 + |U_{\tau 2}|^2 \Delta_1 + |U_{e3}|^2) \cos \Phi_{12} \\
&\quad - \zeta_{32} (|U_{\tau 2}|^2 \Delta_3 + |U_{\tau 3}|^2 \Delta_2 + |U_{e1}|^2) \cos \Phi_{23} \\
&\quad + \mathcal{J} (\zeta_{21} \sin \Phi_{12} - \zeta_{32} \sin \Phi_{23})] , \\
\frac{d\Delta_3}{dt} &= y_\tau^2 [\xi_{31} (|U_{\tau 1}|^2 \Delta_3 + |U_{\tau 3}|^2 \Delta_1 + |U_{e2}|^2) \\
&\quad + \xi_{32} (|U_{\tau 2}|^2 \Delta_3 + |U_{\tau 3}|^2 \Delta_2 + |U_{e1}|^2) \\
&\quad + \zeta_{31} (|U_{\tau 1}|^2 \Delta_3 + |U_{\tau 3}|^2 \Delta_1 + |U_{e2}|^2) \cos \Phi_{13} \\
&\quad + \zeta_{32} (|U_{\tau 2}|^2 \Delta_3 + |U_{\tau 3}|^2 \Delta_2 + |U_{e1}|^2) \cos \Phi_{23} \\
&\quad - \mathcal{J} (\zeta_{31} \sin \Phi_{13} - \zeta_{32} \sin \Phi_{23})] ,
\end{aligned} \tag{3.89}$$

where $\xi_{ij} \equiv (m_i^2 + m_j^2)/\Delta m_{ij}^2$, $\zeta_{ij} \equiv 2m_i m_j/\Delta m_{ij}^2$, $\cos \Phi_{ij} \equiv \text{Re}(U_{\tau i} U_{\tau j}^*)^2/|U_{\tau i} U_{\tau j}^*|^2$ as

well as $\sin \Phi_{ij} \equiv \text{Im}(U_{\tau i} U_{\tau j}^*)^2 / |U_{\tau i} U_{\tau j}^*|^2$ [142]. The two Majorana CP-violating phases ρ and σ affect the evolution of Δ_i via $\cos \Phi_{ij}$ and $\sin \Phi_{ij}$. Given the fact $|\Delta m_{31}^2| \simeq |\Delta m_{32}^2| \sim 30 \Delta m_{21}^2$ with $\Delta m_{21}^2 \simeq 7.5 \times 10^{-5} \text{ eV}^2$ [72], we expect $\xi_{21} \gg |\xi_{31}| \simeq |\xi_{32}|$ to hold in most cases. But this does not necessarily mean that Δ_3 should be more stable against radiative corrections than Δ_1 and Δ_2 , because their running behaviors also depend on the initial inputs of $|U_{\alpha i}|^2$. An appreciable deviation of θ_{23} from $\pi/4$, which is sensitive or equivalent to an appreciable deviation of Δ_3 from zero, requires a sufficiently large value of $\tan \beta$. Hence the resulting octant of θ_{23} is controlled by the neutrino mass ordering (i.e., the sign of Δm_{31}^2 or Δm_{32}^2 , or equivalently the sign of ξ_{31} or ξ_{32}). Since the signs of ξ_{ij} and ζ_{ij} are always the same, one may adjust the evolving direction of Δ_3 without much fine-tuning of the other relevant parameters.

We proceed to present two numerical examples to illustrate radiative corrections to the equalities $|U_{\mu i}| = |U_{\tau i}|$, corresponding to the best-fit results of neutrino oscillation parameters given in Refs. [72] and [73]. We start from $\Delta_i = 0$ (i.e., $\theta_{23} = 45^\circ$ and $\delta = 270^\circ$) at $\Lambda_{\text{FS}} \sim 10^{14} \text{ GeV}$ and run them down to $\Lambda_{\text{EW}} \sim 10^2 \text{ GeV}$ via the RGEs that have been given in Eq. (3.89). Table 3.1 summarizes the typical inputs and outputs of these two examples. Some discussions are in order.

(1) In Example I with the inverted neutrino mass ordering, the best-fit results of Δm_{21}^2 , Δm_{31}^2 , θ_{12} , θ_{13} , θ_{23} and δ at Λ_{EW} [72] can be successfully reproduced from the proper inputs at Λ_{FS} . In this case $\theta_{23}(\Lambda_{\text{FS}}) - \theta_{23}(\Lambda_{\text{EW}}) \simeq 2.6^\circ$ and $\delta(\Lambda_{\text{FS}}) - \delta(\Lambda_{\text{EW}}) \simeq 34^\circ$ hold thanks to the RGE running effects, and thus $\theta_{23}(\Lambda_{\text{EW}})$ lies in the first octant and $\delta(\Lambda_{\text{EW}})$ is located in the third quadrant.

(2) In Example II only the normal neutrino mass ordering allows us to arrive at $\theta_{23}(\Lambda_{\text{EW}}) \simeq 48.9^\circ$ from $\theta_{23}(\Lambda_{\text{FS}}) = 45^\circ$. In this case we obtain $\delta(\Lambda_{\text{EW}}) \simeq 241^\circ$ from $\delta(\Lambda_{\text{FS}}) = 270^\circ$, a result consistent with the best-fit value of δ [73]. The future long- and medium-baseline neutrino oscillation experiments are going to pin down the octant of θ_{23} and the quadrant of δ , and then it will be possible to test the expected correlation between the neutrino mass ordering and the deviation of θ_{23} (or δ) from 45° (or 270°).

(3) The running behaviors of Δ_i from Λ_{FS} down to Λ_{EW} can similarly be understood. In view of $\Delta_3 = \cos^2 \theta_{13} \cos 2\theta_{23}$ given in Eq. (3.88), one must have $\Delta_3(\Lambda_{\text{EW}}) > 0$ for $\theta_{23}(\Lambda_{\text{EW}}) < 45^\circ$ in Example I, and $\Delta_3(\Lambda_{\text{EW}}) < 0$ for $\theta_{23}(\Lambda_{\text{EW}}) > 45^\circ$ in Example II. In comparison, the evolution of Δ_1 or Δ_2 is not so obvious, but $\Delta_1 + \Delta_2 + \Delta_3 = 0$ always holds at any energy scale between Λ_{EW} and Λ_{FS} [142].

(4) One has to adjust the initial values of ρ and σ at Λ_{FS} in a careful way to control the running behaviors of six neutrino oscillation parameters, so that their best-fit results at Λ_{EW} can be correctly reproduced. Nevertheless, we find that it is really possible to resolve the octant of θ_{23} and the quadrant of δ through the radiative breaking of $|U_{\mu i}| = |U_{\tau i}|$ by inputting proper values of the absolute neutrino mass m_1 and the MSSM parameter $\tan \beta$. Once such unknown parameters are measured or constrained to a better degree of accuracy in the future, it will be possible to examine whether the quantum corrections can really accommodate the observed effects of μ - τ symmetry breaking.

Table 3.1. The RGE-triggered corrections to $|U_{\mu i}| = |U_{\tau i}|$ for Majorana neutrinos running from $\Delta_i = 0$ at Λ_{FS} down to Λ_{EW} in the MSSM with $\tan\beta = 31$.

Parameter	$\Lambda_{\text{FS}} \sim 10^{14}$ GeV	$\Lambda_{\text{EW}} \sim 10^2$ GeV
Example I (Capozzi <i>et al</i> [72])		
m_1 (eV)	0.100	0.087
Δm_{21}^2 (eV ²)	1.70×10^{-4}	7.54×10^{-5}
Δm_{31}^2 (eV ²)	-2.98×10^{-3}	-2.34×10^{-3}
θ_{12}	35.2°	33.7°
θ_{13}	10.1°	8.9°
θ_{23}	45.0°	42.4°
δ	270°	236°
ρ	-82°	-66°
σ	19°	27°
\mathcal{J}	-0.040	-0.029
Δ_1	0	0.054
Δ_2	0	-0.142
Δ_3	0	0.088
Example II (Forero <i>et al</i> [73])		
m_1 (eV)	0.100	0.087
Δm_{21}^2 (eV ²)	2.12×10^{-4}	7.60×10^{-5}
Δm_{31}^2 (eV ²)	3.50×10^{-3}	2.48×10^{-3}
θ_{12}	32.1°	34.6°
θ_{13}	6.9°	8.8°
θ_{23}	45.0°	48.9°
δ	270°	241°
ρ	-76°	-45°
σ	17°	29°
\mathcal{J}	-0.027	-0.030
Δ_1	0	0.111
Δ_2	0	0.022
Δ_3	0	-0.133

3.6. Flavor mixing from the charged-lepton sector

So far we have taken the charged-lepton mass matrix M_l to be diagonal. If a specific texture of the neutrino mass matrix M_ν fails to generate the observed pattern of lepton flavor mixing, however, a non-diagonal form of M_l should be taken into account. Of course, M_l is usually non-diagonal in a realistic flavor-symmetry model. In a grand unified theory (GUT), for example, M_l is always associated with the down-type quark mass matrix M_d and thus both of them should be non-diagonal. In all these contexts

it is meaningful to consider the contribution of M_l to the overall lepton flavor mixing matrix $U = O_l^\dagger O_\nu$ [146, 147, 148, 149, 150, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161], where the phases of O_l and O_ν deserve some particular attention. To clarify the issue, we follow the formulas given in Ref. [162] but adopt a slightly different parametrization $O = U_{23}U_{13}U_{12}P_\alpha$ for O_l or O_ν , where $P_\alpha = \text{Diag}\{e^{i\alpha_1}, e^{i\alpha_2}, e^{i\alpha_3}\}$ and

$$\begin{aligned} U_{12} &= \begin{pmatrix} c_{12} & \tilde{s}_{12}^* & 0 \\ -\tilde{s}_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \\ U_{13} &= \begin{pmatrix} c_{13} & 0 & \tilde{s}_{13}^* \\ 0 & 1 & 0 \\ -\tilde{s}_{13} & 0 & c_{13} \end{pmatrix}, \\ U_{23} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & \tilde{s}_{23}^* \\ 0 & -\tilde{s}_{23} & c_{23} \end{pmatrix}, \end{aligned} \quad (3.90)$$

with the definition $\tilde{s}_{ij} = s_{ij}e^{i\delta_{ij}}$ (for $ij = 12, 13, 23$). In comparison, we recall the standard parametrization $O = P_\phi O_{23}U_{13}O_{12}P_\nu$ with $P_\phi = \text{Diag}\{e^{i\phi_1}, e^{i\phi_2}, e^{i\phi_3}\}$ and $P_\nu = \text{Diag}\{e^{i\rho}, e^{i\sigma}, 1\}$. These two descriptions are related to each other as follows:

$$\begin{aligned} \delta_{12} &= \phi_2 - \phi_1, \quad \delta_{13} = \delta + \phi_3 - \phi_1, \quad \delta_{23} = \phi_3 - \phi_2, \\ \alpha_1 &= \phi_1 + \rho, \quad \alpha_2 = \phi_2 + \sigma, \quad \alpha_3 = \phi_3. \end{aligned} \quad (3.91)$$

To be more specific, the lepton flavor mixing matrix U in the new parametrization reads

$$\begin{aligned} U &= O_l^\dagger O_\nu = P_\alpha^{l\dagger} U_{12}^{l\dagger} U_{13}^{l\dagger} U_{23}^{l\dagger} U_{23}^\nu U_{13}^\nu U_{12}^\nu P_\alpha^\nu \\ &= P_\alpha^{l\dagger} U_{23} U_{13} U_{12} P_\alpha^\nu, \end{aligned} \quad (3.92)$$

where the phase matrix P_α^l is irrelevant in physics as it can be rotated away via a redefinition of the phases of three charged-lepton fields. In most cases of the model-building exercises, the three angles of O_l and the (1, 3) angle of O_ν are significantly small \parallel . To the leading order of such small parameters, the three mixing angles of U can approximate to [162]

$$\begin{aligned} \tilde{s}_{12} &\simeq \tilde{s}_{12}^\nu - \tilde{\theta}_{12}^l c_{12}^\nu c_{23}^\nu + \tilde{\theta}_{13}^l c_{12}^\nu \tilde{s}_{23}^{\nu*}, \\ \tilde{s}_{13} &\simeq \tilde{\theta}_{13}^\nu - \tilde{\theta}_{13}^l c_{23}^\nu - \tilde{\theta}_{12}^l \tilde{s}_{23}^\nu, \\ \tilde{s}_{23} &\simeq \tilde{s}_{23}^\nu - \tilde{\theta}_{23}^l c_{23}^\nu. \end{aligned} \quad (3.93)$$

A particularly interesting case is that θ_{13}^l and θ_{13}^ν are both negligibly small as compared with θ_{12}^l , leading us to the approximate results

$$\begin{aligned} \delta &\simeq \delta_{12}^l - \delta_{12}^\nu - \pi, \\ s_{13} &\simeq \theta_{12}^l s_{23}^\nu, \\ s_{12} &\simeq s_{12}^\nu + \theta_{12}^l c_{12}^\nu c_{23}^\nu \cos \delta. \end{aligned} \quad (3.94)$$

\parallel Of course, a given O_l with one or two large angles may correct O_ν in a remarkable way [163]. But this possibility is beyond the scope of our interest in the present article.

In obtaining Eq. (3.94), we have used Eq. (3.91) as well as $\delta_{23} \simeq \delta_{23}^\nu$ and $\delta_{12} \simeq \delta_{12}^\nu$. Given the approximate μ - τ flavor symmetry, θ_{23}^ν is supposed to be around $\pi/4$. If θ_{12}^l approximates to the Cabibbo angle of quark flavor mixing $\theta_C \simeq 0.22$ ¶, then Eq. (3.94) can give rise to an interesting relation $\theta_{13} \simeq \theta_C/\sqrt{2} \simeq 0.15$, which is in good agreement with the present experimental data. Moreover, Eq. (3.94) implies

$$s_{12} \simeq s_{12}^\nu + c_{12}^\nu \theta_{13} \cos \delta. \quad (3.95)$$

If $O_\nu \simeq U_{\text{BM}}$ as given in Eq. (3.11), one will arrive at $s_{12} \gtrsim (1 - \theta_{13})/\sqrt{2}$ from Eq. (3.95). This lower bound is apparently inconsistent with the experimental result of θ_{12} . If $O_\nu \simeq U_{\text{TB}}$ as given in Eq. (3.10), however, the situation will be different because θ_{12}^ν itself is already close to the observed value of θ_{12} . In this case δ is required to approach $\pm\pi/2$ in order to suppress the contribution from the second term in Eq. (3.95).

Although the pattern of lepton flavor mixing appears strikingly different from that of quark flavor mixing, they have been speculated to have a potential link. In this respect a viable GUT model may offer an ideal context where the relevant quarks and leptons reside in the same representations such that their respective Yukawa coupling matrices can be naturally correlated ⁺. For instance, the aforementioned relation $\theta_{13} \simeq \theta_C/\sqrt{2}$ makes a possible unification of quarks and leptons quite appealing. Such a relation will come out if M_ν respects the μ - τ permutation symmetry and O_l has a CKM-like structure (i.e., θ_{12}^l is very close to the Cabibbo angle θ_C while θ_{13}^l and θ_{23}^l are negligibly small). Moreover, the well-known Gatto-Sartori-Tonin (GST) relation $\theta_C \simeq \sqrt{m_d/m_s}$ [164] tempts us to attribute the Cabibbo angle to the down-type quark mass matrix M_d . The latter happens to be related to M_l in the GUT framework [165]. Let us take the SU(5) GUT model, which can be embedded in the SO(10) GUT model, as an example. Accordingly, each family of quarks and leptons are grouped into the SU(5) representations $\bar{\mathbf{5}}$ and $\mathbf{10}$ in the manner

$$\bar{\mathbf{5}} = \begin{pmatrix} d_r^c \\ d_b^c \\ d_g^c \\ e \\ -\nu_e \end{pmatrix}, \quad \mathbf{10} = \begin{pmatrix} 0 & u_g^c & -u_b^c & u_r & d_r \\ \cdot & 0 & u_r^c & u_b & d_b \\ \cdot & \cdot & 0 & u_g & d_g \\ \cdot & \cdot & \cdot & 0 & e^c \\ \cdot & \cdot & \cdot & \cdot & 0 \end{pmatrix} \quad (3.96)$$

with the subscripts “r”, “b” and “g” being the color indices of quarks. Since the right-handed neutrino fields N_i are the SU(5) singlets, their masses can be much larger than the electroweak symmetry breaking scale. In the minimal SU(5) GUT scenario whose Higgs sector only contains the 5-dimensional representations $H_{\mathbf{5}}$ and $H_{\bar{\mathbf{5}}}$, the Yukawa interaction terms include $Y_u^{ij} \mathbf{10}_i \mathbf{10}_j H_{\mathbf{5}}$, $Y_\nu^{ij} \bar{\mathbf{5}}_i N_j^c H_{\mathbf{5}}$ and $Y_d^{ij} \mathbf{10}_i \bar{\mathbf{5}}_j H_{\bar{\mathbf{5}}}$ with i and j being the family indices [33]. After the SU(5) symmetry is broken down to the SM gauge symmetry, the Yukawa coupling matrices of quarks and leptons take the following forms:

$$-\mathcal{L}_{\text{mass}} = Y_u^{ij} Q_i u_j^c H_u + Y_\nu^{ij} L_i N_j^c H_u + Y_d^{ij} Q_i d_j^c H_d$$

¶ This assumption makes sense in some GUT models, where M_l and M_d are usually related to each other in a similar way.

⁺ To see some realistic models with certain flavor symmetries being considered in the GUT framework, we refer the readers to Ref. [33] and references therein.

$$+ Y_d^{ij} e_i^c L_j H_d , \quad (3.97)$$

in which the relevant notations are self-explanatory. It is well known that the last two terms in Eq. (3.97) imply $M_d = M_l^T$ after the electroweak gauge symmetry breaking. Hence it seems natural to expect $\theta_{12}^l \simeq \theta_C$ at the GUT scale if θ_C mainly originates from M_d . On the other hand, the mass relations $m_d = m_e$, $m_s = m_\mu$ and $m_b = m_\tau$ derived from $M_d = M_l^T$ at the GUT scale are difficult to fit current experimental data even though the RGE running effects of those masses are taken into account. Such a problem forces us to extend the Higgs sector with higher dimensional representations. A famous example of this kind is the 45-dimensional Higgs representation $H_{\overline{45}}$ introduced in the Georgi-Jarlskog (GJ) mechanism [166]. This Higgs field contributes to the down-type quark sector and the charged-lepton sector via the Yukawa interaction term

$$Y_d^{ij} \mathbf{10}_i \bar{\mathbf{5}}_j H_{\overline{45}} \implies Y_d^{ij} (Q_i d_j^c - 3e_i^c L_j) H_d , \quad (3.98)$$

in which the factor -3 is a Clebsch-Gordan coefficient. Given such a difference between M_d and M_l^T , the GJ mechanism may lead to a few more realistic mass relations. But a new problem emerges in this case: $\theta_{12}^l \simeq \theta_C/3$, which is too small to enhance θ_{13} from 0 to the observed value.

Following the model-building strategies outlined in Refs. [167, 168], one may certainly consider some more options for the Higgs representations in order to derive $\theta_{12}^l \simeq \theta_C$ as well as the viable mass relations. If only the dimension-four operators are allowed, then only the 5- and 45-dimensional Higgs representations can contribute to lepton and quark masses. When the dimension-five operators are taken into account, more possibilities will arise. The contributions of a given operator to the down-type quark and charged-lepton masses are parametrized as $Y_d^{ij} (Q_i d_j^c + c_{ij} e_i^c L_j) H_d$ with c_{ij} being the Clebsch-Gordan coefficient. A list of all the candidate operators and the associated c_{ij} coefficients can be found in Ref. [169].

Assuming the (1, 3) and (2, 3) rotation angles to be negligible, here we focus on the (1, 2) submatrices of M_d and M_l which are mainly relevant to the first two families. In order to have the GST relation as a natural outcome, let us consider the texture [170, 171]

$$M_d = \begin{pmatrix} 0 & b \\ c & a \end{pmatrix} , \quad M_l = \begin{pmatrix} 0 & c_c c \\ c_b b & c_a a \end{pmatrix} , \quad (3.99)$$

where $b \simeq c$ holds from an empirical observation, and $c_{a,b,c}$ denote the Clebsch-Gordan coefficients of the relevant entries — they can be different from one another, indicating that different entries are generated by different Higgs representations. Given the complex (1, 2) rotation submatrix of U_{12} defined in Eq. (3.90), a diagonalization of M_d or M_l is straightforward. Since a , b and c are related to m_d , m_s and θ_C , one just needs to find out a proper combination of c_a , c_b and c_c which can yield the realistic electron and muon masses as well as the desired relation $\theta_{12}^l \simeq \theta_C$. It is found that only the combination of $c_b = 1/2$ and $c_a = c_c = 6$ can satisfy the above requirement [172, 173]. In this case the relation $\theta_{13} \simeq \theta_C/\sqrt{2}$ can finally be achieved provided M_ν possesses the μ - τ flavor symmetry. But δ_{12}^l and δ_{12}^ν remain undetermined, preventing us from calculating δ and

s_{12}' based on Eq. (3.94). One may follow the idea of spontaneous CP breaking to find a way out. To be specific, the CP symmetry is imposed on the model in the beginning to forbid the presence of CP violation, and it is spontaneously broken by some specific fields so that the resulting CP-violating phases are under control. A good example of this kind has been given in Ref. [174], where a scalar field acquires an imaginary vacuum expectation value and thus breaks the CP symmetry to the maximum level. This vacuum expectation value leads to $\delta_{12}^l = \pm\pi/2$, while δ_{12}' remains vanishing since the neutrino sector has no link to the relevant scalar field. So we arrive at $\delta = \pm\pi/2$ for the PMNS matrix U , and s_{12}' should take a value which is not far away from the experimental value of s_{12} . In such an explicit model-building exercise one may construct $O_\nu = U_{\text{TB}}$ to make things easy.

4. Larger flavor symmetry groups

Before introducing some concrete models to show how to realize the μ - τ flavor symmetry, let us first formulate the connection between neutrino mixing and flavor symmetries from a group-theoretical angle and then outline the general strategy for generating a particular neutrino mixing pattern. In addition, we shall briefly describe some recent works on combining the flavor symmetries and generalized CP (GCP) symmetries. Such a new treatment can not only pave the way for realizing the μ - τ flavor symmetry but also lay the foundation for embedding it in a larger flavor symmetry group. The latter will become relevant especially when one is ambitious to nail down the PMNS matrix U completely, since the μ - τ symmetry itself can only fix a part of the texture of U .

4.1. Neutrino mixing and flavor symmetries

As shown in section 3.1, $\theta_{13} = 0$ and $\theta_{23} = \pi/4$ lead the third column of U to the form $(0, -1, 1)^T/\sqrt{2}$, implying a Z_2 or μ - τ permutation symmetry of M_ν . And the reverse is also true. In fact, a link between the pattern of neutrino mixing and a certain flavor symmetry of M_ν like this always exists, irrespective of any particular form of U in a sense as follows. In the basis where M_l is diagonal, U is identified as the unitary matrix used to diagonalize M_ν (i.e., $U^\dagger M_\nu U^* = D_\nu$). Hence the relation $M_\nu u_i^* = m_i u_i$ is implied, where u_i denotes the i -th column of U and m_i is the i -th neutrino mass eigenvalue corresponding to u_i . Then it is easy to check that M_ν must be invariant under the transformations \mathcal{S}_i (for $i = 1, 2, 3$):

$$\begin{aligned} \mathcal{S}_i^\dagger M_\nu \mathcal{S}_i^* &= \mathcal{S}_i^\dagger (m_i u_i u_i^T - m_j u_j u_j^T - m_k u_k u_k^T) \\ &= m_i u_i u_i^T + m_j u_j u_j^T + m_k u_k u_k^T \\ &= U D_\nu U^T = M_\nu, \end{aligned} \tag{4.1}$$

where \mathcal{S}_i are defined in terms of u_i via

$$\mathcal{S}_i = u_i u_i^\dagger - (u_j u_j^\dagger + u_k u_k^\dagger). \tag{4.2}$$

All the three \mathcal{S}_i are unitary and commute with one another. Thanks to the relation $\mathcal{S}_i\mathcal{S}_j = \mathcal{S}_k$, only two of them are independent. Furthermore, each of \mathcal{S}_i has order 2 as indicated by $\mathcal{S}_i^2 = 1$. Summarizing these features, we reach the conclusion that M_ν always has the Klein symmetry $K_4 = Z_2 \times Z_2$ [175, 176, 177]. The symmetry operators can be explicitly constructed in terms of u_i in an approach described by Eq. (4.2). For instance,

$$\begin{aligned}\mathcal{S}_1^{\text{TB}} &= \frac{1}{3} \begin{pmatrix} 1 & -2 & -2 \\ -2 & -2 & 1 \\ -2 & 1 & -2 \end{pmatrix}, \\ \mathcal{S}_2^{\text{TB}} &= \frac{1}{3} \begin{pmatrix} -1 & 2 & 2 \\ 2 & -1 & 2 \\ 2 & 2 & -1 \end{pmatrix}, \\ \mathcal{S}_3^{\text{TB}} &= \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix},\end{aligned}\tag{4.3}$$

when U takes the well-known flavor mixing pattern U_{TB} as given in Eq. (3.10). Note that $\mathcal{S}_3^{\text{TB}}$ is nothing but the μ - τ permutation operation in Eq. (3.6), up to a sign rearrangement.

The above conclusion can be put another way. If M_ν assumes a flavor symmetry described by \mathcal{S}_i , then the normalized invariant eigenvector of \mathcal{S}_i (defined by $\mathcal{S}_i u_i = u_i$) will constitute one column of U [175, 176, 177] [†]. One may verify this observation as follows. The invariance of M_ν with respect to \mathcal{S}_i yields

$$M_\nu u_i^* = \mathcal{S}_i^\dagger M_\nu \mathcal{S}_i u_i^* \implies \mathcal{S}_i M_\nu u_i^* = M_\nu u_i^*,\tag{4.4}$$

which in turn implies that $M_\nu u_i^*$ is identical with u_i up to a coefficient (i.e., the neutrino mass eigenvalue m_i):

$$M_\nu u_i^* = m_i u_i \implies u_i^\dagger M_\nu u_i^* = m_i.\tag{4.5}$$

It should be noted that the resulting flavor mixing pattern is independent of m_i . This point means that one may obtain a constant pattern of neutrino mixing directly from the flavor symmetry by bypassing a concrete form of M_ν . To do so, however, one has to know all the three symmetry operators \mathcal{S}_i . If only a single \mathcal{S}_i is known, then U can be partially determined. For example, when M_ν respects a symmetry defined by $\mathcal{S}_1^{\text{TB}}$ or $\mathcal{S}_2^{\text{TB}}$ given in Eq. (4.3), one column of U will be $(2, -1, -1)^T/\sqrt{6}$ or $(1, 1, 1)^T/\sqrt{3}$, respectively. We are therefore led to the so-called TM1 or TM2 mixing pattern [175, 178, 179, 180, 181, 182, 183]:

$$U_{\text{TM1}} = \frac{1}{\sqrt{6}} \begin{pmatrix} 2 & \sqrt{2}c & \sqrt{2}\tilde{s}^* \\ -1 & \sqrt{2}c + \sqrt{3}\tilde{s} & \sqrt{2}\tilde{s}^* - \sqrt{3}c \\ -1 & \sqrt{2}c - \sqrt{3}\tilde{s} & \sqrt{2}\tilde{s}^* + \sqrt{3}c \end{pmatrix},$$

[†] But which column it will occupy is a matter of convention and can be determined upon some phenomenological considerations.

$$U_{\text{TM2}} = \frac{1}{\sqrt{6}} \begin{pmatrix} 2c & \sqrt{2} & 2\tilde{s}^* \\ -c + \sqrt{3}\tilde{s} & \sqrt{2} & -\tilde{s}^* - \sqrt{3}c \\ -c - \sqrt{3}\tilde{s} & \sqrt{2} & -\tilde{s}^* + \sqrt{3}c \end{pmatrix}, \quad (4.6)$$

where $\tilde{s} \equiv \sin \theta e^{i\varphi}$ has been defined. In Eq. (4.6) the location of the invariant eigenvector of $\mathcal{S}_1^{\text{TB}}$ or $\mathcal{S}_2^{\text{TB}}$ has been specified. The model-building details and phenomenological consequences of these two particular neutrino mixing patterns will be elaborated in section 5.3. Note that an \mathcal{S}_i can only limit one column of U up to the associated Majorana phase. The reason lies in the fact that when one u_i is rephased as a whole, it remains an invariant eigenvector of \mathcal{S}_i . As for the charged leptons, the diagonal Hermitian matrix $M_l M_l^\dagger$ possesses the symmetry

$$\mathcal{T}^\dagger M_l M_l^\dagger \mathcal{T} = M_l M_l^\dagger, \quad (4.7)$$

in which $\mathcal{T} \equiv \text{Diag}\{e^{i\phi_1}, e^{i\phi_2}, e^{-i(\phi_1+\phi_2)}\}$ is a diagonal phase matrix. Conversely, the Hermitian matrix $M_l M_l^\dagger$ must be diagonal if it satisfies the phase transformation in Eq. (4.7). It is worth pointing out that the above discussions keep valid in the basis of non-diagonal M_l . But in this case one needs to make an appropriate transformation of those basis-dependent quantities. For example, the symmetry operator which is imposed on $M_l M_l^\dagger$ will become $\mathcal{T}' = O_l \mathcal{T} O_l^\dagger$ with O_l being the unitary matrix used to diagonalize $M_l M_l^\dagger$.

Although it is quite possible that the \mathcal{S}_i and \mathcal{T} symmetries just arise by accident, we choose to identify them as the residual symmetries from a larger flavor symmetry group G_F in the following [175, 176, 177]. In other words, the subgroup G_ν (or G_l) generated by \mathcal{S}_i (or \mathcal{T}) remains intact in the neutrino (or charged-lepton) sector when G_F itself is broken down:

$$G_F \implies \begin{cases} G_\nu = \langle \mathcal{S}_i \rangle & \text{for } M_\nu, \\ G_l = \langle \mathcal{T} \rangle & \text{for } M_l M_l^\dagger. \end{cases} \quad (4.8)$$

From the bottom-up point of view, one may get a hold of G_F by demanding that G_l and G_ν close to a group [184]. In order for G_F to be finite, there must exist an n such that $\mathcal{T}^n = 1$. As already remarked, we need a non-degenerate diagonal \mathcal{T} to guarantee the diagonality of $M_l M_l^\dagger$, implying $n \geq 3$. For the sake of simplicity, let us tentatively set $n = 3$ [‡]. This means that \mathcal{T} has 1, ω and ω^2 as its diagonal entries, where $\omega \equiv e^{i2\pi/3}$. But the ordering of these three entries has to be specified, because different options may result in different G_F . Given $G_l = \langle \mathcal{T} = \text{Diag}\{1, \omega, \omega^2\} \rangle$ and $G_\nu = \langle \mathcal{S}_2^{\text{TB}}, \mathcal{S}_3^{\text{TB}} \rangle$, for example, we can arrive at $G_F = S_4$ (the permutation group of four objects). Furthermore, it has been shown that S_4 (or a group containing S_4 as a subgroup) is the unique discrete group that is able to naturally accommodate the flavor mixing pattern U_{TB} from the group-theoretical arguments [177, 185]. Indeed, it is likely that merely one \mathcal{S}_i symmetry is the surviving symmetry from G_F . In this situation G_F is found to be successively S_4 , A_4 (the alternating group of four objects)

[‡] As a matter of fact, another choice of n does not necessarily yield a finite group. For example, it has been checked that no finite group can be obtained by using a \mathcal{T} with $3 < n < 30$ and $\mathcal{S}_{2,3}^{\text{TB}}$ as the generators [33].

or S_3 (the permutation group of three objects) when the generator of G_ν is only one of $\mathcal{S}_1^{\text{TB}}$, $\mathcal{S}_2^{\text{TB}}$ and $\mathcal{S}_3^{\text{TB}}$. Interestingly, the A_4 group that has been popularly considered for realizing U_{TB} [186, 187] actually does not contain a subgroup for the last column of U_{TB} . But an accidental symmetry in such models may elevate the A_4 symmetry to S_4 [188], making U_{TB} available. Roughly speaking, the group generated by \mathcal{T} and \mathcal{S}_i tends to be small when \mathcal{S}_i have some regular forms like those illustrated in Eq. (4.3). If the neutrino mixing pattern and the relevant \mathcal{S}_i lack any regularity, the resulting G_F will become much larger and even not finite. That is why the observed pattern of lepton flavor mixing is more likely to originate from a reasonable flavor symmetry than that of quark flavor mixing [177]. In order to accommodate the experimental value of θ_{13} , which more or less lowers the regularity of U , a much larger flavor symmetry group has to be invoked (an example of this kind is $\Delta(6N^2)$ [189] with N being a big number) [190, 191, 192, 193].

The aforementioned procedure of constructing G_F from certain G_ν and G_l can be reversed: assuming a flavor symmetry group G_F for the charged leptons and neutrinos, one may take some of its subgroups as the residual symmetries and derive the associated flavor mixing pattern [188]. Such an exercise can be done in a purely group-theoretical way with no need of building an explicit dynamic model. First of all, we need to divide the elements of G_F into two categories — one with the elements of order 2 and the other with the elements of order ≥ 3 . Then we may identify three elements \mathcal{S}_i which satisfy $\mathcal{S}_i\mathcal{S}_j = \mathcal{S}_j\mathcal{S}_i = \mathcal{S}_k$ (for $i \neq j \neq k$) in the former category as the generators of G_ν , and fix an element \mathcal{T} in the latter category as the generator of G_l . In the basis where \mathcal{T} is diagonal, U will be composed of the normalized invariant eigenvectors of \mathcal{S}_i . If there are not three such order-2 elements, then only one order-2 element can be identified as the residual symmetry for M_ν (i.e., $G_\nu = Z_2$) in which case U can be partly determined. In fact, some authors have considered the possibility of reducing G_ν from $Z_2 \times Z_2$ to Z_2 so as to leave θ_{13} as a free parameter [194, 195, 196, 197].

4.2. Model building with discrete flavor symmetries

Now let us recapitulate some key issues concerning an application of the group-theoretical arguments to some model-building exercises. The most important issue in building a flavor-symmetry model is how to break G_F while preserving the desired residual symmetries. To serve this purpose, one may introduce a new type of scalar fields — *flavons*, which do not carry any quantum number of the SM gauge symmetry but can constitute some nontrivial representations of G_F . They break a given flavor symmetry via acquiring their appropriate vacuum expectation values (VEVs). In the canonical (type-I) seesaw mechanism the Lagrangian responsible for generating the lepton masses may take a general form as

$$-\mathcal{L}_{\text{mass}} = \sum_{\gamma} y_l^{\gamma} \bar{L}^{\rho} l^{\sigma} H_d \frac{\phi_l^{\gamma}}{\Lambda} + \sum_{\gamma} y_{\nu}^{\gamma} \bar{L}^{\rho} N^{\sigma} H_u \frac{\phi_{\nu}^{\gamma}}{\Lambda}$$

$$+ \sum_{\gamma} y_N^{\gamma} N^{\sigma} N^{\sigma} \frac{\phi_N^{\gamma}}{\Lambda} + \text{h.c.} , \quad (4.9)$$

in which l and N stand respectively for the right-handed charged-lepton and neutrino fields, and the superscripts ρ , σ and γ denote the representations occupied by the fermion and flavon fields. Since the flavor symmetry breaks in different manners in the charged-lepton and neutrino sectors, the flavon fields for these two sectors are distinct. But when ϕ_{ν} and ϕ_N have the same transformation properties, they can be identified as the same field. In order to separate ϕ_l from $\phi_{\nu,N}$, an extra symmetry is usually needed (e.g., an auxiliary Z_2^{aux} symmetry under which ϕ_l and l are odd while the other fields are even). Furthermore, there must be enough independent flavon fields so that the Yukawa coupling parameters present in each sector can fit the lepton masses. Note that Λ represents a high energy scale above which the new fields may become active, and the ratio of a flavon's VEV to Λ (denoted as $\epsilon = \langle \phi \rangle / \Lambda$) is typically assumed to be a small quantity such that the magnitude of a Yukawa coupling parameter can be regulated by the power of ϵ — the spirit of the Froggatt-Nielsen mechanism for explaining the observed quark mass hierarchy [198]. This point will become relevant when one needs to either produce a hierarchical mass matrix like M_l or discuss the soft breaking of a flavor symmetry by higher-order terms. In this connection it is necessary to know what forms the flavon VEVs $\langle \phi_{l,\nu,N} \rangle$ should take in order to break all the flavor symmetries apart from G_l and G_{ν} . One can easily verify that $M_l M_l^{\dagger}$ will be invariant under \mathcal{T} if $\langle \phi_l^{\gamma} \rangle$ satisfies the condition [188]

$$\mathcal{T}^{\gamma} \langle \phi_l^{\gamma} \rangle = \langle \phi_l^{\gamma} \rangle . \quad (4.10)$$

Similarly, the effective neutrino mass matrix resulting from the seesaw mechanism will be unchanged with respect to \mathcal{S}_i provided [188]

$$\mathcal{S}_i^{\gamma} \langle \phi_{\nu}^{\gamma} \rangle = \langle \phi_{\nu}^{\gamma} \rangle , \quad \mathcal{S}_i^{\gamma} \langle \phi_N^{\gamma} \rangle = \langle \phi_N^{\gamma} \rangle . \quad (4.11)$$

If a flavon field has no way to offer the proper VEV required by Eq. (4.10) or (4.11), it has to be forbidden to acquire a VEV and therefore contributes nothing to the lepton masses. With these observations in mind, we are well prepared to assign suitable representations for the fermion and flavon fields in order to build a phenomenologically viable model.

We proceed to illustrate the above points using a simple flavor model based on $G_F = S_4$ [188]. The 24 elements of S_4 belong to five conjugacy classes in which the orders are 1, 2, 2, 3 and 4, respectively. Undoubtedly, the generators \mathcal{S}_i of G_{ν} are contained in the second or third class while the generator \mathcal{T} of G_l resides in the fourth or fifth class. For simplicity, we choose an order-3 element to be \mathcal{T} §. There are five irreducible representations for S_4 : **1**, **1'**, **2**, **3** and **3'**, whose dimensions are self-explanatory. Once the flavor symmetry is specified, a question immediately follows: which representation(s) should the lepton doublets furnish? With an order-2 element of G_F as the residual symmetry for M_{ν} , only two flavors can mix if all the lepton doublets

§ If an order-4 element is chosen as \mathcal{T} , then the flavor mixing pattern U_{BM} shown in Eq. (3.11) will arise.

reside in the one-dimensional representations [176]. In the situation that one lepton doublet is in a one-dimensional representation and the other two form a two-dimensional representation, one column of U will have a vanishing entry [176]. That is why the three lepton doublets are commonly organized into a three-dimensional representation. Here we put them in the representation $\mathbf{3}$, implying that the residual symmetries will be defined in this representation. To be specific, $\mathcal{T}^{\mathbf{3}}$ may be represented by $\text{Diag}\{1, \omega, \omega^2\}$ for a diagonal $M_l M_l^\dagger$. As for $\mathcal{S}_i^{\mathbf{3}}$, they can be represented by the $\mathcal{S}_i^{\text{TB}}$ in Eq. (4.3). Moreover, the explicit forms of \mathcal{T} and \mathcal{S}_i in other irreducible representations are listed in Table 4.1 [188], where σ_1 is the first Pauli matrix, and the flavon VEV alignments (with the identification $\phi_\nu^\gamma = \phi_N^\gamma$) preserving the \mathcal{T} and \mathcal{S}_i symmetries are also shown. Note that $\phi_l^{\mathbf{1}}$ and $\phi_\nu^{\mathbf{1}}$ are always allowed to have VEVs which will never break any symmetry. Furthermore, l and N are also assigned to the representation $\mathbf{3}$. Because of the product rule $\mathbf{3} \times \mathbf{3} = \mathbf{1} + \mathbf{2} + \mathbf{3} + \mathbf{3}'$, only the $\mathbf{1}$, $\mathbf{2}$, $\mathbf{3}$ and $\mathbf{3}'$ flavon representations may contribute to the lepton masses. Among them, $\phi_l^{\mathbf{2}}$ and $\phi_\nu^{\mathbf{3}}$ cannot receive nonzero VEVs which would otherwise break the desired residual symmetries. The VEVs $\langle \phi_l^{\mathbf{1}} \rangle$, $\langle \phi_l^{\mathbf{3}} \rangle$ and $\langle \phi_l^{\mathbf{3}'} \rangle$ help give rise to a diagonal M_l in the following form:

$$M_l = \langle H_d \rangle \begin{pmatrix} y_e & 0 & 0 \\ 0 & y_\mu & 0 \\ 0 & 0 & y_\tau \end{pmatrix}, \quad (4.12)$$

where $y_e = y_l^{\mathbf{1}}/\sqrt{3} + \sqrt{2}y_l^{\mathbf{3}'} / 3$, $y_\mu = y_l^{\mathbf{1}}/\sqrt{3} - \sqrt{2}y_l^{\mathbf{3}'} / 6 + y_l^{\mathbf{3}}/\sqrt{6}$ and $y_\tau = y_l^{\mathbf{1}}/\sqrt{3} - \sqrt{2}y_l^{\mathbf{3}'} / 6 - y_l^{\mathbf{3}}/\sqrt{6}$ containing the proper Clebsch-Gordan coefficients. Note that there are three free parameters in Eq. (4.12), exactly enough to fit the three charged-lepton masses. On the other hand, $\langle \phi_\nu^{\mathbf{1}} \rangle$, $\langle \phi_\nu^{\mathbf{2}} \rangle$ and $\langle \phi_\nu^{\mathbf{3}} \rangle$ will result in a Dirac neutrino mass matrix of the form

$$M_D = \langle H_u \rangle \begin{pmatrix} y_{11} & y_{12} & y_{12} \\ y_{12} & y_{22} & y_{23} \\ y_{12} & y_{23} & y_{22} \end{pmatrix}, \quad (4.13)$$

in which the four independent elements are given by $y_{11} = y_\nu^{\mathbf{1}}/\sqrt{3} + \sqrt{2}y_\nu^{\mathbf{3}'} / 3$, $y_{12} = y_\nu^{\mathbf{2}}/\sqrt{6} - \sqrt{2}y_\nu^{\mathbf{3}'} / 6$, $y_{22} = y_\nu^{\mathbf{2}}/\sqrt{6} + \sqrt{2}y_\nu^{\mathbf{3}'} / 3$ and $y_{23} = y_\nu^{\mathbf{1}}/\sqrt{3} - \sqrt{2}y_\nu^{\mathbf{3}'} / 6$. One can see that the texture of M_D has the same $(2, 3)$ permutation symmetry as the one shown in Eq. (3.7), and $y_{22} + y_{23} = y_{11} + y_{12}$ holds. In addition, the texture of the heavy Majorana neutrino mass matrix M_N is identical to that of M_D with the replacements $y_\nu^{\mathbf{1,2,3}'} \Rightarrow y_N^{\mathbf{1,2,3}'}$. Then it is straightforward to follow the canonical seesaw formula $M_\nu = M_D M_N^{-1} M_D^T$ to show that the neutrino mixing pattern U_{TB} can be achieved. Finally, let us stress that the allowed flavon VEVs and the neutrino mixing pattern will change if just one order-2 element is preserved as the residual symmetry.

As illustrated in Table 4.1, the flavon VEVs have to align in some particular directions to preserve the desired residual symmetries. Whether such special alignments can be naturally achieved matters, as it has something to do with whether the corresponding flavor-symmetry model is convincing or not. In this regard the most popular method of deriving special flavon VEVs is the so-called F -term alignment

Table 4.1. The explicit forms of \mathcal{T} , \mathcal{S}_i , $\langle\phi_l\rangle$ and $\langle\phi_\nu\rangle$ in each irreducible representation with \mathcal{T} being diagonal [188].

	1'	2	3	3'
\mathcal{T}	1	$\text{Diag}\{\omega, \omega^2\}$	$\text{Diag}\{1, \omega, \omega^2\}$	$\text{Diag}\{1, \omega, \omega^2\}$
$\langle\phi_l\rangle$	1	$(0, 0)^T$	$(1, 0, 0)^T$	$(1, 0, 0)^T$
\mathcal{S}_1	-1	σ_1	$\mathcal{S}_1^{\text{TB}}$	$-\mathcal{S}_1^{\text{TB}}$
\mathcal{S}_2	1	$\text{Diag}\{1, 1\}$	$\mathcal{S}_2^{\text{TB}}$	$\mathcal{S}_2^{\text{TB}}$
\mathcal{S}_3	-1	σ_1	$\mathcal{S}_3^{\text{TB}}$	$-\mathcal{S}_3^{\text{TB}}$
$\langle\phi_\nu\rangle$	0	$(1, 1)^T$	$(0, 0, 0)^T$	$(1, 1, 1)^T$

mechanism [199]. The latter invokes the supersymmetry, allowing one to take advantage of the $U(1)$ R-symmetry — $U(1)_R$ (under which the superpotential terms should carry a total charge of 2) \parallel . The charge assignments for this symmetry generally go as follows: the superfields for the SM fermions carry a charge of 1, while the superfields for Higgs and flavons are neutral. Hence the superpotential for generating lepton masses is not affected by this symmetry, but the flavon fields themselves cannot form the superpotential terms any more. To constrain the flavon VEVs, we may introduce the driving fields (denoted by ψ) [199] which transform trivially with respect to the SM gauge symmetry but nontrivially under the relevant flavor symmetry. In particular, they carry a charge of 2 of the $U(1)_R$ symmetry, implying that they must appear linearly in the superpotential terms. To be more explicit, the superpotential involving both ψ and the flavon fields takes the form

$$W = \psi (M_1^2 + M_2\phi + c\phi^2 + \cdots) , \quad (4.14)$$

where M_1 and M_2 are two dimension-one parameters, and c is dimensionless. If the supersymmetry remains intact when the flavor symmetry suffers breakdown, the flavon potential will be given by

$$V(\phi) = \sum_a \left| \frac{\partial W}{\partial \psi_a} \right|^2 , \quad (4.15)$$

where a is used to distinguish different components of the driving fields in multi-dimensional representations. By definition, the flavon VEVs are taken to minimize

\parallel This special symmetry originates from the fact that in the supersymmetry algebra the relations $\{Q, Q^\dagger\} = P$ and $\{Q, Q\} = \{Q^\dagger, Q^\dagger\} = [Q, P] = [Q^\dagger, P] = 0$ keep invariant with respect to the $U(1)$ transformation $Q \rightarrow Q \exp(i\phi)$ [141], where Q and P stand respectively for the generators of supersymmetry and space-time translations, and the curly and square brackets stand respectively for the anticommutation and commutation relations. Since Q itself carries a nonzero $U(1)_R$ charge, the distinct components of a supermultiplet which are connected to one another via the action of Q always have different $U(1)_R$ charges. For a superfield with the $U(1)_R$ charge n , for example, its ϕ , ψ and F components have the n , $n - 1$ and $n - 2$ charges, respectively [141]. The superpotential terms are accordingly required to assume a total charge of 2 in order for their F components to conserve $U(1)$ R-symmetry.

the potential $V(\phi)$:

$$|-F_{\psi_a}^*| = \left| \frac{\partial W}{\partial \psi_a} \right| = 0 . \quad (4.16)$$

For illustration, let us explain how the VEV alignments of $\phi_l^{\mathbf{3}}$ and $\phi_l^{\mathbf{3}'}$ in Table 4.1 ¶ can be derived with the help of three driving fields $\psi^{\mathbf{1}}$, $\psi^{\mathbf{3}'}$ and $\psi^{\mathbf{2}}$. The related superpotential appears as

$$\begin{aligned} W = & \psi^{\mathbf{1}} \left(c_1 \phi_l^{\mathbf{3}} \phi_l^{\mathbf{3}} + c_2 \phi_l^{\mathbf{3}'} \phi_l^{\mathbf{3}'} - M^2 \right) + \psi^{\mathbf{3}'} \left(c_3 \phi_l^{\mathbf{3}} \phi_l^{\mathbf{3}'} \right) \\ & + \psi^{\mathbf{2}} \left(c_4 \phi_l^{\mathbf{3}} \phi_l^{\mathbf{3}} + c_5 \phi_l^{\mathbf{3}'} \phi_l^{\mathbf{3}'} + c_6 \phi_l^{\mathbf{3}} \phi_l^{\mathbf{3}'} \right) . \end{aligned} \quad (4.17)$$

Note that a term like $\psi^{\mathbf{3}'} (M' \phi_l^{\mathbf{3}'})$ has been forbidden by the Z_2^{aux} symmetry mentioned below Eq. (4.9). The conditions for the flavon VEVs $\langle \phi_l^{\mathbf{3}} \rangle = (v_1, v_2, v_3)^T$ and $\langle \phi_l^{\mathbf{3}'} \rangle = (v'_1, v'_2, v'_3)^T$ turn out to be

$$\begin{aligned} c_1 (v_1^2 + 2v_2 v_3) + c_2 (v_1'^2 + 2v_2' v_3') - M^2 &= 0 , \\ c_3 \begin{pmatrix} v_2 v'_3 - v_3 v'_2 \\ v_1 v'_2 - v_2 v'_1 \\ v_3 v'_1 - v_1 v'_3 \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} , \end{aligned} \quad (4.18)$$

as well as

$$\begin{aligned} c_4 \begin{pmatrix} v_2^2 + 2v_1 v_3 \\ v_3^2 + 2v_1 v_2 \end{pmatrix} + c_5 \begin{pmatrix} v_2'^2 + 2v_1' v_3' \\ v_3'^2 + 2v_1' v_2' \end{pmatrix} \\ + c_6 \begin{pmatrix} v_2 v'_2 + v_1 v'_3 + v'_1 v_3 \\ -v_3 v'_3 - v_1 v'_2 - v'_1 v_2 \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix} . \end{aligned} \quad (4.19)$$

The VEV alignments illustrated in Table 4.1 (i.e., $v_{2,3} = v'_{2,3} = 0$) apparently satisfy these equations. In particular, the scale of v_1 and v'_1 is determined by M — the only dimensional parameter in Eq. (4.17), which is supposed to be located at a high energy scale. However, a driving field in the representation $\mathbf{1}$ may not be welcome in some specific models, in which case the mass term like that in Eq. (4.17) will be absent and the trivial vacuum (i.e., all the flavon VEVs are vanishing) offers a solution to the conditions for minimizing $V(\phi)$. To find a way out, the negative and supersymmetry-breaking flavon mass terms $-m_\phi^2 |\phi|^2$ are usually invoked to drive the flavon VEVs from the trivial ones [199].

Last but not least, let us point out that there exists a class of models where no residual symmetries survive. Such models are named the *indirect models* while those featuring residual symmetries are referred to as the *direct models* [201]. This classification is based on how the symmetries for $M_l M_l^\dagger$ and M_ν come about. In the direct models $M_l M_l^\dagger$ and M_ν are constrained by G_l and G_ν — the remnants of G_F , respectively. The relevant flavon VEVs should preserve these residual symmetries. In the indirect models the special forms of $M_l M_l^\dagger$ and M_ν arise accidentally, and the flavon VEVs do not necessarily keep invariant under any symmetry. In a realistic indirect

¶ A possible way of realizing the VEV alignments of $\phi_\nu^{\mathbf{2}}$ and $\phi_\nu^{\mathbf{3}}$ in Table 4.1 can be found in Ref. [200].

model M_l and M_N are typically taken to be diagonal. On the other hand, the Yukawa interaction term of the neutrinos takes the form

$$y_i \overline{L^3} N_i H_u \frac{\phi_i^3}{\Lambda}, \quad (4.20)$$

which can be expressed as

$$y_i (\overline{L}_e v_1^i + \overline{L}_\mu v_2^i + \overline{L}_\tau v_3^i) N_i H_u \frac{1}{\Lambda} \quad (4.21)$$

after the flavon fields have developed their VEVs as $\langle \phi_i^3 \rangle = (v_1^i, v_2^i, v_3^i)^T$. The effective Majorana neutrino mass matrix turns out to be [33]

$$M_\nu = \sum_i \frac{y_i^2 \langle H_u \rangle^2}{M_i \Lambda^2} \langle \phi_i^3 \rangle \langle \phi_i^3 \rangle^T, \quad (4.22)$$

where M_i (for $i = 1, 2, 3$) denote the right-handed neutrino masses. Interestingly, the resulting PMNS matrix U has a special structure: its three columns are proportional to $\langle \phi_1^3 \rangle$, $\langle \phi_2^3 \rangle$ and $\langle \phi_3^3 \rangle$, provided the latter are orthogonal to one another. One is therefore encouraged to choose these column vectors to align with the columns of the desired PMNS matrix U [202]. In order to achieve the flavor mixing pattern U_{TB} , for instance, the following alignments for $\langle \phi_i^3 \rangle$ are favored:

$$\begin{aligned} \langle \phi_1^3 \rangle &= v_1 \begin{pmatrix} 2 \\ -1 \\ -1 \end{pmatrix}, & \langle \phi_2^3 \rangle &= v_2 \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}, \\ \langle \phi_3^3 \rangle &= v_3 \begin{pmatrix} 0 \\ -1 \\ 1 \end{pmatrix}. \end{aligned} \quad (4.23)$$

These VEVs either stay invariant or change their signs under $\mathcal{S}_i^{\text{TB}}$, but none of $\mathcal{S}_i^{\text{TB}}$ is preserved by all the three VEVs. Since $\langle \phi_i^3 \rangle$ appear quadratically in M_ν , the sign changes are actually irrelevant and thus M_ν acquires the $\mathcal{S}_i^{\text{TB}}$ symmetries accidentally. To summarize, in the indirect models the flavor symmetry is mainly responsible for constraining the Yukawa coupling structure of the neutrinos to have the form given in Eq. (4.20). It can also assist the generation of special flavon VEVs like those shown in Eq. (4.23).

4.3. Generalized CP and spontaneous CP violation

As discussed above, the residual-symmetry approach lacks the predictive power for the Majorana phases. In addition, obtaining a reasonable value of θ_{13} at the leading order either needs a large G_F which will complicate the theory or reduces G_ν from $Z_2 \times Z_2$ to Z_2 which will be less predictive. It is found that the GCP symmetry [107, 108] can constrain the Majorana phases and allow a finite θ_{13} to be predicted. That is why the GCP has recently attracted a lot of attention in the model-building exercises, especially since θ_{13} was experimentally determined [203].

A theoretical reason for the introduction of the GCP concept lies in the fact that the canonical CP symmetry is not always consistent with the discrete non-Abelian flavor

symmetry. To consistently define the CP symmetry in the context of a discrete flavor symmetry, a nontrivial condition has to be fulfilled as one can see later on. For the sake of simplicity and without loss of generality, let us consider a scalar multiplet $\Phi = (\phi, \phi^*)^T$ on which G_F acts as

$$\Phi \rightarrow \rho(g)\Phi, \quad g \in G_F. \quad (4.24)$$

Here ρ is used to denote the representation constituted by Φ which is not necessarily irreducible. On the other hand, a CP symmetry is expected to act on Φ in the form

$$\Phi \rightarrow X\Phi^*, \quad (4.25)$$

in which X is unitary to keep the kinetic term $|\partial\Phi|^2$ invariant. Accordingly, a successive implementation of the CP, $g \in G_F$ and *inverse* CP symmetries will transform Φ through the following path:

$$\Phi \rightarrow X\Phi^* \rightarrow X\rho(g)^*\Phi^* \rightarrow X\rho(g)^*X^{-1}\Phi. \quad (4.26)$$

The consistency requirement suggests that $X\rho(g)^*X^{-1}$ should be a transformation belonging to G_F . We are therefore required to combine the CP symmetry with G_F as follows [107, 108]:

$$X\rho(g)^*X^{-1} = \rho(g'), \quad g' \in G_F. \quad (4.27)$$

Thanks to the preservation of the multiplication rules, we just need to assure the consistency of X with the generators of G_F . The canonical CP symmetry (i.e., $X = 1$) always follows when ρ is a real representation. But in a theory involving the complex representations $X = 1$ itself may not satisfy the consistency condition, in which case a GCP symmetry has to be invoked. Note that the GCP transformation may interchange different representations (especially those which are complex conjugate to each other). That is why one chooses to define CP on a vector space spanned by ϕ together with its complex conjugate counterpart ϕ^* . In principle, different real representations can also be connected by a GCP transformation [108]. Hence ρ has to contain all the representations connected by the GCP transformations. Moreover, all the X matrices allowed by Eq. (4.27) constitute a representation of the automorphism group $\text{Aut}(G_F)$ of G_F . Let us consider two elements a_1 and a_2 of $\text{Aut}(G_F)$ whose representation matrices are X_1 and X_2 , respectively. The representation matrix of a_2a_1 is X_2WX_1 , as determined by the reasoning [108]:

$$\begin{aligned} \rho(g'') &= X_2\rho(g')^*X_2^{-1} = X_2W\rho(g')WX_2^{-1} \\ &= X_2WX_1\rho(g)^*X_1^{-1}WX_2^{-1}, \end{aligned} \quad (4.28)$$

where W is the matrix interchanging the complex conjugate components of Φ (i.e., $\Phi^* = W\Phi$), and it has the properties

$$W^2 = 1, \quad \rho(g) = W\rho(g)^*W. \quad (4.29)$$

Note that for an X satisfying Eq. (4.27), $\rho(g)X$ is also a solution but it does not supply any additional physical implications. Hence the GCP transformations of physical interest are given by $\text{Aut}(G_F)$ modding out those equivalent ones and form a group called

H_{CP} . Consequently, the group G_{CP} constituted by G_{F} and H_{CP} is isomorphic to their semi product [107, 108]

$$G_{\text{CP}} = G_{\text{F}} \rtimes H_{\text{CP}} . \quad (4.30)$$

And the multiplication rule for any two elements of G_{CP} is given by

$$(g_1, h_1)(g_2, h_2) = (g_1 g'_2, h_1 h_2) , \quad (4.31)$$

where $g'_2 = h_1 g_2 h_1^{-1}$.

Now let us look at some physical implications of the GCP. By definition, the GCP symmetries constrain $M_l M_l^\dagger$ and M_ν in the following way:

$$X^\dagger M_l M_l^\dagger X = (M_l M_l^\dagger)^* , \quad X^\dagger M_\nu X^* = M_\nu^* . \quad (4.32)$$

Since $M_l M_l^\dagger$ and M_ν are diagonalized by O_l and O_ν , respectively, one may derive the following relations from Eq. (4.32):

$$X_l^\dagger D_l^2 X_l = D_l^2 , \quad X_\nu^\dagger D_\nu X_\nu^* = D_\nu , \quad (4.33)$$

in which

$$X_l = O_l^\dagger X O_l^* , \quad X_\nu = O_\nu^\dagger X O_\nu^* . \quad (4.34)$$

It becomes clear that X_l is a diagonal phase matrix. In comparison, X_ν is also diagonal but its finite entries are ± 1 . These results allow us to obtain a PMNS matrix with the property

$$X_l^\dagger U X_\nu = U^* . \quad (4.35)$$

It is easy to check that this relation will lead us to the trivial CP phases [204, 205]. Hence it is necessary to break the GCP symmetry in order to accommodate CP violation. A phenomenologically interesting and economical way of breaking $G_{\text{F}} \rtimes H_{\text{CP}}$ is to preserve the residual symmetries G_l and $G_\nu \rtimes H_{\text{CP}}^\nu = Z_2 \times H_{\text{CP}}^\nu$ in the charged-lepton and neutrino sectors [107], respectively, as illustrated in Fig. 4.1. This goal can be achieved by requiring the related flavon VEVs to satisfy the conditions

$$\mathcal{T}\langle\phi_l\rangle = \langle\phi_l\rangle , \quad \mathcal{S}\langle\phi_\nu\rangle = X\langle\phi_\nu\rangle^* = \langle\phi_\nu\rangle , \quad (4.36)$$

in which \mathcal{T} , \mathcal{S} and X are the representation matrices for the generators of G_l , Z_2 and H_{CP}^ν , respectively. Furthermore, the commutation between Z_2 and H_{CP}^ν requires

$$X \mathcal{S}^* X^{-1} = \mathcal{S} . \quad (4.37)$$

In light of $\mathcal{S}^2 = 1$, we may diagonalize \mathcal{S} by a unitary matrix $O_{\mathcal{S}}$ (i.e., $O_{\mathcal{S}}^\dagger \mathcal{S} O_{\mathcal{S}} = D_{\mathcal{S}} = \pm \text{Diag}\{-1, 1, -1\}$). One can see that the first and third eigenvalues of \mathcal{S} are degenerate. In this case it is possible to redefine $O_{\mathcal{S}}$ by carrying out a complex $(1, 3)$ rotation. In combination with the condition given in Eq. (4.37), this freedom allows us to have $O_{\mathcal{S}} O_{\mathcal{S}}^T = X$. Taking account of the invariance of M_ν under $Z_2 \times H_{\text{CP}}^\nu$,

$$\mathcal{S}^\dagger M_\nu \mathcal{S}^* = M_\nu , \quad X^\dagger M_\nu X^* = M_\nu^* , \quad (4.38)$$




Figure 4.1. A schematic illustration for the combination of G_F and G_{CP} as well as their breaking pattern which has direct consequence on the PMNS flavor mixing matrix U . Here we focus on the case of $G_\nu = Z_2$.

one finds

$$\begin{aligned} D_S O_S^\dagger M_\nu O_S^* &= O_S^\dagger M_\nu O_S^* D_S, \\ O_S^\dagger M_\nu O_S^* &= \left(O_S^\dagger M_\nu O_S^* \right)^*. \end{aligned} \quad (4.39)$$

This means that $O_S^\dagger M_\nu O_S^*$ can be diagonalized by a real orthogonal matrix

$$R(\theta) = \begin{pmatrix} \cos \theta & 0 & \sin \theta \\ 0 & 1 & 0 \\ -\sin \theta & 0 & \cos \theta \end{pmatrix}. \quad (4.40)$$

The PMNS matrix turns out to be [107]

$$U = U_l^\dagger O_S R(\theta) P_\nu, \quad (4.41)$$

up to possible permutations of its rows or columns. Here U_l is determined by G_l , whereas P_ν is a diagonal matrix (with its entries being ± 1 or $\pm i$) to keep the three neutrino mass eigenvalues positive. In such an approach θ is the only free parameter associated with the flavor mixing angles and CP-violating phases [107], and thus we are led to a few testable correlations among them. Of course, the value of θ can be determined by confronting it with the experimental result of θ_{13} .

In the above direct-model approach, the values of CP-violating phases are purely determined by the group structure of $G_F \rtimes H_{CP}$. So we are left with little room for manoeuvre once the flavor symmetry has been specified. Furthermore, this approach typically leads the CP phases to some simple or special numbers such as $0, \pm\pi/2$ or π [206, 207, 208, 209], although nontrivial CP phases may arise as the flavor symmetry becomes larger [210, 211]. Alternatively, the indirect models can provide various nontrivial CP phases which are easier to control. In such models the canonical CP symmetry is consistent with G_F and all the parameters are real. A nontrivial CP phase is introduced by the complex VEV of a flavon field ϕ which transforms trivially with

respect to G_F [212, 213, 214, 215]. In order to control the phase of $\langle\phi\rangle$, one may introduce an extra Z_n symmetry under which ϕ carries a single charge [216]. When the F -term alignment mechanism [199] is used to shape the flavon VEVs, there will be a superpotential term relevant to ϕ : $\psi(\phi^n/\Lambda^{n-2} \pm M^2)$, where ψ denotes a driving field which is neutral under Z_n and G_F . Therefore, $\langle\phi\rangle$ is required to satisfy the condition

$$|-F_\psi^*| = \left| \frac{\langle\phi\rangle^n}{\Lambda^{n-2}} \pm M^2 \right| = 0, \quad (4.42)$$

which may lead to any nontrivial phase for $\langle\phi\rangle$ with the help of an adjustable n [217, 218, 219].

5. Realization of the μ - τ flavor symmetry

We proceed to discuss how to realize the μ - τ flavor symmetry in different physical contexts. First of all, let us make some general remarks.

(1) $M_l M_l^\dagger$ and M_ν cannot simultaneously possess the same μ - τ flavor symmetry. Otherwise, the resulting lepton flavor mixing pattern would only contain a finite value of θ_{12} and thus be unrealistic. That is why one usually takes M_l to be diagonal and attributes the effects of lepton flavor mixing purely to M_ν in a specific model-building exercise. Of course, one may impose a flavor symmetry on M_l but keep M_ν diagonal, or allow both of them to be structurally non-diagonal and unparallel. Different basis choices are equivalent in principle, but one basis is likely to be advantageous over another in practice when building an explicit model to determine or constrain the flavor structures of both charged leptons and neutrinos. To illustrate this phenomenological point, we assume $M_l M_l^\dagger$ to have the μ - τ permutation symmetry in a form like that given in Eq. (3.7), leading to the eigenvalues m_α^2 (for $\alpha = e, \mu, \tau$) whose expressions are analogous to those given in Eq. (3.9). In this case one has to tolerate some significant cancellations among the parameters of $M_l M_l^\dagger$ in order to generate a strongly hierarchical charged-lepton mass spectrum (i.e., $m_e \ll m_\mu \ll m_\tau$ as shown in Fig. 2.2 [16]). It is certainly possible to conjecture that there exists a flavor symmetry which guarantees the exact cancellations and thus $m_e = m_\mu = 0$ at the lowest order, and it is the effect of flavor symmetry breaking that gives rise to nonzero m_e and m_μ together with the associated flavor mixing parameters. Assuming the strength of symmetry breaking to be $\epsilon \sim \mathcal{O}(0.1)$, one may expect that the finite values of m_μ , θ_{13} and $\Delta\theta_{23} \equiv \theta_{23} - \pi/4$ emerge at the $\mathcal{O}(\epsilon)$ level while nonzero m_e arises at the $\mathcal{O}(\epsilon^3)$ level. To achieve such results, however, a delicate arrangement of the relevant perturbation terms is indispensable [220]. In contrast, the neutrino mass spectrum should not exhibit a strong hierarchy unless m_1 is negligibly small. Another point is also noteworthy: although the lepton flavor mixing matrix is a constant matrix in the symmetry limit, it may be strongly correlated with the mass spectrum of charged leptons or neutrinos (or both of them) after the flavor symmetry is broken. When discussing the effects of μ - τ flavor symmetry breaking, one therefore prefers to work in the basis of M_l being diagonal such that it is more straightforward to infer some information about the neutrino mass spectrum from

the experimental observations of θ_{13} and θ_{23} . In other words, let us simply concentrate on the μ - τ permutation or reflection symmetry of M_ν and its possible breaking schemes in the basis where M_l itself is diagonal.

(2) A $U(1)_e \times U(1)_\mu \times U(1)_\tau$ flavor symmetry or its subgroup should be invoked in the charged-lepton sector to make $M_l M_l^\dagger$ diagonal. Nevertheless, the left-handed charged-lepton and neutrino fields reside in the same $SU(2)_L$ doublets, so the symmetry placed on one sector should equally act on the other sector. Hence the major challenge in model building is to preserve a symmetry in one sector but break it properly in the other sector. To serve this purpose, additional Higgs (or flavon) fields furnishing nontrivial representations of the flavor symmetry are generally needed. In this connection the Higgs (or flavon) fields, which are neutral with respect to the symmetry for keeping the diagonality of $M_l M_l^\dagger$ but transform nontrivially under the μ - τ flavor symmetry, can be employed to break the would-be degeneracy between m_μ and m_τ .

(3) A number of flavor-symmetry models, which are phenomenologically viable, have typically used the canonical seesaw mechanism to explain the smallness of three neutrino masses. Accordingly, at least two right-handed neutrinos are introduced in addition to the SM fields. No matter whether they assume the μ - τ flavor symmetry or not, the effective Majorana neutrino mass matrix M_ν will have this symmetry provided the left-handed neutrinos respect the same symmetry. This interesting point can be explicitly seen from the seesaw formula $M_\nu = M_D M_N^{-1} M_D^T$ [17, 18, 19, 20, 21] by taking the form of M_D as

$$M_D = \langle H \rangle \begin{pmatrix} y_{11} & y_{12} & y_{13} \\ y_{21} & y_{22} & y_{23} \\ y_{21} & y_{22} & y_{23} \end{pmatrix}, \quad (5.1)$$

where M_N is an arbitrary symmetric matrix. However, one might wish that the right-handed neutrinos should also obey some flavor symmetries in order to reduce the number of free parameters [221].

5.1. Models with the μ - τ permutation symmetry

In this subsection we aim to use a series of models, proposed by Grimus and Lavoura [113, 222, 223], to illustrate the essential ingredients for building a phenomenologically viable model based on the μ - τ flavor symmetry. The main idea of their models is that the $U(1)_e \times U(1)_\mu \times U(1)_\tau$ symmetry is conserved by the Yukawa interactions but softly broken by the dimension-three Majorana mass terms of right-handed neutrinos. Thus let us introduce three right-handed neutrinos N_α (for $\alpha = e, \mu, \tau$) and allow each of them to carry an associated lepton number. To properly break the flavor symmetry, three Higgs doublets H_i (for $i = 1, 2, 3$) are needed and each of them develops a VEV v_i (for $\sqrt{v_1^2 + v_2^2 + v_3^2} \simeq 246$ GeV). In addition to the lepton-number symmetries, two Z_2 -type flavor symmetries are also enforced [222]:

$$\begin{aligned} Z_2^{\mu\tau} : & \quad L_\mu \leftrightarrow L_\tau, \quad \mu_R \leftrightarrow \tau_R, \quad N_\mu \leftrightarrow N_\tau, \quad H_3 \rightarrow -H_3; \\ Z_2^{\text{aux}} : & \quad \mu_R, \tau_R, H_2, H_3 \text{ change sign}. \end{aligned} \quad (5.2)$$

Here $Z_2^{\mu\tau}$ is the μ - τ permutation symmetry interchanging all the fermion fields of the μ and τ flavors. Note that H_3 is odd under the $Z_2^{\mu\tau}$ symmetry, and thus it will break this symmetry when obtaining a VEV. Such a requirement is actually necessary for obtaining $m_\mu \neq m_\tau$, as one will see later on. In addition, there is an auxiliary symmetry Z_2^{aux} which separates H_1 from H_2 and H_3 . Because of this symmetry, H_2 and H_3 are responsible for the mass generation of μ and τ while H_1 only couples to e_R as well as N_α . Given the above setting, the Lagrangian for generating lepton masses is expressed as [224]

$$\begin{aligned} -\mathcal{L}_{\text{mass}} = & y_1 \overline{L}_e e_R H_1 + y_2 (\overline{L}_\mu \mu_R + \overline{L}_\tau \tau_R) H_2 \\ & + y_3 (\overline{L}_\mu \mu_R - \overline{L}_\tau \tau_R) H_3 + y_4 \overline{L}_e N_e \tilde{H}_1 \\ & + y_5 (\overline{L}_\mu N_\mu + \overline{L}_\tau N_\tau) \tilde{H}_1 + \text{h.c.} , \end{aligned} \quad (5.3)$$

where $\tilde{H}_1 \equiv i\sigma_2 H_1$. After the electroweak symmetry breaking, M_l and M_D turn out to be diagonal:

$$\begin{aligned} M_l = & \begin{pmatrix} y_1 v_1 & 0 & 0 \\ 0 & y_2 v_2 + y_3 v_3 & 0 \\ 0 & 0 & y_2 v_2 - y_3 v_3 \end{pmatrix} , \\ M_D = & \begin{pmatrix} y_4 v_1 & 0 & 0 \\ 0 & y_5 v_1 & 0 \\ 0 & 0 & y_5 v_1 \end{pmatrix} . \end{aligned} \quad (5.4)$$

Note that it is the coexistence of H_2 and H_3 that renders $m_\mu \neq m_\tau$. The heavy Majorana neutrino mass matrix M_N breaks the lepton-number symmetries in a soft way but maintains the other symmetries, so it also possesses the μ - τ permutation symmetry. Taking account of the seesaw formula, one therefore arrives at the light Majorana neutrino mass matrix M_ν which shares the same symmetry.

Several comments on this simple but instructive model-building approach are in order.

(a) In such a model the right-handed neutrinos necessarily suffer the μ - τ permutation symmetry. The reason is that $U(1)_\mu$ and $U(1)_\tau$ do not commute with $Z_2^{\mu\tau}$. As a result of the $U(1)_\mu$ (or $U(1)_\tau$) symmetry, L_μ (or L_τ) is only allowed to have couplings to μ_R and N_μ (or τ_R and N_τ). In order to keep $\mathcal{L}_{\text{mass}}$ invariant under $Z_2^{\mu\tau}$, μ_R and τ_R have to transform into each other as L_μ and L_τ do. So do N_μ and N_τ . Furthermore, $U(1)_\mu$, $U(1)_\tau$ and $Z_2^{\mu\tau}$ may be unified in a larger non-Abelian group which proves to be the two-dimensional orthogonal group O_2 [225].

(b) Another issue concerns the introduction of so many Higgs doublets. Although the rates of lepton-flavor-changing processes can be under control [226], one has to overcome some other phenomenological problems brought by so many electroweak-scale scalars (see, e.g., Refs. [227, 228, 229] for the discussions on some consequences of the multi-Higgs fields introduced in a number of flavor-symmetry models). To isolate the flavor symmetry issues from the electroweak issues, one can use the flavons which emerge at a high energy scale Λ to substitute for the extra Higgs doublets. As an example, we only keep H_1 while replacing H_2 and H_3 with two flavon fields ϕ_2 and ϕ_3 . And ϕ_2 (or

ϕ_3) has the same transformation properties as H_2 (or H_3) with respect to $Z_2^{\mu\tau}$ and Z_2^{aux} . Correspondingly, the Lagrangian relevant for lepton masses can be obtained from that in Eq. (5.3) with the replacements

$$\begin{aligned} (\overline{L}_\mu \mu_R + \overline{L}_\tau \tau_R) H_2 &\rightarrow (\overline{L}_\mu \mu_R + \overline{L}_\tau \tau_R) H_1 \frac{\phi_2}{\Lambda}, \\ (\overline{L}_\mu \mu_R - \overline{L}_\tau \tau_R) H_3 &\rightarrow (\overline{L}_\mu \mu_R - \overline{L}_\tau \tau_R) H_1 \frac{\phi_3}{\Lambda}. \end{aligned} \quad (5.5)$$

(c) From the point of view of naturalness, the strong mass hierarchy of charged leptons needs some explanations. As for the model under discussion, the smallness of m_e can be easily explained with the help of the Froggatt-Nielsen mechanism [198]. If one assigns a $U(1)_{\text{FN}}$ charge n to e_R , then the Yukawa interaction term for the electron will become

$$y_1 \overline{L}_e e_R H_1 \left(\frac{\langle \phi \rangle}{\Lambda} \right)^n, \quad (5.6)$$

where ϕ is a flavon field carrying a $U(1)_{\text{FN}}$ charge of -1 . Given $\epsilon \equiv \langle \phi \rangle / \Lambda$ as a very small quantity, m_e is suppressed by ϵ^n . However, this model does not permit the same mechanism to explain the smallness of m_μ as compared with m_τ . One finds that a relation $y_2 \langle v_2 \rangle \simeq -y_3 \langle v_3 \rangle$ with an accuracy of about 10% is required for obtaining the realistic values of m_μ and m_τ . Although a fine-tuning of this amount is not unacceptable, it is better to find a more natural way to obtain this kind of approximate relation. The idea is based on a new symmetry K which connects H_2 and H_3 as follows [230]:

$$\mu_R \rightarrow -\mu_R, \quad H_2 \leftrightarrow H_3. \quad (5.7)$$

Hence the coefficients y_2 and y_3 in Eq. (5.3) should be opposite to each other. Furthermore, the K symmetry can constrain the scalar potential to such a form that $v_2 = v_3$ is most likely to take place [230], in which case m_μ is exactly vanishing. When this symmetry is softly broken in the scalar potential, it is natural for us to expect a slight difference between v_2 and v_3 . And thus we arrive at a finite value of m_μ which is much smaller than that of m_τ [230].

(d) The model under discussion can be slightly modified to accommodate the μ - τ reflection symmetry [230]. For this purpose, $Z_2^{\mu\tau}$ is replaced by the μ - τ reflection symmetry while the other symmetries keep unchanged. Note that the μ - τ reflection symmetry can be viewed as a GCP symmetry described by

$$F_\alpha \rightarrow X_{\alpha\beta} F_\beta^c, \quad (5.8)$$

where F stands for all the fermion fields (i.e., L , N and the right-handed charged-lepton fields), $X = S^+$ as given by Eq. (3.6) [†], and the Greek subscripts run over e , μ and τ . On the other hand, the actions of this symmetry on the three Higgs fields are supposed to be

$$H_1 \rightarrow H_1^*, \quad H_2 \rightarrow H_2^*, \quad H_3 \rightarrow -H_3^*. \quad (5.9)$$

[†] A generalization of the μ - τ reflection symmetry can be found in Ref. [231], where X takes a form different from S^+ .

The resulting $\mathcal{L}_{\text{mass}}$ appears as

$$\begin{aligned}
-\mathcal{L}_{\text{mass}} = & y_1 \overline{L}_e e_R H_1 + (y_2 \overline{L}_\mu \mu_R + y_2^* \overline{L}_\tau \tau_R) H_2 \\
& + (y_3 \overline{L}_\mu \mu_R - y_3^* \overline{L}_\tau \tau_R) H_3 + y_4 \overline{L}_e N_e \tilde{H}_1 \\
& + (y_5 \overline{L}_\mu N_\mu + y_5^* \overline{L}_\tau N_\tau) \tilde{H}_1 + \text{h.c.} ,
\end{aligned} \tag{5.10}$$

where y_1 and y_4 are real by definition. As a result, the expressions of M_l and M_D are changed to

$$\begin{aligned}
M_l = & \begin{pmatrix} y_1 v_1 & 0 & 0 \\ 0 & y_2 v_2 + y_3 v_3 & 0 \\ 0 & 0 & y_2^* v_2 - y_3^* v_3 \end{pmatrix} , \\
M_D = & \begin{pmatrix} y_4 v_1 & 0 & 0 \\ 0 & y_5 v_1 & 0 \\ 0 & 0 & y_5^* v_1 \end{pmatrix} .
\end{aligned} \tag{5.11}$$

It should be noted that the phase difference between v_2 and v_3 cannot be $\pm\pi/2$. Otherwise, one would be led to the unrealistic result $m_\mu = m_\tau$. One may easily understand this observation by taking account of the fact that the μ - τ reflection symmetry still holds when v_2 is real and v_3 is purely imaginary. If M_N and M_D take the forms in Eqs. (3.24) and (5.11) respectively, then the seesaw formula assures the texture of M_ν to respect the μ - τ reflection symmetry.

5.2. Models with the μ - τ reflection symmetry

Now we turn to the model-building issues based on the μ - τ reflection symmetry. First of all, let us consider an interesting model developed by Mohapatra and Nishi for the purpose of illustration [232]. In this model the μ - τ reflection symmetry is imposed on all the fermion fields in the manner defined by Eq. (5.8). A $U(1)_{\mu-\tau}$ symmetry, which is helpful to make $M_l M_l^\dagger$ diagonal, is also introduced. As suggested by the notation of $U(1)_{\mu-\tau}$ itself, the e , μ and τ flavor fields carry the $U(1)_{\mu-\tau}$ charges of 0, 1 and -1 , respectively. So the $U(1)_{\mu-\tau}$ symmetry transformation can be described by

$$\mathcal{T} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\theta} & 0 \\ 0 & 0 & e^{-i\theta} \end{pmatrix} , \tag{5.12}$$

where θ is an arbitrary constant. As compared with $U(1)_e \times U(1)_\mu \times U(1)_\tau$, $U(1)_{\mu-\tau}$ has a quite appealing consequence: it commutes with the μ - τ reflection symmetry as indicated by $S^+ \mathcal{T}^* S^+ = \mathcal{T}$. This means that the whole flavor symmetry may be simply a direct product of $U(1)_{\mu-\tau}$ and the μ - τ reflection symmetry. In order to lift the would-be degeneracy between m_μ and m_τ , two extra Higgs doublets $H_{\pm 2}$ carrying the $U(1)_{\mu-\tau}$ charges of ± 2 are introduced in addition to the SM Higgs doublet H_0 . Given this framework, the Yukawa coupling terms relevant for M_l include [232]

$$\lambda_0 \overline{L}_e e_R H_0 + \lambda_{+2} \overline{L}_\mu \tau_R H_{+2} + \lambda_{-2} \overline{L}_\tau \mu_R H_{-2} . \tag{5.13}$$

The invariance of these terms with respect to the μ - τ reflection symmetry requires $\lambda_0 = \text{Re}(\lambda_0)$ and $\lambda_{+2} = \lambda_{-2}^*$ as well as the following transformation properties for the Higgs fields:

$$H_0 \rightarrow H_0^*, \quad H_{+2} \rightarrow H_{-2}^*. \quad (5.14)$$

Apparently, the μ - τ reflection symmetry will be broken if $|\langle H_{+2} \rangle| \neq |\langle H_{-2} \rangle|$ holds. In particular, the relation $|\langle H_{+2} \rangle| \ll |\langle H_{-2} \rangle|$ is required in order to generate $m_\mu \ll m_\tau$. Such a relation can be induced in the scalar potential by a scalar field which changes its sign under the μ - τ reflection symmetry [233]. It should be pointed out that the unwanted terms $\overline{L}_\mu \mu_R H_0$ and $\overline{L}_\tau \tau_R H_0$, which may render $M_l M_l^\dagger$ non-diagonal, have been forbidden by some additional symmetries (e.g., a Z_2^{aux} symmetry under which only μ_R , τ_R and $H_{\pm 2}$ change signs). Note that the Yukawa coupling terms in Eq. (5.13) will give rise to an accidental $U(1)_1 \times U(1)_2$ symmetry under which the related fields transform as

$$\begin{aligned} L_\mu &\rightarrow L_\mu e^{i\theta_1}, & \tau_R &\rightarrow \tau_R e^{i\theta_1}, \\ L_\tau &\rightarrow L_\tau e^{i\theta_2}, & \mu_R &\rightarrow \mu_R e^{i\theta_2}. \end{aligned} \quad (5.15)$$

Although the $U(1)_{\mu-\tau}$ symmetry will be spontaneously broken by $\langle H_{\pm 2} \rangle$, the $U(1)_1 \times U(1)_2$ symmetry promises $M_l M_l^\dagger$ to be diagonal. On the other hand, the Yukawa coupling terms of the neutrinos are given by [232]

$$y_0 \overline{L}_e N_e H_0 + y_2 \overline{L}_\mu N_\mu H_0 + y_3 \overline{L}_\tau N_\tau H_0, \quad (5.16)$$

where $y_0 = \text{Re}(y_0)$ and $y_2 = y_3^*$, as required by the μ - τ reflection symmetry. However, the resulting M_ν cannot be realistic unless M_N contains the $U(1)_{\mu-\tau}$ symmetry breaking terms. The latter can be achieved by including ϕ_1 and ϕ_2 which carry the respective $U(1)_{\mu-\tau}$ charges of 1 and 2 but transform trivially under the μ - τ reflection symmetry. Consequently, the mass or Yukawa-coupling terms contributing to M_N are obtained as

$$\begin{aligned} &M_{11} \overline{N}_e^c N_e + M_{23} \overline{N}_\mu^c N_\tau + y_{12} \overline{N}_e^c N_\mu \phi_1^* + y_{13} \overline{N}_e^c N_\tau \phi_1 \\ &+ y_{22} \overline{N}_\mu^c N_\mu \phi_2^* + y_{33} \overline{N}_\tau^c N_\tau \phi_2, \end{aligned} \quad (5.17)$$

where M_{11} and M_{23} are real mass parameters, whereas $y_{12} = y_{13}^*$ and $y_{22} = y_{33}^*$ hold. After ϕ_1 and ϕ_2 develop their real VEVs, the texture of M_N turns out to exhibit the μ - τ reflection symmetry. Given the form of M_D in Eq. (5.16), the texture of M_ν resulting from the seesaw formula will also share the μ - τ reflection symmetry.

It has recently been pointed out that the μ - τ reflection symmetry is not a necessary condition for obtaining the equality $|U_{\mu i}| = |U_{\tau i}|$ (for $i = 1, 2, 3$) for the PMNS matrix [234, 235]. In fact, this equality may arise in a more general context to be discussed below. Such a statement is based on the observation that an O_l^\dagger of the form in Eq. (3.19) together with a real orthogonal O_ν always results in a PMNS matrix featuring $U_{\mu i} = U_{\tau i}^*$. Moreover, it is found that O_l^\dagger and O_ν will satisfy the above criteria when the residual symmetries G_l and G_ν are real and able to completely fix the neutrino mixing pattern [234, 235]. Consider that G_l is generated by a real \mathcal{T} in which case O_l can be identified as the unitary matrix used to diagonalize \mathcal{T} . To obtain O_l , we first determine

the eigenvalues of \mathcal{T} with the help of [234]

$$\lambda_i^3 - \chi\lambda_i^2 + \chi\lambda_i - 1 = 0, \quad (5.18)$$

where χ is the trace of \mathcal{T} , and $|\lambda_i| = 1$ (for $i = 1, 2, 3$) holds. In addition to the eigenvalue $\lambda_1 = 1$, one finds the other two eigenvalues

$$\lambda_{2,3} = \frac{1}{2} \left[\chi - 1 \pm \sqrt{(\chi - 1)^2 - 4} \right]. \quad (5.19)$$

Apart from the special case of $\chi = 3$ or -1 , λ_2 and λ_3 are generally complex and conjugate to each other. For the general case, one can choose the eigenvectors corresponding to λ_2 and λ_3 to be complex conjugate to each other and the eigenvector for λ_1 to be real, making O_l^\dagger be of the form in Eq. (3.19). In comparison, the generators \mathcal{S}_1 and \mathcal{S}_2 of $G_\nu = Z_2 \times Z_2$ are order 2 and thus only have the eigenvalues 1 and -1 . Since we have required \mathcal{S}_1 and \mathcal{S}_2 themselves to be real, their common eigenvectors can be chosen to be real as well. Then we arrive at $O_\nu = O_S P_\nu$ with O_S being the real orthogonal matrix diagonalizing \mathcal{S}_1 and \mathcal{S}_2 simultaneously. Note that the undetermined diagonal phase matrix P_ν arises from the fact that the \mathcal{S}_1 and \mathcal{S}_2 symmetries can only determine O_ν up to the Majorana phases. Putting all these pieces together, a PMNS matrix $U = O_l^\dagger O_S P_\nu$ fulfilling the relation $|U_{\mu i}| = |U_{\tau i}|$ is finally available. Some immediate comments are in order.

(1) If G_ν only contains one Z_2 factor, then O_S cannot be completely fixed — an example of this kind is the TM1 or TM2 mixing pattern given in Eq. (4.6). In this case the reality of O_S is lost and the relation $|U_{\mu i}| = |U_{\tau i}|$ cannot hold any more. Therefore, G_ν and G_l are required to be able to pin down the lepton flavor mixing pattern in a realistic model-building exercise.

(2) When one considers unifying G_l and G_ν in a larger discrete symmetry G_F , a subgroup of $O(3)$ (e.g., A_4 , S_4 and A_5) will be a promising candidate in which G_l and G_ν are inherently real [234, 235]. Nevertheless, none of A_4 , S_4 or A_5 can give us a satisfactory θ_{13} at the lowest order [234]. Hence higher-order contributions have to be invoked in these symmetry-based models.

(3) This approach may serve as a complementarity to the μ - τ reflection symmetry approach but cannot take over from it. Although both of them can predict $\delta = \pm\pi/2$ and $\theta_{23} = \pi/4$, the former is also able to determine θ_{12} and θ_{13} but the latter fails in this connection. On the other hand, only the μ - τ reflection symmetry dictates the Majorana phases to take trivial values (i.e., 0 or $\pm\pi/2$).

(4) In fact, it is possible to generalize the theorem formulated here to recover the μ - τ reflection symmetry such that the Majorana phases can be fixed. Suppose that M_ν itself, instead of G_ν , should be real in which case O_ν is a real orthogonal matrix without being subject to an undetermined P_ν . The resulting neutrino mixing matrix will have the form in Eq. (3.19) with the Majorana phases being trivial. It is not surprising to find that such results agree with those predicted by the μ - τ reflection symmetry, simply because the neutrino mass matrix will take the form in Eq. (3.24) when one returns to the $M_l M_l^\dagger$ -diagonal basis by an O_l transformation of the form in




Figure 5.1. The possible values of s^2_{23} and $\sin \delta / \sin \varphi$ against φ in the TM1 (red lines) or TM2 (blue lines) neutrino mixing case.

Eq. (3.19) [236, 237, 238, 239]. This scenario provides an alternative way of getting at some interesting consequences of the μ - τ reflection symmetry [235].

5.3. On the TM1 and TM2 neutrino mixing patterns

As emphasized before, a general neutrino mass matrix with the μ - τ permutation symmetry is unable to give a definite prediction for θ_{12} unless some further conditions are placed on it. The most popular example in this regard might be $M_{\mu\mu} + M_{\mu\tau} = M_{ee} + M_{e\mu}$, a condition that can lead us to the flavor mixing pattern U_{TB} [92, 93]. The latter yields $\theta_{12} = \arctan(1/\sqrt{2}) \simeq 35.3^\circ$, which is compatible with the experimental value to a reasonably good degree of accuracy. Before the size of θ_{13} was measured in the Daya Bay experiment [66], the U_{TB} pattern was widely believed to be a good description of neutrino mixing and many discrete flavor symmetry groups (most notably, A_4 [186, 187]) had been employed to build the explicit models for deriving this special flavor mixing matrix. However, the Daya Bay discovery of a relatively large θ_{13} requires a remarkable modification of U_{TB} [240]. The simplest alternative turns out to be the TM1 or TM2 flavor mixing pattern given in Eq. (4.6) [175, 178, 179, 180, 181, 182, 183]. They owe their names to the fact that they keep the first and second columns of the U_{TB} pattern, respectively. In fact, one has [178, 182]

$$\begin{aligned}
 U_{\text{TM1}} &= U_{\text{TB}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & c & se^{-i\varphi} \\ 0 & -se^{i\varphi} & c \end{pmatrix}, \\
 U_{\text{TM2}} &= U_{\text{TB}} \begin{pmatrix} c & 0 & se^{-i\varphi} \\ 0 & 1 & 0 \\ -se^{i\varphi} & 0 & c \end{pmatrix},
 \end{aligned} \tag{5.20}$$

where $c \equiv \cos \theta$ and $s \equiv \sin \theta$. These two patterns are attractive from a phenomenological point of view. First of all, the observed θ_{13} can be easily accommodated thanks to the free parameter θ . Second, the value of s^2_{12} predicted by U_{TB} is only slightly modified in the TM1 or TM2 pattern:

$$s^2_{12} = \frac{1}{3} \cdot \frac{1 - 3s^2_{13}}{1 - s^2_{13}} \simeq 0.318 \quad (\text{for TM1}),$$

$$s_{12}^2 = \frac{1}{3} \cdot \frac{1}{1 - s_{13}^2} \simeq 0.341 \quad (\text{for TM2}) . \quad (5.21)$$

Note that the value of s_{12}^2 given by the TM1 mixing matrix is in better agreement with the experimental result than that given by U_{TB} or the TM2 mixing matrix. Third, θ_{23} is correlated with θ_{13} and the phase parameter φ as follows:

$$\begin{aligned} s_{23}^2 &\simeq \frac{1}{2} \left(1 - 2\sqrt{2}s_{13} \cos \varphi \right) \quad (\text{for TM1}) , \\ s_{23}^2 &\simeq \frac{1}{2} \left(1 + \sqrt{2}s_{13} \cos \varphi \right) \quad (\text{for TM2}) . \end{aligned} \quad (5.22)$$

The dependence of s_{23}^2 on φ , with the value of θ_{13} as an input, is illustrated in Fig. 5.1. One can see that θ_{23} may stay close to $\pi/4$ if φ takes a value around $\pi/2$ or $3\pi/2$. In addition, the correlation between θ_{23} and φ is stronger in the TM1 case and thus easier to be tested. Finally, the Jarlskog invariant of CP violation in the lepton sector [41] is found to be

$$\begin{aligned} \mathcal{J} &= -\frac{1}{6} \sqrt{2 - 6s_{13}^2} s_{13} \sin \varphi \quad (\text{for TM1}) , \\ \mathcal{J} &= -\frac{1}{6} \sqrt{2 - 3s_{13}^2} s_{13} \sin \varphi \quad (\text{for TM2}) , \end{aligned} \quad (5.23)$$

whose maximum magnitude is around 3.8% at $\varphi = \pi/2$ or $3\pi/2$. By comparing the above expression of \mathcal{J} with $\mathcal{J} = c_{12}s_{12}c_{13}^2s_{13}c_{23}s_{23}\sin\delta$ in the standard parametrization of U given below Eq. (2.13), we obtain the Dirac CP-violating phase δ as follows:

$$\begin{aligned} \frac{\sin \delta}{\sin \varphi} &= \frac{1 - s_{13}^2}{\sqrt{1 - 6s_{13}^2 - 4s_{13}^2 \cos 2\varphi}} \quad (\text{for TM1}) , \\ \frac{\sin \delta}{\sin \varphi} &= \frac{1 - s_{13}^2}{\sqrt{1 - 3s_{13}^2 - s_{13}^2 \cos 2\varphi}} \quad (\text{for TM2}) . \end{aligned} \quad (5.24)$$

As shown in Fig. 5.1, δ is approximately equal to φ , especially around $\varphi = \pi/2$ and $3\pi/2$.

Theoretically, exploring some appropriate physical contexts that can justify these two particular mixing patterns makes sense. Besides the residual-symmetry approach discussed in section 4.1 [241, 242, 243, 244, 245, 246, 247], the Friedberg-Lee (FL) symmetry [248, 249, 250, 251, 252, 253, 254, 255, 256] can also lead to the TM1 and TM2 mixing patterns in a natural way. The FL symmetry in the neutrino sector is defined in the sense that the neutrino mass operators should be invariant under the following translational transformations for the neutrino fields:

$$\nu_\alpha \rightarrow \nu_\alpha + \eta_\alpha \xi , \quad (5.25)$$

in which α runs over e, μ and τ , ξ is a spacetime-independent Grassmann-algebra element which anti-commutes with the neutrino fields, and η_α are the complex coefficients. The FL symmetry dictates the neutrino mass terms to be of the form

$$\begin{aligned} -\mathcal{L}_{\text{mass}} &= a \left(\eta_\tau^* \bar{\nu}_\mu - \eta_\mu^* \bar{\nu}_\tau \right) \left(\eta_\tau^* \nu_\mu^c - \eta_\mu^* \nu_\tau^c \right) \\ &\quad + b \left(\eta_\mu^* \bar{\nu}_e - \eta_e^* \bar{\nu}_\mu \right) \left(\eta_\mu^* \nu_e^c - \eta_e^* \nu_\mu^c \right) \\ &\quad + d \left(\eta_\tau^* \bar{\nu}_e - \eta_e^* \bar{\nu}_\tau \right) \left(\eta_\tau^* \nu_e^c - \eta_e^* \nu_\tau^c \right) + \text{h.c.} \end{aligned} \quad (5.26)$$

with a , b and d being the arbitrary complex numbers. When proposing the original version of this effective flavor symmetry, Friedberg and Lee assumed the neutrino fields to transform in a universal way (i.e., $\eta_e = \eta_\mu = \eta_\tau$) [248] in which case the neutrino mass matrix appears as [252]

$$M_\nu = \begin{pmatrix} b+d & -b & -d \\ -b & a+b & -a \\ -d & -a & a+d \end{pmatrix}. \quad (5.27)$$

This matrix can be transformed into the following form by using the U_{TB} matrix:

$$U_{\text{TB}}^\dagger M_\nu U_{\text{TB}}^* = \frac{1}{2} \begin{pmatrix} 3(b+d) & 0 & \sqrt{3}(b-d) \\ 0 & 0 & 0 \\ \sqrt{3}(b-d) & 0 & 4a+b+d \end{pmatrix} \quad (5.28)$$

which can be further diagonalized by a complex $(1, 3)$ rotation \ddagger . So the unitary matrix used to diagonalize M_ν has the same form as U_{TM1} in Eq. (5.20). However, a potential problem associated with M_ν in Eq. (5.28) is its prediction $m_2 = 0$, which is in conflict with $\Delta m_{21}^2 > 0$. A simple way out is to add a universal term $m_0 (\bar{\nu}_e \nu_e^c + \bar{\nu}_\mu \nu_\mu^c + \bar{\nu}_\tau \nu_\tau^c)$ to the above neutrino mass operators [248], leading to $m_2 = m_0$. But this term explicitly breaks the FL symmetry, and hence it needs a convincing explanation. Another way out is to modify the FL symmetry by assuming $\eta_e = -2\eta_\mu = -2\eta_\tau$ [257], and the associated neutrino mass matrix reads

$$M'_\nu = \begin{pmatrix} b+d & 2b & 2d \\ 2b & a+4b & -a \\ 2d & -a & a+4d \end{pmatrix}. \quad (5.29)$$

Similarly, U_{TB} can transform M'_ν into the form

$$U_{\text{TB}}^\dagger M'_\nu U_{\text{TB}}^* = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 3(b+d) & \sqrt{6}(d-b) \\ 0 & \sqrt{6}(d-b) & 2(a+b+d) \end{pmatrix} \quad (5.30)$$

which can be further diagonalized by a complex $(2, 3)$ rotation. Hence the resultant neutrino mixing matrix will be the TM2 pattern. In particular, $m_1 = 0$ comes out from this ansatz, implying that such a modified FL symmetry does not need to be broken. It is worth pointing out that a combination of the modified FL symmetry and the μ - τ reflection symmetry can pin down all the physical parameters of massive neutrinos. Note that $\eta_\mu = \eta_\tau$ allows for this kind of operation which otherwise would not make sense. In this specific scenario the neutrino mass matrix maintains its form in Eq. (5.29) and is subject to some further conditions such as $b = d^*$ and $\text{Im}(a) = 0$. The neutrino masses and flavor mixing quantities can be determined in terms of three real parameters: $\text{Re}(a)$, $\text{Re}(b)$ and $\text{Im}(b)$. While the above results for (m_1, m_2, m_3) and $(\theta_{12}, \theta_{13}, \theta_{23})$ still hold, the CP phases will take specific values ($\varphi = \pi/2$ or $3\pi/2$ and $\rho, \sigma = 0$ or $\pi/2$). To summarize, the modified FL symmetry can lead us to some definite predictions which are compatible with current neutrino oscillation data, but the FL symmetry itself remains puzzling and needs a further study.

\ddagger Taking $b = d$, one is led to the μ - τ permutation symmetry and thus the flavor mixing pattern U_{TB} .

Finally, let us emphasize that the TM1 and TM2 mixing patterns can also be derived via the *indirect model* approach. To achieve the TM1 or TM2 mixing matrix based on an indirect model which has been introduced at the end of section 4.2 for realizing U_{TB} , one has to modify the VEV alignments in Eq. (4.23) or construct M_l and (or) M_N with off-diagonal elements. A possible solution inspired by Eq. (5.29) is to change $\langle\phi_1^3\rangle$ and $\langle\phi_2^3\rangle$ to the following forms [258]:

$$\langle\phi_1^3\rangle = v_1 \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}, \quad \langle\phi_2^3\rangle = v_2 \begin{pmatrix} 1 \\ 0 \\ 2 \end{pmatrix}. \quad (5.31)$$

It is obvious that the effective neutrino mass matrix obtained from Eq. (4.22) with such VEV alignments will have the same form as that given in Eq. (5.29). In this case one neutrino remains massless, although the seesaw mechanism with three right-handed neutrinos has been taken into account. The observation that the TM1 or TM2 mixing pattern can be viewed as U_{TB} multiplied by a complex (1, 3) or (2, 3) rotation matrix from its right-hand side suggests another possible solution: one may keep the regular VEV alignments in Eq. (4.23) and ascribe the TM1 or TM2 mixing to the off-diagonal terms in an approximately diagonal M_N [214, 259]. To be specific, M_N may have the form

$$\begin{aligned} M_N &= \begin{pmatrix} M_1 & 0 & 0 \\ 0 & M_{22} & M_{23} \\ 0 & M_{23} & M_{33} \end{pmatrix} \quad (\text{for TM1}), \\ M_N &= \begin{pmatrix} M_{11} & 0 & M_{13} \\ 0 & M_2 & 0 \\ M_{13} & 0 & M_{33} \end{pmatrix} \quad (\text{for TM2}), \end{aligned} \quad (5.32)$$

where the off-diagonal terms may be traced back to the symmetry-breaking effects and thus relatively small. In this situation one can go back to the mass basis of right-handed neutrinos by means of a complex (2, 3) or (1, 3) rotation which will in turn change the Yukawa coupling matrix of the neutrinos. Then it is easy to check that the seesaw formula allows us to arrive at a particular texture of M_ν , which finally leads us to the TM1 or TM2 neutrino mixing matrix.

5.4. When the sterile neutrinos are concerned

A “sterile” neutrino ν_s , as its name suggests, does not carry any quantum number of the SM gauge symmetry and thus does not directly take part in the standard weak interactions. Although there has not been any convincing evidence for sterile neutrinos, their existence is either theoretically motivated or experimentally hinted. A good example of this kind is the *heavy* right-handed neutrinos introduced for implementing the type-I seesaw mechanism. On the other hand, the long-standing LSND anomaly [260] indicates the possible existence of an $\mathcal{O}(\text{eV})$ sterile neutrino which can more or less mix with ν_e and ν_μ . There are also a few other short-baseline neutrino-oscillation anomalies, such as the reactor [261], Gallium [262] and MiniBooNE [263] anomalies, which could

be explained with the help of active-sterile neutrino mixing §. It is therefore worthwhile to investigate an implementation of the μ - τ flavor symmetry in the presence of sterile neutrinos. One interesting proposal in this respect is that the sterile neutrino sector may be responsible for the μ - τ symmetry breaking [267, 268, 269, 270, 271, 272]. In this subsection we take this possibility seriously in the context of *light* sterile neutrinos.

According to Ref. [273], there is little improvement of the global-fit results in the 3+2 mixing scheme (i.e., the mixing of 3 active neutrinos and 2 sterile neutrinos) with respect to the 3+1 neutrino mixing scheme. For this reason, we restrict our discussion to the 3+1 case although a generalization of our results to the 3+2 or 3+3 case is rather straightforward. In this context the Majorana neutrino mass matrix for ν_e , ν_μ , ν_τ and ν_s can be expressed as [270]

$$M_s = \begin{pmatrix} m_{ee} & m_{e\mu} & m_{e\mu} & m_{es} \\ m_{e\mu} & m_{\mu\mu} & m_{\mu\tau} & m_{\mu s} \\ m_{e\mu} & m_{\mu\tau} & m_{\mu\mu} & m_{\tau s} \\ m_{es} & m_{\mu s} & m_{\tau s} & m_{ss} \end{pmatrix}. \quad (5.33)$$

Here the 3×3 submatrix of M_s for the active neutrinos has already been assumed to have the μ - τ permutation symmetry. But it is unnecessary to assume $m_{\mu s} = m_{\tau s}$ if the origin of sterile neutrinos is different from that of active ones. For instance, the active neutrino sector can be viewed as a consequence of the type-II seesaw mechanism [22, 23, 24, 25, 26] while the other elements of M_s might originate from the canonical seesaw mechanism. A general 4×4 neutrino mass matrix can be diagonalized by a 4×4 unitary matrix U_s to give 4 real mass eigenvalues m_i (for $i = 1, 2, 3, 4$). And U_s contains 6 rotation angles θ_{ij} (for $ij = 12, 13, 23, 14, 24, 34$), 3 Dirac phases and 3 Majorana phases. A quite popular parametrization of U_s reads as follows [274]:

$$U_s = O_{34}U_{24}U_{14}O_{23}U_{13}O_{12}P, \quad (5.34)$$

where O_{ij} is a real orthogonal rotation matrix with the mixing angle θ_{ij} , and U_{ij} denotes a unitary rotation matrix with both the mixing angle θ_{ij} and the phase parameter δ_{ij} as illustrated in Eq. (3.90). In addition, $P = \text{Diag}\{e^{i\rho}, e^{i\sigma}, 1, e^{i\gamma}\}$ denotes the Majorana phase matrix. This parametrization has an advantage that the upper-left 3×3 submatrix of U_s can simply reduce to the PMNS matrix U of the active neutrinos in the limit of vanishing active-sterile mixing.

Now we proceed to study the implications of the neutrino mass matrix in Eq. (5.33) by assuming that it can explain the experimental data concerning the active neutrinos. First of all, we intend to find out the possible values of m_4 and θ_{i4} . To serve this purpose, let us reconstruct the elements of M_s in terms of U_s and the neutrino masses [275]:

$$\begin{aligned} m_{e\mu} &\simeq -\overline{m}_1 c_{12} (s_{12} c_{23} + c_{12} \tilde{s}_{13} s_{23}) + \overline{m}_2 s_{12} (c_{12} c_{23} \\ &\quad - s_{12} \tilde{s}_{13} s_{23}) + m_3 \tilde{s}_{13}^* s_{23} + \overline{m}_4 s_{14} \tilde{s}_{24}^*, \\ m_{e\tau} &\simeq \overline{m}_1 c_{12} (s_{12} s_{23} - c_{12} \tilde{s}_{13} c_{23}) - \overline{m}_2 s_{12} (c_{12} s_{23} \end{aligned}$$

§ In addition, possible keV sterile neutrinos could be a good candidate for warm dark matter in the Universe [264, 265, 266].




Figure 5.2. The profiles of m_4 and s_{34} versus m_1 in the normal neutrino mass hierarchy. The results for $(\rho, \sigma) = (0, 0)$, $(\pi/2, 0)$ and $(\pi/2, \pi/2)$ are shown in red, green and purple colors and denoted as “++”, “-+” and “--”, respectively. Note that we have presented $-s_{34}$ in the $(\rho, \sigma) = (\pi/2, \pi/2)$ case for clarity. The standard neutrino parameters are allowed to take values in their 1σ intervals as listed in Table 2.1.

$$\begin{aligned}
& +s_{12}\tilde{s}_{13}c_{23}) + m_3\tilde{s}_{13}^*c_{23} + \bar{m}_4s_{14}\tilde{s}_{34}^* , \\
m_{\mu\mu} & \simeq \bar{m}_1 (s_{12}c_{23} + c_{12}\tilde{s}_{13}s_{23})^2 + \bar{m}_2 (c_{12}c_{23} \\
& -s_{12}\tilde{s}_{13}s_{23})^2 + m_3s_{23}^2 + \bar{m}_4\tilde{s}_{24}^2 , \\
m_{\tau\tau} & \simeq \bar{m}_1 (s_{12}s_{23} - c_{12}\tilde{s}_{13}c_{23})^2 + \bar{m}_2 (c_{12}s_{23} \\
& +s_{12}\tilde{s}_{13}c_{23})^2 + m_3c_{23}^2 + \bar{m}_4\tilde{s}_{34}^2 ,
\end{aligned} \tag{5.35}$$

where $\bar{m}_1 \equiv m_1 e^{2i\rho}$, $\bar{m}_2 \equiv m_2 e^{2i\sigma}$ and $\bar{m}_4 \equiv m_4 e^{2i\gamma}$. In obtaining these relations, we have made use of the smallness of θ_{i4} as implied by the unitarity of U which has been tested at the percent level [37, 38]. And we are allowed to identify θ_{12} , θ_{13} , θ_{23} and δ_{13} in U_s with their counterparts in U without loss of much accuracy. Taking account of the μ - τ symmetry conditions $m_{e\mu} = m_{e\tau}$ and $m_{\mu\mu} = m_{\tau\tau}$, we arrive at

$$\begin{aligned}
\tilde{s}_{34}^* & = \sqrt{2} \frac{(m_3 - m_{22}) \Delta\theta_{23} - m_{12}\tilde{s}_{13}}{m_{12}\Delta\theta_{23} + m_{11}\tilde{s}_{13} - m_3\tilde{s}_{13}^*} s_{14} - \tilde{s}_{24}^* , \\
\bar{m}_4 & = \sqrt{2} \frac{m_{12}\Delta\theta_{23} + m_{11}\tilde{s}_{13} - m_3\tilde{s}_{13}^*}{s_{14}(\tilde{s}_{34}^* - \tilde{s}_{24}^*)} ,
\end{aligned} \tag{5.36}$$

where m_{11} , m_{12} and m_{22} have already been defined in Eq. (3.5), and $\Delta\theta_{23} \equiv \theta_{23} - \pi/4$. There are so many free parameters that no definite conclusions can be drawn. For the sake of simplicity and illustration, let us consider the CP-conserving case by switching off all the CP phases. Some discussions are in order.

(1) $m_1 < m_2 \ll m_3$. In this case we obtain

$$\begin{aligned}
s_{34} & \sim -\sqrt{2} \frac{\Delta\theta_{23}}{s_{13}} s_{14} - s_{24} , \\
m_4 & \sim \frac{s_{13}^2}{s_{14}(s_{14}\Delta\theta_{23} + \sqrt{2}s_{13}s_{24})} m_3 ,
\end{aligned} \tag{5.37}$$

where s_{14} should not be too small, so as to make the magnitude of m_4 properly suppressed.

(2) $\underline{m_1 \simeq m_2 \gg m_3}$. For $(\rho, \sigma) = (0, 0)$, s_{34} and m_4 have the same expressions as those in Eq. (5.37) if the replacement $m_3 \Rightarrow m_1$ is made. In the $(\rho, \sigma) = (0, \pi/2)$ case, however, s_{34} and m_4 approximate to

$$\begin{aligned} s_{34} &\sim \sqrt{2} \frac{(c_{12}^2 - s_{12}^2) \Delta\theta_{23} - 2c_{12}s_{12}s_{13}}{2c_{12}s_{12}\Delta\theta_{23} + (c_{12}^2 - s_{12}^2)s_{13}} s_{14} - s_{24} , \\ m_4 &\sim \sqrt{2} \frac{2c_{12}s_{12}\Delta\theta_{23} + (c_{12}^2 - s_{12}^2)s_{13}}{s_{14}(s_{34} - s_{24})} m_2 , \end{aligned} \quad (5.38)$$

where the product of s_{14} and $s_{34} - s_{24}$ should not be too small, such that m_4 can be properly suppressed.

(3) $\underline{m_1 \simeq m_2 \simeq m_3}$. For $(\rho, \sigma) = (0, 0)$, s_{34} and m_4 are also the same as those in Eq. (5.37) with the replacement $m_3 \Rightarrow \Delta m_{31}^2/m_1$. On the other hand, the results in the $(\rho, \sigma) = (0, \pi/2)$ case are given by

$$\begin{aligned} s_{34} &\sim \sqrt{2} \frac{c_{12}^2 \Delta\theta_{23} - c_{12}s_{12}s_{13}}{c_{12}s_{12}\Delta\theta_{23} - s_{12}^2 s_{13}} s_{14} - s_{24} , \\ m_4 &\sim 2\sqrt{2} \frac{c_{12}s_{12}\Delta\theta_{23} - s_{12}^2 s_{13}}{s_{14}(s_{34} - s_{24})} m_1 . \end{aligned} \quad (5.39)$$

These analytical approximations are consistent with the numerical results illustrated in Fig. 5.2, in which the possible values of m_4 and s_{34} against m_1 are shown in the normal neutrino mass ordering case. In our numerical calculations we have specified $s_{14} \simeq s_{24} \simeq 0.1$ as a compromise between the unitarity test and short-baseline neutrino-oscillation anomalies. One can see that the allowed range of m_4 is rather wide and covers the region favored by those preliminary anomalies. In agreement with the unitarity test, the allowed range of $|s_{34}|$ is relatively small. Moreover, $|s_{34}|$ is not negligibly small in most of the parameter space, opening an interesting possibility for the short-baseline neutrino oscillations involving ν_τ .

Given the above results, let us return to M_s itself to examine its possible structures and the naturalness issue from a model-building point of view. We focus on the cases that the flavor mixing pattern and mass spectrum of three active neutrinos are mainly determined by the 3×3 submatrix of M_s , although the contribution from the sterile neutrino sector might in principle be dominant. To illustrate, we consider the following three typical cases.

(a) In the $m_1 < m_2 \ll m_3$ case, M_s can be parameterized in the form

$$M_s = m \begin{pmatrix} e\epsilon^2 & d\epsilon^2 & d\epsilon^2 & \epsilon \\ d\epsilon^2 & c\epsilon & -c\epsilon & a\epsilon \\ d\epsilon^2 & -c\epsilon & c\epsilon & b\epsilon \\ \epsilon & a\epsilon & b\epsilon & 1 \end{pmatrix} , \quad (5.40)$$

where $\epsilon \simeq 0.1$, and a, b, c, d and e are all the $\mathcal{O}(1)$ coefficients. The relatively heavy ν_4 as compared with ν_i (for $i = 1, 2, 3$) can be integrated out, like the treatment for the heavy Majorana neutrinos in the seesaw mechanism, leaving us a 3×3 neutrino mass matrix of the form in Eq. (3.59). According to our previous analysis, we can naturally

obtain the phenomenologically acceptable results without having to tune the relevant coefficients.

(b) The overall neutrino mass matrix that can result in $m_1 \simeq m_2 \gg m_3$ together with $(\rho, \sigma) = (0, 0)$ appears as

$$M_s = m \begin{pmatrix} 2c\epsilon & d\epsilon^2 & d\epsilon^2 & \epsilon \\ d\epsilon^2 & c\epsilon & c\epsilon & a\epsilon \\ d\epsilon^2 & c\epsilon & c\epsilon & b\epsilon \\ \epsilon & a\epsilon & b\epsilon & 1 \end{pmatrix}. \quad (5.41)$$

The resulting active neutrino mass matrix is similar to that given in Eq. (3.63) after the sterile neutrino is integrated out. As discussed before, the fine-tuning conditions $a + b - 2d \simeq 0$ and $(a + b)^2 - 2 \simeq 0$ will be required to fit the experimental data. It is therefore difficult to realize such a scenario from the viewpoint of model building.

(c) The $m_1 \simeq m_2 \simeq m_3$ case with $(\rho, \sigma) = (0, 0)$, which allows a relatively light ν_s , can be viewed as a consequence of the following mass matrix:

$$M_s = m \begin{pmatrix} f\epsilon^2 + 1 & e\epsilon^2 & e\epsilon^2 & \epsilon \\ e\epsilon^2 & d\epsilon + 1 & -d\epsilon & a\epsilon \\ e\epsilon^2 & -d\epsilon & d\epsilon + 1 & b\epsilon \\ \epsilon & a\epsilon & b\epsilon & c \end{pmatrix}. \quad (5.42)$$

Like the mass matrix given in Eq. (5.40), the present texture of M_s can naturally fit the experimental results regarding the active neutrino mixing and flavor oscillations. Of course, some higher-order terms in the above three ansätze of the neutrino mass matrix (e.g., those located at the $e\mu$ and $e\tau$ entries) can be neglected in the first place so as to get much more and much simpler predictions.

6. Some consequences of the μ - τ symmetry

The exact or approximate μ - τ flavor symmetry may have some important phenomenological consequences in neutrino physics and cosmology. Here let us look at a few interesting examples of this type, such as neutrino oscillations in matter, flavor distributions of ultrahigh-energy (UHE) cosmic neutrinos, cosmic matter-antimatter asymmetry via leptogenesis, and fermion mass matrices with a common Z_2 symmetry.

6.1. Neutrino oscillations in matter

It is known that the behaviors of neutrino oscillations in matter can be quite different from those in vacuum, simply because the matter-induced *coherent forward scattering* effects modifies the genuine neutrino masses and flavor mixing parameters when a neutrino beam travels through a normal medium (e.g., the Sun or Earth) [42, 43]. If the PMNS matrix U in vacuum possesses the μ - τ flavor symmetry, one can show that the effective PMNS flavor mixing matrix $\tilde{U} = \tilde{V}P_\nu$ in matter must possess the same

symmetry \dagger . In other words, the matter effects respect the μ - τ permutation or reflection symmetry in neutrino oscillations [106, 276]. Let us go into details of this observation in two different ways in the following.

In the basis where the mass and flavor eigenstates of three charged leptons are identical, we denote the effective neutrino mass matrix in matter as \widetilde{M}_ν . No matter whether massive neutrinos are of the Dirac or Majorana nature, the effective Hamiltonian responsible for the propagation of a neutrino beam in matter can be expressed in a similar way as that in Eq. (2.11):

$$\begin{aligned}\widetilde{\mathcal{H}}_{\text{eff}} &= \frac{1}{2E} \widetilde{M}_\nu \widetilde{M}_\nu^\dagger = \frac{1}{2E} \widetilde{V} \widetilde{D}_\nu^2 \widetilde{V}^\dagger \\ &= \frac{1}{2E} (V D_\nu^2 V^\dagger + D_m) ,\end{aligned}\tag{6.1}$$

where $\widetilde{D}_\nu = \text{Diag}\{\widetilde{m}_1, \widetilde{m}_2, \widetilde{m}_3\}$ with \widetilde{m}_i (for $i = 1, 2, 3$) being the effective neutrino masses in matter, and $D_m = \text{Diag}\{A, 0, 0\}$ with $A = 2\sqrt{2} G_F N_e E$ denoting the charged-current contribution to the coherent $\nu_e e^-$ forward scattering (here N_e stands for the background number density of electrons [42]).

Without loss of generality, it is always possible to redefine the phases of three neutrino mass eigenstates such that V_{ei} and \widetilde{V}_{ei} (for $i = 1, 2, 3$) are all real and positive [39]. To be explicit,

$$V = O_{23} O'_\delta O_{13} O_{12} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13} \\ V_{\mu 1} & V_{\mu 2} & c_{13}s_{23} \\ V_{\tau 1} & V_{\tau 2} & c_{13}c_{23} \end{pmatrix}\tag{6.2}$$

with

$$\begin{aligned}V_{\mu 1} &= -s_{12}c_{23}e^{-i\delta} - c_{12}s_{13}s_{23} , \\ V_{\mu 2} &= c_{12}c_{23}e^{-i\delta} - s_{12}s_{13}s_{23} , \\ V_{\tau 1} &= s_{12}s_{23}e^{-i\delta} - c_{12}s_{13}c_{23} , \\ V_{\tau 2} &= -c_{12}s_{23}e^{-i\delta} - s_{12}s_{13}c_{23} ,\end{aligned}\tag{6.3}$$

and $O'_\delta = \text{Diag}\{1, e^{-i\delta}, 1\}$. Such a phase convention is apparently different from that taken in Eq. (2.8). In matter the effective matrix \widetilde{V} takes the same phase convention as V in Eq. (6.2). In this particular basis $V_{ei} = |V_{ei}|$ and $\widetilde{V}_{ei} = |\widetilde{V}_{ei}|$ (for $i = 1, 2, 3$) hold, and thus it will be suitable for discussing the μ - τ flavor symmetry. Thanks to Eq. (6.1), we simply obtain

$$\sum_i \widetilde{m}_i^2 \widetilde{V}_{\alpha i} \widetilde{V}_{\beta i}^* = \sum_i m_i^2 V_{\alpha i} V_{\beta i}^* + \delta_{e\alpha} \delta_{e\beta} A ,\tag{6.4}$$

where α and β run over e , μ and τ . If V possesses the μ - τ symmetry, then it is straightforward to show that \widetilde{V} must have the same symmetry. Let us consider two distinct possibilities.

\dagger Note that the Majorana phase matrix P_ν is not affected by matter effects, as it has nothing to do with neutrino oscillations no matter whether they occur in vacuum or in matter.

(A) *The μ - τ permutation symmetry.* In this limit $\theta_{13} = 0$ and $\theta_{23} = \pi/4$ hold, and thus the phase parameter δ in V can be removed. We are then left with a real V with $V_{e3} = 0$, $V_{\mu 1} = -V_{\tau 1}$, $V_{\mu 2} = -V_{\tau 2}$ and $V_{\mu 3} = V_{\tau 3}$ in the above parametrization. So the elements of \tilde{V} satisfy

$$\begin{aligned} \sum_i \tilde{m}_i^2 |\tilde{V}_{ei}| \tilde{V}_{\mu i}^* &= - \sum_i \tilde{m}_i^2 |\tilde{V}_{ei}| \tilde{V}_{\tau i}^* , \\ \sum_i \tilde{m}_i^2 |\tilde{V}_{\mu i}|^2 &= \sum_i \tilde{m}_i^2 |\tilde{V}_{\tau i}|^2 , \end{aligned} \quad (6.5)$$

as one can see from Eq. (6.4). Because the matrix \tilde{V} is actually real and \tilde{m}_i^2 can be arbitrary for arbitrary values of E , it is always possible to arrive at the results $\tilde{V}_{e3} = 0$, $\tilde{V}_{\mu 1} = -\tilde{V}_{\tau 1}$, $\tilde{V}_{\mu 2} = -\tilde{V}_{\tau 2}$ and $\tilde{V}_{\mu 3} = \tilde{V}_{\tau 3}$ from Eq. (6.5), corresponding to $\tilde{\theta}_{13} = 0$ and $\tilde{\theta}_{23} = \pi/4$ in the parametrization of \tilde{V} similar to Eq. (6.2). Note that it is impossible to simultaneously have $\tilde{V}_{\mu 1} = \tilde{V}_{\tau 1}$, $\tilde{V}_{\mu 2} = \tilde{V}_{\tau 2}$ and $\tilde{V}_{\mu 3} = \tilde{V}_{\tau 3}$ (or $\tilde{V}_{\mu 1} = -\tilde{V}_{\tau 1}$, $\tilde{V}_{\mu 2} = -\tilde{V}_{\tau 2}$ and $\tilde{V}_{\mu 3} = -\tilde{V}_{\tau 3}$), because they violate the unitarity requirement of \tilde{V} . We conclude that \tilde{V} may have the same μ - τ permutation symmetry as V does.

(B) *The μ - τ reflection symmetry.* In this more interesting limit $\theta_{23} = \pi/4$ and $\delta = \pm\pi/2$ hold, leading to $V_{\mu 1} = V_{\tau 1}^*$, $V_{\mu 2} = V_{\tau 2}^*$ and $V_{\mu 3} = V_{\tau 3}$. As both $V_{\mu 3}$ and $V_{\tau 3}$ are real in the phase convention of Eq. (6.2), one may also take $V_{\mu 3} = V_{\tau 3}^*$ and thus $V_{\mu i} = V_{\tau i}^*$ (for $i = 1, 2, 3$) in this case [106]. As a result, Eq. (6.4) leads us to the relations

$$\begin{aligned} \sum_i \tilde{m}_i^2 |\tilde{V}_{ei}| \tilde{V}_{\mu i}^* &= \sum_i \tilde{m}_i^2 |\tilde{V}_{ei}| \tilde{V}_{\tau i} , \\ \sum_i \tilde{m}_i^2 |\tilde{V}_{\mu i}|^2 &= \sum_i \tilde{m}_i^2 |\tilde{V}_{\tau i}|^2 . \end{aligned} \quad (6.6)$$

Hence it is straightforward to obtain $\tilde{V}_{\mu i} = \tilde{V}_{\tau i}^*$ from Eq. (6.6) for arbitrary \tilde{m}_i^2 . Given the parametrization of \tilde{V} similar to Eq. (6.2), the above equalities give rise to $\tilde{\theta}_{23} = \pi/4$ and $\tilde{\delta} = \pm\pi/2$. That is why \tilde{V} possesses the μ - τ reflection symmetry as V does.

Now we turn to another way to show that the matter effects respect the μ - τ reflection symmetry in neutrino oscillations. Taking the parametrization of V in Eq. (6.2), we find

$$\begin{aligned} \tilde{M}_\nu \tilde{M}_\nu^\dagger &= O_{23} O'_\delta \left(O_{13} O_{12} D_\nu^2 O_{12}^\dagger O_{13}^\dagger + D_m \right) O_\delta^\dagger O_{23}^\dagger \\ &= O_{23} O'_\delta \hat{O}_{23} \hat{O}_{13} \hat{O}_{12} \tilde{D}_\nu^2 \hat{O}_{12}^\dagger \hat{O}_{13}^\dagger \hat{O}_{23}^\dagger O_\delta^\dagger O_{23}^\dagger \end{aligned} \quad (6.7)$$

with $\hat{O}_{23} \hat{O}_{13} \hat{O}_{12}$ being a real orthogonal matrix which consists of three rotation angles $(\hat{\theta}_{12}, \hat{\theta}_{13}, \hat{\theta}_{23})$ and has been defined to diagonalize the real bracketed part in Eq. (6.7). Up to the freedom of diagonal phase matrices P_L and P_R denoted below, which are actually irrelevant to neutrino oscillations, the effective neutrino mixing matrix in matter turns out to be

$$\begin{aligned} \tilde{V} &\equiv \tilde{O}_{23} \tilde{O}'_\delta \tilde{O}_{13} \tilde{O}_{12} = P_L O_{23} O'_\delta \hat{O}_{23} \hat{O}_{13} \hat{O}_{12} P_R \\ &= P_L \left(O_{23} O'_\delta \hat{O}_{23} \right) \tilde{O}_{13} \tilde{O}_{12} P_R , \end{aligned} \quad (6.8)$$

where \tilde{O}_{12} and \tilde{O}_{13} have been identified with \hat{O}_{12} and \hat{O}_{13} , respectively. In comparison, $\tilde{\theta}_{23}$ and $\tilde{\delta}$ must arise from a mixture of θ_{23} , δ and $\hat{\theta}_{23}$. In a way analogous to Eq. (2.13), we define the effective Jarlskog invariant $\tilde{\mathcal{J}}$ in matter and calculate it with the help of Eq. (6.8). After a straightforward calculation, we arrive at

$$\begin{aligned}\tilde{\mathcal{J}} &= \tilde{c}_{12}\tilde{s}_{12}\tilde{c}_{13}^2\tilde{s}_{13}\tilde{c}_{23}\tilde{s}_{23}\sin\tilde{\delta} \\ &= \tilde{c}_{12}\tilde{s}_{12}\tilde{c}_{13}^2\tilde{s}_{13}c_{23}s_{23}\sin\delta,\end{aligned}\tag{6.9}$$

in which the formula proportional to $\sin\tilde{\delta}$ is derived from the definition of \tilde{V} in Eq. (6.8), while the one proportional to $\sin\delta$ is obtained from the last equality of Eq. (6.8). Note that the free parameter $\hat{\theta}_{23}$ does not enter $\tilde{\mathcal{J}}$, although it mixes with θ_{23} and δ in the expressions of $\tilde{\theta}_{23}$ and $\tilde{\delta}$. Thus Eq. (6.9) leads us to the interesting Toshev relation [277]

$$\sin 2\tilde{\theta}_{23}\sin\tilde{\delta} = \sin 2\theta_{23}\sin\delta,\tag{6.10}$$

which links $\tilde{\theta}_{23}$ and $\tilde{\delta}$ in matter to θ_{23} and δ in vacuum. In addition to the parametrization of V in Eq. (2.8) or Eq. (6.2), one may also derive the Toshev-like relation from two other parametrizations of V or \tilde{V} [278].

Given $\theta_{23} = \pi/4$ and $\delta = \pm\pi/2$ for the PMNS flavor mixing matrix in vacuum, the Toshev relation in Eq. (6.10) tells us that $\sin 2\tilde{\theta}_{23}\sin\tilde{\delta} = \pm 1$ must hold in matter. As a result, we are left with $\tilde{\theta}_{23} = \theta_{23} = \pi/4$ and $\tilde{\delta} = \delta = \pm\pi/2$. We draw the conclusion that the μ - τ reflection symmetry keeps unchanged for neutrino oscillations in matter. Finally, it is obvious that these discussions are not applicable for the μ - τ permutation symmetry case, in which $\tilde{\delta}$ and δ are irrelevant in physics because of $\tilde{\theta}_{13} = \theta_{13} = 0$.

6.2. Flavor distributions of UHE cosmic neutrinos

An extremely important topic in high-energy neutrino astronomy is to search for the point-like neutrino sources that may help resolve a long-standing problem — the very origin of UHE cosmic rays. The reason is simply that the UHE protons originating in a cosmic accelerator (e.g., the gamma ray burst or active galactic nuclei [40]) unavoidably interact with their ambient photons via $p + \gamma \rightarrow \Delta^+ \rightarrow \pi^+ + n$ or their ambient protons through $p + p \rightarrow \pi^\pm + X$ with X being other particles, producing a large amount of energetic charged pions whose decays can therefore produce copious UHE cosmic muon neutrinos and electron neutrinos or their antiparticles. The so-called neutrino telescope is such a kind of deep underground ice or water Cherenkov detector which can record those rare events induced by the UHE cosmic neutrinos. Today the most outstanding neutrino telescope is the km³-volume IceCube detector at the South Pole, which has successfully identified thirty-seven extraterrestrial neutrino candidate events with deposited energies ranging from 30 TeV to 2 PeV [279, 280]. Among them, the three PeV events represent the highest-energy neutrino interactions ever observed. But where such extraordinary PeV neutrinos came from remains quite mysterious.

Given a distant astrophysical source of either $p\gamma$ or pp collisions, it has the same $\nu_\alpha + \bar{\nu}_\alpha$ flavor distribution $\Phi_e^S : \Phi_\mu^S : \Phi_\tau^S = 1 : 2 : 0$, where $\Phi_\alpha^S \equiv \Phi_{\nu_\alpha}^S + \Phi_{\bar{\nu}_\alpha}^S$ with

$\Phi_{\nu_\alpha}^S$ and $\Phi_{\bar{\nu}_\alpha}^S$ being the fluxes of ν_α and $\bar{\nu}_\alpha$ (for $\alpha = e, \mu, \tau$) at the source. Such an initial flavor distribution is expected to change to $\Phi_e^T : \Phi_\mu^T : \Phi_\tau^T = 1 : 1 : 1$ at a neutrino telescope, where $\Phi_\alpha^T \equiv \Phi_{\nu_\alpha}^T + \Phi_{\bar{\nu}_\alpha}^T$ is similarly defined, because the UHE cosmic neutrinos may oscillate many times on the way to the Earth and finally reach a flavor democracy [281] if the PMNS neutrino mixing matrix U satisfies the $|U_{\mu i}| = |U_{\tau i}|$ condition (for $i = 1, 2, 3$) [83]. As a consequence, the effects of μ - τ symmetry breaking can modify the ratio $\Phi_e^T : \Phi_\mu^T : \Phi_\tau^T = 1 : 1 : 1$ in a nontrivial way.

To be more explicit, the flavor distribution of UHE cosmic neutrinos at a terrestrial neutrino telescope is given by

$$\begin{aligned}\Phi_\alpha^T &= \sum_\beta \left[\Phi_{\nu_\beta}^S P(\nu_\beta \rightarrow \nu_\alpha) + \Phi_{\bar{\nu}_\beta}^S P(\bar{\nu}_\beta \rightarrow \bar{\nu}_\alpha) \right] \\ &= \sum_i \sum_\beta |U_{\alpha i}|^2 |U_{\beta i}|^2 \Phi_\beta^S,\end{aligned}\tag{6.11}$$

where we have used

$$P(\nu_\beta \rightarrow \nu_\alpha) = P(\bar{\nu}_\beta \rightarrow \bar{\nu}_\alpha) = \sum_i |U_{\alpha i}|^2 |U_{\beta i}|^2.\tag{6.12}$$

Note that Eq. (6.12) holds for a very simple reason: the galactic distance that the non-monochromatic UHE cosmic neutrinos have traveled far exceeds the observed neutrino oscillation lengths, and thus $P(\nu_\beta \rightarrow \nu_\alpha)$ and $P(\bar{\nu}_\beta \rightarrow \bar{\nu}_\alpha)$ are averaged over many oscillations and finally become energy- and distance-independent. Given $\Phi_e^S : \Phi_\mu^S : \Phi_\tau^S = 1 : 2 : 0$ and the unitarity of U , we immediately obtain

$$\begin{aligned}\Phi_\alpha^T &= \frac{\Phi_0}{3} \sum_i |U_{\alpha i}|^2 (|U_{ei}|^2 + 2|U_{\mu i}|^2) \\ &= \frac{\Phi_0}{3} \left[1 + \sum_i |U_{\alpha i}|^2 (|U_{\mu i}|^2 - |U_{\tau i}|^2) \right],\end{aligned}\tag{6.13}$$

where $\Phi_0 \equiv \Phi_e^S + \Phi_\mu^S + \Phi_\tau^S$ denotes the total flux of neutrinos and antineutrinos of all flavors. That is why $\Phi_e^T : \Phi_\mu^T : \Phi_\tau^T = 1 : 1 : 1$ can be achieved provided the equalities $|U_{\mu i}| = |U_{\tau i}|$ (for $i = 1, 2, 3$) hold. In fact, the difference between Φ_μ^T and Φ_τ^T is a straightforward signature of the μ - τ symmetry breaking:

$$\Phi_\mu^T - \Phi_\tau^T = \frac{\Phi_0}{3} \sum_i (|U_{\mu i}|^2 - |U_{\tau i}|^2)^2.\tag{6.14}$$

This exact and parametrization-independent result is very useful for us to probe the leptonic flavor mixing structure via the detection of UHE cosmic neutrinos at neutrino telescopes [282].

In the standard parametrization of U , let us define $\varepsilon \equiv \theta_{23} - \pi/4$ to measure a part of the μ - τ symmetry-breaking effects. Then we arrive at

$$\Phi_e^T : \Phi_\mu^T : \Phi_\tau^T = (1 + D_e) : (1 + D_\mu) : (1 + D_\tau)\tag{6.15}$$

with $D_e = -2\Delta$, $D_\mu = \Delta + \bar{\Delta}$ and $D_\tau = \Delta - \bar{\Delta}$, where the first- and second-order perturbation terms Δ [283] and $\bar{\Delta}$ [284] are expressed as

$$\Delta \simeq \frac{1}{2} \sin^2 2\theta_{12} \sin \varepsilon - \frac{1}{4} \sin 4\theta_{12} \sin \theta_{13} \cos \delta,$$

$$\begin{aligned}\overline{\Delta} \simeq & (4 - \sin^2 2\theta_{12}) \sin^2 \varepsilon + \sin^2 2\theta_{12} \sin^2 \theta_{13} \cos^2 \delta \\ & + \sin 4\theta_{12} \sin \varepsilon \sin \theta_{13} \cos \delta ,\end{aligned}\quad (6.16)$$

respectively. So a difference between Φ_μ^T and Φ_τ^T (i.e., a difference between D_μ and D_τ) is actually an effect of the second-order μ - τ symmetry breaking. It is easy to show $\overline{\Delta} \geq 0$ for arbitrary values of δ , but Δ may be either positive or negative (or vanishing in the μ - τ symmetry limit). When the 3σ intervals of θ_{12} , θ_{13} , θ_{23} and δ in Table 2.1 are taken into account, the upper bounds of $|\Delta|$ and $\overline{\Delta}$ can both reach about 0.1, implying that the flavor democracy of Φ_e^T , Φ_μ^T and Φ_τ^T may maximally be broken at the same level.

It is convenient to define three working observables at neutrino telescopes and connect them to the μ - τ symmetry-breaking quantities:

$$\begin{aligned}R_e &\equiv \frac{\Phi_e^T}{\Phi_\mu^T + \Phi_\tau^T} \simeq \frac{1}{2} - \frac{3}{2}\Delta , \\ R_\mu &\equiv \frac{\Phi_\mu^T}{\Phi_\tau^T + \Phi_e^T} \simeq \frac{1}{2} + \frac{3}{4}(\Delta + \overline{\Delta}) , \\ R_\tau &\equiv \frac{\Phi_\tau^T}{\Phi_e^T + \Phi_\mu^T} \simeq \frac{1}{2} + \frac{3}{4}(\Delta - \overline{\Delta}) .\end{aligned}\quad (6.17)$$

The small departure of R_α (for $\alpha = e, \mu, \tau$) from $1/2$ is therefore a clear measure of the overall effects of μ - τ symmetry breaking.

Now we turn to the flavor distribution of UHE cosmic neutrinos at a neutrino telescope by detecting the $\overline{\nu}_e$ flux from a very distant astrophysical source via the famous Glashow-resonance (GR) channel $\overline{\nu}_e e \rightarrow W^- \rightarrow \text{anything}$ [285], which happens over a narrow energy interval around $E_{\overline{\nu}_e}^{\text{GR}} \simeq M_W^2/2m_e \simeq 6.3$ PeV, and its cross section is about two orders of magnitude larger than those of $\overline{\nu}_e N$ interactions of the same $\overline{\nu}_e$ energy [286]. A measurement of the GR phenomenon is also important for another reason: it may serve as a sensitive discriminator of UHE cosmic neutrinos originating from $p\gamma$ and pp collisions [287, 288, 289, 290]. Let us assume that a neutrino telescope is able to observe both the GR-mediated $\overline{\nu}_e$ events and the $\nu_\mu + \overline{\nu}_\mu$ events of charged-current interactions in the vicinity of $E_{\overline{\nu}_e}^{\text{GR}}$. Then their ratio $R_{\text{GR}} \equiv \Phi_{\overline{\nu}_e}^T/\Phi_\mu^T$ can tell us something about the lepton flavor mixing.

To see this point more clearly, we start from the initial flavor distribution of UHE neutrinos at a cosmic accelerator: $\Phi_{\nu_e}^S : \Phi_{\overline{\nu}_e}^S : \Phi_{\nu_\mu}^S : \Phi_{\overline{\nu}_\mu}^S : \Phi_{\nu_\tau}^S : \Phi_{\overline{\nu}_\tau}^S = 1 : 1 : 2 : 2 : 0 : 0$ (pp collisions) or $1 : 0 : 1 : 1 : 0 : 0$ ($p\gamma$ collisions). Thanks to neutrino oscillations, the $\overline{\nu}_e$ flux at a neutrino telescope can be calculated by using the following formula:

$$\Phi_{\overline{\nu}_e}^T = \sum_i \sum_\beta |U_{ei}|^2 |U_{\beta i}|^2 \Phi_{\overline{\nu}_\beta}^S . \quad (6.18)$$

Since the expression of Φ_μ^T has been given in Eq. (6.13), it is straightforward to obtain R_{GR} for two different astrophysical sources:

$$R_{\text{GR}}(pp) \simeq \frac{1}{2} - \frac{3}{2}\Delta - \frac{1}{2}\overline{\Delta} ,$$

$$R_{\text{GR}}(p\gamma) \simeq \frac{\sin^2 2\theta_{12}}{4} - \frac{4 + \sin^2 2\theta_{12}}{4} \Delta - \frac{\sin^2 2\theta_{12}}{4} \overline{\Delta} + \frac{1 + \cos^2 2\theta_{12}}{2} \sin^2 \theta_{13} . \quad (6.19)$$

It is clear that the deviation of $R_{\text{GR}}(pp)$ from $1/2$ and that of $R_{\text{GR}}(p\gamma)$ from $\sin^2 2\theta_{12}/4$ are both governed by the effects of μ - τ symmetry breaking, which can maximally be of $\mathcal{O}(0.1)$. As discussed in the literature [46, 289], the IceCube detector running at the South Pole has a discovery potential to probe $R_{\text{GR}}(pp)$. In comparison, it is more challenging to detect the GR-mediated UHE $\overline{\nu}_e$ events originating from the pure $p\gamma$ collisions at a cosmic accelerator.

In the above discussions we have neglected the uncertainties associated with the initial neutrino fluxes at a given astrophysical source. A careful analysis of the flavor ratios of UHE cosmic neutrinos originating from pp or $p\gamma$ collisions actually leads us to the ratio $\Phi_e^S : \Phi_\mu^S : \Phi_\tau^S \simeq 1 : 2(1 - \eta) : 0$ with $\eta \simeq 0.08$ [291]. If this uncertainty is taken into account, then Eq. (6.13) will accordingly change to

$$\Phi_\alpha^T \simeq \frac{\Phi_0}{3} \left[1 + \sum_i |U_{\alpha i}|^2 (|U_{\mu i}|^2 - |U_{\tau i}|^2) - 2\eta \sum_i |U_{\alpha i}|^2 |U_{\mu i}|^2 \right] \quad (6.20)$$

for $\alpha = e, \mu$ and τ . This result implies that the η -induced correction is in general comparable with (or even larger than) the effect of μ - τ symmetry breaking. On the other hand, there may exist the uncertainties associated with the identification of different flavors at the neutrino telescope. Given the IceCube detector for example, the experimental error for determining the ratio of the muon track to the non-muon shower is typically $\xi \sim 20\%$ [292], depending on the event numbers. Such an estimate means that the ratio R_μ in Eq. (6.17) may in practice be contaminated by ξ , and this contamination is very likely to overwhelm the μ - τ symmetry breaking effect ($\Delta + \overline{\Delta}$) [282].

We have not considered other complexities and uncertainties associated with the origin of UHE cosmic neutrinos, such as their energy dependence, the effect of magnetic fields and possible new physics [293]. It remains too early to say that we have correctly understood the production mechanism of UHE cosmic rays and neutrinos from a given cosmic accelerator. But progress in determining the neutrino mixing parameters is so encouraging that it may finally allow us to well control the error bars from particle physics (e.g., the effect of μ - τ symmetry breaking) and thus focus on the unknowns from astrophysics (e.g., the initial flavor composition of UHE cosmic neutrinos [294]). Needless to say, any constraint on the flavor distribution of UHE cosmic neutrinos to be achieved from a neutrino telescope will be greatly useful in diagnosing the astrophysical sources and in understanding the properties of neutrinos themselves.

6.3. Matter-antimatter asymmetry via leptogenesis

Besides interpreting why the masses of the three known neutrinos are tiny in a qualitatively natural way, the canonical seesaw mechanism has another attractive feature — it can also provide a natural explanation of the observed baryon-antibaryon asymmetry of the Universe via the leptogenesis mechanism [295]: the CP-violating, lepton-number-violating and out-of-equilibrium decays of the hypothetical heavy Majorana neutrinos N_i may give rise to a lepton-antilepton asymmetry; and the latter is subsequently converted to the baryon-antibaryon asymmetry thanks to the sphaleron process [296]. The amount of the lepton-antilepton asymmetry depends on the CP-violating asymmetries ϵ_i between the decays of N_i and their CP-conjugate processes. If the masses of N_i have a strong hierarchy (i.e., $M_1 \ll M_2 \ll M_3$), then only ϵ_1 is expected to really matter at a temperature lower than M_1 . As a good approximation [297],

$$\epsilon_1 = -\frac{3}{16\pi(Y_\nu^\dagger Y_\nu)_{11}} \sum_{i \neq 1} \text{Im} [(Y_\nu^\dagger Y_\nu)_{1i}^2] \frac{M_1}{M_i}, \quad (6.21)$$

where Y_ν denotes the Yukawa coupling matrix of the neutrinos in the mass basis of N_i . With the help of $M_\nu = U D_\nu U^T$ in Eq. (2.10) and the seesaw formula $M_\nu = Y_\nu D_N^{-1} Y_\nu^T \langle H \rangle^2$ in the chosen basis, where $D_N \equiv \text{Diag}\{M_1, M_2, M_3\}$ and $\langle H \rangle = v/\sqrt{2} \simeq 174$ GeV, it is convenient to parametrize Y_ν in the following way [298]:

$$Y_\nu = \frac{1}{\langle H \rangle} U \sqrt{D_\nu} R \sqrt{D_N}, \quad (6.22)$$

in which R is a complex orthogonal matrix satisfying $RR^T = R^T R = 1$. Inserting Eq. (6.22) into Eq. (6.21), we arrive at

$$\epsilon_1 = -\frac{3}{16\pi} \frac{M_1}{\langle H \rangle^2} \frac{\text{Im}(\Delta m_{21}^2 R_{21}^{*2} + \Delta m_{31}^2 R_{31}^{*2})}{\sum_i m_i |R_{i1}|^2}, \quad (6.23)$$

where the orthogonality condition $R_{11}^2 + R_{21}^2 + R_{31}^2 = 0$ has been used. Eq. (6.23) shows that ϵ_1 is essentially independent of the PMNS matrix U , implying that *in general* there is no direct link between CP violation in N_i decays and that in low-energy neutrino oscillations [299]. If a certain flavor symmetry is taken into account to reduce the number of free parameters associated with the heavy and (or) light neutrinos, then R will get constrained to a simpler form so that the above expression of ϵ_1 can be simplified to some extent [300]. For instance, a model which can predict a constant neutrino mixing pattern is expected to result in a real diagonal R and thus a vanishing ϵ_1 [301, 302, 303, 304, 305]. This point is particularly transparent in the so-called “form-dominance” scenarios where Y_ν is simply taken as U multiplied by a diagonal matrix [306].

In fact, it is easy to see that the μ - τ permutation symmetry may have some interesting implications on the leptogenesis picture [307, 308, 309, 310, 311, 312, 313]. In the limit of such a flavor symmetry the Yukawa coupling matrix of the neutrinos

appears as [307, 308]

$$\tilde{Y}_\nu = \begin{pmatrix} y_{11} & y_{12} & y_{12} \\ y_{21} & y_{22} & y_{23} \\ y_{21} & y_{23} & y_{22} \end{pmatrix}, \quad (6.24)$$

and the heavy Majorana neutrino mass matrix M_N has the same form as the light Majorana neutrino mass matrix M_ν given in Eq. (3.7). One can make the transformation $U_N^T M_N U_N = D_N$, where $U_N = O_M U_{12}^N$ with O_M being the maximal mixing matrix:

$$O_M = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2} & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 1 & 1 \end{pmatrix}. \quad (6.25)$$

In this mass basis \tilde{Y}_ν becomes

$$\begin{aligned} Y_\nu &= \tilde{Y}_\nu U_N = U U^\dagger \tilde{Y}_\nu U_N = U U_{12}^\dagger O_M^T \tilde{Y}_\nu O_M U_{12}^N \\ &= U U_{12}^\dagger \begin{pmatrix} y_{11} & \sqrt{2}y_{12} & 0 \\ \sqrt{2}y_{21} & y_{22} + y_{23} & 0 \\ 0 & 0 & y_{22} - y_{23} \end{pmatrix} U_{12}^N, \end{aligned} \quad (6.26)$$

in which $U = O_M U_{12}$ is just the PMNS matrix of the three light Majorana neutrinos consisting of $\theta_{23} = \pi/4$. Comparing Eq. (6.22) with Eq. (6.26), we see that R is now subject to the (1,2) rotation subspace [307, 308]. In this case ϵ_1 is given by

$$\epsilon_1 = -\frac{3}{16\pi} \cdot \frac{M_1}{\langle H \rangle^2} \cdot \frac{\text{Im}(\Delta m_{21}^2 R_{21}^{*2})}{m_1 |R_{11}|^2 + m_2 |R_{21}|^2}. \quad (6.27)$$

With the help of this result and $R_{11}^2 + R_{21}^2 = 1$, it is easy to show that $|\epsilon_1|$ has an upper bound

$$|\epsilon_1| \leq \frac{3}{16\pi} \cdot \frac{M_1}{\langle H \rangle^2} |m_2 - m_1|. \quad (6.28)$$

Note that such an upper bound of ϵ_1 can be used to set a lower bound on M_1 if the leptogenesis mechanism works well in explaining the cosmological matter-antimatter asymmetry. In comparison with the upper bound of ϵ_1 obtained in a more general case, where m_2 in Eq. (6.28) ought to be replaced by m_3 [314], the present upper bound can be lowered by at least a factor $\sqrt{\Delta m_{21}^2 / \Delta m_{31}^2}$. When the effect of μ - τ flavor symmetry breaking is concerned, the situation will become more complicated and thus a careful analysis has to be done [308].

In the literature the so-called minimal seesaw model [315] has received some special attention due to its simplicity and stronger predictive power, so its connection to the leptogenesis mechanism deserves some discussions. Because there are only two heavy Majorana neutrinos in this model, M_N and \tilde{Y}_ν may read as follows [307, 308]:

$$M_N = \begin{pmatrix} M_{22} & M_{23} \\ M_{23} & M_{22} \end{pmatrix}, \quad \tilde{Y}_\nu = \begin{pmatrix} y_{12} & y_{12} \\ y_{22} & y_{23} \\ y_{23} & y_{22} \end{pmatrix}. \quad (6.29)$$

The transformation $Q_M^T M_N Q_M = D_N$, where

$$Q_M = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}, \quad (6.30)$$

leads us to the mass eigenvalues $M_2 = M_{22} + M_{23}$ and $M_3 = M_{22} - M_{23}$. In the mass basis the Yukawa coupling matrix \tilde{Y}_ν becomes

$$Y_\nu = \tilde{Y}_\nu Q_M = \frac{1}{\sqrt{2}} \begin{pmatrix} 2y_{12} & 0 \\ y_{22} + y_{23} & y_{23} - y_{22} \\ y_{22} + y_{23} & y_{22} - y_{23} \end{pmatrix}. \quad (6.31)$$

It is interesting to see that the two columns of Y_ν are orthogonal to each other, and thus $Y_\nu^\dagger Y_\nu$ must be diagonal and ϵ_1 must be vanishing. So a finite value of ϵ_1 has to come from the μ - τ permutation symmetry breaking. Let us take a simple example to illustrate the effect of μ - τ symmetry breaking and its connection to CP violation in the decays of two heavy Majorana neutrinos. To be explicit, we assume M_N and \tilde{Y}_ν to be real and introduce the symmetry breaking term as

$$\tilde{Y}'_\nu = \tilde{Y}_\nu + \begin{pmatrix} -y'_{12} & y'_{12} \\ 0 & 0 \\ 0 & 0 \end{pmatrix}, \quad (6.32)$$

in which y'_{12} is complex so as to accommodate CP violation. In the mass basis we obtain

$$Y'_\nu = \tilde{Y}'_\nu Q_M = \begin{pmatrix} r_1 a & r_2 b \\ a & -b \\ a & b \end{pmatrix}, \quad (6.33)$$

where $a \equiv (y_{22} + y_{23})/\sqrt{2}$, $b \equiv (y_{22} - y_{23})/\sqrt{2}$, $r_1 \equiv \sqrt{2}y_{12}/a$ and $r_2 \equiv \sqrt{2}y'_{12}/b$ are defined. The parameter r_2 is therefore responsible for the generation of θ_{13} and CP violation. It is found that the experimental data at low energies can be reproduced when one assumes $r_1 \simeq 1$, $|r_2| \simeq \sqrt{2}\theta_{13}$, $3a^2\langle H \rangle^2/M_2 = m_2$ and $2b^2\langle H \rangle^2/M_3 = m_3$ [316]. In this case the CP-violating asymmetry between the decay of the lighter heavy Majorana neutrino (with mass M_3) and its CP-conjugate process can approximate to

$$\epsilon = \frac{b^2}{16\pi} \cdot \frac{m_2}{m_3} \text{Im}(r_2^2) \sim 10^{-4} \sin 2\phi, \quad (6.34)$$

where $\phi \equiv \arg(r_2)$ and $b \sim \mathcal{O}(1)$ has been taken into account. This estimate implies that the strength of μ - τ symmetry breaking required by the observed value of θ_{13} is sufficient for a successful leptogenesis mechanism to work in such a minimal seesaw scenario [308].

When the μ - τ reflection symmetry is imposed on both the Yukawa coupling matrix \tilde{Y}_ν and the heavy Majorana neutrino matrix M_N , one has

$$\tilde{Y}_\nu = \begin{pmatrix} y_{11} & y_{12} & y_{12}^* \\ y_{21} & y_{22} & y_{23} \\ y_{21}^* & y_{23}^* & y_{22}^* \end{pmatrix} \quad (6.35)$$

with y_{11} being real, and the form of M_N is the same as that given in Eq. (3.24). In this case M_N can be diagonalized by a unitary matrix U_N , whose complex conjugate is of the form shown in Eq. (3.19). Given the Yukawa coupling matrix $Y_\nu = \tilde{Y}_\nu U_N$ in the mass basis of N_i , a straightforward calculation shows that $Y_\nu^\dagger Y_\nu$ is actually a real matrix which prohibits CP violation [113]. To make the leptogenesis idea viable, one

may choose to softly break the μ - τ reflection symmetry [317]. Another way out is to keep the μ - τ reflection symmetry but take into account the so-called flavor effects to make the leptogenesis mechanism work [232]. If the mass of the lightest heavy Majorana neutrino N_1 lies in the range $10^9 \text{ GeV} < M_1 < 10^{12} \text{ GeV}$, the τ leptons can be in thermal equilibrium, making them distinguishable from the e and μ flavors. In this situation the decay of N_1 to the τ flavor should be treated separately from the decays of N_1 to the other two flavors [318, 319]. Accordingly, the *flavored* CP-violating asymmetries associated with the decays of N_1 are given by

$$\epsilon_1^\alpha = -\frac{1}{8\pi} \cdot \frac{M_1}{\langle H \rangle^2} \sum_{i \neq 1} \text{Im} \left[\frac{3}{2} (Y_\nu^*)_{\alpha 1} (Y_\nu)_{\alpha i} (Y_\nu^\dagger Y_\nu)_{1i} \frac{M_1}{M_i} - (Y_\nu^*)_{\alpha 1} (Y_\nu)_{\alpha i} (Y_\nu^\dagger Y_\nu)_{1i}^* \frac{M_1^2}{M_i^2} \right], \quad (6.36)$$

in which α runs over e , μ and τ . After a straightforward calculation, one can obtain $\epsilon_1^e = 0$ and $\epsilon_1^\mu = -\epsilon_1^\tau$ as a direct consequence of the μ - τ reflection symmetry. This result confirms the vanishing of the overall CP-violating asymmetry (i.e., $\epsilon_1 = \epsilon_1^e + \epsilon_1^\mu + \epsilon_1^\tau = 0$). In spite of $\epsilon_1 = 0$, the lepton-antilepton asymmetry produced from the decays of N_1 is still likely to survive if such processes took place in the temperature range $10^9 \text{ GeV} < T = M_1 < 10^{12} \text{ GeV}$ in which the flavor effects should take effect [232]. When the temperature of the Universe dropped below 10^9 GeV , however, the μ leptons would also be in thermal equilibrium. In this case the simple leptogenesis scenario under discussion would not work anymore if $M_1 < 10^9 \text{ GeV}$ held.

6.4. Fermion mass matrices with the Z_2 symmetry

Being capable of relating the smallness of quark flavor mixing angles to the smallness of quark mass ratios in a way similar to the GST relation, the Fritzsch texture-zero ansatz is a simple and instructive example for studying the possible underlying flavor structures of three quark families [320, 321]:

$$M_q = \begin{pmatrix} 0 & c_q & 0 \\ c_q^* & 0 & b_q \\ 0 & b_q^* & a_q \end{pmatrix}, \quad (6.37)$$

where $q = u$ (up) or d (down), and the Hermiticity has been assumed to reduce the number of free parameters. Given the strong quark mass hierarchies as shown in Fig. 2.2, M_q is expected to have a strongly hierarchical structure (i.e., $|c_q| \ll |b_q| \ll |a_q|$). Unfortunately, such a simple texture has been ruled out mainly because its phenomenological consequences cannot simultaneously fit the small size of V_{cb} and the large value of m_t . One way out is to introduce an additional free parameter to make the (2, 2) entry of M_q nonzero [322, 323, 324]:

$$M_q = \begin{pmatrix} 0 & c_q & 0 \\ c_q^* & d_q & b_q \\ 0 & b_q^* & a_q \end{pmatrix}. \quad (6.38)$$

Then it is easy to check that this new ansatz, which may have a more or less weaker hierarchy than its original counterpart, can be in good agreement with the experimental data on quark flavor mixing and CP violation [325, 326]. Although the Fritzsch-like texture in Eq. (6.38) is apparently different from the pattern of M_ν with a μ - τ flavor symmetry (i.e., the former contains a few texture zeros while the latter is characterized by some linear relations or equalities of different entries), it is possible to find a flavor basis where M_q may essentially have the same form as M_ν . Here we consider a special but interesting possibility that the lepton and quark sectors share a permutation symmetry between their respective second and third families. For simplicity, we assume that the relevant mass matrices M_f (for $f = u, d, l, \nu$) take the following form [327, 328, 329, 330, 331, 332]

$$M_f = P_f^\dagger \widehat{M}_f P_f = P_f^\dagger \begin{pmatrix} A_f & B_f & B_f \\ B_f & C_f & D_f \\ B_f & D_f & C_f \end{pmatrix} P_f \quad (6.39)$$

with \widehat{M}_f being real and $P_f \equiv \text{Diag}\{e^{i\phi_1^f}, e^{i\phi_2^f}, e^{i\phi_3^f}\}$. Obviously, the (2, 3) permutation symmetry requires $\phi_2^f = \phi_3^f$. If the massive neutrinos are the Majorana particles, however, the P_ν matrix on the right-hand side of \widehat{M}_ν in Eq. (6.39) should be replaced by P_ν^* . The unitary matrices used to diagonalize M_f can be universally expressed as

$$O_f = P_f^\dagger O_{23}^f O_{12}^f, \quad (6.40)$$

where O_{23}^f has the same form as O_M in Eq. (6.25), and O_{12}^f contains an unspecified rotation angle θ_{12}^f .

Now let us pay particular attention to the $A_q = 0$ case, where θ_{12}^q can be connected to the quark mass ratio via $\tan \theta_{12}^q = \sqrt{m_1^q/m_2^q}$. Hence it is easy to obtain the GST relation. In view of Eq. (6.40), we find that the CKM quark mixing matrix $V_{\text{CKM}} = O_u^\dagger O_d$ only contains one nonzero mixing angle (i.e., the Cabibbo angle $\theta_{12}^{\text{CKM}} \equiv \theta_C$) in the (2, 3) permutation symmetry limit. The vanishing of θ_{13}^{CKM} and θ_{23}^{CKM} in this scenario qualitatively agrees with the experimental fact that these two angles are very small: $\theta_{13}^{\text{CKM}} \simeq 0.2^\circ$ and $\theta_{23}^{\text{CKM}} \sim 2^\circ$ [16]. So their finite values can be ascribed to the small (2, 3) permutation symmetry breaking [333]. Similar to the μ - τ permutation symmetry, the (2, 3) permutation symmetry can be broken by allowing

$$\widehat{M}_{12}^f \neq \widehat{M}_{13}^f, \quad \widehat{M}_{22}^f \neq \widehat{M}_{33}^f, \quad \phi_2^f \neq \phi_3^f. \quad (6.41)$$

If the symmetry breaking only comes from the phase parameters (i.e., $\phi_2^f \neq \phi_3^f$), then θ_{23}^{CKM} can fit its experimental value by taking $\phi_{32}^d - \phi_{32}^u \sim 0.08$, where $\phi_{32}^q \equiv \phi_3^q - \phi_2^q$ is defined. On the other hand, θ_{13}^{CKM} can be calculated via the correlation $\theta_{13}^{\text{CKM}} \sim \theta_{12}^u \theta_{23}^{\text{CKM}}$, but the result is somewhat smaller than the observed value [326]. In this case one may either give up $A_q = 0$ or break the (2, 3) permutation symmetry of \widehat{M}_q in a slightly stronger way. The former approach may not make a realization of the GST relation self-evident. Following the latter approach, we can obtain nonzero θ_{13}^q by allowing $\widehat{M}_{12}^q \neq \widehat{M}_{13}^q$ but keeping $\widehat{M}_{22}^q = \widehat{M}_{33}^q$. Note that one may simply take $\widehat{M}_{13}^q = 0$ and arrive

at the following zero texture of fermion mass matrices:

$$M_f = P_f^\dagger \widehat{M}_f P_f = P_f^\dagger \begin{pmatrix} 0 & B_f & 0 \\ B_f & C_f & D_f \\ 0 & D_f & C_f \end{pmatrix} P_f. \quad (6.42)$$

Then it is easy to show that θ_{13}^u is vanishingly small and θ_{13}^d can be given by $\theta_{13}^d \simeq \sqrt{m_d m_s / m_b^2}$, which is large enough to generate a phenomenologically acceptable value of θ_{13}^{CKM} [326]. Since the fermion mass matrices in Eq. (6.42) have the same structure as those quark mass matrices in Eq. (6.38) if $a_q = d_q$ is taken, a universal treatment of lepton and quark flavor mixing issues seems quite natural in this sense.

We proceed to discuss some phenomenological consequences of Eq. (6.39) in the lepton sector. Above all, the experimental result $\theta_{23} \simeq \pi/4$ signifies the strong (2,3) permutation symmetry breaking. If the symmetry is only broken by $\phi_2^f \neq \phi_3^f$, then the phase difference $\phi_{32}^l - \phi_{32}^\nu$ has to be very close to $\pm\pi/2$ in order to avoid a significant cancellation between the charged-lepton and neutrino sectors. To be explicit, we focus on the texture in Eq. (6.42) to obtain more definite predictions. We have $\theta_{12}^l \simeq \sqrt{m_e/m_\mu} \simeq 0.07$, and θ_{13}^l is highly suppressed due to $m_e \ll m_\mu \ll m_\tau$. So θ_{12}^ν should take the dominant responsibility for $\theta_{12} \simeq 34^\circ$, implying $\theta_{12}^\nu \simeq \theta_{12}$ as a good approximation. Taking $\tan \theta_{12}^\nu \simeq \sqrt{m_1/m_2}$ in a parallel way, one can roughly fix the three neutrino masses: $m_1 \sim 0.008$ eV, $m_2 \sim 0.011$ eV and $m_3 \sim 0.051$ eV. In addition, $\theta_{13}^\nu \sim \sqrt{m_1 m_2 / m_3^2} \simeq 0.18$, and hence the correct value of θ_{13} can be achieved after the contribution from θ_{12}^l (ranging between -0.05 and 0.05 , as a function of $\phi_{21}^l - \phi_{21}^\nu$) is properly included. Finally, it is also possible to obtain $\delta \simeq \pm\pi/2$.

At this point it is worth mentioning the idea of “flavor democracy” [334, 335] or “universal strength of Yukawa couplings” [336] for the mass matrices of leptons and quarks, which can be regarded as an extreme case of M_f shown in Eq. (6.39) after they receive some proper perturbations respecting the Z_2 permutation symmetry. While the charged-lepton mass hierarchy, the quark mass hierarchy and small quark flavor mixing effects are easily understood in such an approach, one has to constrain the neutrino mass matrix M_ν to be nearly diagonal so as to obtain significant flavor mixing effects in the lepton sector [337]. It is also found that a combination of the flavor democracy idea and the seesaw mechanism allows one to build a phenomenologically viable model [338].

To summarize, the universal texture of fermion mass matrices shown in Eq. (6.39), which possesses the (2,3) or Z_2 permutation symmetry — an analogue of the μ - τ permutation symmetry, can serve as a starting point to describe the flavor structures of quarks and leptons and to identify their similarities and differences. Its variation with some symmetry-breaking effects in Eq. (6.42) contains both the texture zeros and the linear equality of different entries [339, 340]. Such an ansatz is predictive and can essentially fit current experimental data. For simplicity, here we have only discussed the possibility that all the fermion mass matrices share the same texture but their respective parameters are independent of one another. The latter can get correlated if the GUT

framework is taken into account (see, e.g., Refs. [341, 342], where the $(2, 3)$ permutation symmetry is combined with the $SU(5)$ and $SO(10)$ GUT models, respectively).

7. Summary and outlook

In the past two decades we have witnessed a booming period in neutrino physics — an extremely important part of particle physics and cosmology. Especially since the first discovery of atmospheric neutrino oscillations at the Super-Kamiokande detector in 1998, quite a lot of significant breakthroughs have been made in experimental neutrino physics, as recognized by both the 2015 Nobel Prize in Physics and the 2016 Breakthrough Prize in Fundamental Physics. On the one hand, the striking and appealing phenomena of atmospheric, solar, reactor and accelerator neutrino (or antineutrino) oscillations have all been observed in a convincing way, and the oscillation parameters Δm_{21}^2 , $|\Delta m_{31}^2|$, θ_{12} , θ_{13} and θ_{23} have been determined to an impressive degree of accuracy. On the other hand, the unusual geo-antineutrino events and extraterrestrial PeV neutrino events have also been observed, and the sensitivities to neutrino masses in the beta decays, $0\nu 2\beta$ decays and cosmological observations have been improved to a great extent. On the theoretical side, a lot of efforts have been made towards a deeper understanding of the origin of tiny neutrino masses, the secret of flavor mixing and CP violation and a unified picture of leptons and quarks at a much larger framework beyond the SM. Moreover, one has explored various implications of massive neutrinos on the cosmological matter-antimatter asymmetry, warm dark matter and many violent astrophysical or astronomical processes. All these have demonstrated neutrino physics to be one of the most important and exciting frontiers in modern science.

What the present review article has concentrated on is the “minimal flavor symmetry” behind the observed lepton flavor mixing pattern — a partial (or approximate) μ - τ symmetry which is definitely favored by current experimental data, and its various phenomenological implications in neutrino physics. We have discussed both the μ - τ *permutation* symmetry and the μ - τ *reflection* symmetry, and pointed out a few typical ways to slightly break such a symmetry. Some larger discrete flavor symmetry groups, in which the intriguing μ - τ symmetry can naturally arise as a residual symmetry, have been briefly described. Both the bottom-up approach and the top-down approach have been illustrated in this connection, in order to bridge the gap between model building attempts and neutrino oscillation data. To be more explicit, we have summarized the basic strategies of model building with the help of the μ - τ symmetry, either in the presence or in the absence of the seesaw mechanism. The phenomenological consequences of the μ - τ flavor symmetry on some interesting and important topics, such as neutrino oscillations in matter, radiative corrections to the equalities $|U_{\mu i}| = |U_{\tau i}|$ from a superhigh-energy scale down to the electroweak scale, flavor distributions of the UHE cosmic neutrinos at a neutrino telescope, a possible connection between the leptogenesis and low-energy CP violation through the seesaw mechanism and a unified flavor structure of leptons and quarks, have also been discussed.




Figure 7.1. The Fritzsche-Xing “pizza” plot of 28 parameters associated with the SM itself and neutrino masses, lepton flavor mixing angles and CP-violating phases.

Therefore, we hope that this review could serve as a new milestone in the development of neutrino phenomenology, in order to make so many theoretical ideas of ours much deeper and more convergent.

There are still quite a number of open fundamental questions about massive neutrinos and lepton flavor issues. The immediate ones include how small the absolute neutrino mass scale is, whether the neutrino mass spectrum is normal as those of the charged leptons and quarks, whether the antiparticles of massive neutrinos are just themselves, how large the CP-violating phase δ can really be, which octant the largest flavor mixing angle θ_{23} belongs to, whether there are light and (or) heavy sterile neutrinos, what the role of neutrinos is in dark matter, whether the observed matter-antimatter asymmetry of the Universe is directly related to CP violation in neutrino oscillations, etc. Motivated by so many challenging and exciting questions, we are on our way to discovering a new physics world in the coming decades [†].

Last but not least, let us make some concluding remarks with the help of the Fritzsche-Xing “pizza” plot shown in Fig. 7.1. This picture provides a brief summary of 28 fundamental parameters associated with the SM itself and neutrino masses, lepton flavor mixing angles and CP-violating phases. Among them, the five parameters of strong and weak CP violation deserve some special attention. In the quark sector the strong CP-violating phase $\bar{\theta}$ remains a mystery, but the weak CP-violating phase δ_q has been determined to a good degree of accuracy. In the lepton sector none of the CP-violating phases has been directly measured, although a very preliminary hint $\delta_l \sim -\pi/2$ has been achieved from a comparison between the present T2K and Daya Bay data [‡].

[†] See some recent reviews on the future prospects of leptonic CP violation [343], neutrino oscillation experiments [80], searches for the $0\nu 2\beta$ decays [345], electromagnetic properties of massive neutrinos [346] and sterile neutrino physics [266].

[‡] Here we use δ_l to denote the Dirac CP phase in the standard parametrization of the PMNS matrix U , so as to distinguish it from its analogue δ_q in the CKM quark flavor mixing matrix.

The true value of δ_l is expected to be determined in the future long-baseline neutrino oscillation experiments. If massive neutrinos are really the Majorana particles, however, it will be extremely challenging to probe or constrain the Majorana CP-violating phases ρ and σ which can only show up in some extremely rare processes involving lepton number nonconservation.

Perhaps some of the flavor puzzles cannot be resolved unless we finally find out the fundamental flavor theory [347]. But the latter cannot be formulated without a lot of phenomenological and experimental inputs. As Leonardo da Vinci once stressed, “Although nature commences with reason and ends in experience, it is necessary for us to do the opposite. That is, to commence with experience and from this to proceed to investigate the reason.”

We would like to thank Leszek Roszkowski for inviting us to write this review article. One of us (Z.Z.X.) is indebted to Harald Fritzsch, Shu Luo, He Zhang, Shun Zhou and Ye-Ling Zhou for fruitful collaboration on the μ - τ flavor symmetry issues. We are also grateful to Wan-lei Guo, Yu-Feng Li, Jue Zhang and Jing-yu Zhu for their kind helps in plotting several figures. This work is supported in part by the National Natural Science Foundation of China under grant No. 11135009 and grant No. 11375207 (Z.Z.X.); by the National Basic Research Program of China under grant No. 2013CB834300 (Z.Z.X.); by the China Postdoctoral Science Foundation under grant No. 2015M570150 (Z.H.Z.); and by the CAS Center for Excellence in Particle Physics.

8. Appendix

As discussed in section 3.4, the delicate effects of μ - τ reflection symmetry breaking are characterized by the quantities $\Delta\theta_{23}$, $\Delta\delta$, $\Delta\rho$ and $\Delta\sigma$. Their expressions in our analytical approximations can be parametrized as

$$\begin{aligned}\Delta\theta_{23} &= c_{r1}^\theta R_1 + c_{i1}^\theta I_1 + c_{r2}^\theta R_2 + c_{i2}^\theta I_2 , \\ \Delta\delta &= c_{r1}^\delta R_1 + c_{i1}^\delta I_1 + c_{r2}^\delta R_2 + c_{i2}^\delta I_2 , \\ \Delta\rho &= c_{r1}^\rho R_1 + c_{i1}^\rho I_1 + c_{r2}^\rho R_2 + c_{i2}^\rho I_2 , \\ \Delta\sigma &= c_{r1}^\sigma R_1 + c_{i1}^\sigma I_1 + c_{r2}^\sigma R_2 + c_{i2}^\sigma I_2 ,\end{aligned}\tag{8.1}$$

where the relevant coefficients are given by

$$\begin{aligned}c_{r1}^\theta &= 2 (m_{12}^2 - m_{11}^2 s_{13}^2 + m_3^2 s_{13}^2) \Omega_\theta , \\ c_{i1}^\theta &= 4m_{11}m_{12}\tilde{s}_{13}\Omega_\theta , \\ c_{r2}^\theta &= -m_{11-3}m_{22+3}\Omega_\theta , \\ c_{i2}^\theta &= 4m_{11}m_{12}\tilde{s}_{13}\Omega_\theta ;\end{aligned}\tag{8.2}$$

and

$$\begin{aligned}c_{r1}^\delta &= 2 [m_{12} (\overline{m}_1 + \overline{m}_2) m_{22-3} T \\ &\quad + 2m_{11-3}m_{11+3}m_{22-3}s_{13}^2 \\ &\quad - 2m_{12}^2(2m_{11+3} + m_{12}T)s_{13}^2] \Omega_\delta ,\end{aligned}$$

$$\begin{aligned}
c_{i1}^\delta &= 2 \left[(\overline{m}_1 + \overline{m}_2) m_{11+3} m_{22-3} T \right. \\
&\quad \left. - 2m_{12} (\overline{m}_1 - m_3) (\overline{m}_2 - m_3) \right] \tilde{s}_{13} \Omega_\delta , \\
c_{r2}^\delta &= \left[-m_{12} (\overline{m}_1 + \overline{m}_2) m_{22+3} T \right. \\
&\quad \left. - 2m_{12} (m_{12}^2 - 2m_{22} m_{11-3}) T s_{13}^2 \right. \\
&\quad \left. + 2m_{11-3} m_{11+3} m_{22+3} s_{13}^2 \right] \Omega_\delta , \\
c_{i2}^\delta &= m_{22+3} \left[m_{12}^2 - m_{22-3} (2m_{11} \right. \\
&\quad \left. + m_{22-3}) \right] T \tilde{s}_{13} \Omega_\delta ;
\end{aligned} \tag{8.3}$$

and

$$\begin{aligned}
c_{r1}^\rho &= 8m_{11} m_3 \left[m_{12}^2 (m_{11+3} + m_{12} T) \right. \\
&\quad \left. - (\overline{m}_1 - m_3) (\overline{m}_2 - m_3) m_{11+3} \right] c_{12}^2 \tilde{s}_{13} \Omega_\rho , \\
c_{i1}^\rho &= -8m_{11} m_{12} m_3 \left[(\overline{m}_1 - m_3) (\overline{m}_2 - m_3) \right. \\
&\quad \left. - 2m_{11} m_{11+3} s_{13}^2 \right] c_{12}^2 \Omega_\rho , \\
c_{r2}^\rho &= 4m_{11} m_3 \left[m_{12} (m_{12}^2 - 2m_{22} m_{11-3}) T \right. \\
&\quad \left. - m_{11-3} m_{11+3} m_{22+3} \right] c_{12}^2 \tilde{s}_{13} \Omega_\rho , \\
c_{i2}^\rho &= (\overline{m}_1 - m_3) (\overline{m}_2 - m_3) m_{22+3} \left[2m_{11} m_3 c_{12}^2 \right. \\
&\quad \left. + (\overline{m}_1 + \overline{m}_2) (\overline{m}_1 s_{12}^2 - \overline{m}_1 c_{12}^2 \right. \\
&\quad \left. - m_3 s_{12}^2) \right] T \Omega_\rho ;
\end{aligned} \tag{8.4}$$

and

$$\begin{aligned}
c_{r1}^\sigma &= -8m_{11} m_3 \left[m_{12}^2 (m_{11+3} + m_{12} T) \right. \\
&\quad \left. - (\overline{m}_1 - m_3) (\overline{m}_2 - m_3) m_{11+3} \right] s_{12}^2 \tilde{s}_{13} \Omega_\sigma , \\
c_{i1}^\sigma &= 8m_{11} m_{12} m_3 \left[(\overline{m}_1 - m_3) (\overline{m}_2 - m_3) \right. \\
&\quad \left. - 2m_{11} m_{11+3} s_{13}^2 \right] s_{12}^2 \Omega_\sigma , \\
c_{r2}^\sigma &= -4m_{11} m_3 \left[m_{12} (m_{12}^2 - 2m_{22} m_{11-3}) T \right. \\
&\quad \left. - m_{11-3} m_{11+3} m_{22+3} \right] s_{12}^2 \tilde{s}_{13} \Omega_\sigma , \\
c_{i2}^\sigma &= -(\overline{m}_1 - m_3) (\overline{m}_2 - m_3) m_{22+3} \left[2m_{11} m_3 s_{12}^2 \right. \\
&\quad \left. + (\overline{m}_1 + \overline{m}_2) (\overline{m}_1 c_{12}^2 - \overline{m}_1 s_{12}^2 \right. \\
&\quad \left. - m_3 c_{12}^2) \right] T \Omega_\sigma .
\end{aligned} \tag{8.5}$$

In Eqs. (8.2)—(8.5), $T = \tan 2\theta_{12}$, and Ω_θ , Ω_δ , Ω_ρ and Ω_σ are defined through the equations

$$\begin{aligned}
\Omega_\theta^{-1} &= 2 (\overline{m}_1 - m_3) (\overline{m}_2 - m_3) , \\
\Omega_\delta^{-1} &= 2 (\overline{m}_1 + \overline{m}_2) (\overline{m}_1 - m_3) (\overline{m}_2 - m_3) T \tilde{s}_{13} , \\
\Omega_\rho^{-1} &= 4\overline{m}_1 m_3 c_{12} s_{12} (\Omega_\delta T \tilde{s}_{13})^{-1} , \\
\Omega_\sigma^{-1} &= 4\overline{m}_2 m_3 c_{12} s_{12} (\Omega_\delta T \tilde{s}_{13})^{-1} .
\end{aligned} \tag{8.6}$$

- [1] Becquerel H 1896 On the rays emitted by phosphorescence *Compt. Rend. Math.* **122** 420–421
- [2] Chadwick J 1914 The intensity distribution in the magnetic spectrum of beta particles from radium (B+C) *Verh. Deut. Phys. Gesell.* **16** 383–391
- [3] Ellis C D and Wooster W A 1927 Average energy of disintegration of radium E *Proc. Roy. Soc. London* **A117**, 109–
- [4] Fermi E 1934 An attempt of a theory of beta radiation *Z. Phys.* **88** 161–177
- [5] Bethe H A and Bacher R F 1936 Nuclear physics A. Stationary states of nuclei *Rev. Mod. Phys.* **8** 82–229
- [6] Pontecorvo B 1946 Inverse beta process *Chalk River Lab. Report PD-205*
- [7] Cowan C L, Reines F, Harrison F B, Kruse H W and McGuire A D 1956 Detection of the free neutrino: a confirmation *Science* **124** 103–104
- [8] Pontecorvo B 1957 Mesonium and anti-mesonium *Sov. Phys. JETP* **6** 429–431
- [9] Majorana E 1937 Theory of the symmetry of electrons and positrons *Nuovo Cim.* **14** 171–184
- [10] Danby G *et al* 1962 Observation of high-energy neutrino reactions and the existence of two kinds of neutrinos *Phys. Rev. Lett.* **9** 36–44
- [11] Maki Z, Nakagawa M and Sakata S 1962 Remarks on the unified model of elementary particles *Prog. Theor. Phys.* **28** 870–880
- [12] Kodama K *et al* (DONUT Collaboration) 2001 Observation of tau neutrino interactions *Phys. Lett.* **B504** 218–224 (arXiv:hep-ex/0012035)
- [13] Weinberg S 1967 A model of leptons *Phys. Rev. Lett.* **19** 1264–1266
- [14] Davis R, Harmer D S and Hoffman K C 1968 Search for neutrinos from the sun *Phys. Rev. Lett.* **20** 1205–1209
- [15] Gribov V and Pontecorvo B 1969 Neutrino astronomy and lepton charge *Phys. Lett.* **28B** 493–496
- [16] Olive K A *et al* (Particle Data Group) 2014 Review of Particle Physics *Chin. Phys.* **C38** 090001
- [17] Minkowski P 1977 $\mu \rightarrow e\gamma$ at a rate of one out of 10^9 muon decays *Phys. Lett.* **B67** 421–428
- [18] Yanagida T 1979 Horizontal symmetry and masses of neutrinos *Conf. Proc.* **C7902131** 95–99
- [19] Gell-Mann M, Ramond P and Slansky R 1979 Complex spinors and unified theories *Conf. Proc.* **C790927** 315–321
- [20] Glashow S L 1980 The future of elementary particle physics *NATO Sci. Ser.* **B61** 687–713
- [21] Mohapatra R N and Senjanovic G 1980 Neutrino mass and spontaneous parity violation *Phys. Rev. Lett.* **44** 912–915
- [22] Schechter J and Valle J W F 1980 Neutrino masses in $SU(2) \times U(1)$ theories *Phys. Rev.* **D22** 2227–2235
- [23] Cheng T P and Li L F 1980 Neutrino masses, mixings and oscillations in $SU(2) \times U(1)$ models of electroweak interactions *Phys. Rev.* **D22** 2860–2868
- [24] Magg M and Wetterich C 1980 Neutrino mass problem and gauge hierarchy *Phys. Lett.* **B94** 61–64
- [25] Lazarides G, Shafi Q and Wetterich C 1981 Proton lifetime and fermion masses in an $SO(10)$ model *Nucl. Phys.* **B181** 287–300
- [26] Mohapatra R N and Senjanovic G 1981 Neutrino masses and mixings in gauge models with spontaneous parity violation *Phys. Rev.* **D23** 165–180
- [27] Foot R, Lew H, He X G and Joshi G C 1989 Seesaw neutrino masses induced by a triplet of leptons *Z. Phys.* **C44** 441–444
- [28] Ma E 1998 Pathways to naturally small neutrino masses *Phys. Rev. Lett.* **81** 1171–1174
- [29] Weinberg S 1979 Baryon and lepton nonconserving processes *Phys. Rev. Lett.* **43** 1566–1570
- [30] Noether E 1918 Invariant variation problems *Gott. Nachr.* **1918** 235–257
- [31] Fritzsch H and Xing Z Z 2000 Mass and flavor mixing schemes of quarks and leptons *Prog. Part. Nucl. Phys.* **45** 1–81 (arXiv:hep-ph/9912358)
- [32] Altarelli G and Feruglio F 2010 Discrete flavor symmetries and models of neutrino mixing *Rev. Mod. Phys.* **82** 2701–2729 (arXiv:1002.0211)
- [33] King S F and Luhn C 2013 Neutrino mass and mixing with discrete symmetry *Rept. Prog. Phys.* **76** 056201 (arXiv:1301.1340)

- [34] King S F, Merle A, Morisi S and Shimizu Y 2014 Neutrino masses and mixing: from theory to experiment *New J. Phys.* **16** 045018 (arXiv:1402.4271)
- [35] Xing Z Z and Zhou S 2014 A partial μ - τ symmetry and its prediction for leptonic CP violation *Phys. Lett.* **B737** 196–200 (arXiv:1404.7021)
- [36] Xing Z Z 2009 Naturalness and testability of TeV seesaw mechanisms *Prog. Theor. Phys. Suppl.* **180** 112–127 (arXiv:0905.3903)
- [37] Antusch S, Biggio C, Fernandez-Martinez E, Gavela M B and Lopez-Pavon J 2006 Unitarity of the leptonic mixing matrix *JHEP* **0610** 084 (arXiv:hep-ph/0607020)
- [38] Antusch S and Fischer O 2014 Non-unitarity of the leptonic mixing matrix: present bounds and future sensitivities *JHEP* **1410** 94 (arXiv:1407.6607)
- [39] Fritzsch H and Xing Z Z 2001 How to describe neutrino mixing and CP violation *Phys. Lett.* **B517** 363–368 (arXiv:hep-ph/0103242)
- [40] Xing Z Z and Zhou S 2011 *Neutrinos in Particle Physics, Astronomy and Cosmology* (Berlin: Springer-Verlag) p 162
- [41] Jarlskog C 1985 Commutator of the quark mass matrices in the standard electroweak model and a measure of maximal CP violation *Phys. Rev. Lett.* **55** 1039–1042
- [42] Wolfenstein L 1978 Neutrino oscillations in matter *Phys. Rev.* **D17** 2369–2374
- [43] Mikheyev S P and Smirnov A Yu 1985 Resonance amplification of oscillations in matter and spectroscopy of solar neutrinos *Sov. J. Nucl. Phys.* **42** 913–917
- [44] Xing Z Z 2004 Flavor mixing and CP violation of massive neutrinos *Int. J. Mod. Phys.* **A19** 1–80 (arXiv:hep-ph/0307007)
- [45] Bahcall J N, Bahcall N A and Shaviv G 1968 Present status of the theoretical predictions for the Cl-36 solar neutrino experiment *Phys. Rev. Lett.* **20** 1209–1212
- [46] Hampel W *et al* (GALLEX Collaboration) 1999 GALLEX solar neutrino observations: results for GALLEX IV *Phys. Lett.* **B447** 127–133
- [47] Altmann M *et al* (GNO Collaboration) 2000 GNO solar neutrino observations: results for GNO I *Phys. Lett.* **B490** 16–26 (arXiv:hep-ex/0006034)
- [48] Abdurashitov J N *et al* (SAGE Collaboration) 2002 Solar neutrino flux measurements by the Soviet-American Gallium Experiment (SAGE) for half the 22 year solar cycle *J. Exp. Theor. Phys.* **95** 181–193 (arXiv:astro-ph/0204245)
- [49] Fukuda Y *et al* (Super-Kamiokande Collaboration) 1998 Measurements of the solar neutrino flux from Super-Kamiokande’s first 300 days *Phys. Rev. Lett.* **81** 1158–1162 (arXiv:hep-ex/9805021)
- [50] Ahmad Q R *et al* (SNO Collaboration) 2002 Direct evidence for neutrino flavor transformation from neutral current interactions in the Sudbury Neutrino Observatory *Phys. Rev. Lett.* **89** 011301 (arXiv:nucl-ex/0204008)
- [51] Aharmim B *et al* (SNO Collaboration) 2005 Electron energy spectra, fluxes and day-night asymmetry of B-8 solar neutrinos from measurements with NaCl dissolved in the heavy-water detector at the Sudbury Neutrino Observatory *Phys. Rev.* **C72** 055502 (arXiv:nucl-ex/0502021)
- [52] Kayser B 2008 Neutrino mass, mixing and flavor change (arXiv:0804.1497)
- [53] Arpesella C *et al* (Borexino Collaboration) 2008 Direct measurement of the Be-7 solar neutrino flux with 192 days of Borexino data *Phys. Rev. Lett.* **101** 091302 (arXiv:0805.3843)
- [54] Fukuda Y *et al* (Super-Kamiokande Collaboration) 1998 Evidence for oscillation of atmospheric neutrinos *Phys. Rev. Lett.* **81** 1562–1567 (arXiv:hep-ex/9807003)
- [55] Ashie Y *et al* (Super-Kamiokande Collaboration) 2004 Evidence for an oscillatory signature in atmospheric neutrino oscillation *Phys. Rev. Lett.* **93** 101801 (arXiv:hep-ex/0404034)
- [56] Abe K *et al* (Super-Kamiokande Collaboration) 2013 Evidence for the appearance of atmospheric tau neutrinos in Super-Kamiokande *Phys. Rev. Lett.* **110** 181802 (arXiv:1206.0328)
- [57] Ahn M H *et al* (K2K Collaboration) 2003 Indications of neutrino oscillation in a 250 km long baseline experiment *Phys. Rev. Lett.* **90** 041801 (arXiv:hep-ex/0212007)
- [58] Michael D *et al* (MINOS Collaboration) 2006 Observation of muon neutrino disappearance with the MINOS detectors and the NUMI neutrino beam *Phys. Rev. Lett.* **97** 191801

- (arXiv:hep-ex/0607088)
- [59] Abe K *et al* (T2K Collaboration) 2011 Indication of electron neutrino appearance from an accelerator-produced off-axis muon neutrino beam *Phys. Rev. Lett.* **107** 041801 (arXiv:1106.2822)
 - [60] Abe K *et al* (T2K Collaboration) 2013 Measurement of neutrino oscillation parameters from muon neutrino disappearance with an off-axis beam *Phys. Rev. Lett.* **111** 211803 (arXiv:1308.0465)
 - [61] Abe K *et al* (T2K Collaboration) 2014 Precise measurement of the neutrino mixing parameter θ_{23} from muon neutrino disappearance in an off-axis beam *Phys. Rev. Lett.* **112** 181801 (arXiv:1403.1532)
 - [62] Freund M 2001 Analytic approximations for three neutrino oscillation parameters and probabilities in matter *Phys. Rev.* **D64** 053003 (arXiv:hep-ph/0103300)
 - [63] Agafonova N *et al* (OPERA Collaboration) 2015 Discovery of tau neutrino appearance in the CNGS beam with the OPERA experiment *Phys. Rev. Lett.* **115** 121802 (arXiv:1507.01417)
 - [64] Eguchi K *et al* (KamLAND Collaboration) 2003 First results from KamLAND: evidence for reactor anti-neutrino disappearance *Phys. Rev. Lett.* **90** 021802 (arXiv:hep-ex/0212021)
 - [65] An F P *et al* (Daya Bay Collaboration) 2012 Observation of electron-antineutrino disappearance at Daya Bay *Phys. Rev. Lett.* **108** 171803 (arXiv:1203.1669)
 - [66] An F P *et al* (Daya Bay Collaboration) 2013 Improved measurement of electron antineutrino disappearance at Daya Bay *Chin. Phys.* **C37** 011001 (arXiv:1210.6327)
 - [67] Abe S *et al* (KamLAND Collaboration) 2008 Precision measurement of neutrino oscillation parameters with KamLAND *Phys. Rev. Lett.* **100** 221803 (arXiv:0801.4589)
 - [68] Ahn J K *et al* (RENO Collaboration) 2012 Observation of reactor electron antineutrino disappearance in the RENO experiment *Phys. Rev. Lett.* **108** 191802 (arXiv:1204.0626)
 - [69] Abe Y *et al* (Double Chooz Collaboration) 2012 Reactor electron antineutrino disappearance in the Double Chooz experiment *Phys. Rev.* **D86** 052008 (arXiv:1207.6632)
 - [70] An F P *et al* (Daya Bay Collaboration) 2014 Spectral measurement of electron antineutrino oscillation amplitude and frequency at Daya Bay *Phys. Rev. Lett.* **112** 061801 (arXiv:1310.6732)
 - [71] Wang Y F and Xing Z Z 2015 Neutrino masses and flavor oscillations (arXiv:1504.06155)
 - [72] Capozzi F, Fogli G L, Lisi E, Marrone A, Montanino D and Palazzo A 2014 Status of three-neutrino oscillation parameters: circa 2013 *Phys. Rev.* **D89** 093018 (arXiv:1312.2878)
 - [73] Forero D V, Tortola M and Valle J W F 2014 Neutrino oscillations refitted *Phys. Rev.* **D90** 093006 (arXiv:1405.7540)
 - [74] Gonzalez-Garcia M C, Maltoni M and Schwetz T 2014 Updated fit to three neutrino mixing: status of leptonic CP violation *JHEP* **1411** 052 (arXiv:1409.5439)
 - [75] Li Y F and Xing Z Z 2011 Possible capture of keV sterile neutrino dark matter on radioactive β -decaying nuclei *Phys. Lett.* **B695** 205–210 (arXiv:1009.5870)
 - [76] Xing Z Z 2013 On the neutrino mass ordering and flavor mixing structure (arXiv:1309.2102)
 - [77] Li Y F, Cao J, Wang Y and Zhan L 2013 Unambiguous determination of the neutrino mass hierarchy using reactor neutrinos *Phys. Rev.* **D88** 013008 (arXiv:1303.6733)
 - [78] Aartsen M G *et al* (IceCube-PINGU Collaboration) 2014 Letter of intent: the precision IceCube next generation upgrade (PINGU) (arXiv:1401.2046)
 - [79] Patterson R B (for the NO ν A Collaboration) 2013 The NO ν A experiment: status and outlook *Nucl. Phys. Proc. Suppl.* **235–236** 151–157 (arXiv:1209.0716)
 - [80] Adams C *et al* (LBNE Collaboration) 2013 The long-baseline neutrino experiment: exploring fundamental symmetries of the Universe (arXiv:1307.7335)
 - [81] Cabibbo N 1963 Unitary symmetry and leptonic decays *Phys. Rev. Lett.* **10** 531–533
 - [82] Kobayashi M and Maskawa T 1973 CP violation in the renormalizable theory of weak interaction *Prog. Theor. Phys.* **49** 652–657
 - [83] Xing Z Z and Zhou S 2008 Implications of leptonic unitarity violation at neutrino telescopes *Phys. Lett.* **B666** 166–172 (arXiv:0804.3512)
 - [84] Xing Z Z and Zhu J Y 2015 Leptonic unitarity triangles and effective mass triangles of the Majorana

- neutrinos (arXiv:1511.00450)
- [85] Xing Z Z and Zhou Y L 2013 Majorana CP-violating phases in neutrino-antineutrino oscillations and other lepton-number-violating processes *Phys. Rev.* **D88** 033002 (arXiv:1305.5718)
 - [86] Hall L J, Murayama H and Weiner N 2000 Neutrino mass anarchy *Phys. Rev. Lett.* **84** 2572–2575 (arXiv:hep-ph/9911341)
 - [87] Fukuyama T and Nishiura H 1997 Mass matrix of Majorana neutrinos (arXiv:hep-ph/9702253)
 - [88] Ma E and Raidal M 2001 Neutrino mass, muon anomalous magnetic moment and lepton flavor nonconservation *Phys. Rev. Lett.* **87** 011802 (arXiv:hep-ph/0102255)
 - [89] Lam C S 2001 A 2-3 symmetry in neutrino oscillations *Phys. Lett.* **B507** 214–218 (arXiv:hep-ph/0104116)
 - [90] Balaji K R S, Grimus W and Schwetz T 2001 The solar LMA neutrino oscillation solution in the Zee model *Phys. Lett.* **B508** 301–310 (arXiv:hep-ph/0104035)
 - [91] Apollonio M *et al* (CHOOZ Collaboration) 1998 Initial results from the CHOOZ long-baseline reactor neutrino oscillation experiment *Phys. Lett.* **B420** 397–404 (arXiv:hep-ex/9711002)
 - [92] Harrison P F, Perkins D H and Scott W G 2002 Tri-bimaximal mixing and the neutrino oscillation data *Phys. Lett.* **B530** 167–173 (arXiv:hep-ph/0202074)
 - [93] Xing Z Z 2002 Nearly tri-bimaximal neutrino mixing and CP violation *Phys. Lett.* **B533** 85–93 (arXiv:hep-ph/0204049)
 - [94] Vissani F 1997 A study of the scenario with nearly degenerate Majorana neutrinos (arXiv:hep-ph/9708483)
 - [95] Mohapatra R N and Nussinov S 1999 Bimaximal neutrino mixing and neutrino mass matrix *Phys. Rev.* **D60** 013002 (arXiv:hep-ph/9809415)
 - [96] Kajiyama Y, Raidal M and Strumia A 2007 The golden ratio prediction for the solar neutrino mixing *Phys. Rev.* **D76** 117301 (arXiv:0705.4559)
 - [97] Xing Z Z 2003 Wolfenstein-like parametrization of the neutrino mixing matrix *J. Phys.* **G29** 2227–2234 (arXiv:hep-ph/0211465)
 - [98] Giunti C 2003 Current status of neutrino masses and mixings *Nucl. Phys. Proc. Suppl.* **117** 24–28 (arXiv:hep-ph/0209103)
 - [99] Fritzsch H and Xing Z Z 2006 Lepton mass hierarchy and neutrino mixing *Phys. Lett.* **B634** 514–519 (arXiv:hep-ph/0601104)
 - [100] Singh N N, Devi H Z and Patgiri M 2007 Normal versus inverted hierarchical models within μ - τ symmetry (arXiv:0707.2713)
 - [101] Grimus W, Joshipura A S, Kaneko S, Lavoura L, Sawanaka H and Tanimoto M 2005 Non-vanishing U_{e3} and $\cos 2\theta_{23}$ from a broken Z_2 symmetry *Nucl. Phys.* **B713** 151–172 (arXiv:hep-ph/0408123)
 - [102] Kitabayashi T and Yasue M 2005 μ - τ symmetry and maximal CP violation *Phys. Lett.* **B621** 133–138 (arXiv:hep-ph/0504212)
 - [103] Grimus W, Kaneko S, Lavoura L, Sawanaka H and Tanimoto M 2006 μ - τ antisymmetry and neutrino mass matrices *JHEP* **0601** 110 (arXiv:hep-ph/0510326)
 - [104] Ge S F, He H J and Yin F R 2010 Common origin of soft mu-tau and CP breaking in neutrino seesaw and the origin of matter *JCAP* **1005** 017 (arXiv:1001.0940)
 - [105] He H J and Yin F R 2011 Common origin of μ - τ and CP breaking in neutrino seesaw, baryon asymmetry, and hidden flavor symmetry *Phys. Rev.* **D84** 033009 (arXiv:1104.2654)
 - [106] Harrison P H and Scott W G 2002 μ - τ reflection symmetry in lepton mixing and neutrino oscillations *Phys. Lett.* **B547** 219–228 (arXiv:hep-ph/0210197)
 - [107] Feruglio F, Hagedorn C and Ziegler R 2013 Lepton mixing parameters from discrete and CP symmetries *JHEP* **1307** 027 (arXiv:1211.5560)
 - [108] Holthausen M, Lindner M and Schmidt M A 2013 CP and discrete flavour symmetries *JHEP* **1304** 122 (arXiv:1211.6953)
 - [109] Babu K S, Ma E and Valle J W F 2003 Underlying A_4 symmetry for the neutrino mass matrix and the quark mixing matrix *Phys. Lett.* **B552** 207–213 (arXiv:hep-ph/0206292)

- [110] Ma E 2002 Plato’s fire and the neutrino mass matrix *Mod. Phys. Lett.* **A17** 2361–2370 (arXiv:hep-ph/0211393)
- [111] Chankowski P H, Ioannisian A, Pokorski S and Valle J W F 2001 Neutrino unification *Phys. Rev. Lett.* **86** 3488–3491 (arXiv:hep-ph/0011150)
- [112] Chun E and Pokorski S 2000 Slepton flavor mixing and neutrino masses *Phys. Rev.* **D62** 053001 (arXiv:hep-ph/9912210)
- [113] Grimus W and Lavoura L 2004 A nonstandard CP transformation leading to maximal atmospheric neutrino mixing *Phys. Lett.* **B579** 113–122 (arXiv:hep-ph/0305309)
- [114] Zhou Y L 2014 μ - τ reflection symmetry and radiative corrections (arXiv:1409.8600)
- [115] Aizawa I and Yasue M 2005 General property of neutrino mass matrix and CP violation *Phys. Lett.* **B607** 267–275 (arXiv:hep-ph/0409331)
- [116] Aizawa I, Kitabayashi T and Yasue M 2005 Neutrino mass textures with maximal CP violation *Phys. Rev.* **D72** 055014 (arXiv:hep-ph/0504172)
- [117] Aizawa I, Kitabayashi T and Yasue M 2005 Determination of neutrino mass texture for maximal CP violation *Nucl. Phys.* **B728** 220–232 (arXiv:hep-ph/0507332)
- [118] Kitabayashi T and Yasue M 2015 Maximal CP violation in flavor neutrino masses (arXiv:1509.06071)
- [119] Gupta S, Joshipura A S and Patel K M 2013 How good is μ - τ symmetry after results on non-zero θ_{13} ? *JHEP* **1309** 035 (arXiv:1301.7130)
- [120] Abbas M and Smirnov A Yu 2010 Is the tribimaximal mixing accidental? *Phys. Rev.* **D82** 013008 (arXiv:1004.0099)
- [121] Ade P A R *et al* (Planck Collaboration) 2015 Planck 2015 results. XIII. Cosmological parameters (arXiv:1502.01589)
- [122] Mohapatra R N 2004 θ_{13} as a probe of μ - τ symmetry for leptons *JHEP* **0410** 027 (arXiv:hep-ph/0408187)
- [123] Petcov S T 1982 On pseudo-Dirac neutrinos, neutrino oscillations and neutrinoless double β -decay *Phys. Lett.* **B110** 245–249
- [124] Mohapatra R N and Rodejohann W 2005 Broken μ - τ symmetry and leptonic CP violation *Phys. Rev.* **D72** 053001 (arXiv:hep-ph/0507312)
- [125] King S F 2002 Constructing the large mixing angle MNS matrix in seesaw models with right-handed neutrino dominance *JHEP* **0209** 011 (arXiv:hep-ph/0204360)
- [126] Ohlsson T and Zhou S 2014 Renormalization group running of neutrino parameters *Nature Commun.* **5** 5153 (arXiv:1311.3846)
- [127] Chankowski P H and Pluciennik Z 1993 Renormalization group equations for seesaw neutrino masses *Phys. Lett.* **B316** 312–317 (arXiv:hep-ph/9306333)
- [128] Babu K S, Leung C N and Pantaleone J 1993 Renormalization of the neutrino mass operator *Phys. Lett.* **B319** 191–198 (arXiv:hep-ph/9309223)
- [129] Antusch S, Drees M, Kersten J, Lindner M and Ratz M 2001 Neutrino mass operator renormalization revisited *Phys. Lett.* **B519** 238–242 (arXiv:hep-ph/0108005)
- [130] Antusch S, Drees M, Kersten J, Lindner M and Ratz M 2002 Neutrino mass operator renormalization in two Higgs doublet models and the MSSM *Phys. Lett.* **B525** 130–134 (arXiv:hep-ph/0110366)
- [131] Babu K S 1987 Renormalization-group analysis of the Kobayashi-Maskawa matrix *Z. Phys.* **C35** 69–75
- [132] Casas J A, Espinosa J R, Ibarra A and Navarro I 2000 General RG equations for physical neutrino parameters and their phenomenological implications *Nucl. Phys.* **B573** 652–684 (arXiv:hep-ph/9910420)
- [133] Antusch S, Kersten J, Lindner M and Ratz M 2003 Running neutrino masses, mixings and CP phases: analytical results and phenomenological consequences *Nucl. Phys.* **B674** 401–433 (arXiv:hep-ph/0305273)
- [134] Ellis J R and Lola S 1999 Can neutrinos be degenerate in mass? *Phys. Lett.* **B458** 310–321

- (arXiv:hep-ph/9904279)
- [135] Chankowski P H, Krolkowski W and Pokorski S 2000 Fixed points in the evolution of neutrino mixings *Phys. Lett.* **B473** 109–117 (arXiv:hep-ph/9910231)
 - [136] Dighe A, Goswami S and Roy P 2007 Radiatively broken symmetries of nonhierarchical neutrinos *Phys. Rev.* **D76** 096005 (arXiv:0704.3735)
 - [137] Haba N, Matsui Y and Okamura N 2000 The effects of Majorana phases in three-generation neutrinos *Eur. Phys. J.* **C17** 513–520 (arXiv:hep-ph/0005075)
 - [138] Balaji K R S, Dighe A S, Mohapatra R N and Parida M K 2000 Generation of large flavor mixing from radiative corrections *Phys. Rev. Lett.* **84** 5034–5037 (arXiv:hep-ph/0001310)
 - [139] Miura T, Shindou T and Takasugi E 2002 Exploring the neutrino mass matrix at M_R scale *Phys. Rev.* **D66** 093002 (arXiv:hep-ph/0206207)
 - [140] Goswami S, Petcov S T, Ray S and Rodejohann W 2009 Large U_{e3} and tribimaximal mixing *Phys. Rev.* **D80** 053013 (arXiv:0907.2869)
 - [141] Martin S P 2010 A supersymmetry primer *Adv. Ser. Direct. High Energy Phys.* **21** 1–153 (arXiv:hep-ph/9709356)
 - [142] Luo S and Xing Z Z 2014 Resolving the octant of θ_{23} via radiative mu-tau symmetry breaking *Phys. Rev.* **D90** 073005 (arXiv:1408.5005)
 - [143] Luo S, Mei J W and Xing Z Z 2005 Radiative generation of leptonic CP violation *Phys. Rev.* **D72** 053014 (arXiv:hep-ph/0507065)
 - [144] Dighe A, Goswami S and Ray S 2009 Renormalization group evolution of neutrino mixing parameters near $\theta_{13} = 0$ and models with vanishing θ_{13} at the high scale *Phys. Rev.* **D79** 076006 (arXiv:0810.5680)
 - [145] Xing Z Z, Zhang H and Zhou S 2012 Impacts of the Higgs mass on vacuum stability, running fermion masses and two-body Higgs decays *Phys. Rev.* **D86** 013013 (arXiv:1112.3112)
 - [146] Giunti C and Tanimoto M 2002 CP violation in bilarge lepton mixing *Phys. Rev.* **D66** 113006 (arXiv:hep-ph/0209169)
 - [147] Romanino A 2004 Charged lepton contributions to the solar neutrino mixing and θ_{13} *Phys. Rev.* **D70** 013003 (arXiv:hep-ph/0402258)
 - [148] Petcov S T and Rodejohann W 2005 Flavor symmetry $L_e - L_\mu - L_\tau$, atmospheric neutrino mixing and CP violation in the lepton sector *Phys. Rev.* **D71** 073002 (arXiv:hep-ph/0409135)
 - [149] King S F 2005 Predicting neutrino parameters from SO(3) family symmetry and quark-lepton unification *JHEP* **0508** 105 (arXiv:hep-ph/0506297)
 - [150] Hochmuth K A, Petcov S T and Rodejohann W 2007 $U_{\text{PMNS}} = U_l^\dagger U_\nu$ *Phys. Lett.* **B654** 177–188 (arXiv:0706.2975)
 - [151] Chao W and Zheng Y J 2013 Relatively large θ_{13} from modification to the tribimaximal, bimaximal and democratic neutrino mixing matrices *JHEP* **1302** 044 (arXiv:1107.0738)
 - [152] Duarah C, Das A and Singh N N 2012 Dependence of $\tan^2 \theta_{12}$ on Dirac CP phase δ in tribimaximal neutrino mixing under charged lepton correction (arXiv:1210.8265)
 - [153] Marzocca D, Petcov S T, Romanino A and Sevilla M C 2013 Nonzero $|U_{e3}|$ from charged lepton corrections and the atmospheric neutrino mixing angle *JHEP* **1305** 073 (arXiv:1302.0423)
 - [154] Gollu S, Deepthi K N and Mohanta R 2013 Charged lepton correction to tribimaximal lepton mixing and its implications to neutrino phenomenology *Mod. Phys. Lett.* **A28** 1350131 (arXiv:1303.3393)
 - [155] Hall L J and Ross G G 2013 Discrete symmetries and neutrino mass perturbations for θ_{13} *JHEP* **1311** 091 (arXiv:1303.6962)
 - [156] Garg S K and Gupta S 2013 Corrections for tribimaximal, bimaximal and democratic neutrino mixing matrices *JHEP* **1310** 128 (arXiv:1308.3054)
 - [157] Liu Z and Wu Y L 2014 Leptonic CP violation and Wolfenstein parametrization for lepton mixing *Phys. Lett.* **B733** 226–232 (arXiv:1403.2440)
 - [158] Shimizu Y, Tanimoto M and Yamamoto K 2015 Predicting CP violation in deviation from tribimaximal mixing of neutrinos *Mod. Phys. Lett.* **A30** 1550002 (arXiv:1405.1521)

- [159] Petcov S T 2015 Predicting the values of the leptonic CP violation phases *Nucl. Phys.* **B892** 400–428 (arXiv:1405.6006)
- [160] Girardi I, Petcov S T and Titov A V 2015 Determining the Dirac CP violation phase in the neutrino mixing matrix from sum rules *Nucl. Phys.* **B894** 733–768 (arXiv:1410.8056)
- [161] Girardi I, Petcov S T and Titov A V 2015 Predictions for the leptonic Dirac CP violation phase: a systematic phenomenological analysis *Eur. Phys. J.* **C75** 345 (arXiv:1504.00658)
- [162] Antusch S and King S F 2005 Charged lepton corrections to neutrino mixing angles and CP phases revisited *Phys. Lett.* **B631** 42–47 (arXiv:hep-ph/0508044)
- [163] Frampton P H, Petcov S T and Rodejohann W 2004 On deviations from bimaximal neutrino mixing *Nucl. Phys.* **B687** 31–54 (arXiv:hep-ph/0401206)
- [164] Gatto R, Sartori G and Tonin M 1968 Weak selfmasses, Cabibbo angle and broken $SU(2) \times SU(2)$ *Phys. Lett.* **B28** 128–130
- [165] Kile J, Perez M J, Ramond P and Zhang J 2014 θ_{13} and the flavor ring *Phys. Rev.* **D90** 013004 (arXiv:1403.6136)
- [166] Georgi H and Jarlskog C 1979 A new lepton-quark mass relation in a unified theory *Phys. Lett.* **B86** 297–300
- [167] Antusch S, Gross C, Maurer V and Sluka C 2013 $\theta_{13}^{\text{PMNS}} = \theta_c/\sqrt{2}$ from GUTs *Nucl. Phys.* **B866** 255–269 (arXiv:1205.1051)
- [168] King S F 2012 Tribimaximal-Cabibbo mixing *Phys. Lett.* **B718** 136–142 (arXiv:1205.0506)
- [169] Antusch S and Spinrath M 2009 New GUT predictions for quark and lepton mass ratios confronted with phenomenology *Phys. Rev.* **D79** 095004 (arXiv:0902.4644)
- [170] Weinberg S 1977 The problem of mass *Trans. New York Acad. Sci.* **38** 185–201
- [171] Fritzsch H 1977 Calculating the Cabibbo angle *Phys. Lett.* **B70** 436–440
- [172] Antusch S and Maurer V 2011 Large neutrino mixing angle $\theta_{13}^{\text{PMNS}}$ and quark-lepton mass ratios in unified flavour models *Phys. Rev.* **D84** 117301 (arXiv:1107.3728)
- [173] Marzocca D, Petcov S T, Romanino A and Spinrath M 2011 Sizeable θ_{13} from the charged lepton sector in $SU(5)$, (tri)bimaximal neutrino mixing and Dirac CP violation *JHEP* **1111** 009 (arXiv:1108.0614)
- [174] Antusch S, Gross C, Maurer V and Sluka C 2013 A flavour GUT model with $\theta_{13}^{\text{PMNS}} = \theta_c/\sqrt{2}$ *Nucl. Phys.* **B877** 772–791 (arXiv:1305.6612)
- [175] Lam C S 2006 Mass independent textures and symmetry *Phys. Rev.* **D74** 113004 (arXiv:hep-ph/0611017)
- [176] Lam C S 2007 Symmetry of lepton mixing *Phys. Lett.* **B656** 193–198 (arXiv:0708.3665)
- [177] Lam C S 2008 Determining horizontal symmetry from neutrino mixing *Phys. Rev. Lett.* **101** 121602 (arXiv:0804.2622)
- [178] He X G and Zee A 2007 Minimal modification to the tribimaximal neutrino mixing *Phys. Lett.* **B645** 427–431 (arXiv:hep-ph/0607163)
- [179] Xing Z Z and Zhou S 2007 Tribimaximal neutrino mixing and flavor-dependent resonant leptogenesis *Phys. Lett.* **B653** 278–287 (arXiv:hep-ph/0607302)
- [180] Albright C H and Rodejohann W 2009 Comparing trimaximal mixing and its variants with deviations from tribimaximal mixing *Eur. Phys. J.* **C62** 599–608 (arXiv:0812.0436)
- [181] Albright C H, Dueck A and Rodejohann W 2010 Possible alternatives to tribimaximal mixing *Eur. Phys. J.* **C70** 1099–1110 (arXiv:1004.2798)
- [182] He X G and Zee A 2011 Minimal modification to tribimaximal mixing *Phys. Rev.* **D84** 053004 (arXiv:1106.4359)
- [183] Rodejohann W and Zhang H 2012 Simple two parameter description of lepton mixing *Phys. Rev.* **D86** 093008 (arXiv:1207.1225)
- [184] Talbert J 2014 [Re]constructing finite flavour groups: horizontal symmetry scans from the bottom-up *JHEP* **12** 058 (arXiv:1409.7310)
- [185] Lam C S 2008 The unique horizontal symmetry of leptons *Phys. Rev.* **D78** 073015 (arXiv:0809.1185)

- [186] Ma E and Rajasekaran G 2001 Softly broken A_4 symmetry for nearly degenerate neutrino masses *Phys. Rev.* **D64** 113012 (arXiv:hep-ph/0106291)
- [187] Altarelli G and Feruglio F 2005 Tribimaximal neutrino mixing from discrete symmetry in extra dimensions *Nucl. Phys.* **B720** 64–88 (arXiv:hep-ph/0504165)
- [188] Lam C S 2011 Group theory and dynamics of neutrino mixing *Phys. Rev.* **D83** 113002 (arXiv:1104.0055)
- [189] Ishimori H, Kobayashi T, Ohki H, Okada H, Shimizu Y and Tanimoto M 2010 Non-Abelian discrete symmetries in particle physics *Prog. Theor. Phys. Suppl.* **183** 1–163 (arXiv:1003.3552)
- [190] Toorop R A, Feruglio F and Hagedorn C 2011 Discrete flavour symmetries in light of T2K *Phys. Lett.* **B703** 447–451 (arXiv:1107.3486)
- [191] Toorop R A, Feruglio F and Hagedorn C 2012 Finite modular groups and lepton mixing *Nucl. Phys.* **B858** 437–467 (arXiv:1112.1340)
- [192] Lam C S 2013 Finite symmetry of leptonic mass matrices *Phys. Rev.* **D87** 013001 (arXiv:1208.5527)
- [193] Holthausen M, Lim K S and Lindner M 2013 Lepton mixing patterns from a scan of finite discrete groups *Phys. Lett.* **B721** 61–67 (arXiv:1212.2411)
- [194] Ge S F, Dicus D A and Repko W W 2011 Z_2 symmetry prediction for the leptonic Dirac CP phase *Phys. Lett.* **B702** 220–223 (arXiv:1104.0602)
- [195] Ge S F, Dicus D A and Repko W W 2012 Residual symmetries for neutrino mixing with a large θ_{13} and nearly maximal δ_D *Phys. Rev. Lett.* **108** 041801 (arXiv:1108.0964)
- [196] Hernandez D and Smirnov A Yu 2012 Lepton mixing and discrete symmetries *Phys. Rev.* **D86** 053014 (arXiv:1204.0445)
- [197] Hanlon A D, Ge S F and Repko W W 2013 Phenomenological study of residual Z_2^s and \bar{Z}_2^s symmetries *Phys. Lett.* **B729** 185–191 (arXiv:1308.6522)
- [198] Froggatt C D and Nielsen H B 1979 Hierarchy of quark masses, Cabibbo angles and CP violation *Nucl. Phys.* **B147** 277–298
- [199] Altarelli G and Feruglio F 2006 Tribimaximal neutrino mixing, A_4 and the modular symmetry *Nucl. Phys.* **B741** 215–235 (arXiv:hep-ph/0512103)
- [200] King S F and Luhn C 2011 Trimaximal neutrino mixing from vacuum alignment in A_4 and S_4 models *JHEP* **1109** 042 (arXiv:1107.5332)
- [201] King S F and Luhn C 2009 On the origin of neutrino flavour symmetry *JHEP* **0910** 093 (arXiv:0908.1897)
- [202] Chen M C and King S F 2009 A_4 seesaw models and form dominance *JHEP* **0906** 072 (arXiv:0903.0125)
- [203] Zhou Y L 2015 Neutrino mixing and flavor symmetries *PhD Thesis* (Institute of High Energy Physics, Beijing)
- [204] Gronau M, Kfir A and Loewy R 1986 Basis independent tests of CP violation in fermion mass matrices *Phys. Rev. Lett.* **56** 1538
- [205] Branco G C, Lavoura L and Rebelo M N 1986 Majorana neutrinos and CP violation in the leptonic sector *Phys. Lett.* **B180** 264–268
- [206] Ding G J, King S F, Luhn C and Stuart A J 2013 Spontaneous CP violation from vacuum alignment in S_4 models of leptons *JHEP* **1305** 084 (arXiv:1303.6180)
- [207] Feruglio F, Hagedorn C and Ziegler R 2014 A realistic pattern of lepton mixing and masses from S_4 and CP *Eur. Phys. J.* **C74** 2753 (arXiv:1303.7178)
- [208] Ding G J, King S F and Stuart A J 2013 Generalised CP and A_4 family symmetry *JHEP* **1312** 006 (arXiv:1307.4212)
- [209] Li C C and Ding G J 2014 Generalised CP and trimaximal TM1 lepton mixing in S_4 family symmetry *Nucl. Phys.* **B881** 206–232 (arXiv:1312.4401)
- [210] Ding G J and Zhou Y L 2015 Predicting lepton flavor mixing from $\Delta(48)$ and generalized CP symmetries *Chin. Phys.* **C39** 021001 (arXiv:1312.5222)
- [211] Ding G J and Zhou Y L 2014 Lepton mixing parameters from $\Delta(48)$ family symmetry and

- generalised CP *JHEP* **1406** 023 (arXiv:1404.0592)
- [212] Branco G C, Felipe R G, Joaquim F R and Serodio H 2012 Spontaneous leptonic CP violation and nonzero θ_{13} *Phys. Rev.* **D86** 076008 (arXiv:1203.2646)
- [213] Ahn Y H, Kang S K and Kim C S 2013 Spontaneous CP violation in A_4 flavor symmetry and leptogenesis *Phys. Rev.* **D87** 113012 (arXiv:1304.0921)
- [214] Zhao Z H 2014 Minimal modifications to the tribimaximal neutrino mixing *JHEP* **11** 143 (arXiv:1405.3022)
- [215] Karmakar B and Sil A 2016 Spontaneous CP violation in lepton sector: a common origin for θ_{13} , Dirac CP phase and leptogenesis *Phys. Rev.* **D93** 013006 (arXiv:1509.07090)
- [216] Antusch S, King S F, Luhn C and Spinrath M 2011 Right unitarity triangles and tribimaximal mixing from discrete symmetries and unification *Nucl. Phys.* **B850** 477–504 (arXiv:1103.5930)
- [217] Antusch S, King S F and Spinrath M 2013 Spontaneous CP violation in $A_4 \times \text{SU}(5)$ with constrained sequential dominance 2 *Phys. Rev.* **D87** 096018 (arXiv:1301.6764)
- [218] King S F 2013 Minimal predictive seesaw model with normal neutrino mass hierarchy *JHEP* **1307** 137 (arXiv:1304.6264)
- [219] King S F 2013 Minimal seesaw model predicting best fit lepton mixing angles *Phys. Lett.* **B724** 92–98 (arXiv:1305.4846)
- [220] Altarelli G, Feruglio F and Masina I 2004 Can neutrino mixings arise from the charged lepton sector? *Nucl. Phys.* **B689** 157–171 (arXiv:hep-ph/0402155)
- [221] Lam C S 2005 Neutrino 2-3 symmetry and inverted hierarchy *Phys. Rev.* **D71** 093001 (arXiv:hep-ph/0503159)
- [222] Grimus W and Lavoura L 2001 Softly broken lepton numbers and maximal neutrino mixing *JHEP* **0107** 045 (arXiv:hep-ph/0105212)
- [223] Grimus W and Lavoura L 2003 A discrete symmetry group for maximal atmospheric neutrino mixing *Phys. Lett.* **B572** 189–195 (arXiv:hep-ph/0305046)
- [224] Grimus W and Lavoura L 2013 μ - τ interchange symmetry and lepton mixing *Fortsch. Phys.* **61** 535–545 (arXiv:1207.1678)
- [225] Grimus W and Lavoura L 2003 Maximal atmospheric neutrino mixing in an $\text{SU}(5)$ model *Eur. Phys. J.* **C28** 123–131 (arXiv:hep-ph/0211334)
- [226] Grimus W and Lavoura L 2002 Soft lepton-flavor violation in a multi-Higgs-doublet seesaw model *Phys. Rev.* **D66** 014016 (arXiv:hep-ph/0204070)
- [227] Toorop R A, Bazzocchi F, Merlo L and Paris A 2011 Constraining flavour symmetries at the EW scale I: the A_4 Higgs potential *JHEP* **1103** 035 (arXiv:1012.1791)
- [228] Toorop R A, Bazzocchi F, Merlo L and Paris A 2011 Constraining flavour symmetries at the EW scale II: the fermion processes *JHEP* **1103** 040 (arXiv:1012.2091)
- [229] Keus V, King S F and Moretti S 2010 Three-Higgs-doublet models: symmetries, potentials and Higgs boson masses *JHEP* **1401** 052 (arXiv:1310.8253)
- [230] Grimus W and Lavoura L 2004 Maximal atmospheric neutrino mixing and the small ratio of muon to tau mass *J. Phys.* **G30** 73–82 (arXiv:hep-ph/0309050)
- [231] Chen P, Ding G J, Gonzalez-Canales F and Valle J W F 2016 Generalized μ - τ reflection symmetry and leptonic CP violation *Phys. Lett.* **B753** 644–652 (arXiv:1512.01551)
- [232] Mohapatra R N and Nishi C C 2015 Implications of μ - τ flavored CP symmetry of leptons *JHEP* **1508** 092 (arXiv:1506.06788)
- [233] Chang D, Mohapatra R N and Parida M K 1984 Decoupling parity and $\text{SU}(2)_R$ breaking scales: a new approach to left-right symmetric models *Phys. Rev. Lett.* **52** 1072
- [234] Joshipura A S and Patel K M 2015 Generalized μ - τ symmetry and discrete subgroups of $O(3)$ *Phys. Lett.* **B749** 159–166 (arXiv:1507.01235)
- [235] He H J, Rodejohann W and Xu X J 2015 Origin of constrained maximal CP violation in flavor symmetry *Phys. Lett.* **B751** 586–594 (arXiv:1507.03541)
- [236] Mohapatra R N and Nishi C C 2012 S_4 flavored CP symmetry for neutrinos *Phys. Rev.* **D86** 073007 (arXiv:1208.2875)

- [237] Ferreira P M, Grimus W, Lavoura L and Ludl P O 2012 Maximal CP violation in lepton mixing from a model with $\Delta(27)$ flavour symmetry *JHEP* **1209** 128 (arXiv:1206.7072)
- [238] Ma E, Natale A and Popov O 2015 Neutrino mixing and CP phase correlations *Phys. Lett.* **B746** 114–116 (arXiv:1502.08023)
- [239] Ma E 2015 Transformative A_4 mixing of neutrinos with CP violation *Phys. Rev.* **D92** 051301 (arXiv:1504.02086)
- [240] Xing Z Z 2012 Implications of the Daya Bay observation of θ_{13} on the leptonic flavor mixing structure and CP violation *Chin. Phys.* **C36** 281–297 (arXiv:1203.1672)
- [241] Grimus W and Lavoura L 2008 A model for trimaximal lepton mixing *JHEP* **0809** 106 (arXiv:0809.0226)
- [242] Gupta S, Joshipura A S and Patel K M 2012 Minimal extension of tribimaximal mixing and generalized $Z_2 \times Z_2$ symmetries *Phys. Rev.* **D85** 031903(R) (arXiv:1112.6113)
- [243] Varzielas I M and Lavoura L 2013 Flavour models for TM1 lepton mixing *J. Phys.* **G40** 085002 (arXiv:1212.3247)
- [244] Grimus W 2013 Discrete symmetries, roots of unity, and lepton mixing *J. Phys.* **G40** 075008 (arXiv:1301.0495)
- [245] King S F, Neder T and Stuart A J 2013 Lepton mixing predictions from $\Delta(6n^2)$ family symmetry *Phys. Lett.* **B726** 312–315 (arXiv:1305.3200)
- [246] Luhn C 2013 Trimaximal TM1 neutrino mixing in S_4 with spontaneous CP violation *Nucl. Phys.* **B875** 80–100 (arXiv:1306.2358)
- [247] Li C C and Ding G J 2014 Generalised CP and trimaximal TM1 lepton mixing in S_4 family symmetry *Nucl. Phys.* **B881** 206–232 (arXiv:1312.4401)
- [248] Friedberg R and Lee T D 2006 A possible relation between the neutrino mass matrix and the neutrino mapping matrix *High Energy Phys. Nucl. Phys.* **30** 591–598 (arXiv:hep-ph/0606071)
- [249] Friedberg R and Lee T D 2008 Hidden symmetry of the CKM and neutrino mapping matrices *Annals Phys.* **323** 1087–1105 (arXiv:0705.4156)
- [250] Friedberg R and Lee T D 2008 Jarlskog invariant of the neutrino mapping matrix *Annals Phys.* **323** 1677–1691 (arXiv:0709.1526)
- [251] Lee T D 2008 Symmetry and asymmetry *Nucl. Phys.* **A805** 54–71
- [252] Xing Z Z, Zhang H and Zhou S 2006 Nearly tribimaximal neutrino mixing and CP violation from mu-tau symmetry breaking *Phys. Lett.* **B641** 189–197 (arXiv:hep-ph/0607091)
- [253] Luo S and Xing Z Z 2007 Friedberg-Lee symmetry breaking and its prediction for θ_{13} *Phys. Lett.* **B646** 242–247 (arXiv:hep-ph/0611360)
- [254] Jarlskog C 2008 Neutrino sector with Majorana mass terms and Friedberg-Lee symmetry *Phys. Rev.* **D77** 073002 (arXiv:0712.0903)
- [255] Araki T and Geng C Q 2009 Leptogenesis in a model with Friedberg-Lee symmetry *Phys. Lett.* **B680** 343–350 (arXiv:0906.1903)
- [256] He X G and Liao W 2009 The Friedberg-Lee symmetry and minimal seesaw model *Phys. Lett.* **B681** 253–256 (arXiv:0909.1463)
- [257] Zhao Z H 2015 A modified Friedberg-Lee symmetry for the TM1 neutrino mixing *Phys. Rev.* **D92** 113001 (arXiv:1509.06915)
- [258] Antusch S, King S F, Luhn C and Spinrath M 2012 Trimaximal mixing with predicted θ_{13} from a new type of constrained sequential dominance *Nucl. Phys.* **B856** 328–341 (arXiv:1108.4278)
- [259] Karmakar B and Sil A 2015 Nonzero θ_{13} and leptogenesis in a type-I seesaw model with A_4 symmetry *Phys. Rev.* **D91** 013004 (arXiv:1407.5826)
- [260] Aguilar A *et al* (LSND Collaboration) 2001 Evidence for neutrino oscillations from the observation of electron anti-neutrinos in a muon anti-neutrino beam *Phys. Rev.* **D64** 112007 (arXiv:hep-ex/0104049)
- [261] Mention G, Fechner M, Lasserre T, Mueller T A, Lhuillier D, Cribier M and Letourneau A 2011 The reactor antineutrino anomaly *Phys. Rev.* **D83** 073006 (arXiv:1101.2755)
- [262] Abdurashitov J N *et al* (SAGE Collaboration) 2006 Measurement of the response of a

- Ga solar neutrino experiment to neutrinos from an ^{37}Ar source *Phys. Rev.* **C73** 045805 (arXiv:nuc1-ex/0512041)
- [263] Aguilar-Arevalo A A *et al* (MiniBooNE Collaboration) 2010 Event excess in the MiniBooNE search for $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$ oscillations *Phys. Rev. Lett.* **105** 181801 (arXiv:1007.1150)
- [264] Kusenko A 2009 Sterile neutrinos: the dark side of the light fermions *Phys. Rept.* **481** 1–28 (arXiv:0906.2968)
- [265] Merle A 2013 KeV neutrino model building *Int. J. Mod. Phys.* **D22** 1330020 (arXiv:1302.2625)
- [266] Adhikari R *et al.* 2016 A white paper on keV sterile neutrino dark matter (arXiv:1602.04816)
- [267] Goh H S, Mohapatra R N and Ng S P 2002 Testing neutrino mass matrices with approximate $L_e - L_\mu - L_\tau$ symmetry *Phys. Lett.* **B542** 116–122 (arXiv:hep-ph/0205131)
- [268] Gomez-Izquierdo J C and Perez-Lorenzana A 2006 Fixing the solar neutrino parameters with sterile neutrinos *Phys. Rev.* **D74** 013005 (arXiv:hep-ph/0601223)
- [269] Barry J, Rodejohann W and Zhang H 2012 Sterile neutrinos for warm dark matter and the reactor anomaly in flavor symmetry models *JCAP* **1201** 052 (arXiv:1110.6382)
- [270] Merle A, Morisi S and Winter W 2014 Common origin of reactor and sterile neutrino mixing *JHEP* **07** 039 (arXiv:1402.6332)
- [271] Ke H W, Zhou J H, Chen S, Liu T and Li X Q 2015 The hidden symmetries in the PMNS matrix and the light sterile neutrino(s) *Mod. Phys. Lett.* **A30** 1550136 (arXiv:1502.00875)
- [272] Rivera-Agudelo D C and Perez-Lorenzana A 2015 Generating θ_{13} from sterile neutrinos in μ - τ symmetric models (arXiv:1507.07030)
- [273] Gariazzo S, Giunti C, Laveder M, Li Y F and Zavanin E M 2016 Light sterile neutrinos *J. Phys.* **G43** 033001 (arXiv:1507.08204)
- [274] Giunti C and Kim C W 2007 Fundamentals of neutrino physics and astrophysics (Oxford, UK: Oxford University Press)
- [275] Xing Z Z 2012 A full parametrization of the 6×6 flavor mixing matrix in the presence of three light or heavy sterile neutrinos *Phys. Rev.* **D85** 013008 (arXiv:1110.0083)
- [276] Xing Z Z and Zhou Y L 2010 A generic diagonalization of the 3×3 neutrino mass matrix and its implications on the μ - τ flavor symmetry and maximal CP violation *Phys. Lett.* **B693** 584–590 (arXiv:1008.4906)
- [277] Toshev S 1991 On T violation in matter neutrino oscillations *Mod. Phys. Lett.* **A6** 455–460
- [278] Zhou Y L 2011 The Kobayashi-Maskawa parametrization of lepton flavor mixing and its application to neutrino oscillations in matter *Phys. Rev.* **D84** 113012 (arXiv:1110.5023)
- [279] Aartsen M G *et al* (IceCube Collaboration) 2013 Evidence for high-energy extraterrestrial neutrinos at the IceCube detector *Science* **342** 1242856 (arXiv:1311.5238)
- [280] Aartsen M G *et al* (IceCube Collaboration) 2014 Observation of high-energy astrophysical neutrinos in three years of IceCube data *Phys. Rev. Lett.* **113** 101101 (arXiv:1405.5303)
- [281] Learned J G and Pakvasa S 1995 Detecting tau-neutrino oscillations at PeV energies *Astropart. Phys.* **3** 267–274 (arXiv:hep-ph/9405296)
- [282] Xing Z Z 2012 A further study of μ - τ symmetry breaking at neutrino telescopes after the Daya Bay and RENO measurements of θ_{13} *Phys. Lett.* **B716** 220–224 (arXiv:1205.6532)
- [283] Xing Z Z 2006 Neutrino telescopes as a probe of broken μ - τ symmetry *Phys. Rev.* **D74** 013009 (arXiv:hep-ph/0605219)
- [284] Pakvasa S, Rodejohann W and Weiler T J 2008 Flavor ratios of astrophysical neutrinos: implications for precision measurements *JHEP* **0802** 005 (arXiv:0711.4517)
- [285] Glashow S L 1960 Resonant scattering of antineutrinos *Phys. Rev.* **118** 316–317
- [286] Gandhi R, Quigg C, Reno M H and Sarcevic I 1996 Ultrahigh-energy neutrino interactions *Astropart. Phys.* **5** 81–110 (arXiv:hep-ph/9512364)
- [287] Anchordoqui L A, Goldberg H, Halzen F and Weiler T J 2005 Neutrinos as a diagnostic of high energy astrophysical processes *Phys. Lett.* **B621** 18–21 (arXiv:hep-ph/0410003)
- [288] Maltoni M and Winter W 2008 Testing neutrino oscillations plus decay with neutrino telescopes *JHEP* **0807** 064 (arXiv:0803.2050)

- [289] Xing Z Z and Zhou S 2011 The Glashow resonance as a discriminator of UHE cosmic neutrinos originating from $p\gamma$ and pp collisions *Phys. Rev.* **D84** 033006 (arXiv:1105.4114)
- [290] Bhattacharya A, Gandhi R, Rodejohann W and Watanabe A 2011 The Glashow resonance at IceCube: signatures, event rates and pp vs $p\gamma$ interactions *JCAP* **1110** 017 (arXiv:1108.3163)
- [291] Lipari P, Lusignoli M and Meloni D 2007 Flavor composition and energy spectrum of astrophysical neutrinos *Phys. Rev.* **D75** 123005 (arXiv:0704.0718)
- [292] Beacom J F, Bell N F, Hooper D, Pakvasa S and Weiler T J 2003 Measuring flavor ratios of high-energy astrophysical neutrinos *Phys. Rev.* **D68** 093005 (arXiv:hep-ph/0307025)
- [293] Winter W 2012 Neutrinos from cosmic accelerators including magnetic field and flavor effects *Adv. High Energy Phys.* **2012** 586413 (arXiv:1201.5462)
- [294] Xing Z Z and Zhou S 2006 Towards determination of the initial flavor composition of ultrahigh-energy neutrino fluxes with neutrino telescopes *Phys. Rev.* **D74** 013010 (arXiv:astro-ph/0603781)
- [295] Fukugida M and Yanagida T 1986 Baryogenesis without grand unification *Phys. Lett.* **B174** 45–47
- [296] Buchmüller W, Peccei R D and Yanagida T 2005 Leptogenesis as the origin of matter *Ann. Rev. Nucl. Part. Sci.* **55** 311–355 (arXiv:hep-ph/0502169)
- [297] Davidson S, Nardi E and Nir Y 2008 Leptogenesis *Phys. Rept.* **466** 105–177 (arXiv:0802.2962)
- [298] Casas J A and Ibarra A 2001 Oscillating neutrinos and $\mu \rightarrow e, \gamma$ *Nucl. Phys.* **B618** 171–204 (arXiv:hep-ph/0103065)
- [299] Buchmüller W and Plümacher M 1996 Baryon asymmetry and neutrino mixing *Phys. Lett.* **B389** 73–77
- [300] Sierra D A and Varzielas I M 2013 The role of lepton flavor symmetries in leptogenesis *Fortsch. Phys.* **61** 645–665 (arXiv:1205.6134)
- [301] Jenkins E E and Manohar A V 2008 Tribimaximal mixing, leptogenesis and θ_{13} , *Phys. Lett.* **B668** 210–215 (arXiv:0807.4176)
- [302] Hagedorn C, Molinaro E and Petcov S T 2009 Majorana phases and leptogenesis in seesaw models with A_4 symmetry *JHEP* **0909** 115 (arXiv:0908.0240)
- [303] Bertuzzo E, Bari P D, Feruglio F and Nardi E 2009 Flavor symmetries, leptogenesis and the absolute neutrino mass scale *JHEP* **0911** 036 (arXiv:0908.0161)
- [304] Sierra D A, Bazzocchi F, Varzielas I M, Merlo L and Morisi S 2010 Tribimaximal lepton mixing and leptogenesis *Nucl. Phys.* **B827** 34–58 (arXiv:0908.0907)
- [305] Felipe R and Serodio H 2010 Constraints on leptogenesis from a symmetry viewpoint *Phys. Rev.* **D81** 053008 (arXiv:0908.2947)
- [306] Choubey S, King S and Mitra M 2010 On the vanishing of the CP asymmetry in leptogenesis due to form dominance *Phys. Rev.* **D82** 033002 (arXiv:1004.3756)
- [307] Mohapatra R N and Nasri S 2005 Leptogenesis and μ - τ symmetry *Phys. Rev.* **D71** 033001 (arXiv:hep-ph/0410369)
- [308] Mohapatra R N, Nasri S and Yu H B 2005 Leptogenesis, μ - τ symmetry and θ_{13} *Phys. Lett.* **B615** 231–239 (arXiv:hep-ph/0502026)
- [309] Ahn Y H, Kang S K, Kim C S and Lee J 2006 Phased breaking of μ - τ symmetry and leptogenesis *Phys. Rev.* **D73** 093005 (arXiv:hep-ph/0602160)
- [310] Ahn Y H, Kang S K, Kim C S and Lee J 2007 μ - τ symmetry and radiatively generated leptogenesis *Phys. Rev.* **D75** 013012 (arXiv:hep-ph/0610007)
- [311] Hamzaoui C, Nasri S and Toharia M 2014 Partial μ - τ textures and leptogenesis *Phys. Rev.* **D89** 073019 (arXiv:1311.2188)
- [312] Lashin E I, Chamoun N, Hamzaoui C and Nasri S 2014 Neutrino mass textures and partial μ - τ symmetry *Phys. Rev.* **D89** 093004 (arXiv:1311.5869)
- [313] Lashin E I, Chamoun N, Hamzaoui C and Nasri S 2015 Neutrino mixing and leptogenesis in μ - τ symmetry *Phys. Rev.* **D91** 113014 (arXiv:1408.5042)
- [314] Davidson S and Ibarra A 2002 A lower bound on the right-handed neutrino mass from leptogenesis *Phys. Lett.* **B535** 25–32 (arXiv:hep-ph/0202239)

- [315] Frampton P H, Glashow S L and Yanagida T 2002 Cosmological sign of neutrino CP violation *Phys. Lett.* **B548** 119–121
- [316] King S F and Luhn C 2012 A4 models of tribimaximal reactor mixing *JHEP* **1203** 036 (arXiv:1112.1959)
- [317] Ahn Y H, Kang S K, Kim C S and Nguyen T P 2008 Bridges of low energy observables with leptogenesis in μ - τ reflection symmetry (arXiv:0811.1458)
- [318] Abada A, Davidson S, Josse-Michaux F-X, Losada M and Riotto A 2006 Flavour issues in leptogenesis *JCAP* **0604** 004 (arXiv:hep-ph/0601083)
- [319] Nardi E, Nir Y, Roulet E and Racker J 2006 The importance of flavor in leptogenesis *JHEP* **0601** 164 (arXiv:hep-ph/0601084)
- [320] Fritzsh H 1978 Weak interaction mixing in the six-quark theory *Phys. Lett.* **B73** 317–322
- [321] Fritzsh H 1979 Quark masses and flavor mixing *Nucl. Phys.* **B155** 189–207
- [322] Fritzsch H 1987 Hierarchical chiral symmetries and the quark mass matrix *Phys. Lett.* **B184** 391–396
- [323] Fritzsch H and Xing Z Z 1997 Flavor symmetries and the description of flavor mixing *Phys. Lett.* **B413** 396–404 (arXiv:hep-ph/9707215)
- [324] Fritzsch H and Xing Z Z 1998 On the parametrization of flavor mixing in the standard model *Phys. Rev.* **D57** 594–597 (arXiv:hep-ph/9708366)
- [325] Xing Z Z and Zhang H 2004 Complete parameter space of quark mass matrices with four texture zeros *J. Phys.* **G30** 129–136 (arXiv:hep-ph/0309112)
- [326] Xing Z Z and Zhao Z H 2015 On the four-zero texture of quark mass matrices and its stability *Nucl. Phys.* **B897** 302–325 (arXiv:1501.06346)
- [327] Koide Y, Nishiura H, Matsuda K, Kikuchi T and Fukuyama T 2002 Universal texture of quark and lepton mass matrices and a discrete symmetry $Z(3)$ *Phys. Rev.* **D66** 093006 (arXiv:hep-ph/0209333)
- [328] Matsuda K and Nishiura H 2004 Assignments of universal texture components for quark and lepton mass matrices *Phys. Rev.* **D69** 053005 (arXiv:hep-ph/0309272)
- [329] Koide Y 2004 Universal texture of quark and lepton mass matrices with an extended flavor $2 \leftrightarrow 3$ symmetry *Phys. Rev.* **D69** 093001 (arXiv:hep-ph/0312207)
- [330] Matsuda K and Nishiura H 2005 Prediction for quark mixing from universal quark and lepton mass matrices with flavor $2 \leftrightarrow 3$ symmetry *Phys. Rev.* **D71** 073001 (arXiv:hep-ph/0501201)
- [331] Matsuda K and Nishiura H 2005 Large lepton mixing and nonsymmetric mass matrices with flavor $2 \leftrightarrow 3$ symmetry *Phys. Rev.* **D72** 033011 (arXiv:hep-ph/0506192)
- [332] Matsuda K and Nishiura H 2006 Broken flavor $2 \leftrightarrow 3$ symmetry and phenomenological approach for universal quark and lepton mass matrices *Phys. Rev.* **D73** 013008 (arXiv:hep-ph/0511338)
- [333] Joshipura A S 2008 Universal 2-3 symmetry *Eur. Phys. J.* **C53** 77–85 (arXiv:hep-ph/0512252)
- [334] Fritzsch H and Xing Z Z 1996 Lepton mass hierarchy and neutrino oscillations *Phys. Lett.* **B372** 265–270 (arXiv:hep-ph/9509389)
- [335] Fritzsch H and Xing Z Z 1996 Large leptonic flavor mixing and the mass spectrum of leptons *Phys. Lett.* **B440** 313–318 (arXiv:hep-ph/9808272)
- [336] Branco G C, Rebelo M N and Silva-Marcos J I 1998 Quasi-degenerate neutrino masses with universal-strength Yukawa coupling *Phys. Lett.* **B428** 136–142 (arXiv:hep-ph/9802340)
- [337] Barr S M and Dorsner I 2000 A general classification of three neutrino models and U_{e3} *Nucl. Phys.* **B585** 79–104 (arXiv:hep-ph/0003058)
- [338] Rodejohann W and Xing Z Z 2004 Flavor democracy and type-II seesaw realization of bilarge neutrino mixing *Phys. Lett.* **B601** 176–183 (arXiv:hep-ph/0408195)
- [339] Adhikary B, Ghosal A and Roy P 2009 Mu-tau symmetry, tribimaximal mixing and four zero neutrino Yukawa textures *JHEP* **0910** 040 (arXiv:0908.2686)
- [340] Adhikary B, Ghosal A and Roy P 2013 θ_{13} , μ - τ symmetry breaking and neutrino Yukawa textures *Int. J. Mod. Phys.* **A28** 1350118 (arXiv:1210.5328)
- [341] Mohapatra R N, Nasri S and Yu H B 2006 Grand unification of μ - τ symmetry *Phys. Lett.* **B636**

- 114–118 (arXiv:hep-ph/0603020)
- [342] Joshipura A S, Kodrani B P and Patel K M 2009 Fermion masses and mixings in a μ - τ symmetric SO(10) *Phys. Rev.* **D79** 115017 (arXiv:0903.2161)
- [343] Branco G C, Gonzalez Felipe R and Joaquim F R 2012 Leptonic CP violation *Rev. Mod. Phys.* **84** 515–565 (arXiv:1111.5332)
- [344] An F P *et al* 2016 Neutrino physics with JUNO *J. Phys.* **G43** 030401 (arXiv:1507.05613)
- [345] Bilenky S M and Giunti C 2015 Neutrinoless double-beta decay: a probe of physics beyond the standard model *Int. J. Mod. Phys.* **A30** 1530001 (arXiv:1411.4791)
- [346] Giunti C and Studenikin 2015 Neutrino electromagnetic interactions: a window to new physics *Rev. Mod. Phys.* **87** 531–591 (arXiv:1403.6344)
- [347] Xing Z Z 2014 Quark mass hierarchy and flavor mixing puzzles *Int. J. Mod. Phys.* **A29** 1430067 (arXiv:1411.2713)