

# Tree-Structured Clustering in Fixed Effects Models

Moritz Berger & Gerhard Tutz  
Ludwig-Maximilians-Universität München  
Akademiestraße 1, 80799 München

August 8, 2021

## Abstract

Fixed effects models are very flexible because they do not make assumptions on the distribution of effects and can also be used if the heterogeneity component is correlated with explanatory variables. A disadvantage is the large number of effects that have to be estimated. A recursive partitioning (or tree based) method is proposed that identifies clusters of units that share the same effect. The approach reduces the number of parameters to be estimated and is useful in particular if one is interested in identifying clusters with the same effect on a response variable. It is shown that the method performs well and outperforms competitors like the finite mixture model in particular if the heterogeneity component is correlated with explanatory variables. In two applications the usefulness of the approach to identify clusters that share the same effect is illustrated.

**Keywords:** Fixed effects model; random effects model; recursive partitioning; tree-structured regression; regularization.

## 1 Introduction

The analysis of longitudinal data and cross-sectional data that come in clusters requires to take the dependence of observations and the heterogeneity of measurement units into account. Typically, measurements within units tend to be more similar than measurements between units. If the heterogeneity is ignored poor performance of estimators and misleading standard errors are to be expected.

The most popular, widely used model to account for unobserved heterogeneity is the random effects model, see, for example, Verbeke and Molenberghs (2000), Molenberghs and Verbeke (2005) and McCulloch and Searle (2001). Typically in the random effects model it is assumed that the random effects follow a normal

distribution. This strong assumption results in an economical model but inference may be sensitive to the specification of the distribution of random effects, see Heagerty and Kurland (2001), Agresti et al. (2004) and Litière et al. (2007).

Several approaches to weaken the assumption of normally distributed random effects have been proposed. More flexible distributions are obtained, for example, by using mixtures of normals as proposed by Chen and Davidian (2002) and Magder and Zeger (1996). Huang (2009) proposed diagnostic methods for random-effect misspecification and Claeskens and Hart (2009) proposed tests for the assumption of the normal distribution. More recently, Lombardía and Sperlich (2012) proposed the class of semi-mixed effects models, a continuum of models that combine random and fixed effects.

An alternative approach to model heterogeneity uses finite mixtures. In finite mixtures of generalized linear models it is assumed that the density or mass function of the responses given the explanatory variables is determined by a finite mixture of components. Each of the components has its own response distribution and own parameters that determine the influence of explanatory variables. If only part of the parameters, for example the intercepts, are allowed to vary over components one obtains a discrete distribution of the heterogeneity part of the model. Models of that type were considered by Follmann and Lambert (1989) and Aitkin (1999). Follmann and Lambert (1989) investigated the identifiability of finite mixtures of binomial regression models and gave sufficient identifiability conditions for mixing of binary and binomial distributions. Grün and Leisch (2008b) considered identifiability for mixtures of multinomial logit models.

Finite mixture models replace the assumption of a fixed continuous distribution of random effects by the assumption of a discrete distribution. One may see this as an alternative and flexible specification of the heterogeneity component only. However, by assuming a discrete distribution of the intercepts instead of a continuous distribution as in random effects models one also implicitly assumes that there are clusters of units that share the same effect. In some applications it is definitely of interest to identify these units. We will consider an example in which the units are schools and one wants to know which schools are similar in their performance with regard to the education of students.

Here we consider an alternative to finite mixture models with the same objectives, that are use of a flexible discrete distribution and identification of units that share the same effect. However, the starting point is different. We use a fixed effects model in which each unit has its own parameter. An advantage is that no structural assumptions on the unit-specific effects have to be made. Clusters of parameters and therefore units with the same effect are found by tree methodology, although a different one as in classical trees.

Classical recursive partitioning techniques or trees were first introduced by Morgan and Sonquist (1963). Very popular methods are classification and regression trees (CART) by Breiman et al. (1984) and C4.5 by Quinlan (1986) and Quinlan (1993). A newer version of recursive partitioning based on conditional

inference was proposed by Hothorn et al. (2006). An overview on recursive partitioning in health science was given by Zhang and Singer (1999) and with a focus on psychometrics by Strobl et al. (2009). An easily accessible introduction into the basic concepts is found in Hastie et al. (2009).

The tree methodology used here differs from these approaches. In CART and other classical approaches the whole covariate space is recursively partitioned into subspaces. In order to obtain a partitioning in the intercepts (or slopes) only, one has to apply a different form of trees. It has to be designed in a way that the subspaces are built for specific effects only, for example the intercepts, while other parameters that represent common effects of explanatory variables are not partitioned into subspaces. Our main focus is on the clustering of intercepts, however, we will also refer to the case of unit-specific slopes. One big advantage using recursive partitioning techniques is the computational efficiency. The proposed tree-structured model especially enables the evaluation of high-dimensional data. Alternative approaches to identify clusters within a fixed effects model framework as proposed by Tutz and Oelker (2015) fail in high dimensional settings.

The article is organized as follows: In Section 2 we introduce the tree-structured model for unit-specific intercepts and in section 3 we present an illustrative example. Details about the fitting procedure are given in Section 4. After a short introduction of related approaches in Section 5 we give the results of wider simulation studies (Section 6). Finally, Section 7 contains a second application.

## 2 Accounting for Heterogeneity in Clustered Data

Consider clustered data given by  $(y_{ij}, \mathbf{x}_{ij}, \mathbf{z}_{ij})$ ,  $i = 1, \dots, n$ ,  $j = 1, \dots, n_i$ , where  $y_{ij}$  denotes the response of measurement  $j$  for unit  $i$  and two sets of predictive variables  $\mathbf{x}_{ij}^\top = (1, x_{ij1}, \dots, x_{ijp})$  and  $\mathbf{z}_{ij}^\top = (1, z_{ij1}, \dots, z_{ijq})$ . In longitudinal data the units can, for example, represent persons that are measured repeatedly. In the following, we consider alternative methods to account for the potential heterogeneity of units. We start with methods that use random effects, then consider fixed effects model and finite mixtures.

### 2.1 Random Effects Models

In a generalized linear mixed model (GLMM) the mean response  $\mu_{ij} = \mathbb{E}(y_{ij} | \mathbf{b}_i, \mathbf{x}_{ij}, \mathbf{z}_{ij})$  is linked to the explanatory variables by

$$g(\mu_{ij}) = \mathbf{x}_{ij}^\top \boldsymbol{\beta} + \mathbf{z}_{ij}^\top \mathbf{b}_i, \quad (1)$$

where  $\mathbf{x}_{ij}^\top \boldsymbol{\beta}$  is a linear term which contains the fixed effect  $\boldsymbol{\beta}$ . The second term  $\mathbf{z}_{ij}^\top \mathbf{b}_i$  contains the random effects for covariates  $\mathbf{z}_{ij}$  that are varying across units and  $g(\cdot)$  is a known link function. In a GLMM it is assumed that the distribution

of  $y_{ij}|\mathbf{b}_i, \mathbf{x}_{ij}, \mathbf{z}_{ij}$  follows a simple exponential family and that the observations  $y_{ij}$  are conditionally independent. For the random effects  $\mathbf{b}_i$ , which model the heterogeneity of the units, one typically assumes a normal distribution  $\mathbf{b}_i \sim N(\mathbf{0}, \boldsymbol{\Sigma}_{rand})$ .

In a GLMM the distribution of the random effects is used to account for the heterogeneity of the units and the focus is mainly on the parametric term  $\mathbf{x}_{ij}^\top \boldsymbol{\beta}$ . Although the distributional assumption for the random effects makes the estimation of the model very efficient there are also some disadvantages. If the assumed distribution is very different from the real data generating distribution, inference can be biased. The assumption of a continuous distribution also does not allow for the same effects of different units. Hence, clustering of units is not possible. Another crucial point of the GLMM is the assumption that the random effects  $\mathbf{b}_i$  and the covariates  $\mathbf{x}_{ij}$  are uncorrelated. This assumption can lead to poor estimation accuracy, see, for example, Grilli and Rampichini (2011). Functions for the estimation of generalized linear mixed models are provided by the R-package `lme4` (Bates et al., 2015), which we will use for the computations in the applications and simulations.

## 2.2 Fixed Effects Models

In contrast to mixed models, fixed effects models model heterogeneity among units by using one parameter  $\beta_i$  for each unit. The mean response  $\mu_{ij} = \mathbb{E}(y_{ij}|\mathbf{x}_{ij}, \mathbf{z}_{ij})$  is linked to the explanatory variables in the form

$$g(\mu_{ij}) = \eta_{ij} = \mathbf{x}_{ij}^\top \boldsymbol{\beta} + \mathbf{z}_{ij}^\top \boldsymbol{\beta}_i, \quad (2)$$

where  $\mathbf{x}_{ij}$  again is a vector of covariates that have the same effect across all units and  $\mathbf{z}_{ij}$  contains covariates that have different effects over units. Each measurement unit has his own parameter vector  $\boldsymbol{\beta}_i^\top = (\beta_{i0}, \dots, \beta_{iq})$ . The specification of one parameter vector per unit results in a very large number of parameters which can affect estimation accuracy. Moreover, typically there is not enough information to distinguish between all units. To cope with these problems one can assume that there are groups of units that share the same effect on the response. Forming clusters of units leads to a reduced number of parameters and stable estimates. There are several strategies to identify these clusters, the fixed effects model with regularization considered in the next section or the finite mixture model (Section 2.4).

## 2.3 Tree-Structured Clustering

In the approach considered here one assumes that the fixed effects model holds, but not all the unit-specific parameters are assumed to be different. Clusters (or groups) of measurement units are identified by recursive partitioning methods. We first consider unit-specific intercepts only. Let us start with the simplest

case in which all intercepts are equal, that is, the linear predictor has the form  $\eta_{ij} = \mathbf{x}_{ij}^\top \boldsymbol{\beta} + \beta_0$ . If there are two clusters the corresponding linear predictor is given by

$$\eta_{ij} = \mathbf{x}_{ij}^\top \boldsymbol{\beta} + \beta_{i0}^{(k)}, \quad k = 1, 2, \quad (3)$$

where  $k$  denotes if the unit is in the first or the second group. A simple test, for example a likelihood ratio test, for the hypothesis  $H_0 : \beta_{i0}^{(1)} = \beta_{i0}^{(2)}$  can be used to determine if the model with two groups is more adequate for the data than the model in which all the intercepts are equal. By iterative splitting into subsets guided by test statistics one obtains a clustering of units that have to be distinguished with regard to their intercept.

In general, regression trees can be seen as a representation of a partition of the predictor space. A tree is built by successively splitting one node  $A$ , that is already a subset of the predictor space, into two subsets  $A_1$  and  $A_2$  with the split being determined by only one variable. In a fixed effects model, when specifying specific intercepts for each unit, the unit number itself can be seen as a nominal categorical variable with  $n$  categories. The partition has the form  $A \cap S_1, \quad A \cap S_2$ , where  $S_1$  and  $S_2$  are disjoint, non-empty subsets  $S_1 \subset \{1, \dots, n\}$  and its complement  $S_2 = \{1, \dots, n\} \setminus S_1$ . Using this notation another representation of model (3) is given by

$$\eta_{ij} = \mathbf{x}_{ij}^\top \boldsymbol{\beta} + \beta_{i0}^{(1)} I(i \in S_{10}) + \beta_{i0}^{(2)} I(i \in S_{20}),$$

where  $I(\cdot)$  denotes the indicator function with  $I(a) = 1$ , if  $a$  is true and  $I(a) = 0$  otherwise. After several splits one obtains a clustering of the units  $\{1, \dots, n\}$  and the predictor of the resulting model can be represented by

$$\eta_{ij} = \mathbf{x}_{ij}^\top \boldsymbol{\beta} + \sum_{k=1}^{m_0} \beta_{i0}^{(k)} I(i \in S_{k0}) \quad (4)$$

where  $S_{10}, \dots, S_{m_0 0}$  is a partition of  $\{1, \dots, n\}$  consisting of  $m_0$  clusters that have to be distinguished in terms of their individual intercepts.

In the following we will use the model abbreviation *TSC* for tree-structured clustering.

## 2.4 Finite Mixture Models

An alternative approach that also allows to identify clusters of units are finite mixture models. These were, for example, considered by Follmann and Lambert (1989) and Aitkin (1999). The general assumption in finite mixtures of generalized regression models is that the mixture consists of  $K$  components where each component follows a parametric distribution of the exponential family of distributions. The density of the mixture can be given by

$$f(y|\mathbf{x}, \boldsymbol{\beta}, \boldsymbol{\phi}) = \sum_{k=1}^K \pi_k f_k(y|\mathbf{x}, \boldsymbol{\beta}_k, \boldsymbol{\phi}_k)$$

where  $f_k(y|\mathbf{x}, \boldsymbol{\beta}_k, \phi_k)$  denotes the  $k$ -th component of the mixture with parameter vector  $\boldsymbol{\beta}_k$  and dispersion parameter  $\phi_k$ . For the unknown component weights  $\pi_k$   $\sum_{k=1}^K \pi_k = 1$  and  $\pi_k > 0$ ,  $k = 1, \dots, K$  has to hold.

Here we consider models with components that differ in their intercepts. Within the framework of finite mixtures one specifies for the  $k$ -th component of the mixture a model with predictor  $\eta_{ij}^{(k)} = \beta_{i0}^{(k)} + x_{ij}^\top \boldsymbol{\beta}$ . For models with normal response the mixture components are given by  $N(y_{ij}|\eta_{ij}^{(k)}, \sigma^2)$ , where the variance  $\sigma^2$  is fixed for all components. For models with a binary response the mixture components are  $B(y_{ij}|n, \pi_{ij}^{(k)})$ , where  $\pi_{ij}^{(k)} \in (0, 1)$  and  $\text{logit}(\pi_{ij}^{(k)}) = \eta_{ij}^{(k)}$ . For further details, see Grün and Leisch (2007).

Estimation of the mixture model is usually obtained by the EM-algorithm with the number of components  $K$  being specified beforehand. The optimal number of components is chosen afterwards, for example by information criteria like AIC or BIC. Grün and Leisch (2008a) provide the R-package `flexmix`, which is used for the computations in our applications and simulations. Regularization and variable selection for mixture models have been considered by Khalili and Chen (2007) and Städler et al. (2010) but not with the objective of clustering units with regard to their effects.

TABLE 1: *Summary statistics of the test score of the 56 multiple-choice items and covariate gender of the CTB data of the illustrative example (CTB data).*

Variable	Summary statistics					
	$x_{min}$	$x_{0.25}$	$x_{med}$	$\bar{x}$	$x_{0.75}$	$x_{max}$
Test score	21	32	34	34.14	37	46
Gender	male: 761			female: 739		

### 3 An Illustrative Example

Before giving details how to grow trees and estimate the proposed model (4) we want to illustrate the procedure by use of an application. We consider a data set from CTB/McGraw-Hill, a division of the Data Recognition Corporation (DRC). For a description of the original data, see De Boeck and Wilson (2004). The data includes results of an achievement test that measures different objectives and subskills of subjects in mathematics and science. For our investigation we use the results of 1500 grade 8 students from 35 schools. They had to respond to 56 multiple-choice items (31 mathematics, 25 science). The response  $y_{ij}$  is the overall test score of student  $j$  in school  $i$ , defined as the number of correctly solved items. The main objective is to adequately describe the heterogeneity of the 35




FIGURE 1: Paths of coefficients of school-specific intercepts against all splits of the illustrative example (CTB data). The optimal number of splits is marked by a dashed line.




FIGURE 2: Comparison of the estimated distribution of the mixed model and the school-specific intercepts of tree-structured clustering (CTB data).

schools. As additional covariate we include the gender of the students (male: 0, female: 1). The summary statistics of the test scores and the covariate gender is given in Table 1. By using the proposed tree-structured approach the model that was obtained has the form

$$\mu_{ij} = \beta_G \cdot G_{ij} + \sum_{k=1}^{m_0} \beta_{i_0}^{(k)} I(i \in S_{k0}), \quad i = 1, \dots, 35,$$

TABLE 2: *Estimation results of the illustrative example (CTB data) using the classical mixed model, tree-structured clustering and the finite mixture model.*

Predictor	LMM		TSC		FIN	
	Coefficient	95%-CI	Coefficient	95%-CI	Coefficient	95%-CI
gender	-0.106	[-0.475, 0.298]	-0.088	[-0.478, 0.313]	-0.084	[-0.473, 0.309]
$\beta_0$	34.235	[33.964, 34.542]	—	—	—	—
$\sigma_{\text{rand}}^2$	0.416	[0.394, 1.353]	—	—	—	—

School-specific intercept	TSC		FIN	
	Cluster	Coefficient	Cluster	Coefficient
$\beta_{i0}$	1,16	32.384	1,4,6,7,9,16,	33.508
	4,18,19,20,21,22,28	33.434	18,19,20,21,	
	6,7,9,11,29,30	33.904	22,28,30	
	3,5,12,14,15,25,26,31,34	34.517	2,3,5,8,10,11,12,13,14,	34.689
	2,10,13,17,23,24,32	34.999	15,17,23,24,25,26,27,	
8,27,33,35	36.264	29,31,32,33,34,35		

where  $G_{ij} \in \{0, 1\}$  denotes the gender of student  $j$  in school  $i$ ,  $S_{10}, \dots, S_{m_0}$  is a partition of the 35 schools and  $\beta_{i0}^{(k)}$ ,  $k = 1, \dots, m_0$ , denote the effects of the corresponding clusters.

The coefficient paths of the school-specific intercepts obtained by tree-structured clustering are shown in Figure 1. The coefficient paths build a tree that successively partitions the schools in terms of the performance of students. The left end refers to the global intercept estimated as an average over the 35 schools. On the right end of the coefficients paths all possible splits have been performed and the estimated coefficients correspond to those of a simple fixed effects model without clustering. The optimal number of splits that is selected by the algorithm, is marked by the dashed line. It is seen that estimates change strongly in the first steps, but after about ten splits the estimates are very stable.

A graphical comparison of the estimated normal distribution of the random effects using a classical linear mixed model and the distribution of the school-specific intercepts of the tree-structured model is shown in Figure 2. It illustrates the main advantage of the tree-structured model. There is no distributional assumption on the school-specific intercepts, especially no assumption of symmetry. The number of schools in each cluster are quite different and not symmetric. Clustering of similar schools strongly reduces the complexity of the fixed effects model and makes interpretation of school-specific differences very easy. There are two small clusters of schools where the performance in the test considerably deviates upwards or downwards, the differences between the clusters with medium performance are smaller.

Table 2 shows an overview of the estimation results obtained by using the classical linear mixed model (LMM), the proposed tree-structured model (TSC)



and a finite mixture model (FIN), where only the intercepts are allowed to vary over the components. Confidence intervals are obtained by using bootstrap procedures, where the model is fitted repeatedly on sub samples of size  $n$  that are obtained by drawing with replacement. The results here are obtained by 2000 sub samples. It is seen that all of the methods did not find a significant effect for covariate gender. The performance of males and females seems not to differ systematically. The variance obtained by the mixed model is significantly different from zero, which suggests that heterogeneity of schools is definitely present. The lower panel in Table 2 shows the estimated partition of schools obtained by the tree-structured model and the finite mixture model. In the latter case, model selection by AIC and BIC both yield the same result. Tree-structured clustering identifies six clusters of schools until further splits are no longer significant (for details of the algorithm see Section 4). The finite mixture approach identifies only two clusters of schools. This illustrates the tendency of the finite mixture approach to find a small number of clusters, which will be investigated later. For comparison in Table 2 the schools that belong to the two clusters found by the finite mixture model are coloured in black and grey.

## 4 Fitting procedure

In this section we give details of the algorithm that yields the tree-structured model. Let us again consider the model with unit-specific intercepts after the first split, which has the form

$$\eta_{ij} = \mathbf{x}_{ij}^\top \boldsymbol{\beta} + \beta_{i0}^{(1)} I(i \in S_{10}) + \beta_{i0}^{(2)} I(i \in S_{20}). \quad (5)$$

When determining the first split for the nominal predictor  $i \in \{1, \dots, n\}$  one has to consider all possible partitions of the two subsets  $S_{10}$  and  $S_{20}$ . Altogether there are  $2^{n-1} - 1$  possible splits, which can be a very large number. It has been shown in earlier research that it is not necessary to consider all possible partitions, see Breiman et al. (1984) and Ripley (1996) for binary outcomes and Fisher (1958) for quantitative outcomes. It is sufficient to order the predictor categories, here the measurement units, with respect to the means of the response and to treat the predictor as if the categories were ordered. In a first step, units are ordered according to their maximum-likelihood estimates, so that  $\hat{\beta}_{(10)} \leq \hat{\beta}_{(20)} \leq \dots \leq \hat{\beta}_{(n0)}$ . Then one considers splits of adjacent measurement units to obtain the optimal split. To use this simplification one starts with an equivalent representation of model (5) given by

$$\eta_{ij} = \mathbf{x}_{ij}^\top \boldsymbol{\beta} + \beta_0 + \alpha_{i0} I(i > c),$$

with  $\beta_{i0}^{(1)} = \beta_0$  and  $\beta_{i0}^{(2)} = \beta_0 + \alpha_{i0}$ . The set  $C$  of possible thresholds  $c$  is from  $\{1, \dots, n-1\}$ . The fitting procedure considered in the following uses this model as building block. By iterative splitting of adjacent measurement units the searched-for clustering is obtained.

## Basic Algorithm

The basic algorithm for the model with unit-specific intercept is the following.

---

### Tree-Structured Clustering – Unit-specific intercept

#### *Step 1 (Initialization)*

- (a) Estimation: Fit the candidate GLMs with predictors

$$\eta_{ij} = \mathbf{x}_{ij}^\top \boldsymbol{\beta} + \beta_0 + \alpha_{i0} I(i > c_{i0}), \quad i = 1, \dots, n-1$$

- (b) Selection

Select the model that has the best fit. Let  $c_{i_1 0}^*$  denote the best split.

#### *Step 2 (Iteration)*

For  $\ell = 1, 2, \dots$ ,

- (a) Estimation: Fit the candidate models with predictors

$$\eta_{ij} = \mathbf{x}_{ij}^\top \boldsymbol{\beta} + \beta_0 + \sum_{s=1}^{\ell} \alpha_{i_s 0} I(i > c_{i_s 0}^*) + \alpha_{i0} I(i > c_{i0}),$$

for all values  $c_{i0} \in C \setminus \{c_{i_1 0}^*, \dots, c_{i_\ell 0}^*\}$

- (b) Selection

Select the model that has the best fit yielding the split point  $c_{i_{\ell+1} 0}^*$ .

---

In each selection step of the algorithm one has to identify the best split and during the iterations one has to decide when to stop. Common splitting criteria for tree-based methods are impurity measures that have already been introduced by Breiman et al. (1984). An alternative is to use a test statistic to evaluate which split most improves the explanatory power of the predictors. We will draw on the latter concept and use a procedure that is strongly related to the conditional inference framework proposed by Hothorn et al. (2006).

In each iteration one examines the null hypotheses  $H_0 : \alpha_{i0} = 0$  for all remaining possible split points. This can, for example, be tested by a likelihood-ratio test. To determine the best split we simultaneously consider all test statistics  $T_{i0}$  from the set of possible splits  $c_{i0}$  and choose the split point for which  $T_{i0}$  had the largest value. This corresponds to choosing the split with the smallest  $p$ -value obtained from the chi-squared distribution of the test statistic. To determine the optimal number of splits our strategy is to check if the heterogeneity of

measurement units is already modelled sufficiently in each step. Before executing one further split one tests the global null hypothesis that the current model completely captures the heterogeneity of the data against the alternative that the data is more heterogeneous. To decide for the first split one has to examine the null hypothesis  $H_0 : \beta_{10} = \beta_{20} = \dots = \beta_{n0}$ , which corresponds to the case of no heterogeneity. The hypothesis is tested by a likelihood-ratio test with significance level  $\alpha$  and  $n - 1$  degrees of freedom, because  $n - 1$  differences of parameters are tested. Depending on the significance of this global test the selected split or no splitting is performed. After several splits only differences of units within already built clusters are tested. In the  $\ell - th$  step  $n - \ell$  differences have to be tested because  $\ell - 1$  splits are already performed. If a significant effect is found the selected split is performed, otherwise splitting is stopped. The proposed stopping criterion leads to a clear separation of the selection of splits and the splitting decision. In particular the splitting decision is not influenced by the previously identified ordering of measurement units.

The result of the fitting procedure is a sequence of  $m_0 - 1$  selected split points  $c_{i_1 0}^*, \dots, c_{i_{m_0-1} 0}^*$  and corresponding parameter estimates  $\hat{\alpha}_{i_1 0}, \dots, \hat{\alpha}_{i_{m_0-1} 0}$ . Ordering of the selected split points yields the desired clustering of ordered units  $\{1, \dots, c_{i_1 0}^*\}, \{c_{i_1 0}^* + 1, \dots, c_{i_2 0}^*\}, \dots, \{c_{i_{m_0-1} 0}^* + 1, \dots, n\}$ . The corresponding intercepts  $\beta_{i_0}^{(k)}$  for each cluster are then given by

$$\hat{\beta}_{i_0}^{(k)} = \hat{\beta}_0 + \sum_{s=1}^{k-1} \hat{\alpha}_{(i_s 0)}, \quad k = 1, \dots, m_0.$$

During the iterations only the selected split points but no estimates from previous steps are kept. All coefficients of the models, including the parameters  $\beta$  of the linear term, are refitted in each step and the final estimates are those from the last iteration.

## 5 Related Approaches

In the following we will briefly consider alternative methods that account for unobserved heterogeneity and are related to our tree-structured model. One of the approaches is a competitor to the method proposed here and will also be included in the simulations.

Clustering of units can also be obtained by penalized maximum likelihood estimation as proposed more recently by Tutz and Oelker (2015). Let  $\beta_0^T = (\beta_{10}, \dots, \beta_{n0})$  denote the intercepts of the fixed effects model. An estimation procedure that identifies clusters is obtained by maximizing the penalized log-likelihood  $l_p(\beta, \beta_0) = l(\beta, \beta_0) - \lambda J(\beta, \beta_0)$ , where  $l(\beta, \beta_0)$  denotes the unpenalized log-likelihood,  $J(\beta, \beta_0)$  is a specific penalty term and  $\lambda$  is a tuning parameter.

The penalty term that enforces clustering of unit-specific intercepts is given by

$$J(\boldsymbol{\beta}, \boldsymbol{\beta}_0) = \sum_{r>s} |\beta_{r0} - \beta_{s0}|,$$

where only pairwise differences of the unit-specific intercepts are included. If  $\lambda = 0$ , one obtains the unpenalized maximum-likelihood estimates and each unit has his own intercept. If  $\lambda \rightarrow \infty$ , all units are fused to one cluster with the same intercept. For a comparison we use the corresponding R-package `gvcm.cat` proposed by Oelker (2015) in our simulations. The use of such penalties in ANOVA was already proposed by Bondell and Reich (2008) and for variable selection by Gertheiss and Tutz (2010) and Tutz and Gertheiss (2014). A problem with the method is that the penalty contains  $n(n-1)/2$  differences and therefore the algorithm becomes extremely demanding for large values of  $n$ . It typically fails if the number of groups is larger than 50 or 60.

The method proposed here should be distinguished from the mixed effects regression trees (MERT) proposed by Hajjem et al. (2011) and the RE-EM trees, which were independently proposed by Sela and Simonoff (2012). The basic concept is to combine a linear mixed effects model for clustered data and a standard regression tree. The substantial difference is that the tree is not applied to the random or unit-specific effects of the model but to the fixed effects term. The predictor of the estimated model has the form  $\eta_{ij} = f(\mathbf{x}_{ij}) + \mathbf{z}_{ij}^\top \mathbf{b}_i$ , where  $\mathbf{b}_i \sim N(0, \boldsymbol{\Sigma}_{rand})$ . It is the function  $f(\mathbf{x}_{ij})$  that is estimated by a standard regression tree. The model yields random effects that are node-invariant and therefore does not focus on the similarity of units but rather on the dissimilarity of observations within units.

An alternative Bayesian approach to model clustered random effects is based on Dirichlet processes. Dirichlet processes were proposed by Ferguson (1973) and studied, for example, by Sethuraman (1994) and Hjort et al. (2010). The main advantage of Dirichlet processes is their cluster property, which allows to flexibly model discrete distributions. Assuming a Dirichlet process for the distribution of random effects creates ties among the random effects. The resulting Dirichlet process mixture yields clusters of units. Dirichlet process priors have been used within the linear mixed model framework by Bush and MacEachern (1996) and Müller and Rosner (1997). A frequentist approach to linear mixed models with Dirichlet process mixtures was given by Heinzl and Tutz (2013), a combination of Dirichlet processes and fusion penalties was considered in Heinzl and Tutz (2014), Heinzl and Tutz (2015). The approach works for linear models, but extensions to generalized mixed models seem not available.

## 6 Simulations

In the following we investigate the performance of the proposed tree-structured model and compare it to competing methods. The focus is on data settings with

clusters of units that share the same effect on the response and where the strict assumptions of the mixed model do not hold. We are in particular interested in the estimation accuracy and the clustering performance. We will compare the generalized fixed effects model (GFM), the generalized mixed model (GMM), the tree-structured model (TSC), the model based on penalized maximum-likelihood estimation (PENL), the finite mixture model with model selection by AIC (FINA) and the finite mixture model with model selection by BIC (FINB).

We consider several simulation scenarios where the overall number of observations is 800, made up of the components  $n = 200/n_i = 4$ ,  $n = 100/n_i = 8$ ,  $n = 40/n_i = 20$  or  $n = 20/n_i = 40$ . In addition to the unit-specific intercepts we include one continuous covariate  $x_1$  with  $x_{ij1} \sim N(0, 1)$  and one binary covariate  $x_2$  with  $x_{ij2} \sim B(1, 0.5)$ . Unit-specific intercepts  $\beta_{i0}$  are drawn symmetrically from a normal distribution or are drawn from a chi-square distribution that is skewed. In order to obtain clusters of units, the intercepts are sorted according to size and divided into balanced groups. The average over the intercepts of each group is defined as the new unit-specific intercept  $\beta_{i0}^{(k)}$ ,  $k = 1, \dots, m_0$ . We consider scenarios with  $m_0 \in \{5, 10\}$ . Therefore, the true simulated size of clusters varies between 2 for the scenarios with  $n = 20$ ,  $m_0 = 10$ , and 40 for the settings with  $n = 200$ ,  $m_0 = 5$ .

### Correlation between Intercepts and Covariates

An important assumption of the mixed model is that the unit-specific intercepts are independent from the predictors  $\mathbf{x}$ . In order to break this assumption we simulate data with correlations  $\rho = \text{corr}(\beta_{i0}, x_{ij1}) \neq 0$ . For the simulation we use a sequential procedure adopted from Tutz and Oelker (2015). Consider the case of normal distributed intercepts  $\beta_{i0}$ . Here, values are first generated by  $\beta_{i0} \sim N(\mu_b, \sigma_b^2)$  and  $x_{ij1} \sim N(0, 1)$ . Afterwards  $x_{ij1}$  is transformed according to the bivariate normal distribution of  $(\beta_{i0}, x_{ij1})$  with the corresponding correlation. We consider scenarios with  $\rho \in \{0, 0.8\}$ . In the case of chi-squared distributed intercepts the joint distribution of  $(\beta_{i0}, x_{ij1})$  is not bivariate normal, but we can use the same transformation for  $x_{ij1}$  yielding the same empirical correlations.

### Evaluation Criteria

We compare the estimated coefficients to the true parameters by calculating mean squared errors (MSEs). We distinguish between the MSE of the unit-specific intercepts  $\frac{1}{n} \sum_{i=1}^n (\hat{\beta}_{i0} - \beta_{i0})^2$ , referred to as *intercepts*, and the MSE of the effects of the two covariates  $\frac{1}{2} \sum_{d=1}^2 (\hat{\beta}_d - \beta_d)^2$ , referred to as *linear term*. Concerning the mixed model, coefficients  $\hat{\beta}_{i0}$  are computed as the sum of the estimated posteriori modes and the fixed intercept  $\hat{\beta}_0$ . In addition the number of clusters determined by the different approaches are of interest. All the presented evaluations are based on 100 replications.




FIGURE 3: *MSEs of intercepts (upper panel) and the linear term (lower panel) for the settings with normal response, normal intercepts and  $\rho = 0$ .*

## 6.1 Normal Response

We start with simulation scenarios where the responses  $y_{ij}$ ,  $i = 1, \dots, n$ ,  $j = 1, \dots, n_i$  are normally distributed with  $\epsilon_{ij} \sim N(\mu_\epsilon = 0, \sigma_\epsilon^2 = 3^2)$ . Here we set  $\beta_1 = \beta_2 = 2$  as the true parameters of the two covariates. In the first case we consider cluster-specific intercepts that were generated from the fusion of parameters that follow a standard normal distribution.

It is important to mention that in the above setting the effective number of parameters for the mixed model heavily depends on the variance  $\sigma_\epsilon^2$  of the response and the variance  $\sigma_b^2$  of the random intercepts. Following Ruppert et al. (2003), the effective degrees of freedom for the random intercepts for a linear




FIGURE 4: *MSEs of intercepts (upper panel) and the linear term (lower panel) for the settings with normal response, normal intercepts and  $\rho = 0.8$ .*

random intercept model are

$$df_b = \frac{(n-1)n_i}{n_i + \frac{\sigma_\epsilon^2}{\sigma_b^2}}.$$

If  $\sigma_b^2 \rightarrow 0$  or  $\sigma_\epsilon^2 \rightarrow \infty$  the result is a model with only one intercept and if  $\sigma_b^2 \rightarrow \infty$  or  $\sigma_\epsilon^2 \rightarrow 0$  the result is a model with  $n$  intercepts, corresponding to the fixed effects model. With  $\sigma_\epsilon^2 = 9$  and  $\sigma_b^2 = 1$  one obtains the effective degrees of freedom 61.2, 46.5, 26.9 and 15.5 depending on the combination of parameters  $n$  and  $n_i$ . Therefore, one is not too close to the fixed effects model, which allows a fair comparison of the mixed model and the tree-structured model. In the second case with a skewed distribution for the unit-specific intercepts we use  $\beta_{i0} \sim \chi^2(0.5)$  with  $\sigma_b^2/2 = 0.5$  degrees of freedom. After centering of the




FIGURE 5: Selected number of clusters for the settings with normal response, normal intercepts,  $\rho = 0$  (upper panel) and  $\rho = 0.8$  (lower panel). The true number of clusters  $m_0$  is marked by dashed lines.

coefficients one obtains the same empirical values  $\mu_b = 0$  and  $\sigma_b^2 = 1$  as in the standard normal case.

Figure 3 shows the boxplots of the MSEs for the eight different settings generated by normally distributed intercepts and without correlation ( $\rho = 0$ ). As the approach by penalized likelihood estimation is computational unfeasible for a large number of units  $n$ , no results are displayed for the settings with  $n = 200$  and  $n = 100$ . It is seen from the lower panel that all the approaches nearly show the same performance for the linear term. However, distinct differences are seen for the intercepts (upper panel). Although there are clusters of units the mixed model shows good performance for all settings. The fixed effects model performs poorly, especially for the settings with  $n_i = 4$ , the finite mixture model performs poorly for the settings with  $n = 40$  and  $n = 20$ . The estimates of the






FIGURE 6: *MSEs of the linear term for the settings with normal response, chi-squared intercepts and  $\rho = 0.8$ .*

tree-structured model show better performance than the fixed effects model for smaller values of  $n_i$  and comparable performance for larger values. The performance is the same as for the penalty approach if estimates exist. The picture changes in the settings with correlation  $\rho = 0.8$  between covariate  $x_1$  and the unit-specific intercepts (Figure 4). For the linear term (lower panel) the performance of the mixed model and the finite mixture model suffers strongly. In contrast, the estimation accuracy of the fixed effects model, the tree-structured model and the penalized likelihood approach is not affected by the correlation. In particular, the tree-structured model outperforms the penalty approach in all the settings in which the penalty approach works. The results for the intercepts (upper panel) do not change that much but the mixed model and the finite mixture model is now competitive only for small values of  $n_i$ .

Boxplots of the selected number of clusters are given in Figure 5 for  $\rho = 0$  (upper panel) and  $\rho = 0.8$  (lower panel). Since the fixed effects model and the mixed model do not build clusters of units, the given number of clusters for the two approaches is equal to the number of units. There are only minor differences between the settings with and without correlation. The number of clusters identified by the tree-structured model is very close to the true number for the settings with five clusters ( $m_0 = 5$ ) but the true number of clusters is slightly underestimated in the settings with ten clusters. In contrast, the penalty approach selects a distinctly higher number of clusters with a strong variation. The finite mixture model consistently selects only too small number of clusters. On average only about two clusters are selected by AIC as well as by BIC.

The evaluations of the same settings with cluster-specific intercepts that were generated by a chi-squared distribution yield very similar results. In particular the performance of the mixed model seems not to be affected too strongly by the




FIGURE 7: *MSEs of intercepts (upper panel) and the linear term (lower panel) for the settings with binary response, chi-squared intercepts and  $\rho = 0.8$ .*

skewed distribution of the random intercepts. For illustration Figure 6 shows the MSEs of the linear term for the settings with  $\rho = 0.8$ . See the appendix for an overview of all results.

## 6.2 Binary Response

In the following we briefly consider discrete response variables  $y_{ij} \sim B(1, \pi_{ij})$ , where  $\pi_{ij} = \exp(\eta_{ij}) / (1 + \exp(\eta_{ij}))$ . The structure of the simulated data sets remains the same but some modifications to the specifications in Section 6.1 are necessary. The parameters of the linear term are set to  $\beta_1 = \beta_2 = 0.1$ . For the cluster-specific intercepts we chose  $\beta_{i0} \sim N(-0.8, 2^2)$  or as skew counterpart  $\beta_{i0} \sim \chi^2(2)$ , centered such that  $\mu_b = -0.8$ . Since  $n_i = 4$  is a relatively small size when modelling binary responses, we do not consider the corresponding settings.




FIGURE 8: Selected number of clusters for the settings with binary response, chi-squared intercepts and  $\rho = 0.8$ . The true number of clusters  $m_0$  is marked by dashed lines.

Furthermore, we omit the estimates of the fixed effects model because they are very unstable and often do not exist in this case. Accordingly, the order of measurement units used in the algorithm of the tree-structured model is not based on the estimates of the unrestricted model but by adding a small ridge penalty.

In contrast to the settings with normal response, the results for the binary response as a whole seem to be more affected by a skewed distribution of the intercepts. In the following we will focus on the settings with chi-squared distributed intercepts and  $\rho = 0.8$ , and refer to the appendix for further results. Figure 7 shows the MSEs of the unit-specific intercepts (upper panel) and the linear term (lower panel). Again the mixed model and the finite mixture model perform poorly with regard to the linear term, but there are only minor differences for  $n = 20$ . Regarding the intercepts the average results are comparable for all the approaches. It is noticeable that one observes huge outliers for the finite mixture models, especially with model selection by AIC. It is most conspicuous for the settings with  $n = 20$ , where the boxplots have been truncated.

The corresponding boxplots of the selected number of clusters are given in Figure 8. Here the tree-structured model only detects very few clusters (for  $m_0 = 5$  and  $m_0 = 10$ ) and is almost as restrictive as the finite mixture model. As before the penalty approach selects a higher number of clusters and has a stronger variation but the selected number of clusters is closer to the true number of clusters.

TABLE 3: *Description and distribution of the covariates used for the analysis of the Guatemala survey.*

Variable	Description	Categories	Frequency
<b>ethn</b>	Mother’s ethnicity	non-indigenous (Ladino)	612
		indigenous, not speaking Spanish	286
		indigenous, speaking Spanish	313
<b>momEd</b>	Mother’s level of education	not finished primary	571
		finished primary	607
		finished secondary	33
<b>husEd</b>	Husband’s level of education	not finished primary	430
		finished primary	598
		finished secondary	67
		unknown	116
<b>husEmpl</b>	Husband’s employment status	unskilled	45
		professional	120
		agricultural, self-employed	420
		agricultural, employee	407
		skilled service	219
<b>telev</b>	Frequency of TV usage	never	1034
		not daily	52
		daily	125
<b>momAge</b>	Mother 25 years or older	no	583
		yes	628
<b>toilet</b>	Modern toilet in house	no	112
		yes	1099

## 7 A Further Application

As second application we consider data derived from the National Survey of Maternal and Child Health in Guatemala in 1987. The data is available from the R-package `m1mRev` (Bates et al., 2014) and was also analysed by Rodriguez and Goldman (2001). The data contains observations of children that were born in the 5-year period before the survey. In our analysis we include 1211 children living in 45 communities. One observes a minimal number of 20, a maximal number of 50 and an average number of 26.9 pregnancies per community. The response  $y_{ij}$  is a binary outcome with  $y_{ij} = 0$  for traditional prenatal care and  $y_{ij} = 1$  for modern prenatal care, for example by doctors or nurses. The response is modelled by a logistic regression model  $\text{logit}(P(y_{ij} = 1)) = \eta_{ij}$ . The heterogeneity of communities is modelled by the alternative approaches considered here. In total there are 733 pregnancies with traditional and 478 observed pregnancies with modern prenatal care. The two binary and five categorical explanatory variables that characterize the children’s mothers and their families are given in Table 3.

An overview of the estimated coefficients when using a generalized mixed model (GMM), tree-structured clustering (TSC) and a finite mixture model (FIN)

TABLE 4: *Estimation results of the Guatemala survey using the generalized mixed model, tree-structured clustering and the finite mixture model.*

Predictor	GMM		TSC		FIN			
	Coefficient	95%-CI	Coefficient	95%-CI	Coefficient	95%-CI		
<b>ethn</b>								
not spanish	-1.370	[-2.101,-0.774]	-1.090	[-2.469,-0.387]	-0.995	[-2.280,-0.556]		
spanish	-0.720	[-1.235,-0.244]	-0.434	[-1.425, 0.005]	-0.335	[-1.338, 0.011]		
<b>momEd</b>								
primary	0.645	[ 0.331, 1.048]	0.673	[ 0.298, 1.122]	0.646	[ 0.317, 1.078]		
secondary	1.385	[ 0.303, 2.955]	1.405	[ 0.268, 3.046]	1.735	[ 0.364, 2.944]		
<b>husEd</b>								
primary	0.785	[ 0.445, 1.236]	0.817	[ 0.437, 1.303]	0.843	[ 0.444, 1.301]		
secondary	0.194	[-0.809, 1.186]	0.049	[-0.922, 1.286]	0.291	[-0.846, 1.311]		
unknown	0.398	[-0.113, 0.951]	0.520	[-0.101, 1.006]	0.428	[-0.106, 0.962]		
<b>husEmpl</b>								
professional	-0.210	[-1.150, 0.670]	-0.095	[-1.301, 0.820]	-0.408	[-1.336, 0.667]		
agricult, self	-0.119	[-0.975, 0.721]	-0.065	[-1.044, 0.798]	-0.266	[-1.065, 0.716]		
agricult, empl	-0.158	[-1.024, 0.656]	-0.100	[-1.092, 0.750]	-0.238	[-1.103, 0.723]		
skilled	-0.199	[-1.079, 0.606]	-0.125	[-1.123, 0.661]	-0.300	[-1.134, 0.607]		
<b>telev</b>								
not daily	0.355	[-0.497, 1.292]	0.226	[-0.601, 1.286]	0.241	[-0.548, 1.283]		
daily	0.867	[ 0.312, 1.560]	0.928	[ 0.290, 1.570]	0.735	[ 0.307, 1.524]		
momAge	0.099	[-0.208, 0.403]	0.061	[-0.241, 0.411]	0.061	[-0.219, 0.401]		
toilet	-0.869	[-1.833,-0.055]	-1.008	[-1.875, 0.092]	-0.839	[-1.808,-0.154]		
$\beta_0$	-0.011	[-1.223, 1.166]	—	—	—	—		
$\sigma_{\text{rand}}^2$	1.250	[ 1.233, 2.416]	—	—	—	—		
<b>Community-specific intercept</b>			<b>TSC</b>		<b>FIN</b>			
			Cluster	Size	Coefficient	Cluster	Size	Coefficient
$\beta_{i0}$			1	15	-1.286	1	33	-0.696
			2	17	-0.214	2	12	1.465
			3	13	1.448			

is given in Table 4. The 95% confidence intervals were obtained by 2000 bootstrap samples. It can be seen from the results that the age of the mother at the time of the survey as well as the employment status of the husband do not have a significant effect on the form of prenatal care. The educational level of the mother as well as of the husband, however, have a strong impact. For births where the mother at least finished primary or the husband finished primary modern prenatal care was provided more likely compared to births of parents without any graduation. Indigenous mothers (speaking and not speaking Spanish) are also more likely to use traditional prenatal care than non-indigenous mothers. The existence of a modern toilet in the household does not favour the use of modern prenatal care, whereas it is preferred by families using the television regularly.

A comparison of the estimates obtained by the three methods does not show strong distinctions and no clear tendency. Differences occur for variable ethnicity




FIGURE 9: Comparison of the estimated distribution of the mixed model and the community-specific intercepts of tree-structured clustering (Guatemala survey).

(first rows in Table 4), for which the two estimates of the mixed model are larger than for TSC and FIN and for mothers that finished secondary (fourth row) for which the estimate of the finite mixture model is larger than for TSC and GMM.

The estimated community-specific intercepts obtained by tree-structured clustering and the finite mixture model are given in the lower panel of Table 4. Using the tree-structured model results in three clusters of communities that differ in terms of their probability to use modern prenatal care. The finite mixture identifies only two clusters. We prefer to use model selection by BIC as it showed more stable estimates in the simulations with binary response. The detected partitions and the high variance obtained by the mixed model indicate that heterogeneity of communities is definitely present. Nevertheless, only a few clusters of communities have to be distinguished. There is a strong similarity between the third cluster of the tree-structured model ( $\beta_{i0}^{(3)} = 1.448$ ) and the second cluster of the finite mixture model ( $\beta_{i0}^{(2)} = 1.465$ ) but as a whole the partition of tree-structured clustering seems to be more adequate. In Figure 9 the estimated distribution of the community-specific intercepts of the tree-structured model and the estimated normal distribution of the mixed model are graphically illustrated.

## 8 Concluding Remarks

For simplicity we focussed on the most important case of clustered intercepts. However, the general fixed effects model (2) allows for more than one parameter to be unit-specific. It is straightforward to extend the tree-structured model to include a covariate vector  $\mathbf{z}_{ij} = (1, z_{ij1}, \dots, z_{ijq})$ . Then one obtains a model with

predictor

$$\eta_{ij} = \mathbf{x}_{ij}^\top \boldsymbol{\beta} + \sum_{r=0}^q \sum_{k=1}^{m_r} z_{ijr} \beta_{ir}^{(k)} I(i \in S_{kr}), \quad (6)$$

where  $S_{1r}, \dots, S_{m_r r}$  is a partition of the units  $\{1, \dots, n\}$  with respect to the  $r$ -th component of  $\mathbf{z}_{ij}$  and  $\beta_{ir}^{(1)}, \dots, \beta_{ir}^{(m_r)}$  are the corresponding parameters of each cluster. Due to individual splits, the number and form of clusters do not have to be the same for the different components of  $\mathbf{z}_{ij}$ . The fitting procedure given in Section 4 can easily be adapted to this general model. In each iteration one simply has to determine the best split among all covariates and all corresponding splits simultaneously. In a first step the order of the units  $\{1, \dots, n\}$  with respect to single covariates has to be defined. It is not assumed that the order is the same for each of the covariates. The result is one tree for each covariate that represents a partition of units.

The proposed tree structured clustering competes well with the competitors. In particular, it performs better than the finite mixture approach and has the advantage that the number of units is not restricted as in the penalty approach. The applications were chosen to illustrate the potential of the method to find clusters that share the same effect on the covariates. The potential of the method to yield better estimates when the heterogeneity and explanatory variables are correlated is demonstrated in the simulations. The presented results were obtained by an R program that is available from the authors.

## References

- Agresti, A., B. Caffo, and P. Ohman-Strickland (2004). Examples in which misspecification of a random effects distribution reduces efficiency, and possible remedies. *Computational Statistics and Data Analysis* 47, 639–653.
- Aitkin, M. (1999). A general maximum likelihood analysis of variance components in generalized linear models. *Biometrics* 55, 117–128.
- Bates, D., M. Mächler, B. Bolker, and S. Walker (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software* 67(1), 1–48.
- Bates, D., M. Maechler, and B. Bolker (2014). *mlmRev: Examples from Multilevel Modelling Software Review*. R package version 1.0-6.
- Bondell, H. D. and B. J. Reich (2008). Simultaneous regression shrinkage, variable selection and clustering of predictors with oscar. *Biometrics* 64, 115–123.
- Breiman, L., J. H. Friedman, R. A. Olshen, and J. C. Stone (1984). *Classification and Regression Trees*. Monterey, CA: Wadsworth.

- Bush, C. A. and S. N. MacEachern (1996). A semiparametric bayesian model for randomised block designs. *Biometrika* 83(2), 275–285.
- Chen, J. and M. Davidian (2002). A monte carlo EM algorithm for generalized linear models with flexible random effects distribution. *Biostatistics* 3, 347–360.
- Claeskens, G. and J. D. Hart (2009). Goodness-of-fit tests in mixed models. *TEST* 18, 213–239.
- De Boeck, P. and M. Wilson (2004). *Explanatory item response models: A generalized linear and nonlinear approach*. Springer Verlag.
- Ferguson, T. S. (1973). A Bayesian analysis of some nonparametric problems. *The Annals of Statistics* 1, 209–230.
- Fisher, W. D. (1958). On grouping for maximum homogeneity. *Journal of the American Statistical Association* 53(284), 789–798.
- Follmann, D. and D. Lambert (1989). Generalizing logistic regression by non-parametric mixing. *Journal of the American Statistical Association* 84, 295–300.
- Gertheiss, J. and G. Tutz (2010). Sparse modeling of categorical explanatory variables. *Annals of Applied Statistics* 4, 2150–2180.
- Grilli, L. and C. Rampichini (2011). The role of sample cluster means in multi-level models: A view on endogeneity and measurement error issues. *Methodology: European Journal of Research Methods for the Behavioural and Social Sciences* 7(4), 121–133.
- Grün, B. and F. Leisch (2007). Fitting finite mixtures of generalized linear regressions in R. *Computational Statistics & Data Analysis* 51(11), 5247–5252.
- Grün, B. and F. Leisch (2008a). FlexMix version 2: finite mixtures with concomitant variables and varying and constant parameters. *Journal of Statistical Software*, 28(4), 1–35.
- Grün, B. and F. Leisch (2008b). Identifiability of finite mixtures of multinomial logit models with varying and fixed effects. *Journal of Classification* 25(2), 225–247.
- Hajjem, A., F. Bellavance, and D. Larocque (2011). Mixed effects regression trees for clustered data. *Statistics and Probability Letters* 81, 451–459.
- Hastie, T., R. Tibshirani, and J. H. Friedman (2009). *The Elements of Statistical Learning (Second Edition)*. New York: Springer-Verlag.



- Heagerty, P. and B. F. Kurland (2001). Misspecified maximum likelihood estimates and generalised linear mixed models. *Biometrika* 88, 973–985.
- Heinzel, F. and G. Tutz (2013). Clustering in linear mixed models with approximate dirichlet process mixtures using em algorithm. *Statistical Modelling* 13, 41–67.
- Heinzel, F. and G. Tutz (2014). Clustering in linear-mixed models with a group fused lasso penalty. *Biometrical Journal* 56(1), 44–68.
- Heinzel, F. and G. Tutz (2015). Additive mixed models with approximate dirichlet process mixtures: the em approach. *Statistics and Computing*, to appear.
- Hjort, N. L., C. Holmes, P. Müller, and S. G. Walker (2010). *Bayesian nonparametrics*, Volume 28. Cambridge University Press.
- Hothorn, T., K. Hornik, and A. Zeileis (2006). Unbiased recursive partitioning: A conditional inference framework. *Journal of Computational and Graphical Statistics* 15, 651–674.
- Huang, X. (2009). Diagnosis of random-effect model misspecification in generalized linear mixed models for binary response. *Biometrics* 65, 361–368.
- Khalili, A. and J. Chen (2007). Variable selection in finite mixture of regression models. *Journal of the American Statistical Association* 102, 1025–1038.
- Litière, S., A. Alonso, and G. Molenberghs (2007). Type I and Type II Error Under Random Effects Misspecification in Generalized Linear Mixed Models. *Biometrics* 63, 1038–1044.
- Lombardía, M. J. and S. Sperlich (2012). A new class of semi-mixed effects models and its application in small area estimation. *Computational Statistics & Data Analysis* 56(10), 2903–2917.
- Magder, L. and S. Zeger (1996). A smooth nonparametric estimate of a mixing distribution using mixtures of gaussians. *Journal of the American Statistical Association* 91, 1141–1151.
- McCulloch, C. and S. Searle (2001). *Generalized, Linear, and Mixed Models*. New York: Wiley.
- Molenberghs, G. and G. Verbeke (2005). *Models for Discrete Longitudinal Data*. New York: Springer-Verlag.
- Morgan, J. N. and J. A. Sonquist (1963). Problems in the analysis of survey data, and a proposal. *Journal of the American Statistical Association* 58, 415–435.

- Müller, P. and G. L. Rosner (1997). A bayesian population model with hierarchical mixture priors applied to blood count data. *Journal of the American Statistical Association* 92(440), 1279–1292.
- Oelker, M.-R. (2015). *gvcm.cat: Regularized Categorical Effects/Categorical Effect Modifiers/Continuous/Smooth Effects in GLMs*. R package version 1.9.
- Quinlan, J. R. (1986). Induction of decision trees. *Machine Learning* 1, 81–106.
- Quinlan, J. R. (1993). *Programs for Machine Learning*. San Francisco: Morgan Kaufmann Publisher Inc.
- Ripley, B. D. (1996). *Pattern Recognition and Neural Networks*. Cambridge: Cambridge University Press.
- Rodriguez, G. and N. Goldman (2001). Improved estimation procedures for multilevel models with binary response: A case-study. *Journal of the Royal Statistical Society. Series A (Statistics in Society)* 164, 339–355.
- Ruppert, D., M. P. Wand, and R. J. Carroll (2003). *Semiparametric Regression*. Cambridge: Cambridge University Press.
- Sela, R. J. and J. S. Simonoff (2012). Re-em trees: a data mining approach for longitudinal and clustered data. *Machine learning* 86(2), 169–207.
- Sethuraman, J. (1994). A constructive definition of Dirichlet priors. *Statistica Sinica* 4, 639–650.
- Städler, N., P. Bühlmann, and S. van de Geer (2010). L1-penalization for mixture regression models. *Test* 19, 209–256.
- Strobl, C., J. Malley, and G. Tutz (2009). An Introduction to Recursive Partitioning: Rationale, Application and Characteristics of Classification and Regression Trees, Bagging and Random Forests. *Psychological Methods* 14, 323–348.
- Tutz, G. and Gertheiss (2014). Rating scales as predictors – the old question of scale level and some answers. *Psychometrika* 79(3), 357–376.
- Tutz, G. and M. Oelker (2015). Modeling clustered heterogeneity: Fixed effects, random effects and mixtures. *International Statistical Review*, to appear.
- Verbeke and G. Molenberghs (2000). *Linear Mixed Models for longitudinal data*. New York: Springer-Verlag.
- Zhang, H. and B. Singer (1999). *Recursive Partitioning in the Health Sciences*. New York: Springer-Verlag.

## Appendix: Tabular Display of Simulation Results

In the following we give the results of all settings of the simulations described in Section 6. Each table contains the MSEs of the unit-specific intercepts, the MSEs of the linear term and the selected number of clusters as the average of 100 replications, respectively.

TABLE 5: Average results for the settings with normal response, normal distributed intercepts and  $\rho = 0$ .

		MSE - intercepts		MSE - linear term		Number of Clusters	
		$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$
$n = 200$ $n_i = 4$	GFM	2.26	2.26	0.04	0.04	200.00	200.00
	GMM	0.68	0.71	0.03	0.03	200.00	200.00
	TSC	1.56	1.57	0.04	0.04	4.96	5.02
	PEL						
	FINA	1.05	1.10	0.03	0.03	1.89	1.91
	FINB	0.99	1.06	0.03	0.03	1.31	1.36
$n = 100$ $n_i = 8$	GFM	1.14	1.14	0.03	0.03	100.00	100.00
	GMM	0.54	0.56	0.03	0.03	100.00	100.00
	TSC	0.97	0.99	0.03	0.03	5.28	5.38
	PEL						
	FINA	0.82	0.87	0.03	0.03	2.04	2.10
	FINB	0.86	0.91	0.03	0.03	1.67	1.72
$n = 40$ $n_i = 20$	GFM	0.45	0.45	0.03	0.03	40.00	40.00
	GMM	0.31	0.32	0.03	0.03	40.00	40.00
	TSC	0.44	0.46	0.03	0.03	5.82	6.00
	PEL	0.37	0.38	0.03	0.03	15.00	15.06
	FINA	0.53	0.55	0.03	0.03	2.27	2.44
	FINB	0.57	0.61	0.03	0.03	1.86	1.98
$n = 20$ $n_i = 40$	GFM	0.22	0.22	0.03	0.03	20.00	20.00
	GMM	0.19	0.19	0.03	0.03	20.00	20.00
	TSC	0.23	0.24	0.03	0.03	5.76	6.00
	PEL	0.21	0.21	0.03	0.03	9.95	9.99
	FINA	0.32	0.34	0.03	0.03	2.45	2.66
	FINB	0.39	0.43	0.03	0.03	1.96	2.06

TABLE 6: Average results for the settings with normal response, normal distributed intercepts and  $\rho = 0.8$ .

		MSE - intercepts		MSE - linear term		Number of Clusters	
		$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$
$n = 200$ $n_i = 4$	GFM	2.28	2.28	0.05	0.05	200.00	200.00
	GMM	0.88	0.95	0.29	0.32	200.00	200.00
	TSC	1.51	1.53	0.08	0.08	4.86	4.95
	PEL						
	FINA	0.95	1.01	0.30	0.34	1.14	1.10
	FINB	0.92	0.98	0.30	0.34	1.00	1.00
$n = 100$ $n_i = 8$	GFM	1.16	1.16	0.04	0.04	100.00	100.00
	GMM	0.84	0.91	0.25	0.29	100.00	100.00
	TSC	0.96	0.98	0.05	0.06	5.18	5.20
	PEL						
	FINA	0.94	1.00	0.26	0.30	1.25	1.25
	FINB	0.92	0.99	0.28	0.31	1.00	1.02
$n = 40$ $n_i = 20$	GFM	0.48	0.48	0.04	0.04	40.00	40.00
	GMM	0.67	0.76	0.19	0.23	40.00	40.00
	TSC	0.48	0.50	0.04	0.04	5.82	5.93
	PEL	0.39	0.40	0.05	0.06	14.17	14.14
	FINA	0.82	0.89	0.21	0.25	1.53	1.51
	FINB	0.90	0.99	0.26	0.31	1.11	1.02
$n = 20$ $n_i = 40$	GFM	0.25	0.25	0.04	0.04	20.00	20.00
	GMM	0.46	0.54	0.14	0.17	20.00	20.00
	TSC	0.27	0.29	0.05	0.05	5.74	5.97
	PEL	0.25	0.26	0.06	0.06	9.59	9.62
	FINA	0.62	0.71	0.17	0.21	1.80	1.73
	FINB	0.81	0.91	0.25	0.29	1.22	1.16

TABLE 7: Average results for the settings with normal response, chi-squared distributed intercepts and  $\rho = 0$ .

		MSE - intercepts		MSE - linear term		Number of Clusters	
		$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$
$n = 200$ $n_i = 4$	GFM	2.27	2.27	0.04	0.04	200.00	200.00
	GMM	0.50	0.59	0.03	0.03	200.00	200.00
	TSC	1.52	1.59	0.04	0.04	4.60	4.88
	PEL						
	FINA	0.69	0.77	0.03	0.03	1.49	1.80
	FINB	0.63	0.76	0.03	0.03	1.14	1.32
$n = 100$ $n_i = 8$	GFM	1.10	1.10	0.03	0.03	100.00	100.00
	GMM	0.41	0.47	0.02	0.02	100.00	100.00
	TSC	0.91	0.95	0.02	0.03	4.77	5.14
	PEL						
	FINA	0.54	0.50	0.02	0.02	1.72	1.90
	FINB	0.55	0.55	0.02	0.02	1.28	1.53
$n = 40$ $n_i = 20$	GFM	0.45	0.45	0.03	0.03	40.00	40.00
	GMM	0.26	0.28	0.03	0.03	40.00	40.00
	TSC	0.42	0.42	0.03	0.03	4.95	5.15
	PEL	0.30	0.29	0.03	0.03	13.17	13.27
	FINA	0.26	0.28	0.03	0.03	1.85	2.00
	FINB	0.28	0.29	0.03	0.03	1.60	1.68
$n = 20$ $n_i = 40$	GFM	0.23	0.23	0.03	0.03	20.00	20.00
	GMM	0.16	0.16	0.03	0.03	20.00	20.00
	TSC	0.22	0.23	0.03	0.03	4.69	4.92
	PEL	0.15	0.15	0.03	0.03	7.87	8.23
	FINA	0.14	0.18	0.03	0.03	1.88	2.10
	FINB	0.15	0.20	0.03	0.03	1.67	1.81

TABLE 8: Average results for the settings with normal response, chi-squared distributed intercepts and  $\rho = 0.8$ .

		MSE - intercepts		MSE - linear term		Number of Clusters	
		$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$
$n = 200$ $n_i = 4$	GFM	2.30	2.30	0.05	0.05	200.00	200.00
	GMM	0.56	0.73	0.13	0.20	200.00	200.00
	TSC	1.51	1.55	0.05	0.06	4.62	4.85
	PEL						
	FINA	0.64	0.82	0.13	0.20	1.18	1.24
	FINB	0.60	0.77	0.14	0.21	1.01	1.01
$n = 100$ $n_i = 8$	GFM	1.12	1.12	0.04	0.04	100.00	100.00
	GMM	0.53	0.70	0.12	0.18	100.00	100.00
	TSC	0.92	0.95	0.04	0.05	4.72	4.99
	PEL						
	FINA	0.61	0.74	0.12	0.19	1.32	1.33
	FINB	0.60	0.77	0.13	0.20	1.01	1.03
$n = 40$ $n_i = 20$	GFM	0.48	0.48	0.04	0.04	40.00	40.00
	GMM	0.44	0.62	0.11	0.17	40.00	40.00
	TSC	0.45	0.46	0.05	0.05	4.82	5.12
	PEL	0.33	0.32	0.05	0.05	12.85	13.07
	FINA	0.45	0.56	0.11	0.15	1.62	1.56
	FINB	0.51	0.70	0.13	0.20	1.26	1.20
$n = 20$ $n_i = 40$	GFM	0.26	0.26	0.04	0.04	20.00	20.00
	GMM	0.30	0.44	0.08	0.13	20.00	20.00
	TSC	0.26	0.26	0.04	0.04	4.69	4.92
	PEL	0.20	0.19	0.04	0.04	8.04	8.21
	FINA	0.31	0.44	0.08	0.11	1.74	1.77
	FINB	0.38	0.62	0.11	0.17	1.39	1.34

TABLE 9: Average results for the settings with binary response, normal distributed intercepts and  $\rho = 0$ .

		MSE - intercepts		MSE - linear term		Number of Clusters	
		$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$
$n = 100$ $n_i = 8$	GFM						
	GMM	0.74	0.88	0.03	0.03	100.00	100.00
	TSC	1.06	1.29	0.02	0.02	2.96	2.98
	PEL						
	FINA	2.88	2.39	0.03	0.03	2.98	3.03
	FINB	2.11	1.66	0.03	0.02	2.64	2.63
$n = 40$ $n_i = 20$	GFM						
	GMM	0.48	0.56	0.02	0.02	40.00	40.00
	TSC	0.70	0.87	0.02	0.02	3.32	3.50
	PEL	1.23	1.20	0.02	0.02	10.78	14.28
	FINA	10.70	5.26	0.02	0.02	3.49	3.52
	FINB	9.10	3.93	0.02	0.02	3.00	2.97
$n = 20$ $n_i = 40$	GFM						
	GMM	0.71	0.62	0.03	0.03	20.00	20.00
	TSC	2.40	2.18	0.03	0.03	3.44	3.84
	PEL	1.44	1.15	0.03	0.03	5.70	9.15
	FINA	19.94	12.58	0.03	0.03	3.57	3.84
	FINB	15.58	8.71	0.03	0.03	3.12	3.21

TABLE 10: Average results for the settings with binary response, normal distributed intercepts and  $\rho = 0.8$ .

		MSE - intercepts		MSE - linear term		Number of Clusters	
		$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$
$n = 100$ $n_i = 8$	GFM						
	GMM	2.13	2.55	0.48	0.54	100.00	100.00
	TSC	1.59	1.93	0.25	0.29	2.46	2.38
	PEL						
	FINA	3.43	3.89	0.46	0.51	2.35	2.26
	FINB	2.60	2.95	0.50	0.56	1.93	1.85
$n = 40$ $n_i = 20$	GFM						
	GMM	0.92	1.12	0.14	0.15	40.00	40.00
	TSC	0.98	1.16	0.11	0.12	3.04	3.13
	PEL	1.32	1.26	0.05	0.05	10.42	13.19
	FINA	12.51	8.08	0.11	0.14	2.96	2.91
	FINB	8.06	5.39	0.16	0.22	2.45	2.29
$n = 20$ $n_i = 40$	GFM						
	GMM	0.87	0.84	0.07	0.08	20.00	20.00
	TSC	2.67	1.87	0.06	0.07	3.21	3.53
	PEL	1.74	1.26	0.05	0.05	5.61	8.91
	FINA	22.57	13.19	0.06	0.09	3.34	3.41
	FINB	15.15	7.81	0.09	0.14	2.81	2.64

TABLE 11: Average results for the settings with binary response, chi-squared distributed intercepts and  $\rho = 0$ .

		MSE - intercepts		MSE - linear term		Number of Clusters	
		$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$
$n = 100$ $n_i = 8$	GFM						
	GMM	0.68	0.92	0.02	0.02	100.00	100.00
	TSC	0.91	1.39	0.02	0.02	2.79	2.85
	PEL						
	FINA	1.72	2.30	0.02	0.02	2.74	2.90
	FINB	1.42	1.70	0.02	0.02	2.40	2.51
$n = 40$ $n_i = 20$	GFM						
	GMM	0.48	0.61	0.02	0.02	40.00	40.00
	TSC	0.59	0.82	0.02	0.02	3.01	3.37
	PEL	1.60	1.43	0.02	0.02	9.83	12.35
	FINA	5.61	6.94	0.02	0.02	3.04	3.34
	FINB	4.66	4.47	0.02	0.02	2.74	2.91
$n = 20$ $n_i = 40$	GFM						
	GMM	1.61	2.00	0.03	0.03	20.00	20.00
	TSC	2.81	2.96	0.03	0.03	2.94	3.56
	PEL	1.93	2.04	0.03	0.02	5.75	8.04
	FINA	21.18	19.61	0.04	0.03	3.04	3.61
	FINB	19.95	16.30	0.03	0.03	2.77	3.06

TABLE 12: Average results for the settings with binary response, chi-squared distributed intercepts and  $\rho = 0.8$ .

		MSE - intercepts		MSE - linear term		Number of Clusters	
		$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$	$m_0 = 5$	$m_0 = 10$
$n = 100$ $n_i = 8$	GFM						
	GMM	1.55	2.30	0.41	0.50	100.00	100.00
	TSC	1.28	1.91	0.22	0.30	2.50	2.30
	PEL						
	FINA	4.85	4.37	0.33	0.46	2.48	2.24
	FINB	2.68	2.45	0.37	0.50	2.05	1.86
$n = 40$ $n_i = 20$	GFM						
	GMM	0.72	1.15	0.13	0.16	40.00	40.00
	TSC	0.75	1.15	0.09	0.12	2.80	3.01
	PEL	1.72	1.53	0.04	0.05	9.38	11.89
	FINA	9.57	6.76	0.09	0.14	2.85	2.84
	FINB	7.06	4.68	0.11	0.17	2.50	2.47
$n = 20$ $n_i = 40$	GFM						
	GMM	1.66	2.26	0.07	0.07	20.00	20.00
	TSC	3.08	2.92	0.06	0.07	2.81	3.33
	PEL	2.26	2.34	0.05	0.05	5.59	7.72
	FINA	21.87	21.18	0.06	0.08	2.90	3.25
	FINB	21.79	16.13	0.07	0.11	2.68	2.68