

# SEMI-PARAMETRIC EFFICIENCY BOUNDS FOR HIGH-DIMENSIONAL MODELS

BY JANA JANKOVÁ AND SARA VAN DE GEER

*ETH Zürich*

Asymptotic lower bounds for estimation play a fundamental role in assessing the quality of statistical procedures. In this paper we propose a framework for obtaining semi-parametric efficiency bounds for sparse high-dimensional models, where the dimension of the parameter is larger than the sample size. We adopt a semi-parametric point of view: we concentrate on one dimensional functions of a high-dimensional parameter. We follow two different approaches to reach the lower bounds: asymptotic Cramér-Rao bounds and Le Cam's type of analysis. Both these approaches allow us to define a class of asymptotically unbiased or "regular" estimators for which a lower bound is derived. Consequently, we show that certain estimators obtained by de-sparsifying (or de-biasing) an  $\ell_1$ -penalized M-estimator are asymptotically unbiased and achieve the lower bound on the variance: thus in this sense they are asymptotically efficient. The paper discusses in detail the linear regression model and the Gaussian graphical model.

**1. Introduction.** Following the development of numerous methods for high-dimensional estimation, more recently the need for statistical inference has emerged. A number of papers have since studied the problem and proposed constructions of estimators which are asymptotically normally distributed and hence lead to inference. These results naturally give rise to the question of their optimality. This motivates us to study the question whether we can establish asymptotic efficiency bounds in high-dimensional models and whether we can construct an estimator achieving these bounds.

To introduce the setting, suppose that we observe a sample  $X^{(1)}, \dots, X^{(n)}$  which is distributed according to a probability distribution  $P_\beta$  that depends on an unknown high-dimensional parameter  $\beta \in \mathcal{B} \subset \mathbb{R}^p$ . The dimension  $p$  of the parameter can be much larger than the sample size  $n$ . A major structural assumption we consider in this paper is sparsity in the high-dimensional parameter. In these sparse high-dimensional settings, a common approach to estimation is based on regularized M-estimators, where the regularization is in terms of the  $\ell_1$ -penalty. This approach has been studied extensively, and

---

*MSC 2010 subject classifications:* Primary 62J07; secondary 62F12

*Keywords and phrases:* Asymptotic efficiency, high-dimensional, sparsity, Lasso, linear regression, graphical models, Cramér-Rao bound, Le Cam's Lemma

under several settings, it produces near-oracle estimators of  $\beta$  under certain sparsity conditions (and some further conditions). However, the oracle properties of the regularized estimators come at a price: the regularization introduces bias by shrinking the estimated coefficients towards zero. Hence, the regularized approach does not easily yield estimators which are asymptotically normally distributed. This makes it difficult to establish results for statistical inference.

Several streams of work have emerged that studied “post-regularization inference”, which focused on construction of methodology for inference, with some preliminary use of regularized estimators. This was mostly considered for estimation of low-dimensional parameters of the high-dimensional vector. One stream of work concentrates on “de-sparsifying” or “de-biasing” procedures, which were studied for the linear model (Zhang and Zhang (2014), van de Geer et al. (2014), Javanmard and Montanari (2014a), Javanmard and Montanari (2014a), Javanmard and Montanari (2014b), Javanmard and Montanari (2015)), for generalized linear models (van de Geer et al. (2014)) and some special cases of non-linear models, such as undirected graphical models (Janková and van de Geer (2014), Janková and van de Geer (2016a)). This approach uses the  $\ell_1$ -regularized M-estimator as an initial estimator and implements a bias correction step which may be interpreted as one iteration using the Newton-Raphson method. Another stream of work studies the use of orthogonalizing conditions to define a new post-regularization estimator; this approach was considered for general models under high-level conditions in Chernozhukov, Hansen and Spindler (2015). Further examples of high-dimensional inference include the works Ren et al. (2015), Gao, Ma and Zhou (2014) or data splitting methods (Meinshausen and Yu (2009)). The work in essence shows an important result: an asymptotically normal estimator for low-dimensional parameters can be constructed in several of the common models.

Further key questions that were studied concern optimality properties of these de-sparsified estimators. In particular, what are lower bounds on the rate of convergence in the supremum norm? These questions have been investigated for the linear regression with random design (Cai and Guo (2015)) and for Gaussian graphical models (Ren et al. (2015)) and other special cases of non-linear models (Gao, Ma and Zhou (2014)). The results in these settings reveal several important findings, which we discuss for the linear regression and graphical models. The minimax rates for estimation of single elements (of the vector of regression coefficients or the precision

matrix) are shown to satisfy

$$(1) \quad \inf_T \sup_{\beta \in \mathcal{B}} \mathbb{E}_\beta |T(X^{(1)}, \dots, X^{(n)}) - \beta_i| \geq C(1/\sqrt{n} + s \log p/n),$$

for some constant  $C > 0$ , where  $\beta_i \in \mathbb{R}$  is a single regression coefficient or a single entry in a precision matrix and the *unknown* sparsity  $s$  is the number of non-zero entries in the regression vector or, in the case of Gaussian graphical models, in rows of a precision matrix. The infimum in (1) is taken over all estimators  $T$ . The statement (1) further requires some mild regularity conditions (see Cai and Guo (2015), Ren et al. (2015)). Naturally, (1) implies that the parametric rate is optimal: it cannot be improved in order. On the other hand, if there is insufficient sparsity, in particular when the sparsity  $s$  satisfies  $s \gg n/\log p$ , the minimax lower bounds diverge. This is no surprise as the oracle inequalities for certain M-estimators have only been shown under the condition  $s = o(n/\log p)$ . In the intermediate sparsity regime when  $\sqrt{n}/\log p \leq s < n/\log p$ , the parametric rate cannot be achieved.

As for the upper bounds, the parametric rate  $1/\sqrt{n}$  can be achieved for estimation of single entries. This basically follows directly from the asymptotic normality of the de-sparsified estimators, if sparsity of  $\beta$  is of small order  $\sqrt{n}/\log p$ . This sparsity condition is stronger than the condition necessary for oracle inequalities ( $s = o(n/\log p)$ ). However, as we discuss in Section 8.6, the sparsity condition  $s = o(\sqrt{n}/\log p)$  is essentially necessary for asymptotically normal estimation. To summarize the findings, the analysis of the minimax rates revealed that under sufficient sparsity of small order  $\sqrt{n}/\log p$ , the parametric rate of order  $1/\sqrt{n}$  is optimal, and the de-sparsified estimator achieves it (in the above mentioned cases).

In this paper, we attempt to answer further questions that arise concerning the optimality of asymptotically normal estimators in high-dimensional settings. The analysis on minimax rates does not address an important question. The derived lower bound (1) does not reveal any explicit lower bounds on the (asymptotic) variance. The question of efficiency in the spirit of the famous Cramér-Rao result thus remains open in the high-dimensional setting. This motivates us to pose the following questions. Can we establish lower bounds on the variance, similar to the Cramér-Rao bounds in the (semi-)parametric setting, also in the high-dimensional setting? And if yes, can we construct an estimator that achieves these bounds? We give an affirmative answer to these questions.

**2. Our contributions.** Asymptotic efficiency of estimators was thoroughly studied in the traditional settings; we refer the reader to the books

van der Vaart (2000), Bickel et al. (1993) and the references therein. These results are however developed for *fixed* models which do not change with  $n$ , and hence they cannot be applied to high-dimensional settings where the dimension of the parameter may grow with the sample size.

In this paper we develop a framework for establishing asymptotic efficiency of estimators in high-dimensional models changing with  $n$ . We concentrate on two approaches towards deriving the lower efficiency bounds: asymptotic Cramér-Rao bounds and Le Cam’s approach.

Firstly, we develop an asymptotic version of a semi-parametric Cramér-Rao lower bound for sparse high-dimensional linear and graphical models. To this end, we propose a strong asymptotic unbiasedness assumption. Loosely speaking, this unbiasedness assumption measures the rate at which the bias vanishes in shrinking neighbourhoods of the true distribution of “size”  $1/\sqrt{n}$ . We consider the linear model and the Gaussian graphical model and for each of them, we establish lower bounds on the variance of any asymptotically unbiased estimator. The proposed framework might be applicable to other high-dimensional models in a similar spirit.

Consequently, for linear regression and Gaussian graphical models, we show that the de-sparsified estimator is an asymptotically unbiased estimator and is asymptotically efficient, i.e. it reaches the derived lower bound. Thus, compared to previous results, which only showed asymptotic normality or minimaxity (up to order in  $n$ ) of the de-sparsified estimator, we show that it is in terms of variance the best among all asymptotically unbiased estimators: thus in this sense asymptotically efficient.

In the second approach, we extend some of the classical results of Le Cam on local asymptotic normality to the high-dimensional setting. The result underlies a likelihood expansion analysis and involves a careful adjustment of Le Cam’s arguments to the high-dimensional setting. The result obtained gives us the limiting distribution of an asymptotically linear estimator under a small perturbation of the parameter. We next show for the linear model that the de-sparsified estimator is regular: it converges locally uniformly to the limiting normal distribution with zero mean, and among all regular estimators it has the smallest asymptotic variance.

The two approaches above are strongly related, but one does not clearly dominate the other. A more detailed comparison is discussed in Section 11.

As a by-product of our analysis, we establish new oracle results for the Lasso. Typical analysis considers oracle inequalities for the prediction error and the  $\ell_1$ -error which hold with high-probability. We strengthen these oracle inequalities by showing that they also hold for the mean  $\ell_1$ -error and for higher orders of this error. These oracle inequalities are needed to claim

strong asymptotic unbiasedness of the de-sparsified estimators.

**3. Relation to prior work.** As pointed out in Section 2, the traditional results as in, for instance, [van der Vaart \(2000\)](#) or [Bickel et al. \(1993\)](#), are not directly applicable to the high-dimensional setting. We extend the traditional approach to semi-parametric efficiency to the context of high-dimensional models which requires adjustment of the arguments to a model changing with  $n$  and the sparsity of the model is required to keep remainders in approximate expansions under control. Our main results show that the lower bounds for high-dimensional models are analogous to those for parametric models, however, a new message for high-dimensional models is that to obtain the parametric lower bound, we require that the “worst possible sub-direction” is sparse. Without this condition, we are unable to claim asymptotic efficiency of the de-sparsified Lasso estimator.

Regarding the upper bounds, to construct asymptotically efficient estimators, our work follows the methodology from the works [van de Geer et al. \(2014\)](#) and [Janková and van de Geer \(2016a\)](#), where de-sparsified Lasso estimators are proposed for the linear regression and for undirected graphical models. We borrow these constructions with some small adjustments. However, the upper bounds derived for the de-sparsified estimators in the mentioned papers are not sufficient for the present analysis: we need to show a stronger oracle bound which holds in expectation. Moreover, we extend the results for estimation of single entries as considered in [van de Geer et al. \(2014\)](#) and [Janková and van de Geer \(2016a\)](#) to linear functionals.

Asymptotic efficiency of estimators in high-dimensional settings changing with  $n$  was first considered in the paper [van de Geer et al. \(2014\)](#). The paper provides a formulation of asymptotic efficiency of entries of the de-biased lasso. The approach is based on embedding the high-dimensional model into a fixed (i.e. not changing with  $n$ ) infinite-dimensional model, for which semi-parametric efficiency bounds are available (see [van der Vaart \(2000\)](#)). However, such an embedding requires a very special model structure. In the present paper, we do not use an embedding but instead directly develop the theory for models changing with  $n$ .

**4. Organization of the paper.** The particular sections of the paper are divided as follows. In Section 7 we state preliminary results on oracle inequalities for the mean  $\ell_1$ -error of the Lasso estimator. In Section 6 we propose a strong asymptotic unbiasedness assumption. Section 8 gives lower and upper bounds on the variance of asymptotically unbiased estimators in the linear model, considering random design in Section 8.3 and fixed design in Section 8.4. In Section 9 we derive lower and upper bounds on

the variance of asymptotically unbiased estimators in Gaussian graphical models. Section 10 contains an extension of Le Cam's lemma to the high-dimensional setting, which is applicable to general non-linear models. Section 11 summarizes the results, conclusions and some open questions. Finally, the proofs are contained in the supplemental article [Janková and van de Geer \(2016b\)](#).

**5. Notation.** For a vector  $x = (x_1, \dots, x_p) \in \mathbb{R}^p$  we denote its  $\ell_p$  norm by  $\|x\|_p := (\sum_{i=1}^p x_i^p)^{1/p}$  for  $p \geq 1$ . We further let  $\|x\|_\infty := \max_{i=1, \dots, p} |x_i|$  and  $\|x\|_0 = |\{i : i \in \{1, \dots, p\}, x_i \neq 0\}|$ . For a vector  $x \in \mathbb{R}^n$  we denote  $\|x\|_n^2 := \|x\|_2^2/n$  (with some abuse of notation). By  $e_i$  we denote a  $p$ -dimensional vector of zeros with a one at position  $i$ . For a matrix  $A \in \mathbb{R}^{m \times n}$ , we denote its  $(i, j)$ -th entry by  $A_{ij}$ ,  $i = 1, \dots, m, j = 1, \dots, n$ . Further, we let  $\|A\|_\infty := \max_{i=1, \dots, m, j=1, \dots, n} |A_{ij}|$ ,  $\|A\|_1 := \max_{i=1, \dots, m} \sum_{j=1}^n |A_{ij}|$  and we let  $\|A\|_F$  denote the Frobenius norm of  $A$ . We denote its  $j$ -th column by  $A_j$ . By  $\Lambda_{\min}(A)$  and  $\Lambda_{\max}(A)$  we denote the minimum and maximum eigenvalue of a symmetric matrix  $A$ , respectively. We use  $\text{tr}(A)$  to denote the trace of the matrix  $A$ . We recall here that for symmetric matrices  $A, B \in \mathbb{R}^{p \times p}$  it holds that  $\text{vec}(A)^T \text{vec}(B) = \text{tr}(AB)$ , where  $\text{vec}(A)$  is the vectorized version of a matrix  $A$  obtained by stacking columns of  $A$  on each other.

For real sequences  $f_n, g_n$ , we write  $f_n = \mathcal{O}(g_n)$  or  $f_n \lesssim g_n$  if  $|f_n| \leq C|g_n|$  for some  $C > 0$  independent of  $n$  for all  $n$ . We write  $f_n \asymp g_n$  if both  $f_n = \mathcal{O}(g_n)$  and  $1/f_n = \mathcal{O}(1/g_n)$  hold. Finally,  $f_n = o(g_n)$  if  $\lim_{n \rightarrow \infty} f_n/g_n = 0$ . For a sequence of random variables  $X_n$ , we write  $X_n = \mathcal{O}_P(f_n)$  if  $X_n/f_n$  is bounded in probability. We write  $X_n = o_P(1)$  if  $X_n$  converges to zero in probability. We use  $\rightsquigarrow$  to denote the convergence in distribution. By  $1_T$  we denote the indicator function of the set  $T$ . The identity matrix is denoted by  $I$ .

**6. Asymptotic unbiasedness.** This section defines the concept of strong asymptotic unbiasedness that will be needed for the linear and graphical model. We turn to the linear model in the next section. Consider a probability distribution  $P_\beta$  on some observation space  $\mathcal{X}$ , where the parameter  $\beta$  lies in a  $p$ -dimensional parameter space  $\mathcal{B} \subset \mathbb{R}^p$ . We consider the parameter set

$$(2) \quad \mathcal{B}(d_n) := \{\beta \in \mathcal{B} : \|\beta\|_0 \leq d_n, \|\beta\|_2 \leq C\},$$

where  $C > 0$  is some universal constant and  $d_n$  is a *known* sequence that will be specified later. We further define an  $\ell_2$ -neighbourhood of a point  $\beta \in \mathcal{B}(d_n)$  as follows

$$(3) \quad B(\beta, \varepsilon) := \{\tilde{\beta} \in \mathcal{B}(d_n) : \|\tilde{\beta} - \beta\|_2 \leq \varepsilon\}.$$

We remark that all the parameter vectors appearing in this paper are *sequences* depending on  $n$ . In general we omit the index  $n$ , except for situations where omitting the index could lead to confusion.

Let  $g : \mathcal{B} \rightarrow \mathbb{R}$  and let the parameter of interest be  $g(\beta)$ . Our goal is to derive an asymptotic lower bound for the variance of an estimator  $T_n$  of  $g(\beta)$ , which is in some sense asymptotically unbiased. To this end, we define *strong asymptotic unbiasedness* as follows.

**DEFINITION 1.** *Let  $m_n$  be a sequence such that  $n = o(m_n)$ . We say that  $T_n$  is a strongly asymptotically unbiased estimator of  $g(\beta)$  at  $\beta_0$  (in a neighbourhood of size  $c$ ) with a rate  $m_n$  if it holds that  $\text{var}_{\beta_0}(T_n) = \mathcal{O}(1/n)$  and for every  $\beta \in B\left(\beta_0, \frac{c}{\sqrt{m_n}}\right)$  it holds*

$$\lim_{n \rightarrow \infty} \sqrt{m_n}(\mathbb{E}_{\beta} T_n - g(\beta)) = 0.$$

The motivation for Definition 1 comes from the asymptotic unbiasedness assumption for semi-parametric models, which is assumed to hold in a small neighborhood of  $\beta_0$ . Definition 1 implies that the mean squared error of the considered estimator must be of order  $1/n$ .

**7. Strong oracle inequalities for the Lasso.** We present new results on oracle inequalities for the Lasso estimator in linear regression which will be needed in subsequent sections, but can also be of independent interest. Typical high-dimensional analysis derives oracle inequalities for the Lasso which hold with high probability (see [Bühlmann and van de Geer \(2011\)](#) for an overview of such results). The paper [Bellec and Tsybakov \(2016\)](#) derives bounds on the expectation of the prediction error. Here we derive oracle inequalities for the  $\ell_1$ -estimation error that hold in expectation.

Consider the linear model

$$(4) \quad Y = X\beta_0 + \epsilon,$$

where  $X$  is the  $n \times p$  design matrix with independent rows  $X^{(i)}, i = 1, \dots, n$ ,  $Y$  is the  $n \times 1$  vector of observations and  $\epsilon = (\epsilon_1, \dots, \epsilon_n)^T \in \mathbb{R}^n$  is the (unobservable) error. The error satisfies  $\mathbb{E}\epsilon = 0$  and its components  $\epsilon_i$  are independent for  $i = 1, \dots, n$ . Moreover, the error  $\epsilon$  and the design matrix  $X$  are independent. We further denote the Gram matrix by  $\hat{\Sigma} := X^T X/n$ . The vector  $\beta_0 = (\beta_1^0, \dots, \beta_p^0) \in \mathbb{R}^p$  is unknown. The unknown number of non-zero entries of  $\beta_0$  is denoted by  $s := \|\beta_0\|_0$  and is called the sparsity of  $\beta_0$ .



The Lasso estimator with a tuning parameter  $\lambda > 0$  is defined as follows:

$$(5) \quad \hat{\beta} := \arg \min_{\beta \in \mathbb{R}^p} \|Y - X\beta\|_n^2 + 2\lambda\|\beta\|_1.$$

The known results on oracle inequalities for the Lasso (5) give high-probability bounds for the prediction error and the  $\ell_1$ -error (or under some conditions, for the  $\ell_q$ -error for  $1 \leq q \leq 2$ ). In particular, for the tuning parameter  $\lambda \asymp \sqrt{\log p/n}$  and under further conditions that may be found in [Bühlmann and van de Geer \(2011\)](#), it holds

$$\|X(\hat{\beta} - \beta_0)\|_n^2 + \lambda\|\hat{\beta} - \beta_0\|_1 = \mathcal{O}_P(s\lambda^2).$$

[Bellec and Tsybakov \(2016\)](#) show analogous results for the expected prediction error  $\mathbb{E}\|X(\hat{\beta} - \beta_0)\|_n$  for the case of fixed design. We show such results may be obtained for the expected  $\ell_1$ -error, under almost identical conditions. In particular, Theorem 1 presented below implies that the mean  $\ell_1$ -error,  $\mathbb{E}_{\beta_0}\|\hat{\beta} - \beta_0\|_1$ , is up to a logarithmic factor of the same order as the oracle error  $\mathbb{E}_{\beta_0}\|\beta_{ora} - \beta_0\|_1 = \mathcal{O}(s/\sqrt{n})$ , where  $\beta_{ora}$  is the oracle maximum likelihood estimator (i.e. a maximum likelihood estimator applied with the knowledge of true non-zero entries of  $\beta_0$ ). Theorem 1 actually shows a more general result since it considers also higher-order errors, namely the  $k$ -th order error  $\mathbb{E}_{\beta_0}\|\hat{\beta} - \beta_0\|_1^k$  for any fixed  $k \in \{1, 2, \dots\}$ .

We consider the situation when the errors  $\epsilon_i$  are independent and sub-Gaussian (with a universal constant) and the design  $X$  has independent sub-Gaussian rows (with a universal constant). To this end, we recall a sub-Gaussianity assumption on random variables and vectors (see Section 14 in [Bühlmann and van de Geer \(2011\)](#)).

**DEFINITION 2.** *We say that a random vector  $Z \in \mathbb{R}^m$  has sub-Gaussian entries with constants  $K, K_2 > 0$  if*

$$\mathbb{E}e^{Z_j^2/K^2} \leq K_2, \quad j = 1, \dots, m.$$

*We say that a random vector  $Z \in \mathbb{R}^m$  is sub-Gaussian with constants  $K, K_2 > 0$  if for all  $\alpha \in \mathbb{R}^m$  such that  $\|\alpha\|_2 = 1$  it holds that*

$$\mathbb{E}e^{(\alpha^T Z)^2/K^2} \leq K_2.$$

In our further analysis, we typically require that the sub-Gaussianity condition as in Definition 2 is satisfied with universal constants  $K, K_2 > 0$ . A prime example of a sub-Gaussian random vector with a universal constant is a Gaussian random vector with zero mean and covariance matrix  $\Sigma_0$  that satisfies  $\Lambda_{\max}(\Sigma_0) = \mathcal{O}(1)$ . We formulate the conditions on the error and the design in the following.



- (A1) Assume the linear model (4), where the errors  $\epsilon_i$  are independent sub-Gaussian random variables with universal constants and with  $\mathbb{E}\epsilon_i = 0$ .
- (A2) Assume that  $X$  is a random  $n \times p$  matrix independent of  $\epsilon$  with independent rows  $X^{(i)}, i = 1, \dots, n$ , with mean zero and with sub-Gaussian entries with universal constants. We let  $\Sigma_0 := \mathbb{E}\hat{\Sigma}$  and suppose that  $1/\Lambda_{\min}(\Sigma_0) = \mathcal{O}(1)$ .
- (A2\*) Assume that  $X$  is a random  $n \times p$  matrix independent of  $\epsilon$  with independent sub-Gaussian rows  $X^{(i)}, i = 1, \dots, n$ , with universal constants, with mean zero. We let  $\Sigma_0 := \mathbb{E}\hat{\Sigma}$  and suppose that  $1/\Lambda_{\min}(\Sigma_0) = \mathcal{O}(1)$ .

Under conditions (A2) or (A2\*) we denote the inverse covariance matrix by  $\Theta_0 := \Sigma_0^{-1}$  and by  $\Theta_j^0$  we denote its  $j$ -th column ( $j = 1, \dots, p$ ).

**THEOREM 1.** *Suppose that conditions (A1), (A2) are satisfied. Suppose that  $\|\beta_0\|_2 = \mathcal{O}(1)$ ,  $s\sqrt{\log p/n} = o(1)$  and let  $k \in \{1, 2, \dots\}$  be fixed. Consider the Lasso estimator  $\hat{\beta}$  defined in (5) with a tuning parameter  $\lambda \geq c\tau\sqrt{\log p/n}$ , where  $c > 0$  is a sufficiently large universal constant and  $\tau > 1$  satisfies  $\tau^2 > 2k \log((\sqrt{s}\lambda^2)^{-1})/\log p$ . Then there exists a universal constant  $C_1$  such that*

$$(\mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1^k)^{1/k} \leq C_1 s \lambda.$$

Taking  $k = 1$ , under the conditions of Theorem 1 we obtain

$$\mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1 \leq C_1 s \lambda.$$

Theorem 1 can also be easily extended to fixed design, under a compatibility condition (see Section 13) on the Gram matrix  $\hat{\Sigma}$ , which substitutes the condition  $\Lambda_{\min}(\Sigma_0) \geq L > 0$ , and under the condition  $\|\hat{\Sigma}\|_{\infty} = \mathcal{O}(1)$ .

We comment on the conditions (A1), (A2) and  $\|\beta_0\|_2 = \mathcal{O}(1)$ ,  $\|\beta_0\|_0 = o(\sqrt{n/\log p})$  assumed in Theorem 1. Condition  $\|\beta_0\|_0 = o(\sqrt{n/\log p})$  together with conditions (A1), (A2) was used to apply the high-probability oracle results for Lasso as in Bühlmann and van de Geer (2011) to the case of random design. Condition  $\|\beta_0\|_2 = \mathcal{O}(1)$  can be justified under an assumption on the boundedness of the “signal-to-noise ratio”. The “signal-to-noise ratio” is defined as the ratio of the variance of the signal (observations) and the variance of the noise, i.e.  $\sum_{i=1}^n \text{var}_{\beta_0}(Y_i) / \sum_{i=1}^n \text{var}(\epsilon_i) = 1 + \beta_0^T \Sigma_0 \beta_0 / \sigma_{\epsilon}^2$ , where  $\sigma_{\epsilon}^2 := \frac{1}{n} \sum_{i=1}^n \text{var}(\epsilon_i)$ . Hence, under upper-boundedness of  $1/\Lambda_{\min}(\Sigma_0)$ , the signal-to-noise ratio is up to a constant lower-bounded by  $\|\beta_0\|_2^2 / \sigma_{\epsilon}^2$ . If we assume that the signal-to-noise ratio remains bounded

and the variance of the noise  $\sigma_\epsilon^2$  is bounded (as implied by condition (A1)), then the  $\ell_2$ -norm of  $\beta_0$  must also remain bounded.

Finally, the condition  $\tau^2 > 2k \log((s\lambda^2)^{-1})/\log p$  only guarantees that we choose sufficiently large regularization parameter  $\lambda \geq c\tau\sqrt{\log p/n}$  by choosing  $\tau$  large enough compared to the order  $k$  of the error that we want to control. If  $p \geq n$  and  $\lambda = c\tau\sqrt{\log p/n}$ , the condition reduces to  $\tau \geq C\sqrt{k}$  for some constant  $C > 0$ . Then clearly, this condition means that the higher order of error we want to control, the stronger regularization must be chosen.

## 8. The de-sparsified Lasso.

8.1. *Methodology.* As an initial estimator, we consider the Lasso estimator (5). The Lasso estimator is well-understood in terms of prediction and estimation error bounds, and was shown minimax optimal in terms of the prediction error and  $\ell_1$ -error. However, due to the inclusion of the  $\ell_1$ -penalty, the estimator is biased and its limiting distribution can accumulate a positive mass at zero (Knight and Fu (2000)). In view of statistical inference, a de-sparsified or de-biased version of the Lasso was then considered (see Zhang and Zhang (2014), van de Geer et al. (2014), Javanmard and Montanari (2014a), Javanmard and Montanari (2014a), Javanmard and Montanari (2014b), Javanmard and Montanari (2015)), which was shown to be asymptotically normal for estimation of  $\beta_j^0$ .

To construct the de-biased estimator, we further need to construct a surrogate inverse of  $\hat{\Sigma}$ , or in other words we need to construct an estimator of the inverse covariance matrix  $\Theta_0 = \Sigma_0^{-1}$ . We define  $\hat{\Theta}_j$  as an estimate of the column  $\Theta_j^0$  obtained by solving the following program, that will be referred to as *nodewise regression* (see van de Geer et al. (2014)). Recall that  $X$  is the design matrix with rows  $X^{(i)}, i = 1, \dots, n$ . The columns of the design matrix  $X$  will be denoted by  $X_j, j = 1, \dots, p$ , and by  $X_{-j}$  we denote the  $n \times (p-1)$  matrix obtained by removing the  $j$ -th column from  $X$ . For  $j = 1, \dots, p$ , we let

$$(6) \quad \hat{\gamma}_j := \arg \min_{\gamma \in \mathbb{R}^{p-1}} \|X_j - X_{-j}\gamma\|_n^2 + 2\lambda_j \|\gamma\|_1,$$

$$\hat{\tau}_j^2 := \|X_j - X_{-j}\hat{\gamma}_j\|_n^2 + \lambda_j \|\hat{\gamma}_j\|_1,$$

and we denote the  $j$ -th column of the nodewise Lasso estimator by

$$(7) \quad \hat{\Theta}_j := (-\hat{\gamma}_{j,1}, \dots, -\hat{\gamma}_{j,j-1}, 1, -\hat{\gamma}_{j,j+1}, \dots, -\hat{\gamma}_{j,p})^T / \hat{\tau}_j^2,$$

where  $\lambda_j \asymp \sqrt{\log p/n}$  for  $j = 1, \dots, p$ , uniformly in  $j$ . We denote the node-wise Lasso estimator by  $\hat{\Theta} := (\hat{\Theta}_1, \dots, \hat{\Theta}_p)$ . The necessary Karush-Kuhn-

Tucker conditions corresponding to the nodewise regression (obtained by replacing derivatives by sub-differentials) imply the condition  $\|\hat{\Sigma}\hat{\Theta}_j - e_j\|_\infty = \mathcal{O}_P(\lambda_j/\hat{\tau}_j^2)$  (see [van de Geer et al. \(2014\)](#)), which will be needed later. We now define the de-sparsified Lasso introduced in [van de Geer et al. \(2014\)](#),

$$(8) \quad \hat{b} := \hat{\beta} + \hat{\Theta}^T X^T (Y - X\hat{\beta})/n,$$

and we let  $\hat{b}_j$  denote its  $j$ -th entry. The motivation for the definition (8) comes from updating the initial Lasso estimator  $\hat{\beta}$  by removing the bias due to the  $\ell_1$ -penalty. We briefly summarize the main results on  $\hat{b}$  as derived in [van de Geer et al. \(2014\)](#). The estimator  $\hat{b}_j$  can be shown to be asymptotically linear with a remainder term of small order  $1/\sqrt{n}$ , in particular, under the conditions (A1), (A2\*) and

$$s = o(\sqrt{n}/\log p), \quad \max_{j=1,\dots,p} s_j = o(\sqrt{n}/\log p)$$

it holds

$$\hat{b} - \beta_0 = \Theta_0 X^T \epsilon/n + \Delta,$$

where  $s_j := \|\Theta_j^0\|_0$  and  $\|\Delta\|_\infty = o_P(1/\sqrt{n})$ . Thus, after normalization by  $\sqrt{n}$  and by the (estimated) standard deviation, asymptotic normality of entries of  $\hat{b}$  with zero mean and unit variance follows by the central limit theorem. We now investigate the question of “regularity” and asymptotic efficiency of this estimator.

We first show that the de-sparsified estimator  $\hat{b}_j$  satisfies the strong asymptotic unbiasedness condition from Definition 1 in Section 6. We then show that  $\hat{b}_j$  achieves the lower bound on the variance of any strongly asymptotically unbiased estimator. Thus in this sense the de-sparsified estimator is asymptotically efficient. In Section 8.3 we investigate the case of a random Gaussian design matrix and in Section 8.4 the case of a fixed design matrix.

**8.2. Strong asymptotic unbiasedness of the de-sparsified Lasso.** We consider estimation of linear functionals  $g(\beta) = \xi^T \beta$ , where  $\xi \in \mathbb{R}^p$  is a known vector. We define an estimator of  $g(\beta) = \xi^T \beta$  as a linear combination  $\xi$  of the de-sparsified estimator  $\hat{b}$ . This yields

$$(9) \quad \hat{b}_\xi := \xi^T \hat{b} = \xi^T \hat{\beta} + \xi^T \hat{\Theta} X^T (Y - X\hat{\beta})/n.$$

Then we have the following lemma, which shows strong asymptotic unbiasedness of  $\hat{b}_\xi$  for estimation of  $\xi^T \beta$ .

**LEMMA 1.** *Suppose that conditions (A1), (A2\*) are satisfied,  $\beta_0 \in \mathcal{B}(d_n)$  where  $d_n = o(\sqrt{n}/\log p)$ ,  $\max_j s_j \leq d_n$ ,  $\|\xi\|_1 = \mathcal{O}(1)$  and  $\|\Sigma_0\|_\infty = \mathcal{O}(1)$ .*

Let  $\hat{b}_\xi$  be the estimator defined in (9) with tuning parameters of the Lasso and nodewise regression  $\lambda \asymp \lambda_j \asymp \sqrt{\log p/n}$  uniformly in  $j = 1, \dots, p$ . Then  $\hat{b}_\xi$  is a strongly asymptotically unbiased estimator of  $\xi^T \beta$  at  $\beta_0$ .

**8.3. Main results for random design.** We derive lower bounds for the variance of a strongly asymptotically unbiased estimator. We consider the following conditions on the error distribution and the design matrix  $X$ .

- (B1) Assume the linear model (4) with  $\epsilon \sim \mathcal{N}(0, I)$ .
- (B2) Assume that  $X$  is a random  $n \times p$  matrix independent of  $\epsilon$  with independent rows  $X^{(i)} \sim \mathcal{N}(0, \Sigma_0)$  for  $i = 1, \dots, n$ . Suppose that the inverse covariance matrix  $\Theta_0 := \Sigma_0^{-1}$  exists,  $1/\Lambda_{\min}(\Sigma_0) = \mathcal{O}(1)$  and  $\|\Sigma_0\|_\infty = \mathcal{O}(1)$ .

**THEOREM 2.** Suppose that conditions (B1), (B2) are satisfied. Suppose that  $T_n$  is a strongly asymptotically unbiased estimator of  $g(\beta)$  at  $\beta_0 \in \mathcal{B}(d_n)$  with a rate  $m_n$ . Let  $h \in \mathbb{R}^p$  satisfy  $h^T \Sigma_0 h = 1$  and  $\beta_0 + h/\sqrt{m_n} \in B\left(\beta_0, \frac{c}{\sqrt{m_n}}\right)$  for a sufficiently large universal constant  $c$ . Assume moreover that for some  $\dot{g}(\beta_0) \in \mathbb{R}^p$  it holds

$$(10) \quad \sqrt{m_n} (g(\beta_0 + h/\sqrt{m_n}) - g(\beta_0)) = h^T \dot{g}(\beta_0) + o(1).$$

Then

$$\text{nvar}_{\beta_0}(T_n) \geq [h^T \dot{g}(\beta_0)]^2 - o(1).$$

Theorem 2 yields a lower bound  $[h^T \dot{g}(\beta_0)]^2 - o(1)$  on the variance of an estimator which is a strongly asymptotically unbiased estimator in a direction  $h$ , such that  $\beta_0 + h/\sqrt{m_n}$  remains within the model. By maximizing  $[h^T \dot{g}(\beta_0)]^2$  over all feasible  $h$ , we obtain the following corollary.

**COROLLARY 1.** If  $\beta_0 + \Theta_0 \dot{g}(\beta_0) / \sqrt{\dot{g}(\beta_0)^T \Theta_0 \dot{g}(\beta_0) m_n} \in B\left(\beta_0, \frac{c}{\sqrt{m_n}}\right)$ , then the lower bound from Theorem 2 is maximized at the value

$$h_0 := \Theta_0 \dot{g}(\beta_0) / \sqrt{\dot{g}(\beta_0)^T \Theta_0 \dot{g}(\beta_0)},$$

and under the conditions of Theorem 2, we get and under the conditions of Theorem 2, we get

$$\text{nvar}_{\beta_0}(T_n) \geq \dot{g}(\beta_0)^T \Theta_0 \dot{g}(\beta_0) - o(1).$$

DEFINITION 3. Let  $g$  be differentiable at  $\beta_0$  with derivative  $\dot{g}(\beta_0)$ . We call

$$c_0 := \Theta_0 \dot{g}(\beta_0) / \dot{g}(\beta_0)^T \Theta_0 \dot{g}(\beta_0)$$

the worst possible sub-direction for estimating  $g(\beta_0)$ .

The motivation for the terminology *worst possible sub-direction* in Definition 3 is given by Corollary 1. The normalization by  $\dot{g}(\beta_0)^T \Theta_0 \dot{g}(\beta_0)$  is arbitrary but natural from a projection theory point of view.

As a special case, consider estimation of  $g(\beta) = \beta_j$  for some fixed value of  $j \in \{1, \dots, p\}$ . Then  $\dot{g}(\beta) = e_j$ , the  $j$ -th unit vector in  $\mathbb{R}^p$ . Clearly,  $\Theta_0 \dot{g}(\beta_0) = \Theta_0 e_j = \Theta_j^0$  and  $g(\beta_0)^T \Theta_0 \dot{g}(\beta_0) = e_j^T \Theta_0 e_j = \Theta_{jj}^0$ , where  $\Theta_j^0$  is the  $j$ -th column of  $\Theta_0$  and  $\Theta_{jj}^0$  is its  $j$ -th diagonal element. It follows that  $c_j^0 = \Theta_j^0 / \Theta_{jj}^0$  is the worst possible sub-direction for estimating  $\beta_j$ . If  $\beta_0 + \Theta_j^0 / \sqrt{\Theta_{11}^0 m_n} \in B\left(\beta_0, \frac{c}{\sqrt{m_n}}\right)$ , then Corollary 1 implies the lower bound

$$\text{var}_{\beta_0}(T_n) \geq \Theta_{jj}^0 / n + o(1/n).$$

REMARK 1. To establish the lower bound, it is crucial that the worst possible sub-direction lies within the model. For illustration, consider the situation with the parameter of interest being  $g(\beta) = \beta_1$ . When  $\Theta_1^0$  is not sufficiently sparse, we are not allowed to take the global maximizer  $h = \Theta_1^0 / \sqrt{\Theta_{11}^0}$  in the maximum and the lower bound might thus become smaller. In that case, the lower bound is given via a sparse approximation of the (non-sparse) precision matrix. For a set  $M \subset \{1, \dots, p\}$  and a vector  $v \in \mathbb{R}^p$ , we denote  $v_M$  as a  $p$ -dimensional vector with entries not in  $M$  set to zero. Then we may write

$$\begin{aligned} & \max_{\beta_0 + h / \sqrt{m_n} \in B(\beta_0, c / \sqrt{m_n})} \frac{h^T e_1}{h^T \Sigma_0 h} \\ & \geq \max_{\substack{M \subset \{1, \dots, p\}: \\ |M| = d_n - \|\beta_0\|_0}} \max_{\substack{h \in \mathbb{R}^p: \|h_M\|_2 \leq c, \\ \|\beta_M^0 + h_M / \sqrt{m_n}\|_2 \leq C}} \frac{[h_M^T e_1]^2}{h_M^T \Sigma_0 h_M}. \end{aligned}$$

But if  $h := (\Sigma_{M,M}^0)^{-1} e_1$  satisfies  $\|h\|_2 \leq c$  and  $\|\beta_M^0 + h / \sqrt{m_n}\|_2 \leq C$ , then the lower bound is

$$\max_{M \subset \{1, \dots, p\}: |M| = d_n - \|\beta_0\|_0} (\Sigma_{M,M}^0)^{-1}_{11} - o(1),$$

where  $\Sigma_{M,M}^0$  is the reduction of  $\Sigma_0$  obtained by keeping only columns and rows belonging to the set  $M$ . If  $\Theta_1^0$  has sparsity  $d_n - \|\beta_0\|_0$ , then this lower

bound coincides with  $(\Sigma_0)_{11}^{-1} - o(1)$  as before. If  $\Theta_1^0$  is not sufficiently sparse, then the lower bound is given via a sparse approximation of the precision matrix. Finally, as will be seen in the following sections, without assuming the sparsity condition on the worst possible sub-direction, we would not be able to conclude asymptotic efficiency of the de-sparsified Lasso estimator.

Finally we show that the de-sparsified estimator  $\hat{b}_j$  achieves the lower bound on the variance. Thus the de-sparsified estimator is strongly asymptotically unbiased and has the smallest variance among all strongly asymptotically unbiased estimators. We assume Gaussianity of the error and the design matrix, as the lower bounds have only been derived for this case.

**THEOREM 3.** *Suppose that conditions (B1), (B2) are satisfied,  $\beta_0 \in \mathcal{B}(d_n)$  with  $d_n = o(\sqrt{n}/\log p)$  and  $\max_j s_j \leq d_n$ . Assume that  $\|\xi\|_1 = \mathcal{O}(1)$ . Let  $\hat{b}_\xi$  be the estimator defined in (9) with tuning parameters of the Lasso and nodewise regression  $\lambda \asymp \lambda_j \asymp \sqrt{\log p/n}$ , uniformly in  $j = 1, \dots, p$ . Then  $\hat{b}_\xi$  is a strongly asymptotically unbiased estimator of  $\xi^T \beta$  at  $\beta_0$ . Let  $T$  be any strongly asymptotically unbiased estimator of  $\xi^T \beta$  at  $\beta_0$  and assume that  $\beta_0 + \Theta_0 \xi / (\xi^T \Theta_0 \xi n)^{1/2} \in B(\beta_0, c/\sqrt{n})$ . Then it holds*

$$\text{var}_{\beta_0}(T) \geq \frac{\xi^T \Theta_0 \xi + o(1)}{n}, \quad \text{var}_{\beta_0}(\hat{b}_\xi) = \frac{\xi^T \Theta_0 \xi + o(1)}{n}.$$

To obtain the result of Theorem 3, we assumed that  $\beta_0 + \Theta_0 \xi / (\xi^T \Theta_0 \xi n)^{1/2} \in B(\beta_0, c/\sqrt{n})$ , which guarantees that the worst possible sub-direction stays within the model. Further we assumed that the sparsity in  $\beta_0$  satisfies  $s = o(\sqrt{n}/\log p)$  and that the sparsity in the rows of  $\Theta_0$  is of small order  $\sqrt{n}/\log p$ . Thus, to be able to claim asymptotic efficiency of the de-sparsified Lasso, we not only require sparsity in  $\beta_0$ , but also sufficient sparsity in the precision matrix. Note that the sparsity condition on  $\beta_0$  is almost a necessary condition as discussed in Section 8.6 below.

**8.4. Main results for fixed design.** In this section, we assume that the design matrix  $X$  is fixed (non-random). Recall that  $\hat{\Sigma} = X^T X/n$  is the Gram matrix. The following theorem is an analogy of Theorem 2 for fixed design.

**THEOREM 4.** *Let  $X$  be a fixed  $n \times p$  matrix and suppose that condition (B1) is satisfied. Let  $h \in \mathbb{R}^p$  be such that  $h^T \hat{\Sigma} h = \mathcal{O}(1)$  and  $\beta_0 + h/\sqrt{m_n} \in B(\beta_0, c/\sqrt{m_n})$ . Suppose that  $T_n$  is a strongly asymptotically unbiased estimator of  $g(\beta)$  at  $\beta_0$  in the direction  $h$  with rate  $m_n$ . Assume moreover that*

for some  $\dot{g}(\beta_0) \in \mathbb{R}^p$  it holds that

$$(11) \quad \sqrt{m_n}(g(\beta_0 + h/\sqrt{m_n}) - g(\beta_0)) = h^T \dot{g}(\beta_0) + o(1).$$

Then

$$n\text{var}_{\beta_0}(T_n) \geq [h^T \dot{g}(\beta_0)]^2 - o(1).$$

For fixed design, the matrix  $\hat{\Sigma}$  is not invertible, and thus we cannot use the reasoning as in Section 8.3. We can however try to remedy this by proposing an approximate worst possible sub-direction. To this end, we may use an estimator  $\hat{\Theta}$ , which acts as a surrogate inverse of  $\hat{\Sigma}$  in a certain sense. Such an estimate can be obtained in the same way as for the random design, using the nodewise regression (7). The necessary Karush-Kuhn-Tucker conditions of the nodewise regression (obtained by replacing derivatives by sub-differentials) again imply the condition  $\|\hat{\Sigma}\hat{\Theta}_j - e_j\|_\infty = \mathcal{O}_P(\lambda_j/\hat{\tau}_j^2)$ . The de-sparsified estimator can then be defined in the same way as for the random design, as in equation (8).

We consider estimation of  $g(\beta_0) := \beta_j^0$ , although one could further consider estimation of linear functionals, similarly as for the random design. Strong asymptotic unbiasedness of  $\hat{b}_j$  for estimation of  $\beta_j$  then follows similarly as in Lemma 1 (with  $g(\beta) = \beta_j$ ) for all  $\beta \in \mathcal{B}(d_n)$ , under  $d_n = o(\sqrt{n}/\log p)$ , if the compatibility condition is satisfied for  $\hat{\Sigma}$  with a universal constant and  $\|\hat{\Sigma}\|_\infty = \mathcal{O}(1)$ . For the definition of the compatibility condition, see Definition 4 in Section 13 of the supplemental article [Janková and van de Geer \(2016b\)](#). We formulate the asymptotic efficiency of  $\hat{b}_j$  for  $g(\beta) := \beta_j$  in the following theorem.

**THEOREM 5.** *Assume that condition (B1) is satisfied and  $\beta_0 \in \mathcal{B}(d_n)$  with  $d_n = o(\sqrt{n}/\log p)$ . Let  $j \in \{1, \dots, p\}$  and let  $\hat{\Theta}_j$  be obtained using the nodewise regression as in (7) with  $\lambda_j \asymp \sqrt{\log p/n}$ . Suppose that  $\beta_0 + \hat{\Theta}_j/(\hat{\Theta}_{jj}n)^{1/2} \in B(\beta_0, c/\sqrt{n})$ ,  $\|\hat{\Theta}_j\|_2 = \mathcal{O}(1)$ , the compatibility condition is satisfied for  $\hat{\Sigma}$  with a universal constant and  $\|\hat{\Sigma}\|_\infty = \mathcal{O}(1)$ . Then  $\hat{b}_j$  defined in (8) using  $\hat{\Theta}_j$  and with  $\lambda \asymp \sqrt{\log p/n}$  is a strongly asymptotically unbiased estimator of  $\beta_j$  at  $\beta_0$  and for any strongly asymptotically unbiased estimator  $T$  of  $\beta_j$  at  $\beta_0$  it holds*

$$\text{var}_{\beta_0}(T) \geq \frac{\hat{\Theta}_{jj} + o(1)}{n}, \quad \text{var}_{\beta_0}(\hat{b}_j) = \frac{\hat{\Theta}_{jj} + o(1)}{n}.$$

The condition  $\beta_0 + \hat{\Theta}_j/\sqrt{\hat{\Theta}_{jj}n} \in \mathcal{B}(d_n)$  implies that  $\|\hat{\Theta}_j\|_0 = \mathcal{O}(d_n)$ . To this end, we refer to Lemma 12 in Section 14 of the supplemental article [Janková](#)



and van de Geer (2016b), which shows that sparsity in  $\hat{\Theta}_j$  constructed using nodewise regression is guaranteed under random design. The condition  $\|\hat{\Theta}_j\|_2 = \mathcal{O}(1)$  replaces the eigenvalue condition we needed in the case of random design.

8.5. *Le Cam's bounds.* In this section, we provide an alternative approach, which makes another choice in the formulation of asymptotic efficiency. This approach is based on Le Cam's arguments (see e.g. van der Vaart (2000)) rather than the Cramér-Rao bounds, and it allows us to show that the convergence of the de-sparsified estimator to the limiting normal distribution with smallest possible variance is locally uniform in the underlying unknown parameter, and the asymptotic variance of the de-sparsified estimator is smallest among the class of asymptotically linear estimators. Furthermore, the result identifies the asymptotic bias of asymptotically linear estimators. A detailed comparison of the two approaches for deriving the lower bounds is deferred to Section 11.

We consider the setting from Section 8.3, where the design matrix  $X$  is random with the parameter of interest being  $g(\beta) = \beta_j$ .

**THEOREM 6.** *Assume that conditions (B1), (B2) are satisfied,  $\beta_0 \in \mathcal{B}(d_n)$  with  $d_n = o(\sqrt{n}/\log p)$ ,  $\|\Theta_j^0\|_0 \leq d_n$  and  $\Lambda_{\max}(\Sigma_0) = \mathcal{O}(1)$ . Assume that  $\hat{b}_j$  is defined in (8) with tuning parameters  $\lambda \asymp \lambda_j \asymp \sqrt{\log p/n}$ . Then for every  $\tilde{\beta}_n \in B\left(\beta_0, \frac{c}{\sqrt{n}}\right)$  it holds*

$$\frac{\sqrt{n}(\hat{b}_j - \tilde{\beta}_n)}{(\Theta_{jj}^0)^{1/2}} \overset{\tilde{\beta}_n}{\rightsquigarrow} \mathcal{N}(0, 1).$$

Let  $T_n$  be an asymptotically linear estimator with an influence function  $l_{\beta_0}$ :

$$(12) \quad T_n - \beta_j^0 = \frac{1}{n} \sum_{i=1}^n l_{\beta_0}(X^{(i)}, Y^{(i)}) + o_{P_{\beta_0}}(n^{-1/2}),$$

where  $\mathbb{E}l_{\beta_0}(X^{(i)}, Y^{(i)}) = 0$  and  $\text{var}(l_{\beta_0}(X^{(i)}, Y^{(i)})) =: V_{\beta_0} < \infty$ . Assume that for all  $h \in \mathbb{R}^p$  and  $i = 1, \dots, n$  it holds

$$(13) \quad \mathbb{E}l_{\beta_0}(X^{(i)}, Y^{(i)})\epsilon_i h^T X^{(i)} - h_j = o(1).$$

Then

$$V_{\beta_0} \geq \Theta_{jj}^0 + o(1).$$

8.6. *Discussion of the conditions.* We briefly discuss the conditions assumed to obtain the above results. To establish asymptotic efficiency of the de-sparsified estimator, we considered conditions analogous to the conditions assumed in [van de Geer et al. \(2014\)](#). These include a sparsity condition on the parameter  $\beta_0$  of order  $o(\sqrt{n}/\log p)$ , conditions on the covariance matrix  $\Lambda_{\min}(\Sigma_0) = \mathcal{O}(1)$ ,  $\|\Sigma_0\|_{\infty} = \mathcal{O}(1)$ , sparsity of the precision matrix and a Gaussianity assumption on the rows on the precision matrix. Unlike in [van de Geer et al. \(2014\)](#), we assume Gaussianity of the design matrix and the error; this condition was needed to derive the lower bounds. In addition to the conditions from [van de Geer et al. \(2014\)](#), we also assume boundedness of  $\ell_2$ -norm of  $\beta_0$ , which follows if the signal to noise ratio is bounded as argued in Section 7. Condition (13) from Theorem 6 is a variant of asymptotic unbiasedness which is known to be satisfied in many traditional settings (see e.g. [van der Vaart \(2000\)](#)). The condition is discussed in more detail in Section 10 below.

Our analysis requires the sparsity condition  $s = o(\sqrt{n}/\log p)$ . This condition is essentially necessary in the linear regression setting for construction of an asymptotically normal estimator, as argued in the following. First observe that if the (slightly weaker) condition  $s = \mathcal{O}(\sqrt{n}/\log p)$  is not satisfied, then there cannot exist an estimator  $T_n$  of  $\beta_j \in \mathbb{R}$  and a sequence  $\sigma_n = \mathcal{O}(1)$  such that

$$(14) \quad \sqrt{n}(T_n - \beta_j^0)/\sigma_n \rightsquigarrow \mathcal{N}(0, 1).$$

Suppose that there exists an estimator  $T_n$  that satisfies (14). Then necessarily  $\sqrt{n}(T_n - \beta_j^0)/\sigma_n = \mathcal{O}_P(1)$ . By similar reasoning as in [Ren et al. \(2015\)](#), we have under the conditions assumed the minimax rates for  $\mathbb{E}|T_n - \beta_j^0|$  of order  $\frac{1}{\sqrt{n}} + \frac{s \log p}{n}$ . But then necessarily  $s \log p/n = \mathcal{O}(1/\sqrt{n})$ , which gives  $s = \mathcal{O}(\sqrt{n}/\log p)$ . This is only slightly weaker than the condition we require,  $s = o(\sqrt{n}/\log p)$ .

Furthermore, for simplicity of presentation, we assumed that the variance of the noise is fixed at  $\sigma_{\epsilon} = 1$ . In general, we can include the parameter  $\sigma_{\epsilon}$  as an unknown parameter in the model, and by orthogonality of the score corresponding to this parameter and the score corresponding to  $\beta$ , we can easily extend the arguments. The noise variance will then appear in both lower and upper bounds.

**9. Gaussian graphical models.** In this part, we consider efficient estimation of edge weights in undirected Gaussian graphical models. Gaussian graphical models have become a popular tool for representing dependencies within large sets of variables and have found application in areas such

as neuroscience, biology and climate data analysis. In particular, Gaussian graphical models encode conditional dependencies between variables (nodes in the graph) by including an edge between two variables if and only if they are not independent given all the other variables. This corresponds to the problem of estimation of the precision matrix of a multivariate normal distribution, which we now introduce.

(C1) Assume that the  $n \times p$  matrix  $X$  has independent rows  $X^{(i)}$ ,  $i = 1, \dots, n$  which are  $\mathcal{N}_p(0, \Sigma_0)$ -distributed.

Denote the precision matrix by  $\Theta_0 := \Sigma_0^{-1}$ , where the inverse of  $\Sigma_0$  is assumed to exist. The matrix  $\Theta_0 \in \mathbb{R}^{p \times p}$  is unknown, but we assume bounds on its row-sparsity (column-sparsity)  $s_j := \|\Theta_j^0\|_0$ , where  $\Theta_j^0$  is the  $j$ -th column of the precision matrix.

**9.1. Methodology.** There have been several methods proposed for estimation of the precision matrix in the high-dimensional setting when  $p \gg n$  (see [Friedman, Hastie and Tibshirani \(2008\)](#), [Meinshausen and Bühlmann \(2006\)](#)). These methods are based on regularization techniques and lead to estimators that are biased. De-biasing was then studied similarly as in the linear regression, and it was shown that de-biasing leads to estimators which are asymptotically normal. For our further analysis, we consider the de-sparsified nodewise Lasso estimator proposed in [Janková and van de Geer \(2016a\)](#). We show that this estimator is strongly asymptotically unbiased and reaches the lower bound on the variance derived in the previous section.

To introduce the methodology, consider again the nodewise Lasso estimator  $\hat{\Theta} = (\hat{\Theta}_1, \dots, \hat{\Theta}_p)$  defined in (7). Define the de-sparsified nodewise Lasso (see [Janková and van de Geer \(2016a\)](#))

$$(15) \quad \hat{T} := \hat{\Theta} + \hat{\Theta}^T - \hat{\Theta} \hat{\Sigma} \hat{\Theta}.$$

Furthermore, we write  $\hat{T}_{ij} := \hat{\Theta}_{ij} + \hat{\Theta}_{ji} - \hat{\Theta}_i^T \hat{\Sigma} \hat{\Theta}_j$  for  $i, j = 1, \dots, p$ . The method and its asymptotic properties were studied in [Janková and van de Geer \(2016a\)](#). The estimator  $\hat{\Theta}_j$  can be shown to be asymptotically linear with a remainder term of small order  $1/\sqrt{n}$ , in particular, under condition (C1) and under  $\max_{j=1, \dots, p} s_j = o(\sqrt{n}/\log p)$  it holds

$$\hat{T} - \Theta_0 = -\Theta_0^T (\hat{\Sigma} - \Sigma_0) \Theta_0 + \Delta,$$

where  $\|\Delta\|_\infty = o_P(1/\sqrt{n})$ . Thus, after normalization by  $\sqrt{n}$  and by the (estimated) standard deviation, it follows that it is asymptotically standard normal and minimax optimal (see [Ren et al. \(2015\)](#), [Janková and van de Geer \(2016a\)](#)). We investigate the question of “regularity” and asymptotic efficiency of the proposed estimator.

9.2. *Strong asymptotic unbiasedness of the de-sparsified nodewise Lasso.* Suppose that the parameter  $\Theta$  ranges over a parameter space  $T \subset \mathbb{R}^{p \times p}$ . We then define the parameter set

$$\mathcal{G}(d_1, \dots, d_p) := \{\Theta \in T : \Theta = \Theta^T, \|\Theta_j\|_0 \leq C_1 d_j, j = 1, \dots, p, \\ 1/\Lambda_{\min}(\Theta) \leq C_2, \Lambda_{\max}(\Theta) \leq C_3\},$$

for some universal constants  $C_1, C_2, C_3 > 0$ . We also need to readjust the definition of a neighbourhood from (3); hence in this section we let

$$B(\Theta, \epsilon) := \{\tilde{\Theta} \in \mathcal{G}(d_1, \dots, d_p) : \|\tilde{\Theta} - \Theta\|_F \leq \epsilon\}.$$

The following lemma shows that  $\hat{T}_{ij}$  is strongly asymptotically unbiased for estimation of  $\Theta_{ij}^0$ .

LEMMA 2. *Let  $i, j \in \{1, \dots, p\}$ , assume that condition (C1) is satisfied and  $\Theta_0 \in \mathcal{G}(d_1, \dots, d_p)$  with  $\max(d_i, d_j) = o(\sqrt{n}/\log p)$ . Let  $\hat{T}_{ij}$  be defined in (15), where  $\hat{\Theta}_i, \hat{\Theta}_j$  are the  $i$ -th and  $j$ -th columns of the nodewise Lasso estimator with tuning parameters  $\lambda_i \asymp \lambda_j \asymp \sqrt{\log p/n}$ . Then  $\hat{T}_{ij}$  is a strongly asymptotically unbiased estimator for  $\Theta_{ij}^0$ .*

9.3. *Main results.* We first derive an asymptotic lower bound for the variance of  $T_n$  when  $T_n$  is strongly asymptotically unbiased. We restrict our attention to estimation of linear functionals of the precision matrix  $\Theta_0$ ,  $h(\Theta_0) = \text{tr}(\Psi\Theta_0)$ , where  $\Psi \in \mathbb{R}^{p \times p}$  is a known matrix. We shall consider the case when  $\Psi$  is of rank one, say  $\Psi = \xi_1 \xi_2^T$  for some vectors  $\xi_1, \xi_2 \in \mathbb{R}^p$ . This corresponds to estimation of  $g(\Theta_0) = \xi_1^T \Theta_0 \xi_2$ , where  $\xi_1, \xi_2 \in \mathbb{R}^p$  are known vectors.

Contrary to previous sections, the high-dimensional parameter is a matrix, therefore instead of a vector direction  $h$  we shall write the capital letter  $H$  to denote a matrix direction in  $\mathbb{R}^{p \times p}$ .

THEOREM 7. *Assume condition (C1), assume that  $\Theta_0 \in \mathcal{G}(d_1, \dots, d_p)$  where  $\max_{j=1, \dots, p} d_j = o(\sqrt{n}/\log p)$  and  $\Theta_0 + H/\sqrt{n} \in B(\Theta_0, c/\sqrt{n})$ . Suppose that  $T_n$  is a strongly asymptotically unbiased estimator of  $g(\Theta) = \xi_1^T \Theta \xi_2$  at  $\Theta_0 \in \mathcal{G}(d_1, \dots, d_p)$  in the direction  $H := \Theta_0(\xi_1 \xi_2^T + \xi_2 \xi_1^T)\Theta_0/\sigma$ , where*

$$\sigma^2 := \xi_1^T \Theta_0 \xi_1 \xi_2^T \Theta_0 \xi_2 + (\xi_1^T \Theta_0 \xi_2)^2.$$

Then it holds

$$\text{var}_{\Theta_0}(T_n) \geq \frac{\sigma^2 - o(1)}{n}.$$

As a corollary, consider estimation of  $g(\Theta_0) = \Theta_{ij}^0$  for some fixed  $(i, j) \in \{1, \dots, p\}^2$ . Then the worst sub-direction is given by  $H := (\Theta_i^0(\Theta_j^0)^T + \Theta_j^0(\Theta_i^0)^T)/\sigma$  where  $\sigma^2 := (\Theta_{ij}^0)^2 + \Theta_{ii}^0\Theta_{jj}^0$  and the corresponding lower bound is  $((\Theta_{ij}^0)^2 + \Theta_{ii}^0\Theta_{jj}^0)/n + o(1/n)$ .

We now show that the de-sparsified estimator  $\hat{T}_{ij}$  reaches the lower bound on the variance for the parameter of interest  $g(\Theta_0) = \Theta_{ij}^0$ .

**THEOREM 8.** *Suppose that condition (C1) holds,  $\Theta_0 \in \mathcal{G}(d_1, \dots, d_p)$  where  $\max(d_i, d_j) = o(\sqrt{n}/\log p)$ . Suppose that  $\Theta_0 + H/\sqrt{n} \in B(\Theta_0, c/\sqrt{n})$  for  $H := (\Theta_i^0(\Theta_j^0)^T + \Theta_j^0(\Theta_i^0)^T)/\sigma$ . Let  $\hat{T}_{ij}$  be defined in (15), where  $\hat{\Theta}_i, \hat{\Theta}_j$  are the  $i$ -th and  $j$ -th columns of the nodewise Lasso estimator with tuning parameters  $\lambda_i \asymp \lambda_j \asymp \sqrt{\log p/n}$ . Then  $\hat{T}_{ij}$  is a strongly asymptotically unbiased estimator of  $\Theta_{ij}$  at  $\Theta_0$  and for any strongly asymptotically unbiased estimator  $T$  of  $\Theta_{ij}$  at  $\Theta_0$  it holds*

$$\text{var}_{\Theta_0}(T) \geq \frac{\Theta_{ii}^0\Theta_{jj}^0 + (\Theta_{ij}^0)^2 + o(1)}{n}, \quad \text{var}_{\Theta_0}(\hat{T}_{ij}) = \frac{\Theta_{ii}^0\Theta_{jj}^0 + (\Theta_{ij}^0)^2 + o(1)}{n}.$$

The condition  $\Theta_0 + H/\sqrt{n} \in B(\Theta_0, c/\sqrt{n})$  for  $H = (\Theta_i^0(\Theta_j^0)^T + \Theta_j^0(\Theta_i^0)^T)/\sigma$  ensures that perturbation of  $\Theta_0$  along the worst possible sub-direction  $H$  lies within the model. This also implies that  $\|H_k\|_0 \leq 2C_1 d_k, k = 1, \dots, p$ , which in turn implies that necessarily  $\|\Theta_i^0\|_0 = \mathcal{O}(d_k), \|\Theta_j^0\|_0 = \mathcal{O}(d_k)$  for  $k = 1, \dots, p$ . Note that we only require sparsity in the  $i$ -th and  $j$ -th column of the precision matrix. Furthermore, we must have  $\|H\|_F \leq c$ . This is satisfied under the eigenvalue conditions noting that  $\|H\|_F^2 = \text{tr}(H^T H)$  and  $\|\Theta_k^0\|_2 = \mathcal{O}(1)$  for  $k = i, j$ .

**9.4. Discussion of the conditions.** We comment on the conditions used to obtain the above results. The conditions under which we show asymptotic efficiency only include eigenvalue conditions on the true precision matrix, sparsity conditions on columns/rows of the precision matrix and Gaussianity of the observations  $X^{(i)}, i = 1, \dots, n$ . These conditions are almost identical to conditions in van de Geer et al. (2014) and Janková and van de Geer (2016a), with the exception of Gaussianity which was used for deriving the lower bounds. In particular, the condition on row sparsity required is the same as for the linear model:  $s = o(\sqrt{n}/\log p)$ . In view of the results on minimax rates for estimation of elements of precision matrices (which are derived in Ren et al. (2015)), the condition  $s = o(\sqrt{n}/\log p)$  is necessary for asymptotically normal estimation, which follows by similar reasoning as for the linear regression.

**10. Le Cam's bounds for general models.** In this section, we provide an extension to general non-linear models and a general parameter of interest. This is achieved via adjustment of Le Cam's arguments on asymptotic efficiency to the high-dimensional setting. Let  $X^{(1)}, \dots, X^{(n)}$  be i.i.d. with distribution  $P_{\beta_{n,0}} : \beta_{n,0} \in \mathcal{B}$  where  $\mathcal{B}$  is an open convex subset of  $\mathbb{R}^p$ . We consider the parameter set

$$\mathcal{B}(d_n) := \{\beta \in \mathcal{B} : \|\beta\|_0 \leq C_1 d_n, \|\beta\|_2 \leq C_2\},$$

where  $C_1, C_2 = \mathcal{O}(1)$  and  $d_n$  is a known sequence that will be specified later. Suppose that the parameter of interest is  $g(\beta_{n,0})$  for some function  $g : \mathcal{B} \rightarrow \mathbb{R}$ . Assume that for an estimator  $T_n$  of  $g(\beta_{n,0})$ , we can show asymptotic linearity: there exists a real-valued function  $l_{\beta_{n,0}}$  on  $\mathcal{X}$  (an influence function) and some sequence  $\beta_{n,0}$  such that

$$T_n - g(\beta_{n,0}) = \frac{1}{n} \sum_{i=1}^n l_{\beta_{n,0}}(X^{(i)}) + o_{P_{\beta_{n,0}}}(n^{-1/2}),$$

where  $P_{\beta_{n,0}} l_{\beta_{n,0}} = 0$  and the variance  $V_{\beta_{n,0}} := P_{\beta_{n,0}} l_{\beta_{n,0}}^2 < \infty$ . Under the conditions of the central limit theorem, the asymptotic linearity implies that

$$(16) \quad \sqrt{n}(T_n - g(\beta_{n,0}))/V_{\beta_{n,0}}^{1/2} \overset{\beta_{n,0}}{\rightsquigarrow} \mathcal{N}(0, 1).$$

For asymptotically linear estimators, we thus have the ‘‘asymptotic variance’’  $V_{\beta_{n,0}} = P_{\beta_{n,0}} l_{\beta_{n,0}}^2$ . We shall need some conditions on the differentiability of  $g$  and the score function. Furthermore, we shall need a Lindeberg's condition related to the influence and score function. Assume that  $P_\beta$  is dominated by some  $\sigma$ -finite measure for all  $\beta$  in the parameter space and denote the corresponding probability densities by  $p_\beta$ . We denote the log-likelihood by  $\ell_\beta(x) := \log p_\beta(x)$  and the score function by  $s_\beta(x) := \frac{\partial \ell_\beta(x)}{\partial \beta}$  for all  $x \in \mathcal{X}$ .

(D1) (*Differentiability of  $g$* ) Suppose that for a given  $\tilde{\beta}_n \in B(\beta_{n,0}, \frac{c}{\sqrt{n}})$  it holds

$$\sqrt{n}(g(\tilde{\beta}_n) - g(\beta_{n,0})) = h^T \dot{g}(\beta_{n,0}) + o(1),$$

where  $h = \sqrt{n}(\tilde{\beta}_n - \beta_{n,0})$ .

(D2) (*Differentiability of the score*) Suppose that the score function  $\beta \mapsto s_\beta$  is twice differentiable and the second derivative satisfies  $\|\ddot{s}_\beta\|_\infty \leq L$  for some universal constant  $L > 0$  and for all  $\beta \in \mathcal{B}(d_n)$ . Let  $I_{\beta_{n,0}} := P_{\beta_{n,0}} s_{\beta_{n,0}} s_{\beta_{n,0}}^T$  and assume that  $\Lambda_{\max}(I_{\beta_{n,0}}) = \mathcal{O}(1)$ ,  $1/\Lambda_{\min}(I_{\beta_{n,0}}) = \mathcal{O}(1)$  and

$$(17) \quad \left\| \frac{1}{n} \sum_{i=1}^n \dot{s}_{\beta_{n,0}} + I_{\beta_{n,0}} \right\|_\infty = \mathcal{O}_P(\lambda),$$

for some  $\lambda > 0$ . Suppose that  $d_n = o(\max\{1/\lambda, n^{1/3}\})$ .

(D3) (*Lindeberg's condition*) Denote  $f_{\beta_{n,0}}(x) := l_{\beta_{n,0}}(x) + h^T s_{\beta_{n,0}}(x)$  for  $x \in \mathbb{R}^p$ . Suppose that for all  $\epsilon > 0$

$$(18) \quad \lim_{n \rightarrow \infty} P_{\beta_{n,0}} f_{\beta_{n,0}}^2 \mathbf{1}_{|f_{\beta_{n,0}}| > \epsilon \sqrt{n}} = 0,$$

and assume that  $V_{\beta_{n,0}} := P_{\beta_{n,0}} l_{\beta_{n,0}}^2 = \mathcal{O}(1)$  and  $1/V_{\beta_{n,0}} = \mathcal{O}(1)$ .

Condition (D1) is a differentiability condition on  $g$ ; an analogous condition is assumed in the first approach through Cramér-Rao bounds. Condition (D2) is a differentiability condition on the score, which is used to obtain a Taylor expansion of the likelihood. Furthermore, the condition (17) guarantees that  $-\frac{1}{n} \sum_{i=1}^n \dot{s}_{\beta_{n,0}}(X^{(i)})$  is a good estimator of the Fisher information in supremum norm. This can be verified e.g. for linear regression with  $\lambda \asymp \sqrt{\log p/n}$ . Condition (D2) further assumes the sparsity  $d_n = o(\max\{1/\lambda, n^{1/3}\})$ , which guarantees that the likelihood ratio expansion approximately holds. Finally, condition (D3) is a Lindeberg's condition which is needed to conclude asymptotic normality of certain quantities, since in Theorem 9 below we do not require any distributional assumption. This condition can be verified for particular models.

**THEOREM 9.** *Let  $g : \mathcal{B} \rightarrow \mathbb{R}$  and suppose that for some fixed sequence  $\beta_{n,0} \in \mathcal{B}(d_n)$  it holds*

$$(19) \quad T_n - g(\beta_{n,0}) = \frac{1}{n} \sum_{i=1}^n l_{\beta_{n,0}}(X^{(i)}) + o_{P_{\beta_{n,0}}}(n^{-1/2}),$$

where  $P_{\beta_{n,0}} l_{\beta_{n,0}} = 0$ . For some fixed constant  $c > 0$ , let  $\tilde{\beta}_n \in B(\beta_{n,0}, \frac{c}{\sqrt{n}})$  and denote  $h := \sqrt{n}(\tilde{\beta}_n - \beta_{n,0})$ . Suppose that conditions (D1), (D2) and (D3) are satisfied. Then it holds

$$\frac{\sqrt{n}(T_n - g(\beta_{n,0} + \frac{h}{\sqrt{n}})) - (P_{\beta_{n,0}}(l_{\beta_{n,0}} h^T s_{\beta_{n,0}}) - h^T \dot{g}(\beta_{n,0}))}{V_{\beta_{n,0}}^{1/2}} \overset{\beta_{n,0} + \frac{h}{\sqrt{n}}}{\rightsquigarrow} \mathcal{N}(0, 1).$$

The result of Theorem 9 contains a bias term  $P_{\beta_{n,0}}(l_{\beta_{n,0}} h^T s_{\beta_{n,0}}) - h^T \dot{g}(\beta_{n,0})$  which depends on  $h$ . Now consider that the bias term in the result of the theorem above vanishes, i.e. that the following condition on the score function  $s_{\beta_{n,0}}$  and the function  $l_{\beta_{n,0}}$  is satisfied: for every  $h \in \mathbb{R}^p$  it holds that

$$(20) \quad P_{\beta_{n,0}}(l_{\beta_{n,0}} h^T s_{\beta_{n,0}}) - h^T \dot{g}(\beta_{n,0}) = o(1).$$



The condition (20) is a variant of asymptotic unbiasedness which is known to be satisfied in many traditional settings. If condition (20) is satisfied, then the Cauchy-Schwarz inequality implies

$$(h^T \dot{g}(\beta_{n,0}))^2 \leq V_{\beta_{n,0}} h^T I_{\beta_{n,0}} h + o(V_{\beta_{n,0}}^{1/2} (h^T I_{\beta_{n,0}} h)^{1/2}).$$

Hence this implies a lower bound on the asymptotic variance  $V_{\beta_{n,0}}$  of an asymptotically linear estimator as follows

$$(21) \quad V_{\beta_{n,0}} \geq (h^T \dot{g}(\beta_{n,0}))^2 / h^T I_{\beta_{n,0}} h + o(V_{\beta_{n,0}}^{1/2} / (h^T I_{\beta_{n,0}} h)^{1/2}).$$

Assuming that the inverse of  $I_{\beta_{n,0}}$  exists, the right-hand side of (21) is maximized at  $h = I_{\beta_{n,0}}^{-1} \dot{g}(\beta_{n,0})$ , provided that  $\beta_{n,0} + h/\sqrt{n} \in B(\beta_{n,0}, c/\sqrt{n})$ . Hence we obtain the following lower bound on the asymptotic variance

$$V_{\beta_{n,0}} \geq \dot{g}(\beta_{n,0})^T I_{\beta_{n,0}}^{-1} \dot{g}(\beta_{n,0}) + o(V_{\beta_{n,0}}^{1/2} (\dot{g}(\beta_{n,0})^T I_{\beta_{n,0}}^{-1} \dot{g}(\beta_{n,0}))^{1/2}).$$

We summarize this simple claim in the lemma below.

LEMMA 3. *Let  $T_n$  satisfy (19) with  $V_{\beta_{n,0}} = \mathcal{O}(1)$ ,  $1/\Lambda_{\min}(I_{\beta_{n,0}}) = \mathcal{O}(1)$  and for every  $h \in \mathbb{R}^p$  it holds that*

$$(22) \quad P_{\beta_{n,0}}(l_{\beta_{n,0}} h^T s_{\beta_{n,0}}) - h^T \dot{g}(\beta_{n,0}) = o(1),$$

*then if  $\beta_{n,0} + I_{\beta_{n,0}}^{-1} \dot{g}(\beta_{n,0})/\sqrt{n} \in B(\beta_{n,0}, c/\sqrt{n})$ , it holds that*

$$V_{\beta_{n,0}} \geq \dot{g}(\beta_{n,0})^T I_{\beta_{n,0}}^{-1} \dot{g}(\beta_{n,0}) + o(1).$$

Theorem 9 in conjunction with Lemma 3 gives the result summarized in Corollary 2 below.

COROLLARY 2. *Suppose that conditions of Theorem 9 and the condition (22) are satisfied and that  $\beta_{n,0} + I_{\beta_{n,0}}^{-1} \dot{g}(\beta_{n,0})/\sqrt{n} \in B(\beta_{n,0}, c/\sqrt{n})$ . Then*

$$(23) \quad \sqrt{n}(T_n - g(\beta_{n,0} + h/\sqrt{n})) / V_{\beta_{n,0}}^{1/2} \overset{\beta_{n,0} + h/\sqrt{n}}{\rightsquigarrow} \mathcal{N}(0, 1),$$

*where*

$$V_{\beta_{n,0}} \geq \dot{g}(\beta_{n,0})^T I_{\beta_{n,0}}^{-1} \dot{g}(\beta_{n,0}) + o(1).$$

The corollary implies that asymptotic efficiency is attained by an estimator which is asymptotically linear with an influence function  $l_\beta = \dot{g}(\beta)^T I_\beta^{-1} s_\beta$ , provided that it satisfies the condition (22).

We have already shown how these results can be applied to the linear regression setting in Section 8.5. We remark that the result of Theorem 9 is not directly applicable to Gaussian graphical models, where the unknown parameter has overall sparsity  $ps$ , where  $s = o(\sqrt{n}/\log p)$ .

REMARK 2. *The sparsity condition  $d_n = o(n^{1/3})$  arises when considering Taylor expansion of the log-likelihood for general models. Hence, when there is some special structure in the log-likelihood function, weaker sparsity conditions might be possible. For instance, for linear regression setting, the Hessian of the log-likelihood does not depend on the unknown parameter  $\beta_0$ , hence in that case by inspection of the likelihood expansion in the proof of Theorem 9, we see that the condition  $d_n = o(\sqrt{n/\log p})$  is sufficient.*

**11. Conclusions.** In this paper we have proposed a framework for studying asymptotic efficiency in high-dimensional models. We adopted a semi-parametric point of view: we concentrated on one dimensional functions of a high-dimensional parameter for which the lower bounds were derived. The semi-parametric efficiency bounds we obtained correspond to the efficiency bounds for parametric models. However, the treatment for high-dimensional models required more elaborate analysis due to the models changing with  $n$  and assumed sparsity of the model.

We further considered construction of estimators attaining the lower bounds. We showed that indeed construction of asymptotically efficient estimator is possible: a de-sparsified estimator in linear regression and Gaussian graphical models is asymptotically efficient for estimation of certain simple functionals. Our analysis identified the theoretical conditions on the parameter sparsity and further conditions on the model under which asymptotic efficiency may be shown.

*Comparison of the two approaches.* The analysis was done in two ways: in the spirit of asymptotic Cramér-Rao bounds and Le Cam’s bounds (van der Vaart (2000)). These are strongly related: both define a restricted set of estimators which are in some sense asymptotically unbiased and claim lower bounds for any estimator in this class.

However, the two lines of work are not directly comparable as they are different results under different assumptions. Le Cam’s bounds give a lower bound on *asymptotic variance*, while the Cramér-Rao bounds give a bound on the *variance* of an estimator. We formulated Le Cam’s approach for a general sparse model, while the Cramér-Rao bounds were only considered for the linear regression and Gaussian graphical models. Apart from this, the main results arising from the two approaches also present some differences in the assumptions. For the Le Cam’s-type results, we assumed a stronger sparsity condition of order  $d_n = o(n^{1/3}/\log p)$  because of the Taylor expansion of the likelihood. However, for the linear regression setting, the sparsity condition can be improved to  $d_n = o(\sqrt{n}/\log p)$ , which is the same as in the Cramér-Rao bounds. For Gaussian graphical models, Le Cam’s approach

as formulated in this paper cannot be directly used, unlike the approach through the Cramér-Rao bounds.

*Extensions.* Our results on upper bounds are presented for the case when the parameter of interest is a single entry of the high-dimensional parameter or a linear combination with e.g. bounded  $\ell_1$ -norm. It is interesting to note some relations to literature on minimax rates. One question is whether asymptotic efficiency can be attained e.g. for estimation of linear functionals in linear regression when the linear combination  $\xi$  is sparse. Our results needed that  $\|\xi\|_1$  remains bounded. Some recent works on high-dimensional models further consider estimation of more complicated, non-sparse functionals (in linear regression [Cai and Guo \(2015\)](#), for Gaussian sequence models [Collier, Comminges and Tsybakov \(2015\)](#)). These results are however of a different nature. Consider for instance estimation of  $\sum_{i=1}^p \beta_i$  in high-dimensional linear regression. In this case, the parametric rate cannot be achieved ([Cai and Guo \(2015\)](#)) and thus it remains unclear what can be said about “asymptotic efficiency”.

Furthermore, we have treated the case of a one-dimensional parameter of interest, though the analysis might be extended to settings when the parameter of interest is higher-dimensional (of a fixed dimension). Finally, our analysis considered particular examples of de-sparsified estimators, however, other estimators which are in some sense equivalent to these de-sparsified estimators are applicable.

## SUPPLEMENTARY MATERIAL

### Supplement to “Semi-parametric efficiency bounds for high-dimensional models”

(doi: [10.1214/00-AOASXXXXSUPP](https://doi.org/10.1214/00-AOASXXXXSUPP); .pdf). The supplementary material contains proofs.

### References.

- BELLEÇ, P. C. and TSYBAKOV, A. B. (2016). Bounds on the prediction error of penalized least squares estimators with convex penalty. *ArXiv: 1609.06675*.
- BICKEL, P. J., KLAASSEN, C. A., RITOV, Y. and WELLNER, J. A. (1993). *Efficient and adaptive estimation for semiparametric models*. Springer.
- BÜHLMANN, P. and VAN DE GEER, S. (2011). *Statistics for High-Dimensional Data*. Springer.
- CAI, T. and GUO, Z. (2015). Confidence Intervals for High-Dimensional Linear Regression: Minimax Rates and Adaptivity. *ArXiv: 1506.05539*.
- CHERNOZHUKOV, V., HANSEN, C. and SPINDLER, M. (2015). *Valid Post-Selection and Post-Regularization Inference: An Elementary, General Approach*. Annual Review of Economics.

- COLLIER, O., COMMINGES, L. and TSYBAKOV, A. B. (2015). Minimax estimation of linear and quadratic functionals on sparsity classes. *ArXiv: 1502.00665*.
- FRIEDMAN, J., HASTIE, T. and TIBSHIRANI, R. (2008). Sparse inverse covariance estimation with the graphical lasso. *Biostatistics* **9** 432–441.
- GAO, C., MA, Z. and ZHOU, H. H. (2014). Sparse CCA: Adaptive Estimation and Computational Barriers. *ArXiv: 1409.8565*.
- JANKOVÁ, J. and VAN DE GEER, S. (2014). Confidence intervals for high-dimensional inverse covariance estimation. *Electronic Journal of Statistics* **9** 1205–1229.
- JANKOVÁ, J. and VAN DE GEER, S. (2016a). Honest confidence regions and optimality for high-dimensional precision matrix estimation. *TEST*.
- JANKOVÁ, J. and VAN DE GEER, S. (2016b). Supplement to “Semi-parametric efficiency bounds for high-dimensional models”.
- JAVANMARD, A. and MONTANARI, A. (2014a). Confidence intervals and hypothesis testing for high-dimensional regression. *The Journal of Machine Learning Research* **15** 2869–2909.
- JAVANMARD, A. and MONTANARI, A. (2014b). Hypothesis Testing in High-Dimensional Regression under the Gaussian Random Design Model: Asymptotic Theory. *IEEE Trans. on Inform. Theory* **60** 6522 – 6554.
- JAVANMARD, A. and MONTANARI, A. (2015). De-biasing the lasso: Optimal sample size for gaussian designs. *Arxiv: 1508.02757*.
- KNIGHT, K. and FU, W. (2000). Asymptotics for lasso-type estimators. *Annals of Statistics* **28** 1356–1378.
- MEINSHAUSEN, N. and BÜHLMANN, P. (2006). High-dimensional graphs and variable selection with the Lasso. *Annals of Statistics* **34** 1436–1462.
- MEINSHAUSEN, N. and YU, B. (2009). Lasso-type recovery of sparse representations for high-dimensional data. *Annals of statistics* **37** 246–270.
- REN, Z., SUN, T., ZHANG, C. H. and ZHOU, H. H. (2015). Asymptotic normality and optimalities in estimation of large Gaussian graphical model. *Annals of Statistics* **43** 991–1026.
- VAN DE GEER, S., BÜHLMANN, P., RITOV, Y. and DEZEURE, R. (2014). On asymptotically optimal confidence regions and tests for high-dimensional models. *Annals of Statistics* **42** 1166–1202.
- VAN DER VAART, A. (2000). Asymptotic Statistics. *Cambridge University Press*.
- ZHANG, C. H. and ZHANG, S. S. (2014). Confidence intervals for low-dimensional parameters in high-dimensional linear models. *Journal of the Royal Statistical Society: Series B* **76** 217–242.

SEMINAR FÜR STATISTIK  
 ETH ZÜRICH  
 RÄMISTRASSE 101  
 8092 ZÜRICH  
 SWITZERLAND  
 E-MAIL: [jankova@stat.math.ethz.ch](mailto:jankova@stat.math.ethz.ch)  
 E-MAIL: [geer@stat.math.ethz.ch](mailto:geer@stat.math.ethz.ch)

# SUPPLEMENT TO “SEMI-PARAMETRIC EFFICIENCY BOUNDS FOR HIGH-DIMENSIONAL MODELS”

BY JANA JANKOVÁ AND SARA VAN DE GEER

This supplement contains the proofs. Section 12 summarizes some preliminary material on concentration of measure. Section 13 contains the proofs of Section 7: Strong oracle inequalities for the Lasso. Section 14 contains strong oracle inequalities for the nodewise Lasso. Section 15 contains proofs for Section 8. Section 16 contains proofs for Section 9: Gaussian graphical models. In Section 17 we give the proofs for Section 10: Le Cam’s bounds for general models. Proofs for Section 14 are deferred to Appendix A and some technical lemmas are deferred to Appendices B and C.

**12. Concentration inequalities for sub-exponential random variables.** In this preliminary section, we recall some results on concentration results for sub-exponential random variables (for the definition of a sub-exponential random variable, see Section 14.2.1 in Bühlmann and van de Geer (2011)). Lemma 4 below is a version of Lemma 14.13 in Bühlmann and van de Geer (2011).

LEMMA 4. *Let  $Z_1, \dots, Z_n$  be independent random variables with values in some (measurable) space  $\mathcal{Z}$  and  $\gamma_1, \dots, \gamma_p$  be real-valued functions on  $\mathcal{Z}$  satisfying, for  $j = 1, \dots, p$ ,*

$$\mathbb{E}\gamma_j(Z_i) = 0, \quad \mathbb{E}e^{|\gamma_j(Z_i)|/K} \leq M_1, \quad \forall i = 1, \dots, n,$$

*where  $M_1 > 0$  is a universal constant and  $K > 0$ .*

*Then there exists a universal constant  $M_2$  such that for all  $t > 0$  we have with probability at least  $1 - e^{-nt}$  that*

$$\max_{j=1, \dots, p} \left| \frac{1}{n} \sum_{i=1}^n \gamma_j(Z_i) \right| \leq M_2 K t + \sqrt{2t} + \sqrt{\frac{2 \log(2p)}{n}} + \frac{M_2 K \log(2p)}{n}.$$

The following lemma is a version of Corollary 14.1 in Bühlmann and van de Geer (2011).

LEMMA 5. *Assume the conditions of Lemma 4. Then for all  $m = 1, 2, \dots$  it holds*

$$\begin{aligned} & \mathbb{E} \left( \max_{j=1, \dots, p} \left| \frac{1}{n} \sum_{i=1}^n \gamma_j(Z_i) \right|^m \right) \\ & \leq \left( \sqrt{\frac{2 \log(2p + e^{m-1} - p)}{n}} + \frac{M_2 K \log(2p + e^{m-1} - p)}{n} \right)^m, \end{aligned}$$

where  $M_2 > 0$  is a universal constant.

### 13. Proofs for Section 7: Strong oracle inequalities for the Lasso.

In this section we prove the oracle inequality for the Lasso as stated in Theorem 1. We need the following preliminary Lemmas 6, 7 and 8. Lemma 6 below gives sufficient conditions under which the compatibility condition is satisfied. Lemma 7 is a concentration result for sub-Gaussian random variables as in Section 14 in Bühlmann and van de Geer (2011). Lemma 8 is a version of Theorem 6.1 in Bühlmann and van de Geer (2011). Recall that we denote  $\hat{\Sigma} := X^T X/n$  and  $\Sigma_0 := \mathbb{E} X^T X/n$ . We recall the definition of the compatibility condition (see Section 6.13 in Bühlmann and van de Geer (2011)). Let  $S := \{i : \beta_i \neq 0\}$  and let  $s = |S|$ . We denote by  $\beta_S$  the vector obtained from the vector  $\beta \in \mathbb{R}^p$  by replacing entries corresponding to the indices in  $S$  by zeros.

DEFINITION 4. *We say that a matrix  $\Sigma_0$  satisfies the compatibility condition with a constant  $\phi$  if*

$$\phi := \min \left\{ \frac{s \beta^T \Sigma_0 \beta}{\|\beta_S\|_1^2} : \|\beta_{S^c}\|_1 \leq 3 \|\beta_S\|_1 \right\} > 0.$$

LEMMA 6 (Corollary 6.8 in Bühlmann and van de Geer (2011)). *Suppose that  $\Lambda_{\min}(\Sigma_0) \geq L$  for a universal constant  $L > 0$ . Then  $\Sigma_0$  satisfies the compatibility condition with the constant  $L$ . Further suppose that  $s\lambda = o(1)$ . Then on the set  $\|\hat{\Sigma} - \Sigma_0\|_\infty \leq \lambda$ , for all  $n$  sufficiently large,  $\hat{\Sigma}$  satisfies the compatibility condition with the constant  $L/2$ .*

LEMMA 7. *Suppose that  $\epsilon_i, i = 1, \dots, n$  are sub-Gaussian random variables with a universal constant  $K_1$  and that  $X^{(i)}, i = 1, \dots, n$  are independent random vectors with sub-Gaussian entries, with a universal constant  $K_2$ . Suppose that  $\epsilon_i$  and  $X^{(i)}$  are independent for  $i = 1, \dots, n$  and  $\log p/n = o(1)$ . Then there exists a constant  $c_1$  such that for all  $\tau > 1$*

$$P \left( \|\epsilon^T X\|_\infty / n \geq c_1 \tau \sqrt{\frac{\log(2p)}{n}} \right) \leq (2p)^{-\tau^2}.$$

PROOF. We apply Lemma 4 with  $\gamma_j(Z_i) = \epsilon_i X_{ij}$  for  $i = 1, \dots, n$  and  $j = 1, \dots, p$ . Then  $\mathbb{E}\epsilon_i X_{ij} = \mathbb{E}X_{ij}\mathbb{E}(\epsilon_i|X_{ij}) = 0$ , where we used independence of  $\epsilon_i$  and  $X^{(i)}$ . By sub-Gaussianity of  $\epsilon_i$  and  $X_{ij}$  and by the Cauchy-Schwarz inequality it follows

$$\begin{aligned} \mathbb{E}e^{|\epsilon_i X_{ij}|/\max\{K_1, K_2\}} &\leq \mathbb{E}e^{|\epsilon_i|^2/(2\max\{K_1, K_2\}^2) + |X_{ij}|^2/(2\max\{K_1, K_2\}^2)} \\ &\leq (\mathbb{E}e^{|\epsilon_i|^2/\max\{K_1, K_2\}^2})^{1/2} (\mathbb{E}e^{|X_{ij}|^2/\max\{K_1, K_2\}^2})^{1/2} \\ &= \mathcal{O}(1). \end{aligned}$$

Consequently, by Lemma 4 and since  $\log p/n = o(1)$ , there exists a constant  $c_1$  such that for all  $\tau > 1$

$$P\left(\max_{j=1, \dots, p} \left| \frac{1}{n} \sum_{i=1}^n \epsilon_i X_{ij} \right| \geq c_1 \tau \sqrt{\frac{\log(2p)}{n}}\right) \leq (2p)^{-\tau^2}.$$

□

Finally, we give an oracle inequality for the Lasso. The proof may be found in [Bühlmann and van de Geer \(2011\)](#).

LEMMA 8 (a version of Theorem 6.1 in [Bühlmann and van de Geer \(2011\)](#)). *Consider the Lasso estimator  $\hat{\beta}$  defined in (5) with a tuning parameter  $\lambda \geq 2\lambda_0$ . Suppose that  $s\lambda = o(1)$  and that  $\Lambda_{\min}(\Sigma_0) \geq L$  for some universal constant  $L > 0$ . Then on the set*

$$\mathcal{T} := \{\|\epsilon^T X\|_\infty/n \leq \lambda_0, \|\hat{\Sigma} - \Sigma_0\|_\infty \leq \lambda_0\}$$

it holds

$$\|\hat{\beta} - \beta_0\|_1 \leq 16\lambda s/L.$$

We are now ready to prove Theorem 1.

PROOF OF THEOREM 1. First we summarize the oracle inequality for the Lasso which holds with high probability. Let  $\mathcal{T}_1 := \{\|\epsilon^T X\|_\infty/n \leq \tau c_1 \sqrt{\log p/n}\}$  for some  $\tau > 1$  and for some suitable constant  $c_1 > 0$ . By Lemma 7, for the complementary set  $\mathcal{T}_1^c$  it holds that  $P_{\beta_0}(\mathcal{T}_1^c) \leq (2p)^{-\tau^2}$ . Let  $\mathcal{T}_2 := \{\|\hat{\Sigma} - \Sigma_0\|_\infty \leq \tau c_1 \sqrt{\log p/n}\}$ . Then by Lemma 4, taking  $\gamma_{i,j}(X^{(k)}) = e_i^T (X^{(k)}(X^{(k)})^T - \Sigma_0)e_j$  for  $k = 1, \dots, n$  and  $i, j = 1, \dots, p$ , it follows that  $P_{\beta_0}(\mathcal{T}_2^c) \leq (2p)^{-\tau^2}$ . Denote  $\mathcal{T} := \mathcal{T}_1 \cap \mathcal{T}_2$ ; then  $P_{\beta_0}(\mathcal{T}^c) \leq 2(2p)^{-\tau^2}$ . By Lemma 8, when  $\lambda \geq 2\lambda_0 := 2\tau c_1 \sqrt{\log p/n}$ , on the set  $\mathcal{T}$  it holds that  $\|\hat{\beta} - \beta_0\|_1 \leq 16\lambda s/L$ .



We now proceed to show that the oracle inequality for the Lasso holds also in expectation. The definition of  $\hat{\beta}$  gives

$$\|Y - X\hat{\beta}\|_n^2 + \lambda\|\hat{\beta}\|_1 \leq \|\epsilon\|_n^2 + \lambda\|\beta_0\|_1.$$

Consequently,

$$\|\hat{\beta}\|_1 \leq \|\epsilon\|_n^2/\lambda + \|\beta_0\|_1.$$

Then, and by the triangle inequality

$$\|\hat{\beta} - \beta_0\|_1 \leq \|\hat{\beta}\|_1 + \|\beta_0\|_1 \leq \|\epsilon\|_n^2/\lambda + 2\|\beta_0\|_1,$$

and thus for any  $k \in \{1, 2, \dots\}$

$$\mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1^k \leq \mathbb{E}_{\beta_0} (\|\epsilon\|_n^2/\lambda + 2\|\beta_0\|_1)^k.$$

Then by the inequality  $|x + y|^k \leq 2^{k-1}(|x|^k + |y|^k)$ ,  $k \geq 1$ , it follows

$$\mathbb{E}_{\beta_0} (\|\epsilon\|_n^2/\lambda + 2\|\beta_0\|_1)^k \leq \mathbb{E}_{\beta_0} 2^{k-1} \left( (\|\epsilon\|_n^2/\lambda)^k + (2\|\beta_0\|_1)^k \right)$$

By assumption, the random variables  $\epsilon_1, \dots, \epsilon_n$  are sub-Gaussian with universal constants. This also implies that  $\frac{1}{n} \sum_{i=1}^n \text{var}(\epsilon_i^2) = \mathcal{O}(1)$ . Hence by Lemma 5 applied with  $\gamma_j(Z_i) := \epsilon_i - \text{var}(\epsilon_i)$  ( $j = 1, i = 1, \dots, n$ ) we have

$$\mathbb{E}_{\beta_0} (\|\epsilon\|_n^2)^k = \mathcal{O} \left( \left( \frac{1}{n} \sum_{i=1}^n \text{var}(\epsilon_i^2) \right)^k \right) = \mathcal{O}(1).$$

Next observe that by assumption we have  $\|\beta_0\|_2 = \mathcal{O}(1)$  and hence

$$\|\beta_0\|_1^k \leq (\sqrt{s}\|\beta_0\|_2)^k \leq \mathcal{O}(s^{k/2}).$$

We can thus conclude that

$$\begin{aligned} \mathbb{E}_{\beta_0} (\|\epsilon\|_n^2/\lambda + 2\|\beta_0\|_1)^k &\leq 2^{k-1} \left( \mathbb{E}_{\beta_0} (\|\epsilon\|_n^2/\lambda)^k + \mathbb{E}_{\beta_0} (2\|\beta_0\|_1)^k \right) \\ &\leq \mathcal{O}(s^{k/2} \lambda^{-k}). \end{aligned}$$

Hence we obtain a rough bound

$$(\mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1^k)^{1/k} = \mathcal{O}(s^{1/2} \lambda^{-1}).$$

On the set  $\mathcal{T}$  we have the oracle bound  $\|\hat{\beta} - \beta_0\|_1 = \mathcal{O}(s\lambda)$  and thus on the set  $\mathcal{T}$ ,  $\|\hat{\beta} - \beta_0\|_1^k = \mathcal{O}(s^k \lambda^k)$ . Otherwise (so also on the set  $\mathcal{T}^c$ ) we have the rough bound  $\mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1^k = \mathcal{O}(s^{k/2} \lambda^{-k})$ . Denote by  $1_A$  the indicator

function of a set  $A$ . Then it follows using the bounds holding on  $\mathcal{T}$  and  $\mathcal{T}^c$  and by the Cauchy-Schwarz inequality

$$\begin{aligned}
\mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1^k &= \mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1^k 1_{\mathcal{T}} + \mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1^k 1_{\mathcal{T}^c} \\
&\leq \mathcal{O}(s^k \lambda^k) + \sqrt{\mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1^{2k}} \sqrt{\mathbb{E}_{\beta_0} 1_{\mathcal{T}^c}} \\
&= \mathcal{O}(s^k \lambda^k) + \mathcal{O}(\sqrt{s^k \lambda^{-2k}}) \sqrt{P_{\beta_0}(\mathcal{T}^c)} \\
&\leq \mathcal{O}(s^k \lambda^k) + \mathcal{O}(s^{k/2} \lambda^{-k}) \sqrt{2} (2p)^{-\tau^2/2} \\
&= \mathcal{O}(s^k \lambda^k),
\end{aligned}$$

where we used the assumption  $\tau^2 \geq \frac{2k \log(\sqrt{s} \lambda^2)}{\log p}$  which implies

$$\mathcal{O}(s^{k/2} \lambda^{-k}) p^{-\tau^2/2} = \mathcal{O}(s^k \lambda^k).$$

Hence we conclude that there exists a constant  $C_1$  such that

$$(24) \quad (\mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1^k)^{1/k} = C_1 s \lambda.$$

□

**14. Strong oracle inequalities for the nodewise Lasso.** In this section, we derive a strong oracle inequality for the nodewise regression estimator  $\hat{\Theta}$  of the inverse covariance matrix  $\Theta_0$  defined in (7). These results will be needed to show strong asymptotic unbiasedness and upper bounds on the variance of the de-sparsified Lasso. The proofs may be found in Appendix A. We require a sparsity condition on the inverse covariance matrix of the covariates. To this end, denote the sparsity of the  $j$ -th row/column of the matrix  $\Theta_0$  by  $s_j$ , i.e.

$$s_j = \|\Theta_j^0\|_0.$$

First we remark that the paper [van de Geer et al. \(2014\)](#) shows that under conditions (A1), (A2\*) and  $s_j = o(n/\log p)$  it holds  $\|\hat{\Theta}_j - \Theta_j^0\|_1 = \mathcal{O}_P(s_j \lambda_j)$ . We aim to show a stronger claim,  $\mathbb{E} \|\hat{\Theta}_j - \Theta_j^0\|_1 = \mathcal{O}(s_j \lambda_j)$ . This is a more difficult task than for the linear regression, since one has to make sure that the estimate of one over the noise level,  $1/\hat{\tau}_j^2$ , does not blow up in expectation. Moreover, for further results we shall need not only the result for each  $j = 1, \dots, p$  but actually for the maximum over  $j = 1, \dots, p$ , that is, an oracle bound for  $\mathbb{E} \max_{j=1, \dots, p} \|\hat{\Theta}_j - \Theta_j^0\|_1^k$ , where  $k \in \{1, 2, \dots\}$ . We introduce further notation; we let

$$\gamma_j^0 := \arg \min_{\gamma \in \mathbb{R}^{p-1}} \mathbb{E} \|X_j - X_{-j} \gamma\|_n^2$$

and

$$\tau_j^2 := \mathbb{E} \|X_j - X_{-j}\gamma_0\|_n^2.$$

The following lemma is similar to Theorem 1 for the mean  $\ell_1$ -error of the Lasso, however, we consider the  $p$  Lasso estimators obtained from the node-wise regression and derive an upper bound on the maximum mean  $\ell_1$ -error, where the maximum is taken over the  $p$  estimators.

LEMMA 9. *Assume that condition (A2\*) is satisfied, let  $k \in \{1, 2, \dots\}$  be fixed and assume that  $\max_{j=1, \dots, p} s_j \sqrt{\log p/n} = o(1)$ ,  $j = 1, \dots, p$ . Let  $\hat{\gamma}_j$  be defined as in (6) with tuning parameters  $\lambda_j = c\sqrt{\log p/n}$ ,  $j = 1, \dots, p$ , for some sufficiently large constant  $c > 0$ . Then it holds that*

$$[\mathbb{E} \max_{j=1, \dots, p} \|\hat{\gamma}_j - \gamma_j^0\|_1^k]^{1/k} = \mathcal{O}(\max_{j=1, \dots, p} s_j \lambda_j).$$

The following lemma shows that the noise estimator  $\hat{\tau}_j^2$  is a near-oracle estimator of  $\tau_j^2 = 1/\Theta_{jj}^0$ , and  $1/\hat{\tau}_j^2$  is a near-oracle estimator of  $1/\tau_j^2$ .

LEMMA 10. *Assume that condition (A2\*) is satisfied,  $\|\Sigma_0\|_\infty = \mathcal{O}(1)$ ,  $\max_{j=1, \dots, p} s_j \sqrt{\log p/n} = o(1)$  and let  $k \in \{1, 2, \dots\}$  be fixed. Let  $\hat{\gamma}_j$ ,  $j = 1, \dots, p$  be defined as in (6) with tuning parameters  $\lambda_j = c\sqrt{\log p/n}$ ,  $j = 1, \dots, p$ , for some sufficiently large constant  $c > 0$ . Then the following statements hold*

- 1)  $[\mathbb{E} \max_{j=1, \dots, p} |\hat{\tau}_j^2 - \tau_j^2|^k]^{1/k} = \mathcal{O}(\max_{j=1, \dots, p} \sqrt{s_j} \lambda_j),$
- 2)  $[\mathbb{E} \max_{j=1, \dots, p} |\frac{1}{\hat{\tau}_j^2} - \frac{1}{\tau_j^2}|^k]^{1/k} = \mathcal{O}(\max_{j=1, \dots, p} \sqrt{s_j} \lambda_j).$

Combination of the results in Lemmas 9 and 10 gives the following result for the mean  $\ell_1$ -error of the nodewise regression estimator  $\hat{\Theta}$  defined in (7).

LEMMA 11. *Assume that condition (A2\*) is satisfied,  $\|\Sigma_0\|_\infty = \mathcal{O}(1)$ ,  $\max_{j=1, \dots, p} s_j \sqrt{\log p/n} = o(1)$  and let  $k \in \{1, 2, \dots\}$  be fixed. Then for  $\hat{\Theta}_j$ ,  $j = 1, \dots, p$  defined in (7) with tuning parameters  $\lambda_j = c\sqrt{\log p/n}$ ,  $j = 1, \dots, p$ , for some sufficiently large constant  $c > 0$ , it holds*

$$[\mathbb{E} \max_{j=1, \dots, p} \|\hat{\Theta}_j - \Theta_j^0\|_1^k]^{1/k} = \mathcal{O}(\max_{j=1, \dots, p} s_j \lambda_j).$$

We note that the statements of Lemmas 9, 10, 11 would also hold for single estimators for some fixed  $j$  (without taking the maximum over  $j$ ). Then the bounds would only depend on the tuning parameter for that particular estimator and on the sparsity  $s_j$ . We also give the following lemma for on the sparsity in the nodewise Lasso estimator. It shows that under certain conditions, the estimator  $\hat{\gamma}_j$  has sparsity of order  $s_j$  with high probability.

LEMMA 12. Assume that condition (A2\*) is satisfied,  $\max_j s_j \sqrt{\log p/n} = o(1)$ ,  $\Lambda_{\max}(\Sigma_0) = \mathcal{O}(1)$ . Let  $\hat{\gamma}_j$  be defined as in (6) with a tuning parameter  $\lambda_j = c\sqrt{\log p/n}$  for some sufficiently large constant  $c > 0$ . Then it holds that

$$\|\hat{\gamma}_j\|_0 = \mathcal{O}_P(s_j).$$

## 15. Proofs for Section 8.2: The de-sparsified Lasso.

15.1. *Proofs for Section 8.2: Strong asymptotic unbiasedness of the de-sparsified Lasso.*

PROOF OF LEMMA 1. For the de-sparsified estimator  $\hat{b}_\xi$  we have by simple algebra the equality

$$\hat{b}_\xi - \xi^T \beta_0 = \xi^T \hat{\Theta} X^T \epsilon/n + \xi^T (\hat{\Sigma} \hat{\Theta} - I)^T (\hat{\beta} - \beta_0).$$

Consider any  $\beta_0 \in \mathcal{B}(d_n)$ . First note that

$$\mathbb{E}_{\beta_0} \xi^T \hat{\Theta}^T X^T \epsilon/n = \mathbb{E} \mathbb{E}_{\beta_0} (\xi^T \hat{\Theta}^T X^T \epsilon/n | X) = \mathbb{E} \xi^T \hat{\Theta}^T X^T \mathbb{E}_{\beta_0} (\epsilon | X)/n = 0.$$

We then have by the definition of  $\hat{b}_\xi$ , by the Hölder's inequality and the Cauchy-Schwarz inequality

$$\begin{aligned} \mathbb{E}_{\beta_0} (\hat{b}_\xi - \xi^T \beta_0) &= \underbrace{\mathbb{E}_{\beta_0} \xi^T \hat{\Theta} X^T \epsilon/n}_{=0} + \mathbb{E}_{\beta_0} \xi^T (\hat{\Sigma} \hat{\Theta} - I)^T (\hat{\beta} - \beta_0) \\ &\leq \mathbb{E}_{\beta_0} \|\xi\|_1 \|\hat{\Sigma} \hat{\Theta} - I\|_\infty \|\hat{\beta} - \beta_0\|_1 \\ &\leq \|\xi\|_1 (\mathbb{E}_{\beta_0} \|\hat{\Sigma} \hat{\Theta} - I\|_\infty^2)^{1/2} (\mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1^2)^{1/2}. \end{aligned}$$

By the Karush-Kuhn-Tucker conditions for  $\hat{\gamma}_j, j = 1, \dots, p$  (see [van de Geer et al. \(2014\)](#)), we have

$$\|\hat{\Sigma} \hat{\Theta} - I\|_\infty \leq \max_{j=1, \dots, p} \lambda_j / \hat{\tau}_j^2.$$

By Lemma 10 from Section 14 it follows that

$$(\mathbb{E} \|\hat{\Sigma} \hat{\Theta} - I\|_\infty^2)^{1/2} \leq \max_{j=1, \dots, p} \lambda_j (\mathbb{E}_{\beta_0} \max_{j=1, \dots, p} 1/(\hat{\tau}_j^2)^2)^{1/2} = \mathcal{O}(\max_{j=1, \dots, p} \lambda_j).$$

Next we apply Theorem 1. Conditions  $\|\beta_0\|_0 \leq d_n$ ,  $\|\beta_0\|_2 = \mathcal{O}(1)$ , (A1), (A2\*) and sparsity  $d_n = o(\sqrt{n}/\log p)$  imply that conditions of Theorem 1 are satisfied. Hence  $(\mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1^2)^{1/2} = \mathcal{O}(s\lambda)$ . Hence, and since  $\|\xi\|_1 = \mathcal{O}(1)$ , and using the last display we obtain that

$$\mathbb{E}_{\beta_0} (\hat{b}_\xi - \xi^T \beta_0) = \mathcal{O}(s\lambda \max_{j=1, \dots, p} \lambda_j) = o(1/\sqrt{n}),$$

where we used the sparsity condition  $s \leq d_n = o(\sqrt{n}/\log p)$ . Thus we have shown  $\sqrt{n}(\mathbb{E}_{\beta_0}(\hat{b}_\xi - \xi^T \beta_0)) = o(1)$ . But then there exists  $\delta_n \rightarrow 0$  such that  $\sqrt{n/\delta_n}(\mathbb{E}_{\beta_0}(\hat{b}_\xi - \xi^T \beta_0)) = o(1)$  (take e.g.  $\delta_n := \sqrt{\sqrt{n}(\mathbb{E}_{\beta_0}(\hat{b}_\xi - \xi^T \beta_0))}$ ) and hence the estimator  $\hat{b}_\xi$  is strongly asymptotically unbiased with a rate  $m_n := n/\delta_n$ .  $\square$

15.2. *Proofs for Section 8.3: Main results for random design.* Before proving the statement of Theorem 2, we need auxiliary Lemmas 13, 14 and 15. Throughout this section, we denote by  $\phi$  the probability density function of a standard normal random variable.

LEMMA 13. *Let  $Z \sim \mathcal{N}(0, 1)$ . Then for all  $t \in \mathbb{R}$*

$$\mathbb{E} \left[ e^{tZ - t^2/2} - 1 - tZ \right]^2 = e^{t^2} - 1 - t^2.$$

Moreover, for  $2t^2 < 1$  we have

$$\mathbb{E} e^{t^2 Z^2} = \frac{1}{\sqrt{1 - 2t^2}}.$$

PROOF OF LEMMA 13. By direct calculation

$$\mathbb{E} \left[ e^{tZ - t^2/2} \right]^2 = \mathbb{E} e^{2tZ - t^2} = e^{t^2},$$

$$\mathbb{E} e^{tZ - t^2/2} = 1$$

and

$$\mathbb{E} Z e^{tZ - t^2/2} = t \mathbb{E} e^{tZ - t^2/2} = t.$$

The first result of the lemma follows immediately. The second result is also easily found by standard calculations:

$$\mathbb{E} e^{t^2 Z^2} = \int e^{t^2 z^2} \phi(z) dz = \int \phi(z \sqrt{1 - 2t^2}) dz = \frac{1}{\sqrt{1 - 2t^2}}.$$

$\square$

LEMMA 14. *Suppose that  $u \in \mathbb{R}^p$  satisfies  $2u^T \Sigma_0 u < 1$ . Let  $Z = (X, Y)$ , where  $Y = X\beta + \epsilon$ ,  $\epsilon \sim \mathcal{N}(0, I)$  independent of  $X$ , and  $X \sim \mathcal{N}(0, \Sigma_0)$ . Denote the corresponding probability density of  $Z$  by  $p_\beta$  and let  $s_{\beta_0}(Z) := X^T \epsilon$ . Then it holds*

$$\mathbb{E}_{\beta_0} \left( \frac{p_{\beta_0+u}(Z) - p_{\beta_0}(Z)}{p_{\beta_0}(Z)} - s_{\beta_0}(Z)^T u \right)^2 = (1 - 2u^T \Sigma_0 u)^{-n/2} - 1 - nu^T \Sigma_0 u.$$

PROOF OF LEMMA 14. Denote the density of  $Y$  given  $X$  by  $p_{\beta_0}(\cdot|X)$ , i.e. for  $y = (y_1, \dots, y_n)$

$$p_{\beta_0}(y|X) := \prod_{i=1}^n \phi(y_i - (X^{(i)})^T \beta_0) = \frac{1}{(2\pi)^{n/2}} e^{-(y-X\beta_0)^T(y-X\beta_0)/2},$$

where  $\phi$  is the standard normal density.

Given  $X$ , the random variable  $\epsilon^T Xu$  is  $\mathcal{N}(0, nu^T \hat{\Sigma} u)$ -distributed. It follows therefore from the first result of Lemma 13 with  $t^2 = nu^T \hat{\Sigma} u$  that

$$\begin{aligned} & \mathbb{E}_{\beta_0} \left[ \left( \frac{p_{\beta_0+u}(Y - Xu|X) - p_{\beta_0}(Y|X)}{p_{\beta_0}(Y|X)} - s_{\beta_0}(Z)^T u \right)^2 \middle| X \right] \\ &= \mathbb{E} e^{nu^T \hat{\Sigma} u} - 1 - nu^T \hat{\Sigma} u. \end{aligned}$$

Since  $(X^{(i)})^T u \sim \mathcal{N}(0, u^T \Sigma_0 u)$  for  $i = 1, \dots, n$ , we have by the second result of Lemma 13

$$\mathbb{E}_{\beta_0} e^{((X^{(i)})^T u)^2} = \frac{1}{\sqrt{1 - 2u^T \Sigma_0 u}}.$$

Hence

$$\mathbb{E}_{\beta_0} [e^{nu^T \hat{\Sigma} u} - 1 - nu^T \hat{\Sigma} u] = (1 - 2u^T \Sigma_0 u)^{-n/2} - 1 - nu^T \Sigma_0 u,$$

from which the result follows.  $\square$

LEMMA 15. Suppose that  $nu^T \Sigma_0 u = o(1)$ . Then

$$(1 - 2u^T \Sigma_0 u)^{-n/2} - 1 - nu^T \Sigma_0 u = o(nu^T \Sigma_0 u).$$

PROOF. Since  $nu^T \Sigma_0 u = o(1)$ , we can use the following Taylor expansions of log and exp

$$\begin{aligned} (1 - 2u^T \Sigma_0 u)^{-n/2} &= e^{-n \log(1 - 2u^T \Sigma_0 u)/2} = e^{nh^T \Sigma_0 u + o(nh^T \Sigma_0 u)} \\ &= 1 + nh^T \Sigma_0 h + o(nh^T \Sigma_0 h). \end{aligned}$$

Hence

$$(1 - 2u^T \Sigma_0 u)^{-n/2} - 1 - nu^T \Sigma_0 u = o(nu^T \Sigma_0 u).$$

$\square$

PROOF OF THEOREM 2. By assumption (10) on the differentiability of  $g$  and by strong asymptotic unbiasedness of  $T_n$  at  $\beta_0$  and  $\beta_0 + h/\sqrt{m_n}$ , it follows

$$\begin{aligned} h^T \dot{g}(\beta_0) &= \sqrt{m_n} (g(\beta_0 + h/\sqrt{m_n}) - g(\beta_0)) + o(1) \\ &= \sqrt{m_n} \left( \mathbb{E}_{\beta_0 + h/\sqrt{m_n}} T_n(Z) - \mathbb{E}_{\beta_0} T_n(Z) \right) + o(1), \end{aligned}$$

where  $Z := (X, Y)$ . We denote probability density corresponding to  $Z$  by  $p_\beta$ , i.e.

$$p_{\beta_0}(X, Y) := \frac{1}{(2\pi)^{n/2}} e^{-(Y-X\beta_0)^T(Y-X\beta_0)/2}.$$

Let  $s_{\beta_0}(Z) := X^T \epsilon$ . We may rewrite the expressions to obtain

$$\begin{aligned} & \sqrt{m_n} \left( \mathbb{E}_{\beta_0+h/\sqrt{m_n}} T_n(Z) - \mathbb{E}_{\beta_0} T_n(Z) \right) + o(1) \\ &= \sqrt{m_n} \int T_n(z) (p_{\beta_0+h/\sqrt{m_n}}(z) - p_{\beta_0}(z)) dz \\ &= \mathbb{E}_{\beta_0} T_n(Z) \frac{p_{\beta_0+h/\sqrt{m_n}}(Z) - p_{\beta_0}(Z)}{p_{\beta_0}(Z)/\sqrt{m_n}} \\ &= \mathbb{E}_{\beta_0} T_n(Z) \left( \frac{p_{\beta_0+h/\sqrt{m_n}}(Z) - p_{\beta_0}(Z)}{p_{\beta_0}(Z)/\sqrt{m_n}} - s_{\beta_0}(Z)^T h \right) \\ &\quad + \mathbb{E}_{\beta_0} T_n(Z) s_{\beta_0}(Z)^T h \\ &= \mathbb{E}_{\beta_0} (T_n(Z) - g(\beta_0)) \times \\ &\quad \left( \frac{p_{\beta_0+h/\sqrt{m_n}}(Z) - p_{\beta_0}(Z)}{p_{\beta_0}(Z)/\sqrt{m_n}} - s_{\beta_0}(Z)^T h \right) \\ &\quad + \mathbb{E}_{\beta_0} T_n(Z) s_{\beta_0}(Z)^T h, \end{aligned}$$

where we used that  $\mathbb{E}_{\beta_0} s_{\beta_0}(Z) = 0$ ,  $\int p_{\beta_0}(z) dz = 1$  and  $\int p_{\beta_0+h/\sqrt{m_n}}(z) dz = 1$ . Since the variance of  $T_n$  is  $\mathcal{O}(1/n)$  by the definition of strong asymptotic unbiasedness, then

$$\begin{aligned} \mathbb{E}_{\beta_0} (T_n(Z) - g(\beta_0))^2 &= \text{var}(T_n(Z)) + [\mathbb{E}_{\beta_0} (T_n(Z) - g(\beta_0))]^2 \\ &= \mathcal{O}(1/n) + o(1/n) = \mathcal{O}(1/n). \end{aligned}$$

By Lemmas 14 and 15 with  $u := h/\sqrt{m_n}$  and since  $h^T \Sigma_0 h = 1$ ,

$$\begin{aligned} \mathbb{E}_{\beta_0} \left( \frac{p_{\beta_0+h/\sqrt{m_n}}(Z) - p_{\beta_0}(Z)}{p_{\beta_0}(Z)} - s_{\beta_0}(Z)^T h/\sqrt{m_n} \right)^2 &= o(n/m_n h^T \Sigma_0 h) \\ &= o(n/m_n). \end{aligned}$$

Hence, multiplying by  $m_n$

$$\mathbb{E}_{\beta_0} \left( \frac{p_{\beta_0+h/\sqrt{m_n}}(Z) - p_{\beta_0}(Z)}{p_{\beta_0}(Z)/\sqrt{m_n}} - s_{\beta_0}(Z)^T h \right)^2 = o(n).$$

Consequently, and by the Cauchy-Schwarz inequality, we have the upper bound

$$\begin{aligned}
& \left| \mathbb{E}_{\beta_0}(T_n(Z) - g(\beta_0)) \left( \frac{p_{\beta_0+h/\sqrt{m_n}}(Z) - p_{\beta_0}(Z)}{p_{\beta_0}(Z)/\sqrt{m_n}} - s_{\beta_0}(Z)^T h \right) \right| \\
& \leq \sqrt{\mathbb{E}_{\beta_0}(T_n(Z) - g(\beta_0))^2} \sqrt{\mathbb{E} \left( \frac{p_{\beta_0+h/\sqrt{m_n}}(Z) - p_{\beta_0}(Z)}{p_{\beta_0}(Z)/\sqrt{m_n}} - s_{\beta_0}(Z)^T h \right)^2} \\
& = \mathcal{O}(1/\sqrt{n}) \mathcal{O}(\sqrt{n}) = o(1).
\end{aligned}$$

Next observe that by the triangle inequality we have

$$|h^T \dot{g}(\beta_0)| - |\text{cov}_{\beta_0}(T_n, \epsilon^T X h)| \leq |h^T \dot{g}(\beta_0) - \text{cov}_{\beta_0}(T_n, \epsilon^T X h)| = o(1),$$

and hence and by the Cauchy-Schwarz inequality it follows

$$\begin{aligned}
|h^T \dot{g}(\beta_0)| & \leq |\text{cov}_{\beta_0}(T_n, \epsilon^T X h)| + o(1) \\
& \leq \sqrt{\text{var}_{\beta_0}(T_n)} \sqrt{\text{var}_{\beta_0}(\epsilon^T X h)} + o(1). \\
& \leq \sqrt{\text{var}_{\beta_0}(T_n)} \sqrt{n} + o(1),
\end{aligned}$$

where we used that  $\text{var}_{\beta_0}(X^T \epsilon) = n \Sigma_0$  and  $h^T \Sigma_0 h = 1$ . By the strong asymptotic unbiasedness assumption on  $T_n$ , we have  $\text{var}_{\beta_0}(T_n) = \mathcal{O}(1/n)$  and thus taking squares of both sides of the last inequality we obtain

$$|h^T \dot{g}(\beta_0)|^2 \leq n \text{var}_{\beta_0}(T_n) + 2\sqrt{n} \sqrt{\text{var}_{\beta_0}(T_n)} o(1) + o(1) \leq n \text{var}_{\beta_0}(T_n) + o(1).$$

□

**PROOF OF THEOREM 3.** To obtain the lower bound on the variance, we apply Theorem 2 with  $h := \Theta_0 \xi / \sqrt{\xi^T \Theta_0 \xi}$  (note that for  $g(\beta) = \xi^T \beta$ , the condition on  $g$  is satisfied). Then by direct calculation, we see that the condition  $h^T \Sigma_0 h = 1$  is satisfied for this choice of  $h$ . Moreover, since  $\beta_0 + h/\sqrt{n} \in B(\beta_0, c/\sqrt{n})$  by assumption, then it implies that also  $\beta_0 + h/\sqrt{m_n} \in B(\beta_0, c/\sqrt{m_n})$ . Thus by Theorem 2 follows the lower bound  $\text{var}_{\beta_0}(T) \geq \frac{\xi^T \Theta_0 \xi + o(1)}{n}$ , for any strongly asymptotically unbiased estimator  $T$  of  $\xi^T \beta_0$  at  $\beta_0$  with the rate  $m_n$ .

Next we turn to proving the upper bound. By assumptions of this theorem, the conditions of Lemma 1 are also satisfied and thus Lemma 1 implies that  $\hat{b}_\xi$  is strongly asymptotically unbiased at  $\beta_0$ . It remains to calculate the variance of  $\hat{b}_\xi$ . Consider the following decomposition

$$\hat{b}_\xi - \xi^T \beta_0 = \xi^T \Theta_0 X^T \epsilon / n + \xi^T (\hat{\Theta} - \Theta_0)^T X^T \epsilon / n + \xi^T (\hat{\Sigma} \hat{\Theta} - I)^T (\hat{\beta} - \beta_0).$$



Then one can show using the Cauchy-Schwarz inequality that

$$\begin{aligned}
 (25) \quad \text{var}_{\beta_0}(\hat{b}_\xi) &= \underbrace{\text{var}_{\beta_0}(\xi^T \Theta_0 X^T \epsilon / n)}_{r_1} + \underbrace{\text{var}_{\beta_0}(\xi^T (\hat{\Theta} - \Theta_0)^T X^T \epsilon / n)}_{r_2} \\
 &\quad + \underbrace{\text{var}_{\beta_0}(\xi^T (\hat{\Sigma} \hat{\Theta} - I)^T (\hat{\beta} - \beta_0))}_{r_3} \\
 &\quad + \mathcal{O}(r_1^{1/2} r_2^{1/2} + r_1^{1/2} r_3^{1/2} + r_2^{1/2} r_3^{1/2}).
 \end{aligned}$$

First note that as in the proof of Lemma 1, we have  $\mathbb{E} \xi^T \Theta_0 X^T \epsilon / n = 0$  and hence

$$\begin{aligned}
 r_1 = \text{var}_{\beta_0}(\xi^T \Theta_0 X^T \epsilon / n) &= \mathbb{E}_{\beta_0}(\mathbb{E}[(\xi^T \Theta_0 X^T \epsilon / n)^2 | X]) \\
 &= \mathbb{E}_{\beta_0}(\Theta_0 \xi)^T X^T X / n \Theta_0 \xi = \xi^T \Theta_0 \xi / n.
 \end{aligned}$$

For a random variable  $U$ , we have  $\text{var}(U) \leq \mathbb{E} U^2$  and hence by Hölder's inequality and the Cauchy-Schwarz inequality

$$\begin{aligned}
 r_2 &= \text{var}_{\beta_0}(\xi^T (\hat{\Theta} - \Theta_0)^T X^T \epsilon / n) \\
 &\leq \mathbb{E}_{\beta_0}(\xi^T (\hat{\Theta} - \Theta_0)^T X^T \epsilon / n)^2 \\
 &\leq \|\xi\|_1^2 \mathbb{E}_{\beta_0} \left\| \hat{\Theta} - \Theta_0 \right\|_1^2 \|X^T \epsilon / n\|_\infty^2 \\
 &\leq \|\xi\|_1^2 \left( \mathbb{E}_{\beta_0} \left\| \hat{\Theta} - \Theta_0 \right\|_1^4 \right)^{1/2} (\mathbb{E}_{\beta_0} \|X^T \epsilon / n\|_\infty^4)^{1/2} \\
 &= \mathcal{O}(\max_{j=1, \dots, p} \lambda_j^2 s_j^2 \log p / n) = o(1/n),
 \end{aligned}$$

where we used the result of Lemma 11 and applied Lemma 4. For the remainder  $r_3$  we have

$$\begin{aligned}
 r_3 &\leq \mathbb{E}_{\beta_0}(\xi^T (\hat{\Sigma} \hat{\Theta} - I)^T (\hat{\beta} - \beta_0))^2 \\
 &\leq \|\xi\|_1^2 \mathbb{E}_{\beta_0} \|\hat{\Sigma} \hat{\Theta} - I\|_\infty^2 \|\hat{\beta} - \beta_0\|_1^2 \\
 &\leq \|\xi\|_1^2 \left( \mathbb{E}_{\beta_0} \|\hat{\Sigma} \hat{\Theta} - I\|_\infty^4 \right)^{1/2} \left( \mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1^4 \right)^{1/2} \\
 &= \mathcal{O}(\max_{j=1, \dots, p} \lambda_j^2 s_j^2 \lambda^2) = o(1/n),
 \end{aligned}$$

where we used Lemma 10 and Theorem 1.

Thus using the above calculations and using (25) we conclude that

$$\text{var}_{\beta_0}(\hat{b}_\xi) = \xi^T \Theta_0 \xi / n + o(1/n).$$

□

15.3. *Proofs for Section 8.4: Main results for fixed design.*

PROOF OF THEOREM 4. The proof follows the same lines as the proof of Theorem 2. The only difference is that we need to check the condition

$$\mathbb{E}_{\beta_0} \left( \frac{p_{\beta_0+h/\sqrt{m_n}}(Z) - p_{\beta_0}(Z)}{p_{\beta_0}(Z)/\sqrt{m_n}} - s_{\beta_0}(Z)^T h \right)^2 = o(n),$$

for fixed design, where  $p_{\beta_0}$ ,  $Z$  and  $s_{\beta_0}$  are defined identically as in the proof of Theorem 2. We denote  $u := h/\sqrt{m_n}$ . Analogously as in the proof of Lemma 14 in Section 15.2, we obtain

$$\begin{aligned} \mathbb{E}_{\beta_0} \left( e^{-\epsilon^T Xu + \frac{1}{2} u^T X^T Xu} - 1 - \epsilon^T Xu \right)^2 &= e^{u^T X^T Xu} - 1 - u^T X^T Xu \\ &= o(u^T X^T Xu) = o(nu^T \hat{\Sigma} u). \end{aligned}$$

Then by the assumption  $h^T \hat{\Sigma} h = \mathcal{O}(1)$  we obtain

$$o(nu^T \hat{\Sigma} u) = o(nh^T \hat{\Sigma} h/m_n) = o(h^T \hat{\Sigma} h n/m_n) = o(n/m_n).$$

Hence plugging in  $u = h/\sqrt{m_n}$  and multiplying by  $m_n$  we obtain

$$\mathbb{E}_{\beta_0} \left( \frac{p_{\beta_0+h/\sqrt{m_n}}(Z) - p_{\beta_0}(Z)}{p_{\beta_0}(Z)/\sqrt{m_n}} - s_{\beta_0}(Z)^T h \right)^2 = o(n).$$

□

PROOF OF THEOREM 5. The lower bound follows by Theorem 4 (note that  $g(\beta) = \beta_j$  and thus the condition on  $g$  is satisfied) applied with  $h := \hat{\Theta}_j/\sqrt{\hat{\Theta}_{jj}}$ . We only need to check that  $h^T \hat{\Sigma} h = \mathcal{O}(1)$ . By the assumption  $\beta_0 + \hat{\Theta}_j/\sqrt{\hat{\Theta}_{jj}n} \in \mathcal{B}(d_n)$  we obtain that  $s_j := \|\hat{\Theta}_j\|_0 \leq d_n$ . Then

$$\begin{aligned} h^T \hat{\Sigma} h &= \hat{\Theta}_j^T \hat{\Sigma} \hat{\Theta}_j / \hat{\Theta}_{jj} \\ &\leq \|\hat{\Theta}_j^T\|_1 \|\hat{\Sigma} \hat{\Theta}_j - e_j\|_\infty / \hat{\Theta}_{jj} + \hat{\Theta}_j^T e_j / \hat{\Theta}_{jj} \\ &\leq \mathcal{O} \left( \sqrt{s_j} \|\hat{\Sigma} \hat{\Theta}_j \hat{\tau}_j^2 - \hat{\tau}_j^2\|_\infty \right) + 1, \\ (26) \quad &\leq \mathcal{O} \left( \sqrt{s_j} \lambda_j / \hat{\tau}_j^2 \right) + 1 = o(1) + 1, \end{aligned}$$

where we used the KKT condition for  $\hat{\gamma}_j$ ,  $1/\hat{\tau}_j^2 = \hat{\Theta}_{jj} \leq \|\hat{\Theta}_j\|_2 = \mathcal{O}(1)$  and  $s_j \leq d_n = o(\sqrt{n}/\log p)$ . This yields the lower bound  $\hat{\Theta}_{jj} + o(1)$ .

Next we turn to proving the upper bound. Strong asymptotic unbiasedness of  $\hat{b}_j$  follows similarly as in Lemma 1 under the assumptions  $\beta_0 \in \mathcal{B}(d_n)$ ,

$d_n = o(\sqrt{n}/\log p)$ , if  $\hat{\Sigma}$  satisfies the compatibility condition with a universal constant and  $\beta_0 + \hat{\Theta}_j/\sqrt{\hat{\Theta}_{jj}n} \in B(\beta_0, c/\sqrt{n})$ .

For the variance of

$$\hat{b}_j - \beta_j^0 = (\hat{\Theta}_j)^T X^T \epsilon/n + (\hat{\Sigma}\hat{\Theta}_j - e_j)^T (\hat{\beta} - \beta_0),$$

we get (using that  $|\hat{\Theta}_j^T \hat{\Sigma} \hat{\Theta}_j / \hat{\Theta}_{jj} - 1| = o(1)$  as derived in (26))

$$\begin{aligned} \text{var}_{\beta_0}(\hat{b}_j) &= \hat{\Theta}_j^T \hat{\Sigma} \hat{\Theta}_j / n + \mathcal{O}(\mathbb{E} \|\hat{\Sigma} \hat{\Theta}_j - e_j\|_\infty^2 \|\hat{\beta} - \beta_0\|_1^2) \\ &= \hat{\Theta}_j^T \hat{\Sigma} \hat{\Theta}_j / n + \mathcal{O}(\|\hat{\Sigma} \hat{\Theta}_j - e_j\|_\infty^2 \mathbb{E}_{\beta_0} \|\hat{\beta} - \beta_0\|_1^2) \\ &= \hat{\Theta}_j^T \hat{\Sigma} \hat{\Theta}_j / n + \mathcal{O}(\lambda_j^2 / (\hat{\tau}_j^2)^2 s^2 \lambda^2) \\ &= \hat{\Theta}_{jj} / n + o(1/n), \end{aligned}$$

where we used  $1/\hat{\tau}_j^2 = \hat{\Theta}_{jj} \leq \|\hat{\Theta}_j\|_2 = \mathcal{O}(1)$ .  $\square$

#### 15.4. Proofs for Section 8.5: Le Cam's bounds.

PROOF OF THEOREM 6. We apply Theorem 9 from Section 10. In this setting, we have asymptotic linearity of the de-sparsified Lasso (see van de Geer et al. (2014)) with the influence function  $l_{\beta_0}(X_i, Y_i) = (\Theta_j^0)^T X_i \epsilon_i$ , where  $\Theta_j^0$  is the  $j$ -th column of the precision matrix. We first show the bias condition (22) is satisfied with the influence function  $l_\beta$ . By direct calculation, for any  $h \in \mathbb{R}^p$  we have

$$\begin{aligned} P_{\beta_0}(l_{\beta_0} h^T s_{\beta_0}) - h^T e_j &= (\Theta_j^0)^T \mathbb{E} X_1 \epsilon_1^2 X_1^T h - h_j \\ &= (\Theta_j^0)^T \mathbb{E} X_1 X_1^T \mathbb{E}(\epsilon_1^2 | X_1) h - h_j = 0. \end{aligned}$$

Therefore in this case the bias condition holds. Hence we can conclude by Theorem 9 that the de-sparsified estimator

$$\hat{b}_j = \hat{\beta}_j + \hat{\Theta}_j^T X^T (Y - X\hat{\beta})/n$$

satisfies for every  $\tilde{\beta}_n \in B(\beta_0, \frac{c}{\sqrt{n}})$

$$\frac{\sqrt{n}(\hat{b}_j - \tilde{\beta}_n)}{(\Theta_{jj}^0)^{1/2}} \overset{\tilde{\beta}_n}{\rightsquigarrow} \mathcal{N}(0, 1).$$

In addition, no asymptotically linear estimator satisfying the condition (22) (here condition (13)) can have smaller asymptotic variance than  $\Theta_{jj}^0$  as follows by the lower bound on the asymptotic variance in Lemma 3 in Section 10, i.e.

$$V_{\beta_0} \geq \dot{g}(\beta_0)^T I_{\beta_0}^{-1} \dot{g}(\beta_0) + o(1) = \Theta_{jj}^0 + o(1).$$

$\square$

## 16. Proofs for Section 9: Gaussian graphical models.

16.1. *Proofs for Section 9.2: Strong asymptotic unbiasedness of the de-sparsified nodewise Lasso.*

PROOF OF LEMMA 2. By the Karush-Kuhn-Tucker conditions corresponding to the nodewise Lasso estimator, we have  $\|\hat{\Sigma}\hat{\Theta}_i - e_i\|_\infty = \mathcal{O}(\lambda_i/\hat{\tau}_i^2)$ . Hence, and applying a version of Lemma 11 without the maximum over  $j = 1, \dots, p$ , we obtain

$$\begin{aligned}
\mathbb{E}_{\Theta_0}(\hat{T}_{ij} - \Theta_{ij}^0) &= \underbrace{\mathbb{E}_{\Theta_0}(\Theta_i^0)^T(\hat{\Sigma} - \Sigma_0)\Theta_j^0}_{=0} + \mathbb{E}_{\Theta_0}(\hat{\Theta}_i - \Theta_i^0)^T(\hat{\Sigma}\Theta_j^0 - e_j) \\
&\quad + \mathbb{E}_{\Theta_0}(\hat{\Sigma}\hat{\Theta}_i - e_i)^T(\hat{\Theta}_j - \Theta_j^0) \\
&\leq (\mathbb{E}_{\Theta_0}\|\hat{\Theta}_i - \Theta_i^0\|_1^2)^{1/2}(\mathbb{E}_{\Theta_0}\|\hat{\Sigma}\Theta_j^0 - e_j\|_\infty^2)^{1/2} \\
&\quad + (\mathbb{E}_{\Theta_0}\|\hat{\Sigma}\hat{\Theta}_i - e_i\|_\infty^2)^{1/2}(\mathbb{E}_{\Theta_0}\|\hat{\Theta}_j - \Theta_j^0\|_1^2)^{1/2} \\
&\leq \mathcal{O}(s_i\lambda_i\lambda_j(\mathbb{E}_{\Theta_0}1/(\hat{\tau}_j^2)^2)^{1/2}) + \mathcal{O}(s_j\lambda_i\lambda_j(\mathbb{E}_{\Theta_0}1/(\hat{\tau}_i^2)^2)^{1/2}) \\
&= o(1/\sqrt{n}).
\end{aligned}$$

□

16.2. *Proofs for Section 9.3: Main results.* In this section we will give the proof of Theorems 7 and 8. In the proof of Theorem 7, we use Lemma 20 which is stated and proved in Appendix B.

PROOF OF THEOREM 7. The proof is similar to the proof of Theorem 2. By strong asymptotic unbiasedness of  $T_n$  at  $\Theta_0$  in the direction  $H$ , it follows

$$\begin{aligned}
\xi_1^T H \xi_2 &= \sqrt{m_n} (\xi_1^T (\Theta_0 + H/\sqrt{m_n}) \xi_2 - \xi_1^T \Theta_0 \xi_2) \\
&= \sqrt{m_n} (\mathbb{E}_{\Theta_0 + H/\sqrt{m_n}} T_n(X) - \mathbb{E}_{\Theta_0} T_n(X)) + o(1)
\end{aligned}$$

Let  $s_{\Theta_0}(X) := -n(\hat{\Sigma} - \Sigma_0)/2$ . Denoting the probability density correspond-

ing to  $X$  by  $p_\Theta$ , we may further rewrite the expressions to obtain

$$\begin{aligned}
& \sqrt{m_n} \left( \mathbb{E}_{\Theta_0+H/\sqrt{m_n}} T_n(X) - \mathbb{E}_{\Theta_0} T_n(X) \right) \\
&= \sqrt{m_n} \int T_n(x) (p_{\Theta_0+H/\sqrt{m_n}}(x) - p_{\Theta_0}(x)) dz \\
&= \sqrt{m_n} \mathbb{E}_{\Theta_0} T_n(X) \frac{p_{\Theta_0+H/\sqrt{m_n}}(X) - p_{\Theta_0}(X)}{p_{\Theta_0}(X)} \\
&= \mathbb{E}_{\Theta_0} T_n(X) \left( \frac{p_{\Theta_0+H/\sqrt{m_n}}(X) - p_{\Theta_0}(X)}{p_{\Theta_0}(X)/\sqrt{m_n}} - \text{vec}(H)^T \text{vec}(s_{\Theta_0}(X)) \right) \\
&\quad + \mathbb{E}_{\Theta_0} T_n(X) \text{vec}(H)^T \text{vec}(s_{\Theta_0}(X)) \\
&= \mathbb{E}_{\Theta_0} (T_n(X) - \xi_1^T \Theta_0 \xi_2) \times \\
&\quad \times \left( \frac{p_{\Theta_0+H/\sqrt{m_n}}(X) - p_{\Theta_0}(X)}{p_{\Theta_0}(X)/\sqrt{m_n}} - \text{vec}(H)^T \text{vec}(s_{\Theta_0}(X)) \right) \\
&\quad + \mathbb{E}_{\Theta_0} T_n(X) \text{vec}(H)^T \text{vec}(s_{\Theta_0}(X)),
\end{aligned}$$

where in the last equality we used that  $\mathbb{E}_{\Theta_0} s_{\Theta_0}(X) = 0$ ,  $\int p_{\Theta_0}(x) dx = 1$  and  $\int p_{\Theta_0+H/\sqrt{m_n}}(x) dx = 1$ . Since the variance of  $T_n$  is  $\mathcal{O}(1/n)$  by the definition of strong asymptotic unbiasedness, then

$$\begin{aligned}
\mathbb{E}_{\Theta_0} (T_n(X) - \xi_1^T \Theta_0 \xi_2)^2 &= \text{var}_{\Theta_0}(T_n(X)) + [\mathbb{E}_{\Theta_0} (T_n(X) - \xi_1^T \Theta_0 \xi_2)]^2 \\
&= \mathcal{O}(1/n) + o(1/n) = \mathcal{O}(1/n).
\end{aligned}$$

We need to use Lemma 20 in Appendix B to conclude that the remainder is small. Lemma 20 implies

$$\mathbb{E} \left( \frac{p_{\Theta_0+H/\sqrt{m_n}}(X)}{p_{\Theta_0}(X)} - 1 - \text{vec}(H)^T \text{vec}(s_{\Theta_0}(X))/\sqrt{m_n} \right)^2 = o(\delta_n).$$

Consequently, and by the Cauchy-Schwarz inequality, we have the upper bound

$$\begin{aligned}
& \left| \mathbb{E}_{\Theta_0} (T_n(X) - \xi_1^T \Theta_0 \xi_2) \left( \frac{p_{\Theta_0+H/\sqrt{m_n}}(X) - p_{\Theta_0}(X)}{p_{\Theta_0}(X)/\sqrt{m_n}} - \text{vec}(H)^T \text{vec}(s_{\Theta_0}(X)) \right) \right| \\
&\leq \sqrt{\mathbb{E}_{\Theta_0} (T_n(X) - \xi_1^T \Theta_0 \xi_2)^2} \times \\
&\quad \times \sqrt{\mathbb{E}_{\Theta_0} \left( \frac{p_{\Theta_0+H/\sqrt{m_n}}(X) - p_{\Theta_0}(X)}{p_{\Theta_0}(X)/\sqrt{m_n}} - \text{vec}(H)^T \text{vec}(s_{\Theta_0}(X)) \right)^2} \\
&= \mathcal{O}(1/\sqrt{n}) o(\sqrt{n}) = o(1).
\end{aligned}$$

Thus we have

$$\xi_1^T H \xi_2 = \mathbb{E}_{\Theta_0} T_n(X) \text{vec}(H)^T \text{vec}(s_{\Theta_0}(X)) + o(1).$$

Hence, because

$$\begin{aligned} & |\xi_1^T H \xi_2| - |\text{cov}(T_n, \text{vec}(H)^T \text{vec}(s_{\Theta_0}(X)))| \\ & \leq |\xi_1^T H \xi_2 - \text{cov}(T_n, \text{vec}(H)^T \text{vec}(s_{\Theta_0}(X)))| \\ & = o(1) \end{aligned}$$

it follows using the Cauchy-Schwarz inequality that

$$\begin{aligned} |\xi_1^T H \xi_2| &= |\text{cov}_{\Theta_0}(T_n, \text{tr}(-n(\hat{\Sigma} - \Sigma_0)H/2))| + o(1) \\ &\leq \sqrt{\text{var}_{\Theta_0}(T_n)} \sqrt{\text{var}_{\Theta_0}(-n \text{tr}((\hat{\Sigma} - \Sigma_0)H/2))} + o(1). \end{aligned}$$

Now we have

$$\begin{aligned} \text{var}_{\Theta_0}(\text{tr}(n(\hat{\Sigma} - \Sigma_0)H/2)) &= \text{var}_{\Theta_0}(n \text{tr}(\hat{\Sigma}H/2)) \\ &= \text{var}_{\Theta_0}(\xi_1^T \Theta_0 X^T X \Theta_0 \xi_2)/\sigma^2 \\ &= n \text{var}_{\Theta_0}(\xi_1^T \Theta_0 X^{(1)} (X^{(1)})^T \Theta_0 \xi_2)/\sigma^2 \\ &= n(\xi_1^T \Theta_0 \xi_1 \xi_2^T \Theta_0 \xi_2 + (\xi_1^T \Theta_0 \xi_2)^2)/\sigma^2 \\ &= n. \end{aligned}$$

Hence we conclude

$$|\xi_1^T H \xi_2| \leq \sqrt{n} \sqrt{\text{var}_{\Theta_0}(T_n)} + o(1).$$

Plugging in  $H = \Theta_0(\xi_1 \xi_2^T + \xi_2 \xi_1^T) \Theta_0 / \sigma$  we obtain

$$|\xi_1^T H \xi_2|^2 = (\xi_1^T \Theta_0 \xi_1 \xi_2^T \Theta_0 \xi_2 + (\xi_1^T \Theta_0 \xi_2)^2)/\sigma^2 = \sigma^2.$$

By the strong asymptotic unbiasedness assumption on  $T_n$ , we have  $\text{var}_{\Theta_0}(T_n) = \mathcal{O}(1/n)$  and thus taking squares of both sides of the last inequality we obtain

$$\begin{aligned} & \xi_1^T \Theta_0 \xi_1 \xi_2^T \Theta_0 \xi_2 + (\xi_1^T \Theta_0 \xi_2)^2 \\ & \leq n \text{var}_{\Theta_0}(T_n) + 2\sqrt{n} \sqrt{\text{var}_{\Theta_0}(T_n)} o(1) + o(1) \\ & \leq n \text{var}_{\Theta_0}(T_n) + o(1). \end{aligned}$$

□

PROOF OF THEOREM 8. To obtain the lower bound on the variance, we apply Theorem 7 with  $H := (\Theta_i^0(\Theta_j^0)^T + \Theta_j^0(\Theta_i^0)^T)/\sigma$ . Then by direct calculation, see that the condition  $\text{tr}(H^T \Sigma_0 H) = \mathcal{O}(1)$  is satisfied for this choice of  $H$ . Moreover,  $\Theta_0 + H/\sqrt{n} \in B(\Theta_0, c/\sqrt{n})$  by assumption, but then also  $\Theta_0 + H/\sqrt{m_n} \in B(\Theta_0, c/\sqrt{m_n})$ . Hence by Theorem 7 follows the lower bound  $\text{var}_{\Theta_0}(T) \geq \frac{\xi^T \Theta_0 \xi + o(1)}{n}$ .

Next  $\hat{T}_{ij}$  is strongly asymptotically unbiased at  $\Theta_0$  in every direction  $H$  such that  $\Theta_0 + H/\sqrt{n} \in \mathcal{G}(d_1, \dots, d_p)$ , which follows by Lemma 2.

It remains to calculate the variance of  $\hat{T}_{ij}$ . First we have that

$$\text{var}_{\Theta_0}((\Theta_i^0)^T(\hat{\Sigma} - \Sigma_0)\Theta_j^0) = \frac{1}{n} \text{var}_{\Theta_0}((\Theta_i^0)^T X_1 X_1^T \Theta_j^0) = (\Theta_{ii}^0 \Theta_{jj}^0 + (\Theta_{ij}^0)^2)/n.$$

By basic calculations, it follows that

$$\begin{aligned} \text{var}_{\Theta_0}(\hat{T}_{ij}) &= (\Theta_{ii}^0 \Theta_{jj}^0 + (\Theta_{ij}^0)^2)/n + \mathcal{O}(\mathbb{E}_{\Theta_0}((\hat{\Sigma} \Theta_i^0 - e_i)^T(\hat{\Theta}_j - \Theta_j^0))^2) \\ &\quad + \mathcal{O}(\mathbb{E}_{\Theta_0}((\hat{\Theta}_i - \Theta_i^0)^T(\hat{\Sigma} \hat{\Theta}_j - e_j))^2) \\ &= (\Theta_{ii}^0 \Theta_{jj}^0 + (\Theta_{ij}^0)^2)/n + \mathcal{O}(\mathbb{E}_{\Theta_0} \|\hat{\Sigma} \Theta_i^0 - e_i\|_\infty^2 \|\hat{\Theta}_j - \Theta_j^0\|_1^2) \\ &\quad + \mathcal{O}(\mathbb{E}_{\Theta_0} \|\hat{\Theta}_i - \Theta_i^0\|_1^2 \|\hat{\Sigma} \hat{\Theta}_j - e_j\|_1^2). \\ &= (\Theta_{ii}^0 \Theta_{jj}^0 + (\Theta_{ij}^0)^2)/n \\ &\quad + \mathcal{O}((\mathbb{E}_{\Theta_0} \|\hat{\Sigma} \Theta_i^0 - e_i\|_\infty^4)^{1/2} (\mathbb{E}_{\Theta_0} \|\hat{\Theta}_j - \Theta_j^0\|_1^4)^{1/2}) \\ &\quad + \mathcal{O}((\mathbb{E}_{\Theta_0} \|\hat{\Theta}_i - \Theta_i^0\|_1^4)^{1/2} (\mathbb{E}_{\Theta_0} \|\hat{\Sigma} \hat{\Theta}_j - e_j\|_1^4)^{1/2}) \\ &= (\Theta_{ii}^0 \Theta_{jj}^0 + (\Theta_{ij}^0)^2)/n + \mathcal{O}(\lambda_i^2 s_j^2 \lambda_j^2 (\mathbb{E}_{\Theta_0} 1/(\hat{\tau}_i^2)^2)^{1/2}) \\ &\quad + \mathcal{O}(s_i^2 \lambda_i^2 \lambda_j^2 (\mathbb{E}_{\Theta_0} 1/(\hat{\tau}_j^2)^2)^{1/2}) \\ &= (\Theta_{ii}^0 \Theta_{jj}^0 + (\Theta_{ij}^0)^2)/n + o(1/n). \end{aligned}$$

□

### 17. Proofs for Section 10: Le Cam's bounds for general models.

In this section we give the proof of Theorem 9, for which we need Lemma 16 below. Some technical results (contained in Lemmas 21 and 23) are stated and proved in Appendix C.

LEMMA 16. *Assume the conditions of Theorem 9. Suppose that  $Z_n \rightsquigarrow Z$ , where  $Z$  is a random vector with values in  $\mathbb{R}^2$ . Let  $X_n = \psi(Z_n)$  and  $U_n = \psi(Z)$ , where*

$$x \mapsto \psi(x_1, x_2) = \begin{pmatrix} 1/\sqrt{v_{11}} & 0 \\ 0 & 1 \end{pmatrix} \left[ V^{1/2} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} + \begin{pmatrix} -v_{12} \\ -v_{22}/2 \end{pmatrix} \right].$$

*Then the following statements hold.*

1. For any function  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  which is bounded and continuous it holds that

$$\lim_{n \rightarrow \infty} \mathbb{E}f(X_n) - \mathbb{E}f(U_n) = 0.$$

2. Let  $f$  be any bounded and continuous function  $f : \mathbb{R} \rightarrow \mathbb{R}$ . Suppose that

$$\lim_{M \rightarrow \infty} \lim_{n \rightarrow \infty} \mathbb{E} \min(0, M - e^{U_{n,2}}) = 0.$$

Then it holds that

$$\lim_{n \rightarrow \infty} \mathbb{E}f(X_{n,1})e^{X_{n,2}} - \mathbb{E}f(U_{n,1})e^{U_{n,2}} = 0.$$

PROOF OF LEMMA 16. We first prove the first statement. Let  $\epsilon > 0$  and let  $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  be continuous and bounded.

The map  $\psi$  is linear, i.e.  $\psi(x) = Ax + b$  for some  $A \in \mathbb{R}^{2 \times 2}$  and  $b \in \mathbb{R}^2$  ( $A, b$  depending on  $n$ ). Denote

$$D := \begin{pmatrix} 1/\sqrt{v_{11}} & 0 \\ 0 & 1 \end{pmatrix}$$

Observe that for any  $x \in \mathbb{R}^2$

$$\|Ax\|_2^2 = x^T A^T A x = x^T D V D x \leq \Lambda_{\max}(D V D) x^T x.$$

By Lemma 22 we have that  $\Lambda_{\max}(D V D) = \mathcal{O}(1)$  and  $\|b\|_2 = \mathcal{O}(1)$ . Therefore, when  $\|x\|_2 = \mathcal{O}(1)$ , then

$$(27) \quad \|Ax + b\|_2 = \mathcal{O}(1).$$

Take a compact rectangle  $R \subset \mathbb{R}^2$  not depending on  $n$  and such that  $P(Z \notin R) < \epsilon$ .

Divide the rectangle  $R$  into a finite number of non-overlapping rectangles of diameter at most  $\delta/L^{1/2}$ , where  $L$  is a universal constant such that  $L \geq \Lambda_{\max}(D V D)$ . By construction, the number of these rectangles, denote it  $N$ , does not depend on  $n$ . So we have  $R = \cup_{j=1}^N R_j$ , where each  $R_j$  is a rectangle of diameter at most  $\delta/L^{1/2}$ .

For all  $x, y \in R_j$  it holds that  $\|x - y\|_2 \leq \delta/L^{1/2}$  and thus

$$(28) \quad \|\psi(x) - \psi(y)\|_2 = \|A(x - y)\|_2 \leq L^{1/2} \|x - y\|_2 \leq \delta.$$

Note that by (27), there exists a compact set  $S$  not depending on  $n$  such that  $\psi(R) \subset S$  for all  $n$ . The continuous function  $f$  is uniformly continuous on the compact set  $S$ . Hence for the  $\epsilon$  there exists a  $\delta > 0$  such that for all



$z, v \in S$  it holds that if  $\|z - v\|_2 < \delta$  then  $|f(z) - f(v)| < \epsilon$ . But then since for all  $x, y \in R_j$  we have that  $\psi(x), \psi(y) \in S$ , we obtain by (28) and the absolute continuity of  $f$  that

$$|f(\psi(x)) - f(\psi(y))| < \epsilon$$

for all  $n$ . Take a point  $x_j$  from each set  $R_j$  and define  $f_\epsilon = \sum_{j=1}^N f(\psi(x_j)) \mathbf{1}_{R_j}$ . Then  $|f(\psi(x)) - f_\epsilon(x)| < \epsilon$  for all  $x \in R$  (and all  $n$ ) and hence if  $f$  takes values in  $[-K, K]$ , we have the following upper bounds

$$(29) \quad |\mathbb{E}f(\psi(Z)) - \mathbb{E}f_\epsilon(Z)| \leq \epsilon + 2KP(Z \notin R),$$

$$(30) \quad |\mathbb{E}f(\psi(Z_n)) - \mathbb{E}f_\epsilon(Z_n)| \leq \epsilon + 2KP(Z_n \notin R),$$

$$(31) \quad |\mathbb{E}f_\epsilon(Z_n) - \mathbb{E}f_\epsilon(Z)| \leq \sum_{j=1}^N |P(Z_n \in R_j) - P(Z \in R_j)| |f(\psi(x_j))|.$$

Since  $Z_n \rightsquigarrow Z$ , for all  $j = 1, \dots, N$  it holds

$$|P(Z_n \in R_j) - P(Z \in R_j)| \rightarrow 0.$$

Similarly,

$$|P(Z \notin R) - P(Z_n \notin R)| = |P(Z \in R) - P(Z_n \in R)| \rightarrow 0.$$

Finally, by construction we have  $P(Z \notin R) < \epsilon$ . We thus conclude that the upper bounds (29), (30) and (31) can be made smaller than  $C\epsilon$  for  $n$  sufficiently large. The claim follows by combining the three upper bounds.

Next we prove the second statement. Denote  $g(x_1, x_2) = f(x_1)e^{x_2}$ . We write  $g = g^+ - g^-$ , where  $g^+ = \max\{g, 0\}$  is the positive part and  $g^- := \max\{-g, 0\}$  is the negative part. We first prove for the positive part  $g^+$  that

$$(32) \quad \lim_{n \rightarrow \infty} \mathbb{E}g^+(X_n) - \mathbb{E}g^+(U_n) = 0.$$

This will be achieved by first showing that

$$\liminf_{n \rightarrow \infty} \mathbb{E}f^+(X_n)e^{X_{n,2}} - \mathbb{E}f^+(U_n)e^{U_{n,2}} \geq 0$$

and secondly showing that

$$\limsup_{n \rightarrow \infty} \mathbb{E} f^+(X_n) e^{X_{n,2}} - \mathbb{E} f^+(U_n) e^{U_{n,2}} \leq 0.$$

Then combining the two gives (32).

First we prove that  $\liminf_{n \rightarrow \infty} \mathbb{E} f^+(X_n) e^{X_{n,2}} - \mathbb{E} f^+(U_n) e^{U_{n,2}} \geq 0$ . For every  $M$ , since  $g^+$  is non-negative, it holds that  $f^+(x) e^{x^2} \geq f^+(x)(e^{x^2} \wedge M)$ , where the symbol  $\wedge$  denotes the minimum. Hence

$$\begin{aligned} & \mathbb{E} f^+(X_n) e^{X_{n,2}} - \mathbb{E} f^+(U_n) e^{U_{n,2}} \\ & \geq \mathbb{E} f^+(X_n) (e^{X_{n,2}} \wedge M) - \mathbb{E} f^+(U_n) e^{U_{n,2}} \\ & = [\mathbb{E} f^+(X_n) (e^{X_{n,2}} \wedge M) - \mathbb{E} f^+(U_n) (e^{U_{n,2}} \wedge M)] \\ & \quad + [\mathbb{E} f^+(U_n) (e^{U_{n,2}} \wedge M) - \mathbb{E} f^+(U_n) e^{U_{n,2}}] \end{aligned}$$

We have  $f^+(x)(e^{x^2} \wedge M) - f^+(x) e^{x^2} = f^+(x) \min(0, M - e^{x^2})$ . Taking limes inferior of both sides, it follows that

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \mathbb{E} f^+(X_n) e^{X_{n,2}} - \mathbb{E} f^+(U_n) e^{U_{n,2}} \\ & \geq \liminf_{n \rightarrow \infty} [\mathbb{E} f^+(X_n) (e^{X_{n,2}} \wedge M) - \mathbb{E} f^+(U_n) (e^{U_{n,2}} \wedge M)] \\ & \quad + \liminf_{n \rightarrow \infty} -\mathbb{E} f^+(U_n) \min(0, M - e^{U_{n,2}}). \end{aligned}$$

For every fixed  $M$ , the function  $x \mapsto f^+(x)(e^{x^2} \wedge M)$  is bounded and continuous. We may thus apply the first result of the lemma to conclude

$$\liminf_{n \rightarrow \infty} \mathbb{E} f^+(X_n) (e^{X_{n,2}} \wedge M) - \mathbb{E} f^+(U_n) (e^{U_{n,2}} \wedge M) = 0.$$

Therefore, we have

$$\begin{aligned} (33) \quad & \liminf_{n \rightarrow \infty} \mathbb{E} f^+(X_n) e^{X_{n,2}} - \mathbb{E} f^+(U_n) e^{U_{n,2}} \\ & \geq \liminf_{n \rightarrow \infty} -\mathbb{E} f^+(U_n) \min(0, M - e^{U_{n,2}}). \end{aligned}$$

Next since  $|f^+| \leq K$  we have

$$|-\mathbb{E} f^+(U_n) \min(0, M - e^{U_{n,2}})| \leq K \mathbb{E} \min(0, M - e^{U_{n,2}}).$$

Then the assumption

$$\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \mathbb{E} \min(0, M - e^{U_{n,2}}) = 0$$

implies that also

$$\begin{aligned} & \lim_{m \rightarrow \infty} \liminf_n -\mathbb{E}f^+(U_n) \min(0, M - e^{U_{n,2}}) \\ &= - \lim_{m \rightarrow \infty} \limsup_n \mathbb{E}f^+(U_n) \min(0, M - e^{U_{n,2}}) = 0, \end{aligned}$$

so we conclude that

$$(34) \quad \liminf_{n \rightarrow \infty} \mathbb{E}f^+(X_n)e^{X_{n,2}} - \mathbb{E}f^+(U_n)e^{U_{n,2}} \geq 0.$$

Now we prove

$$\limsup_{n \rightarrow \infty} \mathbb{E}f^+(X_n)e^{X_{n,2}} - \mathbb{E}f^+(U_n)e^{U_{n,2}} \leq 0.$$

Similarly as before, since  $K - f^+ \geq 0$  ( $K$  is an upper bound on  $f$ ), we have that

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \mathbb{E}(K - f^+(X_{n,1}))e^{X_{n,2}} - \mathbb{E}(K - f^+(U_{n,1}))e^{U_{n,2}} \\ & \geq \liminf_{n \rightarrow \infty} \mathbb{E}(K - f^+(X_{n,1}))(e^{X_{n,2}} \wedge M) - \mathbb{E}(K - f^+(U_{n,1}))(e^{U_{n,2}} \wedge M) \\ (35) \quad & + \liminf_{n \rightarrow \infty} \mathbb{E}(K - f^+(U_{n,1}))(e^{U_{n,2}} \wedge M) - \mathbb{E}(K - f^+(U_{n,1}))e^{U_{n,2}}. \end{aligned}$$

By the first part of the lemma, we have that for every fixed  $M$  it holds

$$\liminf_n \mathbb{E}(K - f^+(X_{n,1}))(e^{X_{n,2}} \wedge M) - \mathbb{E}(K - f^+(U_{n,1}))(e^{U_{n,2}} \wedge M) = 0,$$

since the function  $(x_1, x_2) \mapsto (K - f^+(x_1))(e^{x_2} \wedge M)$  is bounded and continuous.

For the term in (35), we have since  $|K - f^+| \leq 2K$

$$| -\mathbb{E}(K - f^+(U_{n,1})) \min(0, M - e^{U_{n,2}}) | \leq 2K \mathbb{E}e^{U_{n,2}} \min(0, M - e^{U_{n,2}}).$$

Hence by the assumption  $\lim_{m \rightarrow \infty} \lim_{n \rightarrow \infty} \mathbb{E} \min(0, M - e^{U_{n,2}}) = 0$ , we have that

$$\liminf_{n \rightarrow \infty} -\mathbb{E}(K - f^+(U_{n,1})) \min(0, M - e^{U_{n,2}}) = 0.$$

Thus we conclude that

$$\liminf_{n \rightarrow \infty} \mathbb{E}(K - f^+(X_{n,1}))e^{X_{n,2}} - \mathbb{E}(K - f^+(U_{n,1}))e^{U_{n,2}} \geq 0.$$

Now note that

$$\begin{aligned}
& \liminf_{n \rightarrow \infty} \mathbb{E}(K - f^+(X_{n,1}))e^{X_{n,2}} - \mathbb{E}(K - f^+(U_{n,1}))e^{U_{n,2}} \\
&= \liminf_n -\mathbb{E}f^+(X_{n,1})e^{X_{n,2}} + \mathbb{E}f^+(U_{n,1})e^{U_{n,2}} \\
&= -\limsup_n \mathbb{E}f^+(X_{n,1})e^{X_{n,2}} - \mathbb{E}f^+(U_{n,1})e^{U_{n,2}}.
\end{aligned}$$

So in conclusion we have shown that

$$\begin{aligned}
& \limsup_n \mathbb{E}f^+(X_{n,1})e^{X_{n,2}} - \mathbb{E}f^+(U_{n,1})e^{U_{n,2}} \\
& \leq 0 \leq \liminf_n \mathbb{E}f^+(X_{n,1})e^{X_{n,2}} - \mathbb{E}f^+(U_{n,1})e^{U_{n,2}}.
\end{aligned}$$

This proves (32).

The same procedure can be used for the negative part  $f^-$  (since  $f^-$  is also bounded and positive) to show that

$$\lim_{n \rightarrow \infty} \mathbb{E}f^-(X_{n,1})e^{X_{n,2}} - \mathbb{E}f^-(U_{n,1})e^{U_{n,2}} = 0.$$

We then conclude that

$$\begin{aligned}
& \lim_{n \rightarrow \infty} \mathbb{E}f(X_{n,1})e^{X_{n,2}} - \mathbb{E}f(U_{n,1})e^{U_{n,2}} \\
& \leq \lim_{n \rightarrow \infty} |\mathbb{E}f^+(X_{n,1})e^{X_{n,2}} - \mathbb{E}f^+(U_{n,1})e^{U_{n,2}}| \\
& \quad + \lim_{n \rightarrow \infty} |\mathbb{E}f^-(X_{n,1})e^{X_{n,2}} - \mathbb{E}f^-(U_{n,1})e^{U_{n,2}}| \\
& = 0.
\end{aligned}$$

□

PROOF OF THEOREM 9. We denote  $\ell_\beta(x) := \log p_\beta(x)$  and

$$\Lambda_n := \sum_{i=1}^n \ell_{\beta_{n,0} + h/\sqrt{n}}(X^{(i)}) - \ell_{\beta_{n,0}}(X^{(i)}).$$

Further for  $0 \leq t \leq 1$  denote  $g(t) := \ell((1-t)\beta + t\beta_0)$ . Then by a two-term Taylor expansion of  $g$  we have

$$g(1) - g(0) = \dot{g}(0) + \frac{1}{2}\ddot{g}(\tilde{t}),$$

where  $0 \leq \tilde{t} \leq 1$ . Rewriting this from the definition of  $g$  gives

$$\ell_\beta - \ell_{\beta_0} = \ell_{\beta_0}(\beta - \beta_0) + \frac{1}{2}(\beta - \beta_0)^T \ddot{\ell}_{\tilde{\beta}}(\beta - \beta_0),$$

where  $\tilde{\beta} := (1 - \tilde{t})\beta + \tilde{t}\beta_0$ . Applying the above with  $\tilde{\beta} := \beta_{n,0} + h/\sqrt{n}$  and  $\beta_0 := \beta_{n,0}$  we obtain

$$\Lambda_n := \sum_{i=1}^n \frac{1}{\sqrt{n}} \dot{\ell}_{\beta_{n,0}}(X^{(i)})^T h + \frac{1}{2} h^T \frac{1}{n} \sum_{i=1}^n \ddot{\ell}_{\tilde{\beta}}(X^{(i)}) h,$$

where  $\tilde{\beta} := (1 - \tilde{t})(\beta_{n,0} + h/\sqrt{n}) + \tilde{t}\beta_{n,0} = \beta_{n,0} + (1 - \tilde{t})h/\sqrt{n}$ . We can then write the decomposition of  $\Lambda_n$  as follows

$$\begin{aligned} \Lambda_n &= \frac{1}{\sqrt{n}} \sum_{i=1}^n h^T \dot{\ell}_{\beta_{n,0}}(X^{(i)}) + \frac{1}{2} h^T \frac{1}{n} \sum_{i=1}^n \ddot{\ell}_{\tilde{\beta}}(X^{(i)}) h \\ &= \frac{1}{\sqrt{n}} \sum_{i=1}^n h^T \dot{\ell}_{\beta_{n,0}}(X^{(i)}) + \underbrace{\frac{1}{2} h^T \frac{1}{n} \sum_{i=1}^n (\ddot{\ell}_{\tilde{\beta}}(X^{(i)}) - \ddot{\ell}_{\beta_{n,0}}(X^{(i)})) h}_{rem_1} \\ (36) \quad &+ \underbrace{\frac{1}{2} h^T \left( \frac{1}{n} \sum_{i=1}^n \ddot{\ell}_{\beta_{n,0}}(X^{(i)}) + I(\beta_{n,0}) \right) h}_{rem_2} \\ &\quad - \frac{1}{2} h^T I(\beta_{n,0}) h. \end{aligned}$$

We will now show that the remainders  $rem_1, rem_2$  converge in probability to zero. First observe that since  $\beta_{n,0} + h/\sqrt{n} \in B(\beta_{n,0}, \frac{c}{\sqrt{n}})$ , this implies that  $\|h\|_2 \leq c$  and  $\|h\|_0 \leq 2s$ . Combining these two properties yields  $\|h\|_1 \leq \sqrt{2sc} = \mathcal{O}(\sqrt{s})$ .

Regarding the first remainder we have for each  $j, k = 1, \dots, p$

$$\frac{1}{n} \sum_{i=1}^n (\ddot{\ell}_{\tilde{\beta}}(X^{(i)}) - \ddot{\ell}_{\beta_{n,0}}(X^{(i)}))_{j,k} = \frac{1}{n} \sum_{i=1}^n (\ddot{\ell}_{\tilde{\beta}}(X^{(i)})^T (\tilde{\beta} - \beta_{n,0}))_{j,k},$$

where  $\tilde{\beta} = (1 - \tilde{t})\tilde{\beta} + \tilde{t}\beta_{n,0}$ . By assumption  $\|\ddot{\ell}_{\tilde{\beta}}\|_{\infty} \leq L$  and hence using Hölder's inequality

$$\begin{aligned} \left\| \frac{1}{n} \sum_{i=1}^n \ddot{\ell}_{\tilde{\beta}}(X^{(i)}) - \ddot{\ell}_{\beta_{n,0}}(X^{(i)}) \right\|_{\infty} &\leq \left\| \frac{1}{n} \sum_{i=1}^n \ddot{\ell}_{\tilde{\beta}}(X^{(i)}) \right\|_{\infty} \|\tilde{\beta} - \beta_{n,0}\|_1 \\ &\leq L \|(1 - \tilde{t})h/\sqrt{n}\|_1 = \mathcal{O}(\sqrt{s/n}). \end{aligned}$$

Thus using Hölder's inequality, for the first remainder we get

$$|h^T \frac{1}{n} \sum_{i=1}^n (\ddot{\ell}_{\tilde{\beta}}(X^{(i)}) - \ddot{\ell}_{\beta_{n,0}}(X^{(i)})) h| \leq \mathcal{O}(s^{3/2}/\sqrt{n}) = o(1).$$

For the second remainder we have using assumption  $\|\frac{1}{n} \sum_{i=1}^n \ddot{\ell}_{\beta_{n,0}}(X^{(i)}) + I(\beta_{n,0})\|_\infty = \mathcal{O}_P(\lambda)$  that

$$\begin{aligned} \|h^T (\frac{1}{n} \sum_{i=1}^n \dot{s}_{\beta_{n,0}}(X^{(i)}) + I(\beta_{n,0}))h\|_\infty &\leq \|h\|_1^2 \|\frac{1}{n} \sum_{i=1}^n \dot{s}_{\beta_{n,0}}(X^{(i)}) + I_{\beta_{n,0}}\|_\infty \\ &= \mathcal{O}_P(s\lambda) = o_P(1). \end{aligned}$$

Hence collecting the results, from the decomposition of  $\Lambda_n$  we have

$$(37) \quad \Lambda_n = \frac{1}{\sqrt{n}} \sum_{i=1}^n h^T \dot{\ell}_{\beta_{n,0}}(X^{(i)}) - \frac{1}{2} h^T I(\beta_{n,0}) h + o_P(1).$$

We introduce the following notation. Let

$$V := \begin{pmatrix} V_{\beta_{n,0}} & P_{\beta_{n,0}}(l_{\beta_{n,0}} h^T s_{\beta_{n,0}}) \\ P_{\beta_{n,0}}(l_{\beta_{n,0}} h^T s_{\beta_{n,0}}) & h^T I(\beta_{n,0}) h \end{pmatrix}$$

Furthermore, we denote the entries of the matrix  $V$  by  $v_{ij}, i, j = 1, 2$ .

The condition (D3) implies that for any fixed  $a \in \mathbb{R}^2$  it holds for all  $\epsilon > 0$

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbb{E}(a^T V^{-1/2} (l_{\beta_{n,0}}(X^{(i)}), h^T s_{\beta_{n,0}}(X^{(i)}))^T)^2 \times \\ \times \mathbf{1}_{|a^T V^{-1/2} (l_{\beta_{n,0}}(X^{(i)}), h^T s_{\beta_{n,0}}(X^{(i)}))^T| > \epsilon \sqrt{na^T a}} = 0, \end{aligned}$$

where we used that  $\|V\|_\infty = \mathcal{O}(1)$ , which follows by Lemma 22. By the Lindeberg's central limit theorem we thus have by condition (38) for any  $a \in \mathbb{R}^2$  that

$$\frac{1}{\sqrt{na^T a}} \sum_{i=1}^n a^T V^{-1/2} (l_{\beta_{n,0}}(X^{(i)}), h^T s_{\beta_{n,0}}(X^{(i)}))^T \rightsquigarrow \mathcal{N}(0, 1).$$

Hence, using the likelihood expansion (37) and by the asymptotic linearity (19), we conclude that

$$a^T V^{-1/2} (\sqrt{n}(T_n - g(\beta_{n,0})), \Lambda_n + \frac{1}{2} h^T I(\beta_{n,0}) h) \rightsquigarrow a^T Z, \text{ where } Z \sim \mathcal{N}(0, I_2).$$

Then by the Wold device we have

$$Z_n := V^{-1/2} \begin{pmatrix} \sqrt{n}(T_n - g(\beta_{n,0})) \\ \Lambda_n + \frac{1}{2} h^T I(\beta_{n,0}) h \end{pmatrix} \overset{\beta_{n,0}}{\rightsquigarrow} \mathcal{N}_2(0, I) \sim Z.$$

Now let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be bounded and continuous. We may rewrite

$$\begin{aligned} & \mathbb{E}_{\beta_{n,0}+h/\sqrt{n}} f \left( \frac{\sqrt{n}(T_n - g(\beta_{n,0})) - v_{12}}{\sqrt{v_{11}}} \right) \\ &= \mathbb{E}_{\beta_{n,0}} f \left( \frac{\sqrt{n}(T_n - g(\beta_{n,0})) - v_{12}}{\sqrt{v_{11}}} \right) e^{\Lambda_n} \\ &= \mathbb{E}_{\beta_{n,0}} f(X_{n,1}) e^{X_{n,2}}, \end{aligned}$$

where  $X_n := (X_{n,1}, X_{n,2}) = \psi(Z_n)$  for the function  $\psi$  given by

$$\psi(x_1, x_2) = \begin{pmatrix} 1/\sqrt{v_{11}} & 0 \\ 0 & 1 \end{pmatrix} \left[ V^{1/2}(x_1, x_2)^T + \begin{pmatrix} -v_{12} \\ -v_{22}/2 \end{pmatrix} \right].$$

Similarly, define  $U_n$  as follows

$$\begin{aligned} U_n &:= (U_{n,1}, U_{n,2}) = \psi(Z) \\ &\sim \begin{pmatrix} 1/\sqrt{v_{11}} & 0 \\ 0 & 1 \end{pmatrix} \left[ V^{1/2}\mathcal{N}(0, I_2) + \begin{pmatrix} -v_{12} \\ -v_{22}/2 \end{pmatrix} \right] \\ &= \mathcal{N} \left( \begin{pmatrix} -\frac{v_{12}}{\sqrt{v_{11}}} \\ -\frac{v_{22}}{2} \end{pmatrix}, \begin{pmatrix} 1 & \frac{v_{12}}{\sqrt{v_{11}}} \\ \frac{v_{12}}{\sqrt{v_{11}}} & v_{22} \end{pmatrix} \right). \end{aligned}$$

Since we know that  $Z_n \rightsquigarrow Z$ , we hope that in some sense  $X_n = \psi(Z_n)$  is close to  $U_n = \psi(Z)$ . Note that the function  $\psi$  depends on  $n$ , so we cannot directly apply the Portmanteau Lemma.

We aim to apply Lemma 16 with  $X_n = \psi(Z_n)$  and  $U_n = \psi(Z)$  defined above and with the function  $g(x_1, x_2) = f(x_1)e^{x_2}$ . By Lemma 23, we have that

$$\lim_{M \rightarrow \infty} \lim_{n \rightarrow \infty} \mathbb{E} |\min(0, M - e^{U_{n,2}})| = 0.$$

Hence we get by the second part of Lemma 16

$$\lim_{n \rightarrow \infty} \mathbb{E} g(X_n) - \mathbb{E} g(U_n) = 0.$$

Next we calculate  $\mathbb{E} g(U_n)$ . We have

$$\mathbb{E}_{\beta_{n,0}} g(U_n) = \mathbb{E} f(U_{n,1}) e^{U_{n,2}} = \int_{\mathbb{R}^2} f(u_1) e^{u_2} f_{U_n}(u_1, u_2) du,$$

where  $f_Y$  denotes the density of a random variable  $Y$ . We use Lemma 21 to obtain that  $f_{U_n}(u) e^{u_2} = f_Y(u)$ , where

$$Y \sim \mathcal{N} \left( \begin{pmatrix} 0 \\ v_{22}/2 \end{pmatrix}, \begin{pmatrix} 1 & \frac{v_{12}}{\sqrt{v_{11}}} \\ \frac{v_{12}}{\sqrt{v_{11}}} & v_{22} \end{pmatrix} \right).$$

Hence

$$\mathbb{E}_{\beta_{n,0}} g(U_n) = \int_{\mathbb{R}^2} f(u_1) f_Y(u_1, u_2) du = \mathbb{E} f(Y_1),$$

where  $Y \sim \mathcal{N}(0, 1)$ . Hence for any bounded continuous function  $f$  we have shown

$$\lim_{n \rightarrow \infty} |\mathbb{E}_{\beta_{n,0} + h/\sqrt{n}} f\left(\frac{\sqrt{n}(T_n - g(\beta_{n,0})) - v_{12}}{\sqrt{v_{11}}}\right) - \mathbb{E} f(Y)| = 0.$$

By the Portmanteau Lemma (note that  $Y$  in the above display does not depend on  $n$ ), we thus have

$$\frac{\sqrt{n}(T_n - g(\beta_{n,0})) - v_{12}}{\sqrt{v_{11}}} \xrightarrow{\beta_{n,0} + h/\sqrt{n}} \mathcal{N}(0, 1).$$

Therefore, by the differentiability assumption (D1) on  $g$ , we get

$$\frac{\sqrt{n}(T_n - g(\beta_{n,0} + h/\sqrt{n})) + h^T \dot{g}(\beta_{n,0}) - P_{\beta_{n,0}} l_{\beta_{n,0}} h^T s_{\beta_{n,0}}}{V_{\beta_{n,0}}^{1/2}} \xrightarrow{\beta_{n,0} + h/\sqrt{n}} \mathcal{N}(0, 1).$$

□

## APPENDIX A: PROOFS FOR SECTION 14

Before proving Lemmas 9, 10, 11 and 12 we first recall a version of Theorem 2.4 from van de Geer et al. (2014). We denote  $\eta_j := X_j - X_{-j} \gamma_j^0$ ,  $j = 1, \dots, p$ .

**THEOREM 10** (a version of Theorem 2.4 in van de Geer et al. (2014)). *Suppose that conditions (A2\*) are satisfied and assume that  $s_j \log p/n = o(1)$ . Consider the nodewise regression estimator  $\hat{\Theta}_j$  and the corresponding  $\hat{\tau}_j^2$  with  $\lambda_j = \lambda \geq \tau \sqrt{\log p/n}$  for  $j = 1, \dots, p$ . Then for  $\tau > 1$ , on the set*

$$\begin{aligned} \mathcal{T}_j &:= \{ \|X_{-j}^T \eta_j\|_\infty / n \leq c\tau \sqrt{\log p/n}, \\ &\quad \|\hat{\Sigma}_{-j, -j} - \Sigma_{-j, -j}^0\|_\infty \leq c\tau \sqrt{\log p/n}, \\ &\quad |\eta_j^T \eta_j / n - \tau_j^2| \leq c\tau \sqrt{\log p/n} \}, \end{aligned}$$

(where  $c$  is some sufficiently large constant), we have the following claims for  $j = 1, \dots, p$ ,

$$\|\hat{\gamma}_j - \gamma_j^0\|_1 \leq C_\tau s_j \sqrt{\log p/n}, \quad |\hat{\tau}_j^2 - \tau_j^2| \leq C_\tau \sqrt{s_j \log p/n},$$

$$\|\hat{\Theta}_j - \Theta_j^0\|_1 \leq C_\tau s_j \sqrt{\log p/n},$$

for some constant  $C_\tau > 0$ . Moreover, for some constant  $c_1 > 0$  we have

$$P(\mathcal{T}_j^c) \leq c_1 (2p)^{-\tau^2}.$$



PROOF OF LEMMA 9. The proof of this lemma is essentially the same as the proof of Theorem 1, but we need to make adjustments to obtain an oracle bound for the expectation of the *maximum* of  $\ell_1$ -errors of  $p$  Lasso estimators, i.e. we show an oracle bound for

$$(\mathbb{E} \max_{j=1,\dots,p} \|\hat{\gamma}_j - \gamma_j\|_1^k)^{1/k}.$$

First we summarize the oracle inequality for the nodewise regression which holds with high probability. Let  $\eta_j := X_j - X_{-j}\gamma_j^0$  for  $j = 1, \dots, p$ , then  $\eta_j$  is a sub-Gaussian random vector with a universal constant, since  $\Lambda_{\max}(\Theta_0) = 1/\Lambda_{\min}(\Sigma_0) = \mathcal{O}(1)$ . Under  $1/\Lambda_{\min}(\Sigma_0) = \mathcal{O}(1)$ , one can check that for some universal constant  $L > 0$ , it holds  $\Lambda_{\min}(\Sigma_{-j,-j}^0) \geq L$ . Let  $\mathcal{T}_j$  be as in Theorem 10. Then on  $\mathcal{T} := \cap_{j=1}^p \mathcal{T}_j$  it holds  $\max_{j=1,\dots,p} \|\hat{\gamma}_j - \gamma_j^0\|_1 \leq 16\lambda_j \max_{j=1,\dots,p} s_j/L$ .

We now proceed to show that the oracle inequality for the Lasso holds also in expectation. We follow the steps of the proof of Theorem 1 to get

$$\mathbb{E} \max_{j=1,\dots,p} (\|\epsilon_j\|_n^2/\lambda_j + 2\|\gamma_j^0\|_1)^k \leq \mathbb{E} \max_{j=1,\dots,p} 2^{k-1} \left( (\|\epsilon_j\|_n^2/\lambda_j)^k + (2\|\gamma_j^0\|_1)^k \right)$$

By Lemma 5 we have

$$\mathbb{E} \max_{j=1,\dots,p} (\|\epsilon_j\|_n^2)^k = \mathcal{O}(1).$$

Next observe that by assumption on the eigenvalues of  $\Theta_0$ , we have  $\|\gamma_j^0\|_2 = \mathcal{O}(1)$  and hence

$$\|\gamma_j^0\|_1^k \leq (\max_{j=1,\dots,p} \sqrt{s_j} \|\gamma_j^0\|_2)^k \leq \mathcal{O}(\max_{j=1,\dots,p} s_j^{k/2}).$$

Further steps again follow the steps of the proof of Theorem 1. Thus we obtain

$$\mathbb{E} \max_{j=1,\dots,p} \|\hat{\gamma}_j - \gamma_j^0\|_1^k = \mathcal{O}(\max_{j=1,\dots,p} s_j^k \lambda_j^k),$$

where we chose  $\tau$  sufficiently large (this is possible since  $k$  is fixed) so that

$$\max_{j=1,\dots,p} s_j^{k/2} \lambda_j^{-k} p^{-\tau^2/2} = \mathcal{O}(\max_{j=1,\dots,p} s_j^k \lambda_j^k).$$

Hence we conclude that

$$(38) \quad (\mathbb{E} \max_{j=1,\dots,p} \|\hat{\gamma}_j - \gamma_j^0\|_1^k)^{1/k} = \mathcal{O}(\max_{j=1,\dots,p} s_j \lambda_j).$$

□

PROOF OF LEMMA 10. Before proving statements 1) and 2) of the lemma, we first prove that  $\mathbb{E} \max_{j=1,\dots,p} 1/(\hat{\tau}_j^2)^k = \mathcal{O}(1)$ . Throughout the proof, we use the notation

$$\hat{\Gamma}_j := (-\hat{\gamma}_{j,1}, \dots, -\hat{\gamma}_{j,j-1}, 1, -\hat{\gamma}_{j,j+1}, \dots, -\hat{\gamma}_{j,p}).$$

**Proof of  $\mathbb{E} \max_{j=1,\dots,p} 1/(\hat{\tau}_j^2)^k = \mathcal{O}(1)$ :**

We first show the rough bound  $\mathbb{E} \max_{j=1,\dots,p} \frac{1}{(\hat{\tau}_j^2)^k} = \mathcal{O}(pn^{k/2})$ . First observe that for each  $j = 1, \dots, p$  it holds for  $t \geq 0$

$$\begin{aligned} P(\hat{\tau}_j^2 \leq t) &= P(\hat{\Gamma}_j^T \hat{\Sigma} \hat{\Gamma}_j + \lambda_j \|\hat{\gamma}_j\|_1 \leq t) \\ &\leq P(\hat{\Gamma}_j^T \hat{\Sigma} \hat{\Gamma}_j \leq t \wedge \lambda_j \|\hat{\gamma}_j\|_1 \leq t) \end{aligned}$$

Using the following lower bound

$$\begin{aligned} \hat{\Gamma}_j^T \hat{\Sigma} \hat{\Gamma}_j &= \hat{\Sigma}_{jj} - 2\hat{\Sigma}_{j,-j}^T \hat{\gamma}_j + \hat{\gamma}_j^T \hat{\Sigma}_{-j,-j} \hat{\gamma}_j \\ &\geq \hat{\Sigma}_{jj} - 2|\hat{\Sigma}_{j,-j}^T \hat{\gamma}_j| + \hat{\gamma}_j^T \hat{\Sigma}_{-j,-j} \hat{\gamma}_j \\ &\geq \hat{\Sigma}_{jj} - 2\|\hat{\Sigma}_{j,-j}\|_\infty \|\hat{\gamma}_j\|_1 + \hat{\gamma}_j^T \hat{\Sigma}_{-j,-j} \hat{\gamma}_j \\ &\geq \hat{\Sigma}_{jj} - 2\hat{\Sigma}_{jj} t / \lambda_j \\ &= \hat{\Sigma}_{jj} (1 - 2t / \lambda_j), \end{aligned}$$

we obtain that

$$\begin{aligned} P(\hat{\Gamma}_j^T \hat{\Sigma} \hat{\Gamma}_j \leq t \wedge \lambda_j \|\hat{\gamma}_j\|_1 \leq t) &\leq P(\hat{\Sigma}_{jj} (1 - 2t / \lambda_j) \leq t \wedge \lambda_j \|\hat{\gamma}_j\|_1 \leq t) \\ &\leq P(\hat{\Sigma}_{jj} (1 - 2t / \lambda_j) \leq t). \end{aligned}$$

Next we use concentration results for  $\hat{\Sigma}_{jj}$  around its mean under the sub-Gaussianity assumption on  $X$ . For  $t \leq \lambda_j/4$  it holds that  $1 - 2t/\lambda_j \geq 1/2$  and thus

$$P(\hat{\Sigma}_{jj} (1 - 2t / \lambda_j) \leq t) = P(\hat{\Sigma}_{jj} - \Sigma_{jj}^0 \leq 2t - \Sigma_{jj}^0).$$

For  $0 < t < \lambda_j/4$  and  $n$  sufficiently large it holds that  $2t - \Sigma_{jj}^0 < 0$  (by the minimal eigenvalue condition on  $\Sigma_0$ ) and thus

$$\begin{aligned} P(\hat{\Sigma}_{jj} (1 - 2t / \lambda_j) \leq t) &= P(\hat{\Sigma}_{jj} - \Sigma_{jj}^0 \leq 2t - \Sigma_{jj}^0) \\ &\leq P(|\hat{\Sigma}_{jj} - \Sigma_{jj}^0| \geq |2t - \Sigma_{jj}^0|) \\ &\leq e^{-c(\Sigma_{jj}^0 - 2t)\sqrt{n/\log pn}}, \end{aligned}$$

for some constant  $c > 0$ . Hence collecting the above inequalities, we have so far shown that for any  $0 < t < \lambda_j/4$  and  $n$  sufficiently large it holds

$$(39) \quad P(\hat{\tau}_j^2 \leq t) \leq e^{-c(\Sigma_{jj}^0 - 2t)\sqrt{n/\log p}}.$$

Then by rewriting the expectation as an integral

$$\begin{aligned} \mathbb{E} \max_{j=1,\dots,p} \frac{1}{(\hat{\tau}_j^2)^k} &= \int_0^\infty P\left(\max_{j=1,\dots,p} 1/(\hat{\tau}_j^2)^k > x\right) dx \\ &= \int_0^\infty \max_{j=1,\dots,p} pP(1/(\hat{\tau}_j^2)^k > x) dx \\ &= p \int_0^1 \max_{j=1,\dots,p} P(1/(\hat{\tau}_j^2)^k > x) dx \\ &\quad + p \int_1^{(\lambda_j/4)^{-k}} \max_{j=1,\dots,p} P(1/(\hat{\tau}_j^2)^k > x) dx \\ &\quad + p \int_{(\lambda_j/4)^{-k}}^\infty \max_{j=1,\dots,p} P(1/(\hat{\tau}_j^2)^k > x) dx \\ &\leq p + p(\lambda_j/4)^{-k} + \underbrace{p \int_{(\lambda_j/4)^{-k}}^\infty P(1/(\hat{\tau}_j^2)^k > x) dx}_{ii} \end{aligned}$$

Next we calculate an upper bound on *ii*.

$$\begin{aligned} ii &= \int_{(\lambda_j/4)^{-k}}^\infty P(1/(\hat{\tau}_j^2)^k > x) dx = \int_{(\lambda_j/4)^{-k}}^\infty P(1/\hat{\tau}_j^2 > x^{1/k}) dx \\ &= \int_{(\lambda_j/4)^{-1}}^\infty P(\hat{\tau}_j^2 < x^{-1}) dx \end{aligned}$$

Now we can use the bound (39) since  $x^{-1} \leq \lambda_j/4$ . Using the bound and by standard calculations, we obtain

$$\int_{(\lambda_j/4)^{-1}}^\infty P(\hat{\tau}_j^2 < x^{-1}) dx \leq \int_{(\lambda_j/4)^{-1}}^\infty e^{-c(\Sigma_{11}^0 - 2/x)\sqrt{n/\log pn}} dx = o(1).$$

Hence we obtain the rough bound

$$\mathbb{E} \max_{j=1,\dots,p} \frac{1}{(\hat{\tau}_j^2)^k} = \mathcal{O}\left(p(\lambda_j/4)^{-k}\right) = \mathcal{O}(pn^{k/2}/(\log p)^{k/2}).$$

Define, for  $\tau > 1$  and  $j = 1, \dots, p$ , the sets

$$\begin{aligned} \mathcal{T}_j &:= \{ \|X_{-j}^T \eta_j\|_\infty / n \leq c\tau \sqrt{\log p/n}, \\ &\quad \|\hat{\Sigma}_{-j,-j} - \Sigma_{-j,-j}^0\|_\infty \leq c\tau \sqrt{\log p/n}, \\ &\quad \|\eta_j^T \eta_j / n - \tau_j^2\|_\infty \leq c\tau \sqrt{\log p/n} \}, \end{aligned}$$

(where  $c$  is some sufficiently large constant). Then by Theorem 10, on  $\mathcal{T} = \cap_{j=1}^p \mathcal{T}_j$  we have  $\max_{j=1,\dots,p} 1/(\hat{\tau}_j^2)^k \leq C_\tau$  for some constant  $C_\tau > 0$ . But then and by the Cauchy-Schwarz inequality

$$\begin{aligned} \mathbb{E} \max_{j=1,\dots,p} \frac{1}{(\hat{\tau}_j^2)^k} &= \mathbb{E} \max_{j=1,\dots,p} \frac{1}{(\hat{\tau}_j^2)^k} 1_{\mathcal{T}} + \mathbb{E} \max_{j=1,\dots,p} \frac{1}{(\hat{\tau}_j^2)^k} 1_{\mathcal{T}^c} \\ &\leq \mathcal{O}(1) + \sqrt{\mathcal{O}(pn^{k/2})} \sqrt{2p(2p)^{-\tau^2/2}} = \mathcal{O}(1), \end{aligned}$$

where we chose  $\tau$  sufficiently large.

**Proof of part 1)** First we show that  $\mathbb{E}(\hat{\tau}_j^2)^k = \mathcal{O}(1)$ . We have

$$\begin{aligned} \hat{\tau}_j^2 &= \|X_j - X_{-j}\hat{\gamma}_j\|_n^2 + \lambda_j \|\hat{\gamma}_j\|_1 \\ &= \hat{\Gamma}_j^T \hat{\Sigma} \hat{\Gamma}_j / n + \lambda_j \|\hat{\gamma}_j\|_1 \\ &\leq \|\hat{\Gamma}_j\|_1^2 \|\hat{\Sigma}\|_\infty + \lambda_j \|\hat{\gamma}_j\|_1. \end{aligned}$$

Hence by basic calculations

$$\begin{aligned} \mathbb{E} \max_{j=1,\dots,p} (\hat{\tau}_j^2)^k &\leq \mathbb{E} \max_{j=1,\dots,p} \left[ \|\hat{\Gamma}_j\|_1^2 \|\hat{\Sigma}\|_\infty + \lambda_j \|\hat{\gamma}_j\|_1 \right]^k \\ &\leq 2^{k-1} \mathbb{E} \max_{j=1,\dots,p} \left[ (\|\hat{\Gamma}_j\|_1^2 \|\hat{\Sigma}\|_\infty)^k + (\lambda_j \|\hat{\gamma}_j\|_1)^k \right]. \end{aligned}$$

We have

$$\begin{aligned} \mathbb{E} \max_{j=1,\dots,p} \|\hat{\gamma}_j\|_1^k &\leq \mathbb{E} \max_{j=1,\dots,p} (\|\hat{\gamma}_j - \gamma_j^0\|_1 + \|\gamma_j^0\|_1)^k \\ &\leq \mathbb{E} \max_{j=1,\dots,p} 2^{k-1} (\|\hat{\gamma}_j - \gamma_j^0\|_1^k + \|\gamma_j^0\|_1^k) \\ &= \mathcal{O}(s_j^{k/2}). \end{aligned}$$

Hence

$$\begin{aligned} \mathbb{E} \max_{j=1,\dots,p} (\hat{\tau}_j^2)^k &\leq \mathbb{E} \max_{j=1,\dots,p} \left[ \|\hat{\Gamma}_j\|_1^2 \|\hat{\Sigma}\|_\infty + \lambda_j \|\hat{\gamma}_j\|_1 \right]^k \\ &= \mathcal{O}(s_j^{2k}). \end{aligned}$$

Hence, by Theorem 10, on  $\mathcal{T}$  we have that  $\max_{j=1,\dots,p} \hat{\tau}_j^2 = \mathcal{O}(1)$ , hence it follows

$$\mathbb{E} \max_{j=1,\dots,p} (\hat{\tau}_j^2)^k = \mathbb{E} \max_{j=1,\dots,p} (\hat{\tau}_j^2)^k 1_{\mathcal{T}} + \mathbb{E} \max_{j=1,\dots,p} (\hat{\tau}_j^2)^k 1_{\mathcal{T}^c} = \mathcal{O}(1).$$

We have under  $1/\Lambda_{\min}(\Theta_0) = \mathcal{O}(1)$  that  $\tau_j^2 = 1/\Theta_{jj}^0 = \mathcal{O}(1)$  and hence

$$\mathbb{E} \max_{j=1,\dots,p} |\hat{\tau}_j^2 - \tau_j^2|^k = \mathcal{O}(\mathbb{E} \max_{j=1,\dots,p} (\hat{\tau}_j^2)^k + (\tau_j^2)^k) = \mathcal{O}(1).$$

We can then apply the same procedure as before to get

$$\begin{aligned} \mathbb{E} \max_{j=1,\dots,p} |\hat{\tau}_j^2 - \tau_j^2|^k &= \mathbb{E} \max_{j=1,\dots,p} |\hat{\tau}_j^2 - \tau_j^2|^k 1_{\mathcal{T}} + \mathbb{E} \max_{j=1,\dots,p} |\hat{\tau}_j^2 - \tau_j^2|^k 1_{\mathcal{T}^c} \\ &= \mathcal{O}\left(\max_{j=1,\dots,p} \sqrt{s_j \lambda_j}\right). \end{aligned}$$

**Proof of part 2)** First we have

$$\begin{aligned} &\mathbb{E} \max_{j=1,\dots,p} \left| \frac{1}{\hat{\tau}_j^2} - \frac{1}{\tau_j^2} \right|^k \\ &\leq \mathbb{E} \max_{j=1,\dots,p} \frac{|\hat{\tau}_j^2 - \tau_j^2|^k}{(\hat{\tau}_j^2)^k (\tau_j^2)^k} \\ &\leq \max_{j=1,\dots,p} 1/(\tau_j^2)^k \sqrt{\mathbb{E} \max_{j=1,\dots,p} |\hat{\tau}_j^2 - \tau_j^2|^{2k}} \sqrt{\mathbb{E} \max_{j=1,\dots,p} 1/(\hat{\tau}_j^2)^{2k}}. \end{aligned}$$

Using that  $\mathbb{E} \max_{j=1,\dots,p} 1/(\hat{\tau}_j^2)^{2k} = \mathcal{O}(1)$  and Part 1), we obtain the claim.  $\square$

**PROOF OF LEMMA 11.** For some  $\tau > 0$  and each  $j = 1, \dots, p$  define the sets

$$\begin{aligned} \mathcal{T}_j &:= \{ \|X_{-j}^T \eta_j\|_{\infty}/n \leq c\tau \sqrt{\log p/n}, \\ &\quad \|\hat{\Sigma}_{-j,-j} - \Sigma_{-j,-j}^0\|_{\infty} \leq c\tau \sqrt{\log p/n}, \\ &\quad \|\eta_j^T \eta_j/n - \tau_j^2\|_{\infty} \leq c\tau \sqrt{\log p/n} \}, \end{aligned}$$

By Theorem 10, when  $\lambda_j \geq c\tau \sqrt{\log p/n}$  uniformly in  $j$ , we have on the set  $\cap_{i=1}^p \mathcal{T}_j$  that

$$\max_{j=1,\dots,p} \|\hat{\Theta}_j - \Theta_j^0\|_1 \leq C_{\tau} \max_{j=1,\dots,p} s_j \lambda_j, \quad \max_{j=1,\dots,p} |\hat{\tau}_j^2 - \tau_j^2| \leq C_{\tau} \max_{j=1,\dots,p} \sqrt{s_j \lambda_j},$$

for some  $C_{\tau} > 0$ . Next we rewrite

$$\mathbb{E} \max_{j=1,\dots,p} \|\hat{\Theta}_j - \Theta_j^0\|_1^k = \mathbb{E} \max_{j=1,\dots,p} \|\hat{\Theta}_j - \Theta_j^0\|_1^k 1_{\mathcal{T}} + \mathbb{E} \max_{j=1,\dots,p} \|\hat{\Theta}_j - \Theta_j^0\|_1^k 1_{\mathcal{T}^c}.$$

Then

$$\begin{aligned}
& \mathbb{E} \max_{j=1,\dots,p} \|\hat{\Theta}_j - \Theta_j^0\|_1^k \\
& \leq \mathbb{E} \max_{j=1,\dots,p} [\|\hat{\gamma}_j - \gamma_j^0\|_1 / \hat{\tau}_j^2 + \|\gamma_j^0\|_1 |1/\hat{\tau}_j^2 - 1/\tau_j^2|]^k \\
& \leq \mathbb{E} \max_{j=1,\dots,p} 2^{k-1} [(\|\hat{\gamma}_j - \gamma_j^0\|_1 / \hat{\tau}_j^2)^k + (\|\gamma_j^0\|_1 |1/\hat{\tau}_j^2 - 1/\tau_j^2|)^k] \\
& = \mathcal{O}((s_j \lambda_j)^k),
\end{aligned}$$

where in the last display we used Lemmas 9 and 10.  $\square$

PROOF OF LEMMA 12. Let  $\eta_j := X_j - X_{-j}\gamma_j^0$ . The Karush-Kuhn-Tucker corresponding to the optimization problem (6) give

$$\hat{\Sigma}_{-j,-j}\hat{\gamma}_j + \lambda_j\hat{Z}_j = X_{-j}^T\eta_j/n,$$

where  $\hat{Z}_{j,i} = \text{sign}(\hat{\gamma}_{j,i})$  if  $\hat{\gamma}_{j,i} \neq 0$  and  $\hat{Z}_{j,i} \in [-1, 1]$  otherwise, for  $i = 1, \dots, p$ . Rearranging them, we obtain

$$(40) \quad \Sigma_{-j,-j}^0(\hat{\gamma}_j - \gamma_j^0) + \lambda_j\hat{Z}_j = X_{-j}^T\eta_j/n + (\Sigma_{-j,-j}^0 - \hat{\Sigma}_{-j,-j})(\hat{\gamma}_j - \gamma_j^0).$$

Firstly,

$$\begin{aligned}
\|\Sigma_{-j,-j}^0(\hat{\gamma}_j - \gamma_j^0)\|_2^2 & \leq \Lambda_{\max}(\Sigma_{-j,-j}^0)\|X_{-j}(\hat{\gamma}_j - \gamma_j^0)\|_n^2 \\
& \quad + \Lambda_{\max}(\Sigma_{-j,-j}^0)\|\hat{\gamma}_j - \gamma_j^0\|_1^2\|\hat{\Sigma} - \Sigma_0\|_\infty.
\end{aligned}$$

Secondly,

$$\|(\Sigma_{-j,-j}^0 - \hat{\Sigma}_{-j,-j})(\hat{\gamma}_j - \gamma_j^0)\|_\infty \leq \|\Sigma_{-j,-j}^0 - \hat{\Sigma}_{-j,-j}\|_\infty\|\hat{\gamma}_j - \gamma_j^0\|_1.$$

Denote  $\hat{s}_j := \|\hat{\gamma}_j\|_0$ . On the set

$$\mathcal{T}_j := \{\|X_{-j}^T\eta_j/n\|_\infty \leq \lambda_0, \|(\Sigma_{-j,-j}^0 - \hat{\Sigma}_{-j,-j})(\hat{\gamma}_j - \gamma_j^0)\|_\infty \leq \lambda_0\},$$

we have

$$\begin{aligned}
& \|\lambda_j\hat{Z}_j + X_{-j}^T\eta_j/n + (\Sigma_{-j,-j}^0 - \hat{\Sigma}_{-j,-j})(\hat{\gamma}_j - \gamma_j^0)\|_2^2 \\
& = \sum_{i=1}^p |\lambda_j\hat{Z}_{j,i} + X_i^T\eta_j + e_i^T(\Sigma_{-j,-j}^0 - \hat{\Sigma}_{-j,-j})(\hat{\gamma}_j - \gamma_j^0)|^2 \\
& \geq (\lambda_j - 2\lambda_0)^2\hat{s}.
\end{aligned}$$

Combining the above observations, we obtain

$$\begin{aligned}
& \Lambda_{\max}(\Sigma_{-j,-j}^0) \|X_{-j}(\hat{\gamma}_j - \gamma_j^0)\|_n^2 + \Lambda_{\max}(\Sigma_{-j,-j}^0) \|\hat{\gamma}_j - \gamma_j^0\|_1^2 \|\hat{\Sigma} - \Sigma_0\|_\infty \\
& \geq \|\Sigma_{-j,-j}^0(\hat{\gamma}_j - \gamma_j^0)\|_2^2 \\
& = \|-\lambda \hat{Z}_j + X_{-j}^T \eta_j / n + (\Sigma_{-j,-j}^0 - \hat{\Sigma}_{-j,-j})(\hat{\gamma}_j - \gamma_j^0)\|_2^2 \\
& \geq (\lambda_j - 2\lambda_0)^2 \hat{s}.
\end{aligned}$$

Hence on the set  $\mathcal{T}$ ,

$$\hat{s}_j \leq \frac{\Lambda_{\max}(\Sigma_{-j,-j}^0) \|X_{-j}(\hat{\gamma}_j - \gamma_j^0)\|_n^2 + \Lambda_{\max}(\Sigma_{-j,-j}^0) \|\hat{\gamma}_j - \gamma_j^0\|_1^2 \|\hat{\Sigma} - \Sigma_0\|_\infty}{(\lambda_j - 2\lambda_0)^2}.$$

Finally, taking  $\lambda_0 := c\sqrt{\log p/n}$  for some  $c > 0$  sufficiently large and taking  $\lambda \geq 3\lambda_0$  and under the assumption  $\Lambda_{\max}(\Sigma_0) = \mathcal{O}(1)$ , we obtain

$$\hat{s}_j = \mathcal{O}_P(s_j).$$

□

## APPENDIX B: ADDITIONAL PROOFS FOR SECTION 16

In this section we give Lemma 20 and its proof, but we need the following auxiliary Lemmas 17, 18, 19.

LEMMA 17. *Let  $x \sim \mathcal{N}(0_p, \Sigma_0)$  and let  $\Theta_0 = \Sigma_0^{-1}$ . Then for any  $t \in \mathbb{R}$  and  $A \in \mathbb{R}^{p \times p}$  such that  $\Theta_0 - 2tA$  is symmetric and positive definite it holds*

$$\mathbb{E}_{\Theta_0} e^{tx^T Ax} = \left( \frac{\det(\Theta_0)}{\det(\Theta_0 - 2tA)} \right)^{1/2}.$$

PROOF OF LEMMA 17. By direct calculation, we obtain

$$\begin{aligned}
\mathbb{E}_{\Theta_0} e^{tx^T Ax} &= \int_{\mathbb{R}^p} \frac{\det(\Theta_0)^{1/2}}{(2\pi)^{p/2}} e^{-\frac{1}{2}x^T \Theta_0 x} e^{tx^T Ax} dx \\
&= \int_{\mathbb{R}^p} \frac{\det(\Theta_0)^{1/2}}{(2\pi)^{p/2}} e^{-\frac{1}{2}x^T (\Theta_0 - 2tA)x} dx \\
&= \int_{\mathbb{R}^p} \frac{\det(\Theta_0)^{1/2} \det(\Theta_0 - 2tA)^{1/2}}{(2\pi)^{p/2} \det(\Theta_0 - 2tA)^{1/2}} e^{-\frac{1}{2}x^T (\Theta_0 - 2tA)x} dx \\
&= \frac{\det(\Theta_0)^{1/2}}{\det(\Theta_0 - 2tA)^{1/2}}.
\end{aligned}$$

□

LEMMA 18. *Suppose that  $\Theta_0 + H/\sqrt{m_n}$  is a symmetric positive definite matrix. Let  $p_{\Theta_0}$  be the joint density of the random sample  $x_1, \dots, x_n$ , where each  $x_i$  is an  $\mathcal{N}(0, \Theta_0^{-1})$ -distributed random vector. Then it holds*

$$\begin{aligned} & \mathbb{E}_{\Theta_0} \left( \frac{p_{\Theta_0+H/\sqrt{m_n}}(x)}{p_{\Theta_0}(x)} - 1 - n\text{tr}((\hat{\Sigma} - \Sigma_0)H/\sqrt{m_n}) \right)^2 \\ &= \frac{\det(\Theta_0 + H/\sqrt{m_n})^n}{\det(\Theta_0)^n} \left( \frac{\det(\Theta_0)}{\det(\Theta_0 + 2H)} \right)^{n/2} - 1 + \sum_{i=1}^n \text{var}(x_i^T H x_i) \\ & \quad - 2n\text{tr}[(\Theta_0 + H)^{-1} - \Sigma_0]H. \end{aligned}$$

PROOF OF LEMMA 18. The density is given by

$$p_{\Theta_0}(x_1, \dots, x_n) = \frac{\det(\Theta_0)^{n/2}}{(2\pi)^{np/2}} e^{-\frac{1}{2} \sum_{i=1}^n x_i^T \Theta_0 x_i}$$

For simplicity of notation, denote  $U := H/\sqrt{m_n}$ . Then we have

$$\frac{p_{\Theta_0+U}(x)}{p_{\Theta_0}(x)} - 1 = \frac{\det(\Theta_0 + U)^{n/2} e^{-\frac{1}{2} \sum_{i=1}^n x_i^T U x_i}}{\det(\Theta_0)^{n/2}} - 1$$

The score function is given by  $s_{\Theta_0}(x) = n(\hat{\Sigma} - \Sigma_0)/2$ . Let

$$Z := \text{vec}(U)^T \text{vec}(s_{\Theta_0}(x)) = \text{tr}(n(\hat{\Sigma} - \Sigma_0)U) = \sum_{i=1}^n x_i^T U x_i - n\text{tr}(\Sigma_0 U).$$

First observe that

$$\mathbb{E}_{\Theta_0} Z^2 = \text{var}\left(\sum_{i=1}^n x_i^T U x_i\right) = \sum_{i=1}^n \text{var}(x_i^T U x_i).$$

We have

$$\begin{aligned} & \mathbb{E}_{\Theta_0} \left( \frac{p_{\Theta_0+U}(x)}{p_{\Theta_0}(x)} - 1 - Z \right)^2 \\ &= \mathbb{E}_{\Theta_0} \left( \frac{\det(\Theta_0 + U)^{n/2} e^{-\frac{1}{2} \sum_{i=1}^n x_i^T U x_i}}{\det(\Theta_0)^{n/2}} - 1 - Z \right)^2 \\ &= \frac{\det(\Theta_0 + U)^n}{\det(\Theta_0)^n} \mathbb{E}_{\Theta_0} e^{-\sum_{i=1}^n x_i^T U x_i} + 1 + \mathbb{E}_{\Theta_0} Z^2 + \\ & \quad - 2 \frac{\det(\Theta_0 + U)^{n/2}}{\det(\Theta_0)^{n/2}} \mathbb{E}_{\Theta_0} e^{-\frac{1}{2} \sum_{i=1}^n x_i^T U x_i} + 2 \mathbb{E}_{\Theta_0} Z \\ & \quad - 2 \frac{\det(\Theta_0 + U)^{n/2}}{\det(\Theta_0)^{n/2}} \mathbb{E}_{\Theta_0} Z e^{-\frac{1}{2} \sum_{i=1}^n x_i^T U x_i}. \end{aligned}$$



Using Lemma 17, since  $\Theta_0 + U/2$  is a symmetric positive definite matrix, we obtain

$$\begin{aligned}
\mathbb{E}_{\Theta_0} \left( \frac{p_{\Theta_0+U}(x)}{p_{\Theta_0}(x)} - 1 - Z \right)^2 &= \frac{\det(\Theta_0 + U)^n}{\det(\Theta_0)^n} \left( \frac{\det(\Theta_0)}{\det(\Theta_0 + 2U)} \right)^{n/2} + 1 \\
&\quad + \sum_{i=1}^n \text{var}_{\Theta_0}(x_i^T U x_i) \\
&\quad - 2 \frac{\det(\Theta_0 + U)^{n/2}}{\det(\Theta_0)^{n/2}} \left( \frac{\det(\Theta_0)}{\det(\Theta_0 + U)} \right)^{n/2} \\
&\quad - 2 \frac{\det(\Theta_0 + U)^{n/2}}{\det(\Theta_0)^{n/2}} \mathbb{E}_{\Theta_0} Z e^{-\frac{1}{2} \sum_{i=1}^n x_i^T U x_i} \\
&= \frac{\det(\Theta_0 + U)^n}{\det(\Theta_0)^n} \left( \frac{\det(\Theta_0)}{\det(\Theta_0 + 2U)} \right)^{n/2} - 1 \\
&\quad + \sum_{i=1}^n \text{var}_{\Theta_0}(x_i^T U x_i) \\
&\quad - \underbrace{2 \frac{\det(\Theta_0 + U)^{n/2}}{\det(\Theta_0)^{n/2}} \mathbb{E}_{\Theta_0} Z e^{-\frac{1}{2} \sum_{i=1}^n x_i^T U x_i}}_i.
\end{aligned}$$

Next we calculate  $i$ . We have again by Lemma 17

$$\mathbb{E}_{\Theta_0} e^{tZ} = e^{-n\text{tr}(\Sigma_0 U)t} \mathbb{E}_{\Theta_0} e^{t \sum_{i=1}^n x_i^T U x_i} = e^{-n\text{tr}(\Sigma_0 U)t} \left( \frac{\det(\Theta_0)}{\det(\Theta_0 - 2tU)} \right)^{n/2}.$$

Since  $e^{tZ} \leq e^Z$  for  $t < 0$  and  $\mathbb{E}_{\Theta_0} e^Z < \infty$ , we can interchange differentiation and integration below to obtain

$$\mathbb{E}_{\Theta_0} Z e^{tZ} = \mathbb{E}_{\Theta_0} (e^{tZ})' = (\mathbb{E}_{\Theta_0} e^{tZ})'.$$

Hence

$$\begin{aligned}
\mathbb{E}_{\Theta_0} Z e^{tZ} &= (\mathbb{E}_{\Theta_0} e^{tZ})' \\
&= e^{-n\text{tr}(\Sigma_0 U)t} \left( \frac{\det(\Theta_0)}{\det(\Theta_0 - 2tU)} \right)^{n/2} \times \\
(41) \quad &\quad n [\text{tr}((\Theta_0 - 2tU)^{-1} U) - \text{tr}(\Sigma_0 U)]
\end{aligned}$$

Finally, taking  $t = -1/2$  in (41), we obtain

$$i = -2n\text{tr}[(\Theta_0 + U)^{-1} - \Sigma_0]U.$$

Hence

$$\begin{aligned} \mathbb{E}_{\Theta_0} \left( \frac{p_{\Theta_0+U}(x)}{p_{\Theta_0}(x)} - 1 - Z \right)^2 &= \frac{\det(\Theta_0 + U)^n}{\det(\Theta_0)^n} \left( \frac{\det(\Theta_0)}{\det(\Theta_0 + 2U)} \right)^{n/2} - 1 \\ &\quad + \sum_{i=1}^n \text{var}_{\Theta_0}(x_i^T U x_i) \\ &\quad - 2n \text{tr}[(\Theta_0 + U)^{-1} - \Sigma_0)U] \end{aligned}$$

This finishes the proof.  $\square$

LEMMA 19. *Let  $0 < \delta = \delta_n \rightarrow 0$  and let  $a, b = \mathcal{O}(1)$ . Then*

$$\left( 1 + \frac{a\delta}{\sqrt{n}(b\sqrt{\delta} + \sqrt{n})} \right)^n - 1 = \mathcal{O}(\delta).$$

PROOF OF LEMMA 19.

$$\begin{aligned} \left( 1 + \frac{a\delta}{\sqrt{n}(b\sqrt{\delta} + \sqrt{n})} \right)^n - 1 &= e^{n \log \left( 1 + \frac{a\delta}{\sqrt{n}(b\sqrt{\delta} + \sqrt{n})} \right)} - 1 \\ &= e^{n \left[ \frac{a\delta}{\sqrt{n}(b\sqrt{\delta} + \sqrt{n})} + o\left( \frac{a\delta}{\sqrt{n}(b\sqrt{\delta} + \sqrt{n})} \right) \right]} - 1 \end{aligned}$$

Next

$$\frac{an\delta}{\sqrt{n}(b\sqrt{\delta} + \sqrt{n})} = \mathcal{O}(\delta).$$

Hence, and using that  $e^x - 1 = o(x)$  for  $x \rightarrow 0$ , we obtain

$$e^{n \left[ \frac{a\delta}{\sqrt{n}(b\sqrt{\delta} + \sqrt{n})} + o\left( \frac{a\delta}{\sqrt{n}(b\sqrt{\delta} + \sqrt{n})} \right) \right]} - 1 = \mathcal{O}(\delta).$$

$\square$

LEMMA 20. *Let  $p_{\Theta_0}$  be the joint density of the random sample  $X_1, \dots, X_n$ , where each  $X_i$  is an  $\mathcal{N}(0, \Theta_0^{-1})$ -distributed random vector. Let  $H := \Theta_0(\xi_1 \xi_2^T + \xi_2 \xi_1^T) \Theta_0 / \sigma$ . Then*

$$\mathbb{E}_{\Theta_0} \left( \frac{p_{\Theta_0+H/\sqrt{m_n}}(x)}{p_{\Theta_0}(x)} - 1 - \text{tr}(s_{\Theta_0}(X)H)/\sqrt{m_n} \right)^2 = \mathcal{O}(\delta_n).$$

PROOF OF LEMMA 20. We apply Lemma 18 with

$$H := \Theta_0(\xi_1 \xi_2^T + \xi_2 \xi_1^T) \Theta_0 / \sigma,$$

where  $\sigma^2 := \xi_1^T \Theta^0 \xi_1 \xi_2^T \Theta^0 \xi_2 + (\xi_1^T \Theta^0 \xi_2)^2$ . To apply Lemma 18, we need to show that  $\Theta_0 + H/\sqrt{m_n}$  is a symmetric, positive definite matrix (for  $n$  sufficiently large). This can be seen as follows. First note that  $H := \Theta_0(\xi_1 \xi_2^T + \xi_2 \xi_1^T) \Theta_0 / \sigma$  is symmetric and  $\Theta_0$  is symmetric and hence the symmetry of the sum follows. Next we look at positive definiteness.

For any  $u \in \mathbb{R}^p$  we have by the Cauchy-Schwarz inequality

$$\begin{aligned} u^T H u &= u^T \Theta_0(\xi_1 \xi_2^T + \xi_2 \xi_1^T) \Theta_0 u / \sigma \\ &\leq 2 \sqrt{\xi_1^T \Theta_0 \xi_1 u^T \Theta_0 u} \sqrt{\xi_2^T \Theta_0 \xi_2 u^T \Theta_0 u} / \sigma \leq \Lambda_{\max}(\Theta) u^T u. \end{aligned}$$

Thus

$$\begin{aligned} u^T \Theta_0 u + u^T H u / \sqrt{m_n} &\geq u^T \Theta_0 u - \mathcal{O}(\Lambda_{\max}(\Theta_0) u^T u) / \sqrt{m_n} \\ &\geq [\Lambda_{\min}(\Theta_0) - \mathcal{O}(1) / \sqrt{m_n}] u^T u. \end{aligned}$$

This shows that the matrix  $\Theta_0 + H/\sqrt{m_n}$  is positive definite for  $n$  sufficiently large. But then we can apply Lemma 18 which gives

$$\begin{aligned} &\mathbb{E}_{\Theta_0} \left( \frac{p_{\Theta_0+U}(x)}{p_{\Theta_0}(x)} - 1 - \text{tr}(s_{\Theta_0}(X)H) / \sqrt{m_n} \right)^2 \\ &= \underbrace{\frac{\det(\Theta_0 + U)^n}{\det(\Theta_0)^n} \left( \frac{\det(\Theta_0)}{\det(\Theta_0 + 2U)} \right)^{n/2}}_i - 1 \\ &\quad + \underbrace{\sum_{i=1}^n \text{var}_{\Theta_0}(x_i^T U x_i)}_{ii} \\ &\quad - 2n \underbrace{\text{tr}[(\Theta_0 + U)^{-1} - \Sigma_0] U}_{iii}. \end{aligned}$$

Now we calculate the terms  $i, ii, iii$ .

1.) *Calculation of  $i$ .*

First we calculate  $i$ . Observe that since  $H$  is of rank 2 it follows

$$\begin{aligned} i &= \det(\Theta_0 + H/\sqrt{m_n}) = \det[\Theta_0 + \Theta_0(\xi_1 \xi_2^T + \xi_2 \xi_1^T) \Theta_0 / (2\sigma \sqrt{m_n})] \\ &= (1 + \xi_1^T \Theta_0 \xi_2 / (2\sigma \sqrt{m_n}))^2 \det(\Theta_0). \end{aligned}$$

And hence

$$\begin{aligned} & \frac{\det(\Theta_0 + H/\sqrt{m_n})^n}{\det(\Theta_0)^n} \left( \frac{\det(\Theta_0)}{\det(\Theta_0 + 2H/\sqrt{m_n})} \right)^{n/2} \\ &= \left[ \frac{(1 + \xi_1^T \Theta_0 \xi_2 / (2\sigma \sqrt{m_n}))^2}{(1 + \xi_1^T \Theta_0 \xi_2 / (\sigma \sqrt{m_n}))} \right]^n \end{aligned}$$

2.) *Calculation of ii.*

The following property holds: if  $Y \sim \mathcal{N}_2(0, S)$ , then it holds

$$\text{var}(Y_1 Y_2) = S_{11} S_{22} + S_{12}^2.$$

Further observe that since  $\Theta_0 x_i \sim \mathcal{N}(0, \Theta_0)$  we obtain

$$\begin{aligned} \text{var}_{\Theta_0}(x_i^T H x_i / \sqrt{m_n}) &= \text{var}_{\Theta_0}(\xi_1^T \Theta_0 x_i x_i^T \Theta_0 \xi_2^T) / (\sigma^2 m_n) \\ &= (\xi_1^T \Theta_0 \xi_1 \xi_2^T \Theta_0 \xi_2 + (\xi_1^T \Theta_0 \xi_2)^2) / (\sigma^2 m_n) = 1/m_n. \end{aligned}$$

Thus

$$ii = n/m_n.$$

3.) *Calculation of iii.*

Finally, by inversion of a sum of a matrix with another matrix of rank 2, we get (we omit the calculations)

$$iii = \text{tr}[(\Theta_0 + H/\sqrt{m_n})^{-1} - \Sigma_0] H / \sqrt{m_n} = \mathcal{O}\left(\frac{1}{m_n}\right).$$

By Lemma 18 and the above calculations of  $i, ii, iii$  it then follows

$$\begin{aligned} & \mathbb{E}_{\Theta_0} \left( \frac{p_{\Theta_0 + H/\sqrt{m_n}}(x)}{p_{\Theta_0}(x)} - 1 - \text{tr}(s_{\Theta_0}(X)H) / \sqrt{m_n} \right)^2 \\ &= \underbrace{\left( 1 + \frac{(\xi_1^T \Theta_0 \xi_2)^2 / (4\sigma^2 m_n)}{1 + 2\xi_1^T \Theta_0 \xi_2 / (\sigma m_n)} \right)^n - 1}_I \\ &+ \frac{n}{m_n} - \mathcal{O}\left(\frac{n}{m_n}\right). \end{aligned}$$

For the first term, by Lemma 19 we have that  $I = \mathcal{O}(\delta_n)$ . Hence we conclude that

$$\begin{aligned} & \mathbb{E}_{\Theta_0} \left( \frac{p_{\Theta_0 + H/\sqrt{m_n}}(x)}{p_{\Theta_0}(x)} - 1 - \text{tr}(s_{\Theta_0}(X)H) / \sqrt{m_n} \right)^2 \\ &= \mathcal{O}(\delta_n) + \mathcal{O}\left(\frac{n}{m_n}\right) = \mathcal{O}(\delta_n). \end{aligned}$$

□

## APPENDIX C: ADDITIONAL PROOFS FOR SECTION 17

In this section, we give Lemmas 21, 23.

LEMMA 21. *Let  $Z \in \mathbb{R}^2$  be  $\mathcal{N}(\mu, \Sigma)$ -distributed, where*

$$\mu = \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix}, \Sigma = \begin{pmatrix} \sigma_{11} & \sigma_{12} \\ \sigma_{12} & \sigma_{22} \end{pmatrix}.$$

*Suppose that  $\mu_2 = -\sigma_{22}/2$ . Let  $Y \in \mathbb{R}^2$  be  $\mathcal{N}(\mu + a, \Sigma)$ -distributed, with*

$$a = \begin{pmatrix} \sigma_{12} \\ \sigma_{22} \end{pmatrix}.$$

*Let  $\phi_Z$  be the density of  $Z$  and  $\phi_Y$  be the density of  $Y$ . Then we have the following equality for all  $z = (z_1, z_2) \in \mathbb{R}^2$ :*

$$\phi_Z(z)e^{z_2} = \phi_Y(z).$$

PROOF OF LEMMA 21. The density of  $Z$  is

$$\phi_Z(z) = \frac{1}{2\pi\sqrt{\det(\Sigma)}} e^{-\frac{1}{2}(z-\mu)^T \Sigma^{-1}(z-\mu)}.$$

It holds that

$$\Sigma^{-1}a = (0, 1)^T.$$

Then

$$\frac{1}{2}(z-\mu)^T \Sigma^{-1}(z-\mu) = \frac{1}{2}(z-\mu-a)^T \Sigma^{-1}(z-\mu-a) + a^T \Sigma^{-1}(z-\mu) - \frac{1}{2}a^T \Sigma^{-1}a.$$

We also have

$$a^T \Sigma^{-1}(z-\mu) - \frac{1}{2}a^T \Sigma^{-1}a = (0, 1)^T(z-\mu) - \frac{1}{2}(0, 1)^T a = z_2 - \mu_2 - \frac{1}{2}\sigma_{22} = z_2.$$

□

LEMMA 22. *Let  $\mu$  and  $\Sigma$  be defined as follows*

$$\mu = \begin{pmatrix} -\frac{v_{12}}{\sqrt{v_{11}}} \\ -\frac{v_{22}}{2} \end{pmatrix}, \quad \Sigma = \begin{pmatrix} 1 & \frac{v_{12}}{\sqrt{v_{11}}} \\ \frac{v_{12}}{\sqrt{v_{11}}} & v_{22} \end{pmatrix}.$$

*Suppose that  $V_\beta = \mathcal{O}(1)$ ,  $1/V_\beta = \mathcal{O}(1)$  and  $\Lambda_{\max}(I_\beta) = \mathcal{O}(1)$ . (The relationship between these quantities and the  $v_{ij}$ 's is given in the proof of Theorem 9). Then*

$$\|\mu\|_2^2 = \mathcal{O}(1) \quad \text{and} \quad \Lambda_{\max}(\Sigma) = \mathcal{O}(1).$$

PROOF. First observe that

$$\begin{aligned} v_{12}^2 &= (\mathbb{E}_\beta l_\beta h^T s_\beta)^2 \leq \mathbb{E}_\beta l_\beta^2 \mathbb{E}_\beta (h^T s_\beta)^2 \\ &= V_\beta h \mathbb{E}_\beta s_\beta s_\beta^T h \leq V_\beta \Lambda_{\max}(\mathbb{E}_\beta s_\beta s_\beta^T) h^T h. \end{aligned}$$

Then by assumption  $\Lambda_{\max}(\mathbb{E}_\beta s_\beta s_\beta^T) = \mathcal{O}(1)$ ,  $V_\beta = \mathcal{O}(1)$  and since  $h^T h = \mathcal{O}(1)$ , we have that  $(\mathbb{E}_\beta l_\beta h^T s_\beta)^2 = \mathcal{O}(1)$ . Also observe that  $v_{22} = h^T I_\beta h \leq \Lambda_{\max}(I_\beta) h^T h = \mathcal{O}(1)$  by assumption  $\Lambda_{\max}(I_\beta) = \mathcal{O}(1)$ .

Then, and by  $1/V_\beta = \mathcal{O}(1)$ , it follows that

$$\|\mu\|_2^2 = v_{12}^2/v_{11} + v_{22}^2/4 = (P_\beta l_\beta h^T s_\beta)^2/V_\beta + (h^T I_\beta h)^2/4 = \mathcal{O}(1).$$

We proceed to check that the eigenvalues of  $\Sigma$  are bounded. We have

$$\lambda_{1,2} = \frac{1 + v_{22} \pm \sqrt{D}}{2},$$

where  $D = (1+v_{22})^2 - 4(v_{22} - v_{12}^2/v_{11}) = (1-v_{22})^2 + 4v_{12}^2/v_{11}$ . Clearly,  $D \geq 0$ , and as above, one sees that  $D = \mathcal{O}(1)$ . Hence also  $\Lambda_{\max}(\Sigma) = \mathcal{O}(1)$ .  $\square$

LEMMA 23. *Suppose that*

$$U_n \sim \mathcal{N}\left(\begin{pmatrix} -\frac{v_{12}}{\sqrt{v_{11}}} \\ -\frac{v_{22}}{2} \end{pmatrix}, \begin{pmatrix} 1 & \frac{v_{12}}{\sqrt{v_{11}}} \\ \frac{v_{12}}{\sqrt{v_{11}}} & v_{22} \end{pmatrix}\right).$$

*Suppose that  $V_\beta = \mathcal{O}(1)$ ,  $1/V_\beta = \mathcal{O}(1)$ ,  $\Lambda_{\max}(I_\beta) = \mathcal{O}(1)$  and  $h \in \Theta$ . (The relationship between these quantities and the  $v_{ij}$ 's is given in the proof of Theorem 9).*

*Then it holds that*

$$\lim_{M \rightarrow \infty} \lim_{n \rightarrow \infty} \mathbb{E} \min(0, M - e^{U_{n,2}}) = 0.$$

PROOF OF LEMMA 23. We have

$$\begin{aligned} \mathbb{E} \min(0, M - e^{U_{n,2}}) &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \min(0, M - e^{x_2}) \phi_{U_n}(x) dx \\ &= \int_{-\infty}^{\infty} \int_{\log M}^{\infty} M - e^{x_2} \phi_{U_n}(x) dx \\ &= MP(U_{n,2} > \log M) - P(Y_{n,2} > \log K), \end{aligned}$$

where we applied Lemma 21 and denoted

$$Y \sim \mathcal{N}\left(\begin{pmatrix} 0 \\ v_{22}/2 \end{pmatrix}, \begin{pmatrix} 1 & \frac{v_{12}}{\sqrt{v_{11}}} \\ \frac{v_{12}}{\sqrt{v_{11}}} & v_{22} \end{pmatrix}\right).$$

Thus we have

$$P(Y_{n,2} > \log M) \leq e^{-(\log M)^2/2},$$

and

$$\begin{aligned} P(U_{n,2} > \log M) &= P(U_{n,2} + v_{12}/\sqrt{v_{11}} > \log M + v_{12}/\sqrt{v_{11}}) \\ &\leq e^{-(\log M + v_{12}/\sqrt{v_{11}})^2/2}. \end{aligned}$$

Now observe that

$$\begin{aligned} v_{12}^2 &= (\mathbb{E}_\beta l_\beta h^T s_\beta)^2 \leq \mathbb{E}_\beta l_\beta^2 \mathbb{E}_\beta (h^T s_\beta)^2 \\ &= V_\beta h \mathbb{E}_\beta s_\beta s_\beta^T h \leq V_\beta \Lambda_{\max}(\mathbb{E}_\beta s_\beta s_\beta^T) h^T h. \end{aligned}$$

Then by assumption  $\Lambda_{\max}(\mathbb{E}_\beta s_\beta s_\beta^T) = \mathcal{O}(1)$ ,  $V_\beta = \mathcal{O}(1)$  and since  $h^T h = \mathcal{O}(1)$ , we have that  $(\mathbb{E}_\beta l_\beta h^T s_\beta)^2 = \mathcal{O}(1)$ . Hence  $v_{12} = \mathcal{O}(1)$ . We also have by the assumption  $1/V_\beta = \mathcal{O}(1)$  that  $1/\sqrt{v_{11}} = \mathcal{O}(1)$ . Hence we can conclude that  $v_{12}/\sqrt{v_{11}} \leq L$  for some constant  $L > 0$ . Now without loss of generality, choose  $M$  such that  $\log M > 2L$ . Then we have  $\log M + v_{12}/\sqrt{v_{11}} \geq \log M/2$ . Thus we obtain

$$\begin{aligned} \mathbb{E} \min(0, M - e^{U_{n,2}}) &= MP(U_{n,2} > \log M) - P(Y_{n,2} > \log K) \\ &\leq Me^{-(\log M + v_{12}/\sqrt{v_{11}})^2/2} + e^{-(\log M)^2/2} \\ &\leq Me^{-(\log M)^2/8} + e^{-(\log M)^2/2}. \end{aligned}$$

Finally, taking the limits we obtain

$$\lim_{M \rightarrow \infty} \lim_{n \rightarrow \infty} \mathbb{E} \min(0, M - e^{U_{n,2}}) \leq \lim_{M \rightarrow \infty} \lim_{n \rightarrow \infty} Me^{-(\log M)^2/2} + e^{-(\log M)^2/8} = 0.$$

□