

# On Spectrum Generating Algebra of the Heun Operator

Priyasri Kar<sup>a</sup>, Ritesh K. Singh<sup>b</sup>, Ananda Dasgupta<sup>c</sup>, Prasanta K. Panigrahi<sup>d</sup>

*Department of Physical Sciences,  
Indian Institute of Science Education and Research Kolkata, Mohanpur 741246, India*

<sup>a</sup> pk12rs063@iiserkol.ac.in, <sup>b</sup> ritesh.singh@iiserkol.ac.in,

<sup>c</sup> adg@iiserkol.ac.in, <sup>d</sup> pprasanta@iiserkol.ac.in

February 28, 2022

## Abstract

The Heun operator has been cast, in terms of the elements of an underlying  $su(1, 1)$  algebra, under specific parametric conditions, for the purpose of spectrum generation. These elements are differential operators of *degrees*  $\pm 1/2$  and 0. It is found that the regular singularities at 0 and  $\infty$  of the general Heun equation must be *elementary* under the required parametric conditions. The spectrum generation has been demonstrated through a set of examples.

Differential equations, in general, *do not* admit exact solutions. Several techniques have been devised, to obtain partial, or in certain cases, the complete eigenspace of the differential operator under study [1]. In this regard, the well known factorization method, which at a fundamental level reduces the order of the differential operator involved, has found extensive applications [2]. The harmonic oscillator spectrum generating Heisenberg algebra, the Darboux transformation, factorization of the hypergeometric equation by Schrödinger [3] and subsequent study of a large class of equations by Infeld and Hull [2] serve as classic examples. Supersymmetric quantum mechanics (SUSYQM) uses factorization, to reveal unexpected interconnections amongst apparently different quantal problems [4, 5, 6].

Group theoretical structures also have played significant role in unravelling the symmetry of differential operators [1]. A partial algebraization of the whole solution space has been achieved [7, 8, 9] for a wide class of quantal quasi-exactly solvable (QES) Hamiltonians of one or more dimensions, which are essentially hermitian differential operators of the second order. These Hamiltonians, or their appropriately similarity transformed versions, often take the form of some well known differential operators, e.g., Heun, Lamé etc. Hence, the study of the underlying symmetries of these operators is of immense importance and several attempts have been made to this end. For example, the Lamé equation has been expressed as a bilinear in both  $su(2)$  [10] and  $su(1, 1)$  [11] generators. The latter case relates the non-unitary representations of  $su(1, 1)$ , to the eigenfunction of the periodic Lamé potential. Closed form solutions for confluent hypergeometric and hypergeometric equations have been found [12], using a connection between the space of monomials and the solution space of the above equations [13, 14], which reveals the underlying  $su(1, 1)$  and deformed  $sl(2)$  structures, respectively [15].

Recently, the Heun operator has been cast [16] in terms of the elements of cubic deformations of  $sl(2)$  algebra. This has led to two known solutions of a Heun equation, encountered earlier by Christ and Lee [17]. However, non-linear deformations of  $sl(2)$ , are associated with non-trivial

representation theory [18]. Casting the differential operator in terms of linear  $sl(2)$  algebra enables one to exploit available representation theory. Linear  $su(1, 1)$  symmetry of the Heun operator is present in the literature [7]. In the present work, we use a *different* representation of  $su(1, 1)$  generators. Under certain parametric conditions, this reveals a new symmetry of the equation and leads to a number of solutions, that are unavailable from the methods in [7]. It is found that this new symmetry exists if the singularities at  $z = 0$  and  $z = \infty$  are *elementary*<sup>1</sup>. Under these conditions, an exact map is established between the Heun solution space and the unitary and non-unitary representations of  $su(1, 1)$  algebra.

The Heun equation is given by [20]

$$\mathcal{H}y(z) = \frac{d^2 y(z)}{dz^2} + \left( \frac{\gamma}{z} + \frac{\delta}{z-1} + \frac{\varepsilon}{z-a} \right) \frac{dy(z)}{dz} + \frac{\alpha\beta z - q}{z(z-1)(z-a)} y(z) = 0, \quad (1)$$

with regular singularities at  $z = 0, 1, a(\neq 0, 1)$  and  $\infty$ ; the exponents at these singularities being  $(0, 1-\gamma), (0, 1-\delta), (0, 1-\varepsilon)$  and  $(\alpha, \beta)$ , respectively. Here,  $q$  plays the role of the *eigen parameter*. In this work, we study Heun equations with only real parameters. The equation, being a second order linear homogeneous equation with 4 regular singularities, satisfies the Fuchsian condition:

$$\gamma + \delta + \varepsilon = \alpha + \beta + 1, \quad (2)$$

allowing elimination of  $\varepsilon$  in favor of the others. Eq. (1) can be written as:

$$\mathcal{H}y(z) = \left[ f_1(z) \frac{d^2}{dz^2} + f_2(z) \frac{d}{dz} + f_3(z) \right] y(z) = 0, \quad (3)$$

where,  $f_1(z) = a_0 z^3 + a_1 z^2 + a_2 z$ ,  $f_2(z) = a_3 z^2 + a_4 z + a_5$  and  $f_3(z) = a_6 z + a_7$ . The parameters  $a_i \in \mathbb{R}$ , for  $i = 0, \dots, 7$  are given by,

$$a_0 = 1 \quad a_3 = 1 + \alpha + \beta \quad a_6 = \alpha\beta \quad (4a)$$

$$a_1 = -(a+1) \quad a_4 = -[a\gamma + a\delta - \delta + \alpha + \beta + 1] \quad a_7 = -q \quad (4b)$$

$$a_2 = a \quad a_5 = a\gamma. \quad (4c)$$

Evidently, the Heun equation consists of differential operators of *degrees*<sup>2</sup>  $+1, 0$  and  $-1$ , which are denoted by  $P_+, F(P_0)$  and  $P_-$ , respectively. Eq. (3) can then be rewritten as [16],

$$\mathcal{H}y(z) = [P_+ + F(P_0) + P_-]y(z) = 0 \quad (5)$$

where,

$$P_+ = a_0 z^3 \frac{d^2}{dz^2} + a_3 z^2 \frac{d}{dz} + a_6 z, \quad P_0 = z \frac{d}{dz} - j, \quad P_- = a_2 z \frac{d^2}{dz^2} + a_5 \frac{d}{dz} \quad (6)$$

$$\text{and } F(P_0) = a_1 P_0^2 + ((2j-1)a_1 + a_4) P_0 + (j(j-1)a_1 + ja_4 + a_7). \quad (7)$$

The above operators satisfy cubic deformation of  $sl(2)$  algebra among themselves, which has been exploited [16] to tract a part of the eigenspace of a Heun operator analytically, as mentioned earlier.

With the aim to cast the Heun operator in terms of linear  $su(1, 1)$  algebra, we define the *constituent* operators as,

$$E_+ = 2z^{3/2} \frac{d}{dz} + 2\mu\sqrt{z}, \quad H = 2z \frac{d}{dz} + \mu + \nu \quad \text{and} \quad E_- = 2z^{1/2} \frac{d}{dz} + \frac{2\nu}{\sqrt{z}}; \quad (8)$$

<sup>1</sup>Any regular singularity can be characterized by two exponents  $\rho_1$  and  $\rho_2$ , which are the two roots of the indicial equation. The cases with  $|\rho_1 - \rho_2| = 1/2$  are known as elementary singularity and are of special significance. All regular and irregular singularities can be obtained by the coalescence of two and three or more elementary singularities, respectively [19].

<sup>2</sup>The degree  $d$ , of an operator  $O_d$ , is defined as the change in the power of a monomial, when acted upon by it, i.e.,  $O_d z^p \propto z^{p+d}$ .

where  $\mu, \nu$  and  $\lambda$  are parameters. The operators  $E_+, H$  and  $E_-$ , have degrees  $+1/2, 0$  and  $-1/2$ , respectively. They are found to satisfy  $su(1, 1)$  algebra, with commutators,

$$[H, E_{\pm}] = \pm E_{\pm}, \quad [E_+, E_-] = -2H \quad (9)$$

and the Casimir:

$$C(\mu, \nu) = \frac{1}{2} (E_+ E_- + E_- E_+) - H^2 = -(\mu - \nu)(\mu - \nu - 1). \quad (10)$$

The problem in question, determines the parameters of the constituent operators, which determine the Casimir value. The Casimir value, in turn, determines the representations available for the algebra associated with the problem.

We now proceed to construct the differential operators  $P_+, P_-$  and  $F(P_0)$ , from the constituent  $su(1, 1)$  generators and identify the conditions, under which this is possible. The constituent operators are of degrees  $\pm 1/2$  and  $0$ , whereas, the degrees of operators  $P_{\pm}$  and  $F(P_0)$  are  $\pm 1$  and  $0$ , respectively. Hence, it is clear that the complete Heun operator will comprise of linear and quadratic forms of the constituent operators<sup>3</sup>. Assuming the forms,

$$P_+ = c_+ E_+ E_+ \quad \text{and} \quad P_- = c_- E_- E_- \quad (11)$$

where  $c_{\pm}$  are constants, we obtain,

$$a_0 = 4c_+, \quad a_3 = 2c_+(3 + 4\mu) = \frac{a_0}{2}(3 + 4\mu), \quad a_6 = 2c_+\mu(1 + 2\mu) = \frac{a_0}{2}\mu(1 + 2\mu), \quad (12)$$

$$a_2 = 4c_-, \quad a_5 = 2c_-(1 + 4\nu) = \frac{a_2}{2}(1 + 4\nu), \quad 2c_-\nu(2\nu - 1) = 0. \quad (13)$$

Using Eqs. (12) and (4a), one can solve for  $\mu, \alpha, \beta$ :

$$|\alpha - \beta| = \frac{1}{2}, \quad (14)$$

implying the singularity at  $z = \infty$  is elementary. Similarly, using Eqs. (13) and (4c), one is left with just two values of  $\nu$ :

$$\nu = 0 \implies a_5/a_2 = \gamma = 1/2 \quad \text{or,} \quad \nu = 1/2 \implies a_5/a_2 = \gamma = 3/2. \quad (15)$$

Above solutions of  $\gamma$  imply that the singularity at  $z = 0$  is also elementary. Finally,  $F(P_0)$  can be written in the form

$$F(P_0) = c_2 H H + c_1 H + c_0, \quad (16)$$

where,

$$a_1 = 4c_2, \quad a_4 = 2(c_1 + 2c_2(1 + \mu + \nu)) \quad \text{and} \quad a_7 = -q = c_0 + (\mu + \nu)(c_1 + c_2(\mu + \nu)). \quad (17)$$

Thus, a Heun operator with elementary singularities at  $z = 0$  and  $z = \infty$ , i.e., with parametric conditions Eqs. (14) and (15), becomes,

$$\mathcal{H} = c_+ E_+ E_+ + c_- E_- E_- + c_2 H H + c_1 H + c_0. \quad (18)$$

The choices of  $E_+$  and  $E_-$  fix the values of  $\mu$  and  $\nu$ , respectively, which in turn determines the Casimir,  $C = C(\mu, \nu)$ . Finding the solution to Heun equation, then boils down to identifying a suitable representation space  $V_C$  of  $su(1, 1)$  and finding a  $y \in V_C$ , such that,  $\mathcal{H}y = 0$ . For  $su(1, 1)$  algebra, there are five different classes of representation spaces [21] determined by the values of  $C$  and  $h$ , the eigen value of operator  $H$ . These are shown in Fig. 1 and are listed below.

---

<sup>3</sup>Cubic and quartic combinations, such as  $E_+ E_+ H, E_+ E_+ H H, E_+ E_+ E_+ E_-$  (and similar terms with  $E_+$  and  $E_-$  interchanged), are also of the desired degrees and therefore could be used in the construction of the Heun operators. However, they contain differential operators of higher orders, not present in the Heun equation. Thus, the coefficients of those undesired terms must be put to zero, which revealed no more parametric freedom than the chosen combinations.




Figure 1: Casimir value for  $su(1,1)$  and corresponding representations are listed: positive discrete (PD), negative discrete (ND), principal series (PS), complementary series (CS) and some finite dimensional ones (open circles).

**Principal series (PS):** Infinite dimensional space,  $C \geq 1/4$ ,  $h \in (-\infty, \infty)$  and  $(C, h) \neq (1/4, 1/2)$ ,

**Complementary series (CS):** Infinite dimensional space,  $C \in (0, 1/4)$ ,  $h \in (-\infty, \infty)$ ,  $h \neq 1/2$ ,

**Positive discrete (PD):** Infinite dimensional space,  $C \leq 1/4$ ,  $h \geq 0$ ,

**Negative discrete (ND):** Infinite dimensional space,  $C \leq 1/4$ ,  $h \leq 0$ ,

**Finite dimensions:** Finite dimensional space,  $C = \frac{-(n^2-1)}{4}$ ,  $h \in \left[\frac{-(n-1)}{2}, \frac{(n-1)}{2}\right]$  with  $n \in \mathbb{N}$ .

For a given value of Casimir  $C$ , one of the available representation space  $V_C$  can be chosen. The elements of this space are  $|C, h\rangle \propto z^p$ , with  $p = (h - \mu - \nu)/2$ . We represent the solution  $y$  as a linear combination of  $z^p$  from  $V_C$  and find the co-efficients. Due to the quadratic dependence of  $\mathcal{H}$  on  $E_+$  and  $E_-$ , we notice that, the solution space splits as  $V_C = V_e \oplus V_o$  — the even and odd states of the representation space. The process of finding solutions in  $V_e$  and  $V_o$  is demonstrated below, through a set of examples.

**Example 1:** We choose  $\gamma = 1/2$ ,  $\delta = \varepsilon = -1/2$ ,  $\alpha\beta = 1/2$ . This leads to  $|\alpha - \beta| = 1/2$ , ensuring the singularities at 0 and  $\infty$  to be elementary. The Heun equation takes the form

$$\frac{d^2y}{dz^2} + \frac{1}{2} \left( \frac{1}{z} - \frac{1}{z-1} - \frac{1}{z-a} \right) \frac{dy}{dz} + \frac{z-2q}{2z(z-1)(z-a)} y = 0, \quad (19)$$

with  $\mu = -1$ ,  $\nu = 0$  i.e.,  $C = -2$ . This allows the positive discrete and negative discrete representation spaces along with a triplet representation with  $h \in \{-1, 0, +1\}$  or  $p \in \{0, 1/2, 1\}$ . Choosing the triplet space,  $V_e$  corresponds to  $p \in \{0, 1\}$  and  $V_o$  to  $p \in \{1/2\}$ , leading to the solutions

$$y_1 \in V_e = z + \sqrt{a}, \quad \text{with eigenvalue } q = +\sqrt{a}/2, \quad (20)$$

$$y_2 \in V_e = z - \sqrt{a}, \quad \text{with eigenvalue } q = -\sqrt{a}/2 \quad (21)$$

$$\text{and } y_3 \in V_o = \sqrt{z}, \quad \text{with eigenvalue } q = (a+1)/4. \quad (22)$$

The positive discrete and negative discrete representations also yield solutions, not explicitly shown for this example.

**Example 2:** We choose  $\gamma = 3/2$ ,  $\delta = \varepsilon = -1/2$ ,  $\alpha\beta = 0$ , leading to  $|\alpha - \beta| = 1/2$ . The Heun equation is given by,

$$\frac{d^2y}{dz^2} + \frac{1}{2} \left( \frac{3}{z} - \frac{1}{z-1} - \frac{1}{z-a} \right) \frac{dy}{dz} + \frac{q}{z(z-1)(z-a)} y = 0 \quad (23)$$

and corresponds to  $\mu = -1/2$ ,  $\nu = 1/2$  with  $C = -2$ , as in the previous case. Here, for the triplet space, one obtains  $p \in \{-1/2, 0, 1/2\}$ , with  $p \in \{0\}$  being  $V_o$ . The solutions are:

$$y_1 \in V_e = \sqrt{z} + \sqrt{a/z}, \quad \text{with eigenvalue } q = -(a+1)/4 + \sqrt{a}/2, \quad (24)$$

$$y_2 \in V_e = \sqrt{z} - \sqrt{a/z}, \quad \text{with eigenvalue } q = -(a+1)/4 - \sqrt{a}/2 \quad (25)$$

$$\text{and } y_3 \in V_o = \text{const.}, \quad \text{with eigenvalue } q = 0. \quad (26)$$

Here too, we have positive discrete and negative discrete representations, like the previous example.

**Example 3:** Next, we analyze the Lamé equation, which is a special case of the Heun equation:

$$\frac{d^2 y}{dz^2} + \frac{1}{2} \left( \frac{1}{z} + \frac{1}{z-1} + \frac{1}{z-a} \right) \frac{dy}{dz} - \frac{q + \rho(\rho+1)z/4}{z(z-1)(z-a)} y(z) = 0. \quad (27)$$

Here, the singularities at  $z = 0, 1$  and  $a$  are elementary and choosing  $\rho(\rho+1) = 0$ , makes the singularity at  $z = \infty$  also elementary. With this choice, we have  $\mu = 0 = \nu$ , i.e.,  $C = 0$ , giving us the possibility to choose from singlet, positive discrete and negative discrete representations. The singlet solution is,  $y_0 = \text{const.}$ , with  $q = 0$ . The positive discrete space splits into the even ( $V_e^+$ ) and the odd ( $V_o^+$ ) subspaces and the corresponding solutions, for a chosen eigenvalue  $q \in \mathbb{R}$ , are:

$$y_1 \in V_e^+ = 1 + 2q \left[ \frac{z}{a} + \frac{q+a+1}{3} \left( \frac{z}{a} \right)^2 + \frac{2q^2 + 10q(a+1) + 8(a^2+1) + 7a}{45} \left( \frac{z}{a} \right)^3 + \dots \right] \quad (28)$$

$$y_2 \in V_o^+ = \sqrt{z} \left[ 1 + \frac{4q+a+1}{6} \left( \frac{z}{a} \right) + \frac{16q^2 + 40q(a+1) + 9(a^2+1) + 6a}{120} \left( \frac{z}{a} \right)^2 + \dots \right]. \quad (29)$$

Similarly, for the negative discrete space too, we have the even ( $V_e^-$ ) and the odd ( $V_o^-$ ) subspaces and the corresponding solutions, for a chosen eigenvalue  $q \in \mathbb{R}$ , are:

$$y_3 \in V_e^- = 1 + 2q \left[ \frac{1}{z} + \frac{q+a+1}{3} \left( \frac{1}{z^2} \right) + \frac{2q^2 + 10q(a+1) + 8(a^2+1) + 7a}{45} \left( \frac{1}{z^3} \right) + \dots \right] \quad (30)$$

$$y_4 \in V_o^- = \frac{1}{\sqrt{z}} \left[ 1 + \frac{4q+a+1}{6} \left( \frac{1}{z} \right) + \frac{16q^2 + 40q(a+1) + 9(a^2+1) + 6a}{120} \left( \frac{1}{z^2} \right) + \dots \right]. \quad (31)$$

The domain of convergence for the positive discrete space solutions is  $0 < z < \min\{1, a\}$ , whereas, that for the negative discrete space solutions is  $\max\{1, a\} < z < \infty$ . The solutions in infinite dimensional spaces can always be written as a power series in  $z$  or  $1/z$ , with an overall factor of  $z^p$ ,  $p \in \mathbb{R}$ , like the above four solutions.

For all the above examples, we have obtained solutions, which are *not* available from the previously known  $su(1, 1)$  symmetry of the Heun equation [7]. These solutions are presented here by Eqs. (22), (24), (25), (29) and (31). It may be noted that all of them involve powers of  $\sqrt{z}$ , which reflects the advantage of using a different representation of the  $su(1, 1)$  generators in the present work.

In conclusion, we have found that, the Heun operator can be exactly cast as a quadratic polynomial of elements of  $su(1, 1)$  algebra, if the singularities at  $z = 0$  and  $z = \infty$  are elementary. This allows one to use the representations of  $su(1, 1)$  to find explicit solutions of Heun equation. The finite dimensional representations yield polynomial solutions in  $\sqrt{z}$ , while positive and negative discrete representations give power series solutions in  $z$  and  $1/z$ , respectively, modulo an overall factor of  $z^p$ . Heun equation is connected with a host of physical problems, related to quantum mechanical and non-linear systems. Hence, the relevance of the algebraically generated solution space to these problems needs careful considerations. We intend to return to some of these problems in near future.

**Acknowledgements:** We thank Arunesh Roy for useful discussions.

## References

- [1] Willard Miller Jr., *Symmetry Groups and their Applications*, Academic Press (1972).
- [2] L. Infeld and T. E. Hull, *Rev. Mod. Phys.* **23**, 1 (1951).
- [3] E. Schrödinger, *Proc. Roy. Irish Acad.* **A47**, 53 (1941).
- [4] F. Cooper, A. Khare and U. Sukhatme, *Phys. Rep.* **251**, 267 (1995).
- [5] F. Cooper, A. Khare and U. Sukhatme, *Supersymmetry in Quantum Mechanics*, World Scientific (2001).
- [6] A. Gangopadhyaya, J. V. Mallow, C. Rasinariu, *Supersymmetric Quantum Mechanics: An Introduction*, World Scientific (2010).
- [7] A. V. Turbiner, *Commun. Math. Phys.* **118**, 467 (1988).
- [8] M. A. Shifman and A. V. Turbiner, *Commun. Math. Phys.* **126**, 347 (1989).
- [9] A. G. Ushveridze, *Quasi-exactly Solvable Models in Quantum Mechanics*, Inst. of Physics Publ. (1994).
- [10] Y. Alhassid, F. Gursey and F. Iachello, *Phys. Rev. Lett.* **50**, 873 (1983).
- [11] H. Li and D. Kusnezov, *Phys. Rev. Lett.* **83**, 1283 (1999).
- [12] N. Gurappa, P. K. Jha and P. K. Panigrahi, *SIGMA* **3**, 057 (2007).
- [13] N. Gurappa and P. K. Panigrahi, *Phys. Rev.* **B62**, 1943 (2000).
- [14] N. Gurappa and P. K. Panigrahi, *J. Phys.* **A37**, L605 (2004).
- [15] T. Shreecharan, P. K. Panigrahi and J. Banerji, *Phys. Rev.* **A69**, 012102 (2004).
- [16] A. Roy, A. Sen and P. K. Panigrahi, Preprint [arXiv:1304.2225].
- [17] N. H. Christ and T. D. Lee, *Phys. Rev.* **D12**, 1606 (1975).
- [18] M. Rocek, *Phys. Lett.* **B255**, 554 (1991).
- [19] E. L. Ince, *Ordinary Differential Equations*, Dover Books on Mathematics (1926).
- [20] A. Ronveaux (ed.), *Heun's Differential Equations*, Oxford University Press (1995).
- [21] See for example, W. Groenevelt, *Indag. Math. (N.S.)* **14**, 329 (2003) and references therein.