

Test Flavor SU(3) symmetry in Exclusive Λ_c decays

Cai-Dian Lü¹*, Wei Wang^{2,3}† and Fu-Sheng Yu⁴‡

¹ *Institute of High Energy Physics, P.O. Box 918(4),
Chinese Academy of Sciences, Beijing 100049, People's Republic of China*

² *INPAC, Shanghai Key Laboratory for Particle Physics and Cosmology,
Department of Physics and Astronomy, Shanghai Jiao-Tong University, Shanghai, 200240, China*

³ *State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics,
Chinese Academy of Sciences, Beijing 100190, China*

⁴ *School of Nuclear Science and Technology, Lanzhou University,
Lanzhou 730000, People's Republic of China*

Flavor SU(3) symmetry is a powerful tool to analyze charmed baryon decays, however its applicability remains to be experimentally validated. Since there is not much data on Ξ_c decays, various exclusive Λ_c decays especially the ones into a neutron state are essential for the test of flavor symmetry. These decay modes are also helpful to investigate final state interactions in charmed baryon decays. In this work, we discuss the explicit roles of Λ_c decays into a neutron in testing the flavor symmetry and exploring final state interactions. The involved decay modes include semileptonic decays, two-body and three-body non-leptonic decays, but all of them have not been experimentally observed to date.

I. INTRODUCTION

Charmed baryon decays, in particular Λ_c and Ξ_c decays, are of great interest as they serve as a platform for the study of strong and weak interactions in heavy-to-light baryonic transitions. They can also provide the essential inputs for the Λ_b decay modes into a charmed baryon like Λ_c . On the experimental side, most available results on Λ_c decays are obtained using the old data until recently. In 2014, Belle collaboration provided an measurement of the branching fraction with a very small uncertainty [1],

$$\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+)_{\text{Belle}} = (6.84 \pm 0.24_{-0.27}^{+0.21})\%, \quad (1)$$

but the central value is much larger than the previous measurement by the CLEO-c collaboration [2]:

$$\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+)_{\text{CLEO}} = (5.0 \pm 0.5 \pm 1.2)\%. \quad (2)$$

Based on the large amount of data, Belle collaboration also started to study the doubly Cabibbo-suppressed processes [3]. Making use of the data collected in the e^+e^- collision at the center-of-mass energy of $\sqrt{s} = 4.599$ GeV and adopting the double-tag technique, BES-III collaboration has reported first measurements of absolute hadronic branching fractions of Cabibbo-favored decay

* Email: lucd@ihep.ac.cn

† Email: wei.wang@sjtu.edu.cn

‡ Email: yufsh@lzu.edu.cn

modes [4]. In total, twelve Λ_c decay modes were observed with the significant improvement on the branching fraction in particular for the $\Lambda_c \rightarrow pK^-\pi^+$:

$$\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+)_{\text{BESIII}} = (5.84 \pm 0.27 \pm 0.23)\%. \quad (3)$$

While the uncertainties are comparable with the Belle results in Eq. (1), its central value is much smaller, which is closer to the central value of the CLEO results in Eq. (2). We believe this difference will be clarified in future since the experimental prospect on charmed baryon decays will be very promising [5, 6].

Theoretical description of charmed baryon decays is mostly based on the factorization assumption together with the analysis of some non-factorizable contributions in nonperturbative explicit modes [7–10]. However the factorization scheme does not seem to be supported by experiments, for instance the observed large branching fraction for decays like $\Lambda_c \rightarrow \Sigma^+\pi^0/\Xi^0 K^+$, which are forbidden in the factorization scheme [11]. An alternative and the model-independent approach is to make use of the flavor SU(3) symmetry, which has been argued to work better in charmed baryon decays [12–17] and bottomed baryon decays [18–20].

As the experimental precision is gradually increasing, the time is ripe to validate/invalidate the applicability of the SU(3) symmetry to charmed baryon decays. The SU(3) transformation connects the Λ_c with the Ξ_c . But at this stage and in the foreseeable future there is no experiment which will focus on the study on Ξ_c decays. Thus the Λ_c decays into various final states especially the ones into a neutron are of great value since they will be the only source for the test of the SU(3) symmetry in charmed-baryon decays. The motivation of this work is to discuss the roles of the Λ_c decays into a neutron into the test of SU(3) symmetry and the exploration of final state interactions, including semileptonic decays, two-body and three-body nonleptonic decays. All these exclusive decay modes have not been experimentally measured yet.

This paper is organized as follows. In Sec. II, the semileptonic Λ_c decays are studied. In Sec. III and Sec. IV, we will explore the two-body and three-body nonleptonic decays of the Λ_c , respectively. The last section contains our summary.

II. SEMILEPTONIC Λ_c DECAYS

We start with the semileptonic Λ_c decays. In the flavor SU(3) symmetry limit, the charmed baryons are classified according to the SU(3) irreducible representation, namely as multiplets of the light-quark system: $3 \otimes 3 = \bar{3} \oplus 6$. The Λ_c and Ξ_c forms the charmed-baryon anti-triplet in the initial state:

$$T^a = (\Xi_{c1}^0, -\Xi_{c1}^+, \Lambda_c^+). \quad (4)$$

For the light baryons, we focus on the SU(3) octet which is represented by the matrix:

$$B_b^a = \begin{pmatrix} \frac{1}{\sqrt{6}}\Lambda^0 + \frac{1}{\sqrt{2}}\Sigma^0 & \Sigma^+ & p \\ \Sigma^- & \frac{1}{\sqrt{6}}\Lambda^0 - \frac{1}{\sqrt{2}}\Sigma^0 & n \\ \Xi^- & \Xi^0 & -\sqrt{2/3}\Lambda^0 \end{pmatrix}. \quad (5)$$

The operator responsible for the transition $c \rightarrow qe^+\bar{\nu}_e$ is $[\bar{q}\gamma^\mu(1-\gamma_5)c][\bar{\nu}_e\gamma_\mu(1-\gamma_5)e]$ with $q = d, s$, which forms an SU(3) anti-triplet in the final state. Thus the effective Hamiltonian at hadron level is constructed as

$$H_{\text{eff}} = aH_a(\bar{3})T^b\bar{B}_b^a\bar{\nu}_ee. \quad (6)$$

An implication of the above Hamiltonian is obtained straightforwardly:

$$\mathcal{B}(\Lambda_c \rightarrow ne^+\nu_e) = \frac{3}{2} \frac{|V_{cd}|^2}{|V_{cs}|^2} \mathcal{B}(\Lambda_c \rightarrow \Lambda e^+\nu_e). \quad (7)$$

Measurements of the relevant branching fractions provide a most straightforward test of the flavor SU(3) symmetry in charmed baryon decays. With the most recent data from the BES-III collaboration [21]

$$\mathcal{B}(\Lambda_c \rightarrow \Lambda e^+\nu_e)_{\text{BESIII}} = (3.65 \pm 0.38 \pm 0.20)\%, \quad (8)$$

we can obtain the following result:

$$\mathcal{B}(\Lambda_c \rightarrow ne^+\nu_e)_{\text{SU(3)}} = (2.93 \pm 0.34) \times 10^{-3}, \quad (9)$$

which might be accessible for BES-III and Belle-II collaborations [5, 6].

In semileptonic decays, the neutron can be produced together with a light pseudo-scalar meson. The lowest-lying pseudo-scalar meson can be written as

$$M_b^a = \begin{pmatrix} \frac{1}{\sqrt{2}}\pi^0 + \frac{1}{\sqrt{6}}\eta & \pi^+ & K^+ \\ \pi^- & -\frac{1}{\sqrt{2}}\pi^0 + \frac{1}{\sqrt{6}}\eta & K^0 \\ K^- & \bar{K}^0 & -\sqrt{2/3}\eta \end{pmatrix}. \quad (10)$$

In this case, the effective hadronic interaction Hamiltonian is constructed as

$$H_{\text{eff}} = a[T^a H_a(\bar{3})](\bar{B}_d^c M_c^d)\bar{\nu}_ee + b[T^a \bar{B}_a^b M_b^c H_c(\bar{3})]\bar{\nu}_ee + c[T^a M_a^b \bar{B}_b^c H_c(\bar{3})]\bar{\nu}_ee, \quad (11)$$

where the singlet contribution to η has been neglected. The a, b, c are nonperturbative coefficients. The above Hamiltonian leads to the expectation:

$$\mathcal{B}(\Lambda_c \rightarrow n\bar{K}^0 e^+\nu_e) = \mathcal{B}(\Lambda_c \rightarrow pK^- e^+\nu_e), \quad (12)$$

which is testable in the near future. In fact, the above identity holds in the isospin symmetry, whose breaking effect is much smaller in the charm decays than that of the flavor SU(3) symmetry. In the semi-leptonic decays of $c \rightarrow se^+\nu_e$, the isospins do not change, $\Delta I = 0$. It should be stressed here that this identity is applicable to both resonant and non-resonant contributions.

The branching fraction for the inclusive decay of the Λ_c into an electron has been measured as [11]

$$\mathcal{B}(\Lambda_c \rightarrow e^+ + X) = (4.5 \pm 1.7)\%. \quad (13)$$

Combining the results for the $\Lambda_c \rightarrow \Lambda e^+\nu_e$ in (8), we may expect:

$$\mathcal{B}(\Lambda_c \rightarrow n\bar{K}^0 e^+\nu_e) = \mathcal{B}(\Lambda_c \rightarrow pK^- e^+\nu_e) \sim \mathcal{O}(10^{-3}). \quad (14)$$

III. TWO-BODY NONLEPTONIC Λ_c DECAYS

For two-body nonleptonic decays of the Λ_c , there is no Cabibbo allowed decay mode into a neutron. Two-body decays into a neutron are either singly Cabibbo suppressed,

$$\Lambda_c \rightarrow n\pi^+, \quad \Lambda_c \rightarrow n\rho^+,$$

or doubly Cabibbo suppressed,

$$\Lambda_c \rightarrow nK^+, \quad \Lambda_c \rightarrow nK^{*+}. \quad (15)$$

The nonleptonic Λ_c decays are induced by the operators $[\bar{s}c][\bar{u}d]$ for the Cabibbo-allowed mode and $[\bar{d}c][\bar{u}d]$ for the Cabibbo-suppressed mode. These operators can be decomposed into irreducible representations of flavor SU(3). For instance,

$$(\bar{s}c)(\bar{u}d) = \mathcal{O}_6 + \mathcal{O}_{\overline{15}}, \quad (16)$$

with

$$\begin{aligned} \mathcal{O}_6 &= \frac{1}{2}[(\bar{s}c)(\bar{u}d) - (\bar{u}c)(\bar{s}d)], \\ \mathcal{O}_{\overline{15}} &= \frac{1}{2}[(\bar{s}c)(\bar{u}d) + (\bar{u}c)(\bar{s}d)]. \end{aligned} \quad (17)$$

Perturbative QCD corrections give rise to an enhancement of the coefficient for the \mathcal{O}_6 over the coefficient for the $\mathcal{O}_{\overline{15}}$ by [22, 23]

$$\left[\frac{\alpha_s(m_b)}{\alpha_s(m_W)} \right]^{18/23} \left[\frac{\alpha_s(m_c)}{\alpha_s(m_b)} \right]^{18/25} \sim 2.5. \quad (18)$$

If this is valid, then one has

$$H_{\text{eff}} = eH^{ab}(6)T_{ac}\bar{B}_d^c M_b^d + fH^{ab}(6)T_{ac}M_d^c \bar{B}_b^d + gH^{ab}(6)\bar{B}_a^c M_b^d T_{cd}, \quad (19)$$

with $H^{22}(6) = 1$ for Cabibbo-allowed modes, $H^{23}(6) = H^{32}(6) = -2\sin(\theta_C)$ for singly Cabibbo-suppressed modes, and $H^{33}(6) = +2\sin(\theta_C)^2$ for doubly-Cabibbo-suppressed modes, where θ_C is the Cabibbo angle, and

$$T_{ab} = \epsilon_{abc}T^c. \quad (20)$$

The coefficients e, f, g are the nonperturbative amplitudes.

Using Eq. (19), we find that for the doubly-Cabibbo-suppressed modes:

$$\mathcal{B}(\Lambda_c \rightarrow nK^+) = \mathcal{B}(\Lambda_c \rightarrow pK^0). \quad (21)$$

For the singly-Cabibbo-suppressed modes, we have the decay amplitudes,

$$\mathcal{A}(\Lambda_c \rightarrow n\pi^+) = \sqrt{2}\mathcal{A}(\Lambda_c \rightarrow p\pi^0) = (2f + 2g)\sin(\theta_C), \quad (22)$$

which implies the relation:

$$\mathcal{B}(\Lambda_c \rightarrow n\pi^+) = 2\mathcal{B}(\Lambda_c \rightarrow p\pi^0). \quad (23)$$

Furthermore, we have the amplitudes for Cabibbo-allowed modes:

$$\mathcal{A}(\Lambda_c \rightarrow \Lambda\pi^+) = \frac{1}{\sqrt{6}}(-2e - 2f - 2g), \quad (24)$$

$$\mathcal{A}(\Lambda_c \rightarrow \Sigma^0\pi^+) = \frac{1}{\sqrt{2}}(-2e + 2f + 2g), \quad (25)$$

$$\mathcal{A}(\Lambda_c \rightarrow p\bar{K}^0) = -2e. \quad (26)$$

Thus we can derive the sum rule that can be experimentally examined:

$$\mathcal{B}(\Lambda_c \rightarrow n\pi^+) = \sin^2(\theta_C) \left[3\mathcal{B}(\Lambda_c \rightarrow \Lambda\pi^+) + \mathcal{B}(\Lambda_c \rightarrow \Sigma^0\pi^+) - \mathcal{B}(\Lambda_c \rightarrow p\bar{K}^0) \right]. \quad (27)$$

The recent BES-III data [4] implies

$$\mathcal{B}(\Lambda_c \rightarrow n\pi^+) = \sin^2(\theta_C) [3 \times 1.24\% + 1.27\% - 3.04\%] \sim 0.9 \times 10^{-3}. \quad (28)$$

Measurements in future by BES-III will be able to validate/invalidate the dominance of the sextet assumption in the effective operator.

IV. THREE-BODY NONLEPTONIC Λ_c DECAYS

Compared to two-body decays, three-body Λ_c decays are more involved, since first they can proceed via quasi-two-body process and the non-resonant decays and secondly there are a number of independent amplitudes in SU(3) symmetry. In the following we consider the $NK\pi$ system in the isospin limit:

$$|p\bar{K}^0\pi^0\rangle = \left|\frac{1}{2}\frac{1}{2}\right\rangle\left|\frac{1}{2}\frac{1}{2}\right\rangle|10\rangle = |11\rangle|10\rangle = \frac{1}{\sqrt{2}}|21\rangle + \frac{1}{\sqrt{2}}|11\rangle^{(1)}, \quad (29)$$

$$|pK^-\pi^+\rangle = \left|\frac{1}{2}\frac{1}{2}\right\rangle\left|\frac{1}{2} - \frac{1}{2}\right\rangle|11\rangle = \left(\frac{1}{\sqrt{2}}|10\rangle + \frac{1}{\sqrt{2}}|00\rangle\right)|11\rangle = \frac{1}{2}|21\rangle - \frac{1}{2}|11\rangle^{(1)} + \frac{1}{\sqrt{2}}|11\rangle^{(2)}, \quad (30)$$

$$|n\bar{K}^0\pi^+\rangle = \left|\frac{1}{2} - \frac{1}{2}\right\rangle\left|\frac{1}{2}\frac{1}{2}\right\rangle|11\rangle = \left(\frac{1}{\sqrt{2}}|10\rangle - \frac{1}{\sqrt{2}}|00\rangle\right)|11\rangle = \frac{1}{2}|21\rangle - \frac{1}{2}|11\rangle^{(1)} - \frac{1}{\sqrt{2}}|11\rangle^{(2)}, \quad (31)$$

where the superscripts (1) and (2) are isospin states from $(1-1)$ and $(0-1)$ couplings, respectively, which are independent with each other. Since the Hamiltonian of the $c \rightarrow s\bar{d}u$ transition has $\Delta I = 1$, and the isospin of Λ_c is zero, we can derive the decay amplitudes from the above decompositions:

$$\begin{aligned} \mathcal{A}(\Lambda_c \rightarrow p\bar{K}^0\pi^0) &= \frac{1}{\sqrt{2}}\mathcal{A}^{(1)}, \\ \mathcal{A}(\Lambda_c \rightarrow pK^-\pi^+) &= -\frac{1}{2}\mathcal{A}^{(1)} + \frac{1}{\sqrt{2}}\mathcal{A}^{(2)}, \\ \mathcal{A}(\Lambda_c \rightarrow n\bar{K}^0\pi^+) &= -\frac{1}{2}\mathcal{A}^{(1)} - \frac{1}{\sqrt{2}}\mathcal{A}^{(2)}. \end{aligned} \quad (32)$$

The above amplitudes lead to the sum rule

$$\sqrt{2}\mathcal{A}(\Lambda_c \rightarrow p\bar{K}^0\pi^0) + \mathcal{A}(\Lambda_c \rightarrow pK^-\pi^+) + \mathcal{A}(\Lambda_c \rightarrow n\bar{K}^0\pi^+) = 0. \quad (33)$$

Note that the isospin amplitudes in eq.(32) can be changed if we firstly couple the $K\pi$ states from eq.(29-31), but the sum rule in eq.(33) still holds.

Measurements of branching ratios of the three channels are able to determine the two amplitudes, and in particular investigate the relative strong phases between the two independent decay amplitudes. These phases arise from the final state interactions since if factorization works, the two independent amplitudes are real with vanishing phases at leading order. These amplitudes including phases can provide the essential inputs for the analysis of nonleptonic decays into other baryons like Λ .

From eq.(32), we define the relative strong phase, δ , between $\mathcal{A}^{(1)}$ and $\mathcal{A}^{(2)}$:

$$\frac{\mathcal{A}^{(2)}}{\mathcal{A}^{(1)}} = \left| \frac{\mathcal{A}^{(2)}}{\mathcal{A}^{(1)}} \right| e^{i\delta}. \quad (34)$$

Then the branching fractions can be expressed as

$$\begin{aligned} \mathcal{B}(\Lambda_c \rightarrow p\bar{K}^0\pi^0) &= \frac{1}{2} \left| \mathcal{A}^{(1)} \right|^2, \\ \mathcal{B}(\Lambda_c \rightarrow pK^-\pi^+) &= \frac{1}{4} \left| \mathcal{A}^{(1)} \right|^2 + \frac{1}{2} \left| \mathcal{A}^{(2)} \right|^2 - \frac{1}{\sqrt{2}} \left| \mathcal{A}^{(1)} \right| \left| \mathcal{A}^{(2)} \right| \cos \delta, \\ \mathcal{B}(\Lambda_c \rightarrow n\bar{K}^0\pi^+) &= \frac{1}{4} \left| \mathcal{A}^{(1)} \right|^2 + \frac{1}{2} \left| \mathcal{A}^{(2)} \right|^2 + \frac{1}{\sqrt{2}} \left| \mathcal{A}^{(1)} \right| \left| \mathcal{A}^{(2)} \right| \cos \delta, \end{aligned} \quad (35)$$

where we consider the relative strong phase to understand the final state interaction, and neglect the phase spaces which are actually integrated in the three-body decays. Hence

$$\cos \delta = \frac{\mathcal{B}(n\bar{K}^0\pi^+) - \mathcal{B}(pK^-\pi^+)}{2\sqrt{\mathcal{B}(p\bar{K}^0\pi^0) \left(\mathcal{B}(pK^-\pi^+) + \mathcal{B}(n\bar{K}^0\pi^+) - \mathcal{B}(p\bar{K}^0\pi^0) \right)}}. \quad (36)$$

Defining

$$R_p = \frac{\mathcal{B}(\Lambda_c \rightarrow p\bar{K}^0\pi^0)}{\mathcal{B}(\Lambda_c \rightarrow pK^-\pi^+)}, \quad R_n = \frac{\mathcal{B}(\Lambda_c \rightarrow n\bar{K}^0\pi^+)}{\mathcal{B}(\Lambda_c \rightarrow pK^-\pi^+)}, \quad (37)$$

we have

$$\cos \delta = \frac{R_n - 1}{2\sqrt{R_p(1 + R_n - R_p)}}. \quad (38)$$

From the recent measurement by BESIII [4], $R_p = 0.64 \pm 0.06$. Then $\cos \delta$ can be obtained once the R_n is measured. The relation between $\cos \delta$ and R_n is shown in Fig.1. Since $-1 \leq \cos \delta \leq 1$, we have $0.017 \leq R_n \leq 4.54$, and then the branching fraction of $\Lambda_c \rightarrow n\bar{K}^0\pi^+$ is obtained as,

$$0.04\% \leq \mathcal{B}(\Lambda_c \rightarrow n\bar{K}^0\pi^+)_{\text{Belle}} \leq 33\%, \quad (39)$$

$$0.035\% \leq \mathcal{B}(\Lambda_c \rightarrow n\bar{K}^0\pi^+)_{\text{BESIII}} \leq 28\%. \quad (40)$$

As we can see that this constraint is rather loose, thus the experimental measurements are requested.




FIG. 1: Correlation between $\cos \delta$ and R_n , with δ as the strong phase difference in Eq. (34) and R_n as the ratio of branching fractions in Eq. (37).

V. SUMMARY

Unlike the bottom hadron decays where the momentum transfer is typically large enough to ensure the perturbation theory in QCD, charmed meson and baryon decays are very difficult to understand. Due to the limited energy release, the factorization scheme based on the expansion of $1/m_c$ and $1/E$ is not always valid. Flavor SU(3) symmetry is a powerful tool to analyze the charmed baryon decays, which has been argued to work better than in charmed meson decays, however its validity has to be experimentally examined. Since there is not much data on Ξ_c decays, exclusive Λ_c decays into a neutron are essential for the test of flavor symmetry and investigating final state interactions in charmed baryon decays.

In this work, we have discussed the roles of the exclusive Λ_c decays into a neutron in testing the flavor symmetry and final state interactions. We found that the semileptonic decays into a neutron provide the most-straightforward way to explore the flavor SU(3) symmetry. Two-body nonleptonic decays are capable to examine the assumption of the sextet dominance mechanism. While three-body non-leptonic decays into a neutron are of great interest to explore the final state interactions in Λ_c decays. All these decay modes have not been experimentally observed to date.

Acknowledgements:

The authors are very grateful to Xiao-Gang He, Lei Li, Xiao-Rui Lyu, and Yang-Heng Zheng for enlightening discussions. This work is supported in part by National Natural Science Foundation of China under Grant No.11575110, 11521505, 11347027, 11505083, 11235005 and 11375208, Natural Science Foundation of Shanghai under Grant No.15DZ2272100 and No.15ZR1423100, by the Open Project Program of State Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, China (No.Y5KF111CJ1), and by Scientific Research Foundation

for Returned Overseas Chinese Scholars, State Education Ministry.

-
- [1] A. Zupanc *et al.* [Belle Collaboration], Phys. Rev. Lett. **113**, no. 4, 042002 (2014) doi: 10.1103/PhysRevLett.113.042002 [arXiv:1312.7826 [hep-ex]].
 - [2] D. E. Jaffe *et al.* [CLEO Collaboration], Phys. Rev. D **62**, 072005 (2000) doi: 10.1103/PhysRevD.62.072005 [hep-ex/0004001].
 - [3] S. B. Yang *et al.* [Belle Collaboration], arXiv:1512.07366 [hep-ex].
 - [4] M. Ablikim *et al.* [BESIII Collaboration], arXiv:1511.08380 [hep-ex].
 - [5] D. M. Asner *et al.*, Int. J. Mod. Phys. A **24**, S1 (2009) [arXiv:0809.1869 [hep-ex]].
 - [6] T. Aushev *et al.*, arXiv:1002.5012 [hep-ex].
 - [7] H. Y. Cheng and B. Tseng, Phys. Rev. D **46**, 1042 (1992) [Phys. Rev. D **55**, 1697 (1997)]. doi: 10.1103/PhysRevD.55.1697, 10.1103/PhysRevD.46.1042
 - [8] P. Zenczykowski, Phys. Rev. D **50**, 402 (1994) doi: 10.1103/PhysRevD.50.402 [hep-ph/9309265].
 - [9] T. Uppal, R. C. Verma and M. P. Khanna, Phys. Rev. D **49**, 3417 (1994). doi: 10.1103/PhysRevD.49.3417
 - [10] Fayyazuddin and Riazuddin, Phys. Rev. D **55**, 255 (1997) [Phys. Rev. D **56**, 531 (1997)]. doi: 10.1103/PhysRevD.55.255, 10.1103/PhysRevD.56.531
 - [11] K. A. Olive *et al.* [Particle Data Group Collaboration], Chin. Phys. C **38**, 090001 (2014). doi: 10.1088/1674-1137/38/9/090001
 - [12] M. J. Savage and R. P. Springer, Phys. Rev. D **42**, 1527 (1990). doi:10.1103/PhysRevD.42.1527
 - [13] Y. Kohara, Phys. Rev. D **44**, 2799 (1991). doi:10.1103/PhysRevD.44.2799
 - [14] R. C. Verma and M. P. Khanna, Phys. Rev. D **53**, 3723 (1996) doi:10.1103/PhysRevD.53.3723 [hep-ph/9506394].
 - [15] L. L. Chau, H. Y. Cheng and B. Tseng, Phys. Rev. D **54**, 2132 (1996) doi:10.1103/PhysRevD.54.2132 [hep-ph/9508382].
 - [16] K. K. Sharma and R. C. Verma, Phys. Rev. D **55**, 7067 (1997) doi:10.1103/PhysRevD.55.7067 [hep-ph/9704391].
 - [17] M. J. Savage, Phys. Lett. B **257**, 414 (1991). doi:10.1016/0370-2693(91)91917-K
 - [18] M. Gronau and J. L. Rosner, Phys. Rev. D **89**, no. 3, 037501 (2014) [Phys. Rev. D **91**, no. 11, 119902 (2015)] doi:10.1103/PhysRevD.91.119902, 10.1103/PhysRevD.89.037501 [arXiv:1312.5730 [hep-ph]].
 - [19] X. G. He and G. N. Li, Phys. Lett. B **750**, 82 (2015) doi:10.1016/j.physletb.2015.08.048 [arXiv:1501.00646 [hep-ph]].
 - [20] M. He, X. G. He and G. N. Li, Phys. Rev. D **92**, no. 3, 036010 (2015) doi:10.1103/PhysRevD.92.036010 [arXiv:1507.07990 [hep-ph]].
 - [21] M. Ablikim *et al.* [BESIII Collaboration], Phys. Rev. Lett. **115**, no. 22, 221805 (2015) doi: 10.1103/PhysRevLett.115.221805 [arXiv:1510.02610 [hep-ex]].
 - [22] M. K. Gaillard and B. W. Lee, Phys. Rev. Lett. **33**, 108 (1974). doi: 10.1103/PhysRevLett.33.108
 - [23] G. Altarelli and L. Maiani, Phys. Lett. B **52**, 351 (1974). doi:10.1016/0370-2693(74)90060-4