

Optimal Bounds on Functions of Quantum States under Quantum Channels

Chi-Kwong Li ^{*1}, Diane Christine Pelejo ^{†1,2}, and Kuo-Zhong Wang ^{‡3}

¹Department of Mathematics, College of William and Mary

²Department of Applied Science, College of William and Mary

³Department of Mathematics, National Chiao Tung University

Abstract

Let $(\sigma_1, \sigma_2) \mapsto D(\sigma_1, \sigma_2)$ be a scalar function such as the trace distance, the fidelity, and the relative entropy, etc. For two given quantum states ρ_1 and ρ_2 , we determine optimal bounds for $D(\rho_1, \Phi(\rho_2))$ for $\Phi \in \mathcal{S}$ for different classes of functions $D(\cdot, \cdot)$, where \mathcal{S} is the set of unitary quantum channels, the set of mixed unitary channels, the set of unital quantum channels, and the set of all quantum channels. Furthermore, we determine states $\sigma = \Phi(\rho_2)$ that attain the optimal values and discuss the uniqueness of such states.

Keywords: Quantum states, density matrices, quantum channels, unital channels, mixed unitary channels, fidelity, relative entropy, majorization, eigenvalues.

AMS Classification. 15A18, 15A42, 81P40

1 Introduction

In quantum sciences research, one often compares a pair of quantum states ρ_1, ρ_2 by considering some scalar functions $D(\rho_1, \rho_2)$. For instance, in quantum information and quantum control, one would like to measure the ‘distance’ between a state ρ_1 and another state ρ_2 which go through a quantum channel or a quantum operation Φ . Suppose quantum states are represented as $n \times n$ density matrices, i.e., positive semi-definite matrices with trace one. The following measures are often used [1, 2]:

$$(\text{tr}|\rho_1 - \rho_2|^2)^{1/2}, \quad \frac{1}{2}\text{tr}|\rho_1 - \rho_2|, \quad \sqrt{2}\sqrt{1 - \text{tr}|\sqrt{\rho_1}\sqrt{\rho_2}|}, \quad (1)$$

which are known as the Hilbert-Schmidt (HS) distance, the trace distance and the Bures distance, respectively. Here $|\rho|$ is the positive semidefinite square root of $\rho^\dagger \rho$. In particular, the Bures distance is a function of the fidelity [3]

$$F(\rho_1, \rho_2) = \text{tr}|\sqrt{\rho_1}\sqrt{\rho_2}|. \quad (2)$$

The purpose of this paper is to study the following.

*ckli@math.wm.edu

†dppelejo@email.wm.edu

‡kzwang@math.nctu.edu.tw

Problem 1.1 *Let D be a scalar function on a pair of quantum states. Suppose ρ_1, ρ_2 are two quantum states and \mathcal{S} is a set of quantum channels. Determine the optimal bounds for $D(\rho_1, \Phi(\rho_2))$ for $\Phi \in \mathcal{S}$, and also the states $\sigma = \Phi(\rho_2)$ attaining the optimal bounds.*

These optimal bounds provide insight on the geometry of certain sets of quantum states [4, 5] and play an important role in quantum state discrimination [6, 7, 8]. Physically, if a quantum state ρ_2 goes through some quantum channel Φ , one would like to know $D(\rho_1, \Phi(\rho_2))$ for another fixed quantum state ρ_1 . If Φ is under our control, a solution to this problem can help us select Φ to attain the maximum or minimum value for $D(\rho_1, \Phi(\rho_2))$. On the other hand, if we only know that Φ lies in a certain class of quantum channels, then the solution will tell us the range of values where $D(\rho_1, \Phi(\rho_2))$ lies.

Denote by $\mathcal{M}_n, \mathcal{H}_n, \mathcal{D}_n$ the set of $n \times n$ complex matrices, the set of $n \times n$ Hermitian matrices, and the set of $n \times n$ density matrices, respectively. Then quantum channels are trace preserving completely positive map $\Phi : \mathcal{M}_n \rightarrow \mathcal{M}_n$ with the operator sum representation

$$\Phi(X) = \sum_{j=1}^r F_j X F_j^\dagger \quad \text{for all } X \in \mathcal{M}_n, \quad (3)$$

where $F_1, \dots, F_r \in M_n$ satisfy $\sum_{j=1}^r F_j^\dagger F_j = I_n$. The map Φ is a unitary channel if $r = 1$ and F_1 is unitary; it is a mixed unitary channel if every F_j is a multiple of a unitary matrix; it is unital if $\Phi(I_n) = I_n$.

In the next two sections, we will obtain results for two general classes of functions $D(\cdot, \cdot)$. The first type of functions will cover the Hilbert-Schmidt (HS) distance and the trace distance. The second type will cover the fidelity, the Bures distance, and also the relative entropy defined by

$$S(\rho_1 || \rho_2) = \text{tr} \rho_1 (\log \rho_1 - \log \rho_2). \quad (4)$$

For each class of functions, we will give the complete solution of Problem 1.1 when \mathcal{S} is the set of unitary quantum channels, the set of mixed unitary channels and the set of unital quantum channels. These will be done in the next two sections. We also consider the set of all quantum channels and obtain a complete answer for the first class of functions, and partial results for the second class of functions. Some concluding remarks and future research directions will be mentioned in Section 4.

In our discussion, we will let $\{E_{11}, E_{12}, \dots, E_{nn}\}$ denote the standard basis for \mathcal{M}_n . By the following result [9, Theorem 3.6], the solutions of Problem 1.1 are the same for the set of mixed unitary channels and the set of unital channels.

Lemma 1.2 *Let $\rho, \sigma \in \mathcal{D}_n$. The following are equivalent.*

1. *There exists a mixed unitary quantum channel Φ such that $\Phi(\rho) = \sigma$.*
2. *There exists a unital quantum channel Φ such that $\Phi(\rho) = \sigma$.*
3. $\lambda(\sigma) \prec \lambda(\rho)$.
4. *There are unitary $U_1, \dots, U_n \in M_n$ such that $\sigma = \frac{1}{n}(U_1^\dagger \rho U_1 + \dots + U_n^\dagger \rho U_n)$.*

2 Schur-Convex Functions

For $p \geq 1$, define the Schatten p -norm of a Hermitian matrix A by

$$\|A\|_p = (\text{tr}|A|^p)^{1/p}. \quad (5)$$

It is not hard to see that if A has eigenvalues a_1, \dots, a_n , then

$$\|A\|_p = \ell_p(a_1, \dots, a_n) = \left\{ \sum_{j=1}^n |a_j|^p \right\}^{1/p}. \quad (6)$$

The Hilbert Schmidt norm is $\|\cdot\|_2$ and, up to a multiple, the trace norm is $\|\cdot\|_1$.

In [4, Theorem 4], the authors observed that

$$\max_{U \text{ is unitary}} \|\rho_1 - U\rho_2 U^\dagger\|_1 = \|\Lambda^\downarrow(\rho_1) - \Lambda^\uparrow(\rho_2)\|_1, \quad (7)$$

and

$$\min_{U \text{ is unitary}} \|\rho_1 - U\rho_2 U^\dagger\|_1 = \|\Lambda^\downarrow(\rho_1) - \Lambda^\downarrow(\rho_2)\|_1, \quad (8)$$

where $\Lambda^\downarrow(X)$ (respectively, $\Lambda^\uparrow(X)$) denotes the diagonal matrix having the eigenvalues of X as diagonal entries arranged in descending order (respectively, ascending order).

Actually, the same result holds if one replaces $\|\cdot\|_1$ by any unitary similarity invariant (u.s.i.) norm $\|\cdot\|$, i.e., a norm $\|\cdot\|$ satisfying $\|UXU^\dagger\| = \|X\|$ for all $X \in \mathcal{H}_n$ and unitary $U \in \mathcal{M}_n$. To describe the full generalization of the result, we need the notion of majorization and Schur-convexity. One may see the excellent monograph [10] for the background. We give the basic definition in the following to facilitate our discussion.

Recall that for $x, y \in \mathbf{R}^n$, we say that x is majorized by y , denoted by $x \prec y$ if the sum of the k largest entries of x is not larger than that of y for $j = 1, \dots, n-1$, and the sum of the entries of x equals to that of y . A function $f : \mathbf{R}^n \rightarrow \mathbf{R}$ is Schur-convex if $f(x) \leq f(y)$ whenever $x \prec y$. It is strictly Schur-convex if $f(x) < f(y)$ whenever $x \prec y$ and $x \neq y$.

Denote by $\lambda(X) = (\lambda_1(X), \dots, \lambda_n(X))$ the vector of eigenvalues of $X \in \mathcal{H}_n$ arranged in descending order. In particular, if $X \in \mathcal{D}_n$, then $\lambda(X)$ is a vector in the set

$$\Omega_n = \{(x_1, \dots, x_n) : x_1 \geq \dots \geq x_n \geq 0, x_1 + \dots + x_n = 1\}. \quad (9)$$

We have the following.

Theorem 2.1 *Suppose the function $D : \mathcal{D}_n \times \mathcal{D}_n \rightarrow \mathbf{R}$ is defined by $D(\sigma_1, \sigma_2) = d(\lambda(\sigma_1 - \sigma_2))$ for a Schur-convex function $d : \mathbf{R}^n \rightarrow \mathbf{R}$. Then*

$$\max_{U \text{ is unitary}} D(\rho_1, U\rho_2 U^\dagger) = D(\Lambda^\downarrow(\rho_1), \Lambda^\uparrow(\rho_2)), \quad (10)$$

and

$$\min_{U \text{ is unitary}} D(\rho_1, U\rho_2 U^\dagger) = D(\Lambda^\downarrow(\rho_1), \Lambda^\downarrow(\rho_2)). \quad (11)$$

The maximum is attained at $U\rho_2 U^\dagger$ if there exists a unitary V such that $V\rho_1 V^\dagger = \Lambda^\downarrow(\rho_1)$ and $VU\rho_2 U^\dagger V^\dagger = \Lambda^\uparrow(\rho_2)$. The maximum is attained at $U\rho_2 U^\dagger$ if there exists a unitary V such that $VU\rho_2 U^\dagger V^\dagger = \Lambda^\downarrow(\rho_2)$. The converses of the two preceding statements are also true if d is strictly Schur-convex.

Theorem 2.1 provides a complete solution to Problem 1.1 for the set \mathcal{S} of unitary channels if $D(\sigma_1, \sigma_2) = d(\lambda(\sigma_1 - \sigma_2))$ for a Schur-convex function $d(\cdot)$. In particular, it provides information about the state $\sigma = \Phi(\rho_2)$ that attains the maximum and minimum values. For example, take $\rho_1 = \text{diag}(.55, .45, 0)$ and $\rho_2 = \text{diag}(.35, .33, .32)$, and let $\|\cdot\|$ be any u.s.i. norm. Since all u.s.i. norms are Schur-convex, then $\|\rho_1 - \rho_2\|$ and $\|\rho_1 - \text{diag}(.32, .33, .35)\|$ will yield the minimum and maximum values in the set $\{\|\rho_1 - U\rho_2U^\dagger\| : U \text{ unitary}\}$. Furthermore, if we choose a norm $\|\cdot\|$ that corresponds to a strictly Schur-convex function such as the Schatten p -norm for $p \in (1, \infty)$, then the lower bound and upper bound can only occur at the matrices ρ_2 and $\text{diag}(.32, .33, .35)$, respectively. On the other hand, for the Schatten 1-norm, i.e., the trace norm, the minimum may occur at other matrices such as $U\rho_2U^\dagger = \text{diag}(.33, .35, .32)$ and the maximum may occur at other matrices such as $U\rho_2U^\dagger = \text{diag}(.33, .32, .35)$. Another situation where the optimal is attained by multiple states may arise when ρ_1 has repeated eigenvalues. For example, if $\rho = \frac{1}{n}I_n$, then for any $\Phi \in \mathcal{S}$, $\Phi(\rho_2)$ attains the maximum/minimum.

Next, we turn to Problem 1.1 for the set \mathcal{S} of mixed unitary channels and unital channels. By Lemma 1.2, and the results in [11], we have the following solution of Problem 1.1 if $D(\sigma_1, \sigma_2) = d(\lambda(\sigma_1 - \sigma_2))$ for a Schur-convex function $d(\cdot)$ and \mathcal{S} is the set of mixed unitary channels or the set of unital channels. Furthermore, as shown in Lemma 1.2, we can always construct the mixed unitary channel of the form

$$\sigma \mapsto \frac{1}{n}(U_1\rho U_1^\dagger + \cdots + U_n\rho U_n^\dagger) \quad (12)$$

for some unitary $U_1, \dots, U_n \in M_n$.

Theorem 2.2 Suppose the function $D : \mathcal{D}_n \times \mathcal{D}_n \rightarrow \mathbf{R}$ is defined by $D(\sigma_1, \sigma_2) = d(\lambda(\sigma_1 - \sigma_2))$ for a Schur-convex function $d : \mathbf{R}^n \rightarrow \mathbf{R}$. Let \mathcal{S} be the set of mixed unitary channels or the set of unital channels acting on \mathcal{M}_n . Then

$$\max_{\Phi \in \mathcal{S}} D(\rho_1, \Phi(\rho_2)) = D(\Lambda^\downarrow(\rho_1), \Lambda^\uparrow(\rho_2)) \quad (13)$$

and

$$\min_{\Phi \in \mathcal{S}} D(\rho_1, \Phi(\rho_2)) = D\left(\Lambda^\downarrow(\rho_1), \sum_{j=1}^n d_j E_{jj}\right), \quad (14)$$

where (d_1, \dots, d_n) is determined by the following algorithm:

Step 0. Set $(\Delta_1, \dots, \Delta_n) = \lambda(\rho_1) - \lambda(\rho_2)$.

Step 1. If $\Delta_1 \geq \cdots \geq \Delta_n$, then set $(d_1, \dots, d_n) = \lambda(\rho_1) - (\Delta_1, \dots, \Delta_n)$ and stop.

Else, go to Step 2.

Step 2. Let $1 \leq j < k \leq \ell \leq n$ be such that

$$\Delta_1 \geq \cdots \geq \Delta_{j-1} > \Delta_j = \cdots = \Delta_{k-1} < \Delta_k = \cdots = \Delta_\ell \neq \Delta_{\ell+1}.$$

Replace each $\Delta_j, \dots, \Delta_\ell$ by $(\Delta_j + \cdots + \Delta_\ell)/(\ell - j + 1)$, and go to Step 1.

The maximum is attained at $\Phi \in \mathcal{S}$ if there exists a unitary V satisfying $V\rho_1V^\dagger = \Lambda^\downarrow(\rho_1)$ and $V\Phi(\rho_2)V^\dagger = \Lambda^\uparrow(\rho_2)$. The minimum is attained at $\Phi \in \mathcal{S}$ if there exists a unitary V satisfying $V\rho_1V^\dagger = \Lambda^\downarrow(\rho_1)$ and $V\Phi(\rho_2)V^\dagger = (\sum_{j=1}^n d_j E_{jj})$. The converses of the above two statements also hold if d is strictly Schur-convex.

Here is an example illustrating the construction of the vectors (d_1, \dots, d_n) in the theorem.

Example 2.3 Let $\rho_1 = \frac{1}{10}\text{diag}(4, 3, 3, 0)$ and $\rho_2 = \frac{1}{10}\text{diag}(5, 2, 2, 1)$.

Apply Step 0. Set $(\Delta_1, \dots, \Delta_4) = \frac{1}{10}(4, 3, 3, 0) - \frac{1}{10}(5, 2, 2, 1) = \frac{1}{10}(-1, 1, 1, -1)$.

Apply Step 2. Change $(\Delta_1, \dots, \Delta_4)$ to $\frac{1}{10}(1/3, 1/3, 1/3, -1)$.

Apply Step 1. Set $(d_1, \dots, d_4) = \frac{1}{10}(4, 3, 3, 0) - \frac{1}{10}(1/3, 1/3, 1/3, -1) = \frac{1}{30}(11, 8, 8, 3)$.

Finally, we consider the set \mathcal{S} of all quantum channels. It is known that for any two quantum states, there is a quantum channel sending the first one to the second one. We have the following.

Theorem 2.4 Suppose the function $D : \mathcal{D}_n \times \mathcal{D}_n \rightarrow \mathbf{R}$ is defined by $D(\sigma_1, \sigma_2) = d(\lambda(\sigma_1 - \sigma_2))$ for a Schur-convex function $d : \mathbf{R}^n \rightarrow \mathbf{R}$. Let \mathcal{S} be the set of all quantum channels acting on \mathcal{M}_n . Then

$$\max_{\Phi \in \mathcal{S}} D(\rho_1, \Phi(\rho_2)) = D(\Lambda^\downarrow(\rho_1), E_{nn}) \quad \text{and} \quad \min_{\Phi \in \mathcal{S}} D(\rho_1, \Phi(\rho_2)) = D(\rho_1, \rho_1). \quad (15)$$

The minimum is attained at $\Phi \in \mathcal{S}$ if $\Phi(\rho_2) = \rho_1$. The maximum is attained at $\Phi \in \mathcal{S}$ if there exists a unitary V satisfying $V\rho_1 V^\dagger = \Lambda^\downarrow(\rho_1)$ and $V\Phi(\rho_2)V^\dagger = E_{nn}$. If, in addition, d is strictly Schur-convex, then the converses of the two preceding statements are also true.

Proof. The conclusion on the minimum is clear. For the maximum, note that for any $\sigma \in \mathcal{D}_n$,

$$\sum_{j=1}^k \lambda_j(\rho_1 - \sigma) \leq \sum_{j=1}^k \lambda_j(\rho_1) + \sum_{j=1}^k \lambda_j(-\sigma) \leq \sum_{j=1}^k \lambda_j(\rho_1) + \sum_{j=1}^k \lambda_j(-E_{nn}) = \sum_{j=1}^k \lambda_j(\Lambda^\downarrow(\rho_1) - E_{nn})$$

for $j = 1, \dots, n-1$, and $\sum_{j=1}^n \lambda_j(\rho_1 - \sigma) = 0$. Because $d(\cdot)$ is Schur-convex, the result follows. \square

3 Fidelity, relative entropy, and other functions

In this section, we consider Problem 1.1 for other functions including the fidelity

$$F(\rho_1, \rho_2) = \text{tr} \left(\sqrt{\sqrt{\rho_2} \rho_1 \sqrt{\rho_2}} \right) = \|\sqrt{\rho_1} \sqrt{\rho_2}\|_1 = \text{tr} |\rho_1^{1/2} \rho_2^{1/2}|, \quad (16)$$

and the relative entropy

$$S(\rho || \sigma) = \text{tr}(\rho(\log \rho - \log \sigma)) = \text{tr}(\rho \log \rho) - \text{tr}(\rho \log \sigma). \quad (17)$$

In [12, Theorem 2.1], it was shown that if \mathcal{S} is the set of unitary channels, then

$$\max_{\Phi \in \mathcal{S}} F(\rho_1, \Phi(\rho_2)) = F(\Lambda^\downarrow(\rho_1), \Lambda^\downarrow(\rho_2)) = \sum_{j=1}^n \sqrt{\lambda_j(\rho_1)} \sqrt{\lambda_j(\rho_2)}, \quad (18)$$

and

$$\min_{\Phi \in \mathcal{S}} F(\rho_1, \Phi(\rho_2)) = F(\Lambda^\downarrow(\rho_1), \Lambda^\uparrow(\rho_2)) = \sum_{j=1}^n \sqrt{\lambda_j(\rho_1)} \sqrt{\lambda_{n-j+1}(\rho_2)}. \quad (19)$$

If \mathcal{S} is the set of unital channels, it was also shown in [5, Corollary 2.4] that the above minimum is also valid, but determining the maximum is an open problem.

In the following, we consider different functions f and g on quantum states and study upper bounds and lower bounds for a function $D : \mathcal{D}_n \times \mathcal{D}_n \rightarrow \mathbf{R}$ of the form

$$D(\rho_1, \rho_2) = \text{tr} f(\rho_1) g(\Phi(\rho_2)) \quad \text{and} \quad D(\rho_1, \rho_2) = \text{tr} |f(\rho_1) g(\Phi(\rho_2))| \quad (20)$$

with $\Phi \in \mathcal{S}$ for different sets \mathcal{S} of quantum channels. The results will cover a number of important functions in quantum information research, and the techniques based on the theory of majorization can be further extended to other functions.

To present our results, we need some more definitions and results in majorization (see [10]) to present our general theorem.

A scalar function $f : [0, 1] \rightarrow \mathbf{R}$ can be extended to $f : \mathcal{D}_n \rightarrow \mathcal{H}_n$ such that $f(\sigma) = U^\dagger \text{diag}(f(\mu_1), \dots, f(\mu_n))U$ if $\sigma = U^\dagger \text{diag}(\mu_1, \dots, \mu_n)U$, where U is unitary and $\mu_1 \geq \dots \geq \mu_n \geq 0$.

For two vectors $x, y \in \mathbf{R}^n$, x is weakly majorized by y , denoted by $x \prec_w y$ if the sum of the k largest entries of x is not larger than that of y for $k = 1, \dots, n$. Furthermore, for $x, y \in \mathbf{R}^n$ with nonnegative entries, x is log majorized by y , denoted by $x \prec_{\log} y$ if the product of the entries of x is the same as that of y , and the product of the k largest entries of x is not larger than that of y for $k = 1, \dots, n-1$. It is known that $x \prec_{\log} y$ then $x \prec_w y$.

We have the following.

Theorem 3.1 *Let $f, g : [0, 1] \rightarrow \mathbf{R}$, $\rho_1, \rho_2 \in \mathcal{D}_n$.*

(a) *If $f(\rho_1)$ and $g(\rho_2)$ have eigenvalues $a_1 \geq \dots \geq a_n$ and $b_1 \geq \dots \geq b_n$, then*

$$\min_{U \text{ unitary}} \text{tr}(f(\rho_1)g(U^\dagger \rho_2 U)) = \sum_{j=1}^n a_j b_{n-j+1}, \quad \max_{U \text{ unitary}} \text{tr}(f(\rho_1)g(U^\dagger \rho_2 U)) = \sum_{j=1}^n a_j b_j. \quad (21)$$

The minimum is attained at a unitary U if and only if there exists a unitary V such that $V^\dagger f(\rho_1)V = \text{diag}(a_1, \dots, a_n)$ and $V^\dagger g(U^\dagger \rho_2 U)V = g(V^\dagger U^\dagger \rho_2 UV) = \text{diag}(b_n, \dots, b_1)$; The maximum is attained at a unitary U if and only if there exists a unitary V such that $V^\dagger f(\rho_1)V = \text{diag}(a_1, \dots, a_n)$ and $V^\dagger g(U^\dagger \rho_2 U)V = g(V^\dagger U^\dagger \rho_2 UV) = \text{diag}(b_1, \dots, b_n)$.

(b) *If $f(\rho_1)$ and $g(\rho_2)$ have singular values $\alpha_1 \geq \dots \geq \alpha_n$ and $\beta_1 \geq \dots \geq \beta_n$, then*

$$\min_{U \text{ unitary}} \text{tr}|f(\rho_1)g(U^\dagger \rho_2 U)| = \sum_{j=1}^n \alpha_j \beta_{n-j+1}, \quad \max_{U \text{ unitary}} \text{tr}|f(\rho_1)g(U^\dagger \rho_2 U)| = \sum_{j=1}^n \alpha_j \beta_j. \quad (22)$$

The minimum is attained at a unitary U if and only if there exists a unitary V such that $|V^\dagger f(\rho_1)V| = \text{diag}(\alpha_1, \dots, \alpha_n)$ and $|V^\dagger g(U^\dagger \rho_2 U)V| = |g(V^\dagger U^\dagger \rho_2 UV)| = \text{diag}(\beta_n, \dots, \beta_1)$; the maximum is attained at a unitary U if and only if there exists a unitary V such that $|V^\dagger f(\rho_1)V| = \text{diag}(\alpha_1, \dots, \alpha_n)$ and $|V^\dagger g(U^\dagger \rho_2 U)V| = |g(V^\dagger U^\dagger \rho_2 UV)| = \text{diag}(\beta_1, \dots, \beta_n)$.

Proof. Let $f(\rho_1), g(\rho_2)$ have eigenvalues $a_1 \geq \dots \geq a_n$ and $b_1 \geq \dots \geq b_n$, respectively. For any unitary $V \in M_n$ satisfying $V^\dagger f(\rho_1)V = \text{diag}(a_1, \dots, a_n)$, we have

$$\text{tr}(f(\rho_1)g(U^\dagger \rho_2 U)) = \text{tr}(\text{diag}(a_1, \dots, a_n)V^\dagger g(U^\dagger \rho_2 U)V) = \text{tr}(\text{diag}(a_1, \dots, a_n)g(V^\dagger U^\dagger \rho_2 UV)).$$

By [10, II.9 Theorem H.1.g-h], we have

$$\sum_{j=1}^n a_j b_{n-j+1} \leq \sum_{j=1}^n a_j d_j \leq \sum_{j=1}^n a_j b_j. \quad (23)$$

Evidently, the bounds are attained if the unitary matrices U have the said properties. Assertion (a) follows.

Next, suppose $f(\rho_1), g(\rho_2)$ have singular values $\alpha_1 \geq \dots \geq \alpha_n \geq 0$ and $\beta_1 \geq \dots \geq \beta_n \geq 0$, respectively. Suppose $f(\rho_1)g(U^\dagger \rho_2 U)$ has singular values s_1, \dots, s_n . By [10, II.9 Theorem H.1],

$$(\alpha_1 \beta_n, \dots, \alpha_n \beta_1) \prec_{\log} (s_1, \dots, s_n) \prec_{\log} (\alpha_1 \beta_1, \dots, \alpha_n \beta_n), \quad (24)$$

and $\text{tr}|f(\rho_1)g(U^\dagger \rho_2 U)| = \sum_{j=1}^n s_j$ satisfies

$$\sum_{j=1}^n \alpha_j \beta_{n-j+1} \leq \sum_{j=1}^n s_j \leq \sum_{j=1}^n \alpha_j \beta_j. \quad (25)$$

Suppose $V \in M_n$ is unitary such that $V^\dagger f(\rho_1) V = \text{diag}(\xi_1 \alpha_1, \dots, \xi_n \alpha_n)$ with $\xi_1, \dots, \xi_n \in \{-1, 1\}$. One easily construct the unitary $U \in M_n$ so that $g(U^\dagger \rho_2 U)$ attaining the lower and upper bounds. Evidently, only those unitary matrices having the said properties will yield the optimal bounds. Assertion (b) follows. \square

If \mathcal{S} is the set of all unitary channels, then the lower bounds and upper bounds in Theorem 3.1 are attainable by $\text{tr} f(\sigma_1) g(\Psi(\sigma_2))$ for some $\Psi \in \mathcal{S}$. There are no restrictions to the real valued functions f and g in Theorem 3.1. So, it can be applied to a wide variety of situations. For example, if $f(x) = g(x) = \sqrt{x}$, we obtain the result for the fidelity function $F(\sigma_1, \sigma_2) = \text{tr}|f(\sigma_1)g(\sigma_2)|$ and conclude that for any unitary $U \in M_n$,

$$\sum_{j=1}^n [\lambda_j(\rho_1) \lambda_{n-j+1}(\rho_2)]^{1/2} \leq F(\rho_1, U^\dagger \rho_2 U) \leq \sum_{j=1}^n [\lambda_j(\rho_1) \lambda_j(\rho_2)]^{1/2}. \quad (26)$$

If $f(x) = x$ and $g(x) = \log(x)$, then for any unitary $U \in M_n$,

$$\sum_{j=1}^n \lambda_j(\rho_1) \log \lambda_{n-j+1}(\rho_2) \leq \text{tr}(\rho_1 \log(U^\dagger \rho_2 U)) \leq \sum_{j=1}^n \lambda_j(\rho_1) \log \lambda_j(\rho_2). \quad (27)$$

Here we use the convention that $0 \log 0 = 0$ and $a \log 0 = -\infty$ if $a \in (0, 1]$. Applying this result to $S(\sigma_1 || \sigma_2) = \text{tr} \sigma_1 (\log \sigma_1 - \log \sigma_2)$, we have

$$\sum_{j=1}^n \lambda_j(\rho_1) \log(\lambda_j(\rho_1)/\lambda_j(\rho_2)) \leq S(\rho_1 || U^\dagger \rho_2 U) \leq \sum_{j=1}^n \lambda_j(\rho_1) (\log(\lambda_j(\rho_1)/\lambda_{n-j+1}(\rho_2))) \quad (28)$$

for any unitary $U \in M_n$.

Next, we consider the set \mathcal{S} of mixed unitary channels and the set of unital channels. Given $\rho_1, \rho_2 \in \mathcal{D}_n$, from Lemma 1.2, the following statements are true.

(i) For any $\Phi \in \mathcal{S}$, we have $\lambda(\Phi(\rho_2)) \prec \lambda(\rho_2)$.

(ii) If $f(\rho_1)$ has eigenvalues $a_1 \geq \dots \geq a_n \geq 0$, then for any $(x_1, \dots, x_n) \prec \lambda(\rho_2)$, there is $\Phi \in \mathcal{S}$ such that

$$\text{tr}(f(\rho_1)g(\Phi(\rho_2))) = \sum_{j=1}^n a_j g(x_j), \quad \text{and} \quad \text{tr}|f(\rho_1)g(\Phi(\rho_2))| = \sum_{j=1}^n |a_j g(x_j)|. \quad (29)$$

Hence, we have the following.

Theorem 3.2 *Let $f, g : [0, 1] \rightarrow \mathbf{R}$, $\rho_1, \rho_2 \in \mathcal{D}_n$, and Φ be a unital channel. Suppose $f(\rho_1)$ have eigenvalues $a_1 \geq \dots \geq a_n$, singular values $\alpha_1 \geq \dots \geq \alpha_n$, and ρ_2 has eigenvalues $b_1 \geq \dots \geq b_n$.*

(a) *The best lower upper and upper bounds of $\sum_{j=1}^n \text{tr}(f(\rho_1)g(\Phi(\rho_2)))$ equal*

$$\inf \left\{ \sum_{j=1}^n a_j \lambda_{n-j+1}(g(\sigma)) : \sigma \in \mathcal{D}_n, \lambda(\sigma) \prec (b_1, \dots, b_n) \right\}, \quad (30)$$

and

$$\sup \left\{ \sum_{j=1}^n a_j \lambda_j(g(\sigma)) : \sigma \in \mathcal{D}_n, \lambda(\sigma) \prec (b_1, \dots, b_n) \right\}, \quad \text{respectively.} \quad (31)$$

Suppose the function $g(x)$ is increasing concave. Then the infimum value $\sum_{j=1}^n a_j g(b_{n-j+1})$ is attainable, and a unital channel Φ will attain the infimum value if and only if there is a unitary V satisfying $V^\dagger f(\rho_1) V = \text{diag}(a_1, \dots, a_n)$ and $V^\dagger g(\Phi(\rho_2)) V = \text{diag}(g(b_n), \dots, g(b_1))$. In particular, the infimum can be attained at a unitary channel.

(b) The best lower upper and upper bounds of $\sum_{j=1}^n \text{tr}|f(\rho_1)g(\Phi(\rho_2))|$, respectively, are

$$\inf \left\{ \sum_{j=1}^n \alpha_j \lambda_{n-j+1}(|g(\sigma)|) : \sigma \in \mathcal{D}_n, \lambda(\sigma) \prec (b_1, \dots, b_n) \right\}, \quad (32)$$

and

$$\sup \left\{ \sum_{j=1}^n \alpha_j \lambda_j(|g(\sigma)|) : \sigma \in \mathcal{D}_n, \lambda(\sigma) \prec (b_1, \dots, b_n) \right\}. \quad (33)$$

If the functions $f(x)$ and $g(x)$ have non-negative values on $[0, 1]$, then the lower and upper bounds are the same as those in (a). If in addition that g is increasing concave, then the infimum value $\sum_{j=1}^n a_j g(b_{n-j+1})$ is attainable, and the infimum will occur at $\Phi(\rho_2)$ that satisfy the same conditions described in (a).

Proof. (a) We may assume that $V^\dagger f(\rho_1) V = \text{diag}(a_1, \dots, a_n)$. For any $\Phi \in \mathcal{S}$, we have

$$\sum_{j=1}^n a_j d_j = \text{tr}(f(\rho_1)g(\Phi(\rho_2))), \quad (34)$$

where (d_1, \dots, d_n) are the diagonal entries of $V^\dagger g(\Phi(\rho_2)) V$. Hence, (d_1, \dots, d_n) is majorized by $\lambda(g(\Phi(\rho_2)))$, where $\lambda(\Phi(\rho_2))$ are majorized by (b_1, \dots, b_n) . Similar to the proof of Theorem 3.1,

$$\sum_{j=1}^n a_j \lambda_{n-j+1}(g(\Phi(\rho_2))) \leq \sum_{j=1}^n a_j d_j \leq \sum_{j=1}^n a_j \lambda_j(g(\Phi(\rho_2))) \quad (35)$$

Hence the forms of the best lower upper and upper bounds of $\sum_{j=1}^n \text{tr}(f(\rho_1)g(\Phi(\rho_2)))$ holds. If g is increasing concave, we can apply (vi) of Table 2 in [10, 3.B.2] to the negative of the function $\psi : (x_1, \dots, x_n) \in \Omega_n \mapsto \sum_{j=1}^n a_j g(x_{n-j+1})$ to show that ψ is Schur-concave. Thus the minimum occurs at $(x_1, \dots, x_n) = (b_1, \dots, b_n)$.

(b) Note that the singular values of $g(\Phi(\rho_2))$ are $\gamma_1 \geq \dots \geq \gamma_n$, which is a rearrangement of $|g(x_1)|, \dots, |g(x_n)|$, where x_1, \dots, x_n are the eigenvalues of $\Phi(\rho_2)$ satisfies $(x_1, \dots, x_n) \prec (b_1, \dots, b_n)$. Now, the eigenvalues of $|f(\rho_1)g(\Phi(\rho_2))|$ are the singular values of $f(\rho_1)g(\Phi(\rho_2))$, which is log majorized by $(\alpha_1 \gamma_1, \dots, \alpha_n \gamma_n)$ and log majorizes $(\alpha_1 \gamma_n, \dots, \alpha_n \gamma_1)$. Thus,

$$\sum_{j=1}^n \alpha_j \gamma_{n-j+1} \leq \text{tr}|f(\rho_1)g(\Phi(\rho_2))| \leq \sum_{j=1}^n \alpha_j \gamma_j. \quad (36)$$

If $f(x)$ has nonnegative values, then the eigenvalues of $f(\rho_1)$ are the its singular values, and the same holds for $g(\Phi(\rho_2))$. Thus, the results in (a) applies. \square

We can specialize the result to the function $f(x) = x$ and $g(x) = \log(x)$ to conclude that

$$\sum_{j=1}^n \lambda_j(\rho_1) \log \lambda_{n-j+1}(\rho_2) \leq \text{tr} \rho_1 \log \Phi(\rho_2)$$

for any unital channel Φ , and hence

$$S(\rho_1 || \Phi(\rho_2)) = \text{tr} \rho_1 (\log \rho_1 - \log \Phi(\rho_2)) \leq \sum_{j=1}^n \lambda_j(\rho_1) \log(\lambda_j(\rho_1) / \lambda_{n-j+1}(\rho_2)). \quad (37)$$

For the Fidelity function

$$F(\rho_1, \Phi(\rho_2)) = \text{tr} |\rho_1^{1/2} \Phi(\rho_2)^{1/2}| \quad (38)$$

we can deduce the following result in [5]

$$\min_{\Phi \in \mathcal{S}} F(\rho, \Phi(\sigma)) = F(\Lambda^\downarrow(\rho), \Lambda^\uparrow(\sigma)) = \sum_{i=1}^n \sqrt{\lambda_i(\rho)} \sqrt{\lambda_{n-i+1}(\sigma)}. \quad (39)$$

It was noted in [5] that the maximum value is not easy to determine. As shown in Theorem 3.2, the upper bound of $F(\rho_1, \Phi(\rho_2)) = \text{tr} |\rho_1^{1/2} \Phi(\rho_2)^{1/2}|$ is the same as the upper bound of $\text{tr}(\rho_1^{1/2} \Phi(\rho_2)^{1/2})$, and one needs to determine

$$\sup \left\{ \sum_{j=1}^n \lambda_j(\rho_1)^{1/2} x_j^{1/2} : x_1 \geq \dots \geq x_n \geq 0, (x_1, \dots, x_n) \prec \lambda(\rho_2) \right\}. \quad (40)$$

By the continuity of the function $f(x) = g(x) = \sqrt{x}$ and the compactness of the set $\mathcal{R} = \{(x_1, \dots, x_n) : x_1 \geq \dots \geq x_n \geq 0, (x_1, \dots, x_n) \prec \lambda(\rho_2)\}$, we see that supremum is attainable. On the other hand, the determination of the maximum depends heavily on $\lambda(\rho_1)$ and $\lambda(\rho_2)$. For instance, if $\lambda(\rho_2) = (1/n, \dots, 1/n)$, then \mathcal{R} is a singleton and $F(\rho_1, \Phi(\rho_2)) = \text{tr} |\rho_1^{1/2}| / \sqrt{n}$. If $\rho_2 = \text{diag}(1, 0, \dots, 0)$, then \mathcal{R} contains all quantum states, and $F(\rho_1, \Phi(\rho_2)) = 1$ if $\Phi(\rho_2) = \rho_1$. On the other hand, if $\rho_1 = I_n/n$, then $I_n/n \in \mathcal{R}$ for any ρ_2 so that $F(\rho_1, \Phi(\rho_2)) = 1$ for some unital channel Φ .

In the following, we describe how to determine the unital channel Φ that gives rise to $\max F(\rho_1, \Phi(\rho_2))$ for given $\rho_1, \rho_2 \in \mathcal{D}_n$. The result actually covers a larger class of functions.

Theorem 3.3 *Let $D : \mathcal{D}_n \times \mathcal{D}_n \rightarrow \mathbf{R}$ be defined as follows.*

- (a) $D(\sigma_1, \sigma_2) = \text{tr}(f(\sigma_1)g(\sigma_2))$ or $D(\sigma_1, \sigma_2) = \text{tr}|f(\sigma_1)g(\sigma_2)|$, where $f(x) = x^p$ and $g(x) = x^q$ with $p, q > 0$ such that $p + q = 1$, or
- (b) $D(\sigma_1, \sigma_2) = \text{tr}(f(\sigma_1)g(\sigma_2))$ with $f(x) = x$ and $g(x) = \log x$.

Suppose \mathcal{S} is the set of mixed unitary channels or the set of unital channels acting on \mathcal{M}_n . If $\rho_1, \rho_2 \in \mathcal{D}_n$ have eigenvalues $a_1 \geq \dots \geq a_n \geq 0$ and $b_1 \geq \dots \geq b_n \geq 0$, respectively, then

$$\max_{\Phi \in \mathcal{S}} D(\rho_1, \Phi(\rho_2)) = \sum_{j=1}^n f(a_j)g(d_j), \quad (41)$$

where (d_1, \dots, d_n) is determined by the algorithm below.

If $\Phi \in \mathcal{S}$ such that there exists a unitary V satisfying $V^\dagger \rho_1 V = \Lambda^\downarrow(\rho_1)$ and $V^\dagger \Phi(\rho_2) V = (\sum_{j=1}^n d_j E_{jj})$, then the upper bound is attained.

Algorithm for determining $d_1 \geq \dots \geq d_n$:

Step 0. If $a_r > 0$ and $a_{r+1} = \dots = a_n = 0$, let $a = (a_1, \dots, a_r)$ and $b = (b_1, \dots, b_r)$, and set

$(d_{r+1}, \dots, d_n) = (b_{r+1}, \dots, b_n)$. (Here, if $a_n > 0$, then $r = n$ and (d_{r+1}, \dots, d_n) is vacuous.)

Step 1. Let $k \in \{1, \dots, r\}$ be the largest integer such that

$$\frac{1}{a_1 + \dots + a_k} (a_1, \dots, a_k) \prec \frac{1}{b_1 + \dots + b_k} (b_1, \dots, b_k). \quad (42)$$

Step 2. Set $(d_1, \dots, d_k) = \frac{b_1 + \dots + b_k}{a_1 + \dots + a_k} (a_1, \dots, a_k)$. Stop if $k = r$. Otherwise, change r to $r - k$, $a = (a_{k+1}, \dots, a_r)$, $b = (b_{k+1}, \dots, b_r)$; repeat Steps 1 and 2.

Note that in Step 0 of the algorithm above, we can alternatively choose $(d_{r+1}, \dots, d_n) = (b_{r+1}, \dots, b_n)S$ for any doubly stochastic matrix S . Also, in Step 1, $a_1 + \dots + a_k \neq 0$. This implies that $b_1 + \dots + b_k \neq 0$ because otherwise, the maximality of the choice for k in the previous iteration will be contradicted.

By Theorem 3.3, we see that $S(\rho_1 || \Phi(\rho_2)) \geq \text{tr}(\lambda_j(\rho_1) \log(\lambda_j(\rho_j)/d_j))$, where we use the usual convention that $0 \log 0 = 0$ and $a \log 0 = -\infty$ if $a > 0$. The proof of Theorem 3.3 is quite involved, and will be presented in Section 4. we illustrate the results in Theorem 3.3 and Theorem 3.2 in the following example.

Example 3.4 Let $\rho_1 = \frac{1}{10} \text{diag}(4, 3, 3, 0)$ and $\rho_2 = \frac{1}{10} \text{diag}(5, 2, 2, 1)$.

Apply Step 0. Set $d_4 = 0.1$, $a = (.4, .3, .3)$ and $b = (.5, .2, .2)$.

Apply Step 1. Because $(0.4, 0.3)/0.7 \prec (0.5, 0.2)/0.7$, $(0.4, 0.3, 0.3) \prec (0.5, 0.2, 0.2)/0.9$, we set $(d_1, d_2, d_3) = (0.36, 0.27, 0.27)$, and stop.

Hence, $(d_1, d_2, d_3, d_4) = (0.36, 0.27, 0.27, 0.1)$. For the set \mathcal{S} of unital channels,

$$\min_{\Phi \in \mathcal{S}} F(\rho_1, \Phi(\rho_2)) = (\sqrt{4}, \sqrt{3}, \sqrt{3}, 0)(1, \sqrt{2}, \sqrt{2}, \sqrt{5})^t / 10 = (2 + 2\sqrt{6})/10,$$

$$\max_{\Phi \in \mathcal{S}} F(\rho_1, \Phi(\rho_2)) = (\sqrt{4}, \sqrt{3}, \sqrt{3}, 0)(\sqrt{3.6}, \sqrt{2.7}, \sqrt{2.7}, 1)^t / 10 = 3/\sqrt{10}$$

and

$$\min_{\Phi \in \mathcal{S}} S(\rho_1 || \Phi(\rho_2)) = (4, 3, 3)(\log(10/9), \log(10/9), \log(10/9))^t / 10,$$

$$\max_{\Phi \in \mathcal{S}} S(\rho_1 || \Phi(\rho_2)) = (4, 3, 3)(\log 4, \log(3/2), \log(3/2))^t / 10.$$

Next we consider the set \mathcal{S} of all quantum channels. It is known that for any $\sigma_1, \sigma_2 \in \mathcal{D}_n$, there is a quantum channel Φ such that $\Phi(\sigma_1) = \sigma_2$. Similar to Theorem 3.2, we have the following.

Theorem 3.5 Suppose Ω_n is defined as in (9), $f, g : [0, 1] \rightarrow \mathbf{R}$, and $\rho_1, \rho_2 \in \mathcal{D}_n$ are such that $f(\rho_1)$ have eigenvalues $a_1 \geq \dots \geq a_n$ and singular values $\alpha_1 \geq \dots \geq \alpha_n$. Let Φ be a quantum channel.

(a) The best lower and upper bounds of $\sum_{j=1}^n \text{tr}(f(\rho_1)g(\Phi(\rho_2)))$ equal

$$\inf \left\{ \sum_{j=1}^n a_j \lambda_{n-j+1}(g(\sigma)) : \lambda(\sigma) = (x_1, \dots, x_n) \in \Omega_n \right\} \quad (43)$$

and

$$\sup \left\{ \sum_{j=1}^n a_j \lambda_j(g(\sigma)) : \lambda(\sigma) = (x_1, \dots, x_n) \in \Omega_n \right\}, \quad \text{respectively.} \quad (44)$$

Suppose $g(x)$ is increasing concave, then the infimum value equal to $g(0) \sum_{j=1}^{n-1} a_j + a_n g(1)$ is attainable, and $\Phi \in \mathcal{S}$ attains the infimum if and only if there is a unitary V such that $V^\dagger f(\rho_1) V = \text{diag}(a_1, \dots, a_n)$ and $V^\dagger g(\Phi(\rho_2)) V = \text{diag}(g(0), \dots, g(0), g(1))$.

(b) The best lower and upper bounds of $\sum_{j=1}^n \text{tr}|f(\rho_1)g(\Phi(\rho_2))|$ equal

$$\inf \left\{ \sum_{j=1}^n \alpha_j \lambda_{n-j+1}(|g(\sigma)|) : \lambda(\sigma) = (x_1, \dots, x_n) \in \Omega_n \right\} \quad (45)$$

and

$$\sup \left\{ \sum_{j=1}^n \alpha_j \lambda_j(|g(\sigma)|) : \lambda(\sigma) = (x_1, \dots, x_n) \in \Omega_n \right\}, \quad \text{respectively.} \quad (46)$$

If the functions $f(x)$ and $g(x)$ have non-negative values on $[0, 1]$, then the lower and upper bounds are the same as that in (a). If in addition that g is increasing concave, then the infimum value equals $\text{tr}(f(\rho_1)g(\Phi(\rho_2))) = g(0) \sum_{j=1}^{n-1} a_j + a_n g(1)$ is attainable, and will occur at $\Phi(\rho_2)$ satisfying the same conditions as in (a).

In [5], it was proved that if \mathcal{S} is the set of all quantum channels, then

$$\max_{\Phi \in \mathcal{S}} F(\rho_1, \Phi(\rho_2)) = F(\rho_1, \rho_1) = 1 \quad \text{and} \quad \min_{\Phi \in \mathcal{S}} F(\rho_1, \Phi(\rho_2)) = \lambda_{\min}(\rho_1)^{\frac{1}{2}}. \quad (47)$$

By Theorem 3.5 and Lemma 4.1 in the next section, we have the following.

Corollary 3.6 Suppose \mathcal{S} is the set of all quantum channels, and $\rho_1, \rho_2 \in \mathcal{D}_n$ have eigenvalues $a_1 \geq \dots \geq a_n \geq 0$ and $b_1 \geq \dots \geq b_n \geq 0$, respectively. The the following statements hold.

(a) If $D(\sigma_1, \sigma_2) = \text{tr}(f(\sigma_1)g(\sigma_2))$ or $D(\sigma_1, \sigma_2) = \text{tr}|f(\sigma_1)g(\sigma_2)|$ with $f(x) = x^p, g(x) = x^q$ such that $p, q > 0$ and $p + q = 1$, then

$$\max_{\Phi \in \mathcal{S}} D(\rho_1, \Phi(\rho_2)) = 1 \quad \text{and} \quad \min_{\Phi \in \mathcal{S}} D(\rho_1, \Phi(\rho_2)) = f(a_n).$$

(b) For the relative entropy function,

$$\max_{\Phi \in \mathcal{S}} S(\rho_1 || \Phi(\rho_2)) = \infty \quad \text{and} \quad \min_{\Phi \in \mathcal{S}} S(\rho_1 || \Phi(\rho_2)) = 0. \quad (48)$$

Proof. Similar to the proof of Theorem 3.1, we can focus on

$$\sum_{j=1}^n f(a_j)g(z_j) \quad \text{and} \quad \sum_{j=1}^n a_j \log a_j - \sum_{j=1}^n a_j \log z_j \quad (49)$$

over the set $\Omega_n = \{(x_1, \dots, x_n) : x_1 \geq \dots \geq x_n \geq 0, x_1 + \dots + x_n = 1\}$.

(a) The lower bound follows readily from Theorem 3.2. For the upper bound, by Lemma 4.1(b), we have

$$\sum_{j=1}^n f(a_j)g(z_j) \leq \sum_{j=1}^n f(a_j)g(a_j) = \sum_{j=1}^n a_j^p a_j^q = 1 \quad (50)$$

for all $(z_1, \dots, z_n) \in \Omega_n$.

(b) Choose $(z_1, \dots, z_n) = (0, \dots, 0, 1)$. Since $a_1 > 0$, we have

$$\sum_{j=1}^n a_j \log a_j - \sum_{j=1}^n a_j \log z_j = \infty. \quad (51)$$

From Lemma 4.1(b), $\sum_{j=1}^n a_j \log z_j \leq \sum_{j=1}^n a_j \log a_j$ for all $(z_1, \dots, z_n) \in \Omega_n$. Hence

$$\min_{(z_1, \dots, z_n) \in \Omega_n} \left(\sum_{j=1}^n a_j \log a_j - \sum_{j=1}^n a_j \log z_j \right) = 0. \quad (52)$$

The result follows. \square

4 Proof of Theorem 3.3

To prove Theorem 3.3, we need some auxiliary results. We will focus on the case when $a_1 \geq \dots \geq a_n > 0$.

Lemma 4.1 *Suppose f, g are defined as in Theorem 3.3. Given $p_1, \dots, p_\eta, t_1 \in [0, 1]$ such that $p_1 + \dots + p_\eta > 0$, let*

$$F_{p_1, \dots, p_\eta, t_1}(x_1, \dots, x_{\eta-1}) = f(p_1)g(x_1) + \dots + f(p_{\eta-1})g(x_{\eta-1}) + f(p_\eta)g(t_1 - x_1 - \dots - x_{\eta-1}) \quad (53)$$

for $x_1, \dots, x_{\eta-1} \geq 0$ and $x_1 + \dots + x_{\eta-1} \leq t_1$. Then the following statements are true:

- (a) $F_{p_1, p_2, t_1}(x_1)$ is concave for $x_1 \in [0, t_1]$;
- (b) $F_{p_1, \dots, p_\eta, t_1}(x_1, \dots, x_{\eta-1}) < F_{p_1, \dots, p_\eta, t_1}(\alpha p_1, \dots, \alpha p_{\eta-1})$ for all $(x_1, \dots, x_{\eta-1}) \neq \alpha(p_1, \dots, p_{\eta-1})$, where $\alpha = \frac{t_1}{p_1 + \dots + p_\eta}$.

Proof. For $\eta = 2$, we have $F'_{p_1, p_2, t_1}(\frac{p_1 t_1}{p_1 + p_2}) = 0$ and $F''_{p_1, p_2, t_1}(x_1) < 0$ for all $x_1 \in (0, t_1)$. Hence, (a) holds and in the case $\eta = 2$, (b) is true. Assume that $\eta = k > 2$. F_{p_1, \dots, p_k, t_1} is continuous in $\Gamma_k \equiv \{(x_1, \dots, x_{k-1}) : 0 \leq x_i \leq t_1, x_1 + \dots + x_{k-1} \leq t_1\}$. Since Γ_k is compact, there exists $(\hat{x}_1, \dots, \hat{x}_{k-1}) \in \Gamma_k$ such that

$$F_{p_1, \dots, p_k, t_1}(\hat{x}_1, \dots, \hat{x}_{k-1}) = \max_{\Gamma_k} F_{p_1, \dots, p_k, t_1}. \quad (54)$$

From the case $\eta = 2$, we get $\hat{x}_j = \frac{\hat{x}_j + \hat{x}_i}{p_j + p_i} p_j$ for all i, j and $i \neq j$. This implies that $\hat{x}_j = \alpha p_j$ for all j . Since $\hat{x}_1 + \dots + \hat{x}_k = t_1$, we obtain $\alpha = \frac{t_1}{p_1 + \dots + p_k}$ and then (b) holds. \square

Theorem 4.2 *Let f, g be defined as in Theorem 3.3. Suppose $a = (a_1, \dots, a_n), b = (b_1, \dots, b_n), x = (x_1, \dots, x_n)$ are nonnegative decreasing sequences with $a_n > 0$ and that $x \prec b$ satisfies*

$$\sum_{j=1}^n f(a_j)g(x_j) \equiv f(a)g(x) \geq f(a)g(y) \quad \text{for all } y \prec b. \quad (55)$$

Then the following statements hold.

- (a) There exist $n_0 = 0 < 1 \leq n_1 < n_2 < \dots < n_k = n$ such that for $0 \leq i < k$,

$$\sum_{j=n_i+1}^{n_{i+1}} x_j = \sum_{j=n_i+1}^{n_{i+1}} b_j \quad \text{and} \quad (x_{n_i+1}, \dots, x_{n_{i+1}}) = \alpha_i(a_{n_i+1}, \dots, a_{n_{i+1}}),$$

$$\text{where } \alpha_i = \frac{b_{n_i+1} + \dots + b_{n_{i+1}}}{a_{n_i+1} + \dots + a_{n_{i+1}}}.$$

- (b) The values n_1, \dots, n_k in (a) can be determined as follows:

$$n_1 = \max\{r : \alpha(a_1, \dots, a_r) \prec (b_1, \dots, b_r)\}, \quad (56)$$

and

$$n_j = \max\{r : \alpha(a_{n_{j-1}+1}, \dots, a_r) \prec (b_{n_{j-1}+1}, \dots, b_r)\} \quad \text{for } 1 < j \leq k. \quad (57)$$

Proof. (a) Let $n_1 = \max\{k : (x_1, \dots, x_k) \prec \alpha_0(a_1, \dots, a_k) \text{ for some } \alpha_0\}$. If $n_1 = n$, then by Corollary 3.6, the proof is done.

Suppose that $n_1 < n$. Then $(x_1, \dots, x_{n_1}) = \alpha_0(a_1, \dots, a_{n_1})$ and $x_{n_1+1} \neq \alpha_0 a_{n_1+1}$. We claim that $\sum_{j=1}^{n_1} x_j = \sum_{j=1}^{n_1} b_j$. Suppose that $\sum_{j=1}^{n_1} x_j < \sum_{j=1}^{n_1} b_j$. Let $\beta = \frac{x_{n_1} + x_{n_1+1}}{a_{n_1} + a_{n_1+1}}$. If $x_{n_1} = \beta a_{n_1}$, then $x_{n_1+1} = \beta a_{n_1+1}$. Since $x_{n_1+1} \neq \alpha_0 a_{n_1+1}$ and $x_{n_1} = \alpha_0 a_{n_1}$, $\beta \neq \alpha_0$. Thus, $\beta < \alpha_0$ or $\beta > \alpha_0$.

Case 1. $\beta < \alpha_0$. Let $\hat{x} = (x_1, \dots, x_{n_1-1}, \beta a_{n_1}, \beta a_{n_1+1}, x_{n_1+2}, \dots, x_n)$. We have $\beta a_{n_1} < \alpha_0 a_{n_1} = x_{n_1}$ and $\beta a_{n_1+1} = x_{n_1} + x_{n_1+1} - \beta a_{n_1} > x_{n_1+1}$. Hence \hat{x} is decreasing and $\hat{x} \prec b$. On the other hand,

$$\begin{aligned} & f(a)g(\hat{x}) - f(a)g(x) \\ &= f(a_{n_1})g(\beta a_{n_1}) + f(a_{n_1+1})g(\beta a_{n_1+1}) - (f(a_{n_1})g(x_{n_1}) + f(a_{n_1+1})g(x_{n_1+1})) \\ &= F_{a_{n_1}, a_{n_1+1}, x_{n_1} + x_{n_1+1}} \left(a_{n_1} \frac{x_{n_1} + x_{n_1+1}}{a_{n_1} + a_{n_1+1}} \right) - F_{a_{n_1}, a_{n_1+1}, x_{n_1} + x_{n_1+1}}(x_{n_1}) \\ &> 0 \quad (\text{by Lemma 4.1(b)}). \end{aligned}$$

This is a contradiction.

Case 2. $\beta > \alpha_0$. There exist $m_1 \leq n_1 < m_2$ such that

$$x_{m_1-1} > x_{m_1} = \dots = x_{n_1} \geq x_{n_1+1} = \dots = x_{m_2} > x_{m_2+1}. \quad (58)$$

We will show that $\sum_{j=1}^r x_j < \sum_{j=1}^r b_j$ for $m_1 \leq r < m_2$.

Assertion 1. $\sum_{j=1}^r x_j < \sum_{j=1}^r b_j$ for $n_1 + 1 \leq r < m_2$.

If not, then $\sum_{j=1}^{r_0} x_j = \sum_{j=1}^{r_0} b_j$ for some $n_1 + 1 \leq r_0 < m_2$. Because $\sum_{j=1}^{r_0+1} x_j \leq \sum_{j=1}^{r_0+1} b_j$, we see that $x_{r_0+1} \leq b_{r_0+1}$. Since $\sum_{j=1}^{n_1} x_j < \sum_{j=1}^{n_1} b_j$, we may assume $\sum_{j=1}^r x_j < \sum_{j=1}^r b_j$ for $n_1 \leq r < r_0$. We get $x_{r_0} > b_{r_0} \geq b_{r_0+1}$. But $x_{r_0} = x_{r_0+1} \leq b_{r_0+1}$. This is a contradiction. Thus,

$$\sum_{j=1}^r x_j < \sum_{j=1}^r b_j \quad \text{for } n_1 + 1 \leq r < m_2. \quad (59)$$

Assertion 2. $\sum_{j=1}^r x_j < \sum_{j=1}^r b_j$ for $m_1 \leq r < n_1$.

If not, then $\sum_{j=1}^{r_1} x_j = \sum_{j=1}^{r_1} b_j$ for some $m_1 \leq r_1 < n_1$. Then $x_{r_1} \geq b_{r_1}$. Since $x_{r_1} = \dots = x_{n_1}$ and $b_{r_1} \geq \dots \geq b_{n_1}$, we have $\sum_{j=1}^r x_j \geq \sum_{j=1}^r b_j$ for $r_1 \leq r \leq n_1$. This is impossible since $\sum_{j=1}^{n_1} x_j < \sum_{j=1}^{n_1} b_j$. Hence $\sum_{j=1}^r x_j < \sum_{j=1}^r b_j$ for $m_1 \leq r < n_1$.

By the above argument, $\sum_{j=1}^r x_j < \sum_{j=1}^r b_j$ for $m_1 \leq r < m_2$. Now, let

$$\hat{x} = (x_1, \dots, x_{m_1-1}, x_{m_1} + \delta, x_{m_1+1}, \dots, x_{m_2-1}, x_{m_2} - \delta, x_{m_2+1}, \dots, x_n). \quad (60)$$

For sufficiently small $\delta > 0$, \hat{x} is decreasing and $\hat{x} \prec b$. In fact,

$$\alpha_0 < \beta = \frac{x_{n_1} + x_{n_1+1}}{a_{n_1} + a_{n_1+1}} = \frac{x_{m_1} + x_{m_2}}{a_{n_1} + a_{n_1+1}} = \frac{x_{m_1} + x_{m_2}}{a_{m_1} + a_{n_1+1}} \leq \frac{x_{m_1} + x_{m_2}}{a_{m_1} + a_{m_2}}. \quad (61)$$

The third equality holds because $\alpha_0 a_{m_1} = x_{m_1} = x_{n_1} = \alpha_0 a_{n_1}$. Hence

$$a_{m_1} \frac{x_{m_1} + x_{m_2}}{a_{m_1} + a_{m_2}} > \alpha_0 a_{m_1} = x_{m_1}. \quad (62)$$

Then for sufficiently small $\delta > 0$,

$$\begin{aligned} & f(a)g(\hat{x}) - f(a)g(x) \\ &= f(a_{m_1})g(x_{m_1} + \delta) + f(a_{m_2})g(x_{m_2} - \delta) - (f(a_{m_1})g(x_{m_1}) + f(a_{m_2})g(x_{m_2})) \\ &= F_{a_{m_1}, a_{m_2}, x_{m_1} + x_{m_2}}(x_{m_1} + \delta) - F_{a_{m_1}, a_{m_2}, x_{m_1} + x_{m_2}}(x_{m_1}) \\ &> 0 \quad (\text{by Lemma 4.1}). \end{aligned} \quad (63)$$

This is a contradiction and then $\sum_{j=1}^{n_1} x_j = \sum_{j=1}^{n_1} b_j$. Let

$$n_2 = \max\{k : (x_{n_1+1}, \dots, x_k) = \alpha(a_{n_1+1}, \dots, a_k) \text{ for some } \alpha\}. \quad (64)$$

From the above proof, we also have $\sum_{j=n_1+1}^{n_2} x_j = \sum_{j=n_1+1}^{n_2} b_j$. By induction, we get the desired conclusion.

(b) Suppose $n_1 < \eta \equiv \max\{r : \alpha(a_1, \dots, a_r) \prec (b_1, \dots, b_r)\}$. We have

$$\sum_{j=1}^{n_1} x_j = \sum_{j=1}^{n_1} \alpha_0 a_j = \sum_{j=1}^{n_1} b_j < \sum_{j=1}^{\eta} b_j = \sum_{j=1}^{\eta} \alpha' a_j \quad (65)$$

for some α' . Let $1 < r < k$ with $n_{r-1} < \eta \leq n_r$. Then

$$\sum_{j=1}^{\eta} \alpha' a_j = \sum_{j=1}^{\eta} b_j \geq \sum_{j=1}^{\eta} x_j = \sum_{j=1}^{n_{r-1}} b_j + \sum_{j=n_{r-1}+1}^{\eta} x_j. \quad (66)$$

There is $0 < \alpha'' \leq \alpha'$ such that $\sum_{j=1}^{\eta} \alpha'' a_j = \sum_{j=1}^{\eta} x_j$. Then $\sum_{j=1}^p \alpha'' a_j \leq \sum_{j=1}^p b_j$ for $1 \leq p \leq \eta$. We have $\sum_{j=1}^{n_{r-1}} \alpha'' a_j \leq \sum_{j=1}^{n_{r-1}} b_j = \sum_{j=1}^{n_{r-1}} x_j$. So

$$\sum_{j=n_{r-1}+1}^{\eta} \alpha'' a_j \geq \sum_{j=n_{r-1}+1}^{\eta} x_j = \sum_{j=n_{r-1}+1}^{\eta} \alpha_{n_{r-1}} a_j. \quad (67)$$

Thus $\alpha'' \geq \alpha_{n_{r-1}}$, and hence $\alpha'' a_{\eta} \geq \alpha_{n_{r-1}} a_{\eta} = x_{\eta}$. Let $\hat{x} = (\alpha'' a_1, \dots, \alpha'' a_{\eta}, x_{\eta+1}, \dots, x_n)$. Then \hat{x} is decreasing and $\hat{x} \prec b$. By (a), $n_1 = \max\{k : (x_1, \dots, x_k) = \alpha(a_1, \dots, a_k) \text{ for some } \alpha\}$ and $n_1 < \eta$. Hence, $(x_1, \dots, x_{\eta}) \neq \alpha''(a_1, \dots, a_{\eta})$. We also have $\alpha'' = \frac{x_1 + \dots + x_{\eta}}{a_1 + \dots + a_{\eta}}$. By Lemma 4.1(b),

$$f(a)g(\hat{x}) - f(a)g(x) = \sum_{j=1}^{\eta} f(a_j)g(\alpha'' a_j) - \sum_{j=1}^{\eta} f(a_j)g(x_j) > 0. \quad (68)$$

This is a contradiction. Hence $n_1 = \eta$.

By induction, we only need to show the case n_2 . From the n_1 case, we have $\sum_{j=1}^{n_1} x_j = \sum_{j=1}^{n_1} b_j$. Thus, $(x_{n_1+1}, \dots, x_n) \prec (b_{n_1+1}, \dots, b_n)$, and

$$\sum_{j=n_1+1}^n f(a_j)g(x_j) \leq \max_{(y_{n_1+1}, \dots, y_n) \prec (b_{n_1+1}, \dots, b_n)} \sum_{j=n_1+1}^n f(a_j)g(y_j). \quad (69)$$

On the other hand, for any $(y_{n_1+1}, \dots, y_n) \prec (b_{n_1+1}, \dots, b_n)$, we have $(x_1, \dots, x_{n_1}, y_{n_1+1}, \dots, y_n) \prec b$. Then

$$\begin{aligned} \sum_{j=1}^n f(a_j)g(x_j) &= \max_{y \prec b} f(a)g(y) \\ &\geq \sum_{j=1}^{n_1} f(a_j)g(x_j) + \max_{(y_{n_1+1}, \dots, y_n) \prec (b_{n_1+1}, \dots, b_n)} \sum_{j=n_1+1}^n f(a_j)g(y_j). \end{aligned} \quad (70)$$

This implies that

$$\sum_{j=n_1+1}^n f(a_j)g(x_j) = \max_{(y_{n_1+1}, \dots, y_n) \prec (b_{n_1+1}, \dots, b_n)} \sum_{j=n_1+1}^n f(a_j)g(y_j). \quad (71)$$

From the proof of the case n_1 , the result follows. \square

Proof of Theorem 3.3 From Theorem 3.2, we need only to determine the maximum of $\sum_{j=1}^n f(a_j)g(x_j)$ for $x_1 \geq \dots \geq x_n \geq 0$ and $(x_1, \dots, x_n) \prec (b_1, \dots, b_n)$. Suppose that $a_r > 0$ and $a_{r+1} = \dots = a_n = 0$. Let $\alpha \equiv \sum_{j=1}^n f(a_j)g(d_j)$ attain the maximum for $d_1 \geq \dots \geq d_n \geq 0$ and $(d_1, \dots, d_n) \prec (b_1, \dots, b_n)$. Then $\alpha = \sum_{j=1}^r f(a_j)g(d_j)$ and $(d_1, \dots, d_r) \prec_w (b_1, \dots, b_r)$. Since f is nonnegative and g is increasing,

$$\max\{\sum_{j=1}^r f(a_j)g(x_j) : x_1 \geq \dots \geq x_r \geq 0, (x_1, \dots, x_r) \prec_w (b_1, \dots, b_r)\} \quad (72)$$

$$\leq \max\{\sum_{j=1}^r f(a_j)g(x_j) : x_1 \geq \dots \geq x_r \geq 0, (x_1, \dots, x_r) \prec (b_1, \dots, b_r)\} \equiv \beta. \quad (73)$$

Hence $\alpha \leq \beta$. Given $x_1 \geq \dots \geq x_r \geq 0$ and $(x_1, \dots, x_r) \prec (b_1, \dots, b_r)$, choose $(y_1, \dots, y_n) = (x_1, \dots, x_r, b_{r+1}, \dots, b_n)$. Then $y_1 \geq \dots \geq y_n \geq 0$, $(y_1, \dots, y_n) \prec (b_1, \dots, b_n)$, and $\sum_{j=1}^r f(a_j)g(x_j) = \sum_{j=1}^n f(a_j)g(y_j)$. We obtain $\alpha = \beta$. By Theorem 4.2, we see that the algorithm will produce the state of the form $\Phi(\rho_2)$ attaining the maximum. \square

5 Concluding remarks and further research

Let $(\sigma_1, \sigma_2) \mapsto D(\sigma_1, \sigma_2)$ be a scalar function on quantum states ρ_1, ρ_2 , such as the trace distance, the fidelity function, and the relative entropy. For two given quantum states ρ_1, ρ_2 , we determine optimal bounds for $D(\rho_1, \Phi(\rho_2))$ for $\Phi \in \mathcal{S}$ for different classes of functions $D(\cdot, \cdot)$, where \mathcal{S} is the set of unitary quantum channels, the set of mixed unitary channels, the set of unital quantum channels, and the set of all quantum channels. Specifically, we obtain results for functions of the following form

(a) $D(\sigma_1, \sigma_2) = d(\lambda(\sigma_1 - \sigma_2))$, where $d(X)$ is a Schur-convex function on the eigenvalues of $X \in \mathcal{H}_n$,

(b) $D(\sigma_1, \sigma_2) = \text{tr}(f(\sigma_1)g(\sigma_2))$, and $D(\sigma_1, \sigma_2) = \text{tr}|f(\sigma_1)g(\sigma_2)|$, where $f, g : [0, 1] \rightarrow \mathbf{R}$.

For the class of function in (a), optimal bounds for $D(\rho_1, \Phi(\rho_2))$ are given for $\Phi \in \mathcal{S}$ for the four classes of quantum channels mentioned above. Actually, the results and techniques in Section 2 can be extended to functions of the form

$$D(\sigma_1, \sigma_2) = d(\lambda(\alpha\sigma_1 - \beta\sigma_2)) \quad (74)$$

for given $\alpha, \beta \in \mathbf{R}$, and a Schur-convex function d .

For the class of functions in (b), the optimal lower and upper bounds for $D(\rho_1, \Phi(\rho_2))$ are given for $\Phi \in \mathcal{S}$, where \mathcal{S} is the set of unitary channels. For the set of mixed unitary channels, the set of unital channels, and the set of all quantum channels, we determine the best lower bound if g is an increasing concave function; we also find the best upper bounds for special functions including the fidelity and relative entropy functions. The results and techniques in Section 3 can be extended to cover functions $D : \mathcal{D}_n \times \mathcal{D}_n \rightarrow \mathbf{R}$ of the form $D(\sigma_1, \sigma_2) = \psi(f(\sigma_1)g(\sigma_2))$, where $\psi(X)$ is a Schur concave function on the singular values (eigenvalues or diagonal entries) of the matrix X .

There are many related problems deserving further study. For instance, one may consider Problem 1.1 for a wider class of functions D and different classes of \mathcal{S} . More generally, one may study the optimal bounds for the set

$$\{D(\rho_1, \Phi(\sigma)) : \Phi \in \mathcal{S}, \sigma \in \mathcal{T}\} \quad (75)$$

for a set \mathcal{S} of quantum channels, and a set \mathcal{T} of quantum states. If $\mathcal{T} = \{\sigma_1, \dots, \sigma_k\}$ is a finite set, then one can apply our results to $D(\rho_1, \Phi(\sigma_j))$ for each j to get the optimal bounds for each j , and compare them.

Acknowledgment

The authors would like to thank Professor Jianxin Chen for some helpful discussion. Li is an affiliate member of the Institute for Quantum Computing, University of Waterloo. He is also an honorary professor of the University of Hong Kong, and the University of Shanghai. The research of Li was supported by USA NSF grant DMS 1331021, Simons Foundation Grant 351047, and NNSF of China Grant 11571220. The research of Wang was supported by the Ministry of Science and Technology of the Republic of China under project MOST 104-2918-I-009-001

References

- [1] I. Bengtsson and K. Życzkowski, *Geometry of Quantum States: An Introduction to Quantum Entanglement*, Cambridge University Press, Cambridge (2006).
- [2] A. Fuchs and J. van de Graaf, Cryptographic Distinguishability Measures for Quantum Mechanical States, IEEE Trans. Inf. Th. 45 (1999), 1216 - 1227.
- [3] M.A. Nielsen and I.L. Chuang, *Quantum Computation and Quantum Information*, Cambridge University Press, Cambridge, 2010.
- [4] D. Markham, J.A. Miszczak, Z. Puchala and K. Życzkowski, Quantum State Discrimination: A Geometric Approach, Physics Review A 77 (2008), 042111.
- [5] J. Li, R. Pereira, and S. Plosker, Some Geometric Interpretations of Quantum Fidelity, Linear Algebra Appl. 487 (2015), 158-171.
- [6] C.W. Helstrom, *Quantum Detection and Estimation Theory*, Academic Press, New York (1976).
- [7] C.A. Fuchs, Distinguishability and Accessible Information in Quantum Theory, Ph.D. Thesis, The University of New Mexico, 1996 <http://arxiv.org/pdf/quant-ph/9601020v1.pdf>.
- [8] A. Chefles, Quantum State Discrimination. Authors Issue 6, Contemp. Phys. 41, 401 (2000), 401-424.
- [9] C.-K. Li, Y.-T. Poon, Interpolation by Completely Positive Maps, Linear and Multilinear Algebra 59(2011), 1159–1170
- [10] A.W. Marshall and I. Olkin, and B. Arnold, *Inequalities: Theory of Majorization and Its Application* 2nd edition, Springer, New York, 2011.
- [11] C.-K. Li and N.-K. Tsing, Distance to the Convex Hull of the Unitary Orbit with Respect to Unitary Similarity Invariant Norms, Linear and Multilinear Algebra 25 (1989), 93-103.
- [12] L. Zhang, and S.-M. Fei, Quantum Fidelity and Relative Entropy Between Unitary Orbits, Journal of Physics A: Math. Theor. 47 (2014), 055301.