

Multipartite entanglement in heterogeneous systems

Dardo Goyeneche

Faculty of Physics, Warsaw University, Pasteura 5, 02-093 Warsaw, Poland

Faculty of Applied Physics and Mathematics,

Technical University of Gdańsk, 80-233 Gdańsk, Poland and

Institute of Physics, Jagiellonian University, Kraków, Poland

Jakub Bielawski

Faculty of Physics, Astronomy and Computer Sciences,

Jagiellonian University, Kraków, Poland and

Cracow University of Economics, Kraków, Poland

Karol Życzkowski

Institute of Physics, Jagiellonian University, Kraków, Poland and

Center for Theoretical Physics, Polish Academy of Sciences, Warsaw, Poland

(Dated: Feb. 25, 2016)

Heterogeneous bipartite quantum pure states, composed of two subsystems with a different number of levels, cannot have both reductions maximally mixed. In this work, we demonstrate existence of a wide range of highly entangled states of heterogeneous multipartite systems consisting of $N > 2$ parties such that every reduction to one and two parties is maximally mixed. Two constructions of generating genuinely multipartite maximally entangled states of heterogeneous systems for an arbitrary number of subsystems are presented. Such states are related to quantum error correction codes over mixed alphabets and mixed orthogonal arrays. Additionally, we show that constructed entangled states can be expressed as superpositions of mutually orthogonal GHZ-like states.

Keywords: Genuine multipartite entanglement, quantum error correction codes, mixed orthogonal arrays.

I. INTRODUCTION

Characterization of entanglement in multipartite quantum systems is an important open problem in quantum theory. Entangled states have a fundamental role in quantum teleportation, quantum key distribution, dense coding and error correcting codes [1, 2] and quantum computation [3]. Furthermore, they allow us to generate multipartite Bell inequalities [4] and to formulate an independent proof of the Kochen-Specker theorem [5]. In recent years, multipartite entanglement for homogeneous systems of N qudits systems, i.e., pure states belonging to a Hilbert space $\mathcal{H}_d^{\otimes N}$, has been thoroughly studied [1]. In particular, graph states [6] provide many classes of entangled states, including the maximally entangled ones.

In a more general setup, quantum states belong to a Hilbert space of the form $\mathcal{H}_{d_1}^{\otimes n_1} \otimes \mathcal{H}_{d_2}^{\otimes n_2} \otimes \dots \mathcal{H}_{d_l}^{\otimes n_l}$, where the local Hilbert spaces $\mathcal{H}_{d_1}, \mathcal{H}_{d_2}, \dots, \mathcal{H}_{d_l}$ have possibly different dimensions. Such a Hilbert space contains quantum states associated to *heterogeneous* systems, having different number of levels. For brevity we will refer to such systems as $d_1^{n_1} \times d_2^{n_2} \times \dots \times d_l^{n_l}$. In-

vestigation of entanglement in heterogeneous systems was recently performed in several particular cases, e.g. for tri-partite systems, $2 \times 2 \times n = 2^2 n$ [7, 8] and $2 \times n_1 \times n_2$ [9–11], and for four-partite systems $2^3 \times n$ [12]. A key problem for characterizing entanglement in heterogeneous systems is the lack of a suitable mathematical tool – for instance the Galois fields do not exist in non-prime power dimensions. Therefore, the study of entanglement for heterogeneous systems is more challenging than for the homogeneous case.

Properties of quantum entanglement for heterogeneous bipartite systems are well understood, whereas the complexity of the analogous problem for tripartite systems becomes intractable in general. It has been proven that the problem of calculating the rank of a tensor with three indices over any finite field is NP-complete with respect to the dimension of the tensor [13].

The codification of information in the orbital angular momentum of photons is a resource to generate an unbounded number of discrete levels [14, 15]. This technique has allowed to improve classical [16, 17] and quantum [18, 19] communication protocols. Remarkably, taking into account subsystems consisting of more than two levels each, improves security of some quantum information protocols [20–23], increases capacity of quantum channels [24] and efficiency of quantum gates [25]. Recently, experimentalists paid attention to the entanglement of different degrees of

freedom (e.g. path-polarization) and successfully generated a *qubit-qutrit* hyperentangled state [26]. Additionally, a three partite state composed of one *qubit* and two *qutrits* exhibiting genuinely multipartite entanglement was generated in laboratory [29]. These results provide a motivation to study protection of entanglement in heterogeneous systems under decoherence [27, 28]. It is fair to expect that multipartite entangled states of heterogeneous systems consisting of several qubits and qutrits will be implemented in a near future. However, the theory of quantum entanglement in heterogeneous systems consisting of several subsystems of different sizes is by far not complete. The main aim of this work is to provide a concrete contribution in this direction by considering mixed orthogonal arrays.

A pure multipartite state of a system containing N subsystems is called k -uniform if every density matrix reduced to k subsystems is maximally mixed [1]. Such states are also called maximally multipartite entangled [30] and can be considered as multipartite generalizations of the two-qubit Bell state. Maximally entangled states of homogeneous systems consisting of N subsystems of d levels each are closely related to quantum error correcting codes for messages encoded in an alphabet consisting of d letters [1, 2]. Furthermore, such states are useful for quantum secret sharing protocol [31] and to design holographic quantum codes [32–34].

Highly entangled k -uniform multipartite states can be constructed from graph states [35], orthogonal arrays [36], mutually orthogonal Latin squares and Latin cubes [37] and symmetric matrices [38]. In the present paper, we consider an extension of orthogonal arrays, known as *mixed orthogonal arrays* (MOA) [39], which allows us to construct genuinely multipartite maximally entangled states for heterogeneous systems, related to quantum error correction codes over mixed alphabets [40].

This work is organized as follows: In Section II we establish a link between mixed orthogonal arrays and multipartite entanglement for heterogeneous systems. Additionally, the central concept of *irredundant MOA* is introduced together with basic properties and clarifying examples. In Section III, two powerful constructions to generate irredundant MOA are shown. These constructions allow us to generate maximally entangled states for a wide range of heterogeneous systems. In particular, we prove the existence of 1-uniform and 2-uniform states for heterogeneous systems for any number of parties. In Section IV, we exemplify our constructions by presenting explicitly some new quantum states. Furthermore, we show that every quantum state generated by our proposal can

be written as linear combination involving orthogonal *GHZ*-like states. In Section V, we discuss main results of the work and present some open questions. In Appendix A we show how to find maximally entangled states from redundant MOA and we exemplify this method by finding the simplest 1-uniform state of this work, consisting of two qutrits and one qubit.

II. HETEROGENEOUS ENTANGLEMENT AND MIXED ORTHOGONAL ARRAYS

Orthogonal arrays provide a fundamental tool to design experiments and are known as *Taguchi Designs* [41]. In a previous work we studied orthogonal arrays and its relation to maximally entangled states of homogeneous systems consisting of N subsystems with d levels each [36]. Here, we consider *mixed orthogonal arrays* [39, 42], also called asymmetric orthogonal arrays [43], which form a natural generalization of orthogonal arrays (OA). A mixed orthogonal array $\text{MOA}(r, d_1^{n_1}, d_2^{n_2} \dots d_l^{n_l}, k)$ is an array of r rows and N columns ($r \times N$ array), with $N = n_1 + n_2 + \dots + n_l$ such that the first n_1 columns have symbols from $\{0, 1, \dots, d_1 - 1\}$, the next n_2 columns have symbols from $\{0, 1, \dots, d_2 - 1\}$ and so on, with the property that any $r \times k$ subarray contains every possible combination of k symbols with the same number of appearance. An OA is a particular case of MOA having identical set of symbols in every column, that is, $d_1 = d_2 = \dots = d_l$. The notation considered here for OA and MOA was introduced by Rao [42]. For an excellent introduction to orthogonal arrays and their applications consult the book of Hedayat, Sloane and Stufken [39].

Orthogonal arrays offer a convenient tool to generate highly entangled states for homogeneous multipartite systems composed of N qudits [36], including Dicke states [44], cluster states [45] and graph states [46]. A $\text{MOA}(r, d_1^{n_1}, \dots, d_l^{n_l}, k)$ having $N = n_1 + \dots + n_l$ columns is called *irredundant*, written IrMOA , if every subset of $N - k$ columns contains a different sequence of $N - k$ symbols in every row. It was demonstrated in [36] that an $\text{IrOA}(r, d^N, k)$ leads to a k -uniform state of N subsystems with d levels each. In the present work, we extend the same idea to the mixed alphabets by considering $\text{IrMOA}(r, d_1^{n_1}, d_2^{n_2}, \dots, d_l^{n_l}, k)$

$$\begin{pmatrix} a_{1,1} & a_{1,2} & \dots & a_{1,N} \\ a_{2,1} & a_{2,2} & \dots & a_{2,N} \\ \vdots & \vdots & \dots & \vdots \\ a_{r,1} & a_{r,2} & \dots & a_{r,N} \end{pmatrix}, \quad (1)$$

which lead to k -uniform states of the system $d_1^{n_1} \times$

$d_2^{n_2} \times \cdots \times \cdots, d_l^{n_l}$, consisting of n_1 qubits having d_1 levels, n_2 qubits having n_2 levels and so on. Without loosing of generalization we assume that $d_1 > d_2 > \cdots > d_l$. The generated state, denoted as $|\phi_{d_1^{n_1}, d_2^{n_2}, \dots, d_l^{n_l}}\rangle$, is of the form

$$\begin{aligned} &|a_{1,1}, a_{1,2}, \dots, a_{1,N}\rangle + \\ &|a_{2,1}, a_{2,2}, \dots, a_{2,N}\rangle + \\ &\vdots \\ &|a_{r,1}, a_{r,2}, \dots, a_{r,N}\rangle \end{aligned} \quad (2)$$

For brevity the normalization factors in front of pure states discussed in this work are omitted. In particular, an IrOA(r, d^N, k) (denoted IrOA(r, N, d, k) in [36]) defines a k -uniform state of a homogeneous system consisting of N subsystems with d levels each. This condition means that the partial trace over any selected $N - k$ subsystems forms the maximally mixed state of k qudits.

The classes of states that can be constructed from IrMOA posses two remarkable properties [36]: (A) *Uniformity*: every combination of k symbols has the same number of appearance along the rows. This implies that every reduction to k qudits, $\rho_k = \text{Tr}_{N-k}(|\Phi\rangle\langle\Phi|)$, has a uniform diagonal. (B) *Diagonality*: the irredundancy of the MOA implies that every reduction ρ_k to k qudits forms a diagonal matrix. In this way, OA and irredundancy are the two conditions sufficient to assure uniformity and diagonality of partial traces. This in turn implies that the corresponding pure state is k -uniform. We remark that the same properties hold in the same way for states constructed from IrMOA. Let us present some examples. The array

$$\text{IrOA}(2, 2^2, 1) = \begin{pmatrix} 0 & 0 \\ 1 & 1 \end{pmatrix}, \quad (3)$$

defines the standard Bell state, $|00\rangle + |11\rangle$, while

$$\text{IrOA}(2, 2^3, 1) = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 1 \end{pmatrix}, \quad (4)$$

yields the GHZ state, $|000\rangle + |111\rangle$.

In the case of multilevel systems we have, for instance

$$\text{IrMOA}(4, 4^1 2^2, 1) = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 1 \\ 2 & 1 & 0 \\ 3 & 1 & 1 \end{pmatrix}, \quad (5)$$

which defines a 1-uniform state of a system 4×2^2 consisting of one ququart and two qubits,

$$|\phi_{4^1 2^2}\rangle = |000\rangle + |101\rangle + |210\rangle + |311\rangle. \quad (6)$$

The MOA (5) is listed in the on-line catalog of MOA of strength 2 provided by Kuhfeld [47]. However, state (6) does not capture the aim of our goal. Note that the ququart in the state (6) can be decomposed into two qubits in order to get a 1-uniform state of 4 qubits systems:

$$|\phi_{2^4}\rangle = |0000\rangle + |0101\rangle + |1010\rangle + |1111\rangle, \quad (7)$$

where we considered the identification: $|0\rangle \rightarrow |00\rangle$, $|1\rangle \rightarrow |01\rangle$, $|2\rangle \rightarrow |10\rangle$ and $|3\rangle \rightarrow |11\rangle$. By this reason, in the rest of the work we will be mainly focused on *genuinely heterogeneous systems*, i.e., systems composed of subsystems with coprime levels (e.g. qubits-qutrits or qutrits-ququints). These kind of heterogeneous systems cannot be transformed into homogeneous systems by an identification of symbols, like the one used for the state (6).

III. CONSTRUCTION OF IRREDUNDANT OA AND MOA

In this section, we provide a general framework to construct IrMOA which represents a natural tool to generate genuinely multipartite entangled states for heterogeneous systems with an arbitrary number of parties. In a previous work [36] we used the fact that real Hadamard matrices allow one to construct IrOA($r, 2^N, 2$) for $N > 5$ and suitable values of r . From such arrangements we were able to generate 2-uniform states for $N > 5$ qubits systems. Here, we generalize this result to IrMOA by considering the combinatorial notion of *difference schemes* [39, 48, 49]. A difference scheme $D(s, N, d)$ is an arrangement having s rows, N columns and d different symbols such that the difference (mod d) between every pair of rows contains the d symbols equally often. Thus, there arises the following method to obtain IrMOA of strength one from difference schemes.

CONSTRUCTION 1. *If a difference scheme $D(s, N, d)$ exists then the IrMOA($ds, d^N p_1^1 \dots p_m^1, 1$) exists, where p_1, \dots, p_m are the m distinct prime factors of ds .*

The construction of the IrMOA is simple. From the difference scheme $D(s, N, d)$ we consider the following juxtaposition

$$\begin{aligned} &D(s, N, d) + 0 \\ &D(s, N, d) + 1 \\ &\vdots \\ &D(s, N, d) + d - 1, \end{aligned} \quad (8)$$

where $D(s, N, d) + j$ means that every entry of the difference scheme $D(s, N, d)$ is increased by j and the sum is taken modulo d . This process generates an IrOA($ds, d^N, 1$). In order to obtain the IrMOA($ds, d^N p_1^1 \dots p_m^1, 1$) we have to add m columns C^1, \dots, C^m with entries $(C^l)_j = j \bmod p_l$, where $j = 1, \dots, ds$ and $l = 1, \dots, m$. Let us construct an explicit example from $D(2, 2, 2) = \{00; 01\}$ with help of Construction 1. We obtain then the IrMOA($4, 2^2 2^1, 1$) because the only prime factor of 4 is 2. This IrMOA is in fact the IrOA($4, 2^3, 1$), namely

$$\text{IrOA}(4, 2^3, 1) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 0 & 1 \end{pmatrix}. \quad (9)$$

In order to construct a *genuine IrMOA* we have to consider a difference scheme $D(s, N, d)$ such that ds is not the integer power of a prime number. For example, we can consider $D(2, 3, 3) = \{000; 012\}$ and obtain the genuinely mixed orthogonal array IrMOA($6, 3^3 3^1 2^1, 1$) \equiv IrMOA($6, 3^4 2^1, 1$)

$$\text{IrMOA}(6, 3^4 2^1, 1) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 & 1 \\ 1 & 1 & 1 & 2 & 0 \\ 1 & 2 & 0 & 0 & 1 \\ 2 & 2 & 2 & 1 & 0 \\ 2 & 0 & 1 & 2 & 1 \end{pmatrix}. \quad (10)$$

Let us present a further example. We shall skip the case $D(2, 4, 4) = \{0000; 0123\}$ as $ds = 4 \times 2 = 8$ defines an IrOA instead of a genuine IrMOA. So, we discuss the larger case, $D(2, 5, 5) = \{00000; 01234\}$, which leads to mixed orthogonal array IrMOA($10, 5^5 5^1 2^1, 1$) \equiv IrMOA($10, 5^6 2^1, 1$). That is

$$\text{IrMOA}(10, 5^6 2^1, 1) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 3 & 4 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 2 & 0 \\ 1 & 2 & 3 & 4 & 0 & 3 & 1 \\ 2 & 2 & 2 & 2 & 2 & 4 & 0 \\ 2 & 3 & 4 & 0 & 1 & 0 & 1 \\ 3 & 3 & 3 & 3 & 3 & 1 & 0 \\ 3 & 4 & 0 & 1 & 2 & 2 & 1 \\ 4 & 4 & 4 & 4 & 4 & 3 & 0 \\ 4 & 0 & 1 & 2 & 3 & 4 & 1 \end{pmatrix}. \quad (11)$$

The above examples can be easily generalized to any number of columns N by considering the difference scheme

$$D(2, N, N) = \{0, 0, \dots, 0; 0, 1, \dots, N-1\}. \quad (12)$$

Until now we constructed irredundant mixed orthogonal arrays of strength $k = 1$ from difference

schemes of the form $D(2, N, N)$. However, Construction 1 also works for more general kinds of difference schemes. Here, we have a remarkable observation: From considering Construction 1 and a difference scheme of the form $D(N, N, d)$ we produce a MOA of strength 2 [48] that is irredundant. This property has a central role in this study and it represents the most important contribution of the present paper. We consider this special result as a separate construction:

CONSTRUCTION 2. *If a difference scheme $D(N, N, d)$ exists for $N > 2$ then an IrMOA($dN, d^N p_1^1 \dots p_m^1, 2$) exists, where p_1, \dots, p_m are the m distinct prime factors of dN .*

The case $N = 2$ implies $d=2$ and it does not work here because the number of columns of the resulting IrMOA is three and this array is too small to assure irredundancy. This case generates the 1-uniform state of Eq.(9).

It worth noting that difference schemes of the form $D(p, p, p)$ can be generated from the Fourier matrix $(F_p)_{jk} = \omega_p^{jk} = \omega_p^{D(p,p,p)_{jk}}$ of prime dimension $d = p$, where $\omega_p = e^{2\pi i/p}$. Furthermore, $D(p^m, p^m, p)$ is generated from the tensor product of m Fourier matrices $(F_p^{\otimes m})_{jk} = \omega_p^{D(p^m, p^m, p)_{jk}}$. Let us present some examples of Construction 2 by considering some of these difference schemes. From $D(3, 3, 3) = \{000; 012; 021\}$ we note that there is a single prime factor of $dN = 9$, so we can generate the IrMOA($9, 3^3 3^1, 2$) \equiv IrOA($9, 3^4, 2$). That is,

$$\text{IrOA}^t(9, 3^4, 2) = \begin{pmatrix} 0 & 0 & 0 & 1 & 1 & 1 & 2 & 2 & 2 \\ 0 & 1 & 2 & 1 & 2 & 0 & 2 & 0 & 1 \\ 0 & 2 & 1 & 1 & 0 & 2 & 2 & 1 & 0 \\ 0 & 1 & 2 & 0 & 1 & 2 & 0 & 1 & 2 \end{pmatrix}, \quad (13)$$

where t denotes transposition. Analogously, from difference scheme $D(4, 4, 4) = \{0000; 0123; 0231; 0312\}$ we get IrMOA($16, 4^4 2^1, 2$) given by

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 & 2 & 2 & 2 & 2 & 3 & 3 & 3 & 3 \\ 0 & 1 & 2 & 3 & 1 & 0 & 3 & 2 & 2 & 3 & 0 & 1 & 3 & 2 & 1 & 0 \\ 0 & 2 & 3 & 1 & 1 & 3 & 2 & 0 & 2 & 0 & 1 & 3 & 3 & 1 & 0 & 2 \\ 0 & 3 & 1 & 2 & 1 & 2 & 0 & 3 & 2 & 1 & 3 & 0 & 3 & 0 & 2 & 1 \\ 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 0 & 1 \end{pmatrix}. \quad (14)$$

Here, this IrMOA can be transformed into IrOA($16, 2^9, 2$). Therefore, examples (13) and (14) lead us to OA instead of genuinely mixed OA. The simplest genuine IrMOA arises for difference scheme $D(6, 6, 3)$, as $dN = 18$ is not a power of a prime. This difference scheme can be found in the online catalog of Kuhfeld [52]. To save the space we will not write here the resulting IrMOA($18, 3^6 3^1 2^1$) \equiv IrMOA($18, 3^7 2^1$), but in Section IV we construct the associated quantum state.

IV. QUANTUM STATES FROM IRREDUNDANT MOA

Before constructing k -uniform states related to IrMOA it is interesting to study the maximal values of the strength k allowed for systems $d_1^{n_1} \times \dots \times d_l^{n_l}$. For homogeneous systems consisting of N qudits having d levels each the upper bound is $k \leq N/2$. The states saturating this inequality are called *absolutely maximally entangled* (AME) [31, 50]. For example, Bell states are AME for two qubit systems. For a general $d_1 \times d_2$ system with $d_1 \neq d_2$ AME states do not exist. This is because the von Neumann entropies of the reduced density matrices $\rho_1 = \text{Tr}_2(\rho_{12})$ and $\rho_2 = \text{Tr}_1(\rho_{12})$ are equal, so that both reductions cannot be maximally mixed. Following the same argument, k -uniform states consisting of $N = 2k$ heterogeneous subsystems do not exist [1]. Despite this fact, there is a place for a wide range of k -uniform states with $k < N/2$. To get a more precise upper bound for k note that for a system $d_1 \times \dots \times d_l$, 1-uniform states do not exist if d_1 is larger than the dimension of the complementary system, i.e., $d_2 d_3 \dots d_l$. In general, a k -uniform state do not exist if the product of the size of k local Hilbert spaces is larger than the dimension of the complementary system. This result can be stated as follows: a necessary condition for the existence of a k uniform state of a system $d_1^{n_1} \times \dots \times d_l^{n_l}$ is

$$\left(\prod_{i=1}^k d_i^{n'_i} \right)^2 \leq \prod_{i=1}^l d_i^{n_i}, \quad (15)$$

where $n'_i = \min\{n_i, \max\{k - \sum_{j=1}^{i-1} n_j, 0\}\}$. Here, we recall the already assumed convention, $d_1 > d_2 > \dots > d_l$. In the particular case of N qudits with d levels each Eq.(15) reduces to the standard bound, $k \leq N/2$. Furthermore, a k -uniform state consisting of $n_1 \geq k$ qutrits and n_2 qubits satisfies the bound

$$3^{2k} \leq 3^{n_1} 2^{n_2}, \quad (16)$$

for any n_2 . Let us now to construct 1-uniform and 2-uniform states applying Constructions 1 and 2, respectively. We start by considering the 1-uniform state of 3 qubits arising from IrOA(4, 2³, 1) of Eq.(9) which is accidentally homogeneous,

$$|\phi_{2^3}\rangle = |000\rangle + |011\rangle + |110\rangle + |101\rangle. \quad (17)$$

Here, we followed the identification between IrMOA and quantum states given in Eq.(1) and Eq.(2), where states are not normalized. A genuinely heterogeneous maximally entangled state arises from the array IrMOA(6, 3⁴2¹, 1) of Eq.(10),

which allows us to obtain the 1-uniform state of the system 3⁴ × 2:

$$|\phi_{3^4 2^1}\rangle = |00000\rangle + |01211\rangle + |11120\rangle + |12001\rangle + |22210\rangle + |20121\rangle. \quad (18)$$

Additionally, IrMOA(10, 5⁶2¹, 1), explicitly written in (11) yields the 1-uniform state of the system 5⁶ × 2,

$$|\phi_{5^6 2^1}\rangle = |0000000\rangle + |0123411\rangle + |1111120\rangle + |1234031\rangle + |2222240\rangle + |2340101\rangle + |3333310\rangle + |3401221\rangle + |4444430\rangle + |4012341\rangle. \quad (19)$$

States (18) and (19) can be generalized for any number of parties by considering the difference scheme $(D(s, N, N))_{jk} = jk \pmod{N}$, where $j = 0, \dots, s-1$ and $k = 0, \dots, N-1$. Here, $D(s, N, d)_{jk}$ denotes the entry located at the j -th row and k -th column of the scheme $D(s, N, d)$. Thus, we generate the following 1-uniform state,

$$|\phi_{N^s N}\rangle = \sum_{j=0}^{s-1} \sum_{l=0}^{d-1} \bigotimes_{k=0}^{N-1} |D(s, N, N)_{jk} + l\rangle \otimes |[sl + j]_{p_1} \dots [sl + j]_{p_m}\rangle, \quad (20)$$

where $[X]_a = X \pmod{a}$ and p_1, \dots, p_m are the m distinct prime factors of sN . Furthermore, Construction 2 allows us to generate 2-uniform states. The simplest cases arise from IrOA(9, 3⁴, 2), IrMOA(16, 4⁴2¹, 2) \equiv IrOA(16, 2⁹, 2) and IrOA(25, 5⁶, 2). These arrangements generate 2-uniform states of homogeneous systems: 4 qutrits, 9 qubits and 6 ququints, respectively. Two uniform states characterizing genuine heterogeneous systems are associated to difference schemes $D(N, N, d)$ such that dN is not a power of a prime.

The simplest heterogeneous case arises from $D(6, 6, 3)$ which leads to IrMOA(18, 3⁶3¹2¹, 2) \equiv IrMOA(18, 3⁷2¹, 2) and generates the 2-uniform state of the system 3⁷ × 2:

$$|\phi_{3^7 2^1}\rangle = |00000000\rangle + |00112211\rangle + |01022120\rangle + |01201201\rangle + |02121010\rangle + |02210121\rangle + |11111100\rangle + |11220011\rangle + |12100220\rangle + |12012001\rangle + |10202110\rangle + |10021221\rangle + |22222200\rangle + |22001111\rangle + |20211020\rangle + |20120101\rangle + |21010210\rangle + |21102021\rangle. \quad (21)$$

In the same way, the difference scheme $D(10, 10, 5)$ implies the array IrMOA(50, 5¹⁰5¹2¹, 2) \equiv IrMOA(50, 5¹¹2¹, 2), which produces the state of the system 5¹¹ × 2:

$$\begin{aligned}
|\phi_{5^{11}2^1}\rangle = & |000000000000\rangle + |001312243411\rangle + |012344012320\rangle + |013102434231\rangle + |020443231140\rangle + \\
& |024131302401\rangle + |031421420310\rangle + |032230144121\rangle + |043214321030\rangle + |044023113241\rangle + \\
& |111111111100\rangle + |112423304011\rangle + |123400123420\rangle + |124213040331\rangle + |131004342240\rangle + \\
& |130242413001\rangle + |142032031410\rangle + |143341200221\rangle + |104320432130\rangle + |100134224341\rangle + \\
& |222222222200\rangle + |223034410111\rangle + |234011234020\rangle + |230324101431\rangle + |242110403340\rangle + \\
& |241303024101\rangle + |203143142010\rangle + |204402311321\rangle + |210431043230\rangle + |211240330441\rangle + \\
& |333333333300\rangle + |334140021211\rangle + |340122340120\rangle + |341430212031\rangle + |303221014440\rangle + \\
& |302414130201\rangle + |314204203110\rangle + |310013422421\rangle + |321042104330\rangle + |322301441041\rangle + \\
& |444444444400\rangle + |440201132311\rangle + |401233401220\rangle + |402041323131\rangle + |414332120040\rangle + \\
& |413020241301\rangle + |420310314210\rangle + |421124033021\rangle + |432103210430\rangle + |433412002141\rangle.
\end{aligned} \tag{22}$$

From $D(12, 12, 3)$ we generate and thus a state of the system $3^{13} \times 2$,
 $\text{IrMOA}(36, 3^{12}3^12^1, 2) \equiv \text{IrMOA}(36, 3^{13}2^1, 2)$

$$\begin{aligned}
|\phi_{3^{13}2^1}\rangle = & |00000000000000\rangle + |00001122112211\rangle + |00110011222220\rangle + |00112222001101\rangle + |01020212210110\rangle + \\
& |01022021121021\rangle + |01201201201200\rangle + |01202110022111\rangle + |02121210102020\rangle + |02122101010201\rangle + \\
& |02210120211010\rangle + |02211002120121\rangle + |11111111111100\rangle + |11112200220011\rangle + |11221122000020\rangle + \\
& |11220000112201\rangle + |12101020021210\rangle + |12100102202121\rangle + |12012012012000\rangle + |12010221100211\rangle + \\
& |10202021210120\rangle + |10200212121001\rangle + |10021201022110\rangle + |10022110201221\rangle + |22222222222200\rangle + \\
& |22220011001111\rangle + |22002200111120\rangle + |22001111220001\rangle + |20212101102010\rangle + |20211210010221\rangle + \\
& |20120120120100\rangle + |20121002211011\rangle + |21010102021220\rangle + |21011020202101\rangle + |21102012100210\rangle + \\
& |21100221012021\rangle.
\end{aligned} \tag{23}$$

There exist a useful compact way to write every state arising from Construction 1 and 2. Introducing the shift operator,

$$X|j\rangle = |[j+1]_d\rangle, \tag{24}$$

the GHZ state of N subsystems with d levels each,

$$|GHZ_0\rangle = \sum_{m=0}^{d-1} |m\rangle^{\otimes N}, \tag{25}$$

and its generalization,

$$|GHZ_j\rangle = \bigotimes_{k=0}^{N-1} X^{D(s,N,d)_{jk}} |GHZ_0\rangle |[k]_{p_1} \dots [k]_{p_{m'}}\rangle, \tag{26}$$

we can write

$$|\phi_{d^N p_1^1 \dots p_m^1}\rangle = \sum_{j=0}^{s-1} |GHZ_j\rangle |[k]_{p_{m'+1}} \dots [k]_{p_m}\rangle. \tag{27}$$

where addition in Eq.(24) is understood modulo d , $[k]_{p_i} = k \pmod{p_i}$, $\{p_1, \dots, p_{m'}\}$ are the distinct prime factors of N and $\{p_{m'+1}, \dots, p_m\}$ is the subset of the distinct prime factors of s which are not in common with the prime factors of N .

The superpositions of GHZ-like states (27) should not be confused with a particular class of mixed quantum states called X states [51]. The X states are convex combinations of GHZ-like rank-one projectors $|GHZ\rangle\langle GHZ|$, whereas here we consider superpositions of GHZ-like vectors $|GHZ_j\rangle$.

V. ENDURANCE OF K-UNIFORMITY

In Section IV we generated k -uniform states for heterogeneous systems by considering IrMOA. Here, we show that some columns of IrMOA can be removed without loosing the irredundancy of the orthogonal array in question. As consequence, we are able to generate the family k -uniform states $\{|\phi_{d^x p_1^1 \dots p_m^1}\rangle\}$, where

$$N - \mu_k \leq x \leq N, \quad (28)$$

and m is the number of distinct primes factors of sN . Here, the number μ_k is called *endurance of k -uniformity* and it represents the maximal number of columns of an IrMOA that can be removed so that the remaining MOA preserves both irredundancy and strength k . It is important to stress that the x columns to be removed have to contain d different symbols. Otherwise, the endurance of k -uniformity would be not univocally defined. Observe that the action of removing a column in an IrMOA is not related to a local measurement performed by a given party. Hence, endurance of k -uniformity is not related to the *persistence of entanglement* [45], which represents the minimal number of parties that should make a local measurement in order to assure that the remaining state is a fully separable.

For example, by removing $N - 2$ columns of the IrOA(2,2^N,1) $\{0 \dots 0; 1 \dots 1\}$ we get the IrOA(2,2²,1) $\{00; 11\}$. This operation can be associated to a mapping of the GHZ state of N qubits into the Bell state. Thus, the IrOA(2,2^N,1) has endurance of 1-uniformity $\mu_1 = N - 2$, as 1-uniformity remains after removing $N - 2$ columns of the IrOA. On the other hand, any local measurement on a system prepared in a GHZ state of N parties lead us to a fully separable state, so the GHZ state has persistency of entanglement $P_E = 1$ for any number of parties N . Table I shows the endurance of uniformity for every state explicitly constructed in Section IV

In the on-line catalog of Kuhfeld [52] there are several difference schemes $D(s, N, d)$ for $s \neq 2$ and $s \neq N$ which allow us to generate further 1-uniform and 2-uniform states of heterogeneous systems in the same way as Constructions 1 and 2.

Let us mention that 2-uniform for N qudits having a prime number of levels $d = p$ can be obtained with use of Construction 2 followed by removing of heterogeneous columns. As a final comment, we would like to stress that all the difference schemes $D(s, N, d)$ useful to generate 2-uniform states are such that the $s \times N$ matrix $A_{jk} = \omega^{D(s, N, d)_{jk}}$ has orthogonal rows, where $\omega = e^{2\pi i/d}$, $j = 0, \dots, s - 1$ and $k = 0, \dots, N - 1$. This is not a general prop-

| $ \phi_{d_1^{n_1} d_2^{n_2}}\rangle$ | k | μ_2 | μ_1 | $ \phi_{d_1^{n_1 - \mu_2} d_2^{n_2}}\rangle$ | $ \phi_{d_1^{n_1 - \mu_1} d_2^{n_2}}\rangle$ |
|--------------------------------------|-----|---------|---------|--|--|
| $ \phi_{23}\rangle$ | 1 | - | 0 | - | $ \phi_{23}\rangle$ |
| $ \phi_{3421}\rangle$ | 1 | - | 2 | - | $ \phi_{3221}\rangle$ |
| $ \phi_{5621}\rangle$ | 1 | - | 4 | - | $ \phi_{5221}\rangle$ |
| $ \phi_{3721}\rangle$ | 2 | 2 | 3 | $ \phi_{3521}\rangle$ | $ \phi_{3421}\rangle$ |
| $ \phi_{51121}\rangle$ | 2 | 6 | 7 | $ \phi_{5521}\rangle$ | $ \phi_{5421}\rangle$ |
| $ \phi_{31321}\rangle$ | 2 | 6 | 8 | $ \phi_{3721}\rangle$ | $ \phi_{3521}\rangle$ |

TABLE I: Endurances of uniformity μ_2 and μ_1 for every state $|\phi_{d_1^{n_1} d_2^{n_2}}\rangle$ explicitly presented in Section IV. The first three states are 1-uniform whereas the last three states are 2-uniform. The resulting 2-uniform and 1-uniform states obtained after removing μ_2 and μ_1 columns, respectively, are shown in the 4-th and 5-th column. The 2-uniform state for heterogeneous system belonging to the smallest possible Hilbert space that we found in this work is $|\phi_{3521}\rangle$ (bold).

erty, as we can see from the case $D(8, 8, 4)$ [52] which allows one to generate a 1-uniform state only.

VI. CONCLUDING REMARKS

Multipartite quantum states were usually studied for homogeneous systems consisting of subsystems with the same number of levels. As several experimentally studied physical systems consists of subsystems of different numbers of levels in this work we analyzed quantum entanglement in heterogeneous systems.

Highly entangled quantum states of homogeneous systems consisting of N subsystems with d levels each can be related with irredundant orthogonal arrays [36, 38] containing symbols from a d -letter alphabet. Making use of mixed orthogonal arrays [42, 43] we generalized this relation for quantum states of heterogeneous systems.

In particular, we presented explicit constructions of one and two-uniform states for several heterogeneous quantum systems. Simple cases include, e.g. one uniform state (18) of a system consisting of four qutrits and a qubit and state (19) of a system of six subsystems with five levels each and a single qubit. The state (21) of a system containing seven qutrits and a qubit is two uniform.

It is tempting to believe that some states discussed in this work can be someday realized in laboratory. Furthermore, k -uniform multipartite states for heterogeneous systems can be useful to construct quantum error correction codes. Several questions related to these issues remain open. For example, it is not known for which heterogeneous quantum systems k -uniform states exists. In par-

ticular, it would be interesting to find examples of 3-uniform states for such systems.

Acknowledgements. It is a pleasure to thank P. Horodecki, M. Grassl and A. Sudbery for useful remarks. This work has been supported by the Polish National Science Center under the project number DEC-2011/02/A/ST1/00119, by the European Union FP7 project PhoQuS@UW (Grant Agreement No. 316244) and by the John Templeton Foundation under the project No. 56033.

Appendix A: k -uniform states from redundant MOA

For some systems there exist k -uniform states which are not related to mixed irredundant orthogonal arrays. Such examples were discussed for standard OA, where some minus signs had to be introduced in order to generate certain entangled states for qubits – see Appendix C in Ref. [36]. In this Appendix we construct some k -uniform states by introducing complex phases in the coefficients of the states.

Let us start by considering the state of a system $3 \times 3 \times 2$:

$$|\phi_{3^2 2^1}\rangle = |000\rangle + e^{i\alpha_1}|111\rangle + e^{i\alpha_2}|220\rangle + e^{i\alpha_3}|001\rangle + e^{i\alpha_4}|110\rangle + e^{i\alpha_5}|221\rangle. \quad (\text{A1})$$

For this state, the reductions to the second and third party are maximally mixed for any value of the parameters $\{\alpha_j\}$. That is,

$$\begin{aligned} \rho_B &= \text{Tr}_{AC}(|\phi_{3^2 2^1}\rangle\langle\phi_{3^2 2^1}|) = \frac{1}{3}\mathbb{I}_3 \\ \rho_C &= \text{Tr}_{AB}(|\phi_{3^2 2^1}\rangle\langle\phi_{3^2 2^1}|) = \frac{1}{3}\mathbb{I}_3. \end{aligned}$$

The only restriction to have ρ_A maximally mixed is given by

$$e^{i(\alpha_1+\alpha_2)} + e^{i(\alpha_2+\alpha_3+\alpha_4)} + e^{i(\alpha_4+\alpha_5)} = 0. \quad (\text{A2})$$

Here, the solution is provided by three complex numbers forming an equiangular triangle in the complex plane. A particular solution to this equation is $\alpha_5 = 0$, $\alpha_2 = \alpha_3 = 2\pi/3$ and $\alpha_1 = \alpha_4 = 4\pi/3$. The fact that $\alpha_1 = \dots = \alpha_5 = 0$ is not a solution of Eq.(A2) means that the MOA(6, $3^2 2^1$, 1) generating the state (A1), that is,

$$\begin{pmatrix} 0 & 0 & 0 \\ 1 & 1 & 1 \\ 2 & 2 & 0 \\ 0 & 0 & 1 \\ 1 & 1 & 0 \\ 2 & 2 & 1 \end{pmatrix}, \quad (\text{A3})$$

is not irredundant. Therefore, the following 1-uniform state of systems $3^2 \times 2$ exists

$$|\phi_{3^2 2^1}\rangle = |000\rangle + |221\rangle + \omega(|220\rangle + |001\rangle) + \omega^2(|111\rangle + |110\rangle), \quad (\text{A4})$$

where $\omega = e^{2\pi i/3}$. This state represents the simplest maximally entangled state that we present in this work, in the sense that the dimension of the entire Hilbert space is the smallest possible. We did not find a 1-uniform state for a system 3×2^2 . Finally, note that the classification of entanglement classes for systems $3^2 \times 2$ provided in [10] does not take into account 1-uniform states. To our best knowledge state (A4) did not appear in the literature so far.

-
- [1] A. J. Scott, Multipartite entanglement, quantum-error-correcting codes, and entangling power of quantum evolutions, *Phys. Rev. A* **69**, 052330 (2004).
 - [2] L. Arnaud and N. Cerf, Exploring pure quantum states with maximally mixed reductions, *Phys. Rev. A* **87**, 012319 (2013).
 - [3] R. Jozsa, N. Linden, On the role of entanglement in quantum-computational speed-up, *Proc. R. Soc. Lond. A*, **459**, 2011-2032 (2003).
 - [4] O. Gühne, G. Tóth, P. Hyllus, H. Briegel, Bell Inequalities for Graph States, *Phys. Rev. Lett.* **95**, 120405 (2005).
 - [5] A. Cabello, O. Gühne, D. Rodríguez, Mermin inequalities for perfect correlations, *Phys. Rev. A* **77**, 6, 062106 (2008).
 - [6] M. Hein, W. Dür, J. Eisert, R. Raussendorf, M. Van den Nest, H.-J. Briegel, Entanglement in Graph States and its Applications, *Proc. Int. School of Phys. "Enrico Fermi" on "Quantum Computers, Algorithms and Chaos"*, Varenna, Italy, preprint arXiv:quant-ph/0602096 (2005).
 - [7] C. Yu, H. Song, Y. Wang, Genuine tripartite entanglement semi-monotone for $(2 \times 2 \times n)$ -dimensional systems, *Quant. Inf. & Comp.* **7**, 7, 584 (2007).
 - [8] C. Yu, L. Zhou, H. Song, Genuine tripartite entanglement monotone of $(2 \times 2 \times n)$ -dimensional systems, *Phys. Rev. A* **77**, 2, 022313 (2008).
 - [9] A. Miyake and F. Verstraete, Multipartite entan-

- lement in $2 \times 2 \times N$ quantum systems, Phys. Rev. A **69**, 012101 (2004).
- [10] L. Chen and Y.-X. Chen, Range criterion and classification of true entanglement in a $2 \times M \times N$ system, Phys. Rev. A **73**, 052310 (2006).
 - [11] L. Chen, Y. Chen, Y. Mei, Classification of multipartite entanglement containing infinitely many kinds of states, Phys. Rev. A **74**, 052331 (2006).
 - [12] S. Wang, Y. Lu, G. Long, Entanglement classification of $2 \times 2 \times 2 \times d$ quantum systems via the ranks of the multiple coefficient matrices, Phys. Rev. A **87**, 062305 (2013).
 - [13] J. Hastad, Tensor rank is NP-complete, J. Algorithms **11**, 4, 644-654 (1990).
 - [14] J. Torres, L. Torner, *Twisted Photons: Applications of Light with Orbital Angular Momentum*, Bristol, Wiley (2011).
 - [15] D. Andrews M. Babiker, *The Angular Momentum of Light*, Cambridge University Press, Cambridge (2012).
 - [16] J. Wang et al, Terabit free-space data transmission employing orbital angular momentum multiplexing, Nat. Photonics **6**, 488 (2012).
 - [17] N. Bozinovic et al, Terabit-scale orbital angular momentum mode division multiplexing in fibers, Science **340**, 1545 (2013).
 - [18] S. Groblacher, T. Jennewein, A. Vaziri, G. Weihs, A. Zeilinger, Experimental quantum cryptography with qutrits, New J. Phys. **8**, 75 (2006).
 - [19] M. Mirhosseini, O. Magaña-Loaiza, M. O'Sullivan, B. Rodenburg, M. Malik, D. Gauthier, R. Boyd, High-dimensional quantum cryptography with twisted light, New J. Phys **17**, 033033 (2015).
 - [20] N. Langford et al., Measuring entangled qutrits and their use for quantum bit commitment, Phys. Rev. Lett. **93**, 053601 (2004).
 - [21] G. Molina-Terriza et al., Triggered qutrits for quantum communication protocols, Phys. Rev. Lett. **92**, 167903 (2004).
 - [22] S. Gröblacher et al., Experimental quantum cryptography with qutrits, New J. Phys. **8**, **75** (2006).
 - [23] N. Cerf et al, Security of quantum key distribution using d-level systems, Phys. Rev. Lett. **88**, 127902 (2002).
 - [24] M. Fujiwara, M. Takeoka, J. Mizuno, M. Sasaki, Exceeding the classical capacity limit in a quantum optical channel, Phys. Rev. Lett. **90**, 167906 (2003).
 - [25] T. C. Ralph, K. J. Resch, and A. Gilchrist, Efficient Toffoli gates using qudits, Phys Rev A **75**, 022313 (2007).
 - [26] B. Lanyon, T. Weinhold, N. Langford, J. O'Brien, K. Resch, A. Gilchrist, A. White, Manipulating biphotonic qutrits, Phys. Rev. Lett. **100**, 060504 (2008).
 - [27] X. Xiao and Y. Li, Protecting qutrit-qutrit entanglement by weak measurement and reversal, Eur. J. Phys. D **67**, 204 (2013).
 - [28] X. Xiao, Protecting qubit-qutrit entanglement from amplitude damping decoherence via weak measurement and reversal, Phys. Scr. **89**, 6 (2014).
 - [29] M. Malik, M. Erhard, M. Huber, M. Krenn, R. Fickler, A. Zeilinger, Multi-photon entanglement in high dimensions, preprint arXiv:quant-ph/1509.02561 (2015).
 - [30] P. Facchi, G. Florio, G. Parisi, and S. Pascazio, Maximally multipartite entangled states, Phys. Rev. A **77**, 060304 (2008).
 - [31] W. Helwig, W. Cui, J. I. Latorre, A. Riera, and H.-K. Lo, Absolute maximal entanglement and quantum secret sharing, Phys. Rev. A **86**, 052335 (2012).
 - [32] J. I. Latorre and G. Sierra, Holographic codes, preprint arXiv:quant-ph/1502.06618 (2015).
 - [33] F. Pastawski, B. Yoshida, D. Harlow, J. Preskill, Holographic quantum error-correcting codes: Toy models for the bulk/boundary correspondence, JHEP **06**, 149 (2015).
 - [34] Z. Yanga, P. Hayden and X.-L. Qia, Bidirectional holographic codes and sub-AdS locality, JHEP **175**, 1601 (2016).
 - [35] W. Helwig, Absolutely Maximally Entangled Qudit Graph States, preprint arXiv:quant-ph/1306.2879 (2013).
 - [36] D. Goyeneche, K. Życzkowski, Genuinely multipartite entangled states and orthogonal arrays, Phys. Rev. A **90**, 022316 (2014).
 - [37] D. Goyeneche, D. Alsina, J. I. Latorre, A. Riera, K. Życzkowski, Absolutely Maximally Entangled states, combinatorial designs and multi-unitary matrices, Phys. Rev. A **92**, 032316 (2015).
 - [38] K. Feng, L. Jin, C. Xing, C. Yuan, Multipartite entangled states, symmetric matrices and error-correcting codes, preprint arXiv:quant-ph/1511.07992 (2015).
 - [39] A.S. Hedayat, N.J.A. Sloane, J. Stufken, *Orthogonal Arrays: Theory and Applications*, Springer-Verlag, New York (1999).
 - [40] Z. Wang, S. Yu, H. Fan, C.H. Oh, Quantum Error-Correcting Codes over Mixed Alphabets, preprint arXiv:quant-ph/1205.4253 (2012).
 - [41] R. Roy. *Design of Experiments Using The Taguchi Approach: 16 Steps to Product and Process Improvement*, Wiley, USA, 1st edition (2001).
 - [42] C. Rao, Some combinatorial problems of arrays and applications to design of experiments, Pag. 349-359 of: J. Srivastava, *A Survey of Combinatorial Theory*, Amsterdam, North-Holland (1973).
 - [43] A. Hedayat, K. Pu, J. Stufken, On the construction of asymmetrical orthogonal arrays, Ann. Statist. **20**, 4, 2142-2152 (1992).
 - [44] R. Dicke, Coherence in spontaneous radiation processes, Phys. Rev. **93**, 99 (1954).
 - [45] B. Briegel, R. Raussendorf, Persistent entanglement in arrays of interacting particles, Phys. Rev. Lett. **86**, 910 (2001).
 - [46] M. Hein, J. Eisert, H.J. Briegel, Multi-party entanglement in graph states, Phys. Rev. A **69**, 062311 (2004).
 - [47] <http://support.sas.com/techsup/technote/ts723.html>
 - [48] R. Bose, On the construction of balanced incomplete block designs, Annals of Eugenics **9**, 4, 353-399 (1939).

- [49] C. J. Colbourn, J. H. Dinitz (Eds.) *Handbook of Combinatorial Designs*, 2nd Edition, CRC Press, Boca Raton FL (2006).
- [50] W. Helwig, W. Cui, Absolutely maximally entangled states: existence and applications, preprint arXiv:quant-ph/1306.2536 (2013).
- [51] T. Yu and J. H. Eberly, Evolution from entanglement to decoherence, *Quant. Inf. Comp.* **7**, 459 (2007).
- [52] http://support.sas.com/techsup/technote/ts723_DifferenceSchemes