MULTICLASS CLASSIFICATION, INFORMATION, DIVERGENCE, AND SURROGATE RISK

By John Duchi^{*}, Khashayar Khosravi, and Feng Ruan^{*}

Stanford University

We provide a unifying view of statistical information measures, multi-way Bayesian hypothesis testing, loss functions for multi-class classification problems, and multi-distribution f-divergences, elaborating equivalence results between all of these objects, and extending existing results for binary outcome spaces to more general ones. We consider a generalization of f-divergences to multiple distributions, and we provide a constructive equivalence between divergences, statistical information (in the sense of DeGroot), and losses for multiclass classification. A major application of our results is in multi-class classification problems in which we must both infer a discriminant function γ —for making predictions on a label Y from datum Xand a data representation (or, in the setting of a hypothesis testing problem, an experimental design), represented as a quantizer q from a family of possible quantizers Q. In this setting, we characterize the equivalence between loss functions, meaning that optimizing either of two losses yields an optimal discriminant and quantizer q, complementing and extending earlier results of Nguyen et al. [23] to the multiclass case. Our results provide a more substantial basis than standard classification calibration results for comparing different losses: we describe the convex losses that are consistent for jointly choosing a data representation and minimizing the (weighted) probability of error in multiclass classification problems.

1. Introduction. Consider the multiclass classification problem: a decision maker receives a pair of random variables $(X, Y) \in \mathcal{X} \times \{1, \ldots, k\}$, where Y is unobserved, and wishes to assign the variable X to one of the k classes $\{1, 2, \ldots, k\}$ to minimize the probability of a misclassification. We represent the decision maker via a discriminant function $\gamma : \mathcal{X} \to \mathbb{R}^k$, where each component $\gamma_y(x)$, $y = 1, \ldots, k$, represents the margin (or a score or perceived likelihood) the decision maker assigns to class y for datum x. The goal is then to minimize the expected loss, or L-risk,

(1) $R_L(\gamma) := \mathbb{E}[L(\gamma(X), Y)],$

^{*}Partially supported by NSF-CAREER Award 1553086 and the SAIL-Toyota Center for AI Research. FR additionally supported by the Stanford Graduate Fellowship.

Keywords and phrases: f-divergence, risk, surrogate loss function, hypothesis test

where $L(\gamma(x), y)$ measures the loss of margins $\gamma(x) \in \mathbb{R}^k$ when the true label of x is y and the expectation (1) is taken jointly over (X, Y). When L is the 0-1 loss, $L(\gamma(x), y) = 1 \{\gamma_y(x) \leq \gamma_i(x) \text{ for some } i \neq y\}$, the formulation (1) is the misclassification probability $\mathbb{P}(\operatorname{argmax}_y \gamma_y(X) \neq Y)$. We may also consider the classical k-category Bayesian experiment: given a random variable $X \in \mathcal{X}$ drawn according to one of the k hypotheses

$$H_1: X \sim P_1, \quad H_2: X \sim P_2, \quad \dots, \quad H_k: X \sim P_k$$

with prior probabilities π_1, \ldots, π_k , we wish to choose γ minimizing the expected $\mathbb{E}[L(\gamma(X), Y)]$ or posterior $\sum_y \mathbb{P}(Y = y \mid X = x)L(\gamma(x), y)$ loss.

In many applications, making decisions based on the raw X is undesirable the vector X may be high-dimensional, carry useless information impinging on statistical efficiency, or we may need to store or communicate the sample using limited memory or bandwidth. If all we wish to do is to classify a person as being taller or shorter than 160 centimeters, it makes little sense to track his or her blood type and eye color. With the increase in the number and variety of measurements we collect, such careful design choices are important for maintaining statistical power, interpretability, efficient downstream use, and mitigating false discovery [3]. This desire to give "better" representations of data X has led to a rich body of work in statistics, machine learning, and engineering, highlighting the importance of careful measurement, experimental design, and data representation strategies [28, 11, 26, 32].

As Nguyen et al. [23] note in the binary case, one thus frequently extends the classical formulation (1) to include a stage in which a (data-dependent) $q: \mathcal{X} \to \mathcal{Z}$ maps the vector X into a vector Z. A number of situations suggest such an approach. In most practical classification scenarios [29], an equivalent feature selection reduces the dimension of X or increases its interpretability. As a second motivation, consider the decentralized detection problem [33, 23] in communication applications in engineering, where remote sensors communicate data $X \sim P_i$ through limited bandwidth or memory. In this case, the central processor can infer the distribution P_i only after communication of the transformed vector Z = q(X), and one chooses a quantizer q from a family Q of (low complexity) quantizers. In fuller abstraction, we may treat the problem as a Bayesian experimental design problem, where the mapping $\mathbf{q}: \mathcal{X} \to \mathcal{Z}$ may be stochastic and is chosen from a family Q of possible experiments (observation channels). In each of the preceding examples, the incorporation of a quantizer \mathbf{q} into the classification procedure poses a more complex problem, as one must simultaneously find a data representation q and discriminant γ . The goal, paralleling that for the risk (1),

thus becomes joint minimization of the quantized L-risk

(2)
$$R_L(\gamma \mid \mathbf{q}) := \mathbb{E}\left[L(\gamma(\mathbf{q}(X)), Y)\right]$$

over a prespecified family Q of quantizers $q : \mathcal{X} \to \mathcal{Z}$, where $\gamma : \mathcal{Z} \to \mathbb{R}^k$.

Often—for example, in the zero-one error case—the loss functions $L(\cdot, y)$ are non-convex (even discontinuous), so population or empirical minimization is intractable. It is thus common to replace the loss with a convex surrogate and minimize this surrogate instead. A surrogate is Fisher consistent if its minimization yields a Bayes optimal discriminant γ for the original loss L (for any distribution \mathbb{P} on (X, Y)); researchers have characterized the Fisher consistency of convex surrogates for binary and multiclass classification [36, 21, 2, 31]. A weakness of such results is that they rely strongly on using the unrestricted class of all measurable discriminants $\gamma : \mathcal{X} \to \mathbb{R}^k$, and thus most "natural" convex losses are consistent [36, 2]. In this context, a major difficulty is to understand the consequences of using various surrogate losses, and requiring a restricted quantizer class Q is one approach to discovering nuanced properties of the relationship between surrogate and Bayes risk. Nguyen et al. [23] tackle this in the binary case, considering the problem of joint selection of the discriminant function $\gamma: \mathcal{Z} \to \mathbb{R}$ and quantizer $q: \mathcal{X} \to \mathcal{Z}$. They exhibit a precise correspondence between binary margin-based loss functions and f-divergences—measures of the similarity between two probability distributions developed in information theory and statistics [1, 10, 34]—to give a general characterization of loss equivalence through classes of divergences. An interesting consequence of their results is that, in spite of positive results for Fisher consistency in binary classification problems [36, 21, 2], essentially only hinge-like losses are consistent for the 0-1 loss. We provide the extension of these results to the multiclass case.

Outline and discussion of our contributions. We build on this prior work to provide a unifying framework that relates statistical information measures, loss functions, and generalized notions of entropy in the context of multi-class classification. To begin, we recall a generalization of f-divergences that applies to multiple distributions [16, 13], enumerating analogues of the positivity properties, data-processing inequalities, and discrete approximation available in the binary case, as multi-way f-divergences may be unfamiliar and they motivate our approach (Section 2). We begin our main contributions in Section 3, where we establish a correspondence between loss functions L, generalized entropy on discrete distributions [15], and multiway f-divergences. To make this precise, define the probability simplex $\Delta_k := \{p \in \mathbb{R}^k_+ \mid \mathbf{1}^T p = 1\}$. Let $\pi \in \Delta_k$ be a prior distribution on the

DUCHI, KHOSRAVI, RUAN

class label Y and $\tilde{\pi}(x) \in \Delta_k$ be the posterior probabilities for Y conditional on the observation X = x. For concave $H : \Delta_k \to \mathbb{R}$, DeGroot [11] defines the *information* associated with the experiment (X, Y) as

(3)
$$I_H(X;\pi) := H(\pi) - \mathbb{E}[H(\widetilde{\pi}(X))].$$

The notion of H as a generalized entropy is clear here, as I is the gap between prior and posterior entropy and is always non-negative. In this context, the value $H(\pi)$ measures the *uncertainty* of the experimenter (in some appropriate units) about the unknown class label y when his or her prior belief over y is π , so I is the gap between prior and posterior uncertainty [11].

To relate this type of entropy to loss functions, recall the well-known result [11, 15] that any loss $L : \mathbb{R}^k \times [k] \to \mathbb{R} \cup \{+\infty\}$ induces an entropy $H_L : \Delta_k \to \mathbb{R}$, also called the *pointwise Bayes risk* [15, 27, 13, 35], via

(4)
$$H_L(\pi) = \inf_{\alpha \in \mathbb{R}^k} \left\{ \sum_{i=1}^k \pi_i L(\alpha, i) \right\} - \mathbf{I} \left\{ \pi \in \Delta_k \right\}$$

where $\mathbf{I} \{\cdot\}$ is $+\infty$ if its argument is false (we drop this indicator in the future, defining H_L implicitly on Δ_k). We show an inverse construction, providing an explicit and constructable mapping from any concave function H to a loss L inducing H as the pointwise Bayes risk (4), where for each y the loss $\alpha \mapsto$ $L(\alpha, y)$ is convex. In Section 3.2, we also develop the natural connections between these generalized entropies H and classification calibration [36, 21, 2, 31], in that our explicit L is generally calibrated.

In Section 4, we address the comparison of loss functions—building off of Nguyen et al.'s approach in the binary case [23]—and present our main results on consistency of joint selection of quantizer (data representation) qand discriminant γ . Using our correspondence between concave H, losses L, and f-divergences, we characterize the pairs of losses $L^{(1)}$ and $L^{(2)}$ for which equivalent quantizers and discriminants minimize the quantized risk (2) in the sense that there is a continuous concave h with h(0) = 0 such that

$$R_{L^{(2)}}(\gamma \mid \mathbf{q}) - \inf_{\gamma, \mathbf{q} \in \mathbf{Q}} R_{L^{(2)}}(\gamma \mid \mathbf{q}) \le h\left(R_{L^{(1)}}(\gamma \mid \mathbf{q}) - \inf_{\gamma, \mathbf{q} \in \mathbf{Q}} R_{L^{(1)}}(\gamma \mid \mathbf{q})\right)$$

for any γ and $\mathbf{q} \in \mathbf{Q}$. Another way to understand our results is as providing insight into classification calibration when the Bayes' act (i.e. optimal discriminant γ) does not belong to the class of functions the statistician may choose in a classification problem. A substantial challenge for and criticism of the line of work on classification calibration and surrogate risk consistency [36, 21, 2, 31] is that the results say little for restricted families of classifiers. In this context, a corollary of our main contribution is as follows. The loss $L^{(1)}$ is *calibrated* [36, 2, 31] for $L^{(2)}$ if for any distribution P on $X \times Y$ and sequence $\gamma_n : \mathcal{X} \to \mathbb{R}^k$, $R_{L^{(1)}}(\gamma_n) \to \inf_{\gamma} R_{L^{(2)}}(\gamma)$ implies $R_{L^{(1)}}(\gamma_n) \to \inf_{\gamma} R_{L^{(2)}}(\gamma)$. Now, consider a collection \mathbb{Q} of functions $\mathbf{q} : \mathcal{X} \to \mathcal{Z}$ for some set \mathcal{Z} , and then define the class of functions

$$\mathcal{G}(\mathsf{Q}) := \left\{ \gamma \circ \mathsf{q} \mid \mathsf{q} \in \mathsf{Q} \text{ and } \gamma : \mathcal{Z} \to \mathbb{R}^k \text{ is measurable} \right\}.$$

Translated to this scenario, our main results—Theorems 1 and 2—imply

COROLLARY 1. Assume that the loss $L^{(1)}$ is calibrated for $L^{(2)}$ and let $H_i = H_{L^{(i)}}$ denote the associated pointwise Bayes risk (4). Then

(5)
$$R_{L^{(1)}}(g_n) \to \inf_{g \in \mathcal{G}(\mathbb{Q})} R_{L^{(1)}}(g) \text{ implies } R_{L^{(2)}}(g_n) \to \inf_{g \in \mathcal{G}(\mathbb{Q})} R_{L^{(2)}}(g)$$

for any collection \mathbb{Q} of mappings $\mathcal{X} \to \mathcal{Z}$, any set \mathcal{Z} , any distribution P on $\mathcal{X} \times \{1, \ldots, k\}$, and sequence $g_n \in \mathcal{G}(\mathbb{Q})$ if and only if there exist a > 0, $b \in \mathbb{R}^k$, and $c \in \mathbb{R}$ such that $H_1(\pi) = aH_2(\pi) + b^T \pi + c$ for all $\pi \in \Delta_k$.

This corollary reposes on the connections we develop between losses, uncertainty measures and generalized f-divergences. Such measures of statistical information and divergence have been central to the design of communication and quantization schemes in signal processing [33, 25, 20, 18]; we characterize the divergence measures that, when optimized, yield optimal quantizers and detectors. We also provide a result showing when empirical minimization of a surrogate risk is consistent for the desired (original) risk.

A number of researchers have studied the connections between divergence measures and risk for binary and multi-category experiments; these point to the results we present. Indeed, Blackwell [8] shows that if a quantizer q_1 induces class-conditional distributions with larger divergence than those induced by q_2 , then there are prior probabilities such that q_1 allows tests with lower probability of error than q_2 . Liese and Vajda [19] give a broad treatment of f-divergences, using their representation as the difference between prior and posterior risk in a binary experiment [24] to derive a number of their properties; see also the paper [27]. García-García and Williamson [13] show how multi-distribution f-divergences [16] arise naturally in the context of multi-class classification problems as the gap between prior and posterior risk in classification, as in the work [19]. In the binary case, these results elucidate the characterization of Fisher consistency for quantization and binary classification Nguyen et al. [23] realize. We pursue this line of research to draw the connections between Fisher consistency, information measures, multi-class classification, surrogate losses, and divergences.

Notation. We let **0** and **1** denote the all-zeros and all-ones vectors, respectively. For a vector or collection of objects we define $t_{1:m} = \{t_1, \ldots, t_m\}$. The indicator function $\mathbf{I} \{\cdot\}$ is $+\infty$ if its argument is false, 0 otherwise, while $1 \{\cdot\}$ is 1 if its argument is true, 0 otherwise. We let $\Delta_k = \{v \in \mathbb{R}^k_+ : \mathbf{1}^T v = 1\}$ denote the probability simplex in \mathbb{R}^k . For $m \in \mathbb{N}$, we set $[m] = \{1, \ldots, m\}$. We let aff $A = \{\sum_{i=1}^m \lambda_i x_i \mid \lambda^T \mathbf{1} = 1, x_i \in A, m \in \mathbb{N}\}$ denote the affine hull of a set A, and related the denotes the interior of A relative to aff A. We let $\overline{\mathbb{R}} = \mathbb{R} \cup \{+\infty\}$ and $\underline{\mathbb{R}} = \mathbb{R} \cup \{-\infty\}$. For $f : \mathbb{R}^k \to \overline{\mathbb{R}}$, we let epi $f = \{(x,t) : f(x) \leq t\}$ denote the epigraph of f. We say a convex function f is closed if epi(-f) is closed. For a convex function $f : \mathbb{R}^k \to \overline{\mathbb{R}}$, we say that f is strictly convex at a point $t \in \mathbb{R}^k$ if for all $\lambda \in (0, 1)$ and $t_1, t_2 \neq t$ such that $t = \lambda t_1 + (1 - \lambda)t_2$ we have $f(t) < \lambda f(t_1) + (1 - \lambda)f(t_2)$. The (Fenchel) conjugate of a function $f : \mathbb{R}^k \to \overline{\mathbb{R}}$ is

(6)
$$f^*(s) = \sup_{t \in \mathbb{R}^k} \left\{ s^T t - f(t) \right\}.$$

For any f, the conjugate f^* is closed convex [17, Chapter X]. For measures ν and μ , we let $d\nu/d\mu$ denote the Radon-Nikodym derivative of ν with respect to μ . For random variables X_n , we say $X_n \xrightarrow{L_p} X_\infty$ if $\mathbb{E}[|X_n - X_\infty|^p] \to 0$.

2. Multi-distribution f-divergences. Divergence measures for probability distributions have significant statistical, decision-, and information-theoretic applications, including in optimal testing, minimax rates of convergence, and the design of communication schemes [1, 10, 25, 18]. Central to this work is the f-divergence, introduced by Ali and Silvey [1] and Csiszár [10] (see [19] for an overview). Given distributions P, Q defined on a common set \mathcal{X} , a closed convex function $f : [0, \infty) \to \mathbb{R}$ satisfying f(1) = 0, and any measure μ dominating P and Q, the f-divergence between P and Q is

(7)
$$D_f(P||Q) := \int_{\mathcal{X}} f\left(\frac{p(x)}{q(x)}\right) q(x) d\mu(x) = \int f\left(\frac{dP}{dQ}\right) dQ.$$

Here $p = \frac{dP}{d\mu}$ and $q = \frac{dQ}{d\mu}$ denote the densities of P and Q, respectively, and the value uf(t/u) is defined appropriately for t = 0 and u = 0 (e.g. [19]). A number of classical divergence measures arise out of the f-divergence; taking $f(t) = t \log t$, $f(t) = \frac{1}{2}(\sqrt{t} - 1)^2$, or f(t) = |t - 1| yields (respectively) the KL-divergence, squared Hellinger distance, or total variation distance.

Central to our study of multi-way hypothesis testing and classification is an understanding of relationships between multiple distributions. We use the following generalization [16, 13] of the *f*-divergence to multiple distributions. DEFINITION 2.1. Let P_1, \ldots, P_k be probability distributions on a common σ -algebra \mathcal{F} over a set \mathcal{X} . Let $f : \mathbb{R}^{k-1}_+ \to \mathbb{R}$ be a closed convex function satisfying $f(\mathbf{1}) = 0$. Let μ be any σ -finite measure such that $P_i \ll \mu$ for all i, and let $p_i = dP_i/d\mu$. The f-divergence between P_1, \ldots, P_{k-1} and P_k is

(8)
$$D_f(P_1, \dots, P_{k-1} || P_k) := \int f\left(\frac{p_1(x)}{p_k(x)}, \dots, \frac{p_{k-1}(x)}{p_k(x)}\right) p_k(x) d\mu(x).$$

We must specify the value of the integrand (8) when $p_k(x) = 0$. In this case, the function $\tilde{f} : \mathbb{R}^k_+ \to \overline{\mathbb{R}}$ defined, for an arbitrary $t' \in \text{rel int dom } f$, by

(9)
$$\widetilde{f}(t,u) = \begin{cases} uf(t/u) & \text{if } u > 0\\ \lim_{s \to 0} sf(t'-t+t/s) & \text{if } u = 0\\ +\infty & \text{otherwise} \end{cases}$$

is a closed convex function with value independent of t'; \tilde{f} is the closure of the *perspective* function $\mathbb{R}_+ \times \mathbb{R}^k \ni (u, t) \mapsto uf(t/u)$ of f [17, Prop. IV.2.2.2]. Any time we consider the perspective we implicitly treat it as its closure (9).

We now enumerate a few properties of multi-way f-divergences, showing how they naturally generalize classical binary f-divergences. We focus on basic properties that are useful for our further results on Bayes risk, classification, and hypothesis testing and that parallel results in the binary case (7): they are well-defined, have continuity properties with respect to discrete approximations, and satisfy data-processing inequalities. While Györfi and Nemetz's original work [16] essentially contains the results, we carefully address infinite values (the closure (9)) and strict convexity, and we use them as definitional building blocks; we defer all proofs to Supplement D.

As our first step, we note that Definition 2.1 is independent of the base measure μ . (See Supp. D.1 for a proof generalizing [16, Cor. 1].)

LEMMA 2.1. In expression (8), the value of the divergence does not depend on the choice of the dominating measure μ . Moreover,

$$D_f(P_1,\ldots,P_{k-1}||P_k) \ge 0.$$

The inequality is strict if f is strictly convex at 1 and the P_i are not identical.

Given the importance of quantization to come, we now consider discrete approximations to the divergence. For an at most countable partition \mathcal{P} of \mathcal{X} , we define the partitioned f-divergence

$$D_f(P_1, \dots, P_{k-1} || P_k | \mathcal{P}) = \sum_{A \in \mathcal{P}} f\left(\frac{P_1(A)}{P_k(A)}, \dots, \frac{P_{k-1}(A)}{P_k(A)}\right) P_k(A).$$

DUCHI, KHOSRAVI, RUAN

As in the binary case [34, 19], we have the following approximability result generalizing [16, Thm. 6] to possibly infinite integrands: quantizers give arbitrarily good approximations to f-divergences (see § D.2 for a proof).

PROPOSITION 1. If f is a closed convex function with $f(\mathbf{1}) = 0$, then

$$D_f(P_1, \dots, P_{k-1} || P_k) = \sup_{\mathcal{P}} D_f(P_1, \dots, P_{k-1} || P_k | \mathcal{P})$$

where the supremum is over finite partitions of \mathcal{X} .

In the binary case, f-divergences satisfy data processing inequalities [10, 9, 19] which state that processing or transforming an observation X drawn from the distributions P_1, P_2 , decreases the divergence between them. To formalize this, recall that Q is a Markov kernel from a set \mathcal{X} to \mathcal{Z} if $Q(\cdot | x)$ is a probability distribution on \mathcal{Z} for each $x \in \mathcal{X}$, and for each measurable $A \subset \mathcal{Z}$, the mapping $x \mapsto Q(A | x)$ is measurable. The following general data processing inequality shows that this holds in the multi-distribution case as well, generalizing [16, Thm. 4] to possibly infinite f and the closure (9); we include a proof in Appendix D.3.

PROPOSITION 2. Let f be closed convex with $f(\mathbf{1}) = 0$, Q be a Markov kernel from \mathcal{X} to \mathcal{Z} , and define the marginals $Q_P(A) = \int_{\mathcal{X}} Q(A \mid x) dP(x)$. Then

$$D_f(Q_{P_1},\ldots,Q_{P_{k-1}}||Q_{P_k}) \le D_f(P_1,\ldots,P_{k-1}||P_k).$$

This proposition is related to the relationships between risk, information, and quantization we develop in Sections 3 and 4. Defining a *quantizer* \mathbf{q} to be a measurable mapping $\mathbf{q}: \mathcal{X} \to \mathcal{Z}$ between measurable spaces \mathcal{X} and \mathcal{Z} , the *quantized* f divergence is

$$D_f(P_1, \dots, P_{k-1} \| P_k \mid \mathbf{q}) := \sup_{\mathcal{P}} \sum_{A \in \mathbf{q}^{-1}(\mathcal{P})} f\left(\frac{P_1(A)}{P_k(A)}, \dots, \frac{P_{k-1}(A)}{P_k(A)}\right) P_k(A),$$

where \mathcal{P} ranges over finite partitions of \mathcal{Z} and $q^{-1}(\mathcal{P}) = \{q^{-1}(B) \mid B \in \mathcal{P}\}$. Proposition 2 immediately yields that quantization reduces information: the indicator $Q(A \mid x) = 1 \{q(x) \in A\}$ defines a Markov kernel, yielding

COROLLARY 2. Let f be closed convex, satisfy f(1) = 0, and q be a quantizer of \mathcal{X} . Then

$$D_f(P_1,\ldots,P_{k-1}||P_k||\mathbf{q}) \le D_f(P_1,\ldots,P_{k-1}||P_k).$$

8

We also see that if q_1 and q_2 are quantizers of \mathcal{X} , and q_1 induces a finer partition of \mathcal{X} than q_2 , meaning that for $x, x' \in \mathcal{X}$ the equality $q_1(x) = q_1(x')$ implies $q_2(x) = q_2(x')$, we have

$$D_f(P_1,\ldots,P_{k-1}||P_k||\mathbf{q}_2) \leq D_f(P_1,\ldots,P_{k-1}||P_k||\mathbf{q}_1).$$

This type of ordering is central to this work: any multiclass loss L induces a unique f-divergence, and consistency of discriminants $\gamma : \mathcal{X} \to \mathbb{R}^k$ for a loss L after quantization is intimately tied to the preservation (and relative ordering) of information as related to the quantized risk (2).

3. Risks, information measures, and f-divergences. Having reviewed the basic properties of f-divergences, we turn to a more detailed look at their relationships with multi-way hypothesis tests, multi-class classification, generalized entropies, and statistical informations relating multiple distributions. We build a correspondence between these that parallels that for binary experiments and classification problems [19, 23, 27].

We first recapitulate the probabilistic model for classification and Bayesian hypothesis testing problems from the introduction. We have a prior $\pi \in \Delta_k$ and probability distributions P_1, \ldots, P_k defined on a set \mathcal{X} . The coordinate $Y \in [k]$ is drawn according to a multinomial with probabilities π , and conditional on Y = y, we draw $X \sim P_y$. Following DeGroot [11], we refer to this as an *experiment*. Associated with any experiment is a family of informations as follows. Let $\tilde{\pi}$ be the posterior distribution on Y given observation X = x, $\tilde{\pi}_i(x) = \pi_i dP_i(x)/(\sum_{j=1}^k \pi_j dP_j(x))$. Given any closed concave $H : \mathbb{R}^k_+ \to \mathbb{R}$, which we refer to as *generalized entropy* (see [15, § 3.3]; DeGroot [11] calls H an uncertainty function), the *information* associated with the experiment is the reduction of entropy (uncertainty) from prior to posterior (3),

$$I_H(X;\pi) = H(\pi) - \mathbb{E}[H(\widetilde{\pi}(X))].$$

The expectation is taken over $X \sim \sum_{i=1}^{k} \pi_i P_i$. That $I_H(X; \pi) \ge 0$ is immediate by concavity; DeGroot [11, Thm. 2.1] shows that $I_H(X; \pi) \ge 0$ for all distributions P_1, \ldots, P_k and priors π if and only if H is concave on Δ_k .

In this section we develop equivalence results between multiclass classification losses and risk, multi-way f-divergences, and entropy measures. Concretely, consider $L : \mathbb{R}^k \times [k] \to \mathbb{R}$, and recall the risk (1), defined as $R_L(\gamma) = \mathbb{E}[L(\gamma(X), Y)]$, where $\gamma \in \Gamma$, the set of measurable functions $\gamma : \mathcal{X} \to \mathbb{R}^k$. As in equation (4) in the introduction, each loss L induces the entropy H_L on Δ_k via $H_L(\pi) = \inf_{\alpha \in \mathbb{R}^k} \sum_{i=1}^k \pi_i L(\alpha, i)$, also called the pointwise Bayes risk [15, 13, 27, 35]. In Section 3.1, we give an explicit inverse mapping showing how each generalized entropy H is induced by (at least

DUCHI, KHOSRAVI, RUAN

one) convex loss function L, i.e., $L(\cdot, i)$ is convex for each i. In Section 3.2, we illustrate consistency properties the entropy H implies about the convex loss L inducing it. We connect these results in Section 3.3 with multi-way f-divergences. For any loss L and associated entropy/Bayes risk H_L , for all $\pi \in \Delta_k$ there exists a convex function $f_{L,\pi} : \mathbb{R}^{k-1}_+ \to \mathbb{R}$ with $f_{L,\pi}(1) = 0$ such that the gap between the prior Bayes L-risk—the best expected loss attainable without observing X—and the posterior Bayes risk inf_{γ} $R_L(\gamma)$ is

$$H_L(\pi) - \inf_{\gamma \in \Gamma} R_L(\gamma) = H_L(\pi) - \mathbb{E}[H_L(\widetilde{\pi}(X))] = D_{f_{L,\pi}}(P_1, \dots, P_{k-1} || P_k)$$

(see [15, 13]). The inverse direction is new, and given any closed convex function $f : \mathbb{R}^{k-1}_+ \to \mathbb{R}$ with $f(\mathbf{1}) = 0$, we construct convex losses $L(\cdot, i)$, an associated generalized entropy H_L , and prior $\pi = \mathbf{1}/k \in \Delta_k$ satisfying

$$D_f(P_1,\ldots,P_{k-1}||P_k) = \inf_{\alpha \in \mathbb{R}^k} \sum_{i=1}^k \pi_i L(\alpha,i) - \inf_{\gamma \in \Gamma} R_L(\gamma).$$

3.1. Generalized entropies and losses. We construct a natural bidirectional mapping between losses and generalized entropies, giving a few examples to illustrate. For any loss $L : \mathbb{R}^k \times [k] \to \mathbb{R}$, the construction (4) of H_L yields a closed concave function, as H_L is the infimum of linear functionals of π . It is thus a generalized entropy [15] (or uncertainty function [11]), and the gap $H_L(\pi) - \mathbb{E}[H_L(\tilde{\pi}(X))]$ between prior and posterior entropy is non-negative. The following two examples with zero-one loss are illustrative.

EXAMPLE 1 (Zero-one loss). Consider the zero one loss

$$L^{zo}(\alpha, y) = 1 \{ \alpha_y \le \alpha_j \text{ for some } j \ne y \}$$

where $y \in [k]$. Then we have

$$H_{L^{zo}}(\pi) = \inf_{\alpha} \left\{ \sum_{i=1}^{k} \pi_i \mathbb{1} \left\{ \alpha_i \le \alpha_j \text{ for some } j \ne i \right\} \right\} = 1 - \max_j \pi_j.$$

This generalized entropy is concave, nonnegative, and satisfies $H_{L^{zo}}(\pi) = 0$ if and only if $\pi = e_i$ for a standard basis vector e_i .

EXAMPLE 2 (Cost-weighted classification). In some scenarios, we allow different costs for classifying certain classes y as others; for example, it may be less costly to misclassify a benign tumor as cancerous than the opposite. In this case, we use a matrix $C = [c_{yi}]_{y,i=1}^k \in \mathbb{R}^{k \times k}_+$, where $c_{yi} \ge 0$ is the cost for classifying an observation of class y as class i (i.e. assigning $X \sim P_i$ instead of P_y in the experiment). We assume $c_{yy} = 0$ for each y and define

(10)
$$L^{cw}(\alpha, y) = \max_{i} \{ c_{yi} \mid \alpha_i = \max_{j} \alpha_j \}, \quad \alpha \in \mathbb{R}^k$$

the maximal loss for those indices of α attaining $\max_j \alpha_j$. Let $C = [c_1 \cdots c_k]$ be the column representation of C. If $c_y^T \pi = \min_l c_l^T \pi$, then by choosing any α such that $\alpha_y > \alpha_j$ for all $j \neq y$, we have

$$H_{L^{\mathrm{cw}}}(\pi) = \inf_{\alpha} \left\{ \sum_{y=1}^{k} \pi_y \max_i \{ c_{yi} \mid \alpha_i = \max_j \alpha_j \} \right\} = \min_l \pi^T c_l.$$

The entropy $H_{L^{cw}}$ is concave, nonnegative, and satisfies $H_{L^{cw}}(e_i) = 0$ for standard basis vectors e_i ; Example 1 corresponds to $C = \mathbf{1}\mathbf{1}^T - I_{k \times k}$.

The forward mapping (4) from losses L to entropy H_L is straightforward, though it is many-to-one. Using convex duality and conjugacy arguments, we can show an inverse mapping. This construction is new, though precursors for proper scoring rules and predictions in the probability simplex exist [15, 14, Thm. 2]; these *characterize* proper scoring rules, but it is not always clear how to generate *convex* losses from these. Before stating the proposition, we recall the definition (6) of the Fenchel conjugate $f^*(s) = \sup_t \{s^T t - f(t)\}$.

PROPOSITION 3. For any closed concave $H : \Delta_k \to \mathbb{R}$, the losses

(11)
$$L(\cdot,i): \mathbb{R}^k \to \overline{\mathbb{R}}, \quad L(\alpha,i) = -\alpha_i + (-H)^*(\alpha),$$

 $i \in \{1, \ldots, k\}$, are closed, convex, and satisfy the equality (4) that $H \equiv H_L$.

PROOF. Standard Fenchel conjugacy relationships [17, Chapter X] imply

$$H(\pi) = \inf_{\alpha \in \mathbb{R}^k} \left\{ -\pi^T \alpha + (-H)^*(\alpha) \right\} \text{ where } (-H)^*(\alpha) = \sup_{\pi \in \Delta_k} \left\{ \alpha^T \pi + H(\pi) \right\}$$

Defining $L(\alpha, i) = -\alpha_i + (-H)^*(\alpha)$ for i = 1, ..., k, we can write

$$H(\pi) = \inf_{\alpha \in \mathbb{R}^k} \left\{ -\pi^T \alpha + (-H)^*(\alpha) \right\} = \inf_{\alpha \in \mathbb{R}^k} \left\{ -\pi^T \alpha + \pi^T \mathbf{1} \cdot (-H)^*(\alpha) \right\}$$
$$= \inf_{\alpha \in \mathbb{R}^k} \left\{ \sum_{i=1}^k \pi_i L(\alpha, i) \right\}.$$

Proposition 3 shows that associated with every concave entropy defined on the simplex, there is at least one set of *convex* loss functions $L(\cdot, i)$ generating

DUCHI, KHOSRAVI, RUAN

the entropy via the infimal representation (4), and there is thus a mapping from loss functions to entropies and from entropies to *convex* losses: given any loss L, we may construct a convex loss L^{cvx} with $H_L = H_{L^{\text{cvx}}}$. The mapping from entropies H to loss functions generating H as in (4) is one-tomany, as any losses $L^{(1)}$ and $L^{(2)}$ with the same range satisfy $H_{L^{(1)}} = H_{L^{(2)}}$.

3.2. Surrogate risk consistency and generalized entropies. Our construction (11) of loss functions is a somewhat privileged construction, as it often yields desirable properties of the convex loss function itself, especially as related to the non-convex zero-one loss. Indeed, it is often the case that the convex loss L so generated is Fisher consistent; to make this explicit, we recall the following definition [36, 31].

DEFINITION 3.1. Let $L : \mathbb{R}^k \times [k] \to \overline{\mathbb{R}}$. Then L is classification calibrated for the zero-one loss if for any $\pi \in \Delta_k$ and i^* such that $\pi_{i^*} < \max_i \pi_i$,

(12)
$$\inf_{\alpha \in \mathbb{R}^k} \left\{ \sum_{i=1}^k \pi_i L(\alpha, i) \right\} < \inf_{\alpha \in \mathbb{R}^k} \left\{ \sum_{i=1}^k \pi_i L(\alpha, i) : \alpha_{i^*} \ge \max_j \alpha_j \right\}$$

Given a matrix $C \in \mathbb{R}^{k \times k}_+$ as in Example 2, L is classification calibrated for the cost matrix C if for any $\pi \in \Delta_k$ and any i^* with $c_{i^*}^T \pi > \min_j c_j^T \pi$,

(13)
$$\inf_{\alpha \in \mathbb{R}^k} \left\{ \sum_{i=1}^k \pi_i L(\alpha, i) \right\} < \inf_{\alpha \in \mathbb{R}^k} \left\{ \sum_{i=1}^k \pi_i L(\alpha, i) : \alpha_{i^*} \ge \max_j \alpha_j \right\}$$

Tewari and Bartlett [31, Thm. 2] and Zhang [36, Thm. 3] show the importance of Definition 3.1: let $R(\gamma)$ be the zero-one or cost-weighted risk (Exs. 1–2). If L is lower-bounded, then it is classification calibrated (with respect to zero-one or the cost-weighted loss) if and only if for any sequence $\gamma_n : \mathcal{X} \to \mathbb{R}^k$ and distribution \mathbb{P} on $X \times Y$ we have *Fisher consistency*, i.e.

$$R_L(\gamma_n) \to \inf_{\gamma \in \Gamma} R_L(\gamma) \text{ implies } R(\gamma_n) \to \inf_{\gamma \in \Gamma} R(\gamma).$$

That is, classification calibration (with respect to zero-one-risk or the costweighted risk) is equivalent to surrogate risk consistency of the loss L. Because of the predominance of the zero-one loss in the literature, we use "classification calibration" without any qualification to mean "classification calibration with respect to zero-one loss".

We now show how—under minor restrictions on the generalized entropy function H—the construction (11) yields classification calibrated losses.

DEFINITION 3.2. A convex function $f : \mathbb{R}^k \to \overline{\mathbb{R}}$ is $(\lambda, \kappa, \|\cdot\|)$ -uniformly convex over $C \subset \mathbb{R}^k$ if it is closed and for all $t \in [0, 1]$ and $x_1, x_2 \in C$

$$f(tx_1 + (1-t)x_2) \le tf(x_1) + (1-t)f(x_2) - \frac{\lambda}{2}t(1-t) ||x_1 - x_2||^{\kappa} \left[(1-t)^{\kappa-1} + t^{\kappa-1} \right].$$

We say, without qualification, that f is uniformly convex on C if dom $f \supset C$ and there exist $\lambda > 0$, a norm $\|\cdot\|$, and constant $\kappa < \infty$ such that Definition 3.2 holds; we say f is uniformly concave if -f is uniformly convex. Definition 3.2 is an extension of the usual notion of *strong convexity*, which holds when $\kappa = 2$, and is essentially a quantified notion of strict convexity.

With this definition, we have the following two propositions. These two propositions, whose proofs we provide in Appendix A, show that generalized entropies naturally give rise to classification calibrated loss functions; we provide examples of these results in Section 3.4 to come.

PROPOSITION 4. Assume that H is closed concave, symmetric, and has dom $H = \Delta_k$, and let L have definition (11). Additionally assume that (a) H is strictly concave, and $\inf_{\alpha} \sum_{i=1}^k \pi_i L(\alpha, i)$ is attained for all $\pi \in \Delta_k$, or (b) H is uniformly concave. Then L is classification calibrated.

Even when H is not strictly concave, we can give classification calibration results. Indeed, recall Example 1, which showed that for the zero-one-loss, we have $H_L(\pi) = 1 - \max_i \pi_i$.

PROPOSITION 5. Let $H(\pi) = 1 - \max_j \pi_j$. The loss (11) defined by $L(\alpha, i) = -\alpha_i + (-H)^*(\alpha)$ is classification calibrated. Moreover, we have for any $\pi \in \Delta_k$ and $\alpha \in \mathbb{R}^k$ that

$$\sum_{i=1}^{k} \pi_i L(\alpha, i) - \inf_{\alpha} \sum_{i=1}^{k} \pi_i L(\alpha, i) \ge \frac{1}{k} \bigg(\sum_{i=1}^{k} \pi_i L^{\mathrm{zo}}(\alpha, i) - \inf_{\alpha} \sum_{i=1}^{k} \pi_i L^{\mathrm{zo}}(\alpha, i) \bigg).$$

3.3. Divergences, risk, and generalized entropies. In this section, we show that f-divergences as in Definition 2.1 have a precise correspondence with generalized entropies and losses; García-García and Williamson [13] establish the correspondence between f-divergences and entropy/pointwise Bayes risk H; our results show the important link from f directly to the loss L. We begin as in equation (4) with a concave generalized entropy H and loss L satisfying $H(\pi) = \inf_{\alpha \in \mathbb{R}^k} \sum_{i=1}^k \pi_i L(\alpha, i)$; by Proposition 3 it is no loss of generality to assume this correspondence. Let Γ be the collection of measurable functions $\gamma : \mathcal{X} \to \mathbb{R}^k$. The posterior Bayes risk for L is

(14)
$$H_L(\pi, P_{1:k}) := \inf_{\gamma \in \Gamma} \int_{\mathcal{X}} \sum_{i=1}^k \pi_i L(\gamma(x), i) dP_i(x) = \mathbb{E}[H_L(\widetilde{\pi}(X))],$$

where $\tilde{\pi}(x)$ is the posterior distribution on Y conditional on X = x. The information measure (3) is thus the gap between the prior Bayes L-risk and posterior Bayes L-risk. We may then write

$$\inf_{\alpha \in \mathbb{R}^k} \sum_{i=1}^k \pi_i L(\alpha, i) - \inf_{\gamma \in \Gamma} R_L(\gamma) = H_L(\pi) - H_L(\pi, P_{1:k}) = I_{H_L}(X; \pi)$$
$$= \int_{\mathcal{X}} \sup_{\alpha} \left(H_L(\pi) - \sum_{i=1}^{k-1} \pi_i L(\alpha, i) \frac{dP_i}{dP_k} - \pi_k L(\alpha, k) \right) dP_k = D_{f_{L,\pi}}(P_{1:k-1} \| P_k) + C_{f_{L,\pi}}(P_{1:k-1} \| P_k)$$

where the closed convex function $f_{L,\pi}: \mathbb{R}^{k-1}_+ \to \overline{\mathbb{R}}$ has definition

(15)
$$f_{L,\pi}(t) := \sup_{\alpha \in \mathbb{R}^k} \left(H_L(\pi) - \sum_{i=1}^{k-1} \pi_i L(\alpha, i) t_i - \pi_k L(\alpha, k) \right).$$

As $f_{L,\pi}$ is the supremum of affine functions of its argument t, it is closed convex, and $f_{L,\pi}(\mathbf{1}) = H_L(\pi) - H_L(\pi) = 0$. That is, equation (15) shows that given any loss L or generalized entropy H, the information measure $I_{H_L}(X;\pi)$, gap between prior and posterior L-risk, and $f_{L,\pi}$ -divergence between distributions P_1, \ldots, P_{k-1} and P_k are identical.

We can also give a converse result that shows that every f-divergence can be written as the gap between prior and posterior risks for a convex loss function. We first recall the result that $D_f(P_{1:k-1}||P_k)$ is a statistical information (3) based on an generalized entropy H associated with f. Except for the closure operation, this result is known [13, Thm. 3].

PROPOSITION 6. For closed convex $f : \mathbb{R}^{k-1} \to \overline{\mathbb{R}}$ with $f(\mathbf{1}) = 0$, let

$$H(t_1,\ldots,t_k) = -kt_k f\left(\frac{t_1}{t_k},\ldots,\frac{t_{k-1}}{t_k}\right),$$

where we implicitly use the closure of the perspective (Def. (9)). Then

$$D_f(P_1,\ldots,P_{k-1}||P_k) = H(\mathbf{1}/k) - \mathbb{E}[H(\widetilde{\pi}(X))]$$

where the prior $\pi = 1/k$ and the expectation is taken according to $\sum_i \pi_i P_i$.

By combining Propositions 3 and 6 with the infimal representation (4) of H_L , we immediately obtain the following corollary, which is our explicit construction of a closed convex loss from an f-divergence.

COROLLARY 3. Let $\pi^0 = \mathbf{1}/k$. For any closed and convex function $f : \mathbb{R}^{k-1} \to \overline{\mathbb{R}}$ such that $f(\mathbf{1}) = 0$, the convex losses defined by

$$L(\alpha, i) = -\alpha_i + \sup_{\pi \in \Delta_k} \left\{ \pi^T \alpha - k\pi_k f\left(\frac{\pi_1}{\pi_k}, \dots, \frac{\pi_{k-1}}{\pi_k}\right) \right\}$$

satisfy Eq. (15), i.e. $f(t) = \sup_{\alpha} \{ H_L(\pi^0) - \sum_{i=1}^{k-1} \pi_i^0 L(\alpha, i) t_i - \pi_k^0 L(\alpha, k) \}.$ Additionally,

$$D_f(P_{1:k-1} \| P_k) = \inf_{\alpha \in \mathbb{R}^k} \sum_{i=1}^k \pi_i^0 L(\alpha, i) - \inf_{\gamma} \mathbb{E}[L(\gamma(X), Y)],$$

where the expectation is over $Y \sim \pi^0$ and $X \sim P_y$ conditional on Y = y.

For binary classification problems, Nguyen et al. [23, Thm. 1] provide an explicit construction of a closed convex margin-based loss inducing the f-divergence as in Eq. (15); the binary case allows a complete characterization of all such convex functions, which appears difficult in the multiclass case.

Corollary 3, coupled with the information representation given by the fdivergence (15), shows the complete equivalence between f-divergences, loss functions L, and entropies H. For any f-divergence, there exists a loss function L and prior $\pi = 1/k$ such that $D_f(P_{1:k-1}||P_k) = H_L(\pi) - H_L(\pi, P_{1:k})$. Conversely, for any loss function L and prior π , there exists a multi-way f-divergence such that the gap $H_L(\pi) - H_L(\pi, P_{1:k}) = D_f(P_{1:k-1}||P_k)$.

3.4. Examples of generalized entropies and loss correspondences. To complement our general results, we illustrate the correspondence between (concave) generalized entropies and the loss construction (11) through several examples, using Propositions 4 and 5 to guarantee classification calibration.

EXAMPLE 3 (Zero-one loss, Example 1, continued). We use the generalized entropy $H(\pi) = 1 - \max_j \pi_j$ generated by the zero-one loss to derive a convex loss function L that gives the same entropy via the representation (4). The conjugate to -H is

(16)
$$(-H)^*(\alpha) = 1 + \max\left\{\alpha_{(1)} - 1, \frac{\alpha_{(1)} + \alpha_{(2)}}{2} - \frac{1}{2}, \dots, \frac{\sum_{i=1}^k \alpha_{(i)}}{k} - \frac{1}{k}\right\},\$$

where $\alpha_{(1)} \ge \alpha_{(2)} \ge \cdots$ are the entries of $\alpha \in \mathbb{R}^k$ in sorted order (see § A.5 for a proof). Then the convex "family-wise" loss, named for its similarity to family-wise error control in hypothesis tests,

$$L^{\text{fw}}(\alpha, i) = 1 - \alpha_i + \max_{l \in \{1, \dots, k\}} \left\{ \frac{1}{l} \sum_{j=1}^{l} \alpha_{(j)} - \frac{1}{l} \right\}$$

generates the same entropy $H_{L^{\text{fw}}}$ and associated f-divergence as the zeroone loss. Moreover, Proposition 5 guarantees that L^{fw} is classification calibrated (Def. 3.1). It appears that the loss L^{fw} is a new convex classificationcalibrated loss function. \diamond

Rather than re-considering Example 2, which we do later in the context of showing that distinct convex losses can yield the same generalized entropy, we now consider the multiclass logistic loss. The loss does *not* correspond to the zero-one loss, but it generates Shannon entropy and information.

EXAMPLE 4 (Logistic loss and entropy). For $1 \le i \le k$, define $p_i(\alpha) = e^{\alpha_i} / \sum_{j=1}^k e^{\alpha_j}$. The multi-class logistic loss is then

$$L(\alpha, i) = -\log p_i(\alpha) = \log \left(\sum_{j=1}^k e^{\alpha_j - \alpha_i}\right) \text{ for } 1 \le i \le k.$$

The entropy associated with the loss is the familiar Shannon entropy,

(17)
$$H_L(\pi) = \inf_{\alpha \in \mathbb{R}^k} \left\{ -\sum_{i=1}^k \pi_i \log p_i(\alpha) \right\} = -\sum_{i=1}^k \pi_i \log \pi_i.$$

The conjugacy calculation (11) (i.e. our inverse construction from H to loss L) to generate L also yields the multi-class logistic loss. That the multiclass logistic loss is calibrated for the zero-one loss [36, § 4.4] is now immediate: the negative Shannon entropy is strongly convex over the simplex Δ_k (this is Pinsker's inequality [9, Ch. 17.3]), so the fact that logistic loss and Shannon entropy are dual via Eq. (11) and Proposition 4 yield calibration. The information measure (3) associated with the logistic loss is the mutual information between the observation X and label Y. Indeed, we have

$$I_H(X;\pi) = H(\pi) - \mathbb{E}[H(\widetilde{\pi}(X))] = H(Y) - \int_{\mathcal{X}} H(Y \mid X = x) d\overline{P}(x)$$
$$= H(Y) - H(Y \mid X) = I(X;Y)$$

where *H* denotes the Shannon entropy, $\overline{P} = \sum_{i=1}^{k} \pi_i P_i$, and I(X;Y) is the usual (Shannon) mutual information between *X* and *Y*.

We include one final example to show that in some instances, many different *convex* losses can yield the same generalized entropy H.

EXAMPLE 5 (Hinge losses). Define the pairwise multiclass hinge loss

$$L^{\text{hin}}(\alpha, i) = \sum_{j \neq i} \left[1 + \alpha_j \right]_+ + \mathbf{I} \left\{ \mathbf{1}^T \alpha = 0 \right\}.$$

We also consider the slight extension to weighted loss functions to address asymmetric losses of the form (10) from Example 2. In this case, given the loss matrix $C \in \mathbb{R}^{k \times k}_+$, we set

$$L^{\text{hin}}(\alpha, i) = \sum_{j=1}^{k} c_{ij} \left[1 + \alpha_j \right]_{+} + \mathbf{I} \left\{ \mathbf{1}^T \alpha = 0 \right\}.$$

The loss $L(\alpha, i) = \sum_{j \neq i} c_{ij} [1 + \alpha_j - \alpha_i]_+$ yields a completely identical set of calculations without the restriction $\mathbf{1}^T \alpha = 0$, as it is invariant to shifts. A calculation (see § A.6 for completeness) shows the generalized entropy (4) associated with the hinge loss with loss matrix $C = [c_1 \cdots c_k]$ is

(18)
$$H_{L^{\min}}(\pi) = \inf_{\alpha \in \mathbb{R}^k} \left\{ \sum_{i=1}^k \pi_i L^{\min}(\alpha, i) \right\} = k \min_l \pi^T c_l.$$

Such losses satisfy a number of classification calibration guarantees; we note one, essentially due to Zhang [36, Thm. 8]. For completeness, we include a proof in Appendix A.4.

OBSERVATION 1. Let $\phi : \mathbb{R} \to \mathbb{R}$ be any bounded below convex function, differentiable on $(-\infty, 0]$, with $\phi'(0) < 0$. Then $L(\alpha, y) = \sum_{i=1}^{k} c_{yi}\phi(-\alpha_i)$ is classification calibrated for the cost matrix C (Def. 3.1, Eq. (13)).

Taking $C = \mathbf{1}\mathbf{1}^T - I_{k \times k}$, we see that the hinge loss is calibrated for the zero-one loss (Ex. 1); taking arbitrary $C \in \mathbb{R}^{k \times k}_+$, the weighted hinge loss is calibrated for the cost matrix C. Even more, we have the following quantitative calibration guarantee in analogy with Proposition 5:

$$\sum_{i=1}^k \pi_i L^{\mathrm{hin}}(\alpha, i) - \inf_{\alpha'} \sum_{i=1}^k \pi_i L^{\mathrm{hin}}(\alpha', i) \ge \sum_{i=1}^k \pi_i L^{\mathrm{cw}}(\alpha, i) - \inf_{\alpha'} \sum_{i=1}^k \pi_i L^{\mathrm{cw}}(\alpha', i)$$

for all $\pi \in \Delta_k$ and $\alpha \in \mathbb{R}^k$, strengthening Observation 1. (We prove this as Lemma A.9 in § A.7.) In the binary case [23, Lem. 2], similar quantitative guarantees hold for *any* margin-based classification calibrated loss L for which $H_L = H_{L^{zo}}$; we do not know if this extends to the multiclass case. \diamond

DUCHI, KHOSRAVI, RUAN

4. Comparison of loss functions. In Section 3, we demonstrated the correspondence between loss functions, generalized entropies, statistical information, f-divergences, and (in restricted cases) classification calibration. These correspondences assume that decision makers have access to the entire observation X, which is often not the case; as noted in the introduction, it is often beneficial to pre-process data to make it lower dimensional, communicate or store it efficiently, or to improve statistical behavior. Thus, we now explore the impact quantization has on these concepts.

To motivate this further, consider that each of the family-wise loss L^{fw} of Ex. 3, logistic loss (Ex. 4), and any loss of the form $L(\alpha, y) = \sum_{i \neq y} \phi(-\alpha_i)$ for ϕ convex and decreasing with $\phi'(0) < 0$ (Ex. 5, Observation 1) is classification calibrated. This relates to one of the major criticisms of classification calibration: if the Bayes classifier (minimizer of risk over all functions $\mathcal{X} \to \mathbb{R}^k$) does not belong to the class of functions considered, classification calibration says little. In this context, we shed light on this issue by identifying losses that are consistent (calibrated) even with the additional selection of quantizer or data representation—a restriction of the class of possible functions as in the implication (5) in the introduction.

4.1. A model of quantization and experimental design. Abstractly, we treat the design of an experiment or choice of data representation as a quantization problem, where a quantizer \mathbf{q} maps the space \mathcal{X} to a measurable space \mathcal{Z} . Then, for a loss L, prior $\pi \in \Delta_k$ on the label Y, and discriminant $\gamma: \mathcal{Z} \to \mathbb{R}^k$, we consider the quantized risk (2), which we recall is

$$R_{L,\pi}(\gamma \mid \mathsf{q}) := \mathbb{E}\left[L(\gamma(\mathsf{q}(X)), Y)\right].$$

Given class-conditional distributions $P_{1:k}$ (equivalently, hypotheses $H_i : P_i$ in the Bayesian testing setting) and collection **Q** of quantizers, our criterion is to choose the quantizer **q** that allows the best attainable risk. That is, we consider the quantized Bayes *L*-risk, defined as the infimum of the risk (2) over discriminants $\Gamma = \{\gamma : \mathbb{Z} \to \mathbb{R}^k\},\$

(19)
$$\inf_{\gamma \in \Gamma} R_{L,\pi}(\gamma \mid \mathbf{q}) = \int_{\mathcal{Z}} \inf_{\alpha} \sum_{i=1}^{k} \pi_i L(\alpha, i) dP_i^{\mathbf{q}}(z)$$

where $P^{q}(A) = P(q^{-1}(A))$ denotes the push-forward measure. The risk (19) measures the best attainable risk for a fixed choice of $q \in Q$; one thus seeks the design q giving the lowest quantized Bayes *L*-risk.

Whether for computational or analytic reasons, minimizing the loss (19) is often intractable; the zero-one loss L^{zo} (Ex. 1), for example, is non-

convex and discontinuous. It is thus of interest to understand the asymptotic consequences of using a surrogate loss L in place of the desired loss (say L^{zo}) [36, 21, 2, 31], including the setting in which one incorporates further dimension reduction via the choice $\mathbf{q} \in \mathbf{Q}$. Nguyen et al. [23] introduce and study this problem for binary classification, giving a correspondence between f-divergences, loss functions, and surrogate consistency with quantization. The consequences of using a surrogate for consistency of the resulting quantization and classification procedure in the multiclass case are *a-priori* unclear: we do not know when using such a surrogate can be done without penalty. To that end, we now characterize when two loss functions $L^{(1)}$ and $L^{(2)}$ provide equivalent criteria for choosing quantizers (experimental designs or data representations) according to the Bayes *L*-risk (19).

4.2. Universal Equivalence of Loss Functions. Recalling our arguments in Section 3.3 that statistical information (the gap between prior and posterior risks) is a multi-way f-divergence between distributions P_1, \ldots, P_{k-1} and P_k , we give a quantized version of this construction. In analogy with the results of Section 3.3, the quantized statistical information is

$$(20) \quad I_{H_L}(X; \pi \mid \mathbf{q}) := H_L(\pi) - \mathbb{E}[H_L(\widetilde{\pi}(\mathbf{q}(X)))]$$
$$= \inf_{\alpha \in \mathbb{R}^k} \sum_{i=1}^k \pi_i L(\alpha, i) - \inf_{\gamma} R_{L,\pi}(\gamma \mid \mathbf{q}) = D_{f_{L,\pi}}(P_1, \dots, P_{k-1} || P_k \mid \mathbf{q}),$$

where $H_L(\pi) = \inf_{\alpha \in \mathbb{R}^k} \sum_{i=1}^k \pi_i L(\alpha, i)$ as in (4), the convex function $f_{L,\pi}$ is defined as in expression (15) and does not depend on the quantizer \mathbf{q} , and $\tilde{\pi}(\mathbf{q}(X))$ denotes the posterior distribution on $Y \in [k]$ conditional on observing $\mathbf{q}(X)$. We extend Nguyen et al. [23]'s notion of universal equivalence from the binary case, defining losses as equivalent if they induce the same ordering of quantizers \mathbf{q} under the information measure (20).

DEFINITION 4.1. Loss functions $L^{(1)}$ and $L^{(2)}$ are universally equivalent for the prior π , denoted $L^{(1)} \stackrel{u}{\equiv}_{\pi} L^{(2)}$, if for any distributions P_1, \ldots, P_k on X and quantizers \mathbf{q}_1 and \mathbf{q}_2

$$\begin{split} I\left(X,\pi;H_{L^{(1)}} \mid \mathsf{q}_1\right) &\leq I\left(X,\pi;H_{L^{(1)}} \mid \mathsf{q}_2\right) \quad \text{if and only if} \\ I\left(X,\pi;H_{L^{(2)}} \mid \mathsf{q}_1\right) &\leq I\left(X,\pi;H_{L^{(2)}} \mid \mathsf{q}_2\right). \end{split}$$

Definition 4.1 evidently is equivalent to the ordering condition

(21)
$$\inf_{\gamma} R_{L^{(1)},\pi}(\gamma \mid \mathbf{q}_1) \leq \inf_{\gamma} R_{L^{(1)},\pi}(\gamma \mid \mathbf{q}_2) \text{ if and only if} \\ \inf_{\gamma} R_{L^{(2)},\pi}(\gamma \mid \mathbf{q}_1) \leq \inf_{\gamma} R_{L^{(2)},\pi}(\gamma \mid \mathbf{q}_2),$$

DUCHI, KHOSRAVI, RUAN

for all distributions P_1, \ldots, P_k , on the quantized Bayes *L*-risk (19). This definition is somewhat stringent: losses are universally equivalent only if they induce the same quantizer ordering for all population distributions. If a quantizer \mathbf{q}_1 is finer than \mathbf{q}_2 , all losses yield $I(X, \pi; H_L | \mathbf{q}_2) \leq I(X, \pi; H_L | \mathbf{q}_1)$ by the data processing inequality (Corollary 2 of Section 2). The stronger equivalence notion is important for nonparametric classification settings in which the underlying distribution on (X, Y) is only weakly constrained and neither of a pair of quantizers $\mathbf{q}_1, \mathbf{q}_2 \in \mathbf{Q}$ is finer than the other.

Definition 4.1 and the representation (20) suggest that the entropy function H_L associated with the loss L through the infimal representation (4) and the f-divergence associated with L via the construction (15) are important for the equivalence of two loss functions. This is indeed the case. First, we have the following result on universal equivalence of loss functions based on their associated entropies.

THEOREM 1. Let $L^{(1)}$ and $L^{(2)}$ be bounded below losses and $H_{L^{(1)}}$ and $H_{L^{(2)}}$ be the associated generalized entropies as in the construction (4). Then $L^{(1)}$ and $L^{(2)}$ are universally equivalent with respect to all priors π if and only if there exist $a > 0, b \in \mathbb{R}^k$, and $c \in \mathbb{R}$ such that for all $\pi \in \Delta_k$,

$$H_{L^{(1)}}(\pi) = a H_{L^{(2)}}(\pi) + b^T \pi + c.$$

We can also characterize universal equivalence for a prior π .

THEOREM 2. Let $\pi \in \Delta_k$ and as in Theorem 1, let $L^{(1)}$ and $L^{(2)}$ be bounded below loss functions, with $f_{\pi}^{(1)}$ and $f_{\pi}^{(2)}$ the associated f-divergences as in the construction (15). Then $L^{(1)}$ and $L^{(2)}$ are universally equivalent with respect to the prior π if and only if there exist $a > 0, b \in \mathbb{R}^{k-1}$, and $c \in \mathbb{R}$ such that

(22)
$$f_{\pi}^{(1)}(t) = a f_{\pi}^{(2)}(t) + b^T t + c \quad for \ all \ t \in \mathbb{R}^{k-1}_+.$$

Nguyen et al. [23, Thm. 3] prove Theorem 2 for binary classification problems (k = 2), using convex-conjugacy arguments. We outline our proofs (which apply for arbitrary k and so require a different set of tools) in Section 5.

4.3. Consistency of empirical risk minimization. A major application of these theorems is to show that certain non-convex loss functions (such as the zero-one loss) are universally equivalent to convex loss functions, including variants of the hinge loss, by showing that their associated entropies are scalar multiples. As a first application of Theorems 1 and 2, however, we consider the Bayes consistency of empirical risk minimization for selecting a

discriminant γ and quantizer q (in analogy with [23, Thm. 2]). In this case, we receive a sample $\{(X_1, Y_1), \ldots, (X_n, Y_n)\}$ and define the empirical risk

$$\widehat{R}_{L,n}(\gamma \mid \mathbf{q}) := \frac{1}{n} \sum_{i=1}^{n} L(\gamma(\mathbf{q}(X_i)), Y_i).$$

Now, let $Q_1 \subset Q_2 \subset \cdots \subset Q$ be a non-decreasing collection of quantizers, indexed by sample size n, and similarly let $\Gamma_1 \subset \Gamma_2 \subset \cdots \subset \Gamma$ be a non-decreasing collection of discriminant functions, where we assume the collections satisfy the estimation and approximation error conditions

(23)
$$\mathbb{E}\left[\sup_{\gamma\in\Gamma_{n},\mathsf{q}\in\mathsf{Q}_{n}}\left|\widehat{R}_{L,n}(\gamma\mid\mathsf{q})-R_{L}(\gamma\mid\mathsf{q})\right|\right]\leq\epsilon_{n}^{\mathrm{est}}\right]$$
$$\inf_{\gamma\in\Gamma_{n},\mathsf{q}\in\mathsf{Q}_{n}}R_{L}(\gamma\mid\mathsf{q})-\inf_{\gamma\in\Gamma,\mathsf{q}\in\mathsf{Q}}R_{L}(\gamma\mid\mathsf{q})\leq\epsilon_{n}^{\mathrm{app}}$$

where $\epsilon_n^{\text{est}} \to 0$ and $\epsilon_n^{\text{app}} \to 0$ as $n \to \infty$. Additionally, let R be the risk functional for the cost-weighted misclassification loss L^{cw} (Example 2), where $L^{\text{cw}}(\alpha, y) = \max_i \{c_{yi} \mid \alpha_i = \max_j \alpha_j\}$. Then we have the following result.

THEOREM 3. Assume the conditions (23) and that γ_n and q_n are approximate empirical L-risk minimizers satisfying

$$\epsilon_n^{\text{opt}} := \mathbb{E}\Big[\widehat{R}_{L,n}(\gamma_n \mid \mathsf{q}_n) - \inf_{\gamma \in \Gamma_n, \mathsf{q} \in \mathsf{Q}_n} \widehat{R}_{L,n}(\gamma \mid \mathsf{q})\Big] \to 0 \quad as \ n \to \infty.$$

Let $R^*(\mathbf{Q}) = \inf_{\gamma \in \Gamma, \mathbf{q} \in \mathbf{Q}} R(\gamma \mid \mathbf{q})$. If the loss L is classification calibrated and universally equivalent to the cost-weighted loss L^{cw} , then

$$R(\gamma_n \mid \mathbf{q}_n) - R^{\star}(\mathbf{Q}) \stackrel{L_1}{\to} 0.$$

Theorem 3 guarantees that under the estimation and approximation conditions (23), empirical risk minimization is consistent for minimizing the quantized Bayes risk whenever the loss L is classification calibrated and equivalent to the desired loss. The proof of Theorem 3 reposes on the following risk inequality, which may be of independent interest. The lemma is a consequence of the results on surrogate risk consistency for classification calibration [36, 31, 30] and our universal equivalence guarantees that exhibits the power of calibration and universal equivalence.

LEMMA 4.1. Assume L is classification-calibrated and universally equivalent to the weighted misclassification loss L^{cw} with cost matrix $C \in \mathbb{R}^{k \times k}_+$. Then there exists a continuous concave function h with h(0) = 0 such that

$$R(\gamma \mid \mathbf{q}) - \inf_{\gamma \in \Gamma, \mathbf{q} \in \mathbf{Q}} R(\gamma \mid \mathbf{q}) \le h\left(R_L(\gamma \mid \mathbf{q}) - \inf_{\mathbf{q} \in \mathbf{Q}} R_L^{\star}(\mathbf{q})\right)$$

With the choice $L(\alpha, y) = \sum_{i=1}^{k} c_{yi} [1 + \alpha_i]_+ + \mathbf{I} \{ \mathbf{1}^T \alpha = 0 \}$ or $L(\alpha, y) = \sum_{i=1}^{k} c_{yi} [1 + \alpha_i - \alpha_y]_+$, we may take $h(\epsilon) = (1 + \frac{1}{k})\epsilon$, that is,

$$R(\gamma \mid \mathbf{q}) - \inf_{\gamma \in \Gamma, \mathbf{q} \in \mathbf{Q}} R(\gamma \mid \mathbf{q}) \leq \left(1 + \frac{1}{k}\right) \left[R_L(\gamma \mid \mathbf{q}) - \inf_{\gamma \in \Gamma, \mathbf{q} \in \mathbf{Q}} R_L(\gamma \mid \mathbf{q}) \right].$$

Lemma 4.1 shows that the gap in surrogate risk provides a guaranteed upper bound on the true cost-weighted risk; in the case of the modified hinge losses of Example 5, this gap is linear. In the binary case, even stronger results are possible [23, Lemma 2]—one may take $h(\epsilon) = a\epsilon$ (for some $a < \infty$) in Lemma 4.1 for any margin-based classification-calibrated loss universally equivalent to the 0-1 loss—this relies on the specific form any such binary convex loss must take [23, Eq. (9)]; our Examples 3 (the family-wise loss) and 5 show that fairly different-looking losses can be classification calibrated and universally equivalent to zero-one loss. We provide the proof of Lemma 4.1 in § C.1. Theorem 3, which we prove in § C.2, is then a consequence of this lemma and Theorem 1.

4.4. Examples of universal equivalence. In this section, we give several examples that build off of Theorems 1 and 2, showing that there exist convex losses that allow optimal joint design of quantizers (or measurement strategies) and discriminant functions, opening the way for potentially efficient convex optimization strategies. To that end, we give two hinge-like loss functions that are universally equivalent to the zero-one loss for all prior distributions π . We also give examples of classification calibrated loss functions that are not universally equivalent to the zero-one loss, although minimizing them without quantization yields Bayes-optimal classifiers.

EXAMPLE 6 (Cost-weighted losses). We return to Example 5, where we have $L^{\text{hin}}(\alpha, i) = \sum_{j \neq i} c_{ij} [1 + \alpha_j]_+ + \mathbf{I} \{ \mathbf{1}^T \alpha = 0 \}$. In this case, we have $H_{L^{\text{hin}}}(\pi) = k \min_l \pi^T c_l = k H_{L^{\text{cw}}}(\pi)$, where L^{cw} denotes the cost-weighted misclassification error as in Example 2. Theorem 1 immediately guarantees that the (weighted) hinge loss is universally equivalent to the (weighted) 0-1 loss. The weighted hinge loss L^{hin} is also, as in Example 5, calibrated for the cost-weighted misclassification error. \diamond

EXAMPLE 7 (Max-type losses and zero-one loss). We return to Example 3 and let $L^{\text{fw}}(\alpha, i) = 1 - \alpha_i + \max\{\alpha_{(1)} - 1, \frac{\alpha_{(1)} + \alpha_{(2)}}{2} - \frac{1}{2}, \dots, \frac{\mathbf{1}^T \alpha}{k} - \frac{1}{k}\}$, the

convex family-wise loss. By Example 3, the associated entropy is $H_{L^{\text{fw}}}(\pi) = 1 - \max_j \pi_j = H_{L^{2^o}}$ for $\pi \in \Delta_k$, and Proposition 5 shows that L^{fw} is classification calibrated. We thus have that L^{fw} and the zero-one loss L^{z^o} are universally equivalent by Theorems 1 and 2. \diamond

For our final example, we consider the logistic loss, which is classification calibrated but not universally equivalent to the zero-one loss.

EXAMPLE 8 (Logistic loss). The loss $L^{\log}(\alpha, i) = \log(\sum_{j=1}^{k} e^{\alpha_j - \alpha_i})$ has (Shannon) entropy $H(\pi) = -\sum_{i=1}^{k} \pi_i \log \pi_i$, as in Ex. 4. There are no a, b, csuch that $H_{L^{zo}}(\pi) = 1 - \max_j \pi_j = a H_{L^{\log}}(\pi) + b^T \pi + c$ for all $\pi \in \Delta_k$. Theorem 1 shows that the logistic loss is not universally equivalent to the zero-one loss. That is, in spite of its classification calibration, there are distributions P_1, \ldots, P_k , a collection Q of quantizers $\mathcal{X} \to \mathcal{Z}$, and a sequence $\gamma_n : \mathcal{Z} \to \mathbb{R}^k$ such that $R_{L^{\log}}(\gamma_n \mid \mathbf{q}_n) \to \inf_{\gamma, \mathbf{q} \in \mathbf{Q}} R_{L^{\log}}(\gamma \mid \mathbf{q})$, but $R_{L^{zo}}(\gamma_n \mid \mathbf{q}_n) \not\to \inf_{\gamma, \mathbf{q} \in \mathbf{Q}} R_{L^{zo}}(\gamma \mid \mathbf{q})$.

5. Proof of the Theorems 1 and 2. The remainder of the main body of the paper consists of the major parts of our arguments for Theorems 1 and 2. We divide the proof of the theorems into two parts. The "if" part is straightforward; the "only if" is substantially more complex.

Proof (if direction). We give the proof for Theorem 2; that for Theorem 1 is identical. Assume that dom $f_{\pi}^{(1)} = \text{dom } f_{\pi}^{(2)}$ and there exist $a > 0, b \in \mathbb{R}^{k-1}$, and $c \in \mathbb{R}$ such that Eq. (22) holds. By Definition 2.1 of multi-way *f*-divergences, for any quantizer q, we have

$$D_{f_{\pi}^{(1)}}(P_1,\ldots,P_{k-1}||P_k||\mathbf{q}) = aD_{f_{\pi}^{(2)}}(P_1,\ldots,P_{k-1}||P_k||\mathbf{q}) + b^T\mathbf{1} + c,$$

as $\int_{\mathcal{X}} dP_i = 1$. Applying the relationship (20), we obtain

$$I\left(X,\pi;H_{L^{(1)}}\mid \mathsf{q}\right) = aI\left(X,\pi;H_{L^{(2)}}\mid \mathsf{q}\right) + b^{T}\mathbf{1} + c.$$

As a > 0, the universal equivalence of $L^{(1)}$ and $L^{(2)}$ follows immediately.

We turn to the "only if" part of the proofs of Theorems 1 and 2. A roadmap is as follows: we first define what we call order equivalence of convex functions, which is related to the equivalence of f-divergences and generalized entropies (Def. 5.1). Then, for any two loss functions $L^{(1)}$ and $L^{(2)}$ that are universally equivalent, we show that the associated entropies $H_{L^{(1)}}$ and $H_{L^{(2)}}$, as constructed in the infimal representation (4), and the functions $f^{(1)}$ and $f^{(2)}$ generating the f-divergences via expression (15), are

DUCHI, KHOSRAVI, RUAN

order equivalent (Lemmas 5.1 and 5.2). After this, we provide a characterization of order equivalent closed convex functions (Lemma 5.3), which is the linchpin of our analysis. The lemma shows that for any two order equivalent closed convex functions f_1 and f_2 with dom $f_1 = \text{dom } f_2$, there are parameters $a > 0, b \in \mathbb{R}^k$, and $c \in \mathbb{R}$ such that $f^{(1)}(t) = af^{(2)}(t) + b^T t + c$ for all $t \in \text{dom } f_1 = \text{dom } f_2$. This proves the "only if" part of the Theorems 1 and 2, yielding the desired result. We present the main parts of the proof in the body of the paper, deferring technical nuances to the supplement.

5.1. Universal equivalence and order equivalence. By Definition 4.1 (and its equivalent variant stated (21)), universally equivalent losses $L^{(1)}$ and $L^{(2)}$ induce the same ordering of quantized information measures and f-divergences. The next definition captures this ordering slightly differently.

DEFINITION 5.1. Let $f_1 : \Omega \to \overline{\mathbb{R}}$ and $f_2 : \Omega \to \overline{\mathbb{R}}$ be closed convex functions, where $\Omega \subset \mathbb{R}^k$ is closed convex. Let $m \in \mathbb{N}$ be arbitrary and the matrices $A, B \in \mathbb{R}^{k \times m}$ satisfy $A\mathbf{1} = B\mathbf{1}$, where A has columns $a_i \in \Omega$ and B has columns $b_i \in \Omega$. Then f_1 and f_2 are order-equivalent if for all $m \in \mathbb{N}$ and all such matrices A and B we have

(24)
$$\sum_{j=1}^{m} f_1(a_j) \le \sum_{j=1}^{m} f_1(b_j)$$
 if and only if $\sum_{j=1}^{m} f_2(a_j) \le \sum_{j=1}^{m} f_2(b_j)$.

As the above context suggests, order equivalence has strong connections with universal equivalence of loss functions L and associated f-divergences and generalized entropies. The next two lemmas make this explicit.

LEMMA 5.1. If losses $L^{(1)}$ and $L^{(2)}$ are lower bounded and universally equivalent, then the associated entropies of the construction (4) are order equivalent over $\Delta_k \subset \mathbb{R}^k_+$.

PROOF. Let H_i be the entropy (pointwise Bayes risk) associated with $L^{(i)}$, noting that dom $H_1 = \text{dom } H_2 = \Delta_k$ because $\inf_{\pi \in \Delta_k} H_i(\pi) > -\infty$. Let the matrices $A = [a_1 \cdots a_m] \in \mathbb{R}^{k \times m}_+$ and $B \in \mathbb{R}^{k \times m}_+$ satisfy $a_i, b_i \in \Delta_k$ for each $i = 1, \ldots, m$, and let $v = \frac{1}{m}A\mathbf{1} = \frac{1}{m}B\mathbf{1} \in \Delta_k$. We show that $\sum_{j=1}^m H_1(a_j) \leq \sum_{j=1}^m H_1(b_j)$ if and only if $\sum_{j=1}^m H_2(a_j) \leq \sum_{j=1}^m H_2(b_j)$, that is, expression (24) holds, by constructing appropriate distributions $P_{1:k}$ and π , then applying the universal equivalence of $L^{(1)}$ and $L^{(2)}$.

Let M_0 be any integer large enough that $v_0 = \frac{1}{k}(1 + \frac{1}{M_0})\mathbf{1} - \frac{1}{M_0}v \in \mathbb{R}^k_+$,

so that $v_0 \in \Delta_k$. Then define the vectors $\tilde{a}_1 = v_0, \ldots, \tilde{a}_{mM_0} = v_0$, and let

$$A^{\text{ext}} = [a_1 \cdots a_m \ \widetilde{a}_1 \cdots \widetilde{a}_{mM_0}] \in \mathbb{R}^{k \times M}_+ \text{ and} B^{\text{ext}} = [b_1 \cdots b_m \ \widetilde{a}_1 \cdots \widetilde{a}_{mM_0}] \in \mathbb{R}^{k \times M}_+,$$

where $M = (M_0 + 1)m$. These satisfy $A^{\text{ext}} \mathbf{1} = B^{\text{ext}} \mathbf{1} = \frac{M}{k} \mathbf{1}$. We let a^{ext} and b^{ext} denote the columns of these extended matrices.

Now, let the spaces $\mathcal{X} = [M] \times [M]$ and $\mathcal{Z} = [M]$. Define quantizers $q_1, q_2 : \mathcal{X} \to \mathcal{Z}$ by $q_1(i, j) = i$ and $q_2(i, j) = j$. For $l = 1, \ldots, k$, define the distributions P_l on \mathcal{X} by

$$P_{l}(i,j) = \frac{k^{2}}{M^{2}} \cdot a_{il}^{\text{ext}} b_{jl}^{\text{ext}}, \text{ so } \sum_{j=1}^{M} P_{l}(i,j) = \frac{k}{M} a_{il}^{\text{ext}} \frac{k}{M} \sum_{j=1}^{M} b_{jl}^{\text{ext}} = \frac{k}{M} a_{il}^{\text{ext}}$$

and similarly $\sum_{i} P_l(i, j) = \frac{k}{M} b_{jl}^{\text{ext.}}$. Let $\pi = \frac{1}{k} \mathbf{1}$ be the uniform prior distribution on the label $Y \in \{1, \dots, k\}$, and note that the posterior probability

$$\widetilde{\pi}(\mathsf{q}_{1}^{-1}(\{i\})) = \left[\frac{\pi_{l}\sum_{j}P_{l}(i,j)}{\sum_{l'}\pi_{l'}\sum_{j}P_{l}(i,j)}\right]_{l=1}^{k} = \left[\frac{a_{il}^{\mathrm{ext}}}{\sum_{l'}a_{il'}^{\mathrm{ext}}}\right]_{l=1}^{k} = a_{i}^{\mathrm{ext}} \in \Delta_{k},$$

because $P_l(\mathbf{q}_1^{-1}(i)) = \sum_j P_l(i,j) = \frac{k}{M} a_{il}^{\text{ext}}$, and similarly $\widetilde{\pi}(\mathbf{q}_2^{-1}(\{j\})) = b_j^{\text{ext}} \in \Delta_k$. Taking the expectation over $X \sim \sum_{l=1}^k \pi_l P_l$, we have

$$\mathbb{E}[H_L(\widetilde{\pi}(\mathsf{q}_1^{-1}(\mathsf{q}_1(X))))] = \frac{1}{k} \sum_{i,l} P_l(\mathsf{q}_1^{-1}(i)) H_L(\widetilde{\pi}(\mathsf{q}_1^{-1}(i))) = \frac{1}{M} \sum_{i=1}^M H_L(a_i^{\text{ext}}).$$

because $\sum_{l} a_{il}^{\text{ext}} = 1$. Similarly, $\mathbb{E}[H_L(\widetilde{\pi}(\mathbf{q}_2^{-1}(\mathbf{q}_2(X))))] = \frac{1}{M} \sum_{j=1}^{M} H_{\pi}(b_j^{\text{ext}})$. Recalling the definitions (3) and (20) of the (quantized) information associated with H, we have $I(X, \pi; H \mid \mathbf{q}_1) = H(\pi) - \frac{1}{M} \sum_{i=1}^{M} H(a_i^{\text{ext}})$ and $I(X, \pi; H \mid \mathbf{q}_2) = H(\pi) - \frac{1}{M} \sum_{i=1}^{M} H(b_i^{\text{ext}})$. Then the universal equivalence of losses $L^{(1)}$ and $L^{(2)}$ immediately implies for $\pi = \frac{1}{k}\mathbf{1}$ that

$$H_1(\pi) - \frac{1}{M} \sum_{i=1}^M H_1(a_i^{\text{ext}}) \le H_1(\pi) - \frac{1}{M} \sum_{i=1}^M H_1(b_i^{\text{ext}}) \quad \text{iff}$$
$$H_2(\pi) - \frac{1}{M} \sum_{i=1}^M H_2(a_i^{\text{ext}}) \le H_2(\pi) - \frac{1}{M} \sum_{i=1}^M H_2(b_i^{\text{ext}}).$$

Noting that $a_i^{\text{ext}} = b_i^{\text{ext}}$ for each $i \ge m + 1$, we rearrange the preceding equivalent statements by adding $\frac{1}{M} \sum_{i \ge m+1} H(a_i^{\text{ext}})$ to each side to obtain that the H_i satisfy inequality (24).

DUCHI, KHOSRAVI, RUAN

For f-divergences, a parallel result is possible; as the techniques are similar to those we use to prove Lemma 5.1 (by constructing an explicit discrete space \mathcal{X} and quantizers q), we defer the proof to Supplementary § B.1.

LEMMA 5.2. If losses $L^{(1)}$ and $L^{(2)}$ are universally equivalent for the prior π (Def. 4.1) and lower-bounded, the corresponding f-divergences $f_{L^{(1)},\pi}$ and $f_{L^{(2)},\pi}$ of construction (15) are order equivalent.

5.2. Characterization of the order equivalence of convex functions. Lemmas 5.1 and 5.2 illustrate the intrinsic relationship between the universal equivalence (Def. 4.1) of losses and the order equivalence (Def. 5.1) of their associated generalized entropies and f-divergences. Therefore, it is natural to ask when convex functions are order equivalent. The lemma below characterizes this order equivalence, and coupled with Lemmas 5.1 and 5.2, it immediately implies Theorems 1 and 2.

LEMMA 5.3. Let $f_1, f_2 : \Omega \to \mathbb{R}$ be closed convex functions, where $\Omega \subset \mathbb{R}^k$ is a convex set. Then f_1 and f_2 are order equivalent on Ω if and only if there exist $a > 0, b \in \mathbb{R}^k$, and $c \in \mathbb{R}$ such that for all $t \in \Omega$

(25)
$$f_1(t) = af_2(t) + b^T t + c.$$

While the proof of Lemma 5.3 is complex, we provide a partial proof highlighting the most important parts of the argument, deferring technical details to the supplement. The essential idea is that Lemma 5.3 holds for simplices (and so it certainly holds for H_L); we can then cover any convex set Ω with a number of overlapping simplices to extend the result to all of Ω , which we do fully in Supplement B.2. To demonstrate Lemma 5.3 for simplices, we require

DEFINITION 5.2. Vectors u_0, u_1, \ldots, u_m are affinely independent if

$$u_1 - u_0, \ u_2 - u_0, \ \ldots, \ u_m - u_0,$$

are linearly independent. A set $E \subset \mathbb{R}^k$ is a simplex if $E = \text{Conv}\{u_0, u_1, \ldots, u_k\}$ where u_0, \ldots, u_k are affinely independent.

Then the essential special case of Lemma 5.3 is the following result.

LEMMA 5.4. Let $E = \text{Conv}\{u_0, \ldots, u_k\} \subset \Omega$ where u_0, \ldots, u_k are affinely independent. If f_1 and f_2 are order equivalent, then there exist $a > 0, b \in \mathbb{R}^k$, and $c \in \mathbb{R}$ such that

$$f_1(t) = af_2(t) + b^T t + c \text{ for all } t \in E$$

26

The proof of Lemma 5.4 proceeds in a series of intermediate results, which we provide in turn, deferring proofs to Supplement B.3. Our first step is to argue that we need only prove equivalence results for convex functions on dense subsets of their domains.

LEMMA 5.5 ([17], Prop. IV.1.2.5). Let $f_1, f_2 : \Omega \to \mathbb{R}$ be closed convex and satisfy $f_1(t) = f_2(t)$ for t in a dense subset of Ω . Then $f_1 = f_2$ on Ω .

The first technical lemma we prove is essentially a direct consequence of the definition of order equivalence.

LEMMA 5.6. Let $u_1, \ldots, u_m \in \Omega$, $\alpha \in \mathbb{Q}^m$ satisfy $\mathbf{1}^T \alpha = 1$, and $v \in \Omega$ with $v = \sum_{i=1}^m \alpha_i u_i$. If $f_1, f_2 : \Omega \to \mathbb{R}$ are order equivalent, then

$$\sum_{i=1}^m \alpha_i f_1(u_i) \le f_1(v) \quad if and only if \quad \sum_{i=1}^m \alpha_i f_2(u_i) \le f_2(v).$$

Thus if $\alpha \in \mathbb{Q}^n$ satisfies $\mathbf{1}^T \alpha = 1$ and $u_1, \ldots, u_n \in \Omega$, then

(26)
$$f_1\left(\sum_{i=1}^n \alpha_i u_i\right) = \sum_{i=1}^n \alpha_i f_1(u_i) \text{ iff } f_2\left(\sum_{i=1}^n \alpha_i u_i\right) = \sum_{i=1}^n \alpha_i f_2(u_i).$$

The next lemma shows that we can force equality (25) to hold for the k+1 extreme points and centroid of any simplex in \mathbb{R}^k ; it is intuitive because there are k+2 free parameters in the choices of a > 0, $b \in \mathbb{R}^k$, and $c \in \mathbb{R}$.

LEMMA 5.7. Let $f_1, f_2 : \Omega \to \mathbb{R}$ be closed convex and let $u_0, \ldots, u_k \in \Omega$ be affinely independent. There exist $a > 0, b \in \mathbb{R}^k$, and c such that $f_1(u) = af_2(u) + b^T u + c$ for $u \in \{u_0, \ldots, u_k, u_{cent}\}$, where $u_{cent} = \frac{1}{k+1} \sum_{i=0}^k u_i$.

Lastly, we have the following characterization of the linearity of convex functions over convex hulls.

LEMMA 5.8. Let
$$f : \Omega \to \mathbb{R}$$
 be convex with $u_1, \ldots, u_m \in \Omega$ and $u_{\text{cent}} = \frac{1}{m} \sum_{i=1}^m u_i$. If $f(u_{\text{cent}}) = \frac{1}{m} \sum_{i=1}^m f(u_i)$, then
$$f\left(\sum_{i=1}^m \lambda_i u_i\right) = \sum_{i=1}^m \lambda_i f(u_i) \text{ for all } \lambda \in \mathbb{R}^m_+ \text{ with } \mathbf{1}^T \lambda = 1.$$

With the four lemmas 5.5–5.8, we can now prove Lemma 5.4. By rotating with $u_i - u_0$ and shifting by u_0 , it is no loss of generality to assume that the

functions f_i are defined on $V = \{v \in \mathbb{R}^k_+ \mid \mathbf{1}^T v \leq 1\}$, so that f_1 and f_2 are continuous, defined, convex, and order equivalent on V. We make one further reduction. Let $e_i \in \mathbb{R}^k$ for $1 \leq i \leq k$ be the standard basis for \mathbb{R}^k and $e_0 = \mathbf{0}$ be shorthand for the all-zeros vector. Further, let $e_{\text{center}} = \frac{1}{k+1} \sum_{i=0}^k e_i$ be the centroid of V (so $V = \text{Conv}\{e_0, \ldots, e_k\}$). Lemma 5.7 guarantees the existence of $a > 0, b \in \mathbb{R}^k$, and $c \in \mathbb{R}$ such that

$$f_1(v) = af_2(v) + b^T v + c \text{ for } v \in \{e_0, e_1, \dots, e_k, e_{\text{center}}\}.$$

Now, let $h_1(v) = f_1(v)$ and $h_2(v) = af_2(v) + b^T v + c$, so h_1 and h_2 are convex, order equivalent on V, and satisfy $h_1(v) = h_2(v)$ for $v \in \{e_0, \ldots, e_k, e_{center}\}$. Thus, Lemma 5.4 is equivalent to showing that if h_1, h_2 are convex, order equivalent, and equal on the extreme points and centroid of V, then

(27)
$$h_1(v) = h_2(v) \text{ for } v \in V = \{v \in \mathbb{R}^k_+ \mid \mathbf{1}^T v \le 1\}.$$

We divide our discussion into two cases.

Linear case. Suppose that $h_1(e_{\text{center}}) = \frac{1}{k+1} \sum_{i=0}^k h_1(e_i)$. Then by order equivalence of h_1 and h_2 (Eq. (26)) we have $h_2(e_{\text{center}}) = \frac{1}{k+1} \sum_{i=0}^k h_2(e_i)$. Lemma 5.8 thus implies that h_1 and h_2 are linear on $V = \text{Conv}\{e_0, \ldots, e_k\}$, equal on the vertices of V, and hence equal on its interior.

Nonlinear case. By convexity we have $h_1(e_{\text{center}}) < \frac{1}{k+1} \sum_{i=0}^k h_1(e_i)$, and order equivalence (Lemma 5.6) implies $h_2(e_{\text{center}}) < \frac{1}{k+1} \sum_{i=0}^k h_2(e_i)$. For $v \in V = \text{Conv}\{e_0, \ldots, e_k\}$, we use $v_0 = 1 - \mathbf{1}^T v$ for shorthand, so we may write $v = \sum_{i=0}^k v_i e_i$ and have $[v_0 \ v_1 \ \cdots \ v_k]^T \in \Delta_{k+1}$. Now, fix an arbitrary $v \in V \cap \mathbb{Q}^k$. We wish to show that $h_1(v) = h_2(v)$. To that end we consider consider the gaps due convexity of $h_j(e_{\text{center}})$ to the values of $h_j(e_i)$ relative to those from $h_j(v)$ to $h_j(e_i)$, defining the linear functions $\varphi_j : [0, 1] \to \mathbb{R}$ by

$$\varphi_j(r) := (1-r) \left[h_j(e_{\text{center}}) - \frac{1}{k+1} \sum_{i=0}^k h_j(e_i) \right] + r \left[\sum_{i=0}^k v_i h_j(e_i) - h_j(v) \right]$$

for j = 1, 2. Then

$$\varphi_j(0) = h_j(e_{\text{center}}) - \frac{1}{k+1} \sum_{i=0}^k h_j(e_i) < 0$$

by assumption, and by convexity,

$$\varphi_j(1) = \sum_{i=0}^k v_i h_j(e_i) - h_j(v) \ge 0.$$

The key is that the order equivalence of h_1 and h_2 on V implies that

(28)
$$\operatorname{sign}(\varphi_1(r)) = \operatorname{sign}(\varphi_2(r)) \text{ for } r \in [0,1],$$

so that φ_1 and φ_2 have the same zero crossing $r^* > 0$, i.e. there exists $0 < r^* \le 1$ with $\varphi_1(r^*) = \varphi_2(r^*) = 0$. (We prove equality (28) presently.) At this $r^* > 0$, we find

$$0 = \varphi_1(r^*) - \varphi_2(r^*) = -r^*h_1(v) + r^*h_2(v),$$

where we use that $h_1(e_i) = h_2(e_i)$ for i = 0, ..., k and $h_1(e_{\text{center}}) = h_2(e_{\text{center}})$. That is, $h_1(v) = h_2(v)$, and as $v \in V \cap \mathbb{Q}^k$ is arbitrary and \mathbb{Q}^k is dense, Lemma 5.5 extends the equality $h_1 = h_2$ to all of V. Expression (27) holds.

Returning to the sign equivalence (28), for r > 0, we may divide $\varphi_j(r)$ by r, and we have $\varphi_j(r) \leq 0$ if and only if

$$\frac{1-r}{r} \left[h_j(e_{\text{center}}) - \frac{1}{k+1} \sum_{i=0}^k h_j(e_i) \right] + \sum_{i=0}^k v_i h_j(e_i) \le h_j(v).$$

Defining $\alpha_i = v_i - \frac{1-r}{r(k+1)} \in \mathbb{Q}$ for $i = 0, \ldots, k$ and $\alpha_{k+1} = \frac{1-r}{r}$, the inequality $\varphi_j(r) \leq 0$ is equivalent to $\sum_{i=0}^k \alpha_i h_j(e_i) + \alpha_{k+1} h_j(e_{\text{center}}) \leq h_j(v)$. A calculation yields $\mathbf{1}^T \alpha = 1$ and $\sum_{i=0}^k \alpha_i e_i + \alpha_{k+1} e_{\text{center}} = v$, and applying Lemma 5.6 immediately yields that $\varphi_1(r) \leq 0$ if and only if $\varphi_2(r) \leq 0$ for all $r \in (0, 1] \cap \mathbb{Q}$. Noting that $\varphi_1(0) < 0$ and $\varphi_2(0) < 0$, we obtain equality (28).

6. Discussion. Rather than recapitulating our contributions, we point out a few directions we believe will prove interesting for further study. While Corollary 1 shows that some convex losses are surrogate-risk consistent even with restricted families of classifiers, it does not apply to the practical case in which the collection of discriminants γ is a (convex subset of a) finitedimensional vector space. This longstanding problem certainly deserves further work. Another direction, a bit further afield, is to investigate the links between this work and objective Bayesian approaches and reference priors [5, 4]. In this line of work, one has a family $\{P_{\theta}\}_{\theta \in \Theta}$ of probability models on an observation space \mathcal{X} and before performing inference chooses a prior π on θ to maximize $I_{\pi}(X; \theta)$, the (Shannon) information between $X \sim P_{\theta}$ and $\theta \sim \pi$. For tasks other than minimizing log loss, it may be sensible to use a notion of information and entropy corresponding to the desired loss. Our notions of loss equivalence, including construction of convex losses equivalent to non-convex losses, could provide insight in such situations.

DUCHI, KHOSRAVI, RUAN

References.

- S. M. Ali and S. D. Silvey. A general class of coefficients of divergence of one distribution from another. *Journal of the Royal Statistical Society, Series B*, 28:131–142, 1966.
- [2] P. L. Bartlett, M. I. Jordan, and J. McAuliffe. Convexity, classification, and risk bounds. Journal of the American Statistical Association, 101:138–156, 2006.
- [3] Y. Benjamini and Y. Hochberg. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society, Series* B, 57(1):289–300, 1995.
- [4] J. O. Berger. The case for objective Bayesian analysis. Bayesian Analysis, 1(3): 385–402, 2006.
- [5] J. M. Bernardo. Reference analysis. In D. Day and C. R. Rao, editors, *Bayesian Thinking, Modeling and Computation*, volume 25 of *Handbook of Statistics*, chapter 2, pages 17–90. Elsevier, 2005.
- [6] D. Bertsekas. Nonlinear Programming. Athena Scientific, 1999.
- [7] P. Billingsley. Probability and Measure. Wiley, Second edition, 1986.
- [8] D. Blackwell. Comparison of experiments. In Proceedings of the 2nd Berkeley Symposium on Probability and Statistics, pages 93–102. University of California Press, 1951.
- [9] T. M. Cover and J. A. Thomas. Elements of Information Theory, Second Edition. Wiley, 2006.
- [10] I. Csiszár. Information-type measures of difference of probability distributions and indirect observation. *Studia Scientifica Mathematica Hungary*, 2:299–318, 1967.
- [11] M. H. DeGroot. Uncertainty, information, and sequential experiments. Annals of Mathematical Statistics, 33(2):404-419, 1962.
- [12] J. C. Duchi, L. Mackey, and M. I. Jordan. The asymptotics of ranking algorithms. Annals of Statistics, 41(5):2292–2323, 2013.
- [13] D. García-García and R. C. Williamson. Divergences and risks for multiclass experiments. In Proceedings of the Twenty Fifth Annual Conference on Computational Learning Theory, 2012.
- [14] T. Gneiting and A. Raftery. Strictly proper scoring rules, prediction, and estimation. Journal of the American Statistical Association, 102(477):359–378, 2007.
- [15] P. Grünwald and A. P. Dawid. Game theory, maximum entropy, minimum discrepancy, and robust Bayesian decision theory. Annals of Statistics, 32(4):1367–1433, 2004.
- [16] L. Györfi and T. Nemetz. f-dissimilarity: A generalization of the affinity of several distributions. Annals of the Institute of Statistical Mathematics, 30:105–113, 1978.
- [17] J. Hiriart-Urruty and C. Lemaréchal. Convex Analysis and Minimization Algorithms I & II. Springer, New York, 1993.
- [18] T. Kailath. The divergence and Bhattacharyya distance measures in signal selection. *IEEE Transactions on Communication Technology*, 15(1):52–60, 1967.
- [19] F. Liese and I. Vajda. On divergences and informations in statistics and information theory. *IEEE Transactions on Information Theory*, 52(10):4394–4412, 2006.
- [20] M. Longo, T. D. Lookabaugh, and R. M. Gray. Quantization for decentralized hypothesis testing under communication constraints. *IEEE Transactions on Informa*tion Theory, 36(2):241–255, 1990.
- [21] G. Lugosi and N. Vayatis. On the Bayes-risk consistency of regularized boosting methods. Annals of Statistics, 32(1):30–55, 2004.
- [22] A. W. Marshall, I. Olkin, and B. C. Arnold. Inequalities: Theory of Majorization and

30

Its Applications. Springer, second edition, 2011.

- [23] X. Nguyen, M. J. Wainwright, and M. I. Jordan. On surrogate loss functions and f-divergences. Annals of Statistics, 37(2):876–904, 2009.
- [24] F. Österreicher and I. Vajda. Statistical information and discrimination. IEEE Transactions on Information Theory, 39(3):1036–1039, 1993.
- [25] H. V. Poor and J. B. Thomas. Applications of ali-silvey distance measures in the design of generalized quantizers for binary decision systems. *IEEE Transactions on Communications*, 25(9):893–900, 1977.
- [26] F. Pukelsheim. Optimal Design of Experiments. Classics in Applied Mathematics. SIAM, 1993.
- [27] M. Reid and R. Williamson. Information, divergence, and risk for binary experiments. Journal of Machine Learning Research, 12:731–817, 2011.
- [28] H. Robbins. Some aspects of the sequential design of experiments. Bulletin American Mathematical Society, 55:527–535, 1952.
- [29] R. E. Schapire and Y. Freund. Boosting: Foundations and Algorithms. MIT Press, 2012.
- [30] I. Steinwart. How to compare different loss functions. Constructive Approximation, 26:225–287, 2007.
- [31] A. Tewari and P. L. Bartlett. On the consistency of multiclass classification methods. Journal of Machine Learning Research, 8:1007–1025, 2007.
- [32] N. Tishby, F. Pereira, and W. Bialek. The information bottleneck method. In *The* 37'th Allerton Conference on Communication, Control, and Computing, 1999.
- [33] J. N. Tsitsiklis. Decentralized detection. In Advances in Signal Processing, Vol. 2, pages 297–344. JAI Press, 1993.
- [34] I. Vajda. On the f-divergence and singularity of probability measures. Periodica Mathematica Hungarica, 2(1-4):223-234, 1972.
- [35] R. C. Williamson, E. Vernet, and M. D. Reid. Composite multiclass losses. Journal of Machine Learning Research, to appear, 2016.
- [36] T. Zhang. Statistical analysis of some multi-category large margin classification methods. Journal of Machine Learning Research, 5:1225–1251, 2004.

DUCHI, KHOSRAVI, RUAN

APPENDIX A: PROOFS OF CLASSIFICATION CALIBRATION RESULTS

In this section, we prove Propositions 4 and 5. Before proving the propositions proper, we state several technical lemmas and enumerate continuity properties of Fenchel conjugates that will prove useful. We also collect a few important definitions related to convexity and norms here, which we use without comment in this appendix. For a norm $\|\cdot\|$ on \mathbb{R}^k , we recall the definition of the dual norm $\|\cdot\|_*$ as $\|y\|_* = \sup_{\|u\|\leq 1} u^T y$. For a convex function $f: \mathbb{R}^k \to \overline{\mathbb{R}}$, we let

$$\partial f(u) = \{ g \in \mathbb{R}^k \mid f(v) \ge f(u) + g^T(v - u) \text{ for all } v \in \mathbb{R}^k \}$$

denote the subgradient set of f at the point u. This set is non-empty if $u \in \text{rel int dom } f$ (see [17, Chapter VI]).

A.1. Technical preliminaries. We provide some background on convex functions. We recall Definition 3.2 of uniform convexity, that f is $(\lambda, \kappa, \|\cdot\|)$ -uniformly convex over $C \subset \mathbb{R}^k$ if it is closed and for all $t \in [0, 1]$ and $u_0, u_1 \in C$ we have

(29)
$$f(tu_0 + (1-t)u_1) \leq tf(u_0) + (1-t)f(u_1) - \frac{\lambda}{2}t(1-t) \|u_0 - u_1\|^{\kappa} \left[(1-t)^{\kappa-1} + t^{\kappa-1} \right].$$

We state a related definition of smoothness.

DEFINITION A.1. A function f is $(L, \beta, \|\cdot\|)$ -smooth if it has β -Hölder continuous gradient with respect to the norm $\|\cdot\|$, meaning that

$$\|\nabla f(u_0) - \nabla f(u_1)\|_* \le L \|u_0 - u_1\|^{\beta} \text{ for } u_0, u_1 \in \text{dom } f.$$

Our first technical lemma is an equivalence result for uniform convexity.

LEMMA A.1. Let $f : \Omega \to \mathbb{R}$, where f is closed convex and Ω is a closed convex set. Then f is $(\lambda, \kappa, \|\cdot\|)$ -uniformly convex over Ω if and only if for $u_0 \in \operatorname{rel} \operatorname{int} \Omega$ and all u_1 ,

(30)
$$f(u_1) \ge f(u_0) + s_0^T(u_1 - u_0) + \frac{\lambda}{2} \|u_0 - u_1\|^{\kappa} \text{ for } s_0 \in \partial f(u_0).$$

If inequality (29) holds, then inequality (30) also holds for any points $u_0 \in \Omega$ and s_0 such that $\partial f(u_0) \neq \emptyset$ and $s_0 \in \partial f(u_0)$. See Section A.8.1 for a proof of this lemma. There is also a natural duality between uniform convexity and smoothness of a function's Fenchel conjugate $f^*(v) = \sup_u \{v^T u - f(u)\}$; such dualities are common [17, Ch. X.4].

LEMMA A.2. Let $\Omega \subset \mathbb{R}^k$ be a closed convex set and $f : \Omega \to \mathbb{R}$ be $(\lambda, \kappa, \|\cdot\|)$ -uniformly convex over Ω . Then f^* is $(\lambda^{-\frac{1}{\kappa-1}}, \frac{1}{\kappa-1}, \|\cdot\|_*)$ -smooth (Def. A.1) over dom $f^* = \mathbb{R}^k$.

See Section A.8.2 for a proof of Lemma A.2. We also have two results on the properties of smooth functions, whose proofs we provide in Sections A.8.3 and A.8.4, respectively.

LEMMA A.3. Let f be $(L, \beta, \|\cdot\|)$ smooth. Then $f(u_1) \leq f(u_0) + \nabla f(u_0)^T (u_1 - u_0) + \frac{L}{\beta + 1} \|u_0 - u_1\|^{\beta + 1}.$

LEMMA A.4. Let f be $(L, \beta, \|\cdot\|)$ smooth over \mathbb{R}^k and $\inf_x f(x) > -\infty$. If the sequence u_n satisfies $\lim_n f(u_n) = \inf_x f(x)$, then $\nabla f(u_n) \to 0$.

A.2. Proof of Proposition 4. We state two intermediate lemmas before proving Proposition 4.

LEMMA A.5. If H is symmetric, closed and strictly concave, then $(-H)^*$ is continuously differentiable on \mathbb{R}^k . If $\alpha_i \geq \alpha_j$, then $p = \nabla (-H)^*(\alpha)$ satisfies $p_i \geq p_j$.

PROOF. As H is strictly concave and $(-H)^*(\alpha) = \sup_{\pi \in \Delta_k} \{\pi^T \alpha + H(\pi)\} < \infty$ for all α (suprema of closed concave functions over compact sets are attained), so $(-H)^*$ is continuously differentiable by standard results in convex analysis [17, Thm X.4.1.1], and $\nabla(-H)^*(\alpha) = \operatorname{argmax}_{p \in \Delta_k} \{p^T \alpha + H(p)\}$.

Now let α satisfy $\alpha_i \geq \alpha_j$. As $\nabla(-H)^*(\alpha) = \operatorname{argmax}_{p \in \Delta_k} \{p^T \alpha + H(p)\}$, let us assume for the sake of contradiction that $p_i < p_j$. Then letting A be the permutation matrix swapping entries i and j, the vector p' = Ap satisfies H(p') = H(p) but $H(\frac{1}{2}(p'+p)) > \frac{1}{2}H(p') + \frac{1}{2}H(p) = H(p) = H(p')$, and

$$\alpha^T p - \alpha^T p' = \alpha_i p_i - \alpha_i p_j + \alpha_j p_j - \alpha_j p_i = (\alpha_i - \alpha_j)(p_i - p_j) \le 0.$$

Thus we have $-\alpha^T p \ge -\alpha^T p'$, and so

$$\frac{1}{2}\alpha^{T}(p+p') + H\left(\frac{1}{2}p + \frac{1}{2}p'\right) \ge -\alpha^{T}p + H\left(\frac{1}{2}p + \frac{1}{2}p'\right) > \alpha^{T}p + H(p),$$

a contradiction to the assumed optimality of p. We must have $p_i \geq p_j$ whenever $\alpha_i \geq \alpha_j$.

LEMMA A.6. If H is symmetric and $\sum_{i=1}^{k} \pi_i L(\alpha^*, i) = \inf_{\alpha} \sum_{i=1}^{k} \pi_i L(\alpha, i)$, then $\pi_i > \pi_j$ implies that $\alpha_i^* \ge \alpha_j^*$. If H is strictly concave, then $\alpha_i^* > \alpha_j^*$.

PROOF. Let π satisfy $\pi_i > \pi_j$ as assumed in the lemma, and suppose that $\alpha_i^* < \alpha_j^*$ for the sake of contradiction. Let A be the permutation matrix that swaps α_i^* and α_j^* . Then $(-H)^*(A\alpha^*) = (-H)^*(\alpha^*)$, and $(-H)^*$ is also symmetric, and

$$\sum_{i=1}^{k} \pi_i L(A\alpha^*, i) - \sum_{i=1}^{k} \pi_i L(\alpha^*, i) = -\pi^T A\alpha^* + \pi^T \alpha^* = (\pi_i - \pi_j)(\alpha_i^* - \alpha_j^*) < 0,$$

a contradiction to the optimality of α^* . If H is strictly concave, Lemma A.5 implies that for α^* minimizing $\sum_{i=1}^k \pi_i L(\alpha, i)$, we have $\pi = \nabla (-H)^*(\alpha^*)$. Moreover, by Lemma A.5, if $\pi_i > \pi_j$ we must have $\alpha_i^* > \alpha_j^*$.

PROOF (OF PROPOSITION 4). If the infimum in $\sum_{i=1}^{k} \pi_i L(\alpha, i)$ is attained, then Lemma A.6 gives the result. Otherwise, recall that $H(\pi) = \inf_{\alpha} \{\sum_{i=1}^{k} \pi_i L(\alpha, i)\} > -\infty$, and let $\pi_i > \pi_j$. Let $\alpha^{(m)}$ be any sequence such that $\sum_{i=1}^{k} \pi_i L(\alpha^{(m)}, i) \to H(\pi)$.

Using that H is uniformly concave, we have $\nabla(-H)^*$ is Hölder continuous over \mathbb{R}^k (recall Lemma A.2). This implies that that $\pi - \nabla(-H)^*(\alpha^{(m)}) \to 0$ as $m \to \infty$ (recall Lemma A.4), or

$$\lim_{n \to \infty} \nabla (-H)^* (\alpha^{(m)}) = \pi.$$

Now, by Lemma A.5, for any $p^{(m)} = \nabla (-H)^* (\alpha^{(m)})$, we have that $\alpha_i^{(m)} \ge \alpha_j^{(m)}$ implies $p_i^{(m)} \ge p_j^{(m)}$. Thus, if $\pi_i > \pi_j$, it must be the case that eventually we have $\alpha_i^{(m)} > \alpha_j^{(m)}$. Moreover, if $\liminf |\alpha_i^{(m)} - \alpha_j^{(m)}| = 0$, then we must have $\liminf |p_i^{(m)} - p_j^{(m)}| = 0$, which would contradict that $\pi_i > \pi_j$, as we have $p_i^{(m)} - p_j^{(m)} \to \pi_i - \pi_j$. We thus find that $\liminf (\alpha_i^{(m)} - \alpha_j^{(m)}) > 0$ if the sequence tends to the infimum $H(\pi)$, which implies that

$$\inf_{\alpha} \left\{ \sum_{i=1}^{k} \pi_i L(\alpha, i) : \alpha_i \le \alpha_j \right\} > H(\pi).$$

The loss (11) is thus classification calibrated (Def. 3.1).

A.3. Proof of Proposition 5. Without loss of generality, we assume that $\pi_k < \max_j \pi_j$, so that restricting to $\alpha_k \ge \max_j \alpha_j$ forces α to have larger

34

zero-one risk than $1 - \max_j \pi_j$. We present two lemmas based on convex duality that imply the result. In each, we let $v_{\setminus i} = [v_1 \cdots v_{i-1} v_{i+1} \cdots v_k]^T \in \mathbb{R}^{k-1}$ be the vector $v \in \mathbb{R}^k$ without its *i*th element.

LEMMA A.7. Let $v \in \mathbb{R}^k$. Then

$$\inf_{\alpha_k \ge \max_j \alpha_j} v^T \alpha = \begin{cases} 0 & \text{if } v_k = -\sum_{i=1}^{k-1} v_i, v_{\setminus k} \preceq 0\\ -\infty & \text{otherwise.} \end{cases}$$

PROOF. By introducing Lagrange multipliers $\beta \in \mathbb{R}^{k-1}_+$ for the constraints $\alpha_k \geq \alpha_j$ for $j \neq k$, we have Lagrangian

$$\mathcal{L}(\alpha,\beta) = \left(v + \begin{bmatrix} \beta \\ -\beta^T \mathbf{1} \end{bmatrix}\right)^T \alpha, \text{ so } \inf_{\alpha} \mathcal{L}(\alpha,\beta) = \begin{cases} 0 & \text{if } v_k = \beta^T \mathbf{1}, v_{\setminus k} = -\beta \\ -\infty & \text{otherwise.} \end{cases}$$

Substuting $\beta = -v_{\setminus k}$ and noting that $\beta \succeq 0$ gives the result.

Thus, if we define the matrix $C = \mathbf{1}\mathbf{1}^T - I_{k \times k}$ with columns c_l , we find by strong duality that

$$\inf_{\alpha_k \ge \max_j \alpha_j} \left\{ -\pi^T \alpha + (-H_{L^{zo}})^*(\alpha) \right\}$$
$$= \inf_{\alpha_k \ge \max_j \alpha_j} \left\{ -\pi^T \alpha + \sup_{p \in \Delta_k} \inf_{q \in \Delta_k} \left\{ p^T \alpha + q^T C^T p \right\} \right\}$$
$$= \sup_{p \in \Delta_k} \inf_{\alpha_k \ge \max_j \alpha_j} \left\{ \min_l c_l^T p + (p - \pi)^T \alpha \right\}$$
$$= \sup_{p \in \Delta_k} \left\{ 1 - \max_l p_l \mid \pi_{\setminus k} \succeq p_{\setminus k}, p_k = \pi_k + \mathbf{1}^T (\pi_{\setminus k} - p_{\setminus k}) \right\}$$

by Lemma A.7. The next lemma then immediately implies Proposition 5 once we note that $H_{L^{20}}(\pi) = 1 - \max_{j} \pi_{j}$.

LEMMA A.8. Let
$$G_k(\pi) = \inf_{\alpha_k \ge \max_j \alpha_j} \{-\pi^T \alpha + (-H_{L^{zo}})^*(\alpha)\}$$
. Then
 $G_k(\pi) - (1 - \max_j \pi_j) \ge \frac{1}{k} (\max_j \pi_j - \pi_k).$

PROOF. We essentially construct the optimal $p \in \Delta_k$ vector for the supremum in the definition of G_k . Without loss of generality, we assume that $\pi_1 = \max_j \pi_j > \pi_k$, as the result is trivial if $\pi_k = \max_j \pi_k$. For $t \in \mathbb{R}$, define

$$L(t) = \pi_k + \sum_{j=1}^k [\pi_j - t]_+$$
 and $R(t) = t$.

By defining $t_{\text{low}} = \pi_k$ and $t_{\text{high}} = \pi_1$ we have

$$L(t_{\text{low}}) = \pi_k + \sum_{j=1}^k \left[\pi_j - \pi_k\right]_+ \ge \pi_k + \left[\pi_1 - \pi_k\right]_+ = \pi_1 > \pi_k = R(t_{\text{low}}),$$

$$L(t_{\text{high}}) = \pi_k < \pi_1 = R(t_{\text{high}}),$$

and the fact that L is strictly decreasing in $[t_{\text{low}}, t_{\text{high}}]$ and R is strictly increasing implies that there exists a unique root $t^* \in (t_{\text{low}}, t_{\text{high}})$ such that $L(t^*) = t^* = R(t^*)$. Now, we define the vector p by

$$p_j = \min\{t^*, \pi_j\} \text{ for } j \le k-1, \ p_k = t^*.$$

Then we have $\mathbf{1}^T p = t^* + \sum_{j=1}^{k-1} t^* \wedge \pi_j = \pi_k + \sum_{j=1}^{k-1} ([\pi_j - t^*]_+ + t^* \wedge \pi_j) = 1$, and moreover, we have $G_k(\pi) \ge 1 - \max_l p_l = 1 - t^*$.

It remains to show that $1 - t^* - (1 - \pi_1) \ge \frac{1}{k}(\pi_1 - \pi_k)$; equivalently, we must show that $t^* \le (1 - \frac{1}{k})\pi_1 + \frac{1}{k}\pi_k$. Suppose for the sake of contradiction that this does not hold, that is, that $t^* > (1 - \frac{1}{k})\pi_1 + \frac{1}{k}\pi_k$. We know that $t_{\text{low}} = \pi_k < t^* < \pi_1 = t_{\text{high}}$ by assumption. With t^* satisfying these two constraints, we have that

$$L(t^{\star}) = \pi_k + \sum_{j=1}^k \left[\pi_j - t^{\star}\right]_+ \le \pi_k + \sum_{j=1}^{k-1} \left[\pi_1 - t^{\star}\right]_+ = \pi_k + (k-1)(\pi_1 - t^{\star})$$
$$< \pi_k + (k-1)\left(\pi_1 - \frac{k-1}{k}\pi_1 - \frac{1}{k}\pi_k\right)$$
$$= \frac{k-1}{k}\pi_1 + \frac{1}{k}\pi_k < t^{\star},$$

the two strict inequalities by assumption on t^* . But then we would have $L(t^*) < R(t^*) = t^*$, a contradiction to our choice of t^* . Because L is decreasing, it is thus the case that $t^* \leq (1 - \frac{1}{k})\pi_1 + \frac{1}{k}\pi_k$, giving the result.

A.4. Proof of Observation 1. Our proof is essentially a trivial modification of Zhang [36, Theorem 8]. We assume without loss of generality that $\phi(\cdot) \geq 0$. Let $\pi \in \Delta_k$, and recalling that the cost matrix $C = [c_1 \cdots c_k]$, let

$$L(\pi, \alpha) = \sum_{y=1}^{k} \pi_y \sum_{i=1}^{k} c_{yi} \phi(-\alpha_i) = \sum_{i=1}^{k} \pi^T c_i \phi(-\alpha_i),$$

noting that $\pi^T c_i \geq 0$ for each *i*. Without loss of generality, we may assume that $\pi^T c_1 > \pi^T c_2 = \min_l \pi^T c_l$. If we can show that $\inf_{\alpha} L(\pi, \alpha) < 1$

36

 $\inf_{\alpha_1 \ge \max_j \alpha_j} L(\pi, \alpha)$, then the proof will be complete. Let $\alpha^{(m)} \in \mathbb{R}^k$ be any sequence satisfying $\mathbf{1}^T \alpha^{(m)} = 0$ and $\alpha_1^{(m)} \ge \max_j \alpha_j^{(m)}$ such that $L(\pi, \alpha^{(m)}) \to \inf_{\alpha_1 \ge \max_j \alpha_j} L(\pi, \alpha)$.

We first show that it is no loss of generality to assume that $\alpha^{(m)}$ converges. Suppose for the sake of contradiction that $\limsup_m \sup_m \|\alpha^{(m)}\| = \infty$. Then as $\mathbf{1}^T \alpha^{(m)} = 0$, we must have $\limsup_m \alpha_1^{(m)} = \infty$, so that $\limsup_m \phi(-\alpha_1^{(m)}) = \infty$ because $\phi'(0) < 0$ and ϕ is convex. As it must be the case that $\pi^T c_1 \phi(-\alpha_1^{(m)})$ remains bounded (for the convergence of $L(\pi, \alpha^{(m)})$), we would then have that $\pi^T c_1 = 0$, which is a contradiction because $\pi^T c_1 > \min_l \pi^T c_l \ge 0$. Thus we must have $\limsup_m \|\alpha^{(m)}\| < \infty$, and so there is a subsequence of $\alpha^{(m)}$ converging; without loss of generality, we assume that $\alpha^{(m)} \to \alpha^*$. Then $L(\pi, \alpha^*) = \inf_{\alpha_1 \ge \max \alpha_i} L(\pi, \alpha)$ by continuity of ϕ .

We show that by swapping the value of α_1^{\star} with the value α_2^{\star} (or increasing the latter slightly), we can always improve the value $L(\pi, \alpha^{\star})$. We consider three cases, noting in each that $\alpha_1^{\star} \geq 0$ as $\mathbf{1}^T \alpha^{\star} = 0$.

1. Let $\alpha_1^{\star} = \alpha_2^{\star} \ge 0$. Then $\phi'(-\alpha_1^{\star}) = \phi'(-\alpha_2^{\star}) \le \phi'(0) < 0$, and additionally $c_1^T \pi \phi'(-\alpha_1^{\star}) - c_2^T \pi \phi'(-\alpha_2^{\star}) < 0$ because $c_2^T \pi < c_1^T \pi$. For sufficiently small $\delta > 0$, we thus have

$$c_1^T \pi \phi(-\alpha_1^{\star} + \delta) + c_2^T \pi \phi(-\alpha_2^{\star} - \delta) < c_1^T \pi \phi(-\alpha_1^{\star}) + c_2^T \pi \phi(-\alpha_2^{\star}),$$

whence $L(\pi, \alpha^*) > L(\pi, \alpha^* - \delta(e_1 - e_2)).$

- 2. Let $\alpha_1^{\star} > \alpha_2^{\star}$ and $\phi(-\alpha_1^{\star}) > \phi(-\alpha_2^{\star})$. Then taking $\alpha \in \mathbb{R}^k$ such that $\alpha_1 = \alpha_2^{\star}, \alpha_2 = \alpha_1^{\star}$, and $\alpha_i = \alpha_i^{\star}$ for $i \geq 3$, it is clear that $L(\pi, \alpha) < L(\pi, \alpha^{\star})$.
- 3. Let $\alpha_1^* > \alpha_2^*$ and $\phi(-\alpha_1^*) \le \phi(-\alpha_2^*)$. Using that $-\alpha_1^* < 0$, we see that for sufficiently small $\delta > 0$ we have $\phi(-\alpha_1^* + \delta) < \phi(-\alpha_1^*)$ and $\phi(-\alpha_2^*) \ge \phi(-\alpha_2^* \delta)$, because ϕ must be non-decreasing at the point $-\alpha_2^*$ as $\phi'(0) < 0$. Thus we have $L(\pi, \alpha^* \delta(e_1 e_2)) < L(\pi, \alpha^*)$.

A.5. Biconjugates of the zero-one loss. In this subsection, we calculate the conjugate of the generalized entropy $H(\pi) = 1 - \max_j \pi_j$ (pointwise Bayes risk) associated with the zero-one loss, demonstrating equality (16). Let $(-H)^*(\alpha) = \sup_{\pi \in \Delta_k} \{\pi^T \alpha + 1 - \|\pi\|_\infty\}$. Formulating the Lagrangian for the supremum with dual variables $\theta \in \mathbb{R}$, $\lambda \in \mathbb{R}^k_+$, we have

$$\mathcal{L}(\pi,\theta,\lambda) = \pi^T \alpha - \left\|\pi\right\|_{\infty} - \theta(\mathbf{1}^T \pi - 1) + \lambda^T \pi,$$

which has dual objective

$$\sup_{\pi} \mathcal{L}(\pi, \theta, \lambda) = \begin{cases} \theta & \text{if } \|\alpha + \lambda - \theta \mathbf{1}\|_{1} \leq 1 \\ -\infty & \text{otherwise.} \end{cases}$$

As $\inf_{\lambda \ge 0} \|\alpha + \lambda - \theta \mathbf{1}\|_1 = \sum_{j=1}^k [\alpha_j - \theta]_+$ and strong duality obtains (the problems are linear), we have that the supremum in expression (16) is

$$\sup_{\pi} \left\{ \pi^{T} \alpha - \|\pi\|_{\infty} \mid \pi^{T} \mathbf{1} = 1, \pi \ge 0 \right\} = \inf \left\{ \theta \mid \sum_{j=1}^{k} \left[\alpha_{j} - \theta \right]_{+} \le 1 \right\}.$$

Without loss of generality we may assume that $\alpha_1 \geq \alpha_2 \geq \cdots$ by symmetry, so that over the domain $\theta \in (-\infty, \alpha_1]$, the function $\theta \mapsto \sum_{j=1}^k [\alpha_j - \theta]_+$ is strictly decreasing. Thus, there is a unique smallest θ satisfying $\sum_{j=1}^k [\alpha_j - \theta]_+ \leq$ 1 (attaining the equality), and by inspection, this must be one of

$$\theta \in \left\{ \alpha_1 - 1, \frac{\alpha_1 + \alpha_2}{2} - \frac{1}{2}, \frac{\alpha_1 + \alpha_2 + \alpha_3}{3} - \frac{1}{3}, \dots, \frac{1^T \alpha}{k} - \frac{1}{k} \right\}$$

(Any θ makes some number of the terms $[\alpha_j - \theta]_+$ positive; fixing the number of terms and solving for θ gives the preceding equality.) Expression (16) follows.

A.6. Pointwise infimal risks of hinge losses. We demonstrate equality (18) with $C = [c_1 \cdots c_k]$. We note that

$$H(\pi) = \inf_{\alpha^T \mathbf{1} = 0} \left\{ \sum_{y=1}^k \pi_y \sum_{i=1}^k c_{yi} \left[1 + \alpha_i \right]_+ \right\} = \inf_{\alpha^T \mathbf{1} = 0} \left\{ \sum_{i=1}^k \pi^T c_i \left[1 + \alpha_i \right]_+ \right\},$$

and formulating the Lagrangian by introducing dual variable $\theta \in \mathbb{R}$, we have

$$\mathcal{L}(\alpha, \theta) = \sum_{i=1}^{k} \pi^{T} c_{i} \left[1 + \alpha_{i}\right]_{+} - \theta \mathbf{1}^{T} \alpha.$$

The generalized KKT conditions [6] for this problem are given by taking subgradients of the Lagrangian. At optimum, we must have

$$\nu_i \in \partial [1 + \alpha_i]_+ = \begin{cases} 0 & \text{if } \alpha_i < -1 \\ [0,1] & \text{if } \alpha_i = -1 \\ 1 & \text{if } \alpha_i > -1, \end{cases} \text{ and } \pi^T c_i \nu_i - \theta = 0, \ \mathbf{1}^T \alpha = 0, \ i \in [k].$$

Without loss of generality, assume that $\pi^T c_1 = \min_j \pi^T c_j$. Set $\alpha^* \in \mathbb{R}^k$ and θ^* via

$$\alpha_1^* = (k-1), \quad \alpha_i^* = -1 \text{ for } i \ge 2, \quad \theta^* = \min_j \pi^T c_j = \pi^T c_1.$$

We have $\mathbf{1}^T \alpha^* = (k-1) - (k-1) = 0$, and setting $\nu_i = 1$ for *i* such that $\pi^T c_i = \min_j \pi^T c_j$ and $\nu_i = \frac{\min_j \pi^T c_j}{\pi^T c_i} \in [0,1]$ otherwise, we see that $\pi^T c_i \nu_i - \theta^* = \min_j \pi^T c_j - \min_j \pi^T c_j = 0$ for all *i*; the KKT conditions are satisfied. Thus α^* and θ^* are primal-dual optimal, yielding expression (18).

A.7. Classification calibration of hinge losses. We provide a quantitative guarantee on the classification calibration of hinge-like losses.

LEMMA A.9. Let $C \in \mathbb{R}^{k \times k}_+$ and let $L^{cw}(\alpha, y) = \max_i \{c_{yi} \mid \alpha_i = \max_j \alpha_j\}$ be the cost-weighted loss (10). Let $L(\alpha, y) = \sum_{i=1}^k c_{yi} [1 + \alpha_i]_+ + \mathbf{I} \{\mathbf{1}^T \alpha = 0\}$ or $L(\alpha, y) = \sum_{i=1}^k c_{yi} [1 + \alpha_i - \alpha_y]_+$. Then for any $\pi \in \Delta_k$ and $\alpha \in \mathbb{R}^k$,

$$\sum_{i=1}^k \pi_i L(\alpha, i) - \inf_{\alpha'} \sum_{i=1}^k \pi_i L(\alpha', i) \ge \sum_{i=1}^k \pi_i L^{\mathrm{cw}}(\alpha, i) - \inf_{\alpha'} \sum_{i=1}^k \pi_i L^{\mathrm{cw}}(\alpha', i).$$

PROOF. First, recall from Example 5 that $\inf_{\alpha} \sum_{i=1}^{k} \pi_i L(\alpha, i) = k \min_l \pi^T c_l$. Defining $y(\alpha) = \operatorname{argmax}_j \alpha_j$ (breaking ties in some arbitrary deterministic order), it is sufficient to argue that for $L(\alpha, y) = \sum_{i=1}^{k} c_{yi} [1 + \alpha_i]_+$, we have

(31)
$$\sum_{y} \pi_{y} L(\alpha, y) - k \min_{l} \pi^{T} c_{l} \ge \left[c_{y(\alpha)}^{T} \pi - \min_{l} \pi^{T} c_{l} \right]$$

for any vector $\pi \in \Delta_k$ and α with $\mathbf{1}^T \alpha = 0$.

To show inequality (31), assume without loss of generality that there exists an index $l^* < k$ such that $c_1^T \pi = c_2^T \pi = \cdots = c_{l^*}^T \pi = \min_l c_l^T \pi$, while $c_l^T \pi > c_1^T \pi$ for $l > l^*$. (If $l^* = k$, then inequality (31) is trivial.) We always have $\sum_y \pi_y L(\alpha, y) \ge k \min_l c_l^T \pi$; let us suppose that $\alpha_l \ge \max_j \alpha_j$ for some $l > l^*$; without loss of generality take l = k. Then we have

(32)

$$\sum_{y=1}^{k} \pi_{y} L(\alpha, y) \geq \inf_{\alpha_{k} \geq \max_{j} \alpha_{j}, \mathbf{1}^{T} \alpha = 0} \sum_{y=1}^{k} \pi_{y} \sum_{j=1}^{k} c_{yj} \left[1 + \alpha_{j}\right]_{+}$$

$$= \inf_{\alpha_{k} \geq \max_{j} \alpha_{j}, \mathbf{1}^{T} \alpha = 0} \sum_{l=1}^{k} c_{l}^{T} \pi \left[1 + \alpha_{l}\right]_{+}.$$

Writing the Lagrangian for this problem and introducing variables $\lambda \geq 0$ for the inequality $\alpha_k \geq \max_j \alpha_j$ and $\theta \in \mathbb{R}$ for the equality $\mathbf{1}^T \alpha = 0$, we have

$$\mathcal{L}(\alpha, \lambda, \theta) = \sum_{l=1}^{k} \pi^{T} c_{l} \left[1 + \alpha_{l} \right]_{+} + \theta \mathbf{1}^{T} \alpha + \lambda \left(\max_{j} \alpha_{j} - \alpha_{k} \right).$$

Set $\alpha_1^{\star} = \cdots = \alpha_{l^{\star}}^{\star} = \frac{k - l^{\star} - 1}{l^{\star} + 1}$, $\alpha_k = \frac{k - l^{\star} - 1}{l^{\star} + 1}$, and $\alpha_l = -1$ for $l > l^{\star}$, $l \neq k$. We claim that these are optimal. Indeed, set $\theta^{\star} = -(1 - \epsilon)\pi^T c_1 - \epsilon \pi^T c_k$ for some $\epsilon \in (0, 1)$, whose value we specify later. Taking subgradients of the Lagrangian with respect to α , we have

$$\partial_{\alpha} \mathcal{L}(\alpha^{\star}, \lambda, \theta^{\star}) = \operatorname{diag}(\nu) \begin{bmatrix} c_1^T \\ \vdots \\ c_k^T \end{bmatrix} \pi + \theta^{\star} \mathbf{1} + \lambda \left[\operatorname{Conv} \{ e_1, \dots, e_{l^{\star}}, e_k \} - e_k \right],$$

where $\nu_i \in [0,1]$ are arbitrary scalars satisfying $\nu_i \in \partial [1 + \alpha_i^*]_+$, i.e. $\nu_1 = \cdots = \nu_{l^*} = \nu_k = 1, \ \nu_l \in [0,1]$ for $l \in \{l^* + 1, \dots, k-1\}$. Notably, we have $\pi^T c_l + \theta^* = \epsilon \pi^T (c_1 - c_k) \leq 0$ for $l \in \{1, \dots, l^*\}$ and $\pi^T c_k + \theta^* = (1 - \epsilon)\pi^T (c_k - c_1) \geq 0$. Choosing ϵ small enough that $\epsilon \pi^T c_k + (1 - \epsilon)\pi^T c_1 < \pi^T c_l$ for $l \in \{l^* + 1, \dots, k-1\}$, it is clear that we may take $\nu_l = \frac{(1 - \epsilon)\pi^T c_1 + \epsilon \pi^T c_k}{\pi^T c_l} \in [0, 1]$ for each $l \notin \{1, \dots, l^*, k\}$, and we show how to choose $\lambda^* \geq 0$ so that $\mathbf{0} \in \partial_{\alpha} \mathcal{L}(\alpha^*, \lambda, \theta^*)$. Assume without loss of generality (by scaling of $\lambda \geq 0$) that $\pi^T (c_k - c_1) = 1$. Eliminating extraneous indices in $\partial_{\alpha} \mathcal{L}$, we see that we seek a setting of λ and a vector $v \in \Delta_{l^*+1}$ such that

$$e_{l^{\star}+1} - \epsilon \mathbf{1}_{l^{\star}+1} + \lambda(v - e_{l^{\star}+1}) = 0.$$

This is straightforward: take $v = \frac{1}{l^{\star} \epsilon + (1-\epsilon)} [\epsilon \cdots \epsilon (1-\epsilon)]^T \in \Delta_{l^{\star}+1}$, and set $\lambda^{\star} = l^{\star} \epsilon + (1-\epsilon) \ge 0$.

Summarizing, we find that our preceding choices of α^* were optimal for the problem (32), that is, $\alpha_1^* = \cdots = \alpha_{l^*} = \frac{k-l^*-1}{l^*+1} = \alpha_k$, with $\alpha_l = -1$ for $l^* < l < k$. We thus have

$$\sum_{y=1}^{k} \pi_y L(\alpha, y) \ge \sum_{l=1}^{l^*} \pi^T c_l \left(1 + \frac{k - l^* - 1}{l^* + 1} \right) + \pi^T c_k \left(1 + \frac{k - l^* - 1}{l^* + 1} \right)$$
$$= \min_l \pi^T c_l \left(l^* + \frac{l^*}{l^* + 1} (k - l^* - 1) \right) + \frac{k}{l^* + 1} c_k^T \pi.$$

In particular, we have

$$\sum_{y=1}^{k} \pi_{y} L(\alpha, y) - k \min_{l} \pi^{T} c_{l}$$

$$\geq \min_{l} \pi^{T} c_{l} \left(l^{\star} + \frac{l^{\star}}{l^{\star} + 1} (k - l^{\star} - 1) \right) + \frac{k}{l^{\star} + 1} c_{k}^{T} \pi - k \min_{l} c_{l}^{T} \pi$$

$$= \frac{k}{l^{\star} + 1} \left[c_{k}^{T} \pi - \min_{l} c_{l}^{T} \pi \right].$$

Recalling that we must have had $l^* < k$, we obtain inequality (31).

A.8. Proofs of technical lemmas on smoothness.

A.8.1. Proof of Lemma A.1. Let $t \in (0,1)$ and $u^t = tu_1 + (1-t)u_0$. Assume that inequality (30) holds and let $u_0 \in \operatorname{rel} \operatorname{int} \Omega$. Then

$$f(u_1) \ge f(u^t) + s_t^T(u_1 - u^t) + \frac{\lambda}{2} \|u^t - u_1\|^{\kappa}$$

= $f(u^t) + (1 - t)s_t^T(u_1 - u_0) + \frac{\lambda}{2}(1 - t)^{\kappa} \|u_0 - u_1\|^{\kappa}$

where $s_t \in \partial f(u^t)$, which must exist as $u^t \in \text{rel int } \Omega$ (see Lemma B.3 and [17, Chapter VI]). Similarly, we have

$$f(u_0) \ge f(u^t) + ts_t^T(u_0 - u_1) + \frac{\lambda}{2}t^{\kappa} ||u_0 - u_1||^{\kappa}.$$

Multiplying the preceding inequalities by t and (1 - t), respectively, gives

$$tf(u_1) + (1-t)f(u_0) \ge f(u^t) + \frac{\lambda}{2} \|u_0 - u_1\|^{\kappa} [t(1-t)^{\kappa} + (1-t)t^{\kappa}]$$

= $f(u^t) + \frac{\lambda}{2} t(1-t) \|u_0 - u_1\|^{\kappa} [(1-t)^{\kappa-1} + t^{\kappa-1}]$

for any $u_0 \in \operatorname{rel} \operatorname{int} \Omega$.

We now consider the case that $u_0 \in \Omega \setminus \text{rel int } \Omega$, as the preceding display is equivalent to the uniform convexity condition (29). Let $u_0 \in \Omega \setminus \text{rel int } \Omega$ and $u'_0 \in \text{rel int } \Omega$. For $a \in [0, 1]$ define the functions $h(a) = tf(u_1) + (1 - t)f((1 - a)u_0 + au'_0)$ and

$$h_t(a) = f(tu_1 + (1-t)((1-a)u_0 + au'_0)) + \frac{\lambda}{2}t(1-t) \left\| (1-a)u_0 + au'_0 - u_1 \right\|^{\kappa} \left[(1-t)^{\kappa-1} + t^{\kappa-1} \right].$$

Then h and h_t are closed one-dimensional convex functions, which are thus continuous [17, Chapter I], and we have $h(a) \ge h_t(a)$ for all $a \in (0,1)$ as $(1-a)u_0 + au'_0 \in \text{rel int }\Omega$. Thus

$$tf(u_0) + (1-t)f(u_1) = h(0) = \lim_{a \to 0} h(a) \ge \lim_{a \to 0} h_t(a) = h_t(0)$$

= $f(tu_1 + (1-t)u_0) + \frac{\lambda}{2}t(1-t) ||u_0 - u_1||^{\kappa} [(1-t)^{\kappa-1} + t^{\kappa-1}].$

This is equivalent to the uniform convexity condition (29).

We now prove the converse. Assume the uniform convexity condition (29), which is equivalent to

$$\frac{f(u^t) - f(u_0)}{t} + \frac{\lambda}{2}(1-t) \|u_0 - u_1\|^{\kappa} \left[(1-t)^{\kappa-1} + t^{\kappa-1}\right] \le f(u_1) - f(u_0)$$

for all $u_0, u_1 \in \Omega$ and $t \in (0, 1)$. Let $f'(x, d) = \lim_{t \downarrow 0} \frac{f(x+td) - f(x)}{t}$ be the directional derivative of f in direction d, recalling that if $\partial f(x) \neq \emptyset$ then $f'(x, d) = \sup_{q \in \partial f(x)} \langle g, d \rangle$ (see [17, Ch. VI.1]). Then taking $t \downarrow 0$, we have

$$f'(u_0, u_1 - u_0) + \frac{\lambda}{2} \|u_0 - u_1\|^{\kappa} \le f(u_1) - f(u_0).$$

Because $f'(u_0, u_1 - u_0) = \sup_{g \in \partial f(u_0)} \langle g, u_1 - u_0 \rangle$, this implies the subgradient condition (30).

A.8.2. Proof of Lemma A.2. First, we note that as dom $f = \Omega$ and f is uniformly convex, it is 1-coercive, meaning that $\lim_{\|u\|\to\infty} f(u)/\|u\| = \infty$. Thus dom $f^* = \mathbb{R}^k$; see [17, Proposition X.1.3.8].

As f is strictly convex by assumption, we have that f^* is differentiable [17, Theorem X.4.1.1]. Moreover, as f is closed convex, $f = f^{**}$, and we have for any $s \in \mathbb{R}^k$ that $u_0 = \nabla f^*(s)$ if and only if $s \in \partial f(u_0)$, meaning that f is subdifferentiable on the set $\operatorname{Im} \nabla f^* = {\nabla f^*(s) : s \in \mathbb{R}^k}$, whence $\Omega \supset \operatorname{Im} \nabla f^*$. Now, let $s_0, s_1 \in \mathbb{R}^k$ and $u_0 = \nabla f^*(s_0)$ and $u_1 = \nabla f^*(s_1)$. We must then have $s_0 \in \partial f(u_0)$ and $s_1 \in \partial f(u_1)$ by standard results in convex analysis [17, Corollary X.1.4.4], so that $\partial f(u_0) \neq \emptyset$ and $\partial f(u_1) \neq \emptyset$.

Now we use the uniform convexity condition (30) of Lemma A.1 to see

$$f(u_1) - f(u_0) \ge \langle s_0, u_1 - u_0 \rangle + \frac{\lambda}{2} ||u_0 - u_1||^{\kappa} \text{ and} f(u_0) - f(u_1) \ge \langle s_1, u_0 - u_1 \rangle + \frac{\lambda}{2} ||u_0 - u_1||^{\kappa}.$$

Adding these equations, we find that

$$\langle s_0 - s_1, u_0 - u_1 \rangle \ge \lambda \|u_0 - u_1\|^{\kappa}$$
, so $\|s_0 - s_1\|_* \|u_0 - u_1\| \ge \lambda \|u_0 - u_1\|^{\kappa}$

by Hölder's inequality. Dividing each side by $||u_0 - u_1||$, we obtain $||u_0 - u_1|| \le \lambda^{-\frac{1}{\kappa-1}} ||s_0 - s_1||_*^{\frac{1}{\kappa-1}}$, the desired result once we note that $\nabla f^*(s_i) = u_i$.

A.8.3. Proof of Lemma A.3. Define $u_t = u_0 + t(u_1 - u_0)$ for $t \in [0, 1]$. Then using Taylor's theorem, we have

$$f(u_1) = f(u_0) + \int_0^1 \langle \nabla f(tu_1 + (1-t)u_0), u_1 - u_0 \rangle dt$$

= $f(u_0) + \langle \nabla f(u_0), u_1 - u_0 \rangle + \int_0^1 \langle \nabla f(u_t) - \nabla f(u_0), u_1 - u_0 \rangle dt$
 $\leq f(u_0) + \langle \nabla f(u_0), u_1 - u_0 \rangle + \int_0^1 \| \nabla f(u_t) - \nabla f(u_0) \|_* \| u_0 - u_1 \| dt$
 $\leq f(u_0) + \langle \nabla f(u_0), u_1 - u_0 \rangle + \int_0^1 Lt^{\beta} \| u_0 - u_1 \|^{\beta+1} dt.$

Computing the final integral as $\int_0^1 t^\beta dt = \frac{1}{\beta+1}$ gives the result.

A.8.4. Proof of Lemma A.4. Let $g_n = \nabla f(u_n)$ and suppose for the sake of contradiction that $g_n \not\rightarrow 0$. Then there is a subsequence, which without loss of generality we take to be the full sequence, such that $||g_n||_* \ge c > 0$ for all n. Fix $\delta > 0$, which we will choose later. By Lemma A.3, defining $y_n = u_n - \delta g_n / ||g_n||_*$, we have

$$f(y_n) \le f(u_n) - \delta \|g_n\| + \frac{L}{\beta+1} L \delta^{\beta+1} \le f(u_n) - \delta \left(c - \frac{L}{\beta+1} \delta^{\beta}\right).$$

In particular, we see that if

$$\delta \le \left(\frac{c}{2} \cdot \frac{\beta+1}{L}\right)^{1/\beta}$$
, then $f(y_n) \le f(u_n) - \frac{\delta c}{2}$

for all n, which contradicts the fact that $f(u_n) \to \inf_x f(x) > -\infty$.

APPENDIX B: ORDER EQUIVALENCE OF CONVEX FUNCTIONS

In this appendix, we collect the proofs of our various technical results on order equivalent functions (Definition 5.1).

B.1. Proof of Lemma 5.2. Before proving the lemma, we give a matrix characterization of non-negative vectors with equal sums similar to the characterization of majorization via doubly stochastic matrices (cf. [22]; we temporarily defer the proof of the lemma to Appendix B.1.1).

LEMMA B.1. Vectors $a, b \in \mathbb{R}^m_+$ satisfy $\mathbf{1}^T a = \mathbf{1}^T b$ if and only if there exists a matrix $Z \in \mathbb{R}^{m \times m}_+$ such that $Z\mathbf{1} = a$ and $Z^T\mathbf{1} = b$.

Returning to the proof of Lemma 5.2 proper, let $f_i = f_{L^{(i)},\pi}$ for shorthand. Note that dom $f_1 = \text{dom } f_2 = \mathbb{R}^{k-1}_+$ by the construction (15), because $\min_i \inf_{\alpha \in \mathbb{R}^k} L(\alpha, i) > -\infty$. Now, given matrices $A, B \in \mathbb{R}^{(k-1)\times m}_+$ satisfying $A\mathbf{1} = B\mathbf{1}$ (where $A = [a_1 \cdots a_m]$ and $B = [b_1 \cdots b_m]$), we construct distributions P_i and quantizers \mathbf{q}_1 and \mathbf{q}_2 such that for any f we have

$$D_f(P_1, \dots, P_{k-1} || P_k | \mathbf{q}_1) = C \sum_{j=1}^m f(a_j)$$
 and
 $D_f(P_1, \dots, P_{k-1} || P_k | \mathbf{q}_2) = C \sum_{j=1}^m f(b_j),$

where C > 0 is a constant. We then use Definition 4.1 of loss equivalence to show that f_1 and f_2 are order equivalent.

With that in mind, take M to be any positive integer such that $M > \max\{\|A\mathbf{1}\|_{\infty}, m\}$. We enlarge A and B into matrices $A^{\text{ext}}, B^{\text{ext}} \in \mathbb{R}^{(k-1) \times M}_+$ respectively, adding M-m columns. To construct these matrices, let $a_{i,m+1}^{\text{ext}} = M - \sum_{j=1}^{m} a_{ij}$ and $b_{i,m+1}^{\text{ext}} = M - \sum_{j=1}^{m} b_{ij}$, for $i = 1, \ldots, k-1$, and set $a_{il}^{\text{ext}} = b_{il}^{\text{ext}} = 0$ for all $m+1 < l \leq M$. The enlarged matrices A^{ext} and B^{ext} thus satisfy $\frac{1}{M}A^{\text{ext}}\mathbf{1} = \frac{1}{M}B^{\text{ext}}\mathbf{1} = \mathbf{1}$, and their columns belong to dom $f_1 = \text{dom } f_2 = \mathbb{R}^{k-1}_+$.

Let the spaces $\mathcal{X} = \{(i,j) \mid 1 \leq i \leq M, 1 \leq j \leq M\}$ and $\mathcal{Z} = \{1,2,\ldots,M\}$. Define quantizers $q_1,q_2 : \mathcal{X} \to [M]$ by $q_1(i,j) = i$ and $q_2(i,j) = j$. As $A^{\text{ext}}\mathbf{1} = B^{\text{ext}}\mathbf{1} = M\mathbf{1}$, Lemma B.1 guarantees the existence of matrices $Z^l = [z_{ij}^l] \in \mathbb{R}^{M \times M}_+$ such that

$$Z^{l}\mathbf{1} = \frac{1}{M} [a_{lj}^{\text{ext}}]_{j=1}^{M} \in \mathbb{R}^{M}_{+} \text{ and } (Z^{l})^{T}\mathbf{1} = \frac{1}{M} [b_{lj}^{\text{ext}}]_{j=1}^{M} \in \mathbb{R}^{M}_{+},$$

which implies that for all $l \in [k-1]$, the matrix Z^l satisfies $\sum_{ij} z_{ij}^l = 1$. For each $l \in [k-1]$, define the probability distribution P_l on \mathcal{X} by $P_l((i, j)) = z_{ij}^l$ for $i, j \in [M]$. Let P_k be the distribution defined by $P_k((i, j)) = M^{-2}$.

Under this quantizer design and choice of distributions P_l , we have for any prior $\pi \in \mathbb{R}^k_+$ (upon which the functions f_1 and f_2 implicitly depend) and $f = f_1$ or $f = f_2$ that

$$D_f(P_1, \dots, P_{k-1} || P_k | \mathbf{q}_1) = \frac{1}{M} \sum_{j=1}^M f(a_j^{\text{ext}}) \text{ and}$$
$$D_f(P_1, \dots, P_{k-1} || P_k | \mathbf{q}_2) = \frac{1}{M} \sum_{j=1}^M f(b_j^{\text{ext}}),$$

44

because $\sum_{i=1}^{M} P_l((i,j)) = \sum_{i=1}^{M} z_{ij}^l = a_{lj}^{\text{ext}}/M$ and $\sum_{i=1}^{M} P_k((i,j)) = M^{-1}$, and similarly for B^{ext} . By the loss equivalence of $L^{(1)}$ and $L^{(2)}$ (recall Def. 4.1), we obtain that

$$\sum_{j=1}^{M} f_1(a_j^{\text{ext}}) \le \sum_{j=1}^{M} f_1(b_j^{\text{ext}}) \text{ if and only if } \sum_{j=1}^{M} f_2(a_j^{\text{ext}}) \le \sum_{j=1}^{M} f_2(b_j^{\text{ext}}).$$

Note that $a_j^{\text{ext}} = b_j^{\text{ext}} \in \text{dom } f_i$ for each j > m and i = 1, 2. Moreover, $a_j^{\text{ext}} = a_j$ and $b_j^{\text{ext}} = b_j$ for $1 \le j \le m$, so the preceding display is equivalent to the order equivalence (24).

B.1.1. Proof of Lemma B.1. One direction of the proof is easy: if $Z \in \mathbb{R}^{m \times m}_+$ and $Z\mathbf{1} = a$ while $Z^T\mathbf{1} = b$, then $\mathbf{1}^T Z\mathbf{1} = \mathbf{1}^T a = b^T\mathbf{1}$.

We prove the converse using induction. When m = 1, the result is immediate. We claim that it is no loss of generality to assume that $a_1 \ge a_2 \ge$ $\dots \ge a_m$ and $b_1 \ge \dots \ge b_m$; indeed, let P_a and P_b be permutation matrices such that $P_a a$ and $P_b b$ are in sorted (decreasing) order. Then if we construct $\widetilde{Z} \in \mathbb{R}^{m \times m}_+$ such that $\widetilde{Z}\mathbf{1} = P_a a$ and $\widetilde{Z}^T\mathbf{1} = P_b b$, we have $Z = P_a^T \widetilde{Z} P_b$ satisfies $Z \in \mathbb{R}^{m \times m}_+$ and $Z\mathbf{1} = P_a^T \widetilde{Z}\mathbf{1} = P_a^T P_a a = a$ and $Z^T\mathbf{1} = P_b^T \widetilde{Z}\mathbf{1} = b$.

Now, suppose that the statement of the lemma is true for all vectors of dimension up to m-1; we argue that the result holds for $a, b \in \mathbb{R}^m_+$. It is no loss of generality to assume that $a_m \leq b_m$. Let $z_{m,m} = a_m, z_{i,m} = 0$ for $2 \leq i \leq m-1$, and set $z_{1,m} = b_m - a_m \geq 0$. Additionally, we set $z_{m,i} = 0$ for $1 \leq i \leq m-1$. Now, let $Z_{\text{inner}} \in \mathbb{R}^{m-1 \times m-1}_+$ be the matrix defined by the upper $(m-1) \times (m-1)$ sub-matrix of Z, and define the vectors

$$a^{\text{inner}} = [a_1 + a_m - b_m \ a_2 \ a_3 \ \cdots \ a_{m-1}]^T$$
 and $b^{\text{inner}} = [b_1 \ b_2 \ \cdots \ b_{m-1}]^T$.

Then as $a_1 \ge (1/m)\mathbf{1}^T a = (1/m)\mathbf{1}^T b \ge b_m$, we have $a^{\text{inner}} \ge 0$ and $b^{\text{inner}} \ge 0$, and moreover, $\mathbf{1}^T a^{\text{inner}} = \mathbf{1}^T a - b_m = \mathbf{1}^T b - b_m = \mathbf{1}^T b^{\text{inner}}$. In particular, we have by the inductive hypothesis that we may choose Z_{inner} such that $Z_{\text{inner}}\mathbf{1} = a^{\text{inner}}$ and $Z_{\text{inner}}^T\mathbf{1} = b^{\text{inner}}$. By inspection, setting

$$Z = \begin{bmatrix} & & & b_m - a_m \\ & & & 0 \\ & & & \vdots \\ & & & 0 \\ 0 & \cdots & 0 & & a_m \end{bmatrix}$$

we have $Z \in \mathbb{R}^{m \times m}_+$ and $Z\mathbf{1} = a$ and $Z^T\mathbf{1} = b$.

B.2. Proof of Lemma 5.3. As in our discussion in Section 5.2, we prove the lemma in a sequence of steps and auxiliary lemmas. The roadmap is as follows: first, we show that we may assume the set Ω in Lemma 5.3 has non-empty interior (§ B.2.1). With this done, we can consider any affinely independent subset of k + 1 points from Ω , whence Lemma 5.4 holds. With this done, we can cover Ω with overlapping simplices (§ B.2.2), and extend this to all of Ω (§ B.2.3).

B.2.1. It is no loss of generality to assume Ω has non-empty interior in Lemma 5.3. Let $H = \operatorname{aff} \Omega$ be the affine hull of Ω , where dim $H = l \ge 1$. (If dim H = 0, then Ω is a single point, and Lemma 5.3 is trivial.) We argue that if Lemma 5.3 holds for sets Ω such that int $\Omega \neq \emptyset$ it holds generally; thus we temporarily assume its truth for convex Ω with int $\Omega \neq \emptyset$.

Since dim H = l, we have $H = \{Av + d \mid v \in \mathbb{R}^l\}$ for some full columnrank matrix $A \in \mathbb{R}^{k \times l}$ and $d \in \mathbb{R}^k$. As Ω has non-empty interior relative to H (e.g. [17, Theorem 2.1.3]), we have $\Omega = \{Av + d \mid v \in \Omega_0\}$ for a convex set $\Omega_0 \subset \mathbb{R}^l$ with $\operatorname{int} \Omega_0 \neq \emptyset$. Defining $\tilde{f}_1(v) = f_1(Av + d)$ and $\tilde{f}_2(v) = f_2(Av + d)$, where $\tilde{f}_i(v) = \infty$ for $v \notin \Omega_0$, we have that \tilde{f}_1 and \tilde{f}_2 are order equivalent on $\Omega_0 \subset \mathbb{R}^l$. By assumption, Lemma 5.3 holds for Ω_0 , so there exist $a > 0, \tilde{b} \in \mathbb{R}^l, \tilde{c} \in \mathbb{R}$ such that

$$f_1(Av+d) = \widetilde{f}_1(v) = a\widetilde{f}_2(v) + \langle \widetilde{b}, v \rangle + c, \text{ for } v \in \Omega_0.$$

As A is full column rank, for all $t \in \Omega$ there exists a unique $v_t \in \Omega_0$ such that $t = Av_t + d$, and the mapping $t \mapsto v_t$ is linear, i.e., $v_t = Et + g$ for some $E \in \mathbb{R}^{l \times k}$ and $g \in \mathbb{R}^l$ (we may take $E \in \mathbb{R}^{l \times k}$ to be any left-inverse of A and g = -Ed, so v = Et - Eg). We obtain that $f_i(t) = f_i(Et + g)$ for i = 1, 2, whence

$$f_1(t) = \tilde{f}_1(Et+g) = a\tilde{f}_2(Et+g) + \langle \tilde{b}, Et+g \rangle + \tilde{c} = af_2(t) + \langle E^T\tilde{b}, t \rangle + \langle \tilde{b}, g \rangle + \tilde{c}$$

for all $t \in \Omega$, which is our desired result. From this point forward, we thus assume w.l.o.g. that int $\Omega \neq \emptyset$.

B.2.2. Covering sets with simplices. With Lemma 5.4 in hand, we now show that the special case for simplices is sufficient to show the general Lemma 5.3. First, we show that simplices essentially cover convex sets Ω .

LEMMA B.2. Let v, w be arbitrary points in int $\Omega \subset \mathbb{R}^k$. Then, there exist $u_0, u_1, \ldots, u_k \in \Omega$ such that $v, w \in \text{int } E$, where $E = \text{Conv}\{u_0, u_1, \ldots, u_k\}$. For any points u_2, \ldots, u_k that make v, w, u_2, \ldots, u_k affinely independent (Def. 5.2), there exist $u_0, u_1 \in \Omega$ such that $v, w \in \text{int Conv}\{u_0, u_1, \ldots, u_k\}$. Before proving Lemma B.2, we state a technical lemma about interior points of convex sets.

LEMMA B.3 (Hiriart-Urruty and Lemaréchal [17], Lemma III.2.1.6). Let $C \subset \mathbb{R}^k$ be a convex set, $u \in \text{rel int } C$ and $v \in \text{cl } C$. Then for any $\lambda \in [0, 1)$, we have $\lambda u + (1 - \lambda)v \in \text{rel int } C$.

PROOF (OF LEMMA B.2). Take $u_2, \ldots, u_k \in \Omega$ arbitrarily but in general



Fig 1: The construction in Lemma B.2.

position, so that the points v, w, u_2, \ldots, u_k and w, v, u_2, \ldots, u_k are affinely independent (Def. 5.2). Now, define $u_{\text{cent}} = \frac{1}{k+1}(v+w+\sum_{i=2}^k u_i)$, so that $u_{\text{cent}} \in \text{int }\Omega$ (Lemma B.3), and choose $\epsilon > 0$ small enough that the points $u_0 = v + \epsilon(v - u_{\text{cent}})$ and $u_1 = w + \epsilon(w - u_{\text{cent}})$ are both in Ω . (This is possible as $v, w \in \text{int }\Omega$; see Figure 1.) Then we find that $v = (u_0 + \epsilon u_{\text{cent}})/(1 + \epsilon)$ and $w = (u_1 + \epsilon u_{\text{cent}})/(1 + \epsilon)$, so that

$$u_{\text{cent}} = \frac{1}{k+1} \left(\frac{u_0}{1+\epsilon} + \frac{u_1}{1+\epsilon} + \frac{2\epsilon}{1+\epsilon} u_{\text{cent}} + \sum_{i=2}^k u_i \right),$$

or by rearranging, $u_{\text{cent}} = \lambda_0(u_0 + u_1) + \lambda_1 \sum_{i=2}^k u_i$, where

$$\lambda_0 = \left(1 - \frac{2\epsilon}{(k+1)(1+\epsilon)}\right)^{-1} \frac{1}{(k+1)(1+\epsilon)} \text{ and}$$
$$\lambda_1 = \left(1 - \frac{2\epsilon}{(k+1)(1+\epsilon)}\right)^{-1} \frac{1}{k+1}.$$

Noting that $2\lambda_0 + (k-1)\lambda_1 = 1$ and $\lambda_i > 0$, we obtain that $u_{\text{cent}} \in$ int Conv $\{u_0, \ldots, u_k\}$. As the points u_i are in general position, and as

$$v = \frac{1}{1+\epsilon}u_0 + \frac{\epsilon}{1+\epsilon}u_{\text{cent}}$$
 and $w = \frac{1}{1+\epsilon}u_1 + \frac{\epsilon}{1+\epsilon}u_{\text{cent}}$,

we have $v, w \in int \operatorname{Conv}\{u_0, \ldots, u_k\}$ by Lemma B.3.

B.2.3. Extension from a single simplex to all of Ω . We use Lemma B.2 to show the following lemma, which implies Lemma 5.3.

LEMMA B.4. In addition to the conditions of Lemma 5.3, assume that for all simplices $E \subset \operatorname{int} \Omega$ there exist $a_E > 0$, $b_E \in \mathbb{R}^k$, and $c_E \in \mathbb{R}$ such that $f_1(t) = a_E f_2(t) + b_E^T t + c_E$ for $t \in E$. Then there exist $a > 0, b \in \mathbb{R}^k$, and $c \in \mathbb{R}$ such that

$$f_1(t) = af_2(t) + b^T t + c \text{ for } t \in \Omega.$$

Coupled with Lemma 5.4, Lemma B.4 immediately yields Lemma 5.3; indeed, Lemma 5.4 shows that for any simplex E the conditions of Lemma B.4 holds, so that Lemma 5.3 follows, i.e., $f_1(t) = af_2(t) + b^T t + c$ for all $t \in \Omega$.

PROOF. For $i \in \{1, 2\}$ define the sets

$$S_i = \left\{ \{(u, v) \in \operatorname{int} \Omega \times \operatorname{int} \Omega \mid \nabla f_i(u) \neq \nabla f_i(v) \} \\ \cup \{(u, v) \in \operatorname{int} \Omega \times \operatorname{int} \Omega \mid \nabla f_i(u) \text{ or } \nabla f_i(v) \text{ does not exist} \} \right\} \subset \Omega \times \Omega.$$

We divide our discussion into two cases.

Case 1. First we suppose that $S_1 = S_2 = \emptyset$. Then for $i = 1, 2, f_i$ are differentiable in int Ω in this case, we have that $\nabla f_i(u) = \nabla f_i(v)$ for all $u, v \in \operatorname{int} \Omega$. Then by continuity of the f_i on $\operatorname{int} \Omega$, we must have $f_i(t) = b_i^T t + c_i$ for i = 1, 2. The result follows by taking $a = 1, b = b_1 - b_2$ and $c = c_1 - c_2$ and applying Lemma 5.5.

Case 2. We have that at least one of S_1 and S_2 is non-empty. Without loss of generality, say $S_1 \neq \emptyset$. Choose a pair $(u^*, v^*) \in S_1$ and consider the collection of sets

$$\mathcal{M} = \{ E = \operatorname{Conv}\{u_0, u_1, \dots, u_k\} \mid u^*, v^* \in \operatorname{int} E \text{ and } u_0, u_1, \dots, u_k \in \Omega \}.$$

We show that for any $E_1, E_2 \in \mathcal{M}$, we have $a_{E_1} = a_{E_2}, b_{E_1} = b_{E_2}$ and $c_{E_1} = c_{E_2}$. We know that int $E_1 \cap \text{int } E_2 \neq \emptyset$, as $u^* \in \text{int } E$ for all $E \in \mathcal{M}$.

Now, assume for the sake of contradiction that $a_{E_1} \neq a_{E_2}$. For all $t \in E_1 \cap E_2$ we have

(33)
$$f_1(t) = a_{E_1}f_2(t) + b_{E_1}^T t + c_{E_1}$$
 and $f_1(t) = a_{E_2}f_2(t) + b_{E_2}^T t + c_{E_2}$.

By subtracting the preceding equations from one another after multiplying by a_{E_2} and a_{E_1} , respectively, one obtains that for $t \in E_1 \cap E_2$,

$$f_1(t) = \frac{1}{a_{E_2} - a_{E_1}} \left[(a_{E_2}b_{E_1} - a_{E_1}b_{E_2})^T t + (a_{E_2}c_{E_1} - a_{E_1}c_{E_2}) \right].$$

This yields a contradiction, since we have assumed that either (i) $\nabla f_1(u^*)$ or $\nabla f_2(v^*)$ does not exist or (ii) $\nabla f_1(u^*) \neq \nabla f_1(v^*)$, while $u^*, v^* \in \operatorname{int} E_1 \cap$ int $E_2 = \operatorname{int}(E_1 \cap E_2)$. Thus $a_{E_1} = a_{E_2}$. To obtain $b_{E_1} = b_{E_2}$, note that $\operatorname{int}(E_1 \cap E_2) \neq \emptyset$, as $u^*, v^* \in \operatorname{int}(E_1 \cap E_2)$. Subtracting the equalities (33), we find $0 = f_1(t) - f_1(t) = (b_{E_1} - b_{E_2})^T t + c_{E_1} - c_{E_2}$ for $t \in E_1 \cap E_2$, which has non-empty interior. That $b_{E_1} = b_{E_2}$ immediately implies $c_{E_1} = c_{E_2}$. Hence, there exist some $a > 0, b \in \mathbb{R}^k$, and $c \in \mathbb{R}$ such that for all sets $E \in \mathcal{M}$, we have $a_E = a, b_E = b$, and $c_E = c$.

We complete the proof by showing that $\bigcup \{E \mid E \in \mathcal{M}\}$ is dense in Ω . Define $\Omega^{\circ} \subset \Omega$ as

$$\Omega^{\circ} = \left\{ t \in \Omega \mid \begin{array}{c} \text{there exist } u_3, \dots, u_k \\ \text{such that } u^{\star}, v^{\star}, t, u_3, \dots, u_k \text{ are affinely independent} \end{array} \right\}$$

The set Ω° forms a dense subset of Ω , as $u^{\star} \in \operatorname{int} \Omega \neq \emptyset$. For any $t \in \Omega^{\circ}$, Lemma B.2 guarantees the existence of $E \in \mathcal{M}$ such that $t \in E$ and $u^{\star}, v^{\star} \in \operatorname{int} E$. We thus have $\Omega^{\circ} \subset \bigcup \{E \mid E \in \mathcal{M}\}$. As $f_1(t) = af_2(t) + b^T t + c$ for all $t \in \Omega^{\circ}$ by the previous paragraph, Lemma 5.5 allows us to extend the equality to $f_1(t) = af_2(t) + b^T t + c$ for all $t \in \Omega$.

B.3. Proofs of auxiliary lemmas for Lemma 5.4.

B.3.1. Proof of Lemma 5.6. For each *i*, let $\lambda_i = [\alpha_i]_+$ and $\nu_i = [\alpha_i]_-$ be the positive and negative parts of the α_i , and let $\lambda_i = \frac{r_i}{s}$ and $-\nu_i = \frac{q_i}{s}$ where $q_i, r_i, s \in \mathbb{N}$. Then we have

$$\sum_{i=1}^{m} \lambda_i u_i = v - \sum_{i=1}^{m} \nu_i u_i \text{ or } \sum_{i=1}^{m} r_i u_i = sv + \sum_{i=1}^{m} q_i u_i,$$

where $\mathbf{1}^T r = s + \mathbf{1}^T q$ as $\mathbf{1}^T \alpha = 1$. Defining the matrices

$$A = [\underbrace{u_1 \cdots u_1}_{r_1 \text{ times}} \cdots \underbrace{u_m \cdots u_m}_{r_m \text{ times}}] \text{ and } B = [\underbrace{v \cdots v}_{s \text{ times}} \underbrace{u_1 \cdots u_1}_{q_1 \text{ times}} \cdots \underbrace{u_m \cdots u_m}_{q_m \text{ times}}]$$

we have $A\mathbf{1} = B\mathbf{1}$ with columns in Ω . Then order equivalence (24) implies

$$\sum_{i=1}^{m} r_i f_1(u_i) \le s f_1(v) + \sum_{i=1}^{m} q_i f_1(u_i) \quad \text{iff} \quad \sum_{i=1}^{m} r_i f_2(u_i) \le s f_2(v) + \sum_{i=1}^{m} q_i f_2(u_i)$$

Each of these is equivalent to $\sum_{i=1}^{m} \alpha_i f_j(u_i) \leq f_j(v)$ for j = 1, 2.

B.3.2. Proof of Lemma 5.7. We assume that f_1 and f_2 are non-linear over Conv $\{u_0, \ldots, u_k\}$, as otherwise the result is trivial. Let the vectors g_1 and g_2 and matrix $H \in \mathbb{R}^{k \times k}$ be defined by

$$H = \begin{bmatrix} u_1^T - u_0^T \\ \vdots \\ u_k^T - u_0^T \end{bmatrix}, \quad g_1 = \begin{bmatrix} f_1(u_1) - f_1(u_0) \\ \vdots \\ f_1(u_k) - f_1(u_0) \end{bmatrix}, \quad g_2 = \begin{bmatrix} f_2(u_1) - f_2(u_0) \\ \vdots \\ f_2(u_k) - f_2(u_0) \end{bmatrix}.$$

For any a > 0, define the vector $b(a) \in \mathbb{R}^k$ so that

$$b(a)^{T}(u_{i} - u_{0}) = f_{1}(u_{i}) - f_{1}(u_{0}) - a(f_{2}(u_{i}) - f_{2}(u_{0})) = g_{1,i} - ag_{2,i},$$

that is, as $b(a) = H^{-1}(g_1 - ag_2)$, which is possible as H is full rank. By choosing $c(a) = f_1(u_0) - af_2(u_0) - b(a)^T u_0$, we then obtain that

$$f_1(u_i) = a f_2(u_i) + b(a)^T u_i + c(a)$$

by algebraic manipulations. We now consider u_{cent} . We have

$$b(a)^{T}u_{\text{cent}} = \frac{1}{k+1} \sum_{i=0}^{k} b(a)^{T}u_{i} = b(a)^{T}u_{0} + \frac{1}{k+1} \mathbf{1}^{T}Hb(a)$$
$$= b(a)^{T}u_{0} + \frac{1}{k+1} \mathbf{1}^{T}(g_{1} - ag_{2}),$$

so that

$$af_{2}(u_{\text{cent}}) + b(a)^{T}u_{\text{cent}} + c(a)$$

= $af_{2}(u_{\text{cent}}) + \sum_{i=1}^{k} \frac{f_{1}(u_{i}) - f_{1}(u_{0}) + af_{2}(u_{0}) - af_{2}(u_{i})}{k+1} + f_{1}(u_{0}) - af_{2}(u_{0})$
= $af_{2}(u_{\text{cent}}) + \frac{1}{k+1} \sum_{i=0}^{k} f_{1}(u_{i}) - a\frac{1}{k+1} \sum_{i=0}^{k} f_{2}(u_{i}).$

Thus we may choose an a > 0 such that our desired equalities hold if and only if there exists a > 0 such that

$$f_1(u_{\text{cent}}) - \frac{1}{k+1} \sum_{i=0}^k f_1(u_i) = a f_2(u_{\text{cent}}) - a \frac{1}{k+1} \sum_{i=0}^k f_2(u_i).$$

By the assumption that f_1 and f_2 are non-linear, we have that (by Lemma 5.8) $f_1(u_{\text{cent}}) < \frac{1}{k+1} \sum_{i=0}^k f_1(u_i)$ and $f_2(u_{\text{cent}}) < \frac{1}{k+1} \sum_{i=0}^k f_2(u_i)$. Thus, setting

$$a = \frac{f_1(u_{\text{cent}}) - \frac{1}{k+1} \sum_{i=0}^k f_1(u_i)}{f_2(u_{\text{cent}}) - \frac{1}{k+1} \sum_{i=0}^k f_2(u_i)} > 0$$

gives the desired result.

B.3.3. Proof of Lemma 5.8. Without loss of generality, we assume that $\lambda_1 = \|\lambda\|_{\infty}$ and that $\lambda_1 > 1/m$. Then we have

(34)
$$f\left(\sum_{i=1}^{m} \lambda_{i} u_{i}\right) \leq \sum_{i=1}^{m} \lambda_{i} f(u_{i}) = \lambda_{1} \sum_{i=1}^{m} f(u_{i}) + \sum_{i=1}^{m} (\lambda_{i} - \lambda_{1}) f(u_{i})$$
$$= m \lambda_{1} f(u_{\text{cent}}) + \sum_{i=2}^{m} (\lambda_{1} - \lambda_{i}) f(u_{i}).$$

Let $\Lambda = \sum_{i=1}^{m} (\lambda_1 - \lambda_i) = m\lambda_1 - 1 > 0$. Then

$$u_{\text{cent}} = \frac{1}{m\lambda_1} \sum_{i=1}^m \lambda_i u_i + \frac{\Lambda}{m\lambda_1} \sum_{i=1}^m \frac{\lambda_1 - \lambda_i}{\Lambda} u_i,$$

and we have

$$m\lambda_{1}f(u_{\text{cent}}) \leq n\lambda_{1} \left[\frac{1}{m\lambda_{1}} f\left(\sum_{i=1}^{m} \lambda_{i} u_{i}\right) + \frac{\Lambda}{m\lambda_{1}} f\left(\sum_{i=2}^{m} \frac{\lambda_{1} - \lambda_{i}}{\Lambda} u_{i}\right) \right]$$
$$\leq f\left(\sum_{i=1}^{m} \lambda_{i} u_{i}\right) + \Lambda \sum_{i=2}^{m} \frac{\lambda_{1} - \lambda_{i}}{\Lambda} f(u_{i}).$$

The inequalities (34) must have been equalities, giving the result.

APPENDIX C: PROOFS FOR BAYES CONSISTENCY OF EMPIRICAL RISK MINIMIZATION **C.1. Proof of Lemma 4.1.** With the definition $R^*(q) := \inf_{\gamma} R(\gamma \mid q)$, we have

(35)
$$R(\gamma \mid \mathbf{q}) - \inf_{\gamma \in \Gamma, \mathbf{q} \in \mathbf{Q}} R(\gamma \mid \mathbf{q}) = R(\gamma \mid \mathbf{q}) - R^{\star}(\mathbf{q}) + R^{\star}(\mathbf{q}) - \inf_{\mathbf{q} \in \mathbf{Q}} R^{\star}(\mathbf{q}).$$

For the second two terms, we note that by Theorem 1, we have

$$R^{\star}(\mathsf{q}) = \mathbb{E}[H_{L^{\mathrm{cw}}}(\widetilde{\pi}(\mathsf{q}(X)))] = a\mathbb{E}[H_{L}(\widetilde{\pi}(\mathsf{q}(X)))] + b^{T}\pi + c,$$

and similarly for $\inf_{\mathbf{q}} R^{\star}(\mathbf{q})$, so that

$$R^{\star}(\mathbf{q}) - \inf_{\mathbf{q}\in\mathbf{Q}} R^{\star}(\mathbf{q}) = a \left[R_{L}^{\star}(\mathbf{q}) - \inf_{\mathbf{q}\in\mathbf{Q}} R_{L}^{\star}(\mathbf{q}) \right]$$
$$\leq a \left[R_{L}(\gamma \mid \mathbf{q}) - \inf_{\gamma\in\Gamma,\mathbf{q}\in\mathbf{Q}} R_{L}(\gamma \mid \mathbf{q}) \right]$$

Clearly $t \mapsto at$ is concave, so that it remains to bound $R(\gamma \mid q) - R^*(q)$ in expression (35). To that end, let $L(\alpha) \in \mathbb{R}^k$ denote the vector of losses $L(\alpha, y)$ and define the function

$$H(\epsilon) = \inf_{\pi \in \Delta_k, \alpha} \left\{ \sup_{\alpha'} \pi^T \left(L(\alpha) - L(\alpha') \right) \mid \sup_{\alpha'} \pi^T \left(L^{cw}(\alpha) - L^{cw}(\alpha') \right) \ge \epsilon \right\}$$

and let H^{**} be its Fenchel biconjugate. Then (see [36, Proposition 25 and Corollary 26], as well as the papers [31, 30, 12, Proposition 1]) we have that $H^{**}(\epsilon) > 0$ for all $\epsilon > 0$, and defining $g(\epsilon) = \sup\{\delta : \delta \ge 0, H^{**}(\delta) \le \epsilon\}$ yields the desired concave function by taking h(t) = g(t) + at. In passing, we note that we may w.l.o.g. replace h(t) with min $\{h(t), \max_{ij} c_{ij}\}$.

Now we give the second result. Without loss of generality, we may assume that the vector $\gamma(z) \in \mathbb{R}^k$ has a unique maximal coordinate; we may otherwise assume a deterministic rule for breaking ties. Let $y(\gamma(z)) = \operatorname{argmax}_i \gamma_i(z)$, assumed w.l.o.g. to be unique. Consider that

$$R(\gamma \mid \mathbf{q}) - R^{\star}(\mathbf{q}) = \int \sup_{\alpha} \sum_{i=1}^{k} \left[\widetilde{\pi}_{i}(z) L^{\mathrm{cw}}(\gamma(z), i) - L^{\mathrm{cw}}(\alpha, i) \right] d\overline{P}^{\mathbf{q}}(z)$$

where $\overline{P}^{\mathsf{q}}(A) = \sum_{i=1}^{k} \pi_i P_i(\mathsf{q}^{-1}(A))$ (for measurable $A \subset \mathcal{Z}$) is the pushforward of q and $\widetilde{\pi}(z) = [\pi_i dP_i^{\mathsf{q}}(z)]_{i=1}^k / d\overline{P}^{\mathsf{q}}(z)$ is shorthand for the posterior of Y conditional on observing $z \in \mathsf{q}(\mathcal{X})$. Lemma A.9 immediately implies

$$R(\gamma \mid \mathbf{q}) - R^{\star}(\mathbf{q}) \leq \int \sup_{\alpha} \sum_{i=1}^{k} \left[\widetilde{\pi}_{i}(z) L(\gamma(z), i) - L(\alpha, i) \right] d\overline{P}^{\mathbf{q}}(z)$$
$$= R_{L}(\gamma \mid \mathbf{q}) - R_{L}^{\star}(\mathbf{q})$$

for either of the hinge-type losses, e.g. $L(\alpha, y) = \sum_{i=1}^{k} c_{yi} [1 + \alpha_i - \alpha_y]_+$. In the notation of the previous (general) case, we have a = 1/k because $H_L(\pi) = \frac{1}{k} H_{L^{cw}}(\pi)$, whence we may take h(t) = (1 + 1/k)t.

C.2. Proof of Theorem 3. The proof of Theorem 3 is almost immediate from Lemma 4.1. Indeed, by Lemma 4.1, we have

$$R(\gamma_n \mid \mathsf{q}_n) - R^{\star}(\mathsf{Q}) \le h \left(R_L(\gamma_n \mid \mathsf{q}_n) - R_L^{\star}(\mathsf{Q}) \right)$$

for some *h* concave, bounded, and satisfying h(0) = 0. It is thus sufficient to show that $\mathbb{E}[R_L(\gamma_n | \mathbf{q}_n) - R_L^{\star}(\mathbf{Q})] \to 0$. Now, if we let γ_n^{\star} and \mathbf{q}_n^{\star} minimize $R_L(\gamma | \mathbf{q})$ over the set $\Gamma_n \times \mathbf{Q}_n$ (or be arbitrarily close to minimizing the *L*-risk), we have

$$R_{L}(\gamma_{n} \mid \mathbf{q}_{n}) - R_{L}(\gamma_{n}^{\star} \mid \mathbf{q}_{n}^{\star})$$

$$= R_{L}(\gamma_{n} \mid \mathbf{q}_{n}) - \hat{R}_{L,n}(\gamma_{n} \mid \mathbf{q}_{n}) + \hat{R}_{L,n}(\gamma_{n} \mid \mathbf{q}_{n}) - R_{L}(\gamma_{n}^{\star} \mid \mathbf{q}_{n}^{\star})$$

$$\leq \underbrace{R_{L}(\gamma_{n} \mid \mathbf{q}_{n}) - \hat{R}_{L,n}(\gamma_{n} \mid \mathbf{q}_{n}) + \hat{R}_{L,n}(\gamma_{n}^{\star} \mid \mathbf{q}_{n}^{\star}) - R_{L}(\gamma_{n}^{\star} \mid \mathbf{q}_{n}^{\star})}_{\leq 2 \sup_{\gamma \in \Gamma_{n}, \mathbf{q} \in \mathbf{Q}_{n}} |\hat{R}_{L,n}(\gamma \mid \mathbf{q}) - R_{L}(\gamma \mid \mathbf{q})|}$$

$$+ \hat{R}_{L,n}(\gamma_{n} \mid \mathbf{q}_{n}) - \inf_{\gamma \in \Gamma_{n}, \mathbf{q} \in \mathbf{Q}_{n}} \hat{R}_{L,n}(\gamma \mid \mathbf{q}).$$

Consequently, we have the expectation bound

$$\mathbb{E}\left[R_L(\gamma_n \mid \mathbf{q}_n) - R_L^{\star}(\mathbf{Q})\right] \\ \leq \mathbb{E}\left[R_L(\gamma_n \mid \mathbf{q}_n) - R_L(\gamma_n^{\star} \mid \mathbf{q}_n^{\star})\right] + R_L(\gamma_n^{\star} \mid \mathbf{q}_n^{\star}) - R_L^{\star}(\mathbf{Q}) \leq 2\epsilon_n^{\text{est}} + \epsilon_n^{\text{opt}} + \epsilon_n^{\text{app}},$$

which converges to zero as desired.

APPENDIX D: PROOFS OF BASIC PROPERTIES OF F-DIVERGENCES

In this section, we collect the proofs of the characterizations of generalized f-divergences (Def. 2.1).

D.1. Proof of Lemma 2.1. Let μ_1 and μ_2 be dominating measures; then $\mu = \mu_1 + \mu_2$ also dominates P_1, \ldots, P_k as well as μ_1 and μ_2 . We have for $\nu = \mu_1$ or $\nu = \mu_2$ that

$$\frac{dP_i/d\nu}{dP_k/d\nu} = \frac{dP_i/d\nu}{dP_k/d\nu} \frac{d\nu/d\mu}{d\nu/d\mu} = \frac{dP_i/d\mu}{dP_k/d\mu} \text{ and } \frac{dP_i}{d\nu} \frac{d\nu}{d\mu} = \frac{dP_i}{d\mu},$$

the latter two equalities holding μ -almost surely by definition of the Radon-Nikodym derivative. Thus we obtain for $\nu = \mu_1$ or $\nu = \mu_2$ that

$$\int f\left(\frac{dP_{1:k-1}/d\nu}{dP_k/d\nu}\right) \frac{dP_k}{d\nu} d\nu = \int f\left(\frac{dP_1/d\nu}{dP_k/d\nu}, \dots, \frac{dP_{k-1}/d\nu}{dP_k/d\nu}\right) \frac{dP_k}{d\nu} \frac{d\nu}{d\mu} d\mu$$
$$= \int f\left(\frac{dP_1/d\mu}{dP_k/d\mu}, \dots, \frac{dP_{k-1}/d\mu}{dP_k/d\mu}\right) \frac{dP_k}{d\mu} d\mu,$$

again by definition of the Radon-Nikodym derivative. We have $\frac{p_i}{p_k} = \frac{dP_i/d\mu}{dP_k/d\mu}$ a.s.- μ , which shows that the base measure μ does not affect the integral. To see the positivity, we may take $\mu = \frac{1}{k} \sum_{i=1}^{k} P_i$, in which case Jensen's

inequality implies (the perspective function (9) is convex) that

$$D_f(P_1, \dots, P_{k-1} \| P_k) = \mathbb{E} \left[f\left(\frac{dP_1}{dP_k}(X), \dots, \frac{dP_{k-1}}{dP_k}(X)\right) dP_k(X) \right]$$
$$\geq f\left(\frac{\mathbb{E}[dP_1]}{\mathbb{E}[dP_k]}, \dots, \frac{\mathbb{E}[dP_{k-1}]}{\mathbb{E}[dP_k]}\right) \mathbb{E}[dP_k] = f(\mathbf{1}) = 0,$$

where the expectation is taken under the distribution μ . The inequality is strict for f strictly convex at 1 as long as dP_i/dP_k is non-constant for some *i*, meaning that there exists an *i* such that $P_i \neq P_k$.

D.2. Proof of Proposition 1. Before proving the proposition, we first establish a more general continuity result for f-divergences. This result is a direct generalization of results of [34, Thm. 5]. Given a sub- σ -algebra $\mathcal{G} \subset \mathcal{F}$, we let $P^{\mathcal{G}}$ denote the restriction of the measure P, defined on \mathcal{F} , to \mathcal{G} .

LEMMA D.1. Let $\mathcal{F}_1 \subset \mathcal{F}_2 \subset \ldots$ be a sequence of sub- σ -algebras of \mathcal{F} and let $\mathcal{F}_{\infty} = \sigma(\bigcup_{n \ge 1} \mathcal{F}_n)$. Then $D_f\left(P_1^{\mathcal{F}_n}, \dots, P_{k-1}^{\mathcal{F}_n} \| P_k^{\mathcal{F}_n}\right)$ is non-decreasing in n and

$$\lim_{n \to \infty} D_f \left(P_1^{\mathcal{F}_n}, \dots, P_{k-1}^{\mathcal{F}_n} \| P_k^{\mathcal{F}_n} \right) = D_f \left(P_1^{\mathcal{F}_\infty}, \dots, P_{k-1}^{\mathcal{F}_\infty} \| P_k^{\mathcal{F}_\infty} \right).$$

PROOF. Define the measure $\nu = \frac{1}{k} \sum_{i=1}^{k} P_i$ and vectors V^n via the Radon-Nikodym derivatives

$$V^{n} = \frac{1}{k} \cdot \left(\frac{dP_{1}^{\mathcal{F}_{n}}}{d\nu^{\mathcal{F}_{n}}}, \frac{dP_{2}^{\mathcal{F}_{n}}}{d\nu^{\mathcal{F}_{n}}}, \dots, \frac{dP_{k-1}^{\mathcal{F}_{n}}}{d\nu^{\mathcal{F}_{n}}} \right).$$

Then $(1 - \mathbf{1}^T V^n) d\nu^{\mathcal{F}_n} = \frac{1}{k} dP_k^{\mathcal{F}_n}$, and V^n is a martingale adapted to the filtration \mathcal{F}_n by standard properties of conditional expectation (under the measure ν). Letting $C_k = \{v \in \mathbb{R}^{k-1} \mid \mathbf{1}^T v \leq 1\}$, we define $g : C_k \to \mathbb{R}$ by

$$g(v) = f\left(\frac{v}{1 - \mathbf{1}^T v}\right) (1 - \mathbf{1}^T v)$$

We see that g is convex (it is a perspective function), and we have

$$\mathbb{E}_{\nu}[g(V^n)] = \int f\left(\frac{dP_{1:k-1}^{\mathcal{F}_n}}{dP_k^{\mathcal{F}_n}}\right) \left(1 - \frac{1}{k}\sum_{i=1}^{k-1}\frac{dP_i^{\mathcal{F}_n}}{d\nu^{\mathcal{F}_n}}\right) d\nu = \frac{1}{k}D_f\left(P_{1:k-1}^{\mathcal{F}_n} \| P_k^{\mathcal{F}_n}\right).$$

Because $V^n \in C_k$ for all n, we see that $g(V^n)$ is a submartingale. This gives the first result of the lemma, that $D_f(P_{1:k-1}^{\mathcal{F}_n} \| P_k^{\mathcal{F}_n})$ is non-decreasing in n.

Now, assume that the limit in the second statement is finite, as otherwise the result is trivial. Using $f(\mathbf{1}) = 0$, we have by convexity that for $v \in C_k$,

$$g(v) = f\left(\frac{v}{1-\mathbf{1}^T v}\right) (1-\mathbf{1}^T v) + f(1)\mathbf{1}^T v \ge f\left(v+1(\mathbf{1}^T v)\right)$$
$$\ge \inf_{v \in C_k} f(v+1(\mathbf{1}^T v)) > -\infty,$$

the final inequality a consequence of the fact that f is closed and hence attains its infimum. In particular, the sequence $g(V^n) - \inf_{v \in C_k} g(v)$ is a non-negative submartingale, and thus

$$\sup_{n} \mathbb{E}_{\nu} \left[|g(V^{n}) - \inf_{v \in C_{k}} g(v)| \right] = \lim_{n} \mathbb{E}_{\nu} \left[g(V^{n}) - \inf_{v \in C_{k}} g(v) \right] < \infty.$$

With this integrability guarantee, Doob's second martingale convergence theorem [7, Thm. 35.5] yields the existence of a vector $V^{\infty} \in \mathcal{F}_{\infty}$ such that

$$0 \le \lim_{n} \mathbb{E}_{\nu} \left[g(V^{n}) - \inf_{v \in C_{k}} g(v) \right] = \mathbb{E}_{\nu}[g(V^{\infty})] - \inf_{v \in C_{k}} g(v) < \infty.$$

That $\inf_{v \in C_k} g(v) > -\infty$ implies $\mathbb{E}_{\nu}[|g(V^{\infty})|] < \infty$, giving the lemma. \Box

We now give the proof of Proposition 1 proper. Let the base measure $\mu = \frac{1}{k} \sum_{i=1}^{k} P_i$ and let $p_i = \frac{dP_i}{d\mu}$ be the associated densities of the P_i . Define the increasing sequence of partitions \mathcal{P}^n of \mathcal{X} by sets $A_{\alpha n}$ for vectors $\alpha = (\alpha_1, \ldots, \alpha_{k-1})$ and B with

$$A_{\alpha n} = \left\{ x \in \mathcal{X} \mid \frac{\alpha_j - 1}{2^n} \le \frac{p_j(x)}{p_k(x)} < \frac{\alpha_j}{2^n} \text{ for } j = 1, \dots, k - 1 \right\}$$

where we let each α_j range over $\{-n2^n, -n2^n+1, \ldots, n2^n\}$, and define and $B = (\bigcup_{\alpha} A_{\alpha n})^c = \mathcal{X} \setminus \bigcup_{\alpha} A_{\alpha n}$. Then we have

$$\left(\frac{p_1(x)}{p_k(x)},\ldots,\frac{p_{k-1}(x)}{p_k(x)}\right) = \lim_{n \to \infty} \sum_{\alpha \in \{-n2^n,\ldots,n2^n\}^{k-1}} \frac{\alpha}{2^n} \chi_{A_{\alpha n}}(x) + (n,\ldots,n) \chi_B(x)$$

where χ_A denotes the $\{0,1\}$ -valued characteristic function of the set A. Each term on the right-hand-side of the previous display is \mathcal{F}_n -measurable, where \mathcal{F}_n denotes the sub- σ -field generated by the partition \mathcal{P}^n . Defining $\mathcal{F}_{\infty} = \sigma(\bigcup_{n \ge 1} \mathcal{F}_n)$, we have

$$D_f(P_{1:k-1} \| P_k | \mathcal{P}^n) = D_f\left(P_1^{\mathcal{F}_n}, \dots, P_{k-1}^{\mathcal{F}_n} \| P_k^{\mathcal{F}_n}\right)$$

$$\to D_f\left(P_1^{\mathcal{F}_\infty}, \dots, P_{k-1}^{\mathcal{F}_\infty} \| P_k^{\mathcal{F}_\infty}\right) = D_f(P_1, \dots, P_{k-1} \| P_k),$$

where the limiting operation follows by Lemma D.1 and the final equality because of the measurability containment $(\frac{p_1}{p_k}, \ldots, \frac{p_{k-1}}{p_k}) \in \mathcal{F}_{\infty}$.

D.3. Proof of Proposition 2. Before proving the proposition, we state an inequality that generalizes the classical log-sum inequality (cf. [9, Theorem 2.7.1]).

LEMMA D.2. Let $f : \mathbb{R}^k_+ \to \overline{\mathbb{R}}$ be convex. Let $a : \mathcal{X} \to \mathbb{R}_+$ and $b_i : \mathcal{X} \to \mathbb{R}_+$ \mathbb{R}_+ for $1 \leq i \leq k$ be nonnegative measurable functions. Then for any finite measure μ defined on \mathcal{X} , we have

$$\left(\int ad\mu\right) f\left(\frac{\int b_1 d\mu}{\int ad\mu}, \dots, \frac{\int b_k d\mu}{\int ad\mu}\right) \leq \int a(x) f\left(\frac{b_1(x)}{a(x)}, \dots, \frac{b_k(x)}{a(x)}\right) d\mu(x).$$

PROOF. Recall that the perspective function g(y,t) defined by g(y,t) =tf(y/t) for t > 0 is jointly convex in y and t. The measure $\nu = \mu/\mu(\mathcal{X})$ defines a probability measure, so that for $X \sim \nu$, Jensen's inequality implies

$$\left(\int ad\nu\right) f\left(\frac{\int b_1 d\nu}{\int ad\nu}, \dots, \frac{\int b_k d\nu}{\int ad\nu}\right) \leq \mathbb{E}_{\nu} \left[a(X) f\left(\frac{b_1(X)}{a(X)}, \dots, \frac{b_k(X)}{a(X)}\right)\right]$$
$$= \frac{1}{\mu(\mathcal{X})} \int a(x) f\left(\frac{b_1(x)}{a(x)}, \dots, \frac{b_k(x)}{a(x)}\right) d\mu(x).$$
Noting that $\int b_i d\nu / \int ad\nu = \int b_i d\mu / \int ad\mu$ gives the result.

Noting that $\int b_i d\nu / \int a d\nu = \int b_i d\mu / \int a d\mu$ gives the result.

We use this lemma and Proposition 1 to give Proposition 2. Proposition 1 implies

$$D_f(Q_{P_1},\ldots,Q_{P_{k-1}} \| Q_{P_k}) = \sup_{\mathcal{P}} \left\{ D_f(Q_{P_1},\ldots,Q_{P_{k-1}} \| Q_{P_k} \mid \mathcal{P}) : \mathcal{P} \text{ is a finite partition of } \mathcal{Z} \right\}.$$

It is consequently no loss of generality to assume that \mathcal{Z} is finite and $\mathcal{Z} = \{1, 2, \ldots, m\}$. Let $\mu = \sum_{i=1}^{k} P_i$ be a dominating measure and let $p_i = dP_i/d\mu$. Letting $q(j \mid x) = Q(\{j\} \mid x)$ and $q_{P_i}(j) = Q_{P_i}(\{j\})$, we obtain

$$D_f(Q_{P_1}, \dots, Q_{P_{k-1}} \| Q_{P_k}) = \sum_{i=1}^m q_{P_k}(i) f\left(\frac{q_{P_1}(i)}{q_{P_k}(i)}, \dots, \frac{q_{P_{k-1}}(i)}{q_{P_k}(i)}\right)$$
$$= \sum_{i=1}^m \left(\int_{\mathcal{X}} q(i \mid x) p_k(x) d\mu(x)\right) f\left(\frac{\int_{\mathcal{X}} q(i \mid x) [p_1(x), \dots, p_{k-1}(x)] d\mu(x)}{\int_{\mathcal{X}} q(i \mid x) p_k(x) d\mu(x)}\right)$$
$$\leq \sum_{i=1}^m \int_{\mathcal{X}} q(i \mid x) f\left(\frac{p_1(x)}{p_k(x)}, \dots, \frac{p_{k-1}(x)}{p_k(x)}\right) p_k(x) d\mu(x)$$

by Lemma D.2. Noting that $\sum_{i=1}^{m} q(i \mid x) = 1$, we obtain our desired result.

STANFORD UNIVERSITY STANFORD, CALIFORNIA 94305 E-MAIL: jduchi@stanford.edu khosravi@stanford.edu fengruan@stanford.edu