

Permuting Incomplete Paired Data: A Novel Exact and Asymptotic Correct Randomization Test

Lubna Amro* and Markus Pauly*

July 15, 2018

Abstract

Various statistical tests have been developed for testing the equality of means in matched pairs with missing values. However, most existing methods are commonly based on certain distributional assumptions such as normality, 0-symmetry or homoscedasticity of the data. The aim of this paper is to develop a statistical test that is robust against deviations from such assumptions and also leads to valid inference in case of heteroscedasticity or skewed distributions. This is achieved by applying a novel randomization approach. The resulting test procedure is not only shown to be asymptotically correct but is also finitely exact if the distribution of the data is invariant with respect to the considered randomization group. Its small sample performance is further studied in an extensive simulation study and compared to existing methods. Finally, an illustrative data example is analyzed.

Keywords: Missing Values; Permutation Tests; Randomization Tests; Student's t -test; Welch test.

* University of Ulm, Institute of Statistics, Germany
email: markus.pauly@uni-ulm.de, lubna.amro@uni-ulm.de

1 Motivation and introduction

Matched pairs designs are concerned with scientific experiments in which subjects are observed repeatedly under two different treatments or time points. Testing equality of means of such data, where some of the components are randomly missing is a problem usually encountered in practice. In previous years, several statistical methods have been proposed for dealing with this issue. Here, the most simplest suggestion is to only work with the completely observed pairs and to carry out the well-known paired t -test. The main problem of this approach is that it discards all information from the incomplete observations. This will typically lead to a power loss. Moreover, the test can only be applied when the data are assumed to be missing completely at random; see the monograph of Little and Rubin [2014] for an introduction and discussion on the different missingness mechanisms. One possibility to overcome this problem might be to apply multiple-imputation techniques, see e.g. Verbeke and Molenberghs [2009]. However these techniques often require large sample sizes for being correct, see e.g. the discussion in Akritas et al. [2002]. Other possibilities that use all observed information in the matched pairs design (but not more) include the statistical tests proposed in Lin and Stivers [1974], Ekbohm [1976], Bhoj [1978], Looney and Jones [2003], Kim et al. [2005], and Samawi and Vogel [2014]. All of these tests, however, rely on specific model assumptions such as symmetry or even bivariate normality. Especially the latter is typically not met in practice but crucial for relatively small sample sizes. If these specific assumptions are not met the corresponding tests might possess possibly inflated type-I error control.

Therefore, the aim of the present paper is to provide a statistical test for partially paired data which does not require any parametric assumptions, possesses nice small and large sample properties and (only) uses all observed information. This is achieved by applying a novel studentized randomization approach to a weighted test statistic involving both, the paired t -test as well as the Welch test statistic. The resulting randomization test combines recent findings on studentized permutation tests for complete observations in unpaired and paired designs, see Janssen [1997], Janssen [1999a], Janssen [1999b], Neubert and Brunner [2007], Pauly [2011], Konietzschke and Pauly [2012], Omelka and Pauly [2012], Chung and Romano [2013], DiCiccio and Romano [2015], Pauly et al. [2015] and Chung and Romano [2016]. The application of the studentized randomization approach to incomplete data is original to the present paper.

To formulate the concrete testing problem, let us consider a general matched pairs design given by i.i.d random vectors

$$\mathbf{X}_j = \begin{pmatrix} X_{1j} \\ X_{2j} \end{pmatrix}, \quad j = 1, \dots, n, \quad (1.1)$$

with mean vector $E(\mathbf{X}_1) = \boldsymbol{\mu} = (\mu_1, \mu_2)' \in \mathbb{R}^2$ and an arbitrary covariance matrix $Cov(\mathbf{X}_1) = \boldsymbol{\Sigma} > 0$. Within Model (1.1) we now like to test the null hypothesis $H_0 : \{\mu_1 = \mu_2\}$. Here, some of the random vectors components might be missing completely at random as described in the next section.

This paper is organized as follows: In Section 2 we introduce the statistical model for incomplete paired data and a weighted test statistic. A permutation test based on this statistic is introduced in Section 3, where also its theoretical properties are explained. We review the existing methods for incomplete paired data in Section 4 to choose adequate competitors to investigate together with our novel procedure in extensive simulations given in Section 5. Finally, a real data example is considered in Section 6. All proofs are shifted to the Appendix.

2 Statistical model, hypotheses and statistics

When some of the components of the matched pairs are missing completely at random, we may sort the data into complete and incomplete observed random vectors and rewrite Model (1.1) with independent random variables as

$$\underbrace{\begin{pmatrix} X_{11}^{(c)} \\ X_{21}^{(c)} \end{pmatrix}, \dots, \begin{pmatrix} X_{1n_1}^{(c)} \\ X_{2n_1}^{(c)} \end{pmatrix}}_{\mathbf{X}^{(c)}}, \underbrace{\begin{pmatrix} X_{11}^{(i)} \\ \text{---} \end{pmatrix}, \dots, \begin{pmatrix} X_{1n_2}^{(i)} \\ \text{---} \end{pmatrix}, \begin{pmatrix} \text{---} \\ X_{21}^{(i)} \end{pmatrix}, \dots, \begin{pmatrix} \text{---} \\ X_{2n_3}^{(i)} \end{pmatrix}}_{\mathbf{X}^{(i)}}, \quad (2.1)$$

where $n = n_1 + n_2 + n_3$. As in Model (1.1) we assume that the first components $X_{1j}^{(c)}, X_{1k}^{(i)}$ are i.i.d. with mean μ_1 and variance $\sigma_1^2 \in (0, \infty)$ and the second components $X_{2j}^{(c)}, X_{2l}^{(i)}$ are i.i.d. with mean μ_2 and variance $\sigma_2^2 \in (0, \infty)$ for $j = 1, \dots, n_1, k = 1, \dots, n_2, l = 1, \dots, n_3$. Moreover, the complete pairs $(X_{1j}^{(c)}, X_{2j}^{(c)})'$ are i.i.d. with mean vector $\boldsymbol{\mu} = (\mu_1, \mu_2)'$ and some unstructured covariance matrix $\boldsymbol{\Sigma} > 0$. Thus, the observations are missing completely at random. At the end of Section 3 we comment on how to relax these model assumptions to the case where the incomplete observations are only assumed to be independent.

In the setting (2.1), we like to use all the available data to test the null hypotheses $H_0 : \{\mu_1 = \mu_2\}$ against the one-sided alternative $\{\mu_1 > \mu_2\}$ or the two-sided alternative $\{\mu_1 \neq \mu_2\}$. For ease of convenience, we below focus on the one-sided case and note that two-sided tests can be obtained similarly by, for example, taking absolute values.

To motivate our test statistic, we first consider the most extreme situations. In a completely observed case with $n_1 = n$, the matched pairs would typically be inferred with the usual paired t -test type statistic

$$T_1 = T_1(\mathbf{X}^{(c)}) = \frac{n_1^{-1} \sum_{j=1}^{n_1} D_j}{\sqrt{\hat{\sigma}^2/n_1}} = \frac{\bar{D}}{\sqrt{\hat{\sigma}^2/n_1}}, \quad (2.2)$$

where $D_j = X_{1j}^{(c)} - X_{2j}^{(c)}$ denote the differences of the first and second component for $j = 1, \dots, n_1$ and $\hat{\sigma}^2 = (n_1 - 1)^{-1} \sum_{j=1}^{n_1} (D_j - \bar{D})^2$ is the empirical variance of these differences. If, in comparison, only incomplete vectors were observed, i.e. $n = n_2 + n_3$, we would be in the

situation of an extended Behrens-Fisher problem and could apply a Welch-type test in the test statistic

$$T_2 = T_2(\mathbf{X}^{(i)}) = \frac{\bar{X}_{1\cdot}^{(i)} - \bar{X}_{2\cdot}^{(i)}}{\sqrt{\hat{\sigma}_1^2/n_2 + \hat{\sigma}_2^2/n_3}}, \quad (2.3)$$

where $\bar{X}_{1\cdot}^{(i)} = n_2^{-1} \sum_{k=1}^{n_2} X_{1k}^{(i)}$ and $\bar{X}_{2\cdot}^{(i)} = n_3^{-1} \sum_{\ell=1}^{n_3} X_{2\ell}^{(i)}$ are the sample means and $\hat{\sigma}_1^2 = (n_2 - 1)^{-1} \sum_{k=1}^{n_2} (X_{1k}^{(i)} - \bar{X}_{1\cdot}^{(i)})^2$ and $\hat{\sigma}_2^2 = (n_3 - 1)^{-1} \sum_{\ell=1}^{n_3} (X_{2\ell}^{(i)} - \bar{X}_{2\cdot}^{(i)})^2$ are the corresponding empirical variances. To deal with all other cases of (2.1) adequately, it is then reasonable to propose the usage of a weighted test statistic

$$T = T(\mathbf{X}^{(c)}, \mathbf{X}^{(i)}) = \sqrt{a}T_1(\mathbf{X}^{(c)}) + \sqrt{1-a}T_2(\mathbf{X}^{(i)}), \quad (2.4)$$

where $a \in [0, 1]$ can be used to weight the inference drawn from the complete pairs. We suggest to use $a = 2n_1/(n + n_1)$ or $a = n_1/n$ to cover the usually applied methods in the two extreme cases $n_1 = n$ and $n_1 = 0$. The asymptotic null distribution of T as $\min(n_1, n_2, n_3) \rightarrow \infty$ is stated below. The asymptotic framework is, e.g., reasonable when assuming that each component is missing at random with a certain probability $p \in (0, 1)$.

THEOREM 2.1 *Suppose that $n_2/(n_2 + n_3) \rightarrow \kappa \in (0, 1)$ as $\min(n_1, n_2, n_3) \rightarrow \infty$ then we have under $H_0 : \mu_1 = \mu_2$*

$$\sup_{x \in \mathbb{R}} |P(T \leq x) - \Phi(x)| \rightarrow 0, \quad (2.5)$$

where Φ denotes the cdf of $N(0, 1)$.

From this a consistent asymptotic level α test is given by $\varphi = \mathbf{1}\{T > z_{1-\alpha}\}$ for testing the one-sided case $\{\mu_1 \leq \mu_2\}$ vs $\{\mu_1 > \mu_2\}$ and similar in the two-sided case, see also Samawi and Vogel [2014]. Here $z_{1-\alpha}$ denotes the $(1 - \alpha)$ -quantile of the standard normal distribution, $\alpha \in (0, 1)$. However, large sample sizes of the incomplete paired data are necessary to get an accurate approximation of this test statistic. In particular, if the sample size is small or the underlying distribution shows deviation from symmetry, the test will in general not keep the preassigned type-I error level accurately, see the simulation study in Section 5 below. Therefore, we suggest a novel permutation approach in the next section that is not only asymptotically valid but also finitely exact under certain assumptions.

3 The permutation procedures

Previous to our work, permutation techniques have been applied to the incomplete pairs design by Maritz [1995] and Yu et al. [2012]. However, these methods have the drawback that certain distributional assumptions such as 0-symmetry and equal variances or sample sizes are needed

to obtain a (at least asymptotically) valid level α test.

To overcome this issue, we propose a different permutation test that has the finite sample exactness property under similar invariance assumptions but is also (asymptotically) robust against deviations such as heteroscedasticity or underlying skewed distributions, i.e. valid for our general model (2.1). The procedure is based on separate results about studentized permutation tests for the paired and unpaired case. For example, in the classical matched pairs case with $n_1 = n$ Janssen [1999b] and Konietzschke and Pauly [2014] have already recommended (based on theoretical results and extensive simulation studies) a studentized permutation test in the paired t -test, where the components of each pair are randomly permuted. In comparison, in the general Behrens-Fisher set-up with $n_2 + n_3 = n$, Janssen [1997]; Janssen and Pauls [2003] as well as Janssen [2005] proposed a studentized permutation test in the Welch-type statistic T_2 that is based on randomly permuting the pooled sample, see also Chung and Romano [2013] as well as Pauly et al. [2015] for recent generalizations of this approach to multiple samples and general factorial designs. These studentized permutation procedures have also been implemented in R and SAS, see Placzek et al. [2014] as well as the R package GFD.

Jointly applying the ideas of both approaches we propose a permutation test that is based on the weighted test statistic T as follows: First, we randomly permute the components of each complete observation resulting in

$$\mathbf{X}_\tau^{(c)} = \left(X_{\tau_1(1)1}^{(c)}, \dots, X_{\tau_{n_1}(1)n_1}^{(c)} \right), \left(X_{\tau_1(2)1}^{(c)}, \dots, X_{\tau_{n_1}(2)n_1}^{(c)} \right), \quad (3.1)$$

where $\tau_i, i = 1, \dots, n_1$, are uniformly distributed on the symmetric group \mathcal{S}_2 (i.e. are random permutations of $(1, 2)$) and independent of $\mathbf{X} = (\mathbf{X}^{(c)}, \mathbf{X}^{(i)})$. Second, for the incomplete observations, we randomly permute the pooled incomplete sample $(Z_1, \dots, Z_{n_2+n_3}) = (X_{11}^{(i)}, \dots, X_{1n_2}^{(i)}, X_{21}^{(i)}, \dots, X_{2n_3}^{(i)})$ by means of a random permutation π that is uniformly distributed on $\mathcal{S}_{n_2+n_3}$ and independent of all other variables resulting in

$$\mathbf{X}_\pi^{(i)} = (Z_{\pi(1)}, \dots, Z_{\pi(n_2+n_3)}).$$

Finally, we approximate the null distribution of T by the conditional distribution of the permutation version of the test statistic $T_p = T(\mathbf{X}_\tau^{(c)}, \mathbf{X}_\pi^{(i)}) = \sqrt{a}T_1(\mathbf{X}_\tau^{(c)}) + \sqrt{1-a}T_2(\mathbf{X}_\pi^{(i)})$. The validity of this approach is proven below, where ' \xrightarrow{p} ' denotes convergence in probability.

THEOREM 3.1 *Suppose that $n_2/(n_2 + n_3) \rightarrow \kappa \in (0, 1)$ as $\min(n_1, n_2, n_3) \rightarrow \infty$ then we have*

$$\sup_{x \in \mathbb{R}} |P(T_p \leq x | \mathbf{X}) - \Phi(x)| \xrightarrow{p} 0. \quad (3.2)$$

Moreover, choosing $c_p(\alpha)$ as $(1 - \alpha)$ -quantile of the permutation distribution of T_p given the data, the test $\varphi_p = \mathbf{1}\{T > c_p(\alpha)\} + \gamma_p \mathbf{1}\{T = c_p(\alpha)\}$ is consistent and of asymptotic level α for testing H_0 against one-sided alternatives $\{\mu_1 > \mu_2\}$. Moreover, in the special case that D_1 is 0-symmetric and $X_{11}^{(i)}$ has the same distribution as $X_{21}^{(i)}$ the test is even finitely exact.

By inverting the above one-sided test, we can also construct one-sided confidence interval for the mean difference with asymptotic coverage of probability $(1 - \alpha)$. Moreover, a related two-sided confidence interval for $\mu_1 - \mu_2$ based on this approach is given by

$$[(T \pm c_P(\alpha/2))/S_w],$$

where

$$S_w = \frac{\sqrt{a}}{\hat{\sigma}/\sqrt{n_1}} + \frac{\sqrt{1-a}}{(\hat{\sigma}_1/\sqrt{n_2} + \hat{\sigma}_2/\sqrt{n_3})}$$

is a weighted sum of the inverses of the variance estimates for the paired and unpaired samples, respectively. Finally, applying the intersection-union principle (see e.g. Perlman et al. [1999]) it is straightforward to obtain a permutation TOST test for the bioequivalence hypothesis $H_0^{(b)} : |\mu_1 - \mu_2| \geq \epsilon$ for some given $\epsilon > 0$ from the above result.

REMARK 3.1 [Relaxing the Model Assumptions] *If the data is only missing at random, the i.i.d. assumptions of the incompletely observed first and second components, respectively, in model (2.1) might be violated. However, the above permutation approach can still be valid in such a situation. To accept this consider the relaxed model, where the incomplete components $(X_{1k}^{(i)})_k$ and $(X_{2\ell}^{(i)})_\ell$ are independent of $\mathbf{X}^{(c)}$ and assumed to be (row-wise) independent and infinitesimal with mean μ_1 (first component) and μ_2 (second component), respectively. If both means $\sqrt{n_2}(\bar{X}_{1\cdot}^{(i)} - \mu_1)$ and $\sqrt{n_3}(\bar{X}_{2\cdot}^{(i)} - \mu_2)$ are asymptotically normal with positive asymptotic variance it then follows from results of Janssen and Pauls [2003] and Janssen [2005] that the convergences (3.2) and $T \xrightarrow{d} N(0, 1)$ still hold under the null hypothesis. Thus, the proposed randomization procedure remains an asymptotic exact level α test. Moreover, skipping the assumption of mean equality, i.e. $E(X_{sr}^{(i)}) \neq E(X_{s1}^{(c)})$ may hold for $s = 1, 2, r = 1, \dots, n_{s+1}$, the procedure is even valid for testing the null hypothesis $H_0 : \{E(X_{11}^{(c)}) = E(X_{21}^{(c)}) \text{ and } E(\bar{X}_{1\cdot}^{(i)}) = E(\bar{X}_{2\cdot}^{(i)})\}$.*

4 Alternative approaches

In this subsection, we briefly review the existing literature to select possible competitors for our subsequent simulation study. Several methods that (only) deal with all available data in the incomplete matched pairs design have been suggested. However, most of the available procedures are based on specific assumptions such as normality, 0-symmetry or homoscedasticity of the data. We can summarize the most commonly used methods as follows:

- (a) Tests based on modified maximum likelihood estimators (cf., e.g., Lin and Stivers [1974], Ekbohm [1976, 1981], Woolson et al. [1976], or Hamdan et al. [1978]).
- (b) Tests based on simple mean difference estimator (see, e.g., Lin and Stivers [1974], Ekbohm [1976], Bhoj [1989], Looney and Jones [2003], or Uddin and Hasan [2015]).

- (c) Weighted linear and nonlinear combination tests (cf., e.g., Bhoj [1978, 1984, 1991], Kim et al. [2005], or Samawi and Vogel [2014]).
- (d) Resampling and more specific randomization methods (see, e.g., Maritz [1995] and Yu et al. [2012]).

For ease of presentation, it is not possible to compare our suggested approach with all these procedures. Also, none of them is free from distributional assumptions and at the same time robust against deviations such as heteroscedasticity and skewed distributions. Therefore, we selected a few procedures for the comparisons in our simulation study below. First, the test statistic proposed by Lin and Stivers [1974] has been chosen. It is a studentized mean difference statistic given by

$$T_{LS} = \frac{\overline{X}_{1.}^{(c,i)} - \overline{X}_{2.}^{(c,i)}}{\sqrt{\frac{1}{n_2+n_1} + \frac{1}{n_3+n_1} - \frac{2n_1r}{(n_2+n_1)(n_3+n_1)}} \sqrt{\frac{S_1^2+S_2^2}{n-2}}}.$$

Here, $\overline{X}_{1.}^{(c,i)}$ and $\overline{X}_{2.}^{(c,i)}$ denote the means of the paired and unpaired observations for sample 1 and sample 2 respectively, r is the empirical correlation coefficient of the completely observed pairs and $S_1^2 = \sum_{j=1}^{n_1} (X_{1j}^{(c)} - \overline{X}_{1.}^{(c,i)})^2 + \sum_{k=1}^{n_2} (X_{1k}^{(i)} - \overline{X}_{1.}^{(c,i)})^2$ as well as $S_2^2 = \sum_{l=1}^{n_3} (X_{2l}^{(i)} - \overline{X}_{2.}^{(i)})^2$ are sum of squares. Lin and Stivers motivated this statistic assuming normality of the data, and thus proposed to approximate the distribution of T_{LS} by a t_{n-4} -distribution with $n - 4$ degrees of freedom. In extensive simulation studies (e.g. Dunu [1994], and Bhoj [1989, 1991]), it turned out that the T_{LS} maintained the type-I error rate quite accurately under homoscedasticity. Moreover, it was found to possess a much better power behaviour in case of data with low correlation than the following competitors: The paired t-test only depending on the complete observations, the independent two-sample t-test only depending on the incomplete cases, the test by [Hamdan et al., 1978] based on a modified maximum likelihood estimator and the weighted convex combination test given in [Bhoj, 1991].

From the group of weighted linear and nonlinear combination tests, we selected the recently developed test by Kim et al. [2005] as second competitor. The test statistic is given by the following combination of paired and unpaired mean group differences:

$$t_3 = \frac{n_1(\overline{X}_{1.}^{(c)} - \overline{X}_{2.}^{(c)}) + n_H(\overline{X}_{1.}^{(i)} - \overline{X}_{2.}^{(i)})}{\sqrt{n_1\hat{\sigma}^2 + n_H^2(\hat{\sigma}_1^2/n_2 + \hat{\sigma}_2^2/n_3)}},$$

where n_H is the harmonic mean of n_2 and n_3 . In their paper the authors proposed to approximate the null distribution of t_3 with a standard normal distribution and applied it to data from a colorectal cancer microarray study for identifying differentially expressed genes, see also Yu et al. [2012].




Figure 1: Simulation results for the type-I error level for T_P (---), T (—), T_{LS} (- · -), and t_3 (···) for different distributions under varying correlation values each with sample sizes $(n_1, n_2, n_3) \in (10, 10, 10)$ (left) and $(30, 10, 10)$ (right) and different covariance matrices Σ_1 (a) and Σ_2 (b).

Finally, we selected the asymptotic test T of Section 2 as it was recommended by a simulation study in Samawi and Vogel [2014] in favor of their suggested pooled t-test and the corrected z -test [Looney and Jones, 2003]. A similar conclusion was drawn by Guo and Yuan [2015] who also found out in an extensive simulation study that T controls the type-I error rate quite accurate in case of moderate sample sizes.

Other tests as the corrected z -test by Looney and Jones [2003] or the pooled t-test by Samawi and Vogel [2014] have not been included since they also showed worse performances for small sample sizes in other simulation studies, see e.g. Samawi and Vogel [2014], Guo and Yuan [2015] and Uddin and Hasan [2015]. We thus restricted our simulation study to comparing the small sample behavior of our suggested test with the three competitors from above.

5 Monte-Carlo simulations

In this Section, we investigate the finite sample behavior of our suggested permutation test and the chosen competing procedures as described in Section 4 in extensive simulations. As major assessment criteria all procedures were studied with respect to (i) maintaining the preassigned type-I error level under the null hypothesis and (ii) their powers to detect certain alternatives. The observations were generated using four symmetric and skewed bivariate distribu-

tions: normal, exponential, asymmetric Laplace, and Laplace distribution each with correlation $\rho \in (-1, 1)$ and with the following two covariance matrices

$$\Sigma_1 = \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix}, \quad \Sigma_2 = \begin{pmatrix} 1 & \sqrt{2}\rho \\ \sqrt{2}\rho & 2 \end{pmatrix},$$

representing a homoscedastic and a heteroscedastic setting, respectively. The complete pairs were simulated from the model

$$\begin{pmatrix} X_{1j}^{(c)} \\ X_{2j}^{(c)} \end{pmatrix} = \Sigma^{1/2} \begin{pmatrix} \varepsilon_{1j} \\ \varepsilon_{2j} \end{pmatrix} + \begin{pmatrix} \mu_1 \\ \mu_2 \end{pmatrix}, \quad j = 1, \dots, n_1.$$

Here, the standardized random errors $\varepsilon_{ij}, i = 1, 2$, were generated by $\varepsilon_{ij} = \frac{Y_{ij} - E(Y_{i1})}{\sqrt{\text{var}(Y_{i1})}}$, where Y_{ij} denote normal, exponential, asymmetric Laplace, or Laplace random variables, respectively. Similarly, the incomplete observations were generated as $X_{2j}^{(i)} = \sigma_i \tilde{\varepsilon}_{ij} + \mu_i$, where $\tilde{\varepsilon}_{ij}$ are independent copies of the ε_{ij} and the scaling factors are $\sigma_1 = 1$ and $\sigma_2 \in \{1, \sqrt{2}\}$ depending on the choice of covariance matrix. Moreover, the following sample sizes $(n_1, n_2, n_3) \in \{(10, 10, 10), (30, 10, 10)\}$ were considered and the nominal significance level was taken as $\alpha = 0.05$. Under the null hypothesis, we set $\mu_1 = \mu_2 = 0$. To also evaluate the empirical power, the second samples were shifted by adding a shift $\mu_2 = \delta \in \{0, 0.5, 1\}$. The choice of $\delta = 0$ corresponds to the analysis of the empirical level of significance. All simulations were operated by means of the R computing environment, version 3.2.2 [R Core Team, 2015] and each setting was based on 10,000 simulation runs and $B = 1,000$ permutations runs. The algorithm for the computation of the p -value of our suggested permutation test is as follows:

1. For the given partially paired data, calculate the observed test statistic T with weight $a = 2n_1/(n + n_1)$.
2. Randomly permute the components of the complete pairs $\mathbf{X}^{(c)}$, resulting in $\mathbf{X}_\tau^{(c)}$.
3. Randomly permute the pooled incomplete pairs $\mathbf{X}^{(i)}$, resulting in $\mathbf{X}_\pi^{(i)}$.
4. Calculate the value of the weighted test statistic for the permuted data $T_p = T(\mathbf{X}_\tau^{(c)}, \mathbf{X}_\pi^{(i)})$.
5. Repeat the Steps 2 and 3 Independently $B = 1,000$ times and collect the observed test statistic values in $T_b, b = 1, \dots, B$.
6. Finally, estimate the two sided permutation p -value as follows $p\text{-value} = \min(2p_1, 2 - 2p_1)$, where $p_1 = \frac{\sum_{b=1}^B I(T_{pb} \geq T)}{B}$.

The type-I error level of the proposed procedures for different sample sizes and for homoscedastic as well as heteroscedastic settings are displayed in Figure 1. It can be readily seen from Figure 1 that for all the considered settings, our suggested studentized permutation test T_p




Figure 2: Power simulation results ($\alpha = 0.05$) for T_P (---) and t_3 (···) for Symmetric and Asymmetric distributions under $\rho = 0.9$ for $(n_1, n_2, n_3) \in (30, 10, 10)$ and for Σ_1 (a) and Σ_2 (b).

tends to result in an adequate exact level α test under homoscedasticity as well as heteroscedasticity and over the range of correlations ρ for most settings. Only in the case of the skewed exponential distribution, the control is not adequate. However, in this case all the other competing procedures also failed to control the type-I error rate for the underlying sample sizes. More specific, in the case of homoscedasticity, data with low correlation ($|\rho| \leq 0.5$), and exponential distribution our suggested T_p still results in an accurate test decision and also performs better when the number of complete pairs is increased.

In contrast, the competing tests do not control the type-I error level constantly under heteroscedasticity or even under homoscedasticity in the two considered cases of sample sizes. It can also be seen from Figure 1 that the weighted test T is rather liberal in the case of small number of complete pairs and less liberal for larger number of complete pairs. This behavior of the test does not depend on the homoscedasticity assumption. Moreover, the test that is based on the simple mean difference estimator T_{LS} controls type-I rate in the case of equal variances under all the simulation settings except for the skewed exponential distribution. However, it is very liberal in the case of heteroscedasticity and its behavior is much worse for larger number of complete pairs. Finally, the test t_3 tends to result in a liberal decision in the case of small number of complete pairs and vice versa it is nearly exact level α in case of a larger number of complete pairs; especially under homoscedasticity.

The power of the four tests has also been studied for all settings. Due to the rather liberal behavior of the test based on T_{LS} and the asymptotic test based on T , their power functions are not really comparable to the others. Therefore, we restrict our power simulation study to the analysis of T_p and t_3 . Moreover, since their type-I error rate control is rather similar for $(n_1, n_2, n_3) \in (30, 10, 10)$ we only consider the unbalanced case. Figure 2 shows the power of both methods for highly correlated data involving homoscedastic as well as heteroscedastic settings. It can be easily seen that our suggested studentized permutation test T_p has a larger power under all studied settings. However, both tests yielded an almost similar power behavior when equal variances and symmetric distributions are present. In all other cases (heteroscedasticity or asymmetric distributions) the power of t_3 is considerably smaller. Note, that the power functions are not really comparable under the exponential distribution due to their rather liberal behavior and we have only included it for illustration and completeness. In the power simulations for all the other considered situations (not shown here) both tests performed nearly similar.

6 Analysis of the data example

The Karnofsky Performance Status Scale (KPS) as proposed by Karnofsky [1949], is an important assessment tool for functional impairment. This scale can be used for determining the prognosis of individual patients or for comparing effectiveness of several therapies. The score ranges from 0 to 100. The lower the KPS score, the lower the probability of survival.

A clinical study by Hermann and Looney [2001] has been performed to assess the functional status of hospice patients during their last seven days of life and to discover and examine the different interventions that have been implemented to treat the patients symptoms. Their study consists of 100 hospice patients. KPS scores were used to assess the functional status of the patients on the day before they died and their last day of life resulting in paired data. These pairs were not complete. For example, there were a total of $n_1 = 9$ complete pairs and two unpaired samples for the patients who were only observed on their day before they died or their last day of life of sizes $n_2 = 28$ and $n_3 = 23$ respectively. The data was published in Rempala and Looney [2006]. We applied all considered testing methods T_p , T , T_{LS} , and t_3 to detect the null hypothesis of equality of means of the KPS scores on the day before they died and on the last day of their life ($H_0 : \{\mu_1 = \mu_2\}$) against the one-sided alternative $\{\mu_1 > \mu_2\}$ or the two-sided alternative $\{\mu_1 \neq \mu_2\}$. The results are summarized in Table 1.

Table 1: P-values of the KPS Data

Method	One-sided p-value	Two-sided p-value
T_p	0.0110	0.0220
T	0.0025	0.0050
T_{LS}	0.0077	0.0153
t_3	0.0069	0.0138

All four tests provided in Table 1 reject the null hypothesis and indicate that the mean values of KPS score on the last day and on the day before they died are significantly different (two sided p -value < 0.05). In particular, their KPS score on the day before they died was significantly greater than their score on the last day of life (right-sided p -value < 0.05).

7 Conclusions and discussion

The problem of incomplete paired data occurs frequently in practice. In this work, we have provided a novel randomization test that is not based on any parametric assumptions and uses all the observed data of the matched pairs design. It was shown to be asymptotic valid and even finitely exact if certain invariance properties are met. For the asymptotic considerations, ideas of Janssen [1997], Janssen [1999a], Janssen [1999b], Neubert and Brunner [2007], Pauly [2011], Konietschke and Pauly [2012], Omelka and Pauly [2012], Chung and Romano [2013], DiCiccio and Romano [2015], Pauly et al. [2015] and Chung and Romano [2016] on studentized permutation tests for paired and unpaired two-sample problems were combined. In addition, the theoretical asymptotic findings were also applied to bioequivalence problems and to construct one- and two-sided confidence interval for the two sample mean difference.

In an extensive simulation study, the advantageous theoretical properties of the randomization test were underpinned. There it was seen that our novel test possess nice small sample properties and particularly improves the small sample behavior of the asymptotic test based on the same weighted test statistic T . In particular, the simulations also indicated that our randomization test is robust in terms of error control against heteroscedasticity or skewed distributions in most considered situations. Only in the case of the exponential distribution with rather large correlations its behaviour was rather liberal for small sample sizes. In this setting, however, the other considered competing procedures (Samawi and Vogel [2014], Kim et al. [2005], and Lin and Stivers [1974]) also failed to control the type-I error rate.

To sum up, its simulation results together with the finite exactness property under invariance and its asymptotic correctness make the new permutation procedure recommendable in practice.

Acknowledgement

This work was supported by the German Academic Exchange Service (DAAD) under the project: Research Grants - Doctoral Programmes in Germany, 2015/16 (No. 57129429).

Appendix

Proof of Theorem 2.1:

It follows from the central limit theorem and Slutsky that we have convergence in distribution

$$T_1(\mathbf{X}^{(c)}) \xrightarrow{d} N(0, 1) \quad \text{as well as} \quad T_2(\mathbf{X}^{(i)}) \xrightarrow{d} N(0, 1).$$

Since $T_1(\mathbf{X}^{(c)})$ and $T_2(\mathbf{X}^{(i)})$ are independent, it follows that

$$\sqrt{a}T_1(\mathbf{X}^{(c)}) + \sqrt{1-a}T_2(\mathbf{X}^{(i)}) \xrightarrow{d} N(0, a + 1 - a) = N(0, 1).$$

The stated result thus follows from Polya's Theorem. \square

Proof of Theorem 3.1:

From the results given in Janssen [1999b] and Konietzschke and Pauly [2014] it follows under the given assumptions that

$$\sup_{x \in \mathbb{R}} \left| P(T_1(\mathbf{X}_{\tau}^{(c)}) \leq x | \mathbf{X}^{(c)}) - \Phi(x) \right| \xrightarrow{P} 0.$$

Similarly, the results from [Janssen, 2005], see also [Pauly et al., 2015], show that

$$\sup_{x \in \mathbb{R}} \left| P(T_2(\mathbf{X}_{\pi}^{(i)}) \leq x | \mathbf{X}^{(i)}) - \Phi(x) \right| \xrightarrow{P} 0. \quad (7.1)$$

Since π and $\tau_j, j = 1, \dots, n_1$, are independent the weak convergence result follows from a conditional Slutsky Theorem, see e.g. Chung and Romano [2013] or Pauly [2009]. From this an application of Lemma 1 in Janssen and Pauls [2003] proves the consistency and asymptotic exactness of the permutation test. The finite exactness under the given invariance assumptions follows from classical theory on randomization tests, see e.g. Lehmann and Romano [2006] or Janssen and Völker [2007]. \square

Proof of Remark 3.1:

First, $T \xrightarrow{d} N(0, 1)$ follows from Raikov's Theorem. Thus, to accept the stated results we only have to argue that (7.1) also holds for the more general model under the stated assumptions. But this follows from Theorem 4.1 in Janssen [2005] by applying Lemma 7 in Janssen and Pauls [2003], see also the comments in Example 5(c) of the first paper. \square

References

- Michael G Akritas, Jouni Kuha, and D Wayne Osgood. A nonparametric approach to matched pairs with missing data. *Sociological methods & research*, 30(3):425–454, 2002.
- Dinesh S Bhoj. Testing equality of means of correlated variates with missing observations on both responses. *Biometrika*, 65(1):225–228, 1978.
- Dinesh S Bhoj. On difference of means of correlated variates with incomplete data on both responses. *Journal of Statistical Computation and Simulation*, 19(4):275–289, 1984.
- Dinesh S Bhoj. On comparing correlated means in the presence of incomplete data. *Biometrical Journal*, 31(3):279–288, 1989.
- Dinesh S Bhoj. Testing equality of means in the presence of correlation and missing data. *Biometrical journal*, 33(1):63–72, 1991.
- Eun Yi Chung and Joseph P Romano. Exact and asymptotically robust permutation tests. *The Annals of Statistics*, 41(2):484–507, 2013.
- EunYi Chung and Joseph P Romano. Asymptotically valid and exact permutation tests based on two-sample u-statistics. *Journal of Statistical Planning and Inference*, 168:97–105, 2016.
- Cyrus J DiCiccio and Joseph P Romano. Robust permutation tests for correlation and regression coefficients. 2015.
- Emeka Samuel Dunu. *Comparing the powers of several proposed tests for testing the equality of the means of two populations when some data are missing*. PhD thesis, University of North Texas, 1994.
- G Ekbohm. On testing equality of means in the paired case with incomplete data on both responses. *Biometrical Journal*, 23(3):251–259, 1981.
- Gunnar Ekbohm. On comparing means in the paired case with incomplete data on both responses. *Biometrika*, 63(2):299–304, 1976.
- Beibei Guo and Ying Yuan. A comparative review of methods for comparing means using partially paired data. *Statistical Methods in Medical Research*, 2015.
- MA Hamdan, AI Khuri, and SL Crews. A test for equality of means of two correlated normal variates with missing data on both responses. *Biometrical Journal*, 20:667–674, 1978.
- Carla Hermann and Stephen Looney. The effectiveness of symptom management in hospice patients during the last seven days of life. *Journal of Hospice & Palliative Nursing*, 3(3): 88–96, 2001.

- Arnold Janssen. Studentized permutation tests for non-iid hypotheses and the generalized Behrens-Fisher problem. *Statistics & Probability Letters*, 36(1):9–21, 1997.
- Arnold Janssen. Nonparametric symmetry tests for statistical functionals. *Mathematical Methods of Statistics*, 8(3):320–343, 1999a.
- Arnold Janssen. Testing nonparametric statistical functionals with applications to rank tests. *Journal of Statistical Planning and Inference*, 81(1):71–93, 1999b.
- Arnold Janssen. Resampling student’s t-type statistics. *Annals of the Institute of Statistical Mathematics*, 57(3):507–529, 2005.
- Arnold Janssen and Thorsten Pauls. How do bootstrap and permutation tests work? *Annals of Statistics*, pages 768–806, 2003.
- Arnold Janssen and Dominik Völker. Most powerful conditional tests. *Statistics & Decisions*, 25(1/2007):41–62, 2007.
- David A Karnofsky. The clinical evaluation of chemotherapeutic agents in cancer. *Evaluation of Chemotherapeutic Agents*, 1949.
- Byung Soo Kim, Inyoung Kim, Sunho Lee, Sangcheol Kim, Sun Young Rha, and Hyun Cheol Chung. Statistical methods of translating microarray data into clinically relevant diagnostic information in colorectal cancer. *Bioinformatics*, 21(4):517–528, 2005.
- Frank Konietzschke and Markus Pauly. A studentized permutation test for the nonparametric Behrens-Fisher problem in paired data. *Electronic Journal of Statistics*, 6:1358–1372, 2012.
- Frank Konietzschke and Markus Pauly. Bootstrapping and permuting paired t-test type statistics. *Statistics and Computing*, 24:283–296, 2014.
- Erich L Lehmann and Joseph P Romano. *Testing Statistical Hypotheses*. Springer Science & Business Media, 2006.
- Pi-Erh Lin and Lawrence E Stivers. On difference of means with incomplete data. *Biometrika*, 61(2):325–334, 1974.
- Roderick JA Little and Donald B Rubin. *Statistical analysis with missing data*. John Wiley and Sons, 2014.
- Stephen W Looney and Peter W Jones. A method for comparing two normal means using combined samples of correlated and uncorrelated data. *Statistics in Medicine*, 22(9):1601–1610, 2003.
- Johannes S. Maritz. A permutation paired test allowing for missing values. *Australian Journal of Statistics*, 37(2):153–159, 1995.

- Karin Neubert and Edgar Brunner. A studentized permutation test for the nonparametric Behrens–Fisher problem. *Computational Statistics & Data Analysis*, 51(10):5192–5204, 2007.
- Marek Omelka and Markus Pauly. Testing equality of correlation coefficients in two populations via permutation methods. *Journal of Statistical Planning and Inference*, 142(6):1396–1406, 2012.
- Markus Pauly. Eine Analyse bedingter Tests mit bedingten Zentralen Grenzwertsätzen für Resampling-Statistiken. *Inaugural Dissertation University of Duesseldorf*, 2009.
- Markus Pauly. Discussion about the quality of f-ratio resampling tests for comparing variances. *Test*, 20(1):163–179, 2011.
- Markus Pauly, Edgar Brunner, and Frank Konietschke. Asymptotic permutation tests in general factorial designs. *Journal of the Royal Statistical Society B (Statistical Methodology)*, 77: 461–473, 2015.
- Michael D Perlman, Lang Wu, et al. The emperor’s new tests. *Statistical Science*, 14(4): 355–369, 1999.
- Mariusz Placzek, Frank Konietschke, and Markus Pauly. Studentisierte Permutationstests für verbundene und unverbundene 2-Stichprobenprobleme. *KSFE 2014*, 2014.
- R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria, 2015. URL <https://www.R-project.org/>.
- Grzegorz A Rempala and Stephen W Looney. Asymptotic properties of a two sample randomized test for partially dependent data. *Journal of Statistical Planning and Inference*, 136(1): 68–89, 2006.
- Hani M Samawi and Robert Vogel. Notes on two sample tests for partially correlated (paired) data. *Journal of Applied Statistics*, 41(1):109–117, 2014.
- Nizam Uddin and MS Hasan. Testing equality of two normal means using combined samples of paired and unpaired data. *Communications in Statistics-Simulation and Computation*, (just-accepted), 2015.
- Geert Verbeke and Geert Molenberghs. *Linear mixed models for longitudinal data*. Springer Science & Business Media, 2009.
- RF Woolson, JD Leeper, JW Cole, and WR Clarke. A monte carlo investigation of a statistic for a bivariate missing data problem. *Communications in Statistics-Theory and Methods*, 5 (7):681–688, 1976.
- Donghyeon Yu, Johan Lim, Feng Liang, Kyunga Kim, Byung Soo Kim, and Woncheol Jang. Permutation test for incomplete paired data with application to cdna microarray data. *Computational Statistics & Data Analysis*, 56(3):510–521, 2012.