

Vacuum Rabi Splitting in Nanomechanical QED System with Nonlinear Resonator

M. Y. Zhao,¹ S. M. Yu,¹ X. Xiao,^{1,2} and Y. B. Gao^{1,*}

¹College of Applied Sciences, Beijing University of Technology, Beijing, 100124, China

²Department of Physics, Renmin University of China, Beijing, 100872, China

Considering the intrinsic nonlinearity in a nanomechanical resonator coupled to a charge qubit, vacuum Rabi splitting effect is studied in a nanomechanical QED (qubit-resonator) system. A driven nonlinear Jaynes-Cummings model describes the dynamics of this qubit-resonator system. Using quantum regression theorem and master equation approach, we have calculated the two-time correlation spectrum analytically. In the weak driving limit, these analytical results clarify the influence of the driving strength and nonlinearity parameter on the correlation spectrum. Also, numerical calculations confirm these analytical results.

PACS numbers: 85.85.+j, 85.25.Cp

Keywords: vacuum Rabi splitting, nanomechanical QED, nonlinear resonator

I. INTRODUCTION

In quantum optics [1] and quantum information, [2] the well known Jaynes-Cummings model, [3] one of the most important models, describes the light-matter interaction between a two-level quantum system (qubit) and a boson (resonator). Generally in cavity QED system, [4] nanomechanical QED system [5] and circuit QED system, [6] vacuum Rabi splitting effect has been used to characterize the coupling strength between a qubit and a resonator.

Recently, a nanomechanical resonator with frequency of the order of 1 GHz approaches the quantum regime, [7, 8] it is getting closer to test the basic principles of quantum mechanics. When a superconducting qubit [9, 10] is coupled to a nanomechanical resonator, [11] we can study quantum optical properties, such as, quantum decoherence, [12, 13] Rabi oscillation, [14] vacuum Rabi splitting, [15] classical-quantum transition [16] and phonon blockade. [17]

Increasing the amplitude of driving, the nonlinear response of nanomechanical resonator [18] is not negligible which can be used to detect the classical-quantum transition. [19] When intrinsic nonlinearity of nanomechanical resonator [20] is considered in the qubit-resonator system, we can use superconducting qubit to probe quantum fluctuations of nonlinear resonator. [21] Recently, the nonlinearity can be exploited to generate nonclassical states in mechanical systems [22, 23] and selectively address the nanomechanical qubit transitions in quantum information processing. [24]

In this paper, the dissipative dynamics of the qubit-resonator system is solved by master equation approach and quantum regression theorem. [25] Comparing with previous results, [15] the influence of the nonlinearity of nanomechanical resonator on vacuum Rabi splitting effect is studied analytically and numerically.

The paper is organized as follows. In Sec. II, using a driven nonlinear Jaynes-Cummings model, we describe the dynamics of a qubit-resonator system consisting of a superconducting qubit and a nonlinear nanomechanical resonator. In Sec. III, the two-time correlation spectrum is calculated analytically. In Sec. IV, vacuum Rabi splitting effect is studied. Also numerical simulations confirm our analytical results. Finally, our conclusions are summarized.

II. MODEL

In nanomechanical QED system, consisting a superconducting qubit and a nanomechanical resonator, we can use a driven Jaynes-Cummings model to describe the dynamics of this qubit-resonator system, [15]

$$H_{JC}^{\text{driven}} = \omega_a \sigma_+ \sigma_- + g (a \sigma_+ + a^\dagger \sigma_-) + \omega_c a^\dagger a - \xi \sin(\omega_p t) (a + a^\dagger). \quad (1)$$

Here H_{JC}^{driven} is a driven Jaynes-Cummings type Hamiltonian. The lowering (raising) operator σ_- (σ_+) and the annihilation (creation) operator a (a^\dagger) are defined to describe the qubit with frequency ω_a and the resonator with frequency ω_c respectively. The commutation relations, $[\sigma_-, \sigma_+] = \sigma_z$ and $[a, a^\dagger] = 1$, are satisfied. The last term in Eq. (1) is a classical drive with driving constant ξ and driving frequency ω_p . The g denotes the interaction strength between the qubit and the resonator.

Considering the nonlinearity of nanomechanical resonator, nanomechanical resonator is not assumed to be an ideal resonator again. Moving into a frame with rotating frequency ω_p , the total Hamiltonian for this qubit-resonator system writes [15]

$$H_t = \Delta_a \sigma_+ \sigma_- + g (a \sigma_+ + a^\dagger \sigma_-) + \Delta_c a^\dagger a - \xi (a + a^\dagger) + \chi a^\dagger a + \chi (a^\dagger a)^2, \quad (2)$$

which describes the dynamics of a driven nonlinear Jaynes-Cummings model. The linear part $\chi a^\dagger a$ and nonlinear part $\chi (a^\dagger a)^2$ come from a quartic potential x^4 . [20] Here the nonlinearity parameter χ is small, $\chi \ll g$.

*Corresponding author, E-mail: ybgao@bjut.edu.cn

Some parameters in the Hamiltonian Eq. (2) are

$$\Delta_a = \omega_a - \omega_p, \quad \Delta_c = \omega_c - \omega_p. \quad (3)$$

The detuning δ between the frequencies of the resonator and the qubit is

$$\delta = \Delta_c - \Delta_a. \quad (4)$$

III. TWO-TIME CORRELATION SPECTRUM

In previous results, [15] the induced electromotive force between two ends of nanomechanical resonator is

$$V = iBl\sqrt{\frac{\omega_c}{2M}}(a^\dagger - a). \quad (5)$$

To characterize the vacuum Rabi splitting in this nanomechanical QED system, the two-time correlation spectrum for the V is

$$S_V(\omega) = \frac{1}{\pi} \text{Re} \int_0^\infty d\tau e^{-i\omega\tau} \langle V(\tau)V(0) \rangle. \quad (6)$$

Based on those results in Eqs. (5,6), the two-time correlation function $\langle V(\tau)V(0) \rangle$ writes

$$\begin{aligned} \langle V(\tau)V(0) \rangle &\propto \langle a^\dagger(\tau)a(0) \rangle + \langle a(\tau)a^\dagger(0) \rangle \\ &\quad - \langle a(\tau)a(0) \rangle - \langle a^\dagger(\tau)a^\dagger(0) \rangle. \end{aligned} \quad (7)$$

Considering the interaction with the environment, we use the master equation to describe dissipative dynamics of the qubit-resonator system, [25]

$$\begin{aligned} \dot{\rho} &= -i[H_t, \rho] + \kappa(2a\rho a^\dagger - a^\dagger a\rho - \rho a^\dagger a) \\ &\quad + \frac{\gamma}{2}(2\sigma_- \rho \sigma_+ - \sigma_+ \sigma_- \rho - \rho \sigma_+ \sigma_-), \end{aligned} \quad (8)$$

where the density operator ρ describes the time evolution of the qubit-resonator system. The latter two terms in Eq. (8) describe the decay process of the resonator and the qubit respectively. The parameters κ and γ denote the decay rates of the resonator and the qubit. Here the Markov approximation is satisfied.

In this case, the number operator $N = \sigma_+ \sigma_- + a^\dagger a$ is defined to characterize the total number of excitations of the qubit-resonator system. The operator N satisfies

$$N|j, k\rangle = (j+k)|j, k\rangle,$$

where $|j, k\rangle = |j\rangle \otimes |k\rangle$ is the state of the qubit-resonator system, the index j (k) denotes the state of the qubit (resonator) and satisfies $\sigma_+ \sigma_- |j\rangle = j|j\rangle$ ($a^\dagger a |k\rangle = k|k\rangle$) for $j = 0, 1$ ($k = 0, 1, \dots$).

In this paper, the weak driving limit is adopted, $\xi \rightarrow 0$. When time approaches infinite, $t \rightarrow \infty$, the qubit-resonator system will decay into the vacuum state $|00\rangle$. The weak driving ξ will induce the transitions from the

vacuum state $|00\rangle$ to the other excited states ($|01\rangle$ and $|10\rangle$). Under the lowest order perturbation theory, we need to consider only one-phonon excitation in the qubit-resonator system,

$$N = j + k = 0, 1. \quad (9)$$

The Hilbert space for the reduced density matrix ρ in Eq. (8) reduces into a smaller subspace with a truncated basis

$$\{|j, k\rangle, j + k = 0, 1\}. \quad (10)$$

Thus, in this truncated basis, the corresponding density matrix elements satisfying the master equation in Eq. (8) are

$$\begin{aligned} \frac{d\rho_{00,00}}{d\tau} &= i\xi\rho_{01,00} - i\xi\rho_{00,01} \\ &\quad + 2\kappa\rho_{01,01} + \gamma\rho_{10,10}, \\ \frac{d\rho_{00,01}}{d\tau} &= [i(\Delta_c + 2\chi) - \kappa]\rho_{00,01} \\ &\quad + ig\rho_{00,10} + i\xi(\rho_{01,01} - \rho_{00,00}), \\ \frac{d\rho_{00,10}}{d\tau} &= \left(i\Delta_a - \frac{\gamma}{2}\right)\rho_{00,10} + ig\rho_{00,01} + i\xi\rho_{01,10}, \\ \frac{d\rho_{01,01}}{d\tau} &= -2\kappa\rho_{01,01} + ig(\rho_{01,10} - \rho_{10,01}) \\ &\quad + i\xi(\rho_{00,01} - \rho_{01,00}), \\ \frac{d\rho_{01,10}}{d\tau} &= \left[-i(\delta + 2\chi) - \kappa - \frac{\gamma}{2}\right]\rho_{01,10} \\ &\quad + ig(\rho_{01,01} - \rho_{10,10}) + i\xi\rho_{00,10}, \\ \frac{d\rho_{10,10}}{d\tau} &= -\gamma\rho_{10,10} + ig(\rho_{10,01} - \rho_{01,10}). \end{aligned} \quad (11)$$

In the long-time limit, the system stays in a steady state, we can take $\rho_{00,00} \sim 1$ and $\rho_{00,00} \gg \rho_{01,01}$ ($\rho_{10,10}$). Here can see that $\rho_{00,01}$ ($\rho_{00,10}$) scales as the order of ξ and $\rho_{01,01}$ ($\rho_{10,10}$) scales as the order of ξ^2 . Keeping terms of the ξ and dropping the higher terms of the ξ^2 , we get

$$\begin{aligned} \frac{d\rho_{00,01}}{d\tau} &= [i(\Delta_c + 2\chi) - \kappa]\rho_{00,01} + ig\rho_{00,10} - i\xi\rho_{00,00}, \\ \frac{d\rho_{00,10}}{d\tau} &= \left(i\Delta_a - \frac{\gamma}{2}\right)\rho_{00,10} + ig\rho_{00,01}. \end{aligned} \quad (12)$$

Applying Laplace transformation, we have solved

$$\rho_{00,01}(\tau) = \mu(\tau)\rho_{00,01}^{ss} + \nu(\tau)\rho_{00,10}^{ss} - i\xi C(\tau)\rho_{00,00}^{ss}. \quad (13)$$

Here the coefficients $(\mu(\tau), \nu(\tau)$ and $C(\tau))$ are depen-

dent of time, i.e.,

$$\begin{aligned}\mu(\tau) &= \frac{\Gamma_1 + i\omega_1 + (i\Delta_a - \frac{\gamma}{2})}{2iG} e^{-\Gamma_1\tau} e^{-i\omega_1\tau} \\ &\quad + \frac{\Gamma_2 + i\omega_2 + (i\Delta_a - \frac{\gamma}{2})}{-2iG} e^{-\Gamma_2\tau} e^{-i\omega_2\tau}, \\ \nu(\tau) &= \frac{g}{-2G} e^{-\Gamma_1\tau} e^{-i\omega_1\tau} + \frac{g}{2G} e^{-\Gamma_2\tau} e^{-i\omega_2\tau}, \\ C(\tau) &= \frac{1}{2iG} \frac{\Gamma_1 + i\omega_1 + (i\Delta_a - \frac{\gamma}{2})}{\Gamma_1 + i\omega_1} (1 - e^{-\Gamma_1\tau} e^{-i\omega_1\tau}) \\ &\quad - \frac{1}{2iG} \frac{\Gamma_2 + i\omega_2 + (i\Delta_a - \frac{\gamma}{2})}{\Gamma_2 + i\omega_2} (1 - e^{-\Gamma_2\tau} e^{-i\omega_2\tau}),\end{aligned}$$

$$\Gamma_n = \frac{1}{2} \left(\frac{\gamma}{2} + \kappa \right) + (-1)^n \text{Im}G, \quad (14)$$

$$\omega_n = (-1)^{n+1} \text{Re}G - \frac{1}{2} \Delta_a - \frac{1}{2} (\Delta_c + 2\chi), \quad (15)$$

and

$$G = \sqrt{g^2 - \frac{1}{4} \left[i(\delta + 2\chi) + \left(\frac{\gamma}{2} - \kappa \right) \right]^2}$$

for $n = 1, 2$.

From the above results in Eq. (12), we can calculate the single-time function

$$\langle a(\tau) \rangle = \text{Tr}(a(0) \rho(\tau)) = \rho_{01,00}(\tau) \quad (16)$$

where $\rho(0) = \rho^{ss}$ and the index ss means the steady state. Through some simple calculations, we obtain the steady solution of the density matrix element $\rho_{01,01}(\rho_{10,10})$,

$$\begin{aligned}\rho_{00,01}^{ss} &= \frac{\xi}{(\Delta_c + 2\chi) + i\kappa - \frac{g^2}{(\Delta_a + i\frac{\gamma}{2})}}, \\ \rho_{00,10}^{ss} &= -\frac{g}{(\Delta_a + i\frac{\gamma}{2})} \rho_{00,01}^{ss}.\end{aligned} \quad (17)$$

Using the quantum regression theorem [15, 25, 26], we can obtain the two-time correlation function

$$\langle a(\tau) a^\dagger(0) \rangle = \text{Tr} \{ a(\tau) a^\dagger(0) \rho(0) \} = \mu(\tau)^*. \quad (18)$$

The other two-time functions can be obtained as

$$\langle a^\dagger(\tau) a(0) \rangle = -i\xi C(\tau) \rho_{01,00}^{ss}, \quad (19)$$

$$\langle a^\dagger(\tau) a^\dagger(0) \rangle = 0 \quad (20)$$

and

$$\langle a(\tau) a(0) \rangle = 0. \quad (21)$$

Based on the above results in Eqs. (18,19,20,21), the two-time correlation function for the induced electromotive force is obtained,

$$\langle V(\tau) V(0) \rangle \propto \mu(\tau)^*. \quad (22)$$

In the limit of weak driving, we can neglect the terms ($i\xi\rho_{01,00}(0)$, $\rho_{02,00}(0)$, $\rho_{11,00}(0)$, $\rho_{01,01}(0)$ and $\rho_{01,10}(0)$) which are proportional to the ξ^2 in Eq. (22).

Using the formula,

$$\int_0^\infty d\tau e^{-i(\omega-\omega_0)\tau-\Gamma\tau} = \frac{\Gamma - i(\omega - \omega_0)}{(\omega - \omega_0)^2 + \Gamma^2},$$

the correlation spectrum in Eq. (6) is calculated as

$$S_V(\omega) \propto \frac{\text{Re}\eta_1}{(\omega - \omega_1)^2 + \Gamma_1^2} + \frac{\text{Re}\eta_2}{(\omega - \omega_2)^2 + \Gamma_2^2}. \quad (23)$$

Where some parameters are defined,

$$\eta_n = \frac{\Gamma_n - i\omega_n + (-i\Delta_a - \frac{\gamma}{2})}{(-1)^n 2iG^*} [\Gamma_n - i(\omega - \omega_n)]$$

for $n = 1, 2$.

IV. VACUUM RABI SPLITTING

Generally in vacuum Rabi splitting, the splitting frequency $\Delta\omega$ provides the information of the coupling g between the qubit and the resonator. As seen in Eq. (23), nonlinearity parameter χ modifies the decay rate Γ_1 (Γ_2) and central frequency ω_1 (ω_2) of two peaks in the spectrum $S_V(\omega)$. And in the limit of weak driving, the ξ does not affect the spectrum $S_V(\omega)$, it means that we can use this driven nonlinear Jaynes-Cummings model to characterize vacuum Rabi splitting effect very well. The corresponding splitting frequency between two peaks in the spectrum $S_V(\omega)$ is

$$\Delta\omega = |\omega_1 - \omega_2| = 2\text{Re}G \quad (24)$$

which is independent of the driving strength ξ and determined by the couple strength g , the detuning δ , decay rate κ (γ) and nonlinearity parameter χ .

Assuming nanomechanical resonator as an ideal resonator, $\chi = 0$, the spectrum $S_V(\omega)$ in Eq. (23) will be same as the previous results. [15] When the resonant condition ($\delta = 0$) and the strong-coupling limit ($g \gg [\kappa, \gamma]$) are adopted, we obtain the well-known splitting frequency [4]

$$\Delta\omega \simeq 2g. \quad (25)$$

To further clarify the dependence of correlation spectrum $S_V(\omega)$ on nonlinearity parameter χ and the driving strength ξ more clearly, the resonant condition is adopted, $\Delta_a = \Delta_c = 1.0$. The other parameters are $g = 0.2$, $\kappa = 0.004$, $\gamma = 0.004$ and 1 GHz is taken as the unit for all these parameters. [15, 24, 27]

Numerical calculations by QuTiP [28] are illustrated with some plots in the following. In Fig. 1, we choose the values of $\xi = 0.02$ and $\chi = 0.01$, the increasing of the number of total excitations N tells that the value of




FIG. 1: Correlation spectrum $S_V(\omega)$ versus ω for different values of excitations, we take $N = 1, 5, 10$. The other parameters are $\chi = 0.01, \xi = 0.02$.




FIG. 2: Correlation spectrum $S_V(\omega)$ versus ω for different values of the driving strength, we take $\xi = 0, 0.02, 0.04$. The other parameters are $\chi = 0$.

$N = 10$ is suitable in the other plots of the spectrum $S_V(\omega)$.

Figure 2 shows, in the limit of weak driving ($\xi \ll g$), there are two peaks in the spectrum and the increasing of the driving strength does not affect the spectrum obviously, which confirms the analytical calculations in Eq. (23). Then we take the value of $\xi = 0.02$ in the following plots.

As shown in Fig. 3, the increasing of nonlinearity parameter leads to the shift of central frequency of two peaks dramatically and does not change the distance between the centers of two peaks obviously because nonlinearity parameter is much smaller than the coupling g , we can see it in Eq. (24).

Figure 2 and Figure 3 demonstrate the dependence of the spectrum $S_V(\omega)$ on the driving strength ξ and nonlinearity parameter χ . These numerical results agree with those analytical results in Eq. (23), both of them main-




FIG. 3: Correlation spectrum $S_V(\omega)$ versus ω for different values of the driving strength, we take $\chi = 0, 0.02, 0.04$. The other parameters are $\xi = 0$.

tain that the weak driving strength ξ does not change the splitting frequency $\Delta\omega$, nonlinearity parameter χ changes the heights of two peaks ($\text{Re}\eta_1$ and $\text{Re}\eta_2$) and the shifts of central frequency (ω_1 and ω_2) obviously.

V. CONCLUSIONS

In this paper, vacuum Rabi splitting effect is studied to provide the information of the coupling g . Considering the intrinsic nonlinearity in nanomechanical resonator, a driven nonlinear Jaynes-Cummings model is used to describe the dynamics of the qubit-resonator system. Using quantum regression theorem, the dissipative dynamics of the qubit-resonator system is solved by master equation approach. Here, the two-time correlation spectrum is analytically calculated to clarify the dependence of correlation spectrum on the driving strength ξ and nonlinearity parameter χ . Because of small nonlinearity in nanomechanical resonator, we find that nonlinearity parameter leads to the shifts of central frequency (ω_1 and ω_2) and does not change the splitting frequency $\Delta\omega$ obviously. In Fig. 2 and Fig. 3, numerical results plotted by QuTiP agree with the analytical results in Eq. (23).

Acknowledgments

We thank the discussions with Professor P. Zhang.

[1] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, Cambridge, 1997).

[2] D. DiVincenzo, *Fortschr. Phys.* **48** (2000) 771.

- [3] E. T. Jaynes and F. W. Cummings, Proc. IEEE **51** (1963) 89.
- [4] J. M. Raimond, M. Brune, and S. Haroche, Rev. Mod. Phys. **73** (2001) 565.
- [5] F. Xue, Y. D. Wang, C. P. Sun, H. Okamoto, H. Yamaguchi, and K. Semba, New J. Phys. **9** (2007) 35.
- [6] A. Blais, R. S. Huang, A. Wallraff, S. M. Girvin, and R. J. Schoelkopf, Phys. Rev. A **69** (2004) 062320.
- [7] X. M. H. Huang, C. A. Zorman, M. Mehregany, and M. L. Roukes, Nature (London) **421** (2003) 496.
- [8] M. D. LaHaye, J. Suh, P. M. Echternach, K. C. Schwab, and M. L. Roukes, Nature (London) **459** (2009) 960.
- [9] Y. Makhlin, G. Schoen, and A. Shnirman, Rev. Mod. Phys. **73** (2001) 357.
- [10] J. Q. You and F. Nori, Phys. Today **58** (11) (2005) 42.
- [11] A. N. Cleland, *Foundations of Nanomechanics: From Solid-State Theory to Device Applications* (Springer-Verlag, Berlin, 2002).
- [12] A. D. Armour, M. P. Blencowe, K. C. Schwab, Phys. Rev. Lett. **88** (2002) 148301.
- [13] C. Cheng, Y. B. Gao, Commun. Theor. Phys. **60** (2013) 531.
- [14] X. Xiao, M. Y. Zhao, S. M. Yu, and Y. B. Gao, Commun. Theor. Phys. **65** (2016) 273.
- [15] Y. B. Gao, S. Yang, Y. X. Liu, C. P. Sun, and F. Nori, arxiv: 0902.2512.
- [16] L. F. Wei, Y. X. Liu, C. P. Sun, and F. Nori, Phys. Rev. Lett. **97** (2006) 237201.
- [17] Y. X. Liu, A. Miranowicz, Y. B. Gao, J. Bajer, C. P. Sun, and F. Nori, Phys. Rev. A **82** (2010) 032101.
- [18] L. G. Villanueva, R. B. Karabalin, M. H. Matheny, D. Chi, J. E. Sader, and M. L. Roukes, Phys. Rev. B **87** (2013) 024304.
- [19] I. Katz, A. Retzker, R. Straub, and R. Lifshitz, Phys. Rev. Lett. **99** (2007) 040404. V. Peano and M. Thorwart, Phys. Rev. B **70** (2004) 235401.
- [20] S. Rips, I. WilsonRae, and M. J. Hartmann, Phys. Rev. A **89** (2014) 013854.
- [21] F. R. Ong, M. Boissonneault, F. Mallet, A. C. Doherty, A. Blais, D. Vion, D. Esteve, and P. Bertet, Phys. Rev. Lett. **110** (2013) 047001.
- [22] M. Abdi, M. Pernpeintner, R. Gross, H. Huebl, and M. J. Hartmann, Phys. Rev. Lett. **114** (2015) 173602.
- [23] B. Teklu, A. Ferraro, M. Paternostro, and M. G. A. Paris, EPJ Quantum Technology **2** (2015) 16.
- [24] S. Rips and M. J. Hartmann, Phys. Rev. Lett. **110** (2013) 120503.
- [25] H. J. Carmichael, *Statistical Methods in Quantum Optics 1: Master Equations and Fokker-Planck Equations* (Springer, Berlin, 1999).
- [26] R. J. Brecha, P. R. Rice, and M. Xiao, Phys. Rev. A **59** (1999) 2392.
- [27] A. D. Armour, M. P. Blencowe, and K. C. Schwab, Phys. Rev. Lett. **88** (2002) 148301.
- [28] J. R. Johansson, P. D. Nation, and F. Nori, Comp. Phys. Comm. **184** (2013) 1234.