

ISOTROPIC GEOMETRY OF GRAPH SURFACES ASSOCIATED WITH PRODUCT PRODUCTION FUNCTIONS IN ECONOMICS

Muhittin Evren Aydin, Mahmut Ergut
Firat University and Namik Kemal University, Turkey

Abstract. A production function is a mathematical formalization in economics which denotes the relations between the output generated by a firm, an industry or an economy and the inputs that have been used in obtaining it. In this paper, we study the product production functions of 2 variables in terms of the geometry of their associated graph surfaces in the isotropic 3-space \mathbb{I}^3 . In particular, we derive several classification results for the graph surfaces of product production functions in \mathbb{I}^3 with constant curvature.

Key words: Production function, return to scale, relative curvature, isotropic mean curvature, isotropic space.

2010 Mathematics Subject Classification: 91B38, 53A35, 53A40, 53B25.

1 Introduction

In economics, a *production function* is a non-constant positive function that species the output of a firm, an industry, or an entire economy for all combinations of inputs. Explicitly, it is a map of class C^∞ which has non-vanishing first derivatives defined by

$$\begin{cases} h : \mathbb{R}_+^n \longrightarrow \mathbb{R}_+, & (x_1, x_2, \dots, x_n) \longmapsto h(x_1, x_2, \dots, x_n), \\ \mathbb{R}_+^n = \{(x_1, x_2, \dots, x_n) : x_i > 0, i = 1, \dots, n\}. \end{cases} \quad (1.1)$$

Here h denotes the quantity of output, n is the number of inputs and the variables are the inputs (such as capital, labor, raw materials etc.). Some interesting examples of production functions can be found in [18].

In order for the production functions to model as well economic reality, they are required to get some properties (see e.g. [5, 22]). One of the most important of these properties is the production function f to be homogeneous, i.e. there exist a real number p such that

$$h(\lambda x_1, \lambda x_2, \dots, \lambda x_n) = \lambda^p h(x_1, x_2, \dots, x_n), \quad \lambda \in \mathbb{R}_+. \quad (1.2)$$

(1.2) implies when the inputs are multiplied by same factor, the output is multiplied by some power of the factor.

If $p < 1$ (resp. $p > 1$) in (1.2), then the production function is said to have *decreasing* (resp. *increasing*) *return to scale*. If $p = 1$, then it is said to have *constant return to scale*.

The presence of increasing returns means that a one percent increase in the usage levels of all inputs would result in a greater than one percent increase in

output; the presence of decreasing returns means that it would result in a less than one percent increase in output. Constant returns to scale is the in-between case (cf. [8]).

A.D. Vilcu and G.E. Vilcu [26] completely classified the homogeneous production functions with constant proportional marginal rate of substitution. Further, the homogeneous production functions have been investigated via their associated graph hypersurfaces in [6]-[11], [29].

The most famous one among homogeneous production functions is *Cobb-Douglas production function*, introduced in 1928 by C. W. Cobb and P. H. Douglas [12]. In original form, it is given as

$$Y = bL^k C^{1-k},$$

where b presents the total factor productivity, Y the total production, L the labor input and C the capital input.

The generalized *Cobb-Douglas production function* of n variables is defined by

$$h(\mathbf{x}) = Ax_1^{\alpha_1} x_2^{\alpha_2} \dots x_n^{\alpha_n}, \mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n$$

where $A, \alpha_1, \alpha_2, \dots, \alpha_n > 0$. We note that h has constant return to scale if and only if $\sum_{i=1}^n \alpha_i = 1$.

X. Wang, Y. Fu [28] and A.D. Vilcu, G.E. Vilcu [24, 25] classified the graph hypersurfaces of the generalized Cobb-Douglas production functions with zero Gauss-Kronocker and mean curvature.

On the other hand, there are some non-homogeneous production functions, including the famous Spillman-Mitscherlich and transcendental production functions respectively defined by

$$\begin{cases} h(\mathbf{x}) = A [1 - \exp(-a_1 x_1)] \cdot [1 - \exp(-a_2 x_2)] \cdot \dots \cdot [1 - \exp(-a_n x_n)], \\ \mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n, A > 0, a_i < 0, i = 1, \dots, n \end{cases}$$

and

$$\begin{cases} h(\mathbf{x}) = Ax_1^{a_1} \exp(b_1 x_1) \cdot x_2^{a_2} \exp(b_2 x_2) \cdot \dots \cdot x_n^{a_n} \exp(b_n x_n), \\ \mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n, A > 0, a_i, b_i \in \mathbb{R}, a_i^2 + b_i^2 \neq 0, i = 1, \dots, n. \end{cases}$$

Such production functions, including the generalized Cobb-Douglas production function, belong to a more general class of production functions given by

$$h(\mathbf{x}) = \prod_{j=1}^n g_j(x_j), \mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n,$$

where g_j is a continuous positive real functions with nonzero first derivatives, $j = 1, \dots, n$. H. Alodan et al. [1] called these production functions *product production functions*. A production function is said to be *quasi-product*, if it is of the form

$$h(\mathbf{x}) = F\left(\prod_{j=1}^n g_j(x_j)\right), \mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}_+^n,$$

where F, g_j are continuous positive real functions with nonzero first derivatives, $j = 1, \dots, n$. A lot of classifications of the quasi-product and quasi-sum production functions can be found in [1]-[4], [14, 27] in terms of the geometry of their graph hypersurfaces.

B.-Y. Chen, et al [7, 13] investigated the graph hypersurfaces of the production models via the isotropic geometry. In the present paper, we classify the graph surfaces of product production functions of 2 variables with constant curvature in the isotropic 3-space \mathbb{I}^3 .

2 Basics on isotropic spaces

For later use, we provide a brief review of isotropic geometry from [7], [19]-[21].

The isotropic 3-space \mathbb{I}^3 is a Cayley–Klein space defined from a 3-dimensional projective space $P(\mathbb{R}^3)$ with the absolute figure which is an ordered triple (ω, f_1, f_2) , where ω is a plane in $P(\mathbb{R}^3)$ and f_1, f_2 are two complex-conjugate straight lines in ω . The homogeneous coordinates in $P(\mathbb{R}^3)$ are introduced in such a way that the absolute plane ω is given by $X_0 = 0$ and the absolute lines f_1, f_2 by $X_0 = X_1 + iX_2 = 0$, $X_0 = X_1 - iX_2 = 0$. The intersection point $F(0 : 0 : 0 : 1)$ of these two lines is called the absolute point. The group of motions G_6 of \mathbb{I}^3 is a six-parameter group given in the affine coordinates $x_1 = \frac{X_1}{X_0}$, $x_2 = \frac{X_2}{X_0}$, $x_3 = \frac{X_3}{X_0}$ by

$$(x_1, x_2, x_3) \mapsto (x'_1, x'_2, x'_3) : \begin{cases} x'_1 = a + x_1 \cos \phi - x_2 \sin \phi, \\ x'_2 = b + x_1 \sin \phi + x_2 \cos \phi, \\ x'_3 = c + dx_1 + ex_2 + x_3, \end{cases} \quad (2.1)$$

where $a, b, c, d, e, \phi \in \mathbb{R}$. Such affine transformations are called *isotropic congruence transformations* or *i-motions*. It is easily seen from (2.1) that i-motions are indeed composed by an Euclidean motion in the x_1x_2 -plane (i.e. translation and rotation) and an affine shear transformation in x_3 -direction.

In general, the following terminology is used for the isotropic spaces. Consider the points $\mathbf{x} = (x_1, x_2, x_3)$ and $\mathbf{y} = (y_1, y_2, y_3)$. The projection in \mathbf{z} -direction onto \mathbb{R}^2 , $(x_1, x_2, x_3) \mapsto (x_1, x_2, 0)$, is called the *top view*. In the sequel, many of metric properties in isotropic geometry (invariants under G_6) are Euclidean invariants in the top view such as the isotropic distance, so-called i-distance. *I-distance* of two points \mathbf{x} and \mathbf{y} is defined as the Euclidean distance of their top views, i.e.,

$$\|\mathbf{x} - \mathbf{y}\|_i = \sqrt{\sum_{j=1}^2 (y_j - x_j)^2}. \quad (2.2)$$

The i-metric is degenerate along the lines in \mathbf{z} -direction, and such lines are called *isotropic lines*. The plane containing an isotropic line is called an *isotropic plane*. Therefore, an *isotropic 3-space* \mathbb{I}^3 is the product of the Euclidean 2-space \mathbb{R}^2 and an isotropic line with a degenerate parabolic distance metric.

Let M^2 be a regular surface immersed in \mathbb{I}^3 which has no isotropic tangent planes. Such a surface M^2 is said to be *admissible* and can be parametrized by

$$r : D \subseteq \mathbb{R}^2 \longrightarrow \mathbb{I}^3 : (u_1, u_2) \longmapsto (r_1(u_1, u_2), r_2(u_1, u_2), r_3(u_1, u_2)), \quad (2.3)$$

where r_1, r_2 and r_3 are smooth real-valued functions on a domain $D \subseteq \mathbb{R}^2$. Denote g the metric on M^2 induced from \mathbb{I}^3 . The components of the first fundamental form of M^2 can be calculated via the induced metric g as follows

$$g_{ij} = g(r_{u_i}, r_{u_j}), \quad r_{u_i} = \frac{\partial r}{\partial u_i}, \quad i, j \in \{1, 2\}.$$

The unit normal vector field of M^2 is completely isotropic, i.e. $(0, 0, 1)$. Also, the components of the second fundamental form are (for details, see [20], p. 150-155)

$$t_{ij} = \frac{\det(r_{u_i u_j}, r_{u_1}, r_{u_2})}{\sqrt{\det(g_{ij})}}, \quad r_{u_i u_j} = \frac{\partial^2 r}{\partial u_i \partial u_j}, \quad i, j \in \{1, 2\}. \quad (2.4)$$

Thus the *relative curvature* (so-called the *isotropic curvature* or *isotropic Gaussian curvature*) and the *isotropic mean curvature* are respectively defined by

$$K = \frac{\det(t_{ij})}{\det(g_{ij})}, \quad H = \frac{g_{11}t_{22} - 2g_{12}t_{12} + g_{22}t_{11}}{\det(g_{ij})}. \quad (2.5)$$

A surface is called *isotropic minimal* (resp. *isotropic flat*) if its isotropic mean curvature (resp. relative curvature) vanishes.

In particular, if M^2 is a Monge surface in \mathbb{I}^3 of the form

$$(u_1, u_2) \longmapsto (u_1, u_2, h(u_1, u_2)),$$

then the metric on M^2 induced from \mathbb{I}^3 is given by $g_* = du_1^2 + du_2^2$. This implies that M^2 is always a flat space with respect to the induced metric g_* . Thus its Laplacian is given by

$$\Delta = \sum_{j=1}^2 \frac{\partial^2}{\partial u_j^2}.$$

Also, the matrix of second fundamental form of M^2 becomes the Hessian matrix of h (i.e. the square matrix $(h_{u_i u_j})$ of second-order partial derivatives of the function h). Thereby, the formulas in (2.5) reduce to

$$K = \det(h_{u_i u_j}) = h_{u_1 u_1} h_{u_2 u_2} - (h_{u_1 u_2})^2, \quad H = \Delta h = h_{u_1 u_1} + h_{u_2 u_2}. \quad (2.6)$$

3 Product production functions

Let us consider the product production function of 2 variables given by

$$h : \mathbb{R}_+^2 \longrightarrow \mathbb{R}_+, \quad (x, y) \longmapsto h(x, y) = f(x)g(y),$$

where f, g are continuous positive real functions with $f'(x) = \frac{df}{dx} \neq 0$ and $g'(y) = \frac{dg}{dy} \neq 0$. The graph surface M^2 corresponding to h is of the form

$$r(x, y) = (x, y, h(x, y) = f(x)g(y)), \quad (3.1)$$

which we call *product production surface*.

We remark that the surfaces of the form (3.1) are known as *factorable surfaces* or *homothetical surfaces* in ambient spaces and have been studied in [15, 17, 23, 30].

The following result provides a complete classification of the product production surfaces of 2 variables in \mathbb{I}^3 with constant relative curvature.

Theorem 3.1. *Let M^2 be a product production surface given by (3.1) in \mathbb{I}^3 with constant relative curvature K_0 .*

(A) *If $K_0 = 0$, then one of the following occurs:*

(A.1) *$h(x, y) = c_1 f(x)$ or $h(x, y) = c_2 g(y)$ for nonzero constants c_1, c_2 .*

(A.2) *h is a transcendental production function of 2 variables given by*

$$h(x, y) = A \exp(c_1 x + c_2 y),$$

where A, c_1, c_2 are nonzero constants.

(A.3) *Up to translations of x and y , h is a Cobb-Douglas production function of 2 variables with constant return to scale.*

(B) *If $K_0 \neq 0$, then it is negative ($K_0 < 0$) and, up to translations of x and y , h is a Cobb-Douglas production function of 2 variables given by*

$$h(x, y) = (-K_0)xy.$$

Proof. Assume that M^2 has constant relative curvature K_0 in \mathbb{I}^3 . Then, it follows from (2.6) that

$$(f''g'')fg - (f'g')^2 = K_0 \quad (3.2)$$

where $f' = \frac{df}{dx}$ and $g' = \frac{dg}{dy}$, etc. We divide the proof into two cases:

Case (i) $K_0 = 0$. Then, from (3.2), both situations, f or g constants, are solutions for (3.2). This implies the statement (A.1) of the theorem. Now, let us assume that f and g are non-constant functions. Hence, it follows from (3.2) that f and g cannot be linear functions. Thus the equation (3.2) can be rewritten as

$$\frac{f''f}{(f')^2} - \frac{(g')^2}{g''g} = 0,$$

which yields

$$\frac{ff''}{(f')^2} = \lambda = \frac{(g')^2}{g''g} \quad (3.3)$$

for a nonzero constant λ . In order to solve (3.3) we have to distinguish two situations.

Case (i.1) $\lambda = 1$. Then after solving (2.3) we get

$$f(x) = c_1 \exp(c_2x) \text{ and } g(y) = c_3 \exp(c_4y),$$

where c_i are nonzero constants, $1 \leq i \leq 4$, which gives the the statement (A.2) of the theorem.

Case (i.2) $\lambda \neq 1$. Solving (3.3) yields

$$f(x) = [(1 - \lambda)(c_1x + d_1)]^{\frac{1}{1-\lambda}} \text{ and } g(y) = \left[\left(\frac{\lambda - 1}{\lambda} \right) (c_2y + d_2) \right]^{\frac{\lambda}{\lambda-1}}$$

for nonzero constants c_3, c_4 and some constants d_1, d_2 . Up to suitable translations of x, y , we obtain

$$h(x, y) = f(x)g(y) = Ax^{\frac{1}{1-\lambda}}y^{-\frac{\lambda}{1-\lambda}}$$

for $A = [c_1(1 - \lambda)]^{\frac{1}{1-\lambda}} [c_2(\frac{\lambda-1}{\lambda})]^{\frac{\lambda}{\lambda-1}}$. This proves the statement (A.3) of the theorem.

Case (ii) $K_0 \neq 0$. Suppose that f and g are non-linear functions. Hence, we can rewrite (3.2) as

$$\frac{ff''}{(f')^2} - \frac{(g')^2}{gg''} = \frac{K_0}{(f')^2 gg''}. \quad (3.4)$$

Differentiating of (3.4) with respect to y gives

$$-\left(\frac{(g')^2}{gg''} \right)' = \frac{K_0}{(f')^2} \left(\frac{1}{gg''} \right)'. \quad (3.5)$$

From (3.5), if

$$\left(\frac{1}{gg''} \right)' = 0,$$

i.e. gg'' is a nonzero constant c in (3.5), then we get $\frac{1}{c} \left((g')^2 \right)' = 0$, which is not possible because g is non-linear function. In (3.5), if

$$\left(\frac{(g')^2}{gg''} \right)' = 0, \quad (3.6)$$

then (3.5) reduces to

$$\frac{K_0}{(f')^2} \left(\frac{1}{gg''} \right)' = 0,$$

which yields $gg'' = d$, $d \neq 0$. Considering this in (3.6) implies $\frac{2}{d}g'g'' = 0$ and it is a contradiction. Thereby we can rewrite (3.5) as

$$-\frac{\left(\frac{(g')^2}{gg''} \right)'}{\left(\frac{1}{gg''} \right)'} = \frac{K_0}{(f')^2}. \quad (3.7)$$

Since f is a non-linear function, the right-side of (3.7) is a function of x . However the left-side of (3.7) is either a constant or a function of y . Both cases are not possible.

Now let either f or g be a linear function. Without loss of generality, we may assume that f is a linear function, i.e. $f(x) = c_1x + d_1$, $c_1 \neq 0$, $d_1 \in \mathbb{R}$. Then we get from (3.2)

$$g' = \frac{\sqrt{-K_0}}{c_1}, \quad K_0 < 0.$$

This implies that g is also a linear function, i.e. $g(y) = \frac{\sqrt{-K_0}}{c_1}y + d_2$, $d_2 \in \mathbb{R}$. Thus, up to suitable translations of x and y , we derive

$$h(x, y) = \sqrt{-K_0}xy.$$

This gives of the statement (B) of the theorem.

Therefore, the proof is completed.

Next classifies the product production surfaces of constant isotropic mean curvature in \mathbb{I}^3 .

Theorem 3.2. *Let M^2 be a product production surface given by (3.1) in \mathbb{I}^3 with constant isotropic mean curvature H_0 . Then either*

(A)

$$h(x, y) = \frac{H_0}{g_0}x^2 + d_1x + d_2, \text{ (or } \frac{H_0}{f_0}y^2 + d_1y + d_2)$$

for $f_0, g_0 \in \mathbb{R} - \{0\}$, $d_1, d_2 \in \mathbb{R}$, or

(B) M^2 is isotropic minimal, i.e. $H_0 = 0$, and up translations of x and y , h is a Cobb-Douglas production function of 2 variables given by

$$h(x, y) = Axy, \quad A > 0.$$

Proof. Assume that M^2 has constant isotropic mean curvature H_0 . Then, by (2.6), we get

$$H_0 = f''g + fg''. \quad (3.8)$$

It follows from (3.8) that when g is a nonzero constant g_0 we have

$$f(x) = \frac{H_0}{g_0}x^2 + d_1x + d_2, \quad d_1, d_2 \in \mathbb{R},$$

and analogously if f is a nonzero constant f_0 , we deduce

$$g(y) = \frac{H_0}{f_0}y^2 + d_3y + d_4, \quad d_3, d_4 \in \mathbb{R},$$

which proves the statement (A) of the theorem.

Now suppose that f, g are non-constant functions. Then (3.8) can be rewritten as

$$\frac{f''}{f} + \frac{g''}{g} = \frac{H_0}{fg}. \quad (3.9)$$

After taking the partial derivative of (3.9) with respect to x , we deduce

$$\left(\frac{f''}{f}\right)' \frac{f^2}{f'} = -H_0 \frac{1}{g}. \quad (3.10)$$

The left-side of (3.10) is either a constant or a function of x while the other side is a function of y . This case is only possible when $H_0 = 0$ and

$$\left(\frac{f''}{f}\right)' = 0. \quad (3.11)$$

Similarly, taking the partial derivative of (3.9) with respect to y , we find $H_0 = 0$ and

$$\left(\frac{g''}{g}\right)' = 0. \quad (3.12)$$

This means that M^2 is isotropic minimal, i.e. $H_0 = 0$. Now let us assume that f and g are non-linear functions. By solving (3.11) and (3.12), we derive

$$f(x) = \beta_1 e^{\alpha_1 x} + \beta_2 e^{-\alpha_1 x} \quad \text{and} \quad g(y) = \beta_3 e^{\alpha_2 y} + \beta_4 e^{-\alpha_2 y} \quad (3.13)$$

for nonzero constants α_i, β_j , $i = 1, 2$ and $j = 1, 2, 3, 4$. Substituting (3.13) into (3.9) gives

$$\alpha_1^2 + \alpha_2^2 = 0,$$

which is not possible. Then the functions f and g are linear and up to translations x and y , we obtain $h(x, y) = Axy$, $A \neq 0$. This gives the proof.

Remark 3.3. For the product production function given by $h(x, y) = f(x)g(y)$, we have that f and g are nonconstant functions. Hence, while the statement (A.1) of Theorem 3.1 and the statement (A) of Theorem 3.2 are correct in mathematical perspective, in reality such product production functions do not exist.

Now let us consider Spillman-Mitscherlich and transcendental production functions of 2 variables respectively given by

$$\begin{cases} h(x, y) = A [1 - \exp(ax)] \cdot [1 - \exp(by)], \\ (x, y) \in \mathbb{R}_+^n, A > 0, a, b < 0 \end{cases} \quad (3.14)$$

and

$$\begin{cases} h(x, y) = Ax^{a_1} \exp(b_1 x_1) \cdot x_2^{a_2} \exp(b_2 x_2), \\ (x, y) \in \mathbb{R}_+^n, A > 0, a_i, b_i \in \mathbb{R}, a_i^2 + b_i^2 \neq 0, i = 1, 2. \end{cases} \quad (3.15)$$

From Theorem 3.1 and Theorem 3.2, we obtain the following results for the surfaces corresponding to these production functions.

Corollary 3.4. *Let h be a Spillman-Mitscherlich production function of 2 variables given by (3.14). Then the corresponding graph surface M^2 of h has neither constant relative nor constant isotropic curvature in \mathbb{I}^3 .*

Corollary 3.5. *Let h be a transcendental production function of 2 variables given by (3.15) and M^2 its associated graph surface in \mathbb{I}^3 . Then:*

(A) M^2 has constant relative curvature K_0 in \mathbb{I}^3 if and only if $K_0 = 0$ and one of the following occurs:

(A.1) $\alpha_1 = \alpha_2 = 0$ and $b_1 \neq 0 \neq b_2$, or

(A.2) $\alpha_1 + \alpha_2 = 1$, $a_1 \neq 0 \neq a_2$ and $b_1 = b_2 = 0$.

(B) M^2 has constant isotropic curvature H_0 in \mathbb{I}^3 if and only if $H_0 = 0$ and $a_1 = a_2 = 1$ $b_1 = b_2 = 0$.

References

- [1] H. Alodan, B.-Y. Chen, S. Deshmukh, G.E. Vilcu, *On some geometric properties of quasi-product production models*, arXiv:1512.05190v1 [math.DG], 2015.
- [2] M.E. Aydin, M. Ergut, *Hessian determinants of composite functions with applications for production functions in economics*, Kragujevac J. Math. **38(2)** (2014), 259-268.
- [3] M.E. Aydin, A. Mihai, *Classifications of quasi-sum production functions with Allen determinants*, Filomat **29(6)** (2015), 1351-1359.
- [4] M.E. Aydin, M. Ergut, *Composite functions with Allen determinants and their applications to production models in economics*, Tamkang J. Math. **45(4)** (2014), 427-435.
- [5] R. G. Chambers, *Applied Production Analysis*, Cambridge University Press, 1998.
- [6] B.-Y. Chen, G. E. Vilcu, *Geometric classifications of homogeneous production functions*, Appl. Math. Comput. **225** (2013), 345-351.

- [7] B.-Y. Chen, S. Decu, L. Verstraelen, *Notes on isotropic geometry of production models*, Kragujevac J. Math. **38(1)** (2014), 23–33.
- [8] B.-Y. Chen, *On some geometric properties of h -homogeneous production function in microeconomics*, Kragujevac J. Math. **35(3)** (2011), 343–357.
- [9] B.-Y. Chen, *On some geometric properties of quasi-sum production models*, J. Math. Anal. Appl. **392** (2012), 192–199.
- [10] B.-Y. Chen, *Classification of h -homogeneous production functions with constant elasticity of substitution*, Tamkang J. Math. **43** (2012), 321–328.
- [11] B.-Y. Chen, *Solutions to homogeneous Monge-Ampere equations of homothetic functions and their applications to production models in economics*, J. Math. Anal. Appl. **411** (2014), 223–229.
- [12] C. W. Cobb, P. H. Douglas, *A theory of production*, Amer. Econom. Rev. **18** (1928), 139–165.
- [13] S. Decu, L. Verstraelen, *A note on the isotropical geometry of production surfaces*, Kragujevac J. Math. **37(2)** (2013), 217–220.
- [14] Y. Fu, *Geometric characterizations of quasi-product production models in economics*, Filomat (2016), to be published.
- [15] R. Lopez, M. Moruz, *Translation and homothetical surfaces in Euclidean space with constant curvature*, J. Korean Math. Soc. **52(3)** (2015), 523–535.
- [16] L. Losonczi, *Production functions having the CES property*, Acta Math. Acad. Paedagog. Nyháí. (N.S.) **26(1)** (2010), 113–125.
- [17] H. Meng, H. Liu, *Factorable surfaces in Minkowski space*, Bull. Korean Math. Soc. **46(1)** (2009), 155–169.
- [18] S.K. Mishra, *A brief history of production functions*, IUP J. Manage. Econom. **8(4)** (2010), 6–34.
- [19] H. Pottmann, P. Grohs, N.J. Mitra, *Laguerre minimal surfaces, isotropic geometry and linear elasticity*, Adv. Comput. Math. **31** (2009), 391–419.
- [20] H. Sachs, *Isotrope Geometrie des Raumes*, Vieweg, 1990.
- [21] Z.M. Sipus, *Translation surfaces of constant curvatures in a simply isotropic space*, Period. Math. Hung. **68** (2014), 160–175.
- [22] A. Thompson, *Economics of the Firm, Theory and Practice*, 3rd edition, Prentice Hall, 1981.
- [23] I. Van de Woestyne, *Minimal homothetical hypersurfaces of a semi-Euclidean space*, Results Math. **27** (1995), 333–342.

- [24] G.E. Vilcu, *A geometric perspective on the generalized Cobb–Douglas production functions*, Appl. Math. Lett. 24 (2011), 777–783.
- [25] A. D. Vilcu, G. E. Vilcu, *On some geometric properties of the generalized CES production functions*, Appl. Math. Comput. **218** (2011), 124–129.
- [26] A. D. Vilcu, G. E. Vilcu, *On homogeneous production functions with proportional marginal rate of substitution*, Math. Probl. Eng. 2013 (2013), Article ID 732643, 5 pages.
- [27] A.D. Vilcu, G.E. Vilcu, *Some characterizations of the quasi-sum production models with proportional marginal rate of substitution*, C. R. Math. Acad. Sci. Paris **353** (2015), 1129-1133.
- [28] X. Wang, Y. Fu, *Some characterizations of the Cobb-Douglas and CES production functions in microeconomics*, Abstr. Appl. Anal. 2013 (2013), Article ID 761832, 6 pages.
- [29] X. Wang, *A geometric characterization of homogeneous production models in economics*, Filomat (2016), to be published.
- [30] Y. Yu, H. Liu, *The factorable minimal surfaces*, Proceedings of The Eleventh International Workshop on Diff. Geom. 11 (2007), 33-39.

M.E. Aydin
Department of Mathematics
Firat University
23119 Elazig
Turkey
E-mail: meaydin@firat.edu.tr

M. Ergut
Department of Mathematics
Namik Kemal University
59 000 Tekirdag
Turkey
E-mail: mergut@nku.edu.tr