

Bayesian Quantile Regression for Ordinal Longitudinal Data

Rahim Alhamzawi*

Abstract. Since the pioneering work by [Koenker and Bassett \(1978\)](#), quantile regression models and its applications have become increasingly popular and important for research in many areas. In this paper, a random effects ordinal quantile regression model is proposed for analysis of longitudinal data with ordinal outcome of interest. An efficient Gibbs sampling algorithm was derived for fitting the model to the data based on a location-scale mixture representation of the skewed double exponential distribution. The proposed approach is illustrated using simulated data and a real data example. This is the first work to discuss quantile regression for analysis of longitudinal data with ordinal outcome.

Keywords: Bayesian inference, Cut-points, Longitudinal data, Ordinal regression, Quantile regression .

1 Introduction

The mean value has long been used as a measure of location of the center of a distribution. However, in many applications, there are occasions when analysts are interested to observe and analyze different points of the distribution. The distribution function F of a random variable can be characterized by infinite number of points spanning its support. These points are called quantiles. Thus, quantiles are points taken at regular intervals from F . The θ th quantile of a data distribution, $\theta \in (0, 1)$, is interpreted as the value such that there is $100(1 - \theta)\%$ of mass on its right side and $100\theta\%$ of mass on its left side. In particular, for a continuous random variable Y , the $100\theta\%$ quantile of F is the value y which solves $F(y) = \theta$ (we assume that this value is unique), where $F(y) = P(Y \leq y)$. Thus, the population lower quartile, median and upper quartile are the solutions to the equations $F(y) = \frac{1}{4}$, $F(y) = \frac{1}{2}$ and $F(y) = \frac{3}{4}$, respectively. Compared to mean value, quantiles are useful measures because they are less susceptible to skewed distributions and outliers. This fact form the building block of quantile regression which unlike its standard mean regression counterpart, lies in its flexibility and ability in providing a more complete investigation of the entire conditional distribution of the response variable distribution given its predictors. To this end, quantile regression has become increasingly popular since the pioneering research of [Koenker and Bassett \(1978\)](#).

After its introduction, quantile regression has attracted considerable attention in recent literature. It has been applied in a wide range of fields such as agriculture ([Kostov and Davidova, 2013](#)), body mass index ([Bottai et al., 2014](#)), microarray study

*Statistics Department, College of Administration and Economics, Al-Qadisiyah University, Al Diwaniyah, Iraq

(Wang and He, 2007), financial portfolio (Mezali and Beasley, 2013), economics (Hendricks and Koenker, 1992), ecology (Cade and Noon, 2003), climate change (Reich, 2012), survival analysis (Koenker and Geling, 2001) and so on. A comprehensive account of other recent applications of quantile regression can be found in Yu et al. (2003) and Koenker (2005).

Longitudinal data is encountered in a wide variety of applications, including economics, finance, medicine, psychology and sociology. It is repeatedly measured from independent subjects over time and correlation arises between measures from the same subject. Since the pioneering work by Laird and Ware (1982), the mixed models with random effects have become common and active models to deal with longitudinal data. A number of books and a vast number of research papers published in this area have been motivated by Laird and Ware's mixed models. The majority of these books and research papers focuses on standard mean regression. See for example, Wolfinger and O'connell (1993), Verbeke and Lesaffre (1996), Hedges and Vevea (1998), Tao et al. (1999), McCulloch and Neuhaus (2001), Hedeker and Gibbons (2006) and Baayen et al. (2008), among others. In contrast, limited research papers have been conducted on quantile regression for longitudinal data. For example, Koenker (2004) proposed the l_1 regularization quantile regression model, Lipsitz et al. (1997) studied quantile regression for longitudinal data in different contexts and developed resampling approaches for inference. Geraci and Bottai (2007) suggested a Bayesian quantile regression method for longitudinal data using the skewed Laplace distribution (SLD) for the errors, Reich et al. (2010) proposed a flexible Bayesian quantile regression method for dependent and independent data using an infinite mixture of normals for the errors, Yuan and Yin (2010) studied quantile regression for longitudinal data with nonignorable intermittent missing data and Alhamzawi and Yu (2014) proposed a method for regularization in mixed quantile regression models.

Longitudinal data with ordinal responses routinely appear in many applications, including economics, psychology and sociology. Existing approaches in classical mean regression are typically designed to deal with such data. At present time, the most common method in classical mean regression for modelling such data employs the cumulative logit model. There exists a large literature on the analysis of longitudinal data with ordinal responses, and we refer to Fitzmaurice et al. (2012) for an overview. In contrast, quantile regression approaches for estimating the parameters of ordinal longitudinal data have not been proposed, yet. The goal of this paper is to fill this gap by introducing an ordinal random effects quantile regression model that is appropriate for such data.

In Section 2, we present a random effects ordinal quantile regression model for analysis of longitudinal data with ordinal outcome by using a data augmentation method. We also discuss prior elicitation. In Section 3, we outline the Bayesian estimation method via Gibbs sampler. In Section 4, we carry out simulation scenarios to investigate the performance of the proposed method, and in Section 5, we illustrate our proposed method using a real dataset. We conclude the paper with a brief discussion in Section 6. An appendix contains the Gibbs sampler details.

2 Methods

2.1 Quantile Regression

Given training data $\{(\mathbf{x}_i, y_i), i = 1, \dots, N\}$, with covariate vector $\mathbf{x}_i \in R^p$ and outcome of interest $y_i \in R$. The θ th quantile regression model for the response y_i given the covariate vector \mathbf{x}_i takes the form of

$$Q_{y_i}(\theta|\mathbf{x}_i) = \mathbf{x}_i' \boldsymbol{\beta}, \quad (1)$$

where $Q_{y_i}(\theta|\mathbf{x}_i) = F_{y_i}^{-1}(\theta|\mathbf{x}_i)$ is the inverse Cumulative Distribution Function (CDF) and $\boldsymbol{\beta}$ is the unknown quantile coefficients vector. This is what makes the quantile estimators can be considered nonparametric maximum likelihood estimators (Wasserman, 2006).

Unlike the standard mean regression, the error term does not appear in (1) because all the random variation in the conditional distributions is accounted for by variation in the θ th quantile, $\theta \in (0, 1)$. Consequently, quantile regression does not require any assumption about the distribution of the errors and, unlike standard mean regression, is more robust to outliers and non-normal errors, offering greater statistical efficiency if the data have outliers or is non-normal. It belongs to a robust model family, which can provide a more complete picture of the predictor effects at different quantile levels of the response variable rather than focusing solely on the center of the distribution (Yu et al., 2003). One attractive feature of quantile regression is that the linear quantile regression model (Koenker and Bassett, 1978) can be used to estimate the parameters of the nonlinear model because the quantile regression estimators are equivariant to linear or nonlinear monotonic transformations of the response variable, i.e., $Q_{\log_{10} y_i}(\theta|\mathbf{x}_i) = \log_{10} Q_{y_i}(\theta|\mathbf{x}_i)$. In general, this is a very important property since it tells us that quantile regression provides consistent back transformation and easy in interpretation in the case of transformations such as the logarithm and the square root.

The quantile estimators have the same interpretation as those of a standard mean regression model except for the indexed quantile levels where each is estimated (Cade et al., 2008). For example, if the slope is -0.78 for the response variable y given the predictor x in the 95th quantile would indicate that the 95th quantile of the response variable decreased by 0.78 for each 1 unit increase in x . The unknown quantity $\boldsymbol{\beta}$ is estimated by

$$\min_{\boldsymbol{\beta}} \sum_{i=1}^N \rho_{\theta}(y_i - \mathbf{x}_i' \boldsymbol{\beta}), \quad (2)$$

where $\rho_{\theta}(t) = t\theta - tI(t < 0)$ is the check loss function (CLF) at a quantile θ , $0 < \theta < 1$, and $I(\cdot)$ is the indicator function. By contrast, standard mean regression method based on the quadratic loss t^2 . Koenker and Bassett (1978) observed that the CLF (2) is closely related to the skewed Laplace distribution (SLD) and consequently the unknown quantity $\boldsymbol{\beta}$ can be estimated through exploiting this link. This observation opens new

avenues when dealing with quantile regression and its applications. The density function of a SLD is

$$f(\varepsilon|\theta) = \theta(1 - \theta) \exp \left\{ -\rho_\theta(\varepsilon) \right\}, \quad -\infty < \varepsilon < \infty. \quad (3)$$

Minimizing the CLF (2) is equivalent to maximizing the likelihood function of $\varepsilon_i = y_i - \mathbf{x}'_i \boldsymbol{\beta}$ by assuming ε_i from a SLD. By utilizing the link between the CLF (2) and the SLD, [Yu and Moyeed \(2001\)](#) proposed a Bayesian framework for quantile regression using the SLD for the error distribution and show the propriety of the conditional distribution of $\boldsymbol{\beta}$ under an improper prior distribution. Unfortunately, the joint posterior distribution under this framework does not have a known tractable form and consequently [Yu and Moyeed \(2001\)](#) update the unknown quantity $\boldsymbol{\beta}$ from its posterior using the Metropolis-Hastings (M-H) algorithm. In this context, [Kozumi and Kobayashi \(2011\)](#) proposes a simple and efficient Gibbs sampling algorithm for updating $\boldsymbol{\beta}$ by motivating the SLD as a member of the scale mixture of normals. If $\varepsilon_i \sim N((1-2\theta)v_i, 2v_i)$, then the SLD for ε_i arises when v_i has an exponential distribution with rate parameter $\theta(1-\theta)$. Under this formulation, [Yue and Rue \(2011\)](#) presented a Bayesian framework for structured additive quantile regression models, [Luo et al. \(2012\)](#) developed Bayesian quantile regression for longitudinal data and [Alhamzawi and Yu \(2014\)](#) presented Bayesian Lasso mixed quantile regression.

2.2 Modeling Ordinal Longitudinal Data

Let $\mathbf{y} = \{y_{ij}\}(i = 1, \dots, N; j = 1, \dots, n_i)$, where y_{ij} denote the response for the i th subject measured at the j th time. Then, the θ th quantile regression model for ordinal longitudinal data can be formulated in terms of an ordinal latent variable l_{ij} as follows:

$$Q_{l_{ij}|\alpha_i}(\theta|\mathbf{x}_{ij}, \alpha_i) = \alpha_i + \mathbf{x}'_{ij} \boldsymbol{\beta} \quad (4)$$

where $Q_{l_{ij}|\alpha_i}(\theta|\mathbf{x}_{ij}, \alpha_i)$ is the inverse CDF of the unobserved latent response l_{ij} conditional on a location-shift random effect α_i , $\alpha_i \sim N(0, \phi)$, and \mathbf{x}_{ij} is a vector of predictors. The observed ordinal response y_{ij} is assumed to be related to the unobserved response l_{ij} by

$$y_{ij} = \begin{cases} 1 & \text{if } \delta_0 < l_{ij} \leq \delta_1; \\ c & \text{if } \delta_{c-1} < l_{ij} \leq \delta_c, \quad c = 2, \dots, C-1; \\ C & \text{if } \delta_{C-1} < l_{ij} < \delta_C; \end{cases}$$

where $\delta_0, \dots, \delta_C$ are cut-points whose coordinates satisfy $-\infty = \delta_0 < \delta_1 < \dots < \delta_{C-1} < \delta_C = +\infty$. Here, δ_{c-1} and δ_c are respectively defines the lower and upper bounds of the interval corresponding to observed outcome c .

Assuming that the error ε_{ij} of the unobserved response l_{ij} has a SLD as in (3), we have $\varepsilon_{ij} = (1-2\theta)v_{ij} + \sqrt{2v_{ij}}\epsilon_{ij}$ ([Kozumi and Kobayashi, 2011](#)). Here, the latent variable v_{ij} follows an exponential distribution with rate parameter $\theta(1-\theta)$, and ϵ_{ij} follows the standard normal distribution. Then, the CDF for the c category of the observed response y_{ij} is:

$$P(y_{ij} \leq c | l_{ij}, \delta_c) = P(l_{ij} \leq \delta_c | \boldsymbol{\beta}, \alpha_i, v_{ij}),$$

$$\begin{aligned}
&= P(\alpha_i + \mathbf{x}'_{ij}\boldsymbol{\beta} + (1 - 2\theta)v_{ij} + \sqrt{2v_{ij}}\epsilon_{ij} \leq \delta_c), \\
&= P\left(\epsilon_{ij} \leq \frac{\delta_c - \alpha_i - \mathbf{x}'_{ij}\boldsymbol{\beta} - (1 - 2\theta)v_{ij}}{\sqrt{2v_{ij}}}\right), \\
&= \Phi\left(\frac{\delta_c - \alpha_i - \mathbf{x}'_{ij}\boldsymbol{\beta} - (1 - 2\theta)v_{ij}}{\sqrt{2v_{ij}}}\right),
\end{aligned}$$

where Φ is the standard normal CDF. Using Φ , we can calculate $P(y_{ij} = c | l_{ij}, \delta_{c-1}, \delta_c)$ as follows:

$$\begin{aligned}
P(y_{ij} = c | l_{ij}, \delta_{c-1}, \delta_c) &= P(\delta_{c-1} < l_{ij} \leq \delta_c | \boldsymbol{\beta}, \alpha_i, v_{ij}), \\
&= \Phi\left(\frac{\delta_c - \alpha_i - \mathbf{x}'_{ij}\boldsymbol{\beta} - (1 - 2\theta)v_{ij}}{\sqrt{2v_{ij}}}\right) - \Phi\left(\frac{\delta_{c-1} - \alpha_i - \mathbf{x}'_{ij}\boldsymbol{\beta} - (1 - 2\theta)v_{ij}}{\sqrt{2v_{ij}}}\right).
\end{aligned}$$

2.3 Priors

Prior distribution selection is an essential step in any Bayesian inference; however, in the Bayesian paradigm it is particularly crucial as issues can arise when default prior distributions are used without caution (Kinney and Dunson, 2007). For the fixed effects $\boldsymbol{\beta}$, a typical choice is to assign a zero mean normal prior distribution on each $\beta_k, k = 1, 2, \dots, p$, which leads to the ridge estimator. However, this prior performs poorly if there are big differences in the size of fixed effects (Griffin et al., 2010). An generalization of the ridge prior is a Laplace prior, which is equivalent to the Lasso model (Tibshirani, 1996; Bae and Mallick, 2004). This prior has received considerable attention in the recent literature (for example see, Bae and Mallick (2004); Park and Casella (2008); Hans (2009); Li et al. (2010); Griffin et al. (2010)). In this paper, we assign a Laplace prior on each β_k takes the form of

$$P(\boldsymbol{\beta} | \lambda) = \prod_{k=1}^p \frac{\lambda}{2} e^{-\lambda|\beta_k|}, \quad \lambda \geq 0. \quad (5)$$

According to Andrews and Mallows (1974), the prior (5) can be written as

$$\prod_{k=1}^p \frac{\lambda}{2} e^{-\lambda|\beta_k|} = \prod_{k=1}^p \int_0^{\infty} \text{N}(\beta_k; 0, s_k) \text{Exp}(s_k; \lambda^2/2) ds_k. \quad (6)$$

From (6), it can be seen that we assign a zero-mean normal prior distribution with unknown variance for each β_k . We specify an exponential prior distributions with rate parameter $\lambda^2/2$ for the variances assuming they are independent. Then, we put a gamma prior on λ^2 with shape parameter a_1 and rate parameter a_2 . Since $\alpha_i \sim \text{N}(0, \phi)$, this motivates us to consider an inverse gamma prior on ϕ with shape parameter b_1 and scale parameter b_2 .

Following Montesinos-López et al. (2015) and Sorensen et al. (1995), we consider an order statistics from $U(\delta_0, \delta_C)$ distribution, for the $C - 1$ unknown cut-points:

$$P(\boldsymbol{\delta}) = (C - 1)! \left(\frac{1}{\delta_{\max} - \delta_{\min}} \right)^{C-1} I(\boldsymbol{\delta} \in T), \quad (7)$$

where $\boldsymbol{\delta} = (\delta_0, \delta_1, \dots, \delta_C)$ and $T = \{(\delta_{\min}, \delta_1, \dots, \delta_{\max}) | \delta_{\min} < \delta_1 < \dots < \delta_{C-1} < \delta_{\max}\}$. Because $l_{ij} \sim N(\alpha_i + \mathbf{x}'_{ij}\boldsymbol{\beta} + (1 - 2\theta)v_{ij}, 2v_{ij})$ and we observe $y_{ij} = c$ if $\delta_{c-1} < l_{ij} < \delta_c$, the posterior distribution of all the parameters and latent variables is given by

$$\begin{aligned} P(\boldsymbol{\beta}, \boldsymbol{\alpha}, \mathbf{l}, \boldsymbol{\delta}, \mathbf{v}, \mathbf{s}, \lambda^2, \phi | \mathbf{y}) &\propto P(\mathbf{y} | \mathbf{l}, \boldsymbol{\delta}) P(\mathbf{l} | \boldsymbol{\beta}, \boldsymbol{\alpha}, \mathbf{v}) P(\boldsymbol{\delta}) P(\mathbf{v}) \\ &\times P(\boldsymbol{\beta} | \mathbf{s}) P(\mathbf{s} | \lambda^2) P(\lambda^2) P(\boldsymbol{\alpha} | \phi) P(\phi), \end{aligned} \quad (8)$$

where, $\mathbf{y} = (y_{11}, \dots, y_{N n_N})$, $\mathbf{v} = (v_{11}, \dots, v_{N n_N})$, $\mathbf{l} = (l_{11}, \dots, l_{N n_N})$, $\boldsymbol{\alpha} = (\alpha_1, \dots, \alpha_N)$ and $\mathbf{s} = (s_1, \dots, s_p)$.

The full conditional distributions for $\boldsymbol{\beta}$, $\boldsymbol{\alpha}$, \mathbf{l} , $\boldsymbol{\delta}$, \mathbf{v} , \mathbf{s} , λ^2 and ϕ are summarized below and details of all derivations are provided in Appendix A.

3 Gibbs Sampler

Using the data augmentation procedure as in [Albert and Chib \(1993\)](#), a Gibbs sampling method for the ordinal quantile regression model with longitudinal data is constructed by updating $\boldsymbol{\beta}$, $\boldsymbol{\alpha}$, \mathbf{l} , $\boldsymbol{\delta}$, \mathbf{v} , \mathbf{s} , λ^2 , and ϕ from their full conditional distributions. From (8), we can construct a tractable algorithm for efficient posterior computation that works as follows:

1. Sample v_{ij} ($i = 1, \dots, N, j = 1, \dots, n_i$) from the generalized inverse Gaussian distribution $GIG(\nu, \varrho_1, \varrho_2)$, where $\varrho_1^2 = (l_{ij} - \mathbf{x}'_{ij}\boldsymbol{\beta} - \alpha_i)^2/2$ and $\varrho_2^2 = 1/2$.
2. Sample β_k ($k = 1, 2, \dots, p$) from $N(\mu_{\beta_k}, \sigma_{\beta_k}^2)$ where

$$\sigma_{\beta_k}^2 = \left(\sum_{i=1}^N \sum_{j=1}^{n_i} \frac{x_{ijk}^2}{2v_{ij}} + \frac{1}{s_k} \right)^{-1},$$

and

$$\mu_{\beta_k} = \sigma_{\beta_k}^2 \sum_{i=1}^N \sum_{j=1}^{n_i} \frac{(l_{ijk} - \sum_{h=1, h \neq k}^p x_{ijh}\beta_h - \alpha_i - (1 - 2\theta)v_{ij}) x_{ijk}}{2v_{ij}}.$$

3. Sample s_k , ($k = 1, 2, \dots, p$) from $GIG(0.5, \varrho_1, \varrho_2)$, where $\varrho_1^2 = \beta_k^2$ and $\varrho_2^2 = \lambda^2$.
4. Sample λ^2 from Gamma distribution with shape parameter $p + a_1$ and rate parameter $\sum_{k=1}^p s_k/2 + a_2$.
5. Sample α_i ($i = 1, \dots, N$) from $N(\mu_{\alpha_i}, \sigma_{\alpha_i}^2)$, where

$$\sigma_{\alpha_i}^2 = \left(\sum_{j=1}^{n_i} \frac{1}{2v_{ij}} + \frac{1}{\phi} \right)^{-1},$$

and

$$\mu_{\alpha_i} = \sigma_{\alpha_i}^2 \sum_{j=1}^{n_i} \frac{(l_{ij} - \mathbf{x}'_{ij}\boldsymbol{\beta} - \xi v_{ij})}{2v_{ij}}.$$

6. Sample ϕ from inverse Gamma distribution with shape parameter $\frac{N}{2} + b_1$ and scale parameter $\sum_{i=1}^N \frac{\alpha_i^2}{2} + b_2$.
7. Sample $l_{ij} (i = 1, \dots, N, j = 1, \dots, n_i)$ from truncated normal (TN) distribution $\text{TN}_{(\delta_{c-1}, \delta_c)}(\alpha_i + \mathbf{x}'_{ij}\boldsymbol{\beta} + (1 - 2\theta)v_{ij}, 2v_{ij})$.
8. Sample δ_c from a uniform distribution on the interval $[\min\{\min(l_{ij}|y_{ij} = c + 1), \delta_{c+1}, \delta_{\max}\}, \max\{\max(l_{ij}|y_{ij} = c), \delta_{c-1}, \delta_{\min}\}]$.

The details of the proposed Gibbs sampler algorithm and fully conditional posterior distributions are given in Appendix A.

4 Simulation Studies

We carry out a Monte Carlo simulation studies to assess the performance of the proposed method. We compared the proposed Bayesian quantile regression method for ordinal longitudinal data, referred to as ‘‘BQOL’’, with Bayesian Quantile Regression for Ordinal Models (BQROR) reported by [Rahman \(2016\)](#). The results of Bayesian logistic ordinal regression (BLOR) for longitudinal data and the maximum likelihood logistic ordinal regression (MLE) were also reported. Models were assessed based on the relative average bias and the estimated relative efficiency. Suppose that we are interested in the estimation of a vector of parameters $\boldsymbol{\psi} = (\psi_1, \psi_2, \dots, \psi_m)$. Then, the relative average bias of $\psi_h (h = 1, 2, \dots, m)$ is given by

$$\widehat{\text{bias}}(\hat{\psi}_h) = \frac{1}{M} \sum_{r=1}^M \frac{\hat{\psi}_h^r - \psi_h}{|\psi_h|},$$

and the estimated relative efficiency

$$\widehat{\text{eff}}_{\text{model}}(\hat{\psi}_h) = \frac{S_{\text{model}}^2(\hat{\psi}_h)}{S_{\text{BQOL}}^2(\hat{\psi}_h)},$$

where M denotes the number of replications, $\hat{\psi}_h^r$ is the parameter estimate for the r th replication, ψ_h is the true value, $S^2(\hat{\psi}_h) = \frac{1}{M} \sum_{r=1}^M (\hat{\psi}_h^r - \bar{\psi}_h)^2$ and $\bar{\psi}_h = \frac{1}{M} \sum_{r=1}^M \hat{\psi}_h^r$.

4.1 Simulation 1

Here, we follow the same simulation strategy introduced by [Montesinos-López et al. \(2015\)](#). Specifically, we simulated data from the following liability:

$$l_{ij} = \beta_1 x_{1ij} + \beta_2 x_{2ij} + \beta_3 x_{3ij} + \varepsilon_{ij}, \quad (i = 1, \dots, 40; j = 1, \dots, n_i)$$

where x_{1ij} and x_{2ij} were sampled independently from a uniform distribution on the interval $[-0.1, 0.1]$, $(\beta_1, \beta_2, \beta_3) = (-5, -10, 15)$ and ε_{ij} were sampled independently

Table 1: Estimated relative bias and relative efficiency for the simulated data 1. The proposed model (BQOL) is compared with three other models: the Bayesian quantile regression for ordinal models (BQROR), Bayesian logistic ordinal regression (BLOR) for longitudinal data and the maximum likelihood logistic ordinal regression (MLE)

n_i	Parameter	BQOL $_{\theta=0.5}$		BQROR $_{\theta=0.5}$		BLOR		MLE	
		bias	eff	bias	eff	bias	eff	bias	eff
5	β_1	-0.051	1.000	-0.079	1.103	0.082	1.012	-0.068	1.009
	β_2	-0.002	1.000	0.082	1.113	0.093	1.034	-0.134	1.071
	β_3	0.019	1.000	0.169	1.350	0.110	1.131	0.167	1.025
	δ_1	-0.012	1.000	-0.011	1.091	-0.036	1.113	-0.031	1.004
	δ_2	0.003	1.000	-0.010	1.014	-0.009	1.013	-0.016	1.037
	δ_3	0.003	1.000	-0.024	1.213	-0.001	0.944	0.002	1.056
	δ_4	0.012	1.000	-0.011	1.203	0.013	1.147	0.016	1.007
	β_1	-0.058	1.000	-0.099	1.128	-0.165	1.211	-0.119	1.018
10	β_2	0.022	1.000	0.132	1.059	-0.033	1.067	-0.100	1.009
	β_3	0.020	1.000	0.026	1.056	-0.055	1.029	-0.032	1.035
	δ_1	-0.002	1.000	0.009	1.007	-0.001	1.004	-0.002	1.021
	δ_2	-0.001	1.000	0.004	0.999	-0.003	0.998	0.002	1.048
	δ_3	-0.003	1.000	0.038	1.006	0.002	1.003	-0.003	1.025
	δ_4	-0.001	1.000	0.035	1.014	0.000	0.998	0.006	1.008
	β_1	-0.011	1.000	-0.032	1.117	-0.030	1.071	-0.107	1.051
	β_2	-0.010	1.000	0.077	1.019	-0.162	1.131	-0.116	1.119
20	β_3	0.004	1.000	0.042	1.107	0.029	1.224	0.108	1.103
	δ_1	-0.005	1.000	-0.016	1.008	-0.001	0.999	-0.003	0.999
	δ_2	0.003	1.000	-0.017	1.015	0.014	1.007	-0.004	1.028
	δ_3	0.000	1.000	-0.029	1.073	-0.002	0.996	-0.016	1.008
	δ_4	0.003	1.000	-0.036	1.088	0.002	0.999	-0.006	1.034

from a logistic distribution with location parameter $\mu = 0$ and scale parameter $s = 1$. Three values of n_i were considered, $n_i = 5, 10$, and 20 . The cut-points used were $\delta_1 = -0.8416$, $\delta_2 = -0.2533$, $\delta_3 = 0.2533$ and $\delta_4 = 0.8416$. Then the outcome y_{ij} was sampled according to:

$$y_{ij} = \begin{cases} 1 & \text{if } -\infty < l_{ij} \leq -0.8416, \\ 2 & \text{if } -0.8416 < l_{ij} \leq -0.2533, \\ 3 & \text{if } -0.2533 < l_{ij} \leq 0.2533, \\ 4 & \text{if } 0.2533 < l_{ij} \leq 0.8416, \\ 5 & \text{if } 0.8416 < l_{ij} < \infty. \end{cases}$$

For each choice of n_i ($n_i = 5, 10$, and 20), we generated 200 data sets. We ran the proposed Gibbs sampler algorithm for 20,000 iterations, after a burn-in period of 2000 iterations.

In Table 1, we present the simulation results of Simulation study 1 for $\beta_1, \beta_2, \beta_3, \delta_1$,

δ_2, δ_3 and δ_4 , including the estimated relative bias and the estimated relative efficiency. In general, it can be seen that the absolute bias obtained by the proposed model (BQOL) when $\theta = 0.5$ is much smaller than its competing models. In most cases, BQOL was better than the other methods in terms of bias and the relative efficiency. The results suggest that our method performs well compare to other approaches. We see that the Bayesian quantile regression approach for ordinal data (BQROR) performs poorly compared to the other methods because it ignores the nature of the longitudinal data. We also see that as n_i increases, the Bayesian logistic ordinal regression (BLOR) for longitudinal data yields low bias and more efficiency. In Table 2, we present the simulation results of Simulation study 1 for Bayesian quantile regression methods when $\theta = 0.25$ and 0.75. Again, in most cases, BQOL was better than BQROR in terms of bias and the relative efficiency.

Table 2: Estimated relative bias and relative efficiency of the Bayesian quantile regression methods for the simulated data 1 when $\theta = 0.25$ and 0.75 .

n_i	Parameter	$\theta = 0.25$				$\theta = 0.75$			
		BQOL		BQROR		BQOL		BQROR	
		bias	eff	bias	eff	bias	eff	bias	eff
5	β_1	0.128	1.000	0.194	1.738	-0.174	1.000	-0.191	1.223
	β_2	-0.057	1.000	-0.058	2.014	-0.070	1.000	0.091	2.031
	β_3	0.016	1.000	0.018	1.374	0.083	1.000	0.066	1.319
	δ_1	0.035	1.000	-0.030	1.238	-0.003	1.000	-0.014	1.011
	δ_2	0.018	1.000	-0.039	1.098	0.005	1.000	-0.059	1.015
	δ_3	0.013	1.000	-0.021	1.043	0.010	1.000	-0.040	1.056
	δ_4	0.020	1.000	-0.025	1.051	0.022	1.000	-0.027	1.037
	10	β_1	-0.132	1.000	0.137	1.035	-0.182	1.000	-0.183
β_2		-0.049	1.000	0.064	1.836	-0.054	1.000	-0.074	1.037
β_3		-0.003	1.000	0.002	1.005	0.011	1.000	0.083	1.215
δ_1		-0.016	1.000	0.037	1.016	-0.008	1.000	0.020	1.117
δ_2		-0.009	1.000	0.040	1.142	-0.008	1.000	0.031	1.027
δ_3		-0.008	1.000	0.033	1.115	-0.007	1.000	0.004	1.006
δ_4		0.019	1.000	0.036	1.003	-0.018	1.000	0.028	1.014
20		β_1	-0.026	1.000	-0.037	1.731	0.013	1.000	0.018
	β_2	-0.156	1.000	0.210	1.079	-0.177	1.000	0.235	1.371
	β_3	0.037	1.000	-0.044	1.117	0.013	1.000	0.018	1.009
	δ_1	-0.014	1.000	0.064	1.008	-0.012	1.000	-0.009	1.004
	δ_2	-0.007	1.000	0.005	1.116	-0.009	1.000	0.016	1.032
	δ_3	0.000	1.000	0.030	1.138	-0.008	1.000	0.016	1.098
	δ_4	0.008	1.000	0.070	1.129	-0.006	1.000	0.010	1.007

4.2 Simulation 2

The setup for this simulation study is the same as Simulation 1, except we sampled the latent variable l_{ij} as follows:

$$l_{ij} = \alpha_i + \beta_1 x_{1ij} + \beta_2 x_{2ij} + \beta_3 x_{3ij} + \varepsilon_{ij}, \quad \alpha_i \sim N(0, 1).$$

This allows us to examine the performance of the proposed model in the case of random effects. In this simulation study, we only consider the performance of the Bayesian methods for longitudinal data with ordinal outcome (BQOL and BLOR). In Table 3, we present the estimates of the parameters $\beta_1, \beta_2, \beta_3, \delta_1, \delta_2, \delta_3$ and δ_4 , when $\theta = 0.5$ and 0.25 . From Table 3 we can see that, our approach tends to give less biased parameter estimates for $\beta_1, \beta_2, \beta_3, \delta_1, \delta_2, \delta_3$ and δ_4 compared to BLOR. The convergence of the proposed Gibbs sampling algorithm in this simulation study was monitored using the multivariate potential scale reduction factor (MPSRF) reported by Brooks and Gelman (1998).

Table 3: The parameter estimations for the simulated data 2 when $\theta = 0.50$ and 0.25 .

n_i	Parameter	BQOL $_{\theta=0.50}$		BQOL $_{\theta=0.25}$		BLOR	
		Mean	SD	Mean	SD	Mean	SD
5	β_1	-5.169	0.738	-5.310	1.003	-5.661	1.069
	β_2	-10.028	0.771	-9.838	1.035	-9.682	1.036
	β_3	15.259	0.688	15.392	0.935	14.273	1.053
	δ_1	-0.862	0.061	-0.842	0.083	-0.871	0.088
	δ_2	-0.249	0.066	-0.237	0.064	-0.231	0.071
	δ_3	0.251	0.063	0.261	0.079	0.261	0.069
	δ_4	0.849	0.073	0.857	0.083	0.881	0.093
	δ_5	0.251	0.063	0.261	0.079	0.261	0.069
10	β_1	-5.132	0.935	-5.116	0.998	-5.842	1.014
	β_2	-10.239	0.722	-10.117	1.028	-9.773	0.998
	β_3	15.024	0.776	15.317	0.993	15.235	1.037
	δ_1	-0.848	0.064	-0.842	0.083	-0.863	0.087
	δ_2	-0.258	0.061	-0.244	0.089	-0.255	0.073
	δ_3	0.258	0.073	0.246	0.076	0.259	0.079
	δ_4	0.848	0.059	0.845	0.037	0.868	0.081
	δ_5	0.258	0.073	0.246	0.076	0.259	0.079
20	β_1	-4.992	0.739	-5.162	0.893	-5.337	0.982
	β_2	-10.039	0.812	-10.337	0.926	-10.241	1.013
	β_3	15.040	0.699	14.893	0.794	14.753	1.045
	δ_1	-0.856	0.066	-0.853	0.081	-0.863	0.079
	δ_2	-0.263	0.085	-0.257	0.063	-0.247	0.066
	δ_3	0.249	0.081	0.249	0.066	0.261	0.071
	δ_4	0.842	0.076	0.853	0.087	0.857	0.081
	δ_5	0.249	0.081	0.249	0.066	0.261	0.071




Figure 1: MPSRF for longitudinal ordinal quantile regression in Simulation 2.

From Figure (1), it can be observed that the MPSRF becomes stable and very close to 1 after about the first 5000 iterations for each quantile level under consideration. Hence, the convergence to the posterior distribution was quick and the mixing was good.

5 Longitudinal Data Example

In this section, we consider a data set from the National Institute of Mental Health Schizophrenia Collaborative (NIMHSC) study previously analysed by (Gibbons and Hedeker, 1994). The objective of this study is to assess treatment-related changes in illness severity over time. Specifically, we studied item 79 (imps79o; severity of illness) of the inpatient multidimensional psychiatric scale. This item was measured on a seven point scale

Table 4: Seven point scale for the NIMHSC study

1	normal
2	borderline mentally ill
3	mildly ill
4	moderately ill
5	markedly ill
6	severely ill
7	among the most extremely ill

as in Table 4:

Hedeker and Gibbons (1994) recorded the seven point scale into four: (1) not ill or borderline, (2) mildly or moderately, (3) markedly ill, (4) severely or most extremely ill. Patients were randomized to receive one of four medications, either placebo or one of three different anti-psychotic drugs. This study consists of three predictors: TxDrug a dummy coded drug effect variable (0=Placebo, 1=Drug), the square root of the week (SqrtWeek), and the interaction between TxDrug and SqrtWeek (TxSWeek). At the θ th ordinal quantile regression, we considered

$$Q_{\text{imps79o}_{ij}}(\theta|\cdot) = \alpha_i + \beta_0 + \beta_1 \text{TxDrug}_{ij} + \beta_2 \text{SqrtWeek}_{ij} + \beta_3 \text{TxSWeek}_{ij}$$

Table 5: The parameter estimations for the NIMHSC study when $\theta = 0.50$ and 0.25 .

Parameter	BQOL $_{\theta=0.50}$		BQOL $_{\theta=0.25}$		BLOR	
	Mean	%95 CI	Mean	%95 CI	Mean	%95 CI
β_0	5.663	(5.037, 6.1192)	3.527	(2.442, 4.735)	6.021	(5.571, 7.421)
β_1	-0.073	(-0.523, 0.419)	-0.048	(-0.661, 0.783)	-0.227	(-0.991, 0.771)
β_2	-0.746	(-1.351, -0.1337)	-0.643	(-0.897, -0.437)	-1.247	(-1.562, -0.661)
β_3	-1.206	(-1.663, -0.881)	-1.104	(-1.437, -0.789)	-0.883	(-1.641, -0.291)
δ_1	2.751	(2.538, 2.879)	2.865	(2.479, 3.125)	1.997	(1.087, 3.114)
δ_2	4.173	(3.894, 4.337)	3.984	(3.716, 4.118)	3.764	(3.221, 4.261)
δ_3	5.889	(5.103, 6.709)	5.765	(5.042, 6.327)	5.443	(4.793, 6.431)

Table 5 lists parameter estimations obtained using the Bayesian methods (BQOL and BLOR). The methods are assessed based on 95% credible intervals and the deviance information criterion (DIC; Spiegelhalter et al., 2002). Clearly, it can be seen that the credible intervals (95% CI) obtained using the BQOL when $\theta = 0.50$ are generally shorter than the credible intervals obtained using the BLOR, suggesting an efficiency gain and stable estimation from the posterior distributions. In addition, DIC was computed for our model when $\theta = 0.50$ and $\theta = 0.25$ as well as for BLOR and the numbers were 3311.32, 3615.48 and 3417.39, respectively. Hence, under $\theta = 0.50$, model comparison using DIC indicates that quantile ordinal models can give a better model fit compared to the Bayesian logistic ordinal regression (BLOR) for longitudinal data. This shows that the model used for the errors in (3) is a working model with artificial assumptions, employed on the outcome variable to achieve the equivalence between maximising SLD and the minimising problem in (2).

6 Conclusion

Since Bayesian quantile methods for estimating ordinal models with longitudinal data have not been proposed, yet. This paper fills this gap and presents a random effects ordinal quantile regression model for analysis of longitudinal data with ordinal outcome of interest. An efficient Gibbs sampling algorithm was derived for fitting the model to the data based on a location-scale mixture representation of the skewed double exponential distribution. The proposed approach is illustrated using simulated data and a real data example. Results show that the proposed approach performs well. One of the most desirable features of the proposed method is its model robustness in the sense that makes very minimal assumptions on the form of the error term distribution and thus is able to accommodate non-normal errors and outliers, which are popular in many real world applications.

Appendix A: Gibbs Sampler Details

The full conditional distribution of each v_{ij} , denoted by $P(v_{ij}|l_{ij}, \boldsymbol{\beta}, \alpha_i)$ is proportional to $P(l_{ij}|v_{ij}, \boldsymbol{\beta}, \alpha_i)P(v_{ij})$. Thus, we have

$$\begin{aligned}
P(v_{ij}|l_{ij}, \boldsymbol{\beta}, \alpha_i) &\propto v_{ij}^{-1/2} \exp \left\{ -\frac{1}{2} \left(\frac{l_{ij} - \mathbf{x}'_{ij} \boldsymbol{\beta} - \alpha_i - \xi v_{ij}}{\sqrt{2v_{ij}}} \right)^2 - \zeta v_{ij} \right\} \\
&\propto v_{ij}^{-1/2} \exp \left\{ -\frac{1}{2} \left(\frac{(l_{ij} - \mathbf{x}'_{ij} \boldsymbol{\beta} - \alpha_i)^2 + \xi^2 v_{ij}^2 - 2\xi v_{ij} (l_{ij} - \mathbf{x}'_{ij} \boldsymbol{\beta} - \alpha_i)}{2v_{ij}} + 2\zeta v_{ij} \right) \right\} \\
&\propto v_{ij}^{-1/2} \exp \left\{ -\frac{1}{2} \left[\frac{(l_{ij} - \mathbf{x}'_{ij} \boldsymbol{\beta} - \alpha_i)^2}{2} v_{ij}^{-1} + \left(\frac{\xi^2}{2} + 2\zeta \right) v_{ij} \right] \right\} \\
&\propto v_{ij}^{-1/2} \exp \left\{ -\frac{1}{2} \left[\frac{(l_{ij} - \mathbf{x}'_{ij} \boldsymbol{\beta} - \alpha_i)^2}{2} v_{ij}^{-1} + \left(\frac{1}{2} \right) v_{ij} \right] \right\} \\
&\propto v_{ij}^{-1/2} \exp \left\{ -\frac{1}{2} \left[\varrho_1^2 v_{ij}^{-1} + \varrho_2^2 v_{ij} \right] \right\},
\end{aligned}$$

where $\xi = 1 - 2\theta$ and $\zeta = \theta(1 - \theta)$. Thus, the full conditional distribution of each v_{ij} is a generalized inverse Gaussian distribution GIG $(\nu, \varrho_1, \varrho_2)$, where $\varrho_1^2 = (l_{ij} - \mathbf{x}'_{ij} \boldsymbol{\beta} - \alpha_i)^2/2$ and $\varrho_2^2 = 1/2$. Recall that if $x \sim \text{GIG}(0.5, \varrho_1, \varrho_2)$ then the pdf of x is given by (Barndorff-Nielsen and Shephard, 2001)

$$f(x|\nu, \varrho_1, \varrho_2) = \frac{(\varrho_2/\varrho_1)^\nu}{2K_\nu(\varrho_1\varrho_2)} x^{\nu-1} \exp \left\{ -\frac{1}{2}(x^{-1}\varrho_1^2 + x\varrho_2^2) \right\},$$

where $x > 0$, $-\infty < \nu < \infty$, $\varrho_1, \varrho_2 \geq 0$ and $K_\nu(\cdot)$ is so called “modified Bessel function of the third kind”.

The full conditional distribution of each β_k , denoted by $P(\beta_k|\mathbf{l}, \boldsymbol{\beta}_{-k}, \boldsymbol{\alpha}, \mathbf{v}, s_k)$ is proportional to $P(\mathbf{l}|\boldsymbol{\beta}, \boldsymbol{\alpha}, \mathbf{v}, s_k)P(\beta_k|s_k)$, where $\boldsymbol{\beta}_{-k}$ is the vector $\boldsymbol{\beta}$ excluding the element β_k . Thus, we have

$$\begin{aligned}
P(\beta_k|\mathbf{l}, \boldsymbol{\beta}_{-k}, \boldsymbol{\alpha}, \mathbf{v}, s_k) &\propto P(\mathbf{l}|\boldsymbol{\beta}, \boldsymbol{\alpha}, \mathbf{v}, s_k)P(\beta_k|s_k) \\
&\propto \exp \left\{ -\frac{1}{2} \sum_{i=1}^N \sum_{j=1}^{n_i} \frac{(l_{ij} - \mathbf{x}'_{ij} \boldsymbol{\beta} - \alpha_i - \xi v_{ij})^2}{2v_{ij}} \right\} \exp \left\{ -\frac{\beta_k^2}{2s_k} \right\} \\
&\propto \exp \left\{ -\frac{1}{2} \left[\left(\sum_{i=1}^N \sum_{j=1}^{n_i} \frac{x_{ijk}^2}{2v_{ij}} + \frac{1}{s_k} \right) \beta_k^2 - 2 \sum_{i=1}^N \sum_{j=1}^{n_i} \frac{\tilde{l}_{ijk} x_{ijk}}{2v_{ij}} \beta_k \right] \right\},
\end{aligned}$$

where $\mathbf{x}_{ij} = (x_{ij1}, \dots, x_{ijp})$ and $\tilde{l}_{ijk} = l_{ijk} - \sum_{h=1, h \neq k}^p x_{ijh} \beta_h - \alpha_i - \xi v_{ij}$. Then the full conditional distribution for β_k is normal with mean μ_{β_k} and variance $\sigma_{\beta_k}^2$, where

$$\sigma_{\beta_k}^{-2} = \sum_{i=1}^N \sum_{j=1}^{n_i} \frac{x_{ijk}^2}{2v_{ij}} + \frac{1}{s_k},$$

and

$$\mu_{\beta_k} = \sigma_k^2 \sum_{i=1}^N \sum_{j=1}^{n_i} \frac{\tilde{l}_{ijk} x_{ijk}}{2v_{ij}}.$$

The full conditional distribution of each s_k , denoted by $P(s_k|\beta_k)$ is

$$\begin{aligned} P(s_k|\beta_k) &\propto P(\beta_k|s_k)P(s_k) \\ &\propto \frac{1}{\sqrt{2\pi s_k}} \exp\left\{-\frac{\beta_k^2}{2s_k}\right\} \exp\left\{-\frac{\lambda^2}{2}s_k\right\} \\ &\propto s_k^{-\frac{1}{2}} \exp\left\{-\frac{1}{2}\left(\beta_k^2 s_k^{-1} + \lambda^2 s_k\right)\right\} \end{aligned}$$

Thus, the full conditional distribution of s_k is a GIG(0.5, ϱ_1 , ϱ_2), where $\varrho_1^2 = \beta_k^2$ and $\varrho_2^2 = \lambda^2$.

The full conditional distribution of λ^2 , denoted by $P(\lambda^2|\mathbf{s})$ is

$$P(\lambda^2|\mathbf{s}) = P(\mathbf{s}|\lambda^2)P(\lambda^2) \quad (9)$$

$$\propto \prod_{k=1}^p \frac{\lambda^2}{2} \exp\left\{-\frac{\lambda^2}{2}s_k\right\} (\lambda^2)^{a_1-1} \exp\{-a_2\lambda^2\} \quad (10)$$

$$\propto (\lambda^2)^{p+a_1-1} \exp\left\{-\lambda^2\left(\sum_{k=1}^p s_k/2 + a_2\right)\right\} \quad (11)$$

That is, the full conditional distribution of λ^2 is a Gamma distribution.

The full conditional distribution of each α_i , denoted by $P(\alpha_i|\mathbf{l}, \boldsymbol{\beta}, \mathbf{v}, \phi)$ is proportional to $P(\mathbf{l}|\alpha_i, \boldsymbol{\beta}, \mathbf{v})P(\alpha_i|\phi)$. Thus, we have

$$\begin{aligned} P(\alpha_i|\mathbf{l}, \boldsymbol{\beta}, \mathbf{v}, \phi) &\propto P(\mathbf{l}|\alpha_i, \boldsymbol{\beta}, \mathbf{v})P(\alpha_i|\phi) \\ &\propto \exp\left\{-\frac{1}{2}\sum_{j=1}^{n_i} \frac{(l_{ij} - \mathbf{x}'_{ij}\boldsymbol{\beta} - \alpha_i - \xi v_{ij})^2}{2v_{ij}}\right\} \exp\left\{-\frac{\alpha_i^2}{2\phi}\right\} \\ &\propto \exp\left\{-\frac{1}{2}\left[\left(\sum_{j=1}^{n_i} \frac{1}{2v_{ij}} + \frac{1}{\phi}\right)\alpha_i^2 - 2\sum_{j=1}^{n_i} \frac{\eta_{ij}}{2v_{ij}}\alpha_i\right]\right\}, \end{aligned}$$

where $\eta_{ij} = l_{ij} - \mathbf{x}'_{ij}\boldsymbol{\beta} - \xi v_{ij}$. Then the full conditional distribution for α_i is normal with mean μ_{α_i} and variance $\sigma_{\alpha_i}^2$, where

$$\sigma_{\alpha_i}^{-2} = \sum_{j=1}^{n_i} \frac{1}{2v_{ij}} + \frac{1}{\phi},$$

and

$$\mu_{\alpha_i} = \sigma_{\alpha_i}^2 \sum_{j=1}^{n_i} \frac{\eta_{ij}}{2v_{ij}}.$$

The full conditional distribution of ϕ , denoted by $P(\phi|\boldsymbol{\alpha})$, is proportional to $P(\boldsymbol{\alpha}|\phi)P(\phi)$. Thus, we have

$$\begin{aligned} P(\phi|\boldsymbol{\alpha}) &\propto P(\boldsymbol{\alpha}|\phi)P(\phi), \\ &\propto \left(\prod_{i=1}^N \frac{1}{\sqrt{2\pi\phi}} \right) \exp \left\{ -\frac{\sum_{i=1}^N \alpha_i^2}{2\phi} \right\} \phi^{-b_1-1} \exp \left\{ -\frac{b_2}{\phi} \right\} \\ &\propto \phi^{-\frac{N}{2}-b_1-1} \exp \left\{ -\frac{1}{\phi} \left(\frac{\sum_{i=1}^N \alpha_i^2}{2} + b_2 \right) \right\} \end{aligned}$$

That is, the full conditional distribution of ϕ is a inverse Gamma distribution.

The full conditional distribution of each l_{ij} , denoted by $P(l_{ij}|\boldsymbol{\beta}, \boldsymbol{\delta}, \alpha_i, v_{ij})$ is proportional to $P(y_{ij}|l_{ij}, \boldsymbol{\delta}) P(l_{ij}|\boldsymbol{\beta}, \alpha_i, v_{ij})$. Thus, we have

$$\begin{aligned} P(l_{ij}|\boldsymbol{\beta}, \boldsymbol{\delta}, \alpha_i, v_{ij}) &\propto P(y_{ij}|l_{ij}, \boldsymbol{\delta})P(l_{ij}|\boldsymbol{\beta}, \alpha_i, v_{ij}) \\ &\propto \mathbf{1}\{\delta_{c-1} < l_{ij} \leq \delta_c\} \mathbf{N}(l_{ij}; \mathbf{x}'_{ij}\boldsymbol{\beta} + \alpha_i + \xi v_{ij}, 2v_{ij}) \end{aligned}$$

That is, the full conditional distribution of l_{ij} is a truncated normal distribution.

At last, the full conditional posterior distribution of δ_c , denoted by $P(\delta_c|\mathbf{y}, \mathbf{l})$ is proportional to $p(\mathbf{y}|\mathbf{l}, \boldsymbol{\delta})P(\boldsymbol{\delta})$. Thus, we have

$$P(\delta_c|\mathbf{y}, \mathbf{l}) \propto p(\mathbf{y}|\mathbf{l}, \boldsymbol{\delta})P(\boldsymbol{\delta}) \quad (12)$$

$$\propto \prod_{i=1}^N \prod_{j=1}^{n_i} \sum_{c=1}^C \mathbf{1}(y_{ij} = c) \mathbf{1}(\delta_{c-1} < l_{ij} < \delta_c) \mathbf{1}(\boldsymbol{\delta} \in T) \quad (13)$$

Following [Montesinos-López et al. \(2015\)](#) and [Sorensen et al. \(1995\)](#), the full conditional distribution of δ_c is

$$P(\delta_c|\mathbf{y}, \mathbf{l}) = \frac{1}{\min(l_{ij}|y_{ij} = c+1) - \max(l_{ij}|y_{ij} = c)} \mathbf{1}(\boldsymbol{\delta} \in T)$$

That is, the full conditional distribution of δ_c is a uniform distribution.

References

- Albert, J. H. and Chib, S. (1993). “Bayesian analysis of binary and polychotomous response data.” *Journal of the American statistical Association*, 88(422): 669–679. [6](#)
- Alhamzawi, R. and Yu, K. (2014). “Bayesian Lasso-mixed quantile regression.” *Journal of Statistical Computation and Simulation*, 84(4): 868–880. [2](#), [4](#)
- Andrews, D. F. and Mallows, C. L. (1974). “Scale mixtures of normal distributions.” *Journal of the Royal Statistical Society. Series B (Methodological)*, 99–102. [5](#)
- Baayen, R. H., Davidson, D. J., and Bates, D. M. (2008). “Mixed-effects modeling with crossed random effects for subjects and items.” *Journal of memory and language*, 59(4): 390–412. [2](#)

- Bae, K. and Mallick, B. K. (2004). “Gene selection using a two-level hierarchical Bayesian model.” *Bioinformatics*, 20(18): 3423–3430. [5](#)
- Barndorff-Nielsen, O. E. and Shephard, N. (2001). “Non-Gaussian Ornstein–Uhlenbeck-based models and some of their uses in financial economics.” *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 63(2): 167–241. [15](#)
- Bottai, M., Frongillo, E. A., Sui, X., O’Neill, J. R., McKeown, R. E., Burns, T. L., Liese, A. D., Blair, S. N., and Pate, R. R. (2014). “Use of quantile regression to investigate the longitudinal association between physical activity and body mass index.” *Obesity*, 22(5): E149–E156. [1](#)
- Brooks, S. P. and Gelman, A. (1998). “General methods for monitoring convergence of iterative simulations.” *Journal of computational and graphical statistics*, 7(4): 434–455. [10](#)
- Cade, B. S. and Noon, B. R. (2003). “A gentle introduction to quantile regression for ecologists.” *Frontiers in Ecology and the Environment*, 1(8): 412–420. [2](#)
- Cade, B. S., Terrell, J. W., and Porath, M. T. (2008). “Estimating fish body condition with quantile regression.” *North American Journal of Fisheries Management*, 28(2): 349–359. [3](#)
- Fitzmaurice, G. M., Laird, N. M., and Ware, J. H. (2012). *Applied longitudinal analysis*, volume 998. John Wiley & Sons. [2](#)
- Geraci, M. and Bottai, M. (2007). “Quantile regression for longitudinal data using the asymmetric Laplace distribution.” *Biostatistics*, 8(1): 140–154. [2](#)
- Gibbons, R. D. and Hedeker, D. (1994). “Application of random-effects probit regression models.” *Journal of consulting and clinical psychology*, 62(2): 285. [12](#)
- Griffin, J. E., Brown, P. J., et al. (2010). “Inference with normal-gamma prior distributions in regression problems.” *Bayesian Analysis*, 5(1): 171–188. [5](#)
- Hans, C. (2009). “Bayesian lasso regression.” *Biometrika*, 96(4): 835–845. [5](#)
- Hedeker, D. and Gibbons, R. D. (1994). “A random-effects ordinal regression model for multilevel analysis.” *Biometrics*, 933–944. [13](#)
- (2006). *Longitudinal data analysis*, volume 451. John Wiley & Sons. [2](#)
- Hedges, L. V. and Vevea, J. L. (1998). “Fixed and random-effects models in meta-analysis.” *Psychological methods*, 3(4): 486. [2](#)
- Hendricks, W. and Koenker, R. (1992). “Hierarchical spline models for conditional quantiles and the demand for electricity.” *Journal of the American statistical Association*, 87(417): 58–68. [2](#)
- Kinney, S. K. and Dunson, D. B. (2007). “Fixed and random effects selection in linear and logistic models.” *Biometrics*, 63(3): 690–698. [5](#)
- Koenker, R. (2004). “Quantile regression for longitudinal data.” *Journal of Multivariate*

- Analysis*, 91(1): 74–89. [2](#)
- (2005). *Quantile regression*. 38. Cambridge university press. [2](#)
- Koenker, R. and Bassett, G. (1978). “Regression quantiles.” *Econometrica: Journal of the Econometric Society*, 33–50. [1](#), [3](#)
- Koenker, R. and Geling, O. (2001). “Reappraising medfly longevity: a quantile regression survival analysis.” *Journal of the American Statistical Association*, 96(454): 458–468. [2](#)
- Kostov, P. and Davidova, S. (2013). “A quantile regression analysis of the effect of farmers attitudes and perceptions on market participation.” *Journal of Agricultural Economics*, 64(1): 112–132. [1](#)
- Kozumi, H. and Kobayashi, G. (2011). “Gibbs sampling methods for Bayesian quantile regression.” *Journal of statistical computation and simulation*, 81(11): 1565–1578. [4](#)
- Laird, N. M. and Ware, J. H. (1982). “Random-effects models for longitudinal data.” *Biometrics*, 963–974. [2](#)
- Li, Q., Xi, R., and Lin, N. (2010). “Bayesian regularized quantile regression.” *Bayesian Analysis*, 5(3): 533–556. [5](#)
- Lipsitz, S. R., Fitzmaurice, G. M., Molenberghs, G., and Zhao, L. P. (1997). “Quantile Regression Methods for Longitudinal Data with Drop-outs: Application to CD4 Cell Counts of Patients Infected with the Human Immunodeficiency Virus.” *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 46(4): 463–476. [2](#)
- Luo, Y., Lian, H., and Tian, M. (2012). “Bayesian quantile regression for longitudinal data models.” *Journal of Statistical Computation and Simulation*, 82(11): 1635–1649. [4](#)
- McCulloch, C. E. and Neuhaus, J. M. (2001). *Generalized linear mixed models*. Wiley Online Library. [2](#)
- Mezali, H. and Beasley, J. E. (2013). “Quantile regression for index tracking and enhanced indexation.” *Journal of the Operational Research Society*, 64(11): 1676–1692. [2](#)
- Montesinos-López, O. A., Montesinos-López, A., Crossa, J., Burgueño, J., and Eskridge, K. (2015). “Genomic-enabled prediction of ordinal data with Bayesian logistic ordinal regression.” *G3: Genes— Genomes— Genetics*, 5(10): 2113–2126. [5](#), [7](#), [17](#)
- Park, T. and Casella, G. (2008). “The Bayesian lasso.” *Journal of the American Statistical Association*, 103(482): 681–686. [5](#)
- Rahman, M. A. (2016). “Bayesian Quantile Regression for Ordinal Models.” *Bayesian Analysis*, 11(1): 1–24. [7](#)
- Reich, B. J. (2012). “Spatiotemporal quantile regression for detecting distributional changes in environmental processes.” *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 61(4): 535–553. [2](#)

- Reich, B. J., Bondell, H. D., and Wang, H. J. (2010). “Flexible Bayesian quantile regression for independent and clustered data.” *Biostatistics*, 11(2): 337–352. [2](#)
- Sorensen, D., Andersen, S., Gianola, D., and Korsgaard, I. (1995). “Bayesian inference in threshold models using Gibbs sampling.” *Genetics Selection Evolution*, 27(3): 229–249. [5](#), [17](#)
- Spiegelhalter, D. J., Best, N. G., Carlin, B. P., and Van Der Linde, A. (2002). “Bayesian measures of model complexity and fit.” *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 64(4): 583–639. [14](#)
- Tao, H., Palta, M., Yandell, B. S., and Newton, M. A. (1999). “An estimation method for the semiparametric mixed effects model.” *Biometrics*, 55(1): 102–110. [2](#)
- Tibshirani, R. (1996). “Regression shrinkage and selection via the lasso.” *Journal of the Royal Statistical Society. Series B (Methodological)*, 267–288. [5](#)
- Verbeke, G. and Lesaffre, E. (1996). “A linear mixed-effects model with heterogeneity in the random-effects population.” *Journal of the American Statistical Association*, 91(433): 217–221. [2](#)
- Wang, H. and He, X. (2007). “Detecting differential expressions in Gene Chip microarray studies: a quantile approach.” *Journal of the American Statistical Association*, 102(477): 104–112. [2](#)
- Wasserman, L. (2006). *All of nonparametric statistics*. Springer Science & Business Media. [3](#)
- Wolfinger, R. and O’connell, M. (1993). “Generalized linear mixed models a pseudo-likelihood approach.” *Journal of statistical Computation and Simulation*, 48(3-4): 233–243. [2](#)
- Yu, K., Lu, Z., and Stander, J. (2003). “Quantile regression: applications and current research areas.” *Journal of the Royal Statistical Society: Series D (The Statistician)*, 52(3): 331–350. [2](#), [3](#)
- Yu, K. and Moyeed, R. A. (2001). “Bayesian quantile regression.” *Statistics & Probability Letters*, 54(4): 437–447. [4](#)
- Yuan, Y. and Yin, G. (2010). “Bayesian quantile regression for longitudinal studies with nonignorable missing data.” *Biometrics*, 66(1): 105–114. [2](#)
- Yue, Y. R. and Rue, H. (2011). “Bayesian inference for additive mixed quantile regression models.” *Computational Statistics & Data Analysis*, 55(1): 84–96. [4](#)