

# Randomly Weighted Averages: A Multivariate Case

Hazhir Homei

*Department of Statistics, Faculty of Mathematical Sciences,  
University of Tabriz, P.O.Box 51666-17766, Tabriz, IRAN.*

homei@tabrizu.ac.ir

---

## Abstract

Stochastic linear combinations of some random vectors are studied where the distribution of the random vectors and the joint distribution of their coefficients are Dirichlet. A method is provided for calculating the distribution of these combinations which has been studied before by some authors. Our main result is a generalization of some existing results with a simpler proof.

*Keywords:* Multivariate randomly weighted average; Multivariate Stieltjes transform; Dirichlet Distributions; Multivariate stable distributions; Randomly Linear Transformations; Dependent Components; Lifetime.

---

## 1. Introduction

For given random variables  $X_1, \dots, X_n$  the distribution of the stochastic linear combination  $Z = \sum_{i=1}^n W_i X_i$  is used for the problems in lifetime, stochastic matrices, neural networks and other applications in sociology and biology. Let  $X_i$  ( $1 \leq i \leq n$ ) be the lifetime measured in a lab and  $0 \leq W_i \leq 1$  be the random effect of the environment on it; so  $W_i X_i \leq X_i$  and thus  $\sum_{i=1}^n W_i X_i$  is the average lifetime in the environment (see Homei (2015)). Recently, several authors have focused on computing the lifetime of systems in the real conditions. Indeed, the randomly linear combination of random vectors have

many applications including traditional portfolio selection models, relationship between attitudes and behavior, number of cancer cells in tumor biology, stream flow in hydrology (Nadarajah & Kotz (2005)), branching processes, infinite particle systems and probabilistic algorithms, and vehicle speed and lifetime (cf. Homei (2015), Rezapour & Alamatsaz (2014) and the references therein) so finding their distributions has attracted the attentions of numerous researchers.

In this paper, considering the dependent components and the real environmental conditions, the distribution of lifetimes (in case of the Dirichlet distributions) is calculated showing that the main result obtained here covers several previous results and provides a simpler proof (cf. e.g. Johnson & Kotz (1990a), Sethuraman (1994), van Assche (1987), Volodin & Kotz & Johnson (1993)).

The inner product of two random vectors was introduced in Homei (2014) and the exact distribution of this product was investigated for some random vectors with Beta and Dirichlet distributions. In this paper a new generalization for the inner product of two random vectors is introduced. For a random vector  $\mathbf{W}' = \langle W_1, \dots, W_n \rangle$  and a vector  $\underline{\mathbf{X}} = \langle \mathbf{X}_1, \dots, \mathbf{X}_n \rangle$  of random vectors (each  $\mathbf{X}_i$  being  $k$ -dimensional) the inner product of  $\mathbf{X}$  and  $\mathbf{W}$  is essentially the linear transformation of  $\mathbf{W}$  under the  $k \times n$  matrix  $\underline{\mathbf{X}}$  which is  $\mathbf{Z} = \sum_{i=1}^n W_i \mathbf{X}_i$ . We assume that  $\mathbf{W}$  is independent from  $\mathbf{X}_1, \dots, \mathbf{X}_n$  and has Dirichlet distribution; also  $\mathbf{X}_i$ 's have Dirichlet distributions. Identifying the distribution of  $\mathbf{Z}$  usually requires

- (i) either some long computations with combinatorial identities (see e.g. Volodin & Kotz & Johnson (1993), Johnson & Kotz (1990), Johnson

& Kotz (1990a), Sethuraman (1994), Homei (2014));

- (ii) or advanced techniques requiring performing certain transformations and solving differential equations (see e.g. Homei (2012), Soltani & Homei (2009), van Assche (1987) or Homei (2015) and the references therein).

In this paper a new way is introduced to identify the distribution of randomly linear combinations (of Dirichlet distributions) by which a class of stochastic differential equations can be solved; this new method is much simpler than the existing ones.

## 2. The Main Result

Our main result identifies the distribution of the randomly linear combination when the coefficients come from a Dirichlet distribution.

**Theorem 2.1.** *If  $\mathbf{X}_1, \dots, \mathbf{X}_n$  are independent  $k$ -variate random vectors with respectively  $Dirichlet(\alpha^{(1)}), \dots, Dirichlet(\alpha^{(n)})$  distributions, for some  $k$ -dimensional vectors  $\alpha^{(j)} = \langle \alpha_1^{(j)}, \dots, \alpha_k^{(j)} \rangle$  ( $j = 1, \dots, n$ ), and the random vector  $\mathbf{W} = \langle W_1, \dots, W_n \rangle$  is independent from  $\mathbf{X}_1, \dots, \mathbf{X}_n$  and has the distribution*

$$Dirichlet \left( \sum_{i=1}^k \alpha_i^{(1)}, \dots, \sum_{i=1}^k \alpha_i^{(n)} \right),$$

*then the randomly linear combination  $\mathbf{Z} = \sum_{i=1}^n W_i \mathbf{X}_i$  has the distribution*

$$Dirichlet \left( \sum_{j=1}^n \alpha_1^{(j)}, \dots, \sum_{j=1}^n \alpha_k^{(j)} \right).$$

(The First) Proof. Let  $Y_j$  ( $j = 1, \dots, n$ ) be independent random variables independent from  $(\mathbf{X}_1, \dots, \mathbf{X}_n)$  that have the distribution  $\Gamma(\sum_{i=1}^k \alpha_i^{(j)}, \frac{1}{\mu})$ , respectively. Thus,

$$W_j \stackrel{d}{=} \frac{Y_j}{(\sum_{l=1}^n Y_l)} \text{ for } j = 1, \dots, n.$$

The  $k$ -dimensional vectors  $\mathbf{T}_j = Y_j \mathbf{X}_j$  have independent components and

$$\langle \Gamma(\alpha_1^{(j)}, \frac{1}{\mu}), \dots, \Gamma(\alpha_k^{(j)}, \frac{1}{\mu}) \rangle$$

distribution. Thus, we have

$$\sum_{j=1}^n \left( \frac{\mathbf{T}_j}{\sum_{l=1}^n Y_l} \right) = \sum_{j=1}^n \frac{Y_j}{\sum_{l=1}^n Y_l} \mathbf{X}_j \stackrel{d}{=} \sum_{j=1}^n W_j \mathbf{X}_j.$$

Now,  $\sum_{j=1}^n \left( \frac{\mathbf{T}_j}{\sum_{l=1}^n Y_l} \right)$  has Dirichlet( $\sum_{j=1}^n \alpha_1^{(j)}, \dots, \sum_{j=1}^n \alpha_k^{(j)}$ ) distribution, so  $\sum_{j=1}^n W_j \mathbf{X}_j = \mathbf{Z}$  has the same distribution.  $\boxtimes$

**Corollary 2.2.** *Let the  $k$ -variate random variables  $\mathbf{X}_1, \dots, \mathbf{X}_n$  be independent and with common distribution. If  $\mathbf{W} = \langle W_1, \dots, W_n \rangle$  is independent from  $\mathbf{X}_1, \dots, \mathbf{X}_n$  having Dirichlet( $k\alpha, \dots, k\alpha$ ) distribution, then the randomly linear combination  $\mathbf{Z} = \sum_{i=1}^n W_i \mathbf{X}_i$  has the Dirichlet( $n\alpha, \dots, n\alpha$ ) distribution if and only if  $\mathbf{X}_i$ 's ( $i = 1, \dots, n$ ) have Dirichlet( $\alpha, \dots, \alpha$ ) distributions.*

*Proof.* The “if” part follows from Theorem 2.1 and the “only if” part follows from the theorem of Volodin & Kotz & Johnson (1993).  $\boxtimes$

Some similar results (to Corollary 2.2) can be found in e.g. Alamtasaz (1993). The following results are obtained as special cases of Theorem 2.1 and Corollary 2.2:

- Lemma 1 of Sethuraman (1994) (for  $n=2$ ), and
- Theorem of Volodin & Kotz & Johnson (1993) (when  $n = k$  and  $\alpha_i^{(j)} = a$  for  $j=1, \dots, k, i=1, \dots, k$ ),

which discusses the multivariate case, or the references below which discuss the single-variable case (note that for  $k=2$  the Dirichlet distribution leads to the Beta distribution):

- Theorem of van Assche (1987) (for  $n=k=2, \alpha_1^{(1)} = \alpha_2^{(1)} = \alpha_1^{(2)} = \alpha_2^{(2)} = \frac{1}{2}$ ),
  - Theorem 2 of Johnson & Kotz (1990a) (for  $n=k=2, \alpha_1^{(1)} = \alpha_2^{(1)} = \alpha_1^{(2)} = \alpha_2^{(2)} = a$ ),
  - Theorem 2.4 of Homei (2014) (for  $k=2$  and  $\alpha_1^{(j)} = \alpha, \alpha_2^{(j)} = 1 - \alpha$  for  $j = 1, \dots, n$ ),
  - Theorem 1 of Homei (2013) (for  $k=2$ );
- and others; see the references in Homei (2015).

### 3. Four Other Proofs for Theorem 2.1 and a Variant of It

#### 3.1. Moment Generation Method

*The Second Proof.* The generating moment function of  $\mathbf{T}_j$ 's in the first proof are

$$E(e^{\mathbf{t}'\mathbf{T}_j}) = EE(e^{\mathbf{t}'Y_j\mathbf{X}_j} | X_j) = E\left(\frac{1}{1 - \mathbf{t}'\mathbf{X}_j}\right)^{\sum_{i=1}^k \alpha_i^{(j)}}.$$

By [9, page 77] we have

$$M_{\mathbf{T}_j}(\mathbf{t}) = \left(\frac{1}{1 - t_j}\right)^{\sum_{i=1}^k \alpha_i^{(j)}}.$$

So, the components of the vector  $\mathbf{T}_j$  are independent and have gamma distributions, which proves the theorem.  $\boxtimes$

### 3.2. Applying Basu's Theorem

*The Third Proof.* We can write (for  $j = 1, \dots, n$ )

$$\mathbf{X}_j \sim \left( \frac{\Gamma_{1j}(\alpha_1^{(j)})}{\sum_{i=1}^k \Gamma_{ij}(\alpha_i^{(j)})}, \dots, \frac{\Gamma_{kj}(\alpha_k^{(j)})}{\sum_{i=1}^k \Gamma_{ij}(\alpha_i^{(j)})} \right),$$

and

$$\mathbf{W} \sim \left( \frac{\sum_{i=1}^k \Gamma_{ij}(\alpha_i^{(1)})}{\sum_{j=1}^n \sum_{i=1}^k \Gamma_{ij}(\alpha_i^{(j)})}, \dots, \frac{\sum_{i=1}^k \Gamma_{ij}(\alpha_i^{(n)})}{\sum_{j=1}^n \sum_{i=1}^k \Gamma_{ij}(\alpha_i^{(j)})} \right).$$

So,

$$\mathbf{Z} \stackrel{d}{=} \left( \frac{\sum_{j=1}^k \Gamma_{1j}(\alpha_1^{(j)})}{\sum_{j=1}^n \sum_{i=1}^k \Gamma_{ij}(\alpha_i^{(j)})}, \dots, \frac{\sum_{j=1}^k \Gamma_{kj}(\alpha_k^{(j)})}{\sum_{j=1}^n \sum_{i=1}^k \Gamma_{ij}(\alpha_i^{(j)})} \right).$$

Let us recall that  $\Gamma_{ij}(\alpha_i^{(j)})$ 's (for  $i = 1, \dots, k$  and  $j = 1, \dots, n$ ) are independent random variables with gamma distributions, which implies the independence of the components of  $\mathbf{W}$  from  $\mathbf{X}_j$ 's (Basu's Theorem).  $\boxtimes$

### 3.3. Mathematical Induction

*The Fourth Proof.* For  $n = 2$  the theorem follows from [13, Lemma 1]. Suppose the theorem holds for  $n$  (the induction hypothesis). We prove it for  $n + 1$  (the induction conclusion): By dividing the both sides of

$$\sum_{i=1}^{n+1} Y_i \mathbf{X}_i = \left( \sum_{i=1}^n Y_i \right) \left( \sum_{j=1}^n \frac{Y_j}{\sum_{l=1}^n Y_l} \mathbf{X}_j \right) + Y_{n+1} \mathbf{X}_{n+1}$$

by  $\sum_{i=1}^{n+1} Y_i$  and using the induction hypothesis we have

$$\sum_{i=1}^{n+1} \frac{Y_i}{\sum_{i=1}^{n+1} Y_i} \mathbf{X}_i = \left( \sum_{i=1}^n \frac{Y_i}{\sum_{i=1}^{n+1} Y_i} \right) \left( \sum_{j=1}^n \frac{Y_j}{\sum_{l=1}^n Y_l} \mathbf{X}_j \right) + \frac{Y_{n+1}}{\sum_{i=1}^{n+1} Y_i} \mathbf{X}_{n+1}$$

in which the right hand side holds by [13, Lemma 1] (for  $n = 2$ ).  $\boxtimes$

### 3.4. The Moments Method

*The Fifth Proof.* The general moments  $(s_1, s_2, \dots, s_k)$  of  $\mathbf{Z}$  are as follows:

$$\mathbb{E}\left(\prod_{j=1}^k \left(\sum_{i=1}^n W_j X_{ij}\right)^{s_j}\right) = \mathbb{E}\left(\prod_{j=1}^k \left(\sum_{h_j} \binom{s_j}{h_{1j}, h_{2j}, \dots, h_{nj}}\right) \prod_{i=1}^n (W_j X_{ij})^{h_{ij}}\right)$$

where  $\sum_{h_j}$  denotes summation over all non-negative integers

$$h_j = (h_{1j}, h_{2j}, \dots, h_{nj}) \text{ subject to } \sum_{i=1}^n h_{ij} = s_j \quad (j = 1, 2, \dots, n).$$

This equation can be rearranged as

$$\begin{aligned} &= \mathbb{E}\left(\sum_{h_1} \cdots \sum_{h_k} \left(\prod_{j=1}^k \binom{s_j}{h_{1j}, h_{2j}, \dots, h_{nj}}\right) \prod_{j=1}^k \prod_{i=1}^n (W_i X_{ij})^{h_{ij}}\right) \\ &= \mathbb{E}\left(\sum_{h_1} \cdots \sum_{h_k} \left(\prod_{j=1}^k \binom{s_j}{h_{1j}, h_{2j}, \dots, h_{nj}}\right) \left(\prod_{i=1}^n W_i^{h_{i*}}\right) \prod_{j=1}^k \prod_{i=1}^n X_{ij}^{h_{ij}}\right) \end{aligned}$$

(where  $h_{i*} = \sum_{j=1}^k h_{ij}$ ) and also

$$= \sum_{h_1} \cdots \sum_{h_k} \left(\prod_{j=1}^k \binom{s_j}{h_{1j}, h_{2j}, \dots, h_{nj}}\right) \mathbb{E}\left(\prod_{i=1}^n W_i^{h_{i*}}\right) \mathbb{E}\left(\prod_{j=1}^k \prod_{i=1}^n X_{ij}^{h_{ij}}\right) \quad (\ddagger)$$

By the Dirichlet distribution we have

$$\mathbb{E}\left(\prod_{i=1}^n W_i^{h_{i*}}\right) = \frac{\Gamma(\sum_{i=1}^n \sum_{j=1}^k \alpha_j^{(i)})}{\Gamma(\sum_{i=1}^n \sum_{j=1}^k \alpha_j^{(i)} + \sum_{j=1}^k s_j)} \prod_{i=1}^n \frac{\Gamma(\sum_{j=1}^k \alpha_j^{(i)} + h_{i*})}{\Gamma(\sum_{j=1}^k \alpha_j^{(i)})}$$

and also

$$\mathbb{E}\left(\prod_{j=1}^k \prod_{i=1}^n X_{ij}^{h_{ij}}\right) = \prod_{i=1}^n \mathbb{E}\left(\prod_{j=1}^k X_{ij}^{h_{ij}}\right),$$

and again by the Dirichlet distribution

$$\mathbb{E}\left(\prod_{j=1}^k X_{ij}^{h_{ij}}\right) = \frac{\Gamma(\sum_{j=1}^k \alpha_j^{(i)})}{\Gamma(\sum_{j=1}^k \alpha_j^{(i)} + h_{i*})} \prod_{j=1}^k \frac{\Gamma(\alpha_j^{(i)} + h_{ij})}{\Gamma(\alpha_j^{(i)})}.$$

So, by using (‡)

$$\begin{aligned}
&= \sum_{h_1} \cdots \sum_{h_k} \prod_{j=1}^k \binom{s_j}{h_{1j}, h_{2j}, \dots, h_{nj}} \left( \frac{\Gamma(\sum_{i=1}^n \sum_{j=1}^k \alpha_j^{(i)})}{\Gamma(\sum_{i=1}^n \sum_{j=1}^k \alpha_j^{(i)} + \sum_{j=1}^k s_j)} \right) \\
&\quad \prod_{i=1}^n \frac{\Gamma(\sum_{j=1}^k \alpha_j^{(i)} + h_{i.})}{\Gamma(\sum_{j=1}^k \alpha_j^{(i)})} \left( \prod_{i=1}^n \frac{\Gamma(\sum_{j=1}^k \alpha_j^{(i)})}{\Gamma(\sum_{j=1}^k \alpha_j^{(i)} + h_{i.})} \prod_{j=1}^k \frac{\Gamma(\alpha_j^{(i)} + h_{ij})}{\Gamma(\alpha_j^{(i)})} \right) \\
&= \frac{\Gamma(\sum_{i=1}^n \sum_{j=1}^k \alpha_j^{(i)})}{\Gamma(\sum_{i=1}^n \sum_{j=1}^k \alpha_j^{(i)} + \sum_{j=1}^k s_j)} \sum_{h_1} \cdots \sum_{h_k} \prod_{j=1}^k \binom{s_j}{h_{1j}, h_{2j}, \dots, h_{nj}} \\
&\quad \prod_{j=1}^k \prod_{i=1}^n \frac{\Gamma(\alpha_j^{(i)} + h_{ij})}{\Gamma(\alpha_j^{(i)})}.
\end{aligned}$$

By considering the fact that the sum of the Dirichlet-multinomial distributions on their support equals to one, we have

$$= \frac{\Gamma(\sum_{i=1}^n \sum_{j=1}^k \alpha_j^{(i)})}{\Gamma(\sum_{i=1}^n \sum_{j=1}^k \alpha_j^{(i)} + \sum_{j=1}^k s_j)} \prod_{j=1}^k \frac{\Gamma(\sum_{i=1}^n \alpha_j^{(i)} + s_j)}{\Gamma(\sum_{i=1}^n \alpha_j^{(i)})}$$

which is the general moment of the k-variate

$$Dirichlet \left( \sum_{i=1}^n \alpha_1^{(i)}, \sum_{i=1}^n \alpha_2^{(i)}, \dots, \sum_{i=1}^n \alpha_k^{(i)} \right)$$

distribution, and since  $\mathbf{Z}$  is a bounded random variable, its distribution is uniquely determined by its moments. Thus the proof is complete.  $\boxtimes$

### 3.5. A Variant of Theorem 2.1

**Theorem 3.1.** *The distribution of the randomly linear combination  $\mathbf{Z} = \sum_{i=1}^n W_i \mathbf{X}_i$  is*

$$Dirichlet \left( \frac{1}{2} + \sum_{i=1}^n \alpha_i \right),$$



where  $\mathbf{X}_1, \dots, \mathbf{X}_n$  are two-dimensional independent multivariate random vectors with

$$\text{Dirichlet}\left(\frac{1}{2} + \alpha_1\right), \dots, \text{Dirichlet}\left(\frac{1}{2} + \alpha_n\right)$$

distributions and the random vector  $\mathbf{W} = \langle W_1, \dots, W_n \rangle$  is independent from  $(\mathbf{X}_1, \dots, \mathbf{X}_n)$  and has the distribution

$$\text{Dirichlet}(\alpha_1, \dots, \alpha_n).$$

*Proof.* Let  $Y_j$  ( $j = 1, \dots, n$ ) be independent random variables independent from  $(\mathbf{X}_1, \dots, \mathbf{X}_n)$  that have the distribution  $\Gamma(\alpha_j, \mu)$ , respectively. It can be seen, by some classic ways (e.g.  $E(e^{\mathbf{t}'\mathbf{T}}) = [\Psi(\mathbf{t})]^{\sum_j \alpha_j}$  from Kubo & Kuo & Namli (2013), Table 2), that the distribution of  $\mathbf{T} (= \sum_j \mathbf{T}_j = \sum_j Y_j \mathbf{X}_j)$  is the same distribution of  $\mathbf{T}_j$  with the parameter  $\sum_j \alpha_j$ . We can also write  $\mathbf{T}$  as

$$\mathbf{T} = \sum_j Y_j \mathbf{X}_j = \left( \sum_i Y_i \right) \left( \sum_j \frac{Y_j}{\sum_i Y_i} \mathbf{X}_j \right)$$

and so we have  $\mathbf{T} = Y\mathbf{Z}$  in which  $Y$  has the gamma distribution with the parameter  $\sum_j \alpha_j$ , and  $\mathbf{T}$  has the  $F$  distribution with the parameter  $\sum_j \alpha_j$ , and  $Y$  and  $\mathbf{Z}$  are independent from each other. Of course, one can define  $\mathbf{T}' = Y'\mathbf{X}'$  in such a way that  $\mathbf{T}' \stackrel{d}{=} \mathbf{T}$ ,  $\sum_j Y_j \stackrel{d}{=} Y'$  and  $\mathbf{X}' \sim \text{Dirichlet}\left(\frac{1}{2} + \sum_j \alpha_j\right)$ . One can conclude that  $\mathbf{Z}$  and  $\mathbf{X}'$  have identical distributions by calculating the general moments  $(s_1, s_2)$  of  $\mathbf{T}$  and  $\mathbf{T}'$ , i.e.,  $E(Z_1^{s_1} Z_2^{s_2}) = E((X'_1)^{s_1} (X'_2)^{s_2})$ .  $\boxtimes$

Actually, the above proof also shows that:

**Theorem 3.2.** *Let  $Y_j$  ( $j = 1, \dots, n$ ) be independent random variables independent from  $(\mathbf{X}_1, \dots, \mathbf{X}_n)$  that have the distribution  $\Gamma(\alpha_j, \mu)$ , respectively,*

where  $\mathbf{X}_i$ 's are independent from each other and have Dirichlet distributions. If  $\mathbf{X}$  has a bounded support and the independent random variable  $Y$  has  $\Gamma(\sum_j \alpha_j, \mu)$  distribution such that  $\sum_i Y_i \mathbf{X}_i \stackrel{d}{=} Y \mathbf{X}$ , then  $\mathbf{X}$  and  $\mathbf{Z} = \sum_i W_i \mathbf{X}_i$  have identical distributions, where  $\mathbf{W} = \langle W_1, \dots, W_n \rangle$  is independent from  $\mathbf{X}_i$ 's and has Dirichlet( $\alpha_1, \dots, \alpha_n$ ) distribution.

#### 4. Some Applications in Stochastic Differential Equations

In this section, using Theorem 2.1 and Corollary 2.2, we prove some interesting mathematical facts. As an example consider the following differential equation for each  $n$  (cf. Homei (2014)):

$$(1) \quad \frac{(-1)^{n-1}}{(n-1)!} \frac{n-1}{2} \frac{d^{n-1}}{dz^{n-1}} \int_0^1 (1-t)^{(n-3)/2} (z^2-t)^{-1/2} dt = \left( \frac{1}{\sqrt{z^2-1}} \right)^n,$$

which could be of interest for some authors, who first guess the solution and then, by using techniques like Leibniz differentiations or change of variables or integration by part, prove that it satisfies the equation (by some long inductive arguments).

Let us recall that Theorem 1 of Homei (2015) identifies the distribution of (the 1-dimensional)  $Z$  from the distributions of  $X_i$ 's by means of the differential equation:

$$(2) \quad \frac{(-1)^{n^*-1}}{(n^*-1)!} \frac{d^{n^*-1}}{dz^{n^*-1}} \mathcal{S}(F_Z, z) = \prod_{i=1}^n \frac{(-1)^{m_i-1}}{(m_i-1)!} \frac{d^{m_i-1}}{dz^{m_i-1}} \mathcal{S}(F_{X_i}, z),$$

where  $F_Y$  denotes the cumulative distribution function of a random variable  $Y$  and  $\mathcal{S}(F_Y, z)$  is defined by  $\mathcal{S}(F_Y, z) = \int_{\mathbb{R}} \frac{1}{z-x} F_Y(dx)$  for  $z \in \mathbb{C} \cap (\text{supp} F_Y)^c$  in which  $\text{supp} F_Y$  stands for the support of  $F_Y$ .

Let the distribution of  $\mathbf{X}_i$  ( $i = 1, \dots, n$ ) be Arcsin, that is  $\mathcal{S}(F_{X_i}, z) = \frac{1}{\sqrt{z^2-1}}$ . Also, let  $m_i = 1$  ( $i = 1, \dots, n$ ) and  $n^* = n$ . Then from the equation (2) we will have

$$(3) \quad \frac{(-1)^{n-1}}{(n-1)!} \frac{d^{n-1}}{dz^{n-1}} \mathcal{S}(F_Z, z) = \left( \frac{1}{\sqrt{z^2-1}} \right)^n.$$

The solution of the equation (3) identifies the Stieltjes transformation of the distribution of  $Z$ . Alternatively, from Theorem 2.1 (or Corollary 2.2) the distribution of  $Z$  is power semicircle (see Homei (2014) for more details). Since the Stieltjes transformation of the power semicircle distribution is  $\frac{n-2}{2} \int_0^1 (1-t)^{(n-3)/2} (z^2-t)^{-1/2} dt$  (see e.g. Arizmendi & Perez-Abreu (2010)) then by substituting it in the equation (2) we will get the equation (1) immediately.

As another example consider the moment generating function on the vector  $\mathbf{T}_j$ :

$$M_{\mathbf{T}_j}(\mathbf{t}') = E(\exp(\mathbf{t}'Y_j\mathbf{X}_j)).$$

Using the double conditional expectation and the fact that the components of  $\mathbf{T}_j$  are independent with gamma distributions we have

$$E\left(\frac{1}{1-\mathbf{t}'\mathbf{X}_j}\right)^{\sum_{i=1}^k \alpha_i^{(j)}} = \prod_{i=1}^k \left(\frac{1}{1-t_i}\right)^{\alpha_i^{(j)}}$$

which proves Proposition 4.4 of Kerov & Tsilevich (2004) (cf. also Karlin & Micchelli & Rinott (1986), page 77).

## 5. Conclusions

Some sporadic works of other authors have been unified here; the main result (Theorem 2.1) has several other different proofs (available upon request)

each of which can have various applications (the five proofs presented here are dedicated to all the family members of Professor A.R. SOLTANI). Our method reveals the advantage of the method of Stieltjes transforms for identifying the distribution of stochastic linear combinations, first used by van Assche (1987) and later generalized by others.

## References

## References

- [1] ALAMATSAZ, M.H. (1993). On Characterizations of Exponential and Gamma Distributions, *Statistics and Probability Letters* 17, 315–319.
- [2] ARIZMENDI, O., PEREZ-ABREU, V. (2010). On the Non-classical Infinite Divisibility of Power Semicircle Distributions, *Communications on Stochastic Analysis* 4, 161–178.
- [3] HOMEI, H. (2015). A Novel Extension of Randomly Weighted Averages, *Statistical Papers* 56, 933–946.
- [4] HOMEI, H. (2014). Characterizations of Arcsin and Related Distributions Based on a New Generalized Unimodality, *Communications in Statistics – Theory and Methods*, to appear. doi: 10.1080/03610926.2015.1006788
- [5] HOMEI, H. (2013). Uniform Random Sample and Symmetric Beta Distribution, *arXiv:1309.2779 [math.ST]*.
- [6] HOMEI, H. (2012). Randomly Weighted Averages with Beta Random Proportions. *Statistics and Probability Letters* 82, 1515–1520.

- [7] JOHNSON, N.L., KOTZ, S. (1990). Randomly Weighted Averages: Some Aspects and Extensions, *The American Statistician* 44, 245–249.
- [8] JOHNSON, N.L., KOTZ, S. (1990a). Use of Moments in Deriving Distributions and Some Characterizations, *The Mathematical Scientist* 15, 42–52.
- [9] KARLIN, S., MICHELLI, C. A., RINOTT, Y. (1986). Multivariate Splines: a probabilistic perspective. *Journal of Multivariate Analysis* 20, 69–90.
- [10] KEROV, S. V. E., TSILEVICH, N. V. (2004). The Markov-Krein correspondence in several dimensions. *Journal of Mathematical Sciences* 121, 2345–2359.
- [11] NADARAJAH, S., KOTZ, S. (2005). On the Product and Ratio of Gamma and Beta Random Variables, *Allgemeines Statistisches Archiv* 89, 435-449.
- [12] REZAPOUR, M., ALAMATSAZ, M.H. (2014). Stochastic Comparison of Lifetimes of Two  $(n-k+1)$ -out-of- $n$  Systems with Heterogeneous Dependent Components, *Journal of Multivariate Analysis* 130, 240–251.
- [13] SETHURAMAN, J. (1994). A Constructive Definition of Dirichlet Priors, *Statistica Sinica* 4, 639–650.
- [14] SOLTANI, A.R. AND HOMEI, H. (2009). Weighted Averages with Random Proportions that are Jointly Uniformly Distributed over the Unit Simplex. *Statistics and Probability Letters* 79, 1215–1218.

- [15] VAN ASSCHE, W. (1987). A Random Variable Uniformly Distributed Between two Independent Random Variables, *Sankhyā: The Indian Journal of Statistics—Series A* 49, 207–211.
- [16] VOLODIN, N.A., KOTZ, S., JOHNSON, N.L. (1993). Use of Moments in Distribution Theory: A Multivariate Case, *Journal of Multivariate Analysis* 46, 112–119.