

General immunity and superadditivity of two-way Gaussian quantum cryptography

Carlo Ottaviani* and Stefano Pirandola

Department of Computer Science & York Centre for Quantum Technologies, University of York, YO10 5GH, UK

(Dated: March 7, 2022)

We consider two-way continuous-variable quantum key distribution, studying its security against general eavesdropping strategies. Assuming the asymptotic limit of many signals exchanged, we prove that two-way Gaussian protocols are immune to coherent attacks. More precisely we show the general superadditivity of the two-way security thresholds, which are proven to be higher than the corresponding one-way counterparts in all cases. We perform the security analysis first reducing the general eavesdropping to a two-mode coherent Gaussian attack, and then showing that the superadditivity is achieved by exploiting the random on/off switching of the two-way quantum communication. This allows the parties to choose the appropriate communication instances to prepare the key, accordingly to the tomography of the quantum channel. The random opening and closing of the circuit represents, in fact, an additional degree of freedom allowing the parties to convert, a posteriori, the two-mode correlations of the eavesdropping into noise. The eavesdropper is assumed to have no access to the on/off switching and, indeed, cannot adapt her attack. We explicitly prove that this mechanism enhances the security performance, no matter if the eavesdropper performs collective or coherent attacks.

PACS numbers: 03.67.Dd, 03.65.-w, 42.50.-p, 89.70.Cf

Introduction

Quantum Key Distribution (QKD) [1] is today one of the most advanced quantum technologies among those emerged from the fundamental research in quantum information. Rapidly progressing towards practical implementations [2], the interest in QKD is motivated by the promise of achieving efficient distribution of cryptographic keys over insecure channels. In fact its main goal is to provide an information-theoretic secure strategy to share cryptographic keys in order to replace the current computationally-secure solution [4], which has been proved to be vulnerable [3] to attacks by quantum computers.

The typical scenario involves two parties, Alice and Bob, who want to share a secret message over an insecure channel [5]. To achieve this goal they encode classical information in non-orthogonal quantum states, which are sent over a noisy quantum channel under control of an eavesdropper, Eve. The standard assumptions to analyze the security of QKD protocols are the following: Eve is computationally unbounded, but has no-access to the parties' private spaces [2, 5] and, most importantly, she is restricted by the no-cloning theorem [6]. The distribution of private keys is possible because any attempt to extract the encoded information unavoidably introduces noise on the quantum states. Monitoring this noise the parties can quantify how much Eve has learnt on the secret key and, consequently, apply classical error correction and privacy amplification protocols reducing Eve's information to a negligible amount. Once they have distilled such a key, the parties can safely use the one-time pad protocol. In

case the level of noise is too high, above the security threshold, they can abort the protocol (denial of service).

The first theoretical proposals for QKD protocols have been designed for discrete variables (DV) [1] systems. Today several remarkable implementation of DV-QKD have been achieved in both fibers [7] and free space [8]. Beside this approach, several protocols exploiting quantum continuous-variable (CV) systems have been put forward [9–14]. In CV-QKD [15] the information is encoded in quantum systems with continuous spectra (infinite-dimensional Hilbert space), and a special attention has been devoted to Gaussian CV systems [16].

Gaussian CV-QKD has been achieved in *in-field* implementations [17], with practical performances comparable to those of DV-QKD, despite the latter appears to be more robust for long-distances. The result of Ref. [17] has been possible combining efficient reconciliation protocols [11], post-selection [18] techniques and efficient classical compression codes [19]. The interest in optical CV systems, for quantum information purposes, is now growing, boosted mainly by the natural properties of these systems: relatively cheap experimental implementation, higher rates, broadband detection techniques [20], and the possibility of exploiting a wide range of frequencies [21, 22]. These natural properties make CV-QKD a promising candidate for future practical real-world implementations, especially in the mid-range distances like the metropolitan areas where high rates are desirable [23].

Today, many theoretical efforts are devoted to the design of device independent (DI) QKD protocols [24, 25]. Despite recent remarkable results, the practical implementation of this approach remains still difficult [26–28]. Very likely the next generation of end-to-end quantum networks will use the recently introduced [29, 30] measurement device independent QKD (MDI-QKD) which allows the distribution of cryptographic keys preserving the protection against the most typical side-channel at-

*Electronic address: Carlo.Ottaviani@york.ac.uk

tacks, without the need to pass a Bell test (see Ref. [29] for a general security analysis). Recently a very high-rate CV-MDI QKD protocol has been proposed and tested in a proof of principle experiment [23, 31, 32].

Alongside the study of end-to-end QKD, it is also of great interest the design of more robust *point-to-point* QKD schemes improving the security performances of CV-QKD in noisier environments [14] or able to exploit trusted noise [33] to implement QKD at different frequencies [21]. In this regard, the two-way protocols [14], where the parties make a double use of the quantum channel to improve the tolerance to noise, show higher security thresholds than the one-way counterparts. This idea has been also extended to thermal QKD [34]. Also note that the two-way protocols have been developed for DV-QKD [35, 36] and can be used for direct quantum communication [37, 38].

The main result in this work is the explicit study of the asymptotic security of two-way Gaussian protocols against coherent attacks, and the proof that these schemes are in fact immune to this eavesdropping. The general strategy to achieve this goal follows a previous insight [14] and can be summarized as follows (see Fig. 1): The parties randomly switch ON or OFF the two-way communication line, and they post-select the OFF instances if they detect the presence of coherent attacks, otherwise they use the ON instances. We explicitly study the security threshold of the OFF configuration against two-mode coherent attacks, which are the residual eavesdropping after de Finetti reduction. Our approach allows us to prove that the superadditivity of the two-way thresholds is a general feature. This result can also be understood noting that the ON/OFF switching activates an additional degree of freedom, exclusive to the parties, which can be used to convert (a-posteriori) Eve's correlations into a noise on which Eve has no control.

Results

The Scheme

In Fig. 1 we describe a two-way quantum communication protocol. We focus on use of coherent states, for the encoding, and heterodyne detections for the decoding [10, 14]. Bob prepares a Gaussian-modulated reference coherent state with density matrix $\rho(\beta)$, and use the quantum channel to transmit it to Alice who, randomly, decides to close (case ON) or open (case OFF) the quantum communication. Let discuss the two cases in detail.

Case ON: Alice encoding is performed by applying a Gaussian-modulated displacement $\hat{D}(\alpha)$ on the reference state $\rho(\beta)$, obtaining a new coherent state with density matrix $\rho(\alpha + \beta)$. This is sent back to Bob who performs heterodyne detection on the received state $\rho(\alpha + \beta)$, and applies classical post-processing in order to subtract the reference variable β and derive Bob's estimate $\hat{\beta}$ of Alice's




FIG. 1: Two-way CV-QKD protocol. Steps: *forward*, Bob prepares coherent states of amplitude β and density matrix $\rho(\beta)$ and sends them through the noisy channel. Using the circuit in ON configuration, Alice applies a random displacement $D(\alpha)$ on $\rho(\beta)$ encoding information in the amplitude α . *Backward*, Alice then sends the quantum state $\rho(\alpha + \beta)$ to Bob who applies heterodyne detection with outcome $\gamma \simeq \alpha + \beta$, and applies classical post-processing to subtract the reference amplitude β to recover α . In OFF configuration, the circuit is opened at Alice's station. She applies heterodyne detection on the reference state, obtaining the variable α . She then prepares a new coherent state $\rho(\alpha')$ which is sent back to Bob who heterodynes this state obtaining the variable β' . During the quantum communication Eve, as well as Bob, does not know which setup of the circuit has been adopted. For this reason, she is forced to attack both communication steps, and cannot adapt her attack to the ON/OFF setup.

variable α .

Case OFF. Alice applies heterodyne detection on the reference state $\rho(\beta)$ with outcome α . Then, she prepares a new Gaussian-modulated coherent state $\rho(\alpha')$ which is sent back to Bob, who heterodynes it with outcome β' . After this, the parties can use the two pairs of variables $\{\alpha, \beta\}$ and $\{\alpha', \beta'\}$ to prepare the key.

We note that, during the quantum communication, both Bob and Eve do not know the configuration adopted. This information is shared during the phase of parameter estimation and is part of the classical communication performed by Alice over the public channel. In the following we focus on the use of reverse reconciliation (RR) [11, 16] (direct reconciliation is discussed in the supplemental material). With the quantum communication in ON, the RR corresponds to Alice inferring Bob's final outcome variable $\hat{\beta}$. With the circuit in OFF, the RR corresponds to Bob estimating Alice's variable α during the forward stage, followed by Alice estimating Bob's detection variable β' .

As described in Ref. [14] the advantage of having the ON/OFF switching is that this degree of freedom can be used to post-select the data in order to prepare the key. After the channel tomography they can determine which attack has been performed and in which status of the

circuit. Then they keep data from case ON when they detect a collective attack, while they use data exchanged with the circuit in OFF when the attack is coherent.

Security analysis and attack reduction

We study the security of the scheme assuming the asymptotic limit of many uses of the quantum channel, $N \gg 1$. In the worst-case scenario the eavesdropper attaches ancillary quantum systems, E_k , to each exchanged signal, and process the E_k 's by a global coherent unitary operation. The ancillary output modes are stored in a quantum memory (QM), and coherently measured after the classical communication between Alice and Bob at the end of the protocol. Such an eavesdropping defines a general coherent attack.

The parties can now reduce the complexity of the previous scenario, by applying symmetric random permutations [41] of their classical data. This allows them to get rid of all the correlations between distinct instances of the protocol. It is then clear that, in the case of two-way communication, the de Finetti symmetrization provides a residual two-mode coherent attack, where the only surviving correlations are those between E_1 and E_2 , used by Eve in each single round-trip. These ancillary modes are mixed with the forward and backward signals by means of beam splitters. Note that we can rid of additional modes \mathbf{e} because we work in the asymptotic limit and we bound Eve's accessible information using the Holevo function [39]. Finally, a further simplification comes from the extremality of Gaussian states [16], which means that we can restrict Eve's input $\rho_{E_1 E_2}$ to be a Gaussian state.

The Gaussian attack is collective, when E_1 , E_2 are uncorrelated, or two-mode coherent when they are correlated. Studying this second case with the circuit in ON and in DR, Ref. [40] found that optimal two-mode attacks exist which can reduce the security performances of the two-way protocol below the one-way threshold. Here we show that, using the scheme with the ON/OFF switching, and post-selecting the optimal key-rate accordingly to the attack detected, the parties can overcome this problem. The security analysis is performed according to the setup shown in Fig. 2, where Fig. 2(a) refers to collective attacks, while Fig. 2(b) refers to two-mode coherent attacks. In the latter case, the security analysis is performed in the entanglement based (EB) representation [11, 16].

Description of the two-mode Gaussian attack in the EB representation

In EB representation both Bob and Alice remotely prepare coherent states on the travelling modes B'_1 and A'_2 by using two-mode squeezed vacuum states, described by

covariance matrices (CMs) of the following form

$$V_{B_1 B'_1} = \begin{pmatrix} \mu_B \mathbf{I} & \sqrt{\mu_B^2 - 1} \mathbf{Z} \\ \sqrt{\mu_B^2 - 1} \mathbf{Z} & \mu_B \mathbf{I} \end{pmatrix}, \quad (1)$$

$$V_{A_2 A'_2} = \begin{pmatrix} \mu_A \mathbf{I} & \sqrt{\mu_A^2 - 1} \mathbf{Z} \\ \sqrt{\mu_A^2 - 1} \mathbf{Z} & \mu_A \mathbf{I} \end{pmatrix}, \quad (2)$$

on which they apply heterodyne detections on the respective local modes B_1 and A_2 . The parameters $\mu_A = \mu + 1$, and $\mu_B = \mu + 1$ describe the variance of the thermal state injected by Alice and Bob, respectively. The two travelling modes, B'_1 and A'_2 , are mixed with Eve's modes, E_1 and E_2 , on two identical beam splitters, with transmissivity T . Eve's input state $\rho_{E_1 E_2}$ is a zero mean, two-mode correlated thermal state, with CM

$$V_{E_1 E_2} = \begin{pmatrix} \omega \mathbf{I} & \mathbf{G} \\ \mathbf{G} & \omega \mathbf{I} \end{pmatrix}, \text{ with } \mathbf{G} = \begin{pmatrix} g & \\ & g' \end{pmatrix}, \quad (3)$$

where $\omega \geq 1$ gives the variance of the thermal noise injected, while g and g' describe the correlations between the two ancillas.

Note that the double use of the channel corresponds to a sequential use of the same communication line (optical fibre), so it is reasonable to consider a symmetric channel (T and ω are the same during the forward and backward communication). The correlation parameters g and g' must fulfill the following constraints

$$|g| < \omega, |g'| < \omega, \omega^2 + gg' - 1 \geq \omega |g + g'|, \quad (4)$$

in order to certify that $V_{E_1 E_2}$ is a bona fide CM. If $g' = g$, we must have $1 - \omega \leq g \leq \omega - 1$, with the two extremal conditions corresponding to E_1 and E_2 sharing maximal separable correlations [42, 43]. If $g' = -g$ the ancillas share non-separable correlations. The Eq. (4) provides the bound $-\sqrt{\omega^2 - 1} \leq g \leq \sqrt{\omega^2 - 1}$, with the extremal values corresponding to maximally entangled states. Finally, if $g = g' = 0$, the two ancillas are not correlated, and the two-mode attack collapses to a standard collective one, based on two independent entangling cloners.

Key-rates and security thresholds

We compute now the secret-key rate $R := I - I_E$, where I is Alice-Bob mutual information, and I_E is Eve's accessible information. In the asymptotic case $N \gg 1$, I_E can be replaced by the Holevo information χ . Hence we write

$$R := I - \chi. \quad (5)$$

The goal of the security analysis is the computation of the bound χ , which is defined as

$$\chi := S_E - S_{E|\mathbf{x}}. \quad (6)$$

The functional S_E is the von Neumann entropy, relative to Eve's quantum state ρ_E , and $S_{E|\mathbf{x}}$ is that corresponding to $\rho_{E|\mathbf{x}}$, which describes Eve's state conditioned on the outcomes of the measurements performed by the parties.




FIG. 2: Panel (a) shows the two-way scheme in ON configuration. This is used against collective Gaussian attacks, typically implemented by means of two independent entangling cloners. Each beam-splitter has transmissivity T , and Eve mixes the ancillas E'_1 and E'_2 with the signals modes B_1 and A_2 . Panel (b) describes the OFF configuration, whose security is studied against two-mode Gaussian attacks. In this case we study the security of the scheme in the entanglement based representation.

Against collective attacks, the parties use the protocol in ON, and we have the following ON key-rate

$$R_{ON} := I_{ON}(\alpha : \tilde{\beta}) - \chi_{ON}(\varepsilon : \tilde{\beta}). \quad (7)$$

By contrast, against coherent attacks, they use the circuit in OFF, for which we have the following OFF key-rate

$$R_{OFF} = I_{OFF} - \chi_{OFF}, \quad (8)$$

where

$$I_{OFF} = \frac{I_{OFF}(\alpha : \beta) + I_{OFF}(\alpha' : \beta')}{2} \quad (9)$$

is the mutual information averaged over the forward and backward use, and

$$\chi_{OFF} := S_{AB} - S_{AB|\alpha, \beta'} \quad (10)$$

is computed on Alice and Bob's output state ρ_{AB} .

Thus, for collective attacks we select the ON key-rate R_{ON} , while for coherent attacks we select the OFF key-rate R_{OFF} . Both these key-rates are function of channel parameters, i.e., transmissivity T and excess noise $N := (1 - T)(\omega - 1)/T$ (which gives the extra noise on the channel with respect the vacuum shot-noise). The OFF key-rate, R_{OFF} , is also function of the correlation parameters g and g' . Therefore, once we have R , we find the security thresholds solving the following equation

$$R(T, N, g, g') = 0. \quad (11)$$

This condition provides threshold curves of the type $N = N(T, g, g')$ which simplifies to $N = N(T)$ for collective attacks.

Formulas for the key-rates

The computation of the secret-key rates can be performed using the mathematical tools described in

Ref. [16]. From the knowledge of the CM describing the total and conditional states, we can compute the von Neumann entropies and finally the key rates. For the protocol with coherent states and heterodyne detection we find the following key-rates

$$R_{ON} = \log \frac{2T(1+T)}{e(1-T)(1+\Lambda)} + \sum_{i=1}^3 h(\bar{\nu}_i) - 2h(\omega), \quad (12)$$

$$R_{OFF} = \log \frac{2T}{e(1-T)(1+\tilde{\Lambda})} + \sum_{j=\pm} \frac{h(\bar{\nu}'_j) - h(\nu_j)}{2}, \quad (13)$$

where

$$h(x) := \frac{x+1}{2} \log \frac{x+1}{2} - \frac{x-1}{2} \log \frac{x-1}{2}.$$

In the previous formulas, the symplectic eigenvalues $\bar{\nu}_i$ are computed numerically and we define $\Lambda := T^2 + (1 - T^2)\omega$ and $\tilde{\Lambda} := T + (1 - T)\omega$. It is of particular interest the OFF key-rate of Eq. (13), from which we notice that one can recover the one-way key-rate in the case of collective attacks ($g = g' = 0$). The expressions of the total and conditional symplectic eigenvalues can be computed analytically for large modulation, being equal to

$$\nu_{\pm} \rightarrow \sqrt{(\omega \pm g)(\omega \pm g')}, \quad (14)$$

$$\bar{\nu}'_{\pm} \rightarrow \frac{\sqrt{[\lambda_{\pm} + 1 - T][\lambda'_{\pm} + 1 - T]}}{T}, \quad (15)$$

where $\lambda_{\pm} = T + (\omega \pm g)(1 - T)$ and $\lambda'_{\pm} = T + (\omega \pm g')(1 - T)$.

Protocol with coherent states and homodyne detection

Here we give the key-rate \tilde{R} for the two-way protocol with coherent states and homodyne detection. The only

change with respect to the previous scheme is clearly the use of homodyne detection for decoding. With the circuit in ON, Bob prepares coherent states, randomly displaced by Alice and finally homodyned by Bob. With the protocol in OFF, Bob prepares coherent states and Alice performs homodyne detection. Then she sends back newly prepared coherent states which are homodyned by Bob. After some algebra, we obtain the following analytical expressions for the key-rates

$$\tilde{R}_{ON} = \frac{1}{2} \log \frac{T^2 + \omega + T^3(\omega - 1)}{(1 - T)\Lambda} + h(\tilde{\nu}) - h(\omega), \quad (16)$$

$$\tilde{R}_{OFF} = \frac{1}{2} \log \frac{\sqrt[4]{(\omega^2 - g^2)(\omega^2 - g'^2)}}{(1 - T)\tilde{\Lambda}} - \sum_{i=\pm} \frac{h(\nu_i)}{2}. \quad (17)$$

In the ON key-rate of Eq. (16), used against collective attacks, the asymptotic symplectic eigenvalue $\tilde{\nu}$ can be computed analytically as

$$\tilde{\nu} := \sqrt{\frac{\omega[1 + T^2\omega(1 - T) + T^3]}{T^2 + \omega + T^3(\omega - 1)}}.$$

By contrast, the OFF key-rate of Eq. (17) is exploited under coherent attacks, and the total symplectic eigenvalues ν_i are the same as given in Eq. (14). The details to obtain Eqs. (16) and (17) can be found in the supplemental material, where we also include the secret-key rates computed in DR.

Discussion

The security analysis of the thresholds coming from Eqs. (12) and (13) is summarized in Fig. 3. In particular, the security threshold for the ON configuration is confirmed [14] to be superadditive in Fig. 3(top-left). The black-solid line corresponds to the ON threshold, which is clearly above the threshold of the one-way protocol (dashed line). This comparison is done against collective attacks. The top-right panel of Fig. 3, shows the security threshold for the OFF configuration in the presence of several two-mode attacks with different values of the correlation parameters $g = g'$. The curves labeled (a)-(d) corresponds to the points in Fig. 3 (bottom-left). For example, the red curve (a) describes coherent attacks performed with maximally entangled states. The curve (b) describes attacks with $g' = g = \pm(\omega - 1)$, and the curve (c) corresponds to $g' = -g = \pm(\omega - 1)$. Finally, the dashed curve (d) corresponds to the center of the correlation plane, where the OFF threshold coincides with the one-way threshold. Thus, we see that for any value of Eve's correlation parameters, g and g' , Alice and Bob can always post-select an instance of the two-way protocol (ON or OFF) whose threshold is strictly greater than that of the corresponding one-way protocol.




FIG. 3: This figure summarizes the results for the protocol with coherent states and heterodyne detection, whose rates are given in Eqs. (12) and (13). The top panels describe the security thresholds, in terms of tolerable excess noise N versus transmissivity T . In the top-left panel, we consider collective attacks and we compare the ON two-way threshold $R_{ON} = 0$ (black solid line) with the threshold of the one-way protocol (dashed line). In the pink region the two-way protocol is secure, while the one-way counterpart is not. In the top-right panel, we consider coherent attacks and we compare the OFF two-way threshold (a)-(c) with respect to the one-way threshold (d). In particular, curve (a) is obtained for $g = \pm\sqrt{\omega^2 - 1}$, i.e., Eve using maximally entangled states; curve (b) considers the case $g' = g$ with $g = \pm(\omega - 1)$; and curve (c) refers to $g' = -g$ and $g = \pm(\omega - 1)$. Note that curve (d) coincides with the OFF threshold against collective attacks, in which case the protocol is used in ON. The same labels (a)-(d) are used in the bottom-left panel, which describes the various attacks on the correlation plane (g, g') , obtained setting $\omega \simeq 1.097$ in the constraint of Eq. (4). Finally, in the bottom-right panel, we plot the OFF key-rate against coherent attacks (red lines), compared to the quantum mutual information (black lines) describing the correlations of Eve's ancillas. We set $T = 0.3$ and $\omega \simeq 1.097$, so that the one-way rate is $\simeq 0$. We see that the OFF key-rate is always strictly positive and it increases for increasing correlations in the attack.

Finally, Fig. 3 (bottom-right) describes the connection between the OFF key rate and the amount of correlations in Eve's ancillas, as quantified by the quantum mutual information. We can see that the OFF key rate not only is positive (while the one-way rate is always zero) but it also increases with Eve's correlations, which are converted into noise by the OFF configuration. Thus, the ON/OFF switching, together with the post-selection of the correct instances, allows one to implement two-way CV-QKD in a way which is not only secure, but also more robust to excess noise with respect to one-way protocols

under completely general attacks.

Methods

A detailed description of the methods can be found in the supplementary material. The security analysis of the protocol has been performed in the entanglement based representation for the case OFF, so that we could compute the Holevo bound from the study of Alice-Bob CM. For the case ON in RR, we started from the output covariance matrix of Eve, to compute the total von Neumann entropy. We then computed the conditional von Neumann entropy completing Eve's covariance matrix with the Bob's mode on which we applied the heterodyne or the homodyne measurement, accordingly with the case studied.

Conclusions

In this work we have studied the security of two-way CV-QKD addressing, explicitly, the superadditivity of its security threshold against coherent attack. To the best of our knowledge this represents the first attempt of such

a complete study for direct point-to-point two-way protocols. Our security analysis is obtained assuming the asymptotic limit, i.e., large number of signals exchanged and large modulation. This allowed us to find closed formulas for the secret-key rates, from which we have proved that the two-way Gaussian protocols are more robust to excess noise than their one-way counterparts in both collective and coherent attacks.

For this property, it is crucial the random ON/OFF switching of the protocol, so that the eavesdropper's correlations are under control of the parties and they are transformed, if needed, into useless noise. Our analysis contributes to the general understanding of the security properties of two-way protocols and is useful to extend CV-QKD to regime with high excess noise. Future developments could involve the study of this ON/OFF switching strategy in more complex quantum communication scenarios.

Acknowledgments

We acknowledge financial support from the EPSRC via the 'UK Quantum Communications HUB' (Grant no. EP/M013472/1).

-
- [1] Bennet, C.H. & Brassard G. Quantum cryptography: public key distribution and coin tossing, Proceedings of the IEEE International Conference on Computers, Systems, and Signal Processing, Bangalore, p. 175 (1984)
 - [2] Scarani, V., et al. The security of practical quantum key distribution, *Rev. Mod. Phys.* **81**, 1301 (2009).
 - [3] Shor, P.W. Polynomial-time algorithms for prime factorization and discrete logarithms on a quantum computer, *SIAM J. Sci. Statist. Comput.* **26**, 1484, (1997), quant-ph/9508027
 - [4] Rivest, R., Shamir, A., Adleman, L. A method for obtaining digital signatures and public-key cryptosystems, *Communications of the ACM* **21** (2): 120–126, (1978)
 - [5] Gisin N. et al. Quantum cryptography, *Rev. Mod. Phys.* **74**, 145–195 (2002).
 - [6] Wootters, W. & Zurek, W. A single quantum cannot be cloned, *Nature* **299**: 802–803, (1982).
 - [7] Korzh, B. et al. Provably secure and practical quantum key distribution over 307 km of optical fibre, *Nature Phot.* **9**, 163–168, (2015).
 - [8] Ursin, R. et al. Entanglement-based quantum communication over 144 km, *Nature Phys.* **3**, 481 - 486 (2007).
 - [9] Grosshans, F. & Grangier, P. Continuous variable quantum cryptography using coherent states, *Phys. Rev. Lett.* **88**, 057902 (2002).
 - [10] Weedbrook, C., et al. Quantum cryptography without switching, *Phys. Rev. Lett.* **93** (17), 170504 (2004).
 - [11] Grosshans, F., et al. Virtual entanglement and reconciliation protocols for quantum cryptography with continuous variables, *Quantum Information & computation* (2003).
 - [12] Garcia-Patron, R., Cerf, N.J., Continuous-variable quantum key distribution protocols over noisy channels, *Phys. Rev. Lett.* **102**, 130501 (2009)
 - [13] Grosshans, F. et al. High-rate quantum cryptography using Gaussian-modulated coherent states, *Nature* **421**, 238–241 (2003).
 - [14] Pirandola, S. et al. Continuous variable quantum cryptography using two-way quantum communication, *Nature Phys.*, **4**, 726, (2008).
 - [15] Diamanti, E., & Leverrier, E. Distributing secret keys with quantum continuous variables: principle, security and implementations, *Entropy*, **17**, 6072, (2015).
 - [16] Weedbrook, C., et al. Gaussian quantum information, *Rev. Mod. Phys.*, **84**, 621, (2012).
 - [17] Jouguet, P., et al. Experimental demonstration of long-distance continuous-variable quantum key distribution, *Nature Photonics* **7**, 378–381 (2013).
 - [18] Silberhorn, C. et al. Continuous variable quantum cryptography: beating the 3 dB Loss Limit, *Phys. Rev. Lett.* **89**, 167901 (2002).
 - [19] Leverrier, A., et al. Multidimensional reconciliation for a continuous-variable quantum key distribution, *Phys. Rev. A*, **77**, 042325, (2008).
 - [20] Braunstein, S.L. & van Loock, P. Quantum information with continuous variables, *Rev. Mod. Phys.* **77**, 513, (2005).
 - [21] Weedbrook, C., et al. Quantum cryptography approaching the classical limit, *Phys. Rev. Lett.* **105**, 110501 (2010).
 - [22] Weedbrook, C., et al. Continuous-variable quantum key distribution using thermal states, *Phys. Rev. A* **86**, 022318 (2012).
 - [23] Pirandola, S. et al. Reply to 'Discrete and continuous variables for measurement-device-independent quantum cryptography', *Nature Phot.* **9**, 773, (2015).

- [24] Ekert, A., Quantum cryptography based on the Bell's theorem, Phys. Rev. Lett. **69**, 1293, (1992).
- [25] Gisin N. et al. Proposal for implementing device-independent quantum key distribution based on a heralded qubit amplifier, Phys. Rev. Lett. **105**, 070501 (2010).
- [26] Vazirani, U. & Vidick, T. Fully device-independent quantum key distribution, Phys. Rev. Lett. **113**, 140501 (2014).
- [27] Colbeck, R., Victory for the quantum code maker?, Physics **7**, 99 (2014).
- [28] Hensen, B., et al. Experimental loophole-free violation of a Bell inequality using entangled electron spins separated by 1.3 km, arXiv 1508.05949, (2015).
- [29] Braunstein, S.L. & S. Pirandola, Side-channel-free quantum key distribution, Phys. Rev. Lett. **108**, 130502 (2012).
- [30] Lo, H.-K., Curty, M. & Qi, B. Measurement-device-independent quantumkey distribution, Phys. Rev. Lett. **108**, 130503 (2012).
- [31] Pirandola, S. et al. High-rate measurement-device-independent quantum cryptography, Nature. Phot. **9**, 396 (2015). See also arXiv.1312.4104 (2013).
- [32] Ottaviani, C., Spedalieri G., Braunstein S.L. & Pirandola S. Continuous-variable quantum cryptography with an untrusted relay: Detailed security analysis of the symmetric configuration, Phys. Rev. A **91**, 022320 (2015).
- [33] Pirandola, S. Quantum discord as a resource for quantum cryptography, Sci. Rep. **4**, 6956 (2014).
- [34] Weedbrook, C., Ottaviani, C. & Pirandola, S. Two-way quantum cryptography at different wavelengths, Phys. Rev. A, **89**, 012309, (2014).
- [35] Boström, K., Felbinger, T. Deterministic secure direct communication using entanglement, Phys. Rev. Lett. **89**, 187902, (2002).
- [36] Lucamarini, M., & Mancini S. secure deterministic communication without entanglement, Phys. Rev. Lett. **94**, 140501 (2005).
- [37] Shapiro, J.H., Defeating passive eavesdropping with quantum illumination, Phys. Rev. A **80**, 022320 (2009).
- [38] Zhang, Z., et al. Entanglement's benefit survives an entanglement-breaking channel, Phys. Rev. Lett. **111**, 010501 (2013).
- [39] Pirandola, S., Braunstein, S.L. and Lloyd, S., Characterization of collective gaussian attacks and security of coherent-state quantum cryptography, Phys. Rev. Lett. **101**, 200504, (2008).
- [40] Ottaviani, C., Mancini S. and Pirandola, S. Two-way Gaussian quantum cryptography against coherent attacks in direct reconciliation, Phys. Rev. A, **92**, 062323, (2015).
- [41] Renner, R. & Cirac, J.I. de Finetti representation theorem for infinite-dimensional quantum systems and applications to quantum cryptography, Phys. Rev. Lett. **102**, 110504 (2009).
- [42] Pirandola, S. Entanglement reactivation in separable environments, New J. Phys. **15**, 113046, (2013).
- [43] Pirandola, S., Serafini, A., Lloyd, S., Correlation matrices of two-mode bosonic systems, Phys. Rev. A **79**, 052327 (2009).

Supplemental Material for “General immunity and superadditivity of two-way Gaussian quantum cryptography”

This supplemental material gives the details of the calculations for the security analysis of the protocols described in this work, and includes also the key-rates calculated for two-way protocol used in direct reconciliation (DR), which has not been discussed in the main body. We underline that compared to the one-way protocol, the two-way presents a richer number of cases which need to be analyzed. The protocols are named with respect to the *preparation* and *detection* scheme adopted. Here we discuss the protocol with coherent states and heterodyne detection, and with coherent states and homodyne detection. Each one of previous cases can be implemented in DR as well as reverse reconciliation (RR), and here we give the results for both reconciliation schemes.

Secret-Key Rate and symplectic analysis

The secret-key rate quantifies the gap between Alice and Bob's mutual information and the information shared between Eve and the parties. Which parties' variable(s) has(have) to be considered depends on the setup of the protocol (one-way, two-way, ON or OFF) and, in general, from the reconciliation protocol employed.

For instance consider the one-way protocol. We assume that Alice sends a modulated coherent state with amplitude α to Bob, who receives a noisy version of this state, whose amplitude is β . The parties can then obtain two distinct secret-key rates defined as follows

$$R^{\blacktriangleright} := I(\alpha : \beta) - \chi(\varepsilon : \alpha), \quad (\text{S1})$$

$$R^{\blacktriangleleft} := I(\alpha : \beta) - \chi(\varepsilon : \beta). \quad (\text{S2})$$

The first describes the key-rate in DR, while the second the RR. The function I is the classical mutual information quantifying correlations between Alice's variable, α , and Bob's variable, β . For each quadrature measured, and used to encode information, the mutual information is given by the following general signal-to-noise ratio

$$I = \frac{1}{2} \log \frac{V}{V_C}, \quad (\text{S3})$$

where V is the variance of the variable used to prepare the key, and V_C the conditioned variance of this statistical variable after the measurement performed by the parties.

In the asymptotic limit of many uses of the quantum channel we can bound Eve's accessible information by the Holevo function, which is given by

$$\chi(\varepsilon : x) := S(\varepsilon) - S(\varepsilon|x). \quad (\text{S4})$$

The function $S(\cdot)$ describes the von Neumann entropy which, for Gaussian quantum systems, has a simple form

given by

$$S = \sum_k h(\nu_k), \quad (\text{S5})$$

with the entropic function $h(\cdot)$ is defined as follows

$$h(\nu_k) := \frac{\nu_k + 1}{2} \log \frac{\nu_k + 1}{2} - \frac{\nu_k - 1}{2} \log \frac{\nu_k - 1}{2}, \quad (\text{S6})$$

and where the ν_k 's are the symplectic eigenvalues of the CM which describes the dynamics of the studied Gaussian quantum system [S1].

The expression of the von Neumann entropy of Eq. (S6) can be further simplified exploiting the limit of large signal modulation, in which case we can write [S1]

$$h(\nu_k) = \log \frac{e}{2} \nu_k + O\left(\frac{1}{\nu_k}\right). \quad (\text{S7})$$

The computation of the symplectic spectra can be done in prepare and measure configuration, in which case the ν_k 's are obtained from the symplectic analysis of Eve's output CM or, in case we use the equivalent EB representation, from Alice-Bob's CM. This second approach is used in the following, to study the OFF configurations, i.e., when we consider coherent attacks.

To compute the symplectic spectrum, we first compute the appropriate CM \mathbf{V} and then, from matrix

$$\mathbf{M} = i\Omega\mathbf{V},$$

where $\Omega = \oplus_{i=1}^n \tilde{\omega}_i$, with $\tilde{\omega}_i$ the single-mode symplectic form given by

$$\tilde{\omega}_i = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},$$

we compute the ordinary eigenvalues, which come in pairs. The symplectic spectrum is obtained taking their absolute value.

Protocol with coherent states and heterodyne detection

We start showing how we obtain the ON key-rate for the protocol with coherent states and heterodyne detection, which is described in Fig. 2 (a) of the main text. The security analysis is performed using Eve's CM, obtained from the outputs $\{E_1, E_1'', E_2, E_2''\}$. From this we obtain the total von Neumann entropy and, by simple conditioning procedure, one can also compute Eve's conditional CM. This describes the conditional state $\rho_{E_1, E_1'', E_2, E_2''|\alpha}$, for the protocol in DR. By contrast, to study the protocol in RR, we complete Eve's output CM with Bob's post-processed output mode B , on which we apply the heterodyne detection in order to obtain the conditional CM for this case.

Case ON

Bob sends modulated coherent states to Alice providing, on average, a thermal state with variance $\mu_B = \mu + 1$, where μ accounts for the classical Gaussian modulation on the top of the vacuum shot-noise. Alice applies an additional random displacement, $D(\alpha)$, on the states received from Bob with modulation variance $\mu_{ON} = \mu \geq 0$.

Mutual information Alice-Bob mutual information can be computed from the expression of the variance of post-processed mode $\langle B^2 \rangle$, given by

$$\langle B^2 \rangle = [T^2 + T\mu_{ON} + (1 - T^2)\omega]I, \quad (\text{S8})$$

from which, in the limit of large modulation $\mu_{ON} = \mu \rightarrow \infty$, we obtain the signal variance

$$V = T\mu. \quad (\text{S9})$$

We then compute the conditional variance from Eq. (S8) by setting Alice modulation $\mu_{ON} = 0$, obtaining

$$V_C = T^2 + (1 - T^2)\omega. \quad (\text{S10})$$

Finally, using Eqs. (S9) and (S10) with the expression of the mutual information in case of heterodyne detection

$$I := \log \frac{V + 1}{V_C + 1}, \quad (\text{S11})$$

we obtain the following Alice-Bob mutual information in the limit of large modulation

$$I_{ON} = \log \frac{T\mu}{1 + T^2 + (1 - T^2)\omega}. \quad (\text{S12})$$

Total Covariance Matrix We compute now the CM of Eve's output quantum state $\rho_{E_1', E_1'', E_2', E_2''}$. We arrange it in the following normal form

$$\mathbf{V}_E = \begin{pmatrix} \mathbf{A} & \mathbf{C} \\ \mathbf{C}^T & \mathbf{B} \end{pmatrix}, \quad (\text{S13})$$

where

$$\begin{aligned} \mathbf{A} &:= \begin{pmatrix} \omega \mathbf{I} & \sqrt{T(\omega^2 - 1)} \mathbf{Z} \\ \sqrt{T(\omega^2 - 1)} \mathbf{Z} & \Psi \mathbf{I} \end{pmatrix}, \\ \mathbf{B} &:= \begin{pmatrix} \omega \mathbf{I} & \sqrt{T(\omega^2 - 1)} \mathbf{Z} \\ \sqrt{T(\omega^2 - 1)} \mathbf{Z} & \tilde{\Psi} \mathbf{I} \end{pmatrix}, \\ \mathbf{C} &:= \begin{pmatrix} 0 & \Xi \mathbf{Z} \\ 0 & \Phi \mathbf{Z} \end{pmatrix}, \end{aligned} \quad (\text{S14})$$

with

$$\begin{aligned} \tilde{\Psi} &= [T\omega + (1 - T)^2\omega + T(1 - T)\mu_B] + (1 - T)\mu_{ON}, \\ \Psi &= T(\omega - \mu_B) + \mu_B, \\ \Phi &= (1 - T)\sqrt{T}(\mu_B - \omega), \\ \Xi &= -(1 - T)\sqrt{(\omega^2 - 1)}. \end{aligned} \quad (\text{S15})$$

From Eq. (S13) we easily obtain the total symplectic spectrum by taking the limits for $\mu_{ON} = \mu \rightarrow \infty$ and $\mu_b = \mu + 1 \rightarrow \infty$

$$\{\nu_1, \nu_2, \nu_3, \nu_4\} \rightarrow \{\omega, \omega, (1-T)^2\mu^2\}. \quad (\text{S16})$$

The latter, used with Eqs. (S5) and (S7), gives the total von Neumann entropy

$$S_E = \log \frac{e^2}{4} (1-T)^2 \mu^2 + 2h(\omega). \quad (\text{S17})$$

Conditional CM and Key-rate in Direct Reconciliation For the DR the conditional CM can be obtained straightforwardly from Eq. (S13) setting $\mu_{ON} = 0$ on both quadratures in the block describing Eve's output E_2'' , i.e., $\tilde{\Psi}$ in Eq. (S15). The resulting conditional CM has the following asymptotic symplectic spectrum

$$\{\bar{\nu}_1, \bar{\nu}_2, \bar{\nu}_3, \bar{\nu}_4\} \rightarrow \{1, 1, \omega, (1-T^2)\mu\}. \quad (\text{S18})$$

Using Eq. (S18) with Eq. (S5) and (S7), we compute the conditional von Neumann entropy

$$S_{E|\alpha} = \log \frac{e}{2} (1-T^2)\mu + h(\omega). \quad (\text{S19})$$

Now, using Eqs. (S17) and (S19) in Eq. (S4), one obtains the expression of the Holevo bound

$$\chi_{ON}^{\blacktriangleright} = \log \frac{e}{2} \frac{(1-T)}{(1+T)} \mu. \quad (\text{S20})$$

Finally, by subtracting the Holevo function of Eq. (S20) from the mutual information of Eq. (S12) we get the ON key-rate in DR

$$R_{ON}^{\blacktriangleright} = \log \frac{2}{e} \frac{T(1+T)}{(1-T)[1+T^2+(1-T^2)\omega]} - h(\omega).$$

Reverse Reconciliation To study the security of the protocol in RR, we need to re-compute the conditional von Neumann entropy for this case. We complete the CM of Eq. (S13) adding the blocks describing Bob's output mode B and its correlations with the rest of Eve's modes. Then we apply a heterodyne detection on B obtain Eve's conditional CM after Bob's measurements. We then write

$$\mathbf{V}^{\blacktriangleleft} = \begin{pmatrix} \mathbf{V}_E & \bar{\mathbf{C}} \\ \bar{\mathbf{C}}^T & \bar{\mathbf{B}} \end{pmatrix}, \quad (\text{S21})$$

where

$$\bar{\mathbf{B}} = [T^2 + T\mu + (1-T^2)\omega]\mathbf{I},$$

$$\bar{\mathbf{C}} = \sqrt{1-T} \begin{pmatrix} \sqrt{T(\omega^2-1)}\mathbf{Z} \\ T(\omega-1)\mathbf{I} \\ \sqrt{(\omega^2-1)}\mathbf{Z} \\ \sqrt{T}[T(\omega-1)-\mu]\mathbf{I} \end{pmatrix}.$$

We then apply the formula for heterodyne detection [S1] obtaining the following conditional CM

$$\mathbf{V}_C^{\blacktriangleleft} = \mathbf{V}_E + \bar{\mathbf{C}}(\bar{\mathbf{B}} + \mathbf{I})^{-1}\bar{\mathbf{C}}^T, \quad (\text{S22})$$

which gives the following conditional symplectic spectrum

$$\{\bar{\nu}_1, \bar{\nu}_2, \bar{\nu}_3, \bar{\nu}_4\} \rightarrow \{\bar{\nu}_1, \bar{\nu}_2, \bar{\nu}_3, (1-T^2)\mu\}. \quad (\text{S23})$$

Notice that the eigenvalues $\bar{\nu}_1, \bar{\nu}_2, \bar{\nu}_3$ are asymptotically depending only on the channel parameters (ω, T) , and are related by the following expression

$$\bar{\nu}_1 \bar{\nu}_2 \bar{\nu}_3 = \frac{[1+T^3+(1-T)(1+T^2)\omega]\omega}{T(1+T)}.$$

From the eigenvalues of Eq. (S23), used with Eqs. (S4), (S5) and (S7) we obtain the Holevo bound for the RR

$$\chi_{ON}^{\blacktriangleleft} = \log \frac{e}{2} (1-T^2)\mu + h(\bar{\nu}_1) + h(\bar{\nu}_2) + h(\bar{\nu}_3),$$

which, used with Eq. (S12) in the definition of Eq. (7) in the main text, gives the secret-key rate for the protocol used in RR given in Eq. (12) of the main text.

Case OFF

We now describe the details of the calculations for the protocol used in OFF, as described in Fig. 2-(b) of the main text. In this case we perform the security analysis considering two-mode coherent attacks, in the EB representation.

Total Covariance Matrix and von Neumann entropy

Bob starts from a two-mode squeezed vacuum state, described by the CM of Eq. (1) in the main text. Applying a local heterodyne detection on mode B_1 , he projects the traveling mode B_1' in a coherent state. In the same way, Alice applies a local heterodyne detection on mode A_2 , projecting the traveling mode A_2' in a coherent state. Finally, we assume that Eve injects the general Gaussian state described by Eq. (3) in the main text. Since the total state of Alice, Bob and Eve is pure, we can reduce ourselves to compute Alice and Bob's state (having the same entropy of Eve's). We then order Alice's and Bob's the output modes as follows $\{B_1, A_2, A_1, B_2\}$, and obtain the following expression

$$\mathbf{V}_{AB}^{OFF} = \begin{pmatrix} \tilde{\mathbf{A}} & \tilde{\mathbf{C}} \\ \tilde{\mathbf{C}}^T & \tilde{\mathbf{B}} \end{pmatrix}, \quad (\text{S24})$$

where the matrix blocks have been defined as follows,

$$\begin{aligned}\tilde{\mathbf{A}} &= \begin{pmatrix} \mu_B \mathbf{I} & \tilde{\delta} \mathbf{I} \\ \tilde{\delta} \mathbf{I} & \mu_A \mathbf{I} \end{pmatrix}, \\ \tilde{\mathbf{B}} &= \tilde{\tau}(\mu_A) \mathbf{I}, \\ \tilde{\mathbf{C}} &= \begin{pmatrix} \mathbf{0} \\ \tilde{\delta} \mathbf{I} \\ (1-T)\mathbf{G} \end{pmatrix},\end{aligned}$$

where

$$\mathbf{G} := \begin{pmatrix} g \\ g' \end{pmatrix},$$

the coefficients $\tilde{\delta}$ and $\tilde{\tau}(y)$ have been defined as follows,

$$\begin{aligned}\tilde{\delta} &:= \sqrt{T[\mu_B^2 - 1]}, \\ \tilde{\tau}(y) &:= (1-T)\omega + Ty.\end{aligned}\quad (\text{S25})$$

We compute the symplectic spectrum of CM (S24) and taking the asymptotic limit, $\mu_A = \mu_B \rightarrow \infty$, we obtain the following analytical expressions

$$\{\nu_{\pm}, \nu_3, \nu_4\} \rightarrow \{\sqrt{(\omega \pm g)(\omega \pm g')}, (1-T)\mu, (1-T)\mu\}, \quad (\text{S26})$$

which gives the total von Neumann entropy for the case OFF

$$S_{AB} = \log \left(\frac{e}{2} \right)^2 (1-T)^2 \mu^2 + h(\nu_-) + h(\nu_+). \quad (\text{S27})$$

Conditional covariance matrix and Alice-Bob mutual information

To obtain the conditional CM in DR we set $\mu_A = \mu_B = 1$, in modes B_1 and A_2 . It is easy to verify that the resulting CM has the following symplectic spectrum

$$\{\bar{\nu}_1, \bar{\nu}_2, \bar{\nu}_+, \bar{\nu}_-\} \rightarrow \{1, 1, \sqrt{\lambda_+ \lambda'_+}, \sqrt{\lambda_- \lambda'_-}\}, \quad (\text{S28})$$

where, $\lambda_{\pm} = T + (\omega \pm g)(1-T)$ and $\lambda'_{\pm} = T + (\omega \pm g')(1-T)$. Using these eigenvalues, we compute the following conditional von Neumann entropy

$$S_{AB|\alpha', \beta} = h(\bar{\nu}_+) + h(\bar{\nu}_-).$$

The previous equation and Eq. (S27) are then used to obtain the asymptotic expression of the Holevo function in DR, which is given by

$$\chi_{OFF}^{\blacktriangleright} := \frac{S_{AB} - S_{AB|\alpha, \beta'}}{2}. \quad (\text{S29})$$

$$= \log \frac{e}{2} (1-T)\mu + \frac{1}{2} \sum_{k=\pm} [h(\nu_k) - h(\bar{\nu}_k)]. \quad (\text{S30})$$

The conditional CM corresponding to the RR, is obtained by applying two consecutive heterodyne detections, starting from CM of Eq. (S24). We first measure

mode B_2 and then we apply another heterodyne detection on mode A_2 (the order of these two local measurements is of course irrelevant). The resulting conditional CM has the symplectic spectrum

$$\bar{\nu}'_{\pm} \rightarrow \frac{\sqrt{[\lambda_{\pm} + 1 - T][\lambda'_{\pm} + 1 - T]}}{T}.$$

These are used to compute the Holevo bound which is given by

$$\chi_{OFF}^{\blacktriangleleft} = \log \frac{e}{2} (1-T)\mu + \frac{1}{2} \sum_{k=\pm} [h(\nu_k) - h(\bar{\nu}'_k)]. \quad (\text{S31})$$

Alice-Bob Mutual Information and Secret-key rate.

Alice-Bob mutual information is easily computed from the coefficient $\tilde{\tau}$, given in Eq. (S25). Taking the limit of large modulation, using the formula defining the mutual information for heterodyne detections (S11), and averaging over the double use of the quantum channel we obtain the following expression

$$I_{OFF} = \log \frac{T\mu}{1 + \tilde{\Lambda}}, \quad (\text{S32})$$

where $\tilde{\Lambda} = T + (1-T)\omega$. Notice that, differently from the Holevo bound, the mutual information is independent from the correlation parameters described by the matrix \mathbf{G} .

Using Eq. (S30) and Eq. (S32), and after some simple algebra, we get the analytical expression of the key-rate in DR

$$R_{OFF}^{\blacktriangleright} = \log \frac{2T}{e(1-T)(1 + \tilde{\Lambda})} + \frac{1}{2} \sum_{k=\pm} [h(\bar{\nu}_k) - h(\nu_k)], \quad (\text{S33})$$

The key-rate in RR of Eq. (13) in the main text, is obtained using previous Eq. (S31) and Eq. (S32).

Protocol with coherent states and homodyne detection

In contrast to the protocol analysed in previous section, here the decodings are performed by means of homodyne detections. This modifies the expression of the mutual information and these of the conditional von Neumann entropies.

Case ON

Direct Reconciliation: conditional covariance matrix The conditional CM in DR is obtained as before. We start from the total CM of Eq. (S13) and we apply the




Figure S1: This figure summarizes the results for the protocol with coherent states and homodyne detection, whose rates are given in Eqs. (S39) and (S42) which are the same as Eqs. (16) and (17) in the main text. The top panels give the security thresholds in terms of tolerable excess noise N versus transmissivity T . In the top-left panel, we consider collective attacks and we compare the ON two-way threshold $\tilde{R}_{ON} = 0$ (black solid line) with the threshold of the one-way protocol (dashed line). In the pink region the two-way protocol is secure, while the one-way counterpart is not. In the top-right panel, we consider coherent attacks and we compare the OFF two-way threshold (a)-(c) with respect to the one-way threshold (d). In particular, curve (a) is obtained for $g = \pm\sqrt{\omega^2 - 1}$, i.e., Eve using maximally entangled states; curve (b) considers the case $g' = g$ with $g = \pm(\omega - 1)$; and curve (c) refers to $g' = -g$ and $g = \pm(\omega - 1)$. Note that curve (d) coincides with the OFF threshold against collective attacks, in which case the protocol is used in ON. The same labels (a)-(d) are used in the bottom-left panel, which describes the various attacks on the correlation plane (g, g') , obtained setting $\omega \simeq 1.049$ in the constraint of Eq. (4) in the main text. Finally, in the bottom-right panel, we plot the OFF key-rate against coherent attacks (red lines), compared to the quantum mutual information (black lines) describing the correlations of Eve's ancillas. We set $T = 0.2$ and $\omega \simeq 1.049$, so that the one-way rate is $\simeq 0$. We see that the OFF key-rate is always strictly positive and it increases for increasing correlations in the attack.

following conditioning procedure

$$\begin{aligned}\bar{\mu}_{ON}^q &= 1/\mu \xrightarrow{\mu \rightarrow \infty} 0, \\ \bar{\mu}_{ON}^p &= \mu,\end{aligned}\quad (\text{S34})$$

which describes Alice's effective modulation in order to describe the measurement of only one quadrature during the decoding stage (homodyne detection).

Direct Reconciliation: mutual information, Holevo bound and key-rate The conditioning procedure, de-

scribed by Eqs. (S34), can clearly also be used to determine Alice-Bob mutual information. In the present case only one-quadrature is used to encode the key so Alice-Bob mutual information is given by the following expression

$$\tilde{I}_{ON} = \frac{1}{2} \log \frac{T\mu}{T^2 + (1 - T^2)\omega}. \quad (\text{S35})$$

Eve's conditional CM is obtained from Eq. (S13) applying the recipe of Eqs. (S34). One easily obtains the conditional symplectic spectrum which, in the asymptotic limit, is given by

$$\{\nu_1^\blacktriangleleft, \nu_2^\blacktriangleleft, \nu_3^\blacktriangleleft, \nu_4^\blacktriangleleft\} \rightarrow \{1, \omega, \sqrt{(1 - T)^2(1 - T^2)\omega\mu^3}\}. \quad (\text{S36})$$

After some algebra we obtain the following Holevo bound

$$\tilde{\chi}_{ON}^\blacktriangleleft = \frac{1}{2} \log \frac{(1 - T)^2\mu}{(1 - T^2)\omega} + h(\omega),$$

which, used with Eq. (S35), provides the ON key-rate for the protocol in DR

$$\tilde{R}_{ON}^\blacktriangleleft = \frac{1}{2} \log \frac{T(1 + T)\omega}{(1 - T)[T^2 + (1 - T^2)\omega]} - h(\omega). \quad (\text{S37})$$

It is interesting to note that plotting the security threshold of the key-rate of Eq. (S37), one finds that it provides a positive key-rate even below 3 dB, which sets the limit performance of the one-way version of this protocol in DR.

Reverse Reconciliation: conditional covariance matrix The security of the protocol is performed repeating the steps described in previous sections, replacing the heterodyne detections with homodyne measurements on B . Indeed, we apply the following formula

$$\tilde{\mathbf{V}}_C^\blacktriangleleft = \mathbf{A} - \mathbf{C}(\Pi\bar{\mathbf{B}}\Pi)\mathbf{C}^T,$$

to Eq. (S21). Note that $\Pi := \text{diag}(1, 0)$ ($\text{diag}(0, 1)$) for heterodyne on quadrature \hat{q} (\hat{p}). We then compute the symplectic spectrum of CM $\tilde{\mathbf{V}}_C^\blacktriangleleft$, which we rewrite here in the following form

$$\begin{aligned}\tilde{\nu} &\rightarrow \sqrt{\frac{\omega[1 + T^2\omega - T^3(\omega - 1)]}{T^2 + \omega + T^3(\omega - 1)}}, \\ \nu_2^\blacktriangleleft &\rightarrow \omega, \\ \nu_3^\blacktriangleleft \nu_4^\blacktriangleleft &\rightarrow \sqrt{\frac{(1 - T)^3[T^2 + \omega + T^3(\omega - 1)]\mu^3}{T}}.\end{aligned}\quad (\text{S38})$$

We then obtain the Holevo bound $\tilde{\chi}_{ON}^\blacktriangleleft$

$$\tilde{\chi}_{ON}^\blacktriangleleft = h(\omega) - h(\tilde{\nu}) + \frac{1}{2} \log \frac{T(1 - T)\mu}{T^2 + \omega + T^3(\omega - 1)},$$

which combined with the mutual information of Eq. (S35) gives the ON key-rate in RR of Eq. (16) of the main text, i.e.,

$$\tilde{R}_{ON}^\blacktriangleleft = \frac{1}{2} \log \frac{T^2 + \omega + T^3(\omega - 1)}{(1 - T)\Lambda} + h(\tilde{\nu}) - h(\omega), \quad (\text{S39})$$

with $\Lambda := T^2 + (1 - T^2)\omega$.

Case OFF

This case is studied in the EB representation, following the same steps of the previous OFF case, for both the DR and RR, replacing the final heterodyne with homodyne detections. The mutual information is computed averaging over the double use of the quantum channel, i.e., using the following definition of the mutual information

$$\tilde{I}_{OFF} := \frac{1}{2} \left(\frac{1}{2} \sum_{i=A,B} \log \frac{T\mu_i}{(1-T)\omega + T\mu_i} \right)_{\mu_i=\mu \rightarrow \infty} \frac{1}{2} \log \frac{T\mu}{\tilde{\Lambda}}, \quad (\text{S40})$$

where $\tilde{\Lambda} := T + (1-T)\omega$.

Direct Reconciliation

The steps to compute the conditional CM have been discussed previously, so here we just provide the analytical expressions of the conditional symplectic spectra

$$\begin{aligned} \tilde{\nu}_{\pm}^{\blacktriangleright} &= \sqrt{(1-T)\Gamma_{\pm}\mu}, \\ \tilde{\eta}_{\pm}^{\blacktriangleright} &= \sqrt{\frac{(\omega \pm g')[T + (\omega \pm g)(1-T)]}{\Gamma_{\pm}}}, \end{aligned}$$

where we define

$$\Gamma_{\pm} := 1 - T + T(\omega \pm g').$$

Notice that, in previous spectra, the role of the correlation parameters depends on the quadrature measured by the homodyne detection of the decoding stage. We obtain the following Holevo bound for the DR

$$\tilde{\chi}_{OFF}^{\blacktriangleright} = \frac{1}{2} \log \frac{(1-T)\mu}{\sqrt{[1 + t(\omega - 1)]^2 - T^2 g^2}},$$

and the key-rate in direct reconciliation is given by

$$\begin{aligned} \tilde{R}_{OFF}^{\blacktriangleright} &= \frac{1}{2} \log \frac{T\sqrt{[1 + T(\omega - 1)]^2 - T^2 g^2}}{(1-T)[T + (1-T)\omega]} \\ &\quad - \sum_{k=\pm} \frac{h(\tilde{\eta}_k^{\blacktriangleright}) - h(\nu_k)}{2}, \end{aligned}$$

where the eigenvalues ν_{\pm} are defined in Eq. (S26).

Reverse Reconciliation

For the RR, when the homodyne detection is performed on the quadrature \hat{q} , we obtain the following conditional symplectic eigenvalues

$$\tilde{\nu}_{\pm}^{\blacktriangleleft} = \sqrt{\frac{(1-T)(\omega \pm g)\mu}{T}}. \quad (\text{S41})$$

By contrast, in case of homodyne detection on \hat{p} , the corresponding eigenvalues can be obtained from Eq. (S41) by exchanging $g \longleftrightarrow g'$. Averaging over the two detections, we find the following Holevo bound

$$\tilde{\chi}_{OFF}^{\blacktriangleleft} = \sum_{k=\pm} \frac{h(\nu_k)}{2} + \frac{1}{2} \log \frac{T(1-T)\mu}{\sqrt[4]{(\omega^2 - g'^2)(\omega^2 - g^2)}},$$

which subtracted to the mutual information of Eq. (S40) gives the key-rate of Eq. (17) of the main text, i.e.,

$$\tilde{R}_{OFF}^{\blacktriangleleft} = \frac{1}{2} \log \frac{\sqrt[4]{(\omega^2 - g^2)(\omega^2 - g'^2)}}{(1-T)\tilde{\Lambda}} - \sum_{k=\pm} \frac{h(\nu_k)}{2}. \quad (\text{S42})$$

[1] Weedbrook, C., et al. Gaussian quantum information, Rev. Mod. Phys. **84**, 621 (2012).