

OPTION SPANNING BEYOND L_p -MODELS

N. GAO AND F. XANTHOS

ABSTRACT. The aim of this paper is to study the spanning power of options in a static financial market that allows non-integrable assets. Our findings extend and unify the results in [13, 14, 24] for L_p -models. We also apply the spanning power properties to the pricing problem. In particular, we show that prices on call and put options of a limited liability asset can be uniquely extended by arbitrage to all marketed contingent claims written on the asset.

1. INTRODUCTION

Throughout this paper, Ω stands for the state space of a financial market, Σ stands for the σ -algebra modelling the market information structure, and \mathbb{P} stands for a probability over (Ω, Σ) . The space of contingent claims, X , is modelled as an ideal (i.e., solid subspace) of $L_0(\Sigma)$ containing the constant functions, which represent investments in the riskless asset. A claim displays limited liabilities if it is positive. For a limited liability claim f , its option space is the collection of all portfolios of call and put options written on f , which can be identified as follows:

$$O_f = \text{Span} \{1, (f - k)^+ : k \in \mathbb{R}\}.$$

The space of all contingent claims written on f is identified as the space of all functions measurable with respect to $\sigma(f)$, the sub- σ -algebra

Date: April 5, 2024.

2010 Mathematics Subject Classification. Primary: 91G20, 91B25, Secondary: 46B42.

Key words and phrases. Spanning of options, market completeness, arbitrage, Kreps-Yan Theorem, order continuous dual.

The authors are partially supported by an NSERC grant.

generated by f , i.e.,

$$L_0(\sigma(f)).$$

A stream of research has been devoted to the study of spanning power of options on f , i.e., the size of O_f . In the seminal paper [30], Ross showed that if the state space Ω is finite then the options on f span the space of contingent claims written on f , i.e.,

$$O_f = L_0(\sigma(f)),$$

and if, in addition, f is one-to-one, then the option space of f completes the market, i.e.,

$$O_f = L_0(\Sigma).$$

These elegant results of Ross have inspired many successive contributions to the study of options. See e.g. [7, 20, 22] for related results on finite state spaces. In particular, they have also been examined for financial markets with infinite state spaces

Nachman proved in [24] that if $X = L_p(\Sigma)$ ($1 \leq p < \infty$), then the options on f span the space of contingent claims written on the asset in two ways: approximating by a.e. convergence or by p -th mean convergence. Precisely, it was proved that an asset $x \in L_p(\Sigma)$ is a contingent claim on $f \in L_p(\Sigma)$ iff there exists a sequence of portfolios of options on f converging a.e. to x iff there exists a sequence of portfolios of options on f converging in the p -th mean to x . That is,

$$\overline{O_f}^{a.e.} \cap L_p(\Sigma) = \overline{O_f}^{\|\cdot\|_p} = L_0(\sigma(f)) \cap L_p(\Sigma). \quad (1)$$

Galvani ([13]) and Galvani and Troitsky ([14]) proved further that if Ω is a Polish space equipped with the Borel σ -algebra and f is one-to-one and bounded, then O_f completes the market $X = L_p(\Sigma)$ ($1 \leq p \leq \infty$). That is, for $1 \leq p < \infty$,

$$\overline{O_f}^{a.e.} \cap L_p(\Sigma) = \overline{O_f}^{\|\cdot\|_p} = L_p(\Sigma), \quad (2)$$

and

$$\overline{O_f}^{a.e.} \cap L_\infty(\Sigma) = \overline{O_f}^{w^*} = L_\infty(\Sigma). \quad (3)$$

In this paper, we explore the spanning power of options in general spaces of contingent claims. Our contributions here are two-fold. Firstly, the spaces of contingent claims in our setting can be modelled as any ideal of $L_0(\Sigma)$ which contains the constant functions and admits a strictly positive order continuous linear functional. This framework includes not only the L_p -space ($1 \leq p \leq \infty$) models, but also the much wider class of Orlicz space models as well as many non-integrable space models which have been extensively used in the theory of risk measures (see e.g. [5, 6, 9, 11, 16, 25]).

Secondly, we provide an approach to unify the norm and w^* -topologies used in the results of Nachman, Galvani and Troitsky, and thus give more comprehensive insight into the general structures of option spaces; see Theorem 3.1 and Remark 3.3. The unification in our approach is due to the use of the topology $\sigma(X, X_n^\sim)$, where X_n^\sim is the set of all order continuous linear functionals on X .

Observe that $(X_n^\sim)_+$ has a natural connection with linear pricing functionals. Recall that a linear pricing functional ϕ on X is given by a state-price density $y \geq 0$ via integration, i.e.

$$\phi(x) = \int_{\Omega} xy \, d\mathbb{P} \text{ for all } x \in X,$$

where y is measurable and satisfies $\int_{\Omega} |xy| \, d\mathbb{P} < \infty$ for all $x \in X$. By Dominated Convergence Theorem, it is easily seen that ϕ is order continuous on X . Conversely, by Radon-Nikodym theorem, one can easily see that each positive order continuous linear functional on X has a positive density, and thus is a linear pricing functional. Therefore, $(X_n^\sim)_+$ is just the collection of linear pricing functionals on X .

Because of this link, we are able to apply Theorem 3.1 and shed light on the following general problem, raised in [8]: “Under what circumstances can prices on the marketed assets or basic derivative assets be uniquely extended by arbitrage to prices on all derivative assets in a large class and when is such an extension unique?” In Theorem 3.4, we prove that when the arbitrage condition is understood

as a no-free lunch condition (see [21]), one can extend uniquely the prices on O_f to the marketed contingent claims written on f .

Finally, we mention that there is a stream of works studying market completion using options in a continuous time setting. In this framework, the model is said to be complete, if any contingent claim payoff can be obtained as the terminal value of a self-financing trading strategy. We refer the reader to the recent papers [10, 18, 26, 29, 32].

2. PRELIMINARY RESULTS

We refer to [3, 4] for all unexplained terminology and standard facts on vector and Banach lattices. A vector subspace Y of a vector lattice X is called a **sublattice** if $|y| \in Y$ whenever $y \in Y$; in this case, $y_1 \wedge y_2$ and $y_1 \vee y_2$ both belong to Y whenever $y_1, y_2 \in Y$. A subspace Y is called an **ideal** (or a **solid subspace**) of X , if $|x| \leq |y|$ and $y \in Y$ imply $x \in Y$. A linear functional ϕ on a vector lattice X is said to be **order continuous** if $\phi(x_\alpha) \rightarrow 0$ whenever $x_\alpha \xrightarrow{o} 0$ in X . The collection of all order continuous linear functionals on X is denoted by X_n^\sim and is called the **order continuous dual** of X . A linear functional ϕ on X is said to be **positive** if $\phi(x) \geq 0$ whenever $x \geq 0$, and is said to be **strictly positive** if $\phi(x) > 0$ whenever $x > 0$.

The following lemma will be used. Recall first that a vector lattice is said to be **order complete** (or **Dedekind complete**) if every order bounded above subset has a supremum, and is said to have the **countable sup property** if any subset having a supremum possesses a countable subset with the same supremum. A subset A of a vector lattice X is said to be **order closed** if $x \in A$ whenever there exists a net (a_α) in A such that $a_\alpha \xrightarrow{o} x$ in X .

Lemma 2.1. *Let X be an order complete vector lattice with the countable sup property and Y be a sublattice of X . Then Y is order closed in X iff for any increasing sequence in Y which is order bounded above in X , its supremum in X also lies in Y .*

Given a probability space $(\Omega, \Sigma, \mathbb{P})$, denote by $L_0(\Sigma)$ the space of all real-valued measurable functions (*modulo a.e. equality*). We use 1 to denote the constant one function. Recall that $L_0(\Sigma)$ is a vector lattice, endowed with the natural order: $f \leq g$ iff $f(\omega) \leq g(\omega)$ for a.e. $\omega \in \Omega$. By [23, Lemma 2.6.1], it is easily seen that any ideal of $L_0(\Sigma)$ is order complete and has the countable sup property. Hence, Lemma 2.1 is applicable to them. Recall also that $f_n \xrightarrow{o} 0$ in an ideal X of $L_0(\Sigma)$ if and only if $f_n \xrightarrow{a.e.} 0$ and $(f_n)_{n=1}^\infty$ is *order bounded* in X , i.e., there exists $f \in X$ such that $|f_n| \leq f$ a.e. for each $n \geq 1$. We remark that the class of ideals of $L_0(\Sigma)$ which admit strictly positive order continuous linear functionals is very large. For example, by [15, Proposition 5.19], all Banach function spaces (i.e., ideals of $L_0(\Sigma)$ endowed with complete lattice norm), including all Orlicz spaces, are as such.

For a subset Y of $L_0(\Sigma)$, define $\sigma(Y)$ to be the smallest sub- σ -algebra of Σ which makes all members in Y measurable and contains all \mathbb{P} -null sets. Denote by $L_0(\sigma(Y))$ the set of all functions in $L_0(\Sigma)$ which are measurable with respect to $\sigma(Y)$. Clearly, $Y \subset L_0(\sigma(Y))$. If $Y = \{f\}$, we write $\sigma(f)$ instead of $\sigma(\{f\})$, for the sake of simplicity. The following result is an improved and generalized market completeness theorem in the sense of Green and Jarrow ([17, Theorem 1]).

Lemma 2.2. *Let X be an ideal of $L_0(\Sigma)$ and Y be a sublattice of X such that $1 \in Y$. Then the following are equivalent:*

- (a) Y is order closed in X ,
- (b) $Y = L_0(\sigma(Y)) \cap X$.

3. MAIN RESULTS

In this section, the space of contingent claims, X , is always modelled as an ideal of $L_0(\Sigma)$ over a given probability space $(\Omega, \Sigma, \mathbb{P})$ that contains the constant functions and admits a strictly positive order continuous linear functional. Our first main result is as follows.

Theorem 3.1. *Let f be a limited liability claim in X . For a claim $g \in X$, the following are equivalent:*

- (a) *g is a contingent claim written on f , i.e., $g \in L_0(\sigma(f)) \cap X$,*
- (b) *g can be approximated by portfolios of options on f in the $\sigma(X, X_n^\sim)$ -topology, i.e., $g \in \overline{O_f}^{\sigma(X, X_n^\sim)}$,*
- (c) *There exists a sequence (g_n) in O_f such that $g_n \xrightarrow{a.e.} g$.*

The following corollary is immediate.

Corollary 3.2. *Let f be a limited liability claim in X such that $\sigma(f) = \Sigma$. Then we have the following:*

- (a) *The option space of f completes the market in the $\sigma(X, X_n^\sim)$ -topology, i.e., $\overline{O_f}^{\sigma(X, X_n^\sim)} = X$,*
- (b) *The option space of f completes the market by approximating via a.e. convergence, i.e., for any $g \in X$, there exists a sequence (g_n) in O_f such that $g_n \xrightarrow{a.e.} g$.*

Remark 3.3. Note that our Theorem 3.1 and Corollary 3.2 imply both the aforementioned results of Nachman, Galvani and Troitsky. Indeed, recall first that, for $1 \leq p < \infty$, $X = L_p(\Sigma)$ is **order continuous**, i.e., $x_\alpha \downarrow 0$ in X implies $\|x_\alpha\| \downarrow 0$. In this case, one has $X_n^\sim = X^*$, so that $\sigma(X, X_n^\sim)$ is just the weak topology on X . Thus, by Mazur's theorem, $\overline{C}^{\sigma(X, X_n^\sim)} = \overline{C}^w = \overline{C}^{\|\cdot\|}$ for any convex subset C of X . Consequently, it follows that

$$\overline{O_f}^{\sigma(X, X_n^\sim)} = \overline{O_f}^{\|\cdot\|_p}.$$

Now it is clear that Equation (1) follows from Theorem 3.1. If, in addition, Ω is a Polish space with Σ being the Borel algebra, and f is one-to-one, then it is easily seen that $\sigma(f) = \Sigma$ by [4, Theorem 12.29]. Thus, Equation (2) follows from Corollary 3.2. Equation (3) also follows from Corollary 3.2, since $L_\infty(\Sigma)_n^\sim = L_1(\Sigma)$ and thus $\sigma(L_\infty(\Sigma), L_\infty(\Sigma)_n^\sim)$ is just the w^* -topology.

We now turn to discuss the pricing problem. Our notation and terminology are in accordance with [21, 31].

Let f be a fixed asset in X and π be a positive linear functional on the option space $M := O_f$, which is interpreted as a linear pricing functional on M . We denote by $M_0 := \{x \in M \mid \pi(x) = 0\}$ the set of all portfolios of options on f that can be bought or sold with zero price. We say that (M, π) admits **no free lunches** (cf. [31, Definition 1.3]), if the following holds

$$C \cap X_+ = \{0\}, \text{ where } C = \overline{M_0 - X_+}^{\sigma(X, X_n^\sim)}.$$

We say that a price, p , of an asset $g \in X$ is **consistent** with (M, π) if there exists a strictly positive functional $x^* \in X_n^\sim$ such that $x^*|_M = \pi$ and $x^*(g) = p$ ([21, Definition, pp. 29]). The price of $g \in X$ is said to be **determined by arbitrage from** (M, π) if there is a single price p for g that is consistent with (M, π) ([21, Definition, pp. 30]).

Theorem 3.4. *Suppose that the space X of contingent claims is a Banach function space in $L_0(\Sigma)$. Let f be a limited liability asset in X and π be a positive linear functional on the option space $M = O_f$. If (M, π) admits no free lunches, then the price of any contingent claim $g \in L_0(\sigma(f)) \cap X$ is determined by arbitrage from (M, π) .*

The proof of this result essentially depends on the following version of the Kreps-Yan Theorem, which is of independent interest.

Proposition 3.5. *Let X be a Banach function space in $L_0(\Sigma)$. Then the Kreps-Yan theorem holds true for $(X, \sigma(X, X_n^\sim))$. That is, for each $\sigma(X, X_n^\sim)$ -closed cone C in X such that $C \supset -X_+$ and $C \cap X_+ = \{0\}$, there exists a strictly positive functional $\phi \in X_n^\sim$ such that $\phi|_C \leq 0$.*

The proof of this result (see Section 4) relies on [19, Theorem 3.1]. For more results in this direction, we refer the reader to [27, 28]. For no-arbitrage results, we refer the reader to the monograph [12] and the references therein.

4. PROOFS OF RESULTS

Proof of Lemma 2.1. Let (y_n) be an increasing sequence in Y that is order bounded above in X . Since X is order complete, it follows that (y_n) has a supremum, x , in X . Since (y_n) is increasing, it follows that $y_n \uparrow x$ in X , so that $y_n \xrightarrow{o} x$ in X . Thus, if Y is order closed in X , then $x \in Y$. This proves the “only if” part.

For the “if” part, observe first that, in this case, for any sequence (y_n) in Y which is *order bounded* in X , its supremum and infimum in X also lie in Y . Indeed, denote by x the supremum of (y_n) in X . Put $z_n = \bigvee_{k=1}^n y_k$. Then $z_n \in Y$ as Y is a sublattice of X , and moreover, the supremum of (z_n) in X is still x . Since (z_n) is increasing, it follows from the “if” assumption that $x \in Y$. Replacing (y_n) with $(-y_n)$, one sees easily that the infimum of (y_n) in X also lies in Y . Now let $(y_\alpha) \subset Y$ and $x \in X$ be such that $y_\alpha \xrightarrow{o} x$ in X . By passing to a tail, we may assume that (y_α) is order bounded in X . Then since X is order complete, we have

$$\inf_{\alpha} \sup_{\beta \geq \alpha} |y_\beta - x| = 0,$$

where all the suprema and infima are taken in X . By the countable sup property of X , we can choose $\{\alpha_n\}_{n=1}^\infty$ such that $\inf_n \sup_{\beta \geq \alpha_n} |y_\beta - x| = 0$. Without loss of generality, we can assume that (α_n) is increasing. It follows that

$$\inf_n \sup_{m \geq n} |y_{\alpha_m} - x| = 0,$$

or equivalently, $y_{\alpha_n} \xrightarrow{o} x$, so that $x = \inf_n \sup_{m \geq n} y_{\alpha_m}$; cf. [4, Theorem 8.16]. Applying the preceding observation to the suprema and then to the infimum, we obtain that $x \in Y$. \square

Proof of Lemma 2.2. Assume first (b) holds. Let (f_n) be an increasing sequence in Y and f be its supremum in X . Then $f_n \uparrow f$ a.e. Since each f_n is $\sigma(Y)$ -measurable, we have that f is also $\sigma(Y)$ -measurable, so that $f \in L_0(\sigma(Y)) \cap X = Y$. Thus since X is order complete and has the countable sup property, Lemma 2.1 implies that (a) holds.

Conversely, assume that (a) holds. We first claim that $\sigma(Y) = \{A \in \Sigma : \chi_A \in Y\}^1$. Denote the right hand side by \mathcal{G} . We first show that it is a σ -algebra. Indeed, it is clear that $\emptyset \in \mathcal{G}$, and that if $A \in \mathcal{G}$, then $\chi_{A^c} = 1 - \chi_A \in Y$, so that $A^c \in \mathcal{G}$. Now let $(A_k)_1^\infty$ be a sequence of sets in \mathcal{G} . Then $\chi_{\cup_{k=1}^n A_k} = \bigvee_{k=1}^n \chi_{A_k} \in Y$, and from $\chi_{\cup_{k=1}^n A_k} \uparrow \chi_{\cup_{k=1}^\infty A_k}$ in X , it follows that $\chi_{\cup_{k=1}^\infty A_k} \in Y$, since Y is order closed. Therefore, $\bigcup_{k=1}^\infty A_k \in \mathcal{G}$. This concludes the proof of that \mathcal{G} is a σ -algebra. Next, we show that each $f \in Y$ is measurable with respect to \mathcal{G} . Indeed, for any real number r , it follows from $Y \ni n(f - r)^+ \wedge 1 \uparrow \chi_{\{f > r\}}$ in X that $\chi_{\{f > r\}} \in Y$, so that $\{f > r\} \in \mathcal{G}$, and f is \mathcal{G} -measurable. Clearly, \mathcal{G} contains all \mathbb{P} -null sets². Thus we have $\sigma(Y) \subset \mathcal{G}$. The reverse inclusion being clear, this completes the proof of the claim.

It is clear that $Y \subset L_0(\sigma(Y)) \cap X$. Now take $f \in L_0(\sigma(Y)) \cap X$. By considering f^\pm , we may assume that f is non-negative. Then we can find a sequence (f_n) of simple functions which are measurable with respect to $\sigma(Y)$ such that $f_n \uparrow f$ everywhere, so that $f_n \uparrow f$ in X . By the preceding claim, we have that $f_n \in Y$. Therefore, $f \in Y$, and thus $L_0(\sigma(Y)) \cap X \subset Y$. It follows that $Y = L_0(\sigma(Y)) \cap X$. \square

Proof of Theorem 3.1. We claim that O_f is a sublattice of X . Indeed, put $Z = \text{Span}\{b, s, (s - kb)^+ : k \in \mathbb{R}\}$, where $b = f + 1$ and $s = f$. Note that, being an ideal of $L_0(\Sigma)$, X is order complete, and thus it is uniformly complete (cf. [2, Lemma 1.56]). By [8, Theorem (1)], it follows that Z is a sublattice of X . Now simply observe that $Z = O_f$. Indeed, the inclusion $O_f \subset Z$ is immediate as $f, 1 \in Z$ and Z is closed under lattice operations. For the reverse inclusion, note that $s = f = (f - 0)^+ \in O_f$ so that $b \in O_f$ as well. Also, for $k \geq 1$ we have $(s - kb)^+ = 0$, and for $k < 1$ we have $(s - kb)^+ = (1 - k)(f - \frac{k}{1 - k}1)^+ \in O_f$.

Assume that (c) holds. Since Z is a sublattice of X , by considering the positive and negative parts, respectively, we may assume that $g \geq 0$ and $g_n \geq 0$ for all n . For any $k \geq 1$, since $g_k \wedge g_n \xrightarrow{a.e.} g_k \wedge g$ and

¹This is essentially contained in [17, Theorem 1].

²Keep in mind that χ_A is identified as 0 in $L_0(\Sigma)$ if $\mathbb{P}(A) = 0$.

$(g_k \wedge g_n)_n$ is order bounded in X , it follows that $g_k \wedge g_n \xrightarrow{o} g_k \wedge g$ in X , and therefore, $g_k \wedge g_n \xrightarrow{\sigma(X, X_n^\sim)} g_k \wedge g$ as $n \rightarrow \infty$. By the fact that Z is a sublattice again, we have $g_k \wedge g_n \in Z$ for all $k, n \geq 1$. Hence,

$$g_k \wedge g \in \overline{Z}^{\sigma(X, X_n^\sim)}$$

for any $k \geq 1$. Now $g_k \wedge g \xrightarrow{a.e.} g$ and $(g_k \wedge g)$ is order bounded in X , we have $g_k \wedge g \xrightarrow{o} g$ in X , so that $g_k \wedge g \xrightarrow{\sigma(X, X_n^\sim)} g$. Therefore, since $\overline{Z}^{\sigma(X, X_n^\sim)}$ is $\sigma(X, X_n^\sim)$ -closed, we have

$$g \in \overline{Z}^{\sigma(X, X_n^\sim)}.$$

This proves that (c) \Rightarrow (b).

Suppose now (b) holds. Recall that X_n^\sim is a band (i.e., order closed ideal) of the order dual X^\sim ([3, Theorem 1.57]). It follows from [3, Theorem 3.50] that the dual of X under the topology $|\sigma|(X, X_n^\sim)$ is just X_n^\sim . Therefore, by Mazur's theorem (cf. [3, Theorem 3.13]), since Z is convex, $\overline{Z}^{\sigma(X, X_n^\sim)} = \overline{Z}^{|\sigma|(X, X_n^\sim)}$. Consequently, there exists a net (g_α) in Z such that $g_\alpha \xrightarrow{|\sigma|(X, X_n^\sim)} g$. In particular, if x_0^* is any strictly positive order continuous functional on X , then

$$x_0^*(|g_\alpha - g|) \rightarrow 0.$$

Take (α_n) such that $x_0^*(|g_{\alpha_n} - g|) \leq \frac{1}{2^n}$. Then since $\bigvee_{m=n}^k |g_{\alpha_m} - g| \wedge 1 \uparrow_k \sup_{m \geq n} |g_{\alpha_m} - g| \wedge 1$, it follows from order continuity of x_0^* that

$$\begin{aligned} x_0^*\left(\sup_{m \geq n} |g_{\alpha_m} - g| \wedge 1\right) &= \lim_k x_0^*\left(\bigvee_{m=n}^k |g_{\alpha_m} - g| \wedge 1\right) \\ &\leq \lim_k x_0^*\left(\sum_{m=n}^k |g_{\alpha_m} - g| \wedge 1\right) \leq \frac{1}{2^{n-1}}. \end{aligned}$$

Therefore, we have

$$x_0^*\left(\inf_{n \geq 1} \sup_{m \geq n} |g_{\alpha_m} - g| \wedge 1\right) = 0,$$

and thus by strict positivity of x_0^* , we have

$$\inf_{n \geq 1} \sup_{m \geq n} |g_{\alpha_m} - g| \wedge 1 = 0.$$

If $g_{\alpha_n} \not\xrightarrow{a.e.} g$, then there exist $\varepsilon > 0$ and a measurable set A of positive measure such that $\limsup_n |g_n(\omega) - g(\omega)| \geq \varepsilon$ for any $\omega \in A$. Therefore, it is easily seen that

$$\inf_{n \geq 1} \sup_{m \geq n} |g_{\alpha_n} - g| \wedge 1 \geq (\varepsilon \chi_A) \wedge 1 > 0.$$

This contradiction concludes the proof of (b) \Rightarrow (c).

Now put $Y = \overline{Z}^{\sigma(X, X_n^\sim)}$. Then Y is clearly order closed in X . Moreover, by the preceding paragraph, $Y = \overline{Z}^{|\sigma|(X, X_n^\sim)}$, implying that it is also a sublattice of X by [3, Theorem 3.46]. Thus by Lemma 2.2, $Y = L_0(\sigma(Y)) \cap X$. Since $f \in Y$, it is clear that $\sigma(Y) \supset \sigma(f)$, so that

$$Y = L_0(\sigma(Y)) \cap X \supset L_0(\sigma(f)) \cap X.$$

For the reverse inclusion, note that, by definition of O_f , it is easily seen that each $g \in O_f$ is measurable with respect to $\sigma(f)$. Now for an arbitrary $g \in Y$, we can take, by the implication (b) \Rightarrow (c), a sequence (g_n) in O_f such that $g_n \xrightarrow{a.e.} g$. Clearly, g is also $\sigma(f)$ -measurable. Therefore, it follows that

$$Y \subset L_0(\sigma(f)) \cap X,$$

and hence $Y = L_0(\sigma(f)) \cap X$. This proves (a) \Leftrightarrow (b). \square

Proof of Proposition 3.5. We apply [19, Theorem 3.1] to $(X, \sigma(X, X_n^\sim))$, and verify that the following Assumptions (C) and (L) are satisfied.

Assumption (C): For every sequence (x_n^*) in X_n^\sim , there exist strictly positive numbers (α_n) such that $\sum_{n=1}^\infty \alpha_n x_n^*$ converges in X_n^\sim with respect to the $\sigma(X_n^\sim, X)$ -topology.

Assumption (L): Any family $\{x_\gamma^*\}_{\gamma \in \Gamma}$ in $(X_n^\sim)_+$ admits a countable subfamily $\{x_{\gamma_n}^*\}_{n \geq 1}$ such that, for any $x \in X_+$, $x_{\gamma_n}^*(x) = 0$ for all $n \geq 1$ implies $x_\gamma^*(x) = 0$ for all $\gamma \in \Gamma$.

We first verify that Assumption (C) is satisfied. Indeed, since X is a Banach lattice, we know that the order dual X^\sim equals the norm dual X^* ([3, Corollary 4.5]) and is thus a Banach lattice. By [3, Theorem 1.57], X_n^\sim is a band (i.e., order closed ideal) in $X^\sim = X^*$, and

is thus norm closed in X^* by [3, Theorem 3.46]. Now for a sequence (x_n^*) in X_n^\sim , put $\alpha_n = \frac{1}{2^n \|x_n^*\| + 1}$ for each $n \geq 1$. Then α_n 's are strictly positive, and $\sum_1^\infty \alpha_n x_n^*$ converges in norm to some $x^* \in X^*$. Since X_n^\sim is norm closed in X^* , it follows that $x^* \in X_n^\sim$. Clearly, $\sum_1^\infty \alpha_n x_n^*$ also converges to x^* in the $\sigma(X_n^\sim, X)$ -topology.

We now verify that Assumption (L) is also satisfied. For the given family $\{x_\gamma^*\}_{\gamma \in \Gamma}$ in $(X_n^\sim)_+$, put $N_\gamma := \{x \in X : x_\gamma^*(|x|) = 0\}$ and $C_\gamma := N_\gamma^d := \{x \in X : |x| \wedge |y| = 0 \text{ for all } y \in N_\gamma\}$ for each γ . Observe that N_γ is a band. Indeed, it is clearly an ideal. If a net (x_α) in N_γ converges in order to some $x \in X$, then $|x_\alpha - x| \xrightarrow{o} 0$ implies that $x_\gamma^*(|x|) = |x_\gamma^*(|x_\alpha|) - x_\gamma^*(|x|)| \leq x_\gamma^*(||x_\alpha| - |x||) \leq x_\gamma^*(|x_\alpha - x|) \rightarrow 0$, and consequently, $x_\gamma^*(|x|) = 0$, i.e., $x \in N_\gamma$. This yields the *band decomposition* $X = N_\gamma \oplus C_\gamma$ by [3, Theorem 1.42]. Recall from [1, Corollary 5.22] that X has a weak unit $u > 0$, i.e., any function $x \in X$ is supported in $\{\omega : u(\omega) > 0\}$ off a null set. Write $u = f_\gamma + e_\gamma$ where $f_\gamma \in N_\gamma$ and $e_\gamma \in C_\gamma$. Since $f_\gamma \wedge e_\gamma = 0$, it is easily seen that there exists $A_\gamma \in \Sigma$ such that $e_\gamma = u\chi_{A_\gamma}$ and $f_\gamma = u\chi_{A_\gamma^c}$. Each function in N_γ is disjoint with e_γ and is thus supported in A_γ^c off a null set; each function in C_γ is disjoint with f_γ and is thus supported in A_γ off a null set. By countable sup property of X , we choose $\{\gamma_n\}_{n=1}^\infty$ such that $\sup_n e_{\gamma_n} = \sup_\gamma e_\gamma$. If $x_{\gamma_n}^*(x) = 0$ for all $n \geq 1$ and some $x \in X_+$, then we have $x \in N_{\gamma_n}$, so that $x \wedge e_{\gamma_n} = 0$, for all $n \geq 1$. It follows that $x \wedge \sup_\gamma e_\gamma = x \wedge \sup_n e_{\gamma_n} = \sup_n (x \wedge e_{\gamma_n}) = 0$, and consequently, $x \wedge e_\gamma = 0$ for any γ . This implies that x is supported in A_γ^c off a null set and hence belongs to N_γ , i.e., $x_\gamma^*(x) = 0$. \square

Proof of Theorem 3.4. It is clear that $C := \overline{M_0 - X_+}^{\sigma(X, X_n^\sim)}$ is a $\sigma(X, X_n^\sim)$ -closed cone with $-X_+ \subset C$ and $C \cap X_+ = \{0\}$ because of no free lunches. Thus by Proposition 3.5, there exists a strictly positive linear functional $x^* \in X_n^\sim$ such that $x^*|_C \leq 0$. This last condition implies that $x^*|_{M_0} = 0$, so that $\ker \pi = M_0 \subset \ker(x^*|_M)$. Hence, there exists $\lambda > 0$ such that $\pi = \lambda x^*|_M$. Therefore, for each $g \in L_0(\sigma(f)) \cap X =$

$\overline{\mathcal{O}}_f^{\sigma(X, X_n^\sim)}$, it is easily seen that the price $p := \lambda x^*(g)$ is consistent with (M, π) . \square

Acknowledgements. The authors are grateful to the anonymous referee and editors for their many suggestions, which greatly improved the accessibility of the paper. The first author also thanks the Faculty of Science at Ryerson University for the hospitality received during his visit there.

REFERENCES

- [1] Y. Abramovich, C.D. Aliprantis, *An Invitation to Operator Theory*, Graduate Studies in Mathematics Vol. 50, American Mathematical Society, 2002.
- [2] C.D. Aliprantis, O. Burkinshaw, *Locally solid Riesz spaces with applications to economics*, Mathematical Surveys and Monographs No. 105, American Mathematical Society, 2003.
- [3] C.D. Aliprantis, O. Burkinshaw, *Positive Operators*, Springer 2006.
- [4] C.D. Aliprantis, K.C. Border, *Infinite dimensional analysis*, Springer, 3rd Edition, 2007.
- [5] S. Biagini, M. Frittelli, A unified framework for utility maximization problems: An Orlicz space approach, *Ann. Appl. Probab.* 18(3), 2008, 929–966.
- [6] S. Biagini, M. Frittelli, On the extension of the Namioka-Klee theorem and on the Fatou property for risk measures, In: *Optimality and Risk—Modern Trends in Mathematical Finance*, Springer Berlin Heidelberg, 2010, 1–28.
- [7] A.M. Baptista, Options and efficiency in multivariate security markets, *Math. Finance* 15(4), 2005, 569–587.
- [8] D.J. Brown, S.A. Ross, Spanning, valuation and options, *Economic Theory* 1(1), 1991, 3–12.
- [9] P. Cheridito, T. Li, Risk measures on Orlicz hearts, *Math. Finance* 19(2), 2009, 189–214.
- [10] M. Davis, J. Oblój, Market Completion using Options. In: *Advances in Mathematics of Finance*, Volume 83 of Banach Center Publications, 2002, 4960.
- [11] F. Delbaen, Risk measures for non-integrable random variables, *Math. Finance* 19(2), 2009, 329–333.
- [12] F. Delbaen, W. Schachermayer. The mathematics of arbitrage. Springer Science & Business Media, 2006.

- [13] V. Galvani, Option spanning with exogenous information structure, *Journal of Mathematical Economics* 45(1), 2009, 73–79.
- [14] V. Galvani, V.G Troitsky, Options and efficiency in spaces of bounded claims, *Journal of Mathematical Economics* 46(4), 2010, 616–619.
- [15] N. Gao, V. Troitsky, F. Xanthos, Unbounded order convergence and application to Cesáro means in Banach lattices, *Israel J. Math.*, to appear, arXiv:1509.07914.
- [16] N. Gao, F. Xanthos, On the C-property and w^* -representations of risk measures, *Math. Finance*, to appear, arXiv:1511.03159.
- [17] R. Green, R. Jarrow, Spanning and completeness in markets with contingent claims, *Journal of Economic Theory* 41, 1987, 202–210.
- [18] J. Hugonnier, S. Malamud, E. Trubowitz, Endogenous completeness of diffusion driven equilibrium markets, *Econometrica* 80(3), 2012, 1249–1270.
- [19] E. Jouini, C. Napp, W. Schachermayer, Arbitrage and state price deflators in a general intertemporal framework. *J. Math. Econom.* 41(6), 2005, 722–734.
- [20] C. Kountzakis, I.A Polyrakis, F. Xanthos, Nonreplication of options, *Math. Finance* 22(3), 2012, 569–584.
- [21] D.M. Kreps, Arbitrage and equilibrium in economies with infinitely many commodities, *Journal of Mathematical Economics* 8(1), 1981, 15–35.
- [22] I.A Polyrakis, F. Xanthos, Maximal submarkets that replicate any option, *Annals of Finance* 7(3), 2011, 407–423.
- [23] P. Meyer-Nieberg, *Banach lattices*, Universitext, Springer-Verlag, Berlin, 1991.
- [24] D.C. Nachman, Spanning and completeness with options, *Review of Financial Studies* 1(3), 1988, 311–328.
- [25] J. Orihuela, M.R. Galán, Lebesgue property for convex risk measures on Orlicz spaces, *Mathematics and Financial Economics* 6(1), 2012, 15–35.
- [26] F. Riedel, F. Herzberg, Existence of financial equilibria in continuous time with potentially complete markets, *Journal of Mathematical Economics* 49(5), 2013, 398–404.
- [27] D. B. Rokhlin, The Kreps-Yan theorem for L_∞ , *Int. J. Math. Math. Sci.* 17, 2005, 2749–2756.
- [28] D. B. Rokhlin, The Kreps-Yan theorem for Banach ideal spaces. (Russian) *Sibirsk. Mat. Zh.* 50 (2009), no. 1, 199–204; translation in *Sib. Math. J.* 50 (2009), no. 1, 162–166.
- [29] M. Romano, N. Touzi, Contingent Claims and Market Completeness in a Stochastic Volatility Model, *Math. Finance* 7(4), 1997, 399–412.

- [30] S.A Ross, Options and efficiency, *The Quarterly Journal of Economics* 90(1), 1976, 75-89.
- [31] W. Schachermayer, No arbitrage: on the work of David Kreps: Arbitrage and equilibrium in economies with infinitely many commodities, *Positivity* 6(3), 2002, 359–368.
- [32] D. C. Schwarz, Market Completion with Derivative Securities, preprint, arXiv:1506.00188.

SCHOOL OF MATHEMATICS, SOUTHWEST JIAOTONG UNIVERSITY, CHENGDU, SICHUAN, 610000, CHINA.

E-mail address: ngao@home.swjtu.edu.cn

DEPARTMENT OF MATHEMATICS, RYERSON UNIVERSITY, 350 VICTORIA ST., TORONTO, ON, M5B 2K3, CANADA.

E-mail address: foivos@ryerson.ca