

Nondeterministic quantum communication complexity: the cyclic equality game and iterated matrix multiplication

Harry Buhrman¹, Matthias Christandl², Jeroen Zuiddam¹

January 7, 2022

Abstract

We study nondeterministic multiparty quantum communication with a quantum generalization of broadcasts. We show that, with number-in-hand classical inputs, the communication complexity of a Boolean function in this communication model equals the logarithm of the *support rank* of the corresponding tensor, whereas the approximation complexity in this model equals the logarithm of the *border support rank*. This characterisation allows us to prove a log-rank conjecture posed by Villagra et al. for nondeterministic multiparty quantum communication with message passing.

The support rank characterization of the communication model connects quantum communication complexity intimately to the theory of asymptotic entanglement transformation and algebraic complexity theory. In this context, we introduce the *graphwise equality problem*. For a cycle graph, the complexity of this communication problem is closely related to the complexity of the computational problem of multiplying matrices, or more precisely, it equals the logarithm of the support rank of the iterated matrix multiplication tensor. We employ Strassen's laser method to show that asymptotically there exist nontrivial protocols for every odd-player cyclic equality problem. We exhibit an efficient protocol for the 5-player problem for small inputs, and we show how Young flattenings yield nontrivial complexity lower bounds.

1. Introduction

Let $f : X \times Y \times Z \rightarrow \{0,1\}$ be a function on a product of finite sets X , Y and Z . Alice, Bob and Charlie have to compute f in the following sense. Alice receives an $x \in X$, Bob receives a $y \in Y$ and Charlie receives a $z \in Z$, and each player receives a private random bit string. Then the players communicate in rounds. Each round, one player communicates by broadcasting a bit to the other players. After these rounds of communication, each player has to output a bit, such that if $f(x,y,z) = 1$, then with some nonzero probability all players output 1 and if $f(x,y,z) = 0$, then with probability zero all players output 1. The complexity of such a protocol is the number of broadcasts in the protocol, and we denote the minimum complexity of all such protocols by $N(f)$.

¹QuSoft, CWI Amsterdam and University of Amsterdam, Science Park 123, 1098 XG Amsterdam, Netherlands. Email: buhrman@cwi.nl, j.zuiddam@cwi.nl

²Department of Mathematical Sciences, University of Copenhagen, Universitetsparken 5, 2100 Copenhagen Ø, Denmark. Email: christandl@math.ku.dk

Now we allow the players to be quantum, as follows. Alice receives an $x \in X$, Bob receives a $y \in Y$ and Charlie receives a $z \in Z$. Then, in rounds, the players communicate by creating a GHZ-like state

$$|\text{GHZ}\rangle = \alpha |000\rangle + \beta |111\rangle$$

and sharing this state among each other, a quantum broadcast. Moreover, the players can do any local quantum computations. Again, after these rounds of communication, each player has to output a bit, such that if $f(x, y, z) = 1$, then with some nonzero probability all players output 1 and if $f(x, y, z) = 0$, then with probability zero all players output 1. The quantum complexity of such a quantum protocol is the number of broadcasts in the protocol, and we denote the minimum complexity of all quantum protocols by $\text{NQ}(f)$. We will make this definition more precise and more general in Section 2. Note that the quantum model can simulate the classical model. Also note that, nondeterministically, one quantum broadcast can be used to send a qubit from one player to another by using teleportation (see Theorem 9); the quantum model can thus simulate a message-passing model.

Our results.

- Our main technical result is that the quantum complexity of a function in the above model equals the logarithm of the so-called *support rank* of the tensor $\sum_{x,y,z} f(x, y, z) |x\rangle|y\rangle|z\rangle$ corresponding to f . We prove this in Section 2.
- Modifying the quantum model such that the players can only communicate by message passing — that is, in each communication round one player sends a qubit to one other player — increases the complexity by at most a factor $k - 1$, and this relationship is tight. However, asymptotically in the input size, the increase is only $k/2$ and this relationship is tight. This solves a *nondeterministic multiplayer quantum log-rank conjecture* in the message-passing model of Villagra et al. [VNYN13]. This topic is covered in Section 3.
- We define the k -player *graphwise equality* problem to be the problem in which k players are identified with vertices in a graph G , and each player has to compute the equality function with his neighbours in G . Of particular interest is the cycle graph $G = C_k$ and the corresponding *cyclic equality* problem. For this cyclic equality problem, in the classical broadcast model, the naïve protocol in which every player broadcasts his inputs is the optimal protocol. The same holds in the quantum model when k is even. Interestingly, we show with Strassen’s laser method that for all odd $k \geq 3$ there is a nontrivial quantum protocol. Moreover, for all odd $k \geq 3$ we give nontrivial lower bounds on the value of NQ by use of Young flattenings. These results are related to the complexity of matrix multiplication and iterated matrix multiplication. A consequence of our work is that finding new protocols for the cyclic equality problem for three players yields new algorithms for matrix multiplication. Section 4 covers the classical case, the even quantum case, an explicit quantum protocol for $k = 5$, and the Young flattening lower bound. Section 5 covers the Strassen laser method.

Related work. The two-player nondeterministic quantum communication model was introduced by De Wolf [Wol03]. He shows that the communication complexity in this

model is characterized by the logarithm of the support rank of the communication matrix. Quantum broadcast channels have been studied by e.g. Ambainis et al. [ABDR04]. Multiparty nondeterministic quantum communication with message passing has been studied by Villagra et al. [VNYN13]. They show that the logarithm of the support rank of the communication tensor is a lower bound for the message-passing complexity and conjecture that this lower bound is polynomially related to the message-passing complexity.

The support rank of 3-tensors has been studied by Cohn and Umans in the context of the complexity of matrix multiplication [CU13]. They give nontrivial upper bounds on the support rank of the matrix multiplication tensor that do not come from upper bounds on the tensor rank. As an interesting fact, we note that given a matrix A and a number k , deciding whether the support rank of A is at least k is NP-hard [BK15].

The complexity of matrix multiplication plays a central role in algebraic complexity theory. We refer to [BCS97] for general background information. Connections between algebraic complexity theory and entanglement transformations have been studied before, see for example [CDS08]. The iterated matrix multiplication tensor has been studied in the context of arithmetic circuit complexity and the VP versus VNP problem, see for example [Ges15]. To the knowledge of the authors, the tensor rank or support rank of the iterated matrix multiplication tensor has not been studied before.

Acknowledgements. We thank Peter Bürgisser, Péter Vrana, Florian Speelman and Teresa Piovesan for helpful discussions. Part of this work was done while MC and JZ were visiting the Simons Institute for the Theory of Computing, UC Berkeley. HB was partially funded by the European Commission, through the SIQS project and by the Netherlands Organisation for Scientific Research (NWO) through gravitation grant Networks. MC acknowledges financial support from the European Research Council (ERC Grant Agreement no 337603), the Danish Council for Independent Research (Sapere Aude) and the Swiss National Science Foundation (project no PP00P2_150734). Part of this work was done while MC was with ETH Zurich. JZ is supported by NWO through the research programme 617.023.116 and by the European Commission through the SIQS project.

2. Support rank characterization of the quantum broadcast model

We refer to Nielsen and Chuang [NC10] for background information on the quantum computation model.

Quantum multiparty communication protocol. We will give two definitions of a quantum broadcast model, which are equivalent in the nondeterministic setting. The first model clearly generalizes the classical broadcast model, while the second model is easier to analyse. For any natural number m , denote by $[m]$ the set $\{1, 2, \dots, m\}$. Let k be a positive integer and let f be a Boolean function on $[2^n]^k = [2^n] \times [2^n] \times \dots \times [2^n]$,

$$f : [2^n]^k \rightarrow \{0, 1\}.$$

We define a k -player quantum communication protocol as follows. Each player i has a local Hilbert space H_i with a register initialised in the input state $|x_i\rangle$. The players have access to a quantum broadcast channel, which, given a qubit state $\alpha|0\rangle + \beta|1\rangle$, will create the state $\alpha|0\rangle^{\otimes k} + \beta|1\rangle^{\otimes k}$ and distribute this state among the k players.

The players proceed in communication rounds; each round a designated player uses the broadcast channel. Let R_i be the first qubit of H_i and let $R = R_1 \otimes \cdots \otimes R_k$. After the communication is finished, we apply a projection onto $|11 \cdots 1\rangle$ in R . If the resulting tensor is 0 then the output of the protocol is 0, otherwise the output of the protocol is 1. The complexity of the protocol is the number of communication rounds. We say the protocol *nondeterministically computes* f if the probability that the output equals 1 is nonzero if $f(x_1, \dots, x_k) = 1$ and the probability that the output equals 0 is one if $f(x_1, \dots, x_k) = 0$.

We will now give an equivalent definition of the quantum broadcast model. This is the definition that we will use in the rest of the paper. Each player i has a finite-dimensional Hilbert space H_i . The protocol thus takes place in the space $H_1 \otimes \cdots \otimes H_k$. The space is initialised in the state $|x_1 \cdots x_k\rangle |\text{GHZ}_r^k\rangle$, where

$$|\text{GHZ}_r^k\rangle := \sum_{a=1}^r |a\rangle |a\rangle \cdots |a\rangle \in (\mathbb{C}^r)^{\otimes k}$$

is the k -party GHZ-state of rank r , shared among the k players, and $x_i \in [2^n]$ is the classical input to player i . (For clarity we will suppress any normalizations in quantum states when possible.) The players now apply local quantum operations. Let R_i be the first qubit of H_i and let $R = R_1 \otimes \cdots \otimes R_k$. We apply a projection onto $|11 \cdots 1\rangle$ in R . If the resulting tensor is 0 then the output of the protocol is 0, otherwise the output of the protocol is 1. The *complexity* of the protocol is $\log_2(r)$. We say the protocol *nondeterministically computes* f if the probability that the output equals 1 is nonzero if $f(x_1, \dots, x_k) = 1$ and the probability that the output equals 0 is one if $f(x_1, \dots, x_k) = 0$.

Definition 1. Let k be a positive integer and let f be a function $[2^n]^k \rightarrow \{0, 1\}$. The *k -player nondeterministic quantum communication complexity of f* is the minimal complexity of a k -player quantum communication protocol that nondeterministically computes f , and is denoted by $\text{NQ}(f)$.

Approximating protocols. Let f be a function $[2^n]^k \rightarrow \{0, 1\}$. Let $(\Pi_j)_{j \in \mathbb{N}}$ be a sequence of protocols, such that when $f(x_1, \dots, x_k) = 1$, the probability that Π_j outputs 1 on input x converges to a nonzero number as j goes to infinity, and when $f(x_1, \dots, x_k) = 0$, the probability that Π_j outputs 0 on input x converges to 1 as j goes to infinity. Then we say that the sequence $(\Pi_j)_{j \in \mathbb{N}}$ *approximately nondeterministically computes* f . The complexity of an approximating sequence is the maximum complexity of any protocol Π_j in the sequence.

Definition 2. The *k -player approximate nondeterministic quantum communication complexity of f* is the minimal complexity of a sequence (Π_j) that approximately nondeterministically computes f , and is denoted by $\underline{\text{NQ}}(f)$.

Classical protocol. We define a *k -player classical communication protocol* as follows. Each player receives a classical input and a private random bit string. The protocol proceeds in rounds. Each round we let a single predetermined player communicate by broadcasting a bit to all the other players. After the last communication round, every player presents an output bit. If all the output bits are 1, then the output of the protocol is 1; otherwise the output of the protocol is 0. Again, we say the classical protocol *nondeterministically computes* f if the probability that the output equals 1

is nonzero if $f(x_1, \dots, x_k) = 1$ and the probability that the output equals 0 is one if $f(x_1, \dots, x_k) = 0$.

Definition 3. The k -player *nondeterministic classical communication complexity* of f is the minimal complexity of a k -player classical communication protocol that nondeterministically computes f , and is denoted by $N(f)$.

Remark 4. For simplicity, we have taken the input set for each of the k players to be the same set $[2^n]$. We note that the definitions in this section and most of the results in this paper naturally generalize to the situation where the players get inputs from sets of different sizes.

Support rank and border support rank. Let t be a tensor in $(\mathbf{C}^m)^{\otimes k}$. The *tensor rank* of t is the smallest number r such that t can be written as a sum of r simple tensors, that is, $t = \sum_{i=1}^r u_i^1 \otimes u_i^2 \otimes \dots \otimes u_i^k$ for some vectors $u_i^j \in \mathbf{C}^m$. We denote the tensor rank of t by $R(t)$. Fix a basis for $(\mathbf{C}^m)^{\otimes k}$ and define the support of a tensor t in $(\mathbf{C}^m)^{\otimes k}$ to be the set of basis element that occur with nonzero coefficient in t . The *support rank* or *nondeterministic rank* of t is the smallest number r such that there exists a tensor in the space $(\mathbf{C}^m)^{\otimes k}$ with the same support as t and tensor rank r . We denote the support rank of t by $R_s(t)$. Note that support rank is basis *dependent*.

The *border rank* of t is the smallest number r such that there exists a sequence of tensors $(t_j)_{j \in \mathbf{N}}$ converging to t in the Euclidean topology (or equivalently in the Zariski topology) such that $R(t_j)$ is at most r for every j . We denote the border rank of t by $\underline{R}(t)$. The *border support rank* of t is the smallest number r such that there exists a tensor in $(\mathbf{C}^m)^{\otimes k}$ with the same support as t and border rank r . We denote the border support rank of t by $\underline{R}_s(t)$.

Theorem 5. Let $f : [2^n]^k \rightarrow \{0, 1\}$ be a function and let t be the tensor in $(\mathbf{C}^{2^n})^{\otimes k}$ with entries given by f , that is, $t = \sum_{i \in [2^n]^k} f(i) |i_1\rangle |i_2\rangle \dots |i_k\rangle$. Then $\text{NQ}(f) = \log_2 R_s(t)$ and $\underline{\text{NQ}}(f) = \log_2 \underline{R}_s(t)$.

Lemma 6 (Cleanup lemma). Let $\{|\psi_i\rangle : i \in [q]\} \subseteq (\mathbf{C}^m)^{\otimes k}$ be a set of k -tensors, for some natural number q . Then there exists a k -partite rank-1 linear map $\langle \ell| := \langle \ell_1| \otimes \dots \otimes \langle \ell_k|$ with $\langle \ell_j| \in (\mathbf{C}^m)^*$ such that $\langle \ell|\psi_i\rangle \neq 0$ for every $i \in [q]$.

Proof. We will give a proof by recursively constructing $\langle \ell|$. Let Id be the identity map on \mathbf{C}^m . If $j \leq k$, $\langle a| \in ((\mathbf{C}^m)^*)^{\otimes j}$ and $|b\rangle \in (\mathbf{C}^m)^{\otimes k}$, then we denote by $\langle a|b\rangle$ the contraction of $\langle a|$ and $|b\rangle$, that is, $\langle a|b\rangle = (\langle a| \otimes \text{Id}^{\otimes k-j}) |b\rangle$.

The base case is $\langle \ell| = 1$. For the recursion, suppose we are given an element $\langle \ell'| \in ((\mathbf{C}^m)^*)^{\otimes j}$ such that $\langle \ell'|\phi_i\rangle := \langle \ell|\psi_i\rangle$ is nonzero for every $i \in [q]$. We will construct an element $\langle \ell| \in ((\mathbf{C}^m)^*)^{\otimes j+1}$ such that $\langle \ell|\psi_i\rangle$ is nonzero for every $i \in [q]$. Since $\langle \ell'|\phi_i\rangle$ is nonzero for every $i \in [q]$, there is an element $\langle u_i| \in (\mathbf{C}^m)^*$ such that $\langle u_i|\phi_i\rangle$ is nonzero. Consider the the maps $(\langle u_1| + x \langle u_2|) |\phi_i\rangle$ for $i \in \{1, 2\}$, in the variable x . Each map only has a single root. Therefore, there exists a value α_2 for x such that both maps evaluate to a nonzero number. Next, consider the maps $(\langle u_1| + \alpha_2 \langle u_2| + x \langle u_3|) |\phi_i\rangle$ for $i \in \{1, 2, 3\}$, in variable x . Again, each of the three maps has only a single root. Therefore, there exists a value α_3 for x such that all three maps evaluate to a nonzero number. Repeat this construction to obtain an element $\langle u| \in (\mathbf{C}^m)^*$ such that $\langle u|\phi_i\rangle$ is nonzero for every $i \in [q]$. Let $\langle \ell|$ be $\langle \ell'| \otimes \langle u|$. \square

Proof of Theorem 5. We first show $\text{NQ}(f) \leq \log_2 R_s(t)$. Let r be the support rank of t . Then there exists a unit vector $\psi \in (\mathbf{C}^{2^n})^{\otimes k}$ with rank r and support equal to the support of f . This means that there are vectors $|u_i^j\rangle \in \mathbf{C}^{2^n}$ such that $\psi = \sum_{i=1}^r |u_i^1\rangle \cdots |u_i^k\rangle$. For every player j define a matrix

$$A_j := \alpha_j \sum_{i=1}^r |u_i^j\rangle\langle i|$$

where α_j is a nonzero complex number such that $A_j^\dagger A_j$ has eigenvalue at most 1. The matrix $I - A_j^\dagger A_j$ is thus positive semidefinite and hence there exists a matrix A'_j such that $A_j'^\dagger A'_j = I - A_j^\dagger A_j$. Define for every player j a quantum operation

$$\mathcal{E}_j : \rho \mapsto A_j \rho A_j^\dagger \otimes |1\rangle\langle 1| + A'_j \rho A_j'^\dagger \otimes |0\rangle\langle 0|.$$

Note that this operation introduces a new control qubit register which player j can measure to see whether he applied A_j or A'_j .

The protocol for the k players is as follows. Let x_1, \dots, x_k be the inputs given to the players. The players share a k -party GHZ-state of rank r . Player j applies \mathcal{E}_j to his part of the GHZ-state. If his control qubit is $|0\rangle$ then he sets his output qubit R_i to $|0\rangle$. Otherwise, he measures the rest of the system. If the outcome equals $|x_j\rangle$, then he sets R_j to $|1\rangle$, otherwise he sets R_j to $|0\rangle$.

The above protocol uses a GHZ-state of rank r , so it has complexity $\log_2(r)$. We claim that the protocol nondeterministically computes f . If the players in the first measurement each get outcome $|1\rangle$, then the state of the total system is $|\psi\rangle$. Because $|\psi\rangle$ has norm 1, this happens with nonzero probability $|\alpha_1|^2 \cdots |\alpha_k|^2$. If $f(x_1, \dots, x_k) = 0$, then $|x_1 \cdots x_k\rangle$ does not occur in the support of ψ , so the probability that the players measure $|x_1\rangle, \dots, |x_k\rangle$ respectively is zero. Hence in this case the register R is not in state $|11 \cdots 1\rangle$. On the other hand, if $f(x_1, \dots, x_k) \neq 0$, then $|x_1 \cdots x_k\rangle$ does occur in the support of ψ , so the probability that the players measure $|x_1\rangle, \dots, |x_k\rangle$ respectively is nonzero. Hence with nonzero probability the register R is in state $|11 \cdots 1\rangle$.

We now show $\text{NQ}(f) \geq \log_2 R_s(t)$. Suppose we have a protocol that nondeterministically computes f with complexity r . This means that the players perform *local* quantum operations that together form a linear map L which transforms, for any $x_1, \dots, x_k \in [2^n]$, the state

$$|x_1 \cdots x_k\rangle |\text{GHZ}_r\rangle$$

to a state of the form

$$|x_1 \cdots x_k\rangle \sum_{a \in A} |\psi_x^a\rangle |a_1\rangle |a_2\rangle \cdots |a_k\rangle,$$

where the sum is over $A := \{a \in \{0, 1\}^k \mid f(x_1, \dots, x_k) = a_1 \cdot a_2 \cdots a_k\}$ and where $|\psi_x^a\rangle$ is some nonzero vector, representing the state of the *work space* of the players. Since the map L is *linear*, it maps the tensor

$$s_1 := \sum_{x_1, \dots, x_k} |x_1 \cdots x_k\rangle |\text{GHZ}_r\rangle$$

to the tensor

$$s_2 := \sum_{x_1, \dots, x_k} |x_1 \cdots x_k\rangle \sum_{a \in S} |\psi_x^a\rangle |a_1 \cdots a_k\rangle.$$

The tensor rank of $\sum_x |x_1 \cdots x_k\rangle$ is 1 and hence the tensor rank of s_1 is r . Because L is a local map, the tensor rank of s_2 is at most r . By applying the cleanup lemma Lemma 6 and projecting on states with $|a_1 \cdots a_k\rangle = |1 \cdots 1\rangle$, we obtain a tensor

$$s_3 := \sum_{x_1, \dots, x_k} |x_1 \cdots x_k\rangle c_x$$

where $c_x \in \mathbf{C}$ is zero if $f(x) = 0$ and nonzero if $f(x) = 1$. The rank of the tensor s_3 is at most r . The support of s_3 equals the support of f , so the support rank of f is at most r .

The statement about the approximate complexity of f follows from the definition of border support rank. \square

Remark 7. We note that having a NQ-protocol for f of complexity n is the same as having an SLOCC protocol for transforming $\text{GHZ}_{2^n}^k$ to a tensor with the same support as f . We will use the SLOCC paradigm in some parts of the text.

3. Nondeterministic log-rank conjecture for message-passing protocols

Definition 8. Let $\text{NQ}_0(f)$ be the minimal complexity of a protocol that nondeterministically computes f , without preshared entanglement but with the added ability for players to send a qubit to another player. The complexity of such a protocol is the total number of qubits sent.

Villagra et al. [VNYN13] show that $\text{NQ}_0(f)$ is at least the logarithm of the support rank of f . They furthermore conjecture that $\text{NQ}_0(f)$ is upper bounded by a polynomial in the logarithm of the support rank. The following theorem proves this conjecture.

Theorem 9 (“Nondeterministic log-rank conjecture”). *Let $f : [2^n]^k \rightarrow \{0, 1\}$. Then we have $\text{NQ}(f) \leq \text{NQ}_0(f) \leq (k - 1) \text{NQ}(f)$.*

Proof. For the first inequality, suppose we have an NQ_0 -protocol for f . We replace the communication of a qubit by the nondeterministic teleportation of that qubit. Beforehand, all players agree on the basis in which the teleportation should happen. If any teleportation during the protocol does not happen in this basis, then the player that notices this sets his output register R_i to $|0\rangle$.

For the second inequality, suppose we have an NQ-protocol for f which uses a GHZ-state of rank r . Then we can construct a NQ_0 -protocol for f as follows. The players start with no shared entanglement. Player 1 constructs a GHZ-state of rank r locally. In the first $k - 1$ communication rounds, player 1 distributes the GHZ-state over the other $k - 1$ players. After that, the players perform the NQ-protocol. The resulting NQ_0 -protocol has complexity at most $(k - 1) \text{NQ}(f)$. \square

To say something about the ‘tightness’ of Theorem 9 we consider the natural easy function in the NQ-model, namely $f(x_1, \dots, x_k) = [x_1 = x_2 = \cdots = x_k]$ with $x_i \in [2^n]$.

Proposition 10 (Single bit inputs). *Let $f : [2]^k \rightarrow \{0, 1\}$ be the function defined by $f(x_1, \dots, x_k) = [x_1 = x_2 = \cdots = x_k]$ for $x_i \in [2]$. Then we have $\text{NQ}(f) = 1$ and $\text{NQ}_0(f) = (k - 1) \text{NQ}(f)$.*

Proof. Note that the tensor of this function is GHZ_2^k , so $\text{NQ}(f) = 1$. Now consider a protocol that nondeterministically computes f without preshared entanglement and r rounds of communication. We may assume, without loss of generality, that the protocol consists of a first phase in which the players communicate and a second phase in which the players only do local quantum operations. After the first phase the players are sharing some state E consisting of EPR-pairs shared among certain pairs of the players. We thus obtain a local linear map which maps $\sum_x |x\rangle E$ to a tensor with the same support as GHZ_2^k . However, if $r < k - 1$, then, viewing E as a graph, E is disconnected. Therefore there is a grouping of the players into two groups such that there are no EPR-pairs between the groups. Such a state cannot be converted to a GHZ_2^k state by SLOCC. \square

Asymptotically, we can improve the relationship stated in Theorem 9, as follows.

Theorem 11 (Asymptotic upper bound). *For any $\varepsilon > 0$, there is an n_0 such that for all $f : [m]^k \rightarrow \{0, 1\}$, if $\text{NQ}(f) > n_0$, then*

$$\text{NQ}_0(f) \leq \frac{(k + \varepsilon)}{2} \text{NQ}(f).$$

To prove Theorem 11 we use the theory of asymptotic SLOCC conversion rates.

Definition 12. Given tensors $\psi \in V_1 \otimes \cdots \otimes V_k$ and $\phi \in W_1 \otimes \cdots \otimes W_k$, we say that ψ can be transformed into ϕ via SLOCC operations, if there exist linear transformations $A_i : V_i \rightarrow K_i$ such that $\phi = (A_1 \otimes \cdots \otimes A_k)\psi$; and we write $\psi \xrightarrow{\text{SLOCC}} \phi$. Define

$$\omega_n(\psi, \phi) = \frac{1}{n} \inf\{m \in \mathbb{N}_{\geq 1} \mid \psi^{\otimes m} \xrightarrow{\text{SLOCC}} \phi^{\otimes n}\}$$

and

$$\omega(\psi, \phi) = \lim_{n \rightarrow \infty} \omega_n(\psi, \phi).$$

Lemma 13. *The limit $\omega(\psi, \phi)$ exists and for all n the inequality $\omega_n(\psi, \phi) \geq \omega(\psi, \phi)$ holds; in other words, $\omega_n = \omega + o(1)$.*

Theorem 14 (Vrana-Christandl [VC16]). *Let $\text{GHZ}_2^{K_k}$ be the k -party tensor consisting of EPR-pairs between any parties. Then*

$$\omega(\text{GHZ}_2^{K_k}, \text{GHZ}_2^k) = \frac{1}{k-1}.$$

In other words, for any $\varepsilon > 0$, there is an n_0 such that for all $n > n_0$,

$$(\text{GHZ}_2^{K_k})^{\otimes n(\frac{1}{k-1} + \varepsilon)} \xrightarrow{\text{SLOCC}} (\text{GHZ}_2^k)^{\otimes n}.$$

Proof of Theorem 11. Creating $\text{GHZ}_2^{K_k}$ in the NQ_0 -model costs $\binom{k}{2}$ messages. Asymptotically, we can transform $1/(k-1)$ copies of $\text{GHZ}_2^{K_k}$ to one copy of GHZ_2^k by SLOCC. More precisely, by Theorem 14, for any $\varepsilon > 0$, there is an n_0 such that for all $n > n_0$,

$$(\text{GHZ}_2^{K_k})^{\otimes \frac{n}{k-1} + \varepsilon n} \xrightarrow{\text{SLOCC}} (\text{GHZ}_2^k)^{\otimes n}.$$

We conclude that, for any $\varepsilon > 0$, there is an n_0 such that for all $n > n_0$, $\binom{k}{2}(\frac{n}{k-1} + \varepsilon n) = ((k + \varepsilon')n)/2$ messages are sufficient to generate $(\text{GHZ}_2^k)^{\otimes n}$ by SLOCC.

To prove the theorem, suppose we have an NQ -protocol for f which uses a GHZ state of rank 2^n and no communication. Consider the following NQ_0 -protocol for f . Create a GHZ -state of rank 2^n by sending $\frac{(k + \varepsilon')n}{2}$ messages and then continue with the NQ -protocol. \square

The following proposition says that the asymptotic relationship of Theorem 11 is tight.

Proposition 15 (*n*-bit inputs). *Let $f : [2^n]^k \rightarrow \{0, 1\}$ be the function defined by $f(x_1, \dots, x_k) = [x_1 = x_2 = \dots = x_k]$ for $x_i \in [2^n]$. Then we have $\text{NQ}(f) = n$ and $\text{NQ}_0(f) \geq \frac{k}{2} \text{NQ}(f)$.*

Proof. As in the previous proof, note that the tensor corresponding to f is $\text{GHZ}_{2^n}^k$. Suppose there is an NQ_0 protocol using r messages. View the communication pattern of this protocol as an undirected multigraph G (i.e. parallel edges are allowed) on k vertices. Note that G has r edges. Let $E = \text{GHZ}_2^G$ be the tensor that has an EPR pair at every edge in G . The protocol yields an SLOCC transformation of E to $\text{GHZ}_{2^n}^k$. Let ℓ be the minimal number of edges across any cut of G . Then ℓ is at most the minimal degree d of G . The sum of all degrees in G equals $2r$, so $k\ell \leq kd \leq 2r$, which implies the inequality $r \geq k\ell/2$. The number ℓ is equal to $\min_{S \subseteq [k]} \log_2 \text{rk}_S(E)$, where $\text{rk}_S(E)$ denotes the rank of E after flattening according to the set S . This value cannot increase under any SLOCC transformation. Now note that $\log_2 \text{rk}_{\{i\}}(\text{GHZ}_{2^n}^k) = n$ for any $i \in [k]$, so $\ell \geq n$. We conclude that $r \geq kn/2$. \square

Remark 16. Another way to prove Proposition 15 is to first symmetrize the protocol to obtain an SLOCC transformation of a state E with $\log_2 \text{rk}_{\{i\}}(E) = (k-1)!2r$ to the state $\text{GHZ}_{2^{k!n}}^k$. We have $\log_2 \text{rk}_{\{i\}}(\text{GHZ}_{2^{k!n}}^k) = k!n$. Since $\log_2 \text{rk}_{\{i\}}$ is an SLOCC-monotone, we obtain the inequality $(k-1)!2r \geq k!n$ and hence $r \geq kn/2$.

4. Cyclic equality problem

The two-player equality problem EQ_n is the problem of Alice and Bob having to decide whether their n -bit inputs are equal. Since the identity matrix has full support rank, we have $\text{NQ}(\text{EQ}_n) = n$. We generalize EQ_n to multiple players as follows. Let G be an undirected graph. Let EQ_n^G be the problem of $|G|$ players having to solve the n -bit equality problem between players connected by edges. (Note that this definition naturally generalizes to hypergraphs.) If G is a bipartite graph, one easily sees that by grouping the players we can transform the problem into an equality problem on en bits EQ_{en} , where e is the number of edges in the graph. Therefore $\text{NQ}(\text{EQ}_n^G) = en$, that is, the trivial protocol is optimal for bipartite graphs. On the other hand, if G contains an odd cycle, then this argument fails. In the rest of this paper we will focus on the extreme case of G being an odd cycle and investigate the complexity of the corresponding equality problem.

Definition 17. The k -player *cyclic equality problem* on n bits $\text{EQ}_n^{C_k}$ is the function

$$\text{EQ}_n^{C_k} : ([2^n] \times [2^n])^k \rightarrow \{0, 1\} : (a_1 b_1, \dots, a_k b_k) \mapsto \begin{cases} 1 & \text{if } b_1 = a_2, b_2 = a_3, \dots, b_k = a_1 \\ 0 & \text{otherwise,} \end{cases}$$

that is, the players are arranged in a circle; player i receives two n -bit inputs a_i, b_i and has to decide whether $a_i = b_{i-1}$ and $b_i = a_{i+1}$, where the indices are taken modulo k .

It turns out that the tensor corresponding to this function is a generalisation of the *matrix multiplication tensor*, one of the central objects of study in algebraic complexity

theory. This tensor arises as follows in algebraic complexity theory. Consider the bilinear map

$$\mathbf{C}^{m \times m} \times \mathbf{C}^{m \times m} \rightarrow \mathbf{C}^{m \times m} : (A, B) \mapsto AB$$

which multiplies two complex $m \times m$ matrices. Any bilinear map $U \times V \rightarrow W$ corresponds canonically to a tensor in $U \otimes V \otimes W$. The number of multiplications in the field \mathbf{C} necessary to perform the bilinear map is equal to the tensor rank of the corresponding tensor, up to a factor 2. The tensor corresponding to the matrix multiplication map is

$$\langle m, m, m \rangle := \sum_{x \in [m]^3} |x_1 x_2\rangle |x_2 x_3\rangle |x_3 x_1\rangle.$$

A natural generalisation of the tensor $\langle m, m, m \rangle$ to a k -party tensor is the so-called *iterated matrix multiplication* tensor

$$\text{IMM}_m^k := \sum_{x \in [m]^k} |x_1 x_2\rangle |x_2 x_3\rangle \cdots |x_k x_1\rangle.$$

Clearly, $\text{IMM}_m^3 = \langle m, m, m \rangle$. The tensor IMM_m^k corresponds to the multilinear map

$$(\mathbf{C}^{m \times m})^{\times k} \rightarrow \mathbf{C} : (A_1, A_2, \dots, A_k) \mapsto \text{tr}(A_1 A_2 \cdots A_k)$$

which computes the trace of the product of k matrices. We note that, when viewed as a polynomial in the matrix entries, IMM_m^k plays a special role in the field of arithmetic circuits and geometric complexity theory. Namely, IMM_3^k is complete for the class VP_e of families of polynomials computable by small formulas [BOC92], and IMM_k^k is complete for the class VQP, for which the determinant is also complete [Blä01]. The following connection between iterated matrix multiplication and cyclic equality is readily observed.

Proposition 18. *The tensor corresponding to the cyclic equality function $\text{EQ}_n^{C_k}$ on n bits is the iterated matrix multiplication tensor $\text{IMM}_{2^n}^k$ with $2^n \times 2^n$ matrices. Therefore, we have the equalities $\text{NQ}(\text{EQ}_n^{C_k}) = \log_2 \text{R}_s(\text{IMM}_{2^n}^k)$ and $\underline{\text{NQ}}(\text{EQ}_n^{C_k}) = \log_2 \underline{\text{R}}_s(\text{IMM}_{2^n}^k)$*

The remainder of this paper is organized as follows. In the following four paragraphs we do the following: (1) we show that in the classical model, the naïve protocol in which every player broadcasts his input is optimal; (2) we show that when k is even the naïve protocol is optimal quantumly; (3) we exhibit nontrivial protocols when $n = 1$ and $k = 3$ or $k = 5$; (4) we show nontrivial lower bounds on the quantum complexity by use of Young flattenings. Finally, in the last section, we show that the Strassen laser method yields nontrivial protocols for all odd $k \geq 3$, asymptotically.

Classical lower bound with the fooling set method. We will show that in the classical situation the trivial protocol is always optimal. To prove a lower bound on the classical complexity of the cyclic equality problem we use the fooling set method.

Theorem 19. *The classical nondeterministic communication complexity $\text{N}(\text{EQ}_n^{C_k})$ of the cyclic equality problem equals kn .*

Proof. Let $S \subseteq [2^{2n}]^k$ be the set of 1-inputs of the function $\text{EQ}_n^{C_k}$. This set has size 2^{kn} . Let Π be a classical protocol for $\text{EQ}_n^{C_k}$ and denote by $\Pi_r(x_1, \dots, x_k)$ the sequence of messages sent by the players in the protocol Π on input $x \in [2^{2n}]^k$ and private randomness $r \in [m]^k$. Suppose there are distinct 1-inputs $x, y \in S$ and private randomnesses

$r, s \in [m]^k$ such that $\Pi_r(x_1, \dots, x_k) = \Pi_s(y_1, \dots, y_k)$. There is an i such that $x_i \neq y_i$, say $i = 1$. We have $\Pi_r(x_1, \dots, x_k) = \Pi_{(r_1, s_2, \dots, s_k)}(x_1, y_2, \dots, y_k)$, so the protocol outputs 1 on input x_1, y_2, \dots, y_k with randomness (r_1, s_2, \dots, s_k) . However, x_1, y_2, \dots, y_k is a 0-input, a contradiction. Therefore, $\Pi_r(x_1, \dots, x_k) \neq \Pi_s(y_1, \dots, y_k)$. We conclude that $N(\text{EQ}_n^{C_k}) \geq \log_2(|S|)$. \square

An even number of quantum players. When k is even, the cycle graph C_k is bipartite, and, as mentioned above, the best protocol for an equality problem on a bipartite graph is the trivial protocol. We record this statement in terms of border support rank in the following proposition.

Proposition 20. *For even k , $m^k \leq \underline{R}_s(\text{IMM}_m^k)$. As a consequence, we have the equalities $\underline{NQ}(\text{EQ}_n^{C_k}) = NQ(\text{EQ}_n^{C_k}) = kn$.*

Proof. Let t be a tensor with the same support as $\text{IMM}_m^k \in (\mathbf{C}^{m^2})^{\otimes k}$. Label the players with the numbers $1, 2, \dots, k$. Group the *even* players together and group the *odd* players together and flatten the tensor t accordingly into a matrix A in $(\mathbf{C}^{m^2})^{\otimes k/2} \otimes (\mathbf{C}^{m^2})^{\otimes k/2}$. The matrix A has the same support as the identity matrix in $(\mathbf{C}^{m^2})^{\otimes k/2} \otimes (\mathbf{C}^{m^2})^{\otimes k/2}$ and thus has rank m^k . \square

Note that for odd k the above proof yields the lower bound $m^{k-1} \leq \underline{R}_s(\text{IMM}_m^k)$. We will show in Theorem 22 that this lower bound is not tight.

Nontrivial 3-player and 5-player quantum protocols. In the 3-player situation, Strassen's celebrated decomposition of the tensor $\text{IMM}_2^3 = \langle 2, 2, 2 \rangle$ into a sum of 7 simple tensors [Str69] gives a nontrivial protocol for $\text{EQ}_1^{C_3}$, and thus $NQ(\text{EQ}_1^{C_3}) \leq \log_2(7)$. We show that for 5 players there also exists a nontrivial protocol for $\text{EQ}_1^{C_5}$, as follows. Recall that we have defined $\text{IMM}_2^5 = \sum_{i \in [2]^5} |i_1 i_2\rangle |i_2 i_3\rangle |i_3 i_4\rangle |i_4 i_5\rangle |i_5 i_1\rangle$. Observe that an upper bound $R(\text{IMM}_2^5) \leq r$ implies $R(\text{IMM}_n^5) \leq \mathcal{O}(n^{\log_2(r)})$ by taking tensor powers of IMM_2^5 .

Theorem 21. $R(\text{IMM}_2^5) \leq 31$, and thus $NQ(\text{EQ}_1^{C_5}) \leq \log_2(31)$.

Proof. Let $|-\rangle := |1\rangle - |2\rangle$, $|+\rangle := |1\rangle + |2\rangle$ and $|\Phi^+\rangle = |11\rangle + |22\rangle$. Let $\text{Cyc}_5 := \sum_{\sigma \in C_5} \sigma$ be the cyclic symmetrizer acting on $(\mathbf{C}^4)^{\otimes 5}$ by permuting the 5 parties, and moreover let $\text{Sym}_2 := \sum_{\sigma \in S_2} \sigma$ be a 'local symmetrizer' acting diagonally on $(\mathbf{C}^2)^{\otimes 10}$ by permuting the basis states $|1\rangle$ and $|2\rangle$ of each \mathbf{C}^2 . Let

$$\begin{aligned} t := & - | -1 \rangle | 11 \rangle | 11 \rangle | 1+ \rangle | 22 \rangle \\ & - | -1 \rangle | 12 \rangle | 21 \rangle | 1+ \rangle | 22 \rangle \\ & - | \Phi^+ \rangle | 22 \rangle | -1 \rangle | 1+ \rangle | 22 \rangle. \end{aligned}$$

By direct computation, we see that $\text{IMM}_2^5 = \text{Cyc}_5(\text{Sym}_2(t)) + |\Phi^+\rangle^{\otimes 5}$. We observe that the right hand side yields a sum of 31 simple tensors. \square

We have a proof generalizing Theorem 21 to $R(\text{IMM}_2^k) \leq 2^k - 1$ for all odd k , which will appear in a forthcoming paper.

Quantum lower bound with Young flattenings. Let $t \in V_1 \otimes V_2 \otimes V_3$ be some 3-tensor. By grouping V_1 and V_2 , the tensor t can be viewed as a matrix $A \in (V_1 \otimes V_2) \otimes V_3$; this is called a *flattening*. The rank of the flattening A is a lower bound for the border rank of t and thus we obtain lower bounds on the border rank of tensors by computing the rank of their flattenings. However, this type of lower bound can never be bigger than the dimension of any local space V_i , and there do exist tensors with border rank larger than the local dimensions, for example the matrix multiplication tensor $\langle 2, 2, 2 \rangle$.

One approach to overcome this ‘local dimension limitation’ is as follows. We let $\phi : V_2 \rightarrow W_1 \otimes W_2$ be a linear map such that $R(\phi(v)) \leq e$ for all $v \in V_2$. By applying ϕ to the central tensor leg of t , we transform t into a 4-tensor $s \in V_1 \otimes W_1 \otimes W_2 \otimes V_3$. Next, we flatten s to a matrix $A \in (V_1 \otimes W_1) \otimes (W_2 \otimes V_3)$. The rank of A divided by e is a lower bound for the border rank of t . We will be using a specific linear map ϕ which originates from the representation theory of the general linear group. When one takes such representation theoretic maps ϕ to construct a flattening as above one speaks of a *Young flattening* [LO11]. An early appearance of this type of flattening can be recognized in the work of Strassen [Str83]. The following lower bound is obtained with a Young flattening.

Theorem 22. For odd $k \geq 3$, $(2n^2 - n)n^{k-3} \leq \underline{R}_s(\text{IMM}_n^k)$. As a consequence, we have the lower bound $(k-1)n + \log_2(2 - \frac{1}{n}) \leq \underline{\text{NQ}}(\text{EQ}_n^{C_k})$.

Proof. Let $k = 3$. The proof for odd $k > 3$ goes similarly after having grouped the k parties appropriately to 3 parties. For a vector space V , let $\wedge^a V$ be the a th exterior power of V . Define the linear map

$$\begin{aligned} \phi : \mathbf{C}^{2n-1} &\rightarrow \wedge^p \mathbf{C}^{2n-1} \otimes \wedge^{p+1} \mathbf{C}^{2n-1} \\ |j\rangle &\mapsto \sum_{j_1 < \dots < j_p} |j_1\rangle \wedge \dots \wedge |j_p\rangle \otimes |j_1\rangle \wedge \dots \wedge |j_p\rangle \wedge |j\rangle, \end{aligned}$$

and note that the rank of the matrix $\phi(v)$ equals $\binom{2n-2}{p}$ for any $v \in \mathbf{C}^{2n-1}$. We consider the tensor

$$t_1 := \sum_i \alpha_{i_1, i_2, i_3} |i_1 i_2\rangle |i_2 i_3\rangle |i_3 i_1\rangle \in \mathbf{C}^{n^2} \otimes \mathbf{C}^{n^2} \otimes \mathbf{C}^{n^2},$$

where i runs over $[n]^3$ and the α_{i_1, i_2, i_3} are nonzero complex numbers. The border rank of t_1 is at least the border rank of

$$t_2 := \sum_i \alpha_{i_1, i_2, i_3} |i_1 i_2\rangle |i_2 + i_3 - 1\rangle |i_3 i_1\rangle \in \mathbf{C}^{n^2} \otimes \mathbf{C}^{2n-1} \otimes \mathbf{C}^{n^2}.$$

Apply ϕ to the central tensor leg of t_2 and then flatten to obtain

$$A := \sum_i \sum_{j_1 < \dots < j_p} \alpha_{i_1, i_2, i_3} |i_1 i_2\rangle |j_1\rangle \wedge \dots \wedge |j_p\rangle \otimes |j_1\rangle \wedge \dots \wedge |j_p\rangle \wedge |i_2 + i_3 - 1\rangle |i_3 i_1\rangle.$$

View A as a direct sum of n matrices $A_{i_1} \in (\mathbf{C}^n \otimes \wedge^p \mathbf{C}^{2n-1}) \otimes (\wedge^{p+1} \mathbf{C}^{2n-1} \otimes \mathbf{C}^n)$; the matrix A_{i_1} corresponds to the linear map

$$f_{i_1} := |i_2\rangle |j_1\rangle \wedge \dots \wedge |j_p\rangle \mapsto \sum_{i_3} \alpha_{i_1, i_2, i_3} |j_1\rangle \wedge \dots \wedge |j_p\rangle \wedge |i_2 + i_3 - 1\rangle |i_3\rangle.$$


Let $p = n - 1$. We claim that every matrix A_{i_1} is upper triangular with elements α_{i_1, i_2, i_3} on the diagonal, up to permutations of the rows and columns. Assuming the claim is

true, we get that $R(A) = \sum_{i_1} R(A_{i_1}) = n \dim(\mathbf{C}^n \otimes \wedge^{n-1} \mathbf{C}^{2n-1}) = n^2 \binom{2n-1}{n-1}$. This implies the lower bound $\underline{R}_s(\text{IMM}_n^3) \geq n^2 \binom{2n-1}{n-1} / \binom{2n-2}{n-1} = 2n^2 - n$.

To prove this claim we define a partial order on the basis elements $|j_1\rangle \wedge \cdots \wedge |j_n\rangle \otimes |\ell\rangle$ of the target space of A_{i_1} . We will use the same partial order as Landsberg and Michałek [LM16]. Denote the basis elements of the target space by (P, ℓ) with P an n -subset of $[2n-1]$ and $\ell \in [n]$. Let (P_1, ℓ_1) and (P_2, ℓ_2) be two such basis elements and define $\ell := \min(\ell_1, \ell_2)$. We say $(P_1, \ell_1) < (P_2, \ell_2)$

1. if the ordered sequence of the ℓ smallest elements in P_2 is lexicographically smaller than the ordered sequence of the ℓ smallest elements in P_1 ;
2. or if the sequences of ℓ smallest elements are equal and $\ell_1 < \ell_2$.

One checks that this defines a partial order and that the unique minimal element in this order is $(\{n, \dots, 2n-1\}, 1)$. For example, with $n = 2$ the partial order has the following Hasse diagram.



We prove the claim by induction on $<$, with the minimal element as a base case. For now let all the α_{i_1, i_2, i_3} be 1. First, under A_{i_1} we have

$$|n\rangle \otimes |n+1\rangle \wedge \cdots \wedge |2n-1\rangle \mapsto |n\rangle \wedge \cdots \wedge |2n-1\rangle \otimes |1\rangle,$$

so the minimal element $(\{n, \dots, 2n-1\}, 1)$ is in the image of A_{i_1} . Let (P, ℓ) be in the target space of A_{i_1} and assume that every (P', ℓ') with $(P', \ell') < (P, \ell)$ is in the image. Write $P = (p_1, \dots, p_n)$ with $p_1 \leq \cdots \leq p_n$. Under A_{i_1} we have,

$$|p_1\rangle \wedge \cdots \wedge \widehat{p_\ell} \cdots \wedge |p_n\rangle \otimes |1 + p_\ell - \ell\rangle \mapsto \sum_m |p_1\rangle \wedge \cdots \wedge \widehat{p_\ell} \cdots \wedge |p_n\rangle \wedge |p_\ell - \ell + m\rangle \otimes |m\rangle.$$

Taking $m = \ell$, one sees that the basis element (P, ℓ) is present in the sum. Moreover, for any other (P', m) appearing in the sum we have $(P', m) < (P, \ell)$. Indeed, if $m > \ell$, then $p_\ell - \ell + m > p_\ell$, so the smallest ℓ elements in P' are lexicographically larger than the smallest ℓ elements in P , meaning $(P', m) < (P, \ell)$ by rule 1; if $m < \ell$, then $p_m \leq p_\ell - \ell + m < p_\ell$, so the m smallest elements of P' and P are equal, meaning $(P', m) < (P, \ell)$ by rule 2. Therefore, the basis element (P, ℓ) is in the image. This argument shows that A_{i_1} has full rank. Moreover, this argument shows that, up to a permutation of the rows and columns, the matrix A_{i_1} is upper triangular with ones on the diagonal. Repeating this argument with general values for α_{i_1, i_2, i_3} proves the claim. \square

Remark 23. The lower bound in Theorem 22 improves a lower bound of Ikenmeyer on the border support rank of IMM_n^3 [Ike13, 8.2.17].

5. Strassen's laser method for iterated matrix multiplication

In this section we show that $\text{NQ}(\text{EQ}_n^{C_k}) < kn$ for odd k . We will prove this result in the language of algebraic complexity theory.

Definition 24. Define $\omega_k := \inf\{\alpha \in \mathbf{R} \mid \text{R}(\text{IMM}_n^k) \in \mathcal{O}(n^\alpha)\}$. We call this the *exponent* of iterated matrix multiplication. Define $\omega_{s,k} := \inf\{\alpha \in \mathbf{R} \mid \text{R}_s(\text{IMM}_n^k) \in \mathcal{O}(n^\alpha)\}$. We call this the *support rank exponent* of iterated matrix multiplication.

Asymptotically, we have $\text{NQ}(\text{EQ}_n^{C_k}) \leq \omega_{s,k} n + \mathcal{O}(1) \leq \omega_k n + \mathcal{O}(1)$. The exponents ω_3 and $\omega_{s,3}$ are known as ω and ω_s in the literature. The support rank exponent of matrix multiplication was first studied by Cohn and Umans [CU13]. The best upper bound on ω_s comes from the upper bound $\omega \leq 2.3728639$ of Le Gall [LG14]. Interestingly, Cohn and Umans show the relationship

$$\omega \leq (3\omega_s - 2)/2.$$

Therefore, one way of finding upper bounds on ω is to construct an efficient quantum communication protocol for the cyclic equality problem EQ_n^3 .

For any k we have $k - 1 \leq \omega_k \leq k$, and if k is even, then $\omega_k = k$ (Proposition 20). The aim of this section will be to show: if $k \geq 3$ is odd, then

$$\omega_k < k.$$

Schönhage τ -theorem. In this section we will generalize some tools for obtaining upper bounds on the exponent of ω_3 to all exponents ω_k , in particular, we generalize the Schönhage τ -theorem. The proofs in this section are straightforward generalizations of the proofs for $k = 3$ which can be found in [Blä13]. In the next paragraph, we will use Strassen's laser method to show that $\omega_k < k$ for all odd k .

First we recall an important relationship between border rank and rank. We use the following more precise notion of border rank. Let $h \in \mathbf{N}$ and let t be a tensor in $\mathbf{C}^{\otimes m_1} \otimes \dots \otimes \mathbf{C}^{\otimes m_k}$. Define $R_h(t)$ to be the minimum number r such that there exist vectors $v_i^j \in (\mathbf{C}[\varepsilon])^{m_j}$ that satisfy $\sum_{i=1}^r v_i^1 \otimes \dots \otimes v_i^k = \varepsilon^h t + \mathcal{O}(\varepsilon^{h+1})$. A well-known but nontrivial result is that $\underline{R}(t) = \min_h R_h(t)$. It is not hard to show that $R_{h+h'}(t \otimes t') \leq R_h(t) R_{h'}(t')$. The relationship we are talking about is the following.

Proposition 25. *For every $h, k \in \mathbf{N}$, there is a number c_h such that for all tensors $t \in \mathbf{C}^{m_1} \otimes \dots \otimes \mathbf{C}^{m_k}$, $R(t) \leq c_h R_h(t)$. The number c_h depends polynomially on h .*

Proof. Let t be a tensor in $\mathbf{C}^{m_1} \otimes \dots \otimes \mathbf{C}^{m_k}$ with $R_h(t) = r$. Then there are $v_i^j \in (\mathbf{C}[\varepsilon])^{m_j}$ such that

$$\sum_{i=1}^r v_i^1 \otimes \dots \otimes v_i^k = \varepsilon^h t + \mathcal{O}(\varepsilon^{h+1}).$$

Decomposing every v_i^j into ε -homogeneous components $v_i^j = \sum_{a_j=0}^h \varepsilon^{a_j} v_i^j(a_j)$, and collecting powers of ε gives

$$\sum_{i=1}^r \sum_{a_1, \dots, a_k \in [h]} \varepsilon^{a_1 + \dots + a_k} v_i^1(a_1) \otimes \dots \otimes v_i^k(a_k) = \varepsilon^h t + \mathcal{O}(\varepsilon^{h+1}).$$

Taking only the summands such that $a_1 + \dots + a_k = h$ gives a rank decomposition of t . There are $\binom{h+k-1}{k-1} r$ such summands. \square

Next, we show that an upper bound on the border rank of ‘unbalanced’ iterated matrix multiplication tensors yields an upper bound on ω_k .

Proposition 26. *If $\underline{R}(\langle n_1, n_2, \dots, n_k \rangle) \leq r$, then $\omega_k \leq k \log_{n_1 \dots n_k} r$.*

Proof. Let $N = n_1 \dots n_k$. There is an h such that $R_h(\langle n_1, \dots, n_k \rangle) \leq r$. By taking the tensor product of all cyclic shifts of $\langle n_1, \dots, n_k \rangle$, we get $R_{kh}(\langle N, \dots, N \rangle) \leq r^k$ and thus $R_{khs}(\langle N^s, \dots, N^s \rangle) \leq r^{ks}$ for all s . Hence $\underline{R}(\langle N^s, \dots, N^s \rangle) \leq c_{khs} r^{ks}$ for some number c_{khs} which is constant in N . Therefore,

$$\omega \leq \log_{N^s}(c_{khs} r^{ks}) = ks \log_{N^s}(r) + \log_{N^s}(c_{khs}).$$

If s goes to infinity then $\log_{N^s}(c_{khs})$ goes to zero, so $\omega_k \leq k \log_N(r)$. \square

The real workhorse is the following straightforward generalization of a theorem of Schönhage [Sch81].

Proposition 27 (k -party Schönhage τ -theorem). *Suppose that $r > p$ and*

$$\underline{R}\left(\bigoplus_{i=1}^p \langle n_1^i, n_2^i, \dots, n_k^i \rangle\right) \leq r.$$

Define τ by $\sum_{i=1}^p (\prod_{j=1}^k n_j^i)^\tau = r$. Then $\omega_k \leq k\tau$

We follow the proof of [Blä13]. We first prove two lemmas. For tensors $s, t \in \mathbf{C}^{m_1} \otimes \dots \otimes \mathbf{C}^{m_k}$, let $s \leq t$ denote the existence of an SLOCC transformation mapping t to s . Let $a, b \in \mathbf{N} + 1$.

Lemma 28. *Let t be a tensor such that $\underline{R}(t^{\oplus a}) \leq b$. Then for all s , $\underline{R}((t^{\otimes s})^{\oplus a}) \leq \lceil b/a \rceil^s a$.*

Proof. We prove the lemma by induction over s . The base case $s = 1$ follows from the assumption. For the induction step, we have

$$(t^{\otimes s+1})^{\oplus a} = t^{\oplus a} \otimes t^{\otimes s} \leq \text{GHZ}_b \otimes t^{\otimes s} = (t^{\otimes s})^{\oplus b},$$

and thus, by the induction hypothesis,

$$\underline{R}((t^{\otimes s+1})^{\oplus a}) \leq \underline{R}((t^{\otimes s})^{\oplus b}) \leq \underline{R}((t^{\otimes s})^{\oplus (\lceil b/a \rceil a)}) \leq \lceil \frac{b}{a} \rceil \lceil \frac{b}{a} \rceil^s a \leq \lceil \frac{b}{a} \rceil^{s+1} a,$$

proving the lemma. \square

Lemma 29. *If $\underline{R}(\langle n_1, n_2, \dots, n_k \rangle^{\oplus a}) \leq b$, then $\omega \leq k \log_{n_1 \dots n_k} \lceil b/a \rceil$.*

Proof. The inequality $\underline{R}(\langle n_1, n_2, \dots, n_k \rangle^{\oplus a}) \leq b$ implies by Lemma 28 the inequality $\underline{R}(\langle n_1^s, n_2^s, \dots, n_k^s \rangle) \leq \underline{R}(\langle n_1^s, n_2^s, \dots, n_k^s \rangle^{\oplus a}) \leq \lceil b/a \rceil^s a$ which by Proposition 26 yields

$$\omega_k \leq k \frac{s \log \lceil \frac{b}{a} \rceil + \log(a)}{s \log(n_1 \dots n_k)},$$

which goes to $k \log \lceil b/a \rceil / \log(n_1 \dots n_k)$ when s goes to infinity. \square

Proof of Proposition 27. We assume $\underline{R}(\bigoplus_{i=1}^p \langle n_1^i, n_2^i, \dots, n_k^i \rangle) \leq r$. This implies that there is an $h \in \mathbb{N}$ such that $R_h(\bigoplus_{i=1}^p \langle n_1^i, n_2^i, \dots, n_k^i \rangle) \leq r$. Taking the s th tensor power gives $R_{hs}((\bigoplus_{i=1}^p \langle n_1^i, n_2^i, \dots, n_k^i \rangle)^{\otimes s}) \leq r^s$. We expand the tensor power to get

$$R_{hs} \left(\bigoplus_{\sigma} \left(\bigotimes_{i=1}^p \langle (n_1^i)^{\sigma_i}, (n_2^i)^{\sigma_i}, \dots, (n_k^i)^{\sigma_i} \rangle \right)^{\oplus \binom{s}{\sigma_1, \dots, \sigma_p}} \right) \leq r^s,$$

where the first direct sum is over all p -tuples σ of nonnegative integers with sum s . We can also write this inequality as

$$R_{hs} \left(\bigoplus_{\sigma} \langle \prod_i (n_1^i)^{\sigma_i}, \dots, \prod_i (n_k^i)^{\sigma_i} \rangle \right)^{\oplus \binom{s}{\sigma_1, \dots, \sigma_p}} \leq r^s.$$

There exists a number c_{hs} depending polynomially on h and s such that

$$R \left(\bigoplus_{\sigma} \langle \prod_i (n_1^i)^{\sigma_i}, \dots, \prod_i (n_k^i)^{\sigma_i} \rangle \right)^{\oplus \binom{s}{\sigma_1, \dots, \sigma_p}} \leq c_{hs} r^s.$$

Define τ by $\sum_{i=1}^p (\prod_{j=1}^k n_j^i)^{\tau} = r$. Then $\sum_{\sigma} \binom{s}{\sigma_1, \dots, \sigma_p} (\prod_i (n_1^i)^{\sigma_i} \dots \prod_i (n_k^i)^{\sigma_i})^{\tau} = r^s$. In this sum, consider the maximum summand and fix the corresponding σ . Define the numbers $n_j := \prod_i (n_j^i)^{\sigma_j}$. Let $a := \binom{s}{\sigma_1, \dots, \sigma_p}$ and $b := r^s c_{hs}$. We apply Lemma 29 to the inequality $R(\langle n_1', \dots, n_k' \rangle^{\oplus a}) \leq b$ to obtain

$$\omega_k \leq k\tau + \frac{(p-1)\log(s+1) + \log(c_{hs})}{\log(n_1 \dots n_k)},$$

which goes to $k\tau$ when s goes to infinity. (See [Blä13] for more details.) \square

Strassen's laser method. We will now use Strassen's laser method to prove the main result of this section.

Theorem 30. *For any odd k we have $\omega_k < k$.*

We will give a proof for the case $k = 5$, the other cases being similar. Define the 5-tensor $\text{Str}_q^5 = \sum_{i=1}^q |ii000\rangle + |0ii00\rangle$ in $\mathbf{C}^{q+1} \otimes \mathbf{C}^q \otimes \mathbf{C}^{q+1} \otimes \mathbf{C} \otimes \mathbf{C}$.

Proposition 31. $\underline{R}(\text{Str}_q^5) \leq q + 1$.

Proof. Expanding $\sum_{i=1}^q (|0\rangle + \varepsilon|i\rangle) |i\rangle (|0\rangle + \varepsilon|i\rangle) |0\rangle |0\rangle$ gives

$$\sum_{i=1}^q |0i000\rangle + \varepsilon \sum_{i=1}^q |ii000\rangle + |0ii00\rangle + \mathcal{O}(\varepsilon^2).$$

Subtracting $|0\rangle (\sum_{i=1}^q |i\rangle) |000\rangle$ yields $\varepsilon \text{Str}_q^5 + \mathcal{O}(\varepsilon^2)$. \square

Define the tensor $\langle n_1, n_2, n_3, n_4, n_5 \rangle$ to be

$$\sum_{x \in [n_1] \times \dots \times [n_5]} |x_1 x_2\rangle |x_2 x_3\rangle |x_3 x_4\rangle |x_4 x_5\rangle |x_5 x_1\rangle.$$

So $\text{IMM}_n^5 = \langle n, n, n, n, n \rangle$.

Proposition 32. $\text{GHZ}_2^5 \leq \langle 2, 2, 2, 2, 2 \rangle$.

Proof. Let ϕ be the map $|ab\rangle \mapsto \delta_{[a=b]} |a\rangle$. Apply $\phi^{\otimes 5}$ to $\langle 2, 2, 2, 2, 2 \rangle$. This yields one copy of $\text{GHZ}_2^{[5]}$. \square

Remark 33. We mention that the subrank result of Proposition 32 can be improved asymptotically in the sense that rate $\omega(\langle 2, 2, 2, 2, 2 \rangle, \text{GHZ}^5) = 1/2$ [VC16]. Using this fact in the proof of Theorem 30 gives the slightly better upper bound $\omega_k \leq \log_q((q+1)^k/4)$.

For the proof of Theorem 30 we have to define the notion of the decomposition of the support of a tensor and the corresponding inner and outer structure of a tensor. Let I_1, \dots, I_k be finite sets. A *decomposition* \mathcal{D} of $I_1 \times \dots \times I_k$ is a collection of sets I_i^j such that

$$I_i = \bigsqcup_j I_i^j,$$

meaning that for every i , $\cap_j I_i^j = \emptyset$ and $\cup_j I_i^j = I_i$. Let t be a tensor in $\mathbf{C}^{m_1} \otimes \dots \otimes \mathbf{C}^{m_k}$ and index the basis elements in this space by elements of $[m_1] \times \dots \times [m_k]$. Let \mathcal{D} be a decomposition of $[m_1] \times \dots \times [m_k]$. We view \mathcal{D} as a ‘cut’ of $[m_1] \times \dots \times [m_k]$ into smaller product sets and thus as a ‘cut’ of t into smaller tensors. We define $t|_{I_1^{j_1}, I_2^{j_2}, \dots, I_k^{j_k}}$ to be the restriction of t to the basis elements in $I_1^{j_1} \times I_2^{j_2} \times \dots \times I_k^{j_k}$. These smaller tensors we think of as the ‘inner structure’ of t . We define the ‘outer structure’ of t with respect to \mathcal{D} to be the tensor $t_{\mathcal{D}}$ indexed by sequences (j_1, \dots, j_k) such that $t_{\mathcal{D}}$ has a 1 at position (j_1, \dots, j_k) if t restricted to $I_1^{j_1} \times \dots \times I_k^{j_k}$ is not the zero tensor, and a 0 otherwise.

Proof of Theorem 30. We will give a proof for the case $k = 5$, the other cases being similar. Define a block decomposition \mathcal{D} of the support $I_1 \times \dots \times I_5$ of Str_q^5 by

$$\begin{aligned} I_1 &= \{0\} \cup \{1, \dots, q\} \\ I_2 &= \{1, \dots, q\} \\ I_3 &= \{0\} \cup \{1, \dots, q\} \\ I_4 &= \{0\} \\ I_5 &= \{0\}. \end{aligned}$$

We have the outer structure $(\text{Str}_q^5)_{\mathcal{D}} = |11000\rangle + |01100\rangle \cong |10100\rangle + |00000\rangle$. Note that this is just an EPR pair between party 1 and 3. The inner structures are $\sum_{i=1}^q |ii000\rangle$ and $\sum_{i=1}^q |0ii00\rangle$, which are also known as $\langle 1, q, 1, 1, 1 \rangle$ and $\langle 1, 1, q, 1, 1 \rangle$. Let Cyc_5 be the map $t \mapsto t \otimes \sigma t \otimes \sigma^2 t \otimes \sigma^3 t \otimes \sigma^4 t$ with $\sigma = (12345)$. Let $\hat{\mathcal{D}} = \text{Cyc}_5 \mathcal{D}$ be the naturally corresponding decomposition. Then

$$\langle 2, 2, 2, 2, 2 \rangle^{\otimes s} = (\text{Cyc}_5 \text{Str}_q^5)_{\hat{\mathcal{D}}^{\otimes s}}^{\otimes s} \quad \text{and} \quad \underline{\mathbf{R}}((\text{Cyc}_5 \text{Str}_q^5)^{\otimes s}) \leq (q+1)^{5s}. \quad (1)$$

Note how the first statement relies on 5 being odd.

The inner structure of $(\text{Cyc}_5 \text{Str}_q^5)_{\hat{\mathcal{D}}^{\otimes s}}^{\otimes s}$ consists of tensors from $I := \{ \langle n_1, n_2, n_3, n_4, n_5 \rangle \mid n_1 \dots n_5 = q^{5s} \}$. Combining equation (1) with Proposition 32 gives that there are 2^s elements $t_1, t_2, \dots \in I$ such that

$$\underline{\mathbf{R}}(t_1 \oplus t_2 \oplus \dots) \leq (q+1)^{5s}.$$

Now the τ -theorem says that if we define τ by

$$2^s(q^{5s})^\tau = (q+1)^{5s}$$

then $\omega_5 \leq 5\tau$. Therefore,

$$\omega_5 \leq 5\tau \leq \log_q \frac{(q+1)^5}{2}$$

which gives $\omega_5 \leq 4.84438$. In general, one gets $\omega_k \leq \log_q \frac{(q+1)^k}{2}$ which is strictly smaller than k for q large enough. \square

References

- [ABDR04] Andris Ambainis, Harry Buhrman, Yevgeniy Dodis, and Hein Röhrig. Multiparty quantum coin flipping. In *Computational Complexity, 2004. Proceedings. 19th IEEE Annual Conference on*, pages 250–259. IEEE, 2004.
- [BCS97] Peter Bürgisser, Michael Clausen, and M Amin Shokrollahi. Algebraic complexity theory, volume 315 of Grundlehren der Mathematischen Wissenschaften, 1997.
- [BK15] Amey Bhangale and Swastik Kopparty. The complexity of computing the minimum rank of a sign pattern matrix. *arXiv preprint arXiv:1503.04486*, 2015.
- [Blä01] Markus Bläser. Complete problems for Valiant’s class of qp-computable families of polynomials. In *Computing and Combinatorics*, pages 1–10. Springer, 2001.
- [Blä13] Markus Bläser. Fast matrix multiplication. *Theory of Computing, Graduate Surveys*, 5:1–60, 2013.
- [BOC92] Michael Ben-Or and Richard Cleve. Computing algebraic formulas using a constant number of registers. *SIAM Journal on Computing*, 21(1):54–58, 1992.
- [CDS08] Eric Chitambar, Runyao Duan, and Yaoyun Shi. Tripartite entanglement transformations and tensor rank. *Physical review letters*, 101(14):140502, 2008.
- [CU13] Henry Cohn and Christopher Umans. Fast matrix multiplication using coherent configurations. In *Proceedings of the Twenty-Fourth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1074–1086. SIAM, 2013.
- [Ges15] Fulvio Gesmundo. Geometric aspects of iterated matrix multiplication. *arXiv preprint arXiv:1512.00766*, 2015.
- [Ike13] Christian Ikenmeyer. *Geometric complexity theory, tensor rank, and Littlewood-Richardson coefficients*. PhD thesis, Universität Paderborn, 2013.

- [LG14] François Le Gall. Powers of tensors and fast matrix multiplication. In *Proceedings of the 39th international symposium on symbolic and algebraic computation*, pages 296–303. ACM, 2014.
- [LM16] JM Landsberg and Mateusz Michałek. On the geometry of border rank algorithms for matrix multiplication and other tensors with symmetry. *arXiv preprint arXiv:1601.08229*, 2016.
- [LO11] Joseph M Landsberg and Giorgio Ottaviani. New lower bounds for the border rank of matrix multiplication. *arXiv preprint arXiv:1112.6007*, 2011.
- [NC10] Michael A Nielsen and Isaac L Chuang. *Quantum computation and quantum information*. Cambridge university press, 2010.
- [Sch81] Arnold Schönhage. Partial and total matrix multiplication. *SIAM Journal on Computing*, 10(3):434–455, 1981.
- [Str69] Volker Strassen. Gaussian elimination is not optimal. *Numerische Mathematik*, 13(4):354–356, 1969.
- [Str83] Volker Strassen. Rank and optimal computation of generic tensors. *Linear algebra and its applications*, 52:645–685, 1983.
- [VC16] Péter Vrana and Matthias Christandl. Entanglement distillation from Greenberger-Horne-Zeilinger shares. *arXiv preprint arXiv:1603.03964*, 2016.
- [VNYN13] Marcos Villagra, Masaki Nakanishi, Shigeru Yamashita, and Yasuhiko Nakashima. Tensor rank and strong quantum nondeterminism in multiparty communication. *IEICE TRANSACTIONS on Information and Systems*, 96(1):1–8, 2013.
- [Wol03] Ronald de Wolf. Nondeterministic quantum query and communication complexities. *SIAM Journal on Computing*, 32(3):681–699, 2003.