

A Selection of Giant Radio Sources from NVSS

D. D. Proctor

*Visiting Scientist, Lawrence Livermore National Laboratory, L-405,
7000 East Avenue, Livermore, CA, 94550; proctor1@llnl.gov*

ABSTRACT

Results of the application of pattern recognition techniques to the problem of identifying Giant Radio Sources (GRS) from the data in the NVSS catalog are presented and issues affecting the process are explored. Decision-tree pattern recognition software was applied to training set source pairs developed from known NVSS large angular size radio galaxies. The full training set consisted of 51,195 source pairs, 48 of which were known GRS for which each lobe was primarily represented by a single catalog component. The source pairs had a maximum separation of $20'$ and a minimum component area of 1.87 square arc minutes at the 1.4 mJy level. The importance of comparing resulting probability distributions of the training and application sets for cases of unknown class ratio is demonstrated. The probability of correctly ranking a randomly selected (GRS, non-GRS) pair from the best of the tested classifiers was determined to be $97.8 \pm 1.5\%$. The best classifiers were applied to the over 870,000 candidate pairs from the entire catalog. Images of higher ranked sources were visually screened and a table of over sixteen hundred candidates, including morphological annotation, is presented. These systems include doubles and triples, Wide-Angle Tail (WAT) and Narrow-Angle Tail (NAT), S- or Z-shaped systems, and core-jets and resolved cores. While some resolved lobe systems are recovered with this technique, generally it is expected that such systems would require a different approach.

Subject headings: astronomical data bases: miscellaneous — astronomical data bases: catalogs — galaxies: general — methods: data analysis — methods: statistical — techniques: image processing

1. INTRODUCTION

Using the Karl G. Jansky Very Large Array (VLA), the NRAO VLA Sky Survey (NVSS) (Condon et al. 1998) is a 1.4 GHz continuum survey covering the entire sky north of -40°

declination with a resolution of $45''$ FWHM. The associated catalog of discrete sources from this survey contains over 1.8 million entries. This catalog was generated by fitting flux densities of all significant peaks with elliptical Gaussians. Thus it is a catalog of components, where a single physical system may have multiple catalog entries.

While the NVSS survey and catalog have been out since 1998, few large scale attempts to identify GRS in the data have been reported. Lara et al. (2001) examined NVSS images above $+60^\circ$ (0.842 steradians or approximately 8% of the NVSS survey area) and presented a sample of 84 large angular size radio galaxies. This suggests that approximately one thousand similar systems should be present in the entire NVSS survey area. Machalski et al. (2001) compiled a list of 40 candidates of Fanaroff & Riley (1974) type II (FRII) or FRI/II with 1.4 GHz flux density on the NVSS maps of $S_{1.4} < 500$ mJy and sky area of 0.47 steradians. A recent attempt by Solovyov & Verkhodanov (2011) to develop an algorithm for selection of large, faint sources from the catalog was limited to sources between $4'$ and $6'$ separation, major axis of source lobes greater than $1'$, and integrated flux density $\lesssim 100$ mJy for at least one of the components. They reported 61 candidates for the initial stage of their work and subsequently reported on radio and optical identifications of 50 Giant Radio Galaxies (GRG) from that candidate list (Solovyov & Verkhodanov 2014). Andernach et al. (2012) has made a first visual inspection of approximately 3500 images of NVSS, SUMSS and WENSS radio surveys to search for such sources. Those results are pending.

For this report, GRS will be interpreted broadly to include any physically associated radio system with projected angular size $\geq 4'$. Thus GRS may include star-forming regions of, presumably nearer, galaxies as well as the other usual types. GRG are typically defined as having an overall projected size of ≥ 1 Mpc, thus identification of a GRS as GRG depends upon a distance determination, a determination beyond the scope of this paper.

Lara et al. (2001) confirmed 37 of the 84 GRS from their sample to be GRG. GRG are of interest in the study of evolution of galaxies, jet interaction with intergalactic medium, and testing consistency with the unified scheme for radio galaxies and quasars, as well as verifying cosmological hydrodynamical simulations. See for example, Ishwara-Chandra & Saikia (1999) and Malarecki et al. (2013) for more details.

The task of identifying GRS from the catalog data will be considered and the feasibility of the pattern recognition process to significantly expand the number of GRS and GRS candidates will be explored. Reported here are results of an initial stage of work to apply pattern recognition techniques to the problem of identifying large radio galaxy systems for which each lobe is primarily represented by a single component in the NVSS catalog. GRS with multi-component lobes are only found serendipitously with this procedure. Finding systems with multi-component lobes could be attempted with procedures similar to that

used for finding groups in FIRST (Proctor 2011) or the morphological-operator approach employed by Santiago-Bautista et al. (2013). It is expected that multiple approaches will be necessary to fully mine the information in the database.

The remainder of the paper is organized as follows. Section 2 contains general comments and background on pattern recognition procedures, including training set construction, feature set selection, decision-tree classifiers, cross validation process and classifier evaluation. Section 3 discusses training set results for variations in the training set, feature set, and classifier options. Section 4 discusses results of applying the best classifier ensembles found to the pairlist from the entire catalog, shows consistency between certain training and application sets, and gives a table and mosaic of resulting candidates. Finally, discussion and summary are presented in Section 5.

It should be noted the intensity scaling of the figures shown in this paper have often been chosen to enhance features of interest. Unless otherwise indicated, the cutouts shown in the mosaics are 20' in size. Note that only the tabulated coordinates that are associated with an NVSS catalog source have the significant figures of the NVSS catalog itself. The remainder are given to the same number of digits as the NVSS catalog for tabular consistency and convenience. The key to the annotations used in the candidate table is given in the appendix.

2. PATTERN RECOGNITION PROCEDURES

This section presents an overview of pattern recognition procedures. The pattern recognition process involves four steps: creation of a training set, selection of features, selection and training of the classifier and evaluation of the classifier. For those unfamiliar with the process, these steps are discussed in detail in the remainder of this section.

2.1. *Training set construction*

For this study, a training set was constructed from the population of NVSS catalog sources above $+60^\circ$ declination since that was what was available for a verifiable training set. The GRS target class was developed from the sample of 84 large angular size radio galaxies presented by Lara et al. (2001). In constructing their sample, they visually inspected all NVSS maps above $+60^\circ$ declination and pre-selected 122 map features apparently related to a single physical source, for which the total flux density was ≥ 100 mJy and the angular extension was larger than 4', as measured along the 'spine' of the source. They then observed

these sources with the VLA to confirm large angular size radio galaxies and reject those objects which were result of superposition of multiple adjacent sources. This resulted in 84 confirmed GRS and generally consisted of galaxies of angular size $\geq 4'$ and total flux density greater than 100 mJy at 1.4 GHz. It is noted that although a few sources fulfilling the requirements of the sample may have been missed due the the existence of "holes" in the maps available at the time, the list was felt to be sufficiently complete to demonstrate proof of principle for the pattern recognition process. From the Lara et al. (2001) list of 84 sources, 48 were selected for the pattern recognition training set. These 48 sources were those for which each lobe was primarily represented by single catalog component. The fitted version of the catalog was used both to avoid deconvolution issues and to have a direct correspondence to the images. Comparison with deconvolved parameters is left for future examination. Using the catalog parameters, the model silhouette size at the 1.4 mJy level was determined for each GRS component. The silhouette size is defined for this study as the area enclosed by the contour of the fitted model at the 1.4 mJy level. The minimum size noted for the larger-area component of each pair was 2.43 square arc minutes and minimum size for the smaller component was 1.87 square arc minutes. Components of the remaining resolved-lobe GRS were excluded from the training set. Using the minimum silhouette sizes found for the Lara GRS sample, a list of source pairs for the training set was constructed such that the first member of the pair was at least 2.43 square arc minutes and the second member was at least 1.87 square arc minutes, with the second member of the pair within $20'$ of the first and less than or equal to first in area. Aside from components of the resolved GRS, every source pair that satisfied the stated conditions were included in the training set. This resulted in a list of 51195 pairs for the training set, of which 48 were the GRS selected from Lara et al. (2001) sample.

Normally it is desired to construct a training set from a random sample from the entire population. This ensures that the statistics generated for the training set will be extendable to the remaining population (the application set) and that the resulting classifier produces results representative of the population. As the case here, this is not always feasible and subsequent evaluation of results, as discussed below, is required. We note that there is considerable discussion in the literature concerning the effect of skew (the class ratio, non-target to target), on classifier performance. For cases where one class is heavily outnumbered by the the other, as the case for this study, where it is over a thousand-to-one, the classifier may have problems learning the minority class. Several methods have been proposed to counter that effect, including over-sampling of the minority class and under-sampling of the majority class (Chawla et.al. 2002; Batista et al. 2004; Chawla et.al. 2008; Cieslak & Chawla 2008). This topic will be explored and examples of under-sampling the majority class will be presented and discussed. Comparisons of the resulting classifications for the training set

and application set (discussed in Section 4, below) should be made to verify that the results are self-consistent.

2.2. *Feature Set Selection*

The NVSS catalog provides position, peak flux, major and minor axes size, and position angle of the elliptic Gaussian fit for each source. Preliminary decision-tree training runs used five features - the silhouette size and peak flux of each pair member and the source pair separation. (It was noted in Proctor (2006) that the OC1 classifier was well able to adapt to different functional forms for variables in the feature set assuming all relevant information was available.) Given the relatively small number of GRS in the training set, it was decided to use silhouette size and peak flux for the components, contributing only four features to the feature set versus using the semi-major and semi-minor axes and peak flux, which would require six features, in addition to the pair separation. The number of source neighbors within 20' of the larger source, with silhouette size greater than 1.871 square arc minutes, was added for later tests as an attempt to include the source environment in the feature set. Table 1 is a list of features chosen. Most of the results reported here used this set of six features, the exception being an eight feature set discussed briefly in Section 3.7. Larger feature sets may be explored further in future work.

2.3. *Classifier Selection: Background, Overview, and Algorithm Options*

This section describes background relating to the choice of classifier and gives an overview of the chosen classifier with its options. For this work, the Oblique Classifier One (OC1) decision tree software of Murthy, Kasif, & Salzberg (1994) was chosen. OC1 has been used for a number of astronomical applications, including cosmic ray identification, star-galaxy separation, quasar candidate selection, X-ray source classification and sidelobe flagging in radio surveys. See White (2008) for a review. In that reference oblique decision trees were found to represent a good compromise between the demands of computational efficiency, classification accuracy, and analytical value of the results. It has also been used by the author for sorting triples in the FIRST database, (Proctor 2003) and comparing pattern recognition feature sets (Proctor 2006).

2.3.1. Classifier Overview

In general, a classifier is an algorithm that uses a set of numerical features describing members of a population, to separate those members into different classes. Decision trees are a supervised pattern recognition technique in which statistical methods are applied to typical data of known classification (the training set) to generate rules that can then be applied to the feature values of unknown objects (the application or test set) to determine their classification. A decision tree is a data structure that contains tests at branches and classifications at leaf nodes. Typically a binary tree is constructed in which the test at a branch compares a linear combination of the feature values to zero. If the test is true, a sample is sent down one branch, if false the other branch. The process continues until a leaf node is reached that contains a class label. The job of the algorithm is to use the training set to determine the coefficients of the features in the linear combination at each branch. Geometrically, this results in the feature space being partitioned by a set of hyperplanes into separate volumes, each with the appropriate label. OC1, while not the first oblique decision tree algorithm, is freely available¹, open source, and has been found easy to use. See Murthy, Kasif, & Salzberg (1994) for more detail. While more complex to construct, in general oblique trees are expected to provide a more compact representation.

Application of the resulting classifier to a sample member of unknown classification, results in that member being assigned a probability of being a particular class. To determine the probability estimate at a particular leaf node for the positive class, the so-called Laplace estimate has been adopted. If an object ends at a node with N_{node} training set objects, of which N_{pos} are of the positive class, then the tree's estimated probability in favor of the positive class for the object is $(N_{pos} + 1) / (N_{node} + 2)$. This form is applicable both for pure nodes, where $N_{node} = N_{pos}$ or $N_{node} = N_{neg}$ and nodes of mixed class resulting from any pruning of the tree. This was the form adopted by White et al. (2000) and Provost & Domingos (2003) and considered by the latter to be the de facto standard, though others still use maximum-likelihood estimates. Provost & Domingos (2003) found that, while not surprising statistically, the uniformity and magnitude of the improvement using this estimate to be remarkable.

Five fold cross-validation was adopted. Cross-validation is a standard pattern recognition technique used to avoid bias that would occur if points used in testing were the same as those used in training. For five fold cross validation the training set members are divided into five folds (groups) and training set members from four folds are used to generate classifiers for the remaining fold, each fold being classified in succession.

¹See <http://ccb.jhu.edu/software/oc1/oc1.tar.gz>

If multiple classifiers are generated using some randomization procedure, as in the search for the best hyperplane, the resulting probability estimate is expected to be of improved accuracy and its standard deviation can be calculated. Such improvement was demonstrated by Provost & Domingos (2003). For the current work, ten decision trees were generated with random seeds for each fold, resulting in an ensemble of ten decision trees contributing to probability estimates for each fold for the training set and 50 decision trees contributing to the probability estimates for the entire catalog application set members. It should be noted that this is a conditional probability, depending on the classifier, training set, and feature set. Rankings based on probability of class membership are needed so that cases most likely to belong to the target class can be considered first. The ensemble average of the resulting probability estimates for a member will be designated the normalized score or vote. Ordering this vote, say high to low, for the entire sample, results in a sort-ordered distribution, with index 1 to N, where N is the number of members in the sample. If the index is divided by the number of sample members it becomes the sample-size normalized index. The plot of normalized score or vote versus the sample-size normalized index is designated the vote curve.

For each source i , if $P_i(GRS)$ is the estimate in favor of the source being GRS and $P_i(nonGRS)$ is the estimate against, then

$$P_i(GRS) + P_i(nonGRS) = 1. \quad (1)$$

Thus,

$$\sum_{i=1}^N P_i(GRS) + \sum_{i=1}^N P_i(nonGRS) = N, \quad (2)$$

where N is the number of points in the sample under consideration. Normalizing by N gives

$$[\sum_{i=1}^N P_i(GRS) + \sum_{i=1}^N P_i(nonGRS)]/N = 1. \quad (3)$$

The normalized total in favor is the 'area under the curve' of the sort-ordered, sample-size normalized vote plot. It allows comparison of vote curves for the training set and application set, as will be shown in Section 4. See Proctor (2003) and Proctor (2006) for other examples of vote curve comparisons.

2.3.2. Classifier Options

OC1 allows, among other options, choice of pruning portion, impurity measure, order of coefficient perturbation, number of jumps attempted to get out of a local minimum, and the number of restarts at each node to test for the best hyperplane. Each will be discussed in more detail in the following paragraphs. The other OC1 program options should not affect solution results.

Pruning portion: OC1 prunes decision trees by default, to avoid the problem of overfitting. The only pruning method implemented is error complexity pruning using a separate pruning set. This option specifies that a chosen fraction of the training set will be used exclusively for pruning. If the pruning portion = 0, no pruning is done.

Impurity measures: The impurity measure is the metric used to determine the goodness of a hyperplane location. New nodes are added so as to minimize the "impurity" of the training set members at the node. Impurity measures Gini Index, Twoing Rule, Max Minority, Sum Minority and Information Gain are distributed with the OC1 software. Early tests showed performance of Max Minority and Sum Minority for lower skew training sets was quite poor, and they were dropped from further comparisons. Two other impurity measures were tested, since they were developed specifically to be skew insensitive. The impurity measure, designated Flach in tables below, was developed by Flach (2003) as a skew insensitive splitting criterion, in particular

$$1 - \frac{2tpr \cdot fpr}{tpr + fpr} - \frac{2(1 - tpr)(1 - fpr)}{1 - tpr + 1 - fpr} \quad (4)$$

is optimized in selecting the split at a node. Here tpr is the ratio of true positive to total positive instances for the trial split and fpr is the ratio of false positive to true negative. For the impurity measure denoted Hellinger, the form for relative impurity for the Hellinger distance developed by Cieslak & Chawla (2008) as a skew insensitive measure is

$$\sqrt{(\sqrt{tpr} - \sqrt{fpr})^2 + (\sqrt{1 - tpr} - \sqrt{1 - fpr})^2}. \quad (5)$$

Order of coefficient perturbation: At each node OC1 adjusts the coefficients of the hyperplane under consideration. The default order of adjustment is sequential, which means all the coefficients of a hyperplane are perturbed in order to search for the best value to be found for that coefficient. The alternatives are best first and random. For best first, the coefficient that provides the greatest improvement in the impurity measure is done first. For random coefficient perturbation, a coefficient is selected at random and the number of times to pick a random coefficient can be specified.

Number of jumps to escape local minimum: When the algorithm cannot improve any of the coefficients of a hyperplane deterministically, it is stuck in a local minimum. It then attempts a number of jumps out of this minimum by choosing a random direction and sliding the hyperplane in that direction to improve the impurity measure.

Number of restarts: The number of restarts is the number of random hyperplanes at each node examined for best position. This includes the best axis parallel split, if desired.

2.4. Classifier Evaluation

The confusion matrix, also called the contingency table, is useful for defining evaluation metrics. For the two class problem under consideration, the confusion matrix becomes as shown in Table 2, where TP, FP, FN, TN stand for true positive, false positive, false negative, and true negative counts respectively; PP and PN stand for predicted positive and predicted negative counts; and POS and NEG stand for the number of positive and negative class members respectively, with N_{ts} the training set size, ($POS + NEG = N_{ts}$). Lower case is used for relative frequencies, e.g. $tp = TP/N_{ts}$. The true positive rate (hit rate) is defined as $tpr = TP/POS$ and the false positive rate (false alarm rate) $fpr = FP/NEG$. Accuracy is defined as $Acc = (TP + TN)/N_{ts}$ and error rate as $Err = (FP + FN)/N_{ts}$. Another important quantity is the class ratio or skew, defined here as $c = NEG/POS$. Without loss of generality, the GRS class is considered the positive class for this paper.

Until recently, classification accuracy has been the primary metric used to evaluate classifiers (Demšar 2006). As class distribution becomes more skewed, evaluation based on accuracy breaks down, as it is biased toward the dominate class. It has become common to use Receiver Operating Characteristic (ROC) analysis to make more general comparisons (Ferri et al. 2002). A ROC graph depicts trade offs between tpr and fpr, typically shown as fpr on the x-axis and tpr on the y-axis. The Area Under the ROC Curve (AUC) gives the classifier performance as a single scalar and has been shown to be equivalent to the probability that a randomly chosen (positive class member, negative class member) pair will be correctly ranked (Hanley & McNeil (1982), Batista et al. (2004), and references therein). An algorithm for generating a ROC curve from a set of ranked examples was given by Provost & Fawcett (2001) and was adopted for this work. It generates a segment that bisects the area that would have resulted from the most optimistic and most pessimistic orderings of examples with the same score or vote. Note that a specific classifier instantiation is represented by a single point on the ROC curve, thus only when one classifier is better over entire performance space can it be declared the better classifier. Example ROC curves will be shown below. Comparing classifiers using ROC analysis enables comparison of ranking quality across the entire range of possible class thresholds. Hanley & McNeil (1982) discuss the three way equivalence between the AUC, the Wilcoxon-Mann-Whitney non-parametric test statistic ("the Wilcoxon"), and the probability of a correct ranking of, in the case here, a (GRS, non-GRS) pair. They also developed a method for estimating the standard error of

the difference between the areas of two ROC curves for case of applying different treatments to the same data. They give

$$SE(\hat{AUC}_1 - \hat{AUC}_2) = \sqrt{SE^2(\hat{AUC}_1) + SE^2(\hat{AUC}_2) - 2 \cdot r \cdot SE(\hat{AUC}_1) \cdot SE(\hat{AUC}_2)} \quad (6)$$

where $\hat{AUC}_1$ and $\hat{AUC}_2$ are the respective area estimators of the areas for the two cases, $SE(\hat{AUC}_1)$ and $SE(\hat{AUC}_2)$ are their respective standard errors, and r is a quantity representing the correlation introduced between the two areas from using the same sample training set. The standard errors are given by

$$SE(\hat{AUC}) = \sqrt{\frac{\hat{AUC}(1 - \hat{AUC}) + (N_{GRS} - 1)(Q_1 - \hat{AUC}^2) + (N_{nonGRS} - 1)(Q_2 - \hat{AUC}^2)}{N_{GRS} \cdot N_{nonGRS}}}$$

where estimators for Q_1 and Q_2 are $Q_1 \simeq \hat{AUC}/(2 - \hat{AUC})$ and $Q_2 \simeq 2 \cdot \hat{AUC}^2/(1 + \hat{AUC})$ and N_{GRS} and N_{nonGRS} are the number of GRS and non-GRS in the training set, respectively. A table in the reference then gives r as a function of the average of two areas vs. the average of r_{GRS} and r_{nonGRS} , where r_{GRS} is the correlation coefficient between the estimated probabilities for the GRS in each area and r_{nonGRS} is the correlation coefficient between the non-GRS for each area. They suggest Kendall's tau (see Press et al. (1992) for an algorithm) as the appropriate correlation statistic for results obtained from an ordinal scale, as the case here. To test the null hypothesis that the areas are the same for both cases, the ratio z is calculated as

$$z = \frac{\hat{AUC}_1 - \hat{AUC}_2}{SE(\hat{AUC}_1 - \hat{AUC}_2)} \quad (7)$$

and compared to the value of z_c , the critical value for the probability of Type I error chosen.

3. TRAINING SET RESULTS

This section presents training set results for variations in pruning, skew, impurity measures, order of coefficient perturbation, number of hyperplanes per node examined, and number of jumps attempted to escape local minimum. A feature set comparison is also made and GRS outliers are discussed.

3.1. Pruning portion

Initial attempts with training sets of larger skew showed that it was necessary to avoid pruning to obtain a classifier solution. Subsequently, it was learned that Provost & Domingos

(2003), in a study of decision tree induction for probability-based rankings, found larger trees may be better for probability estimation. Thus, subsequent results have all been generated without pruning.

3.2. *Training Set Skew-Ratio Comparisons*

While experience suggested OC1 is fairly robust, there was concern about its ability to construct decision trees with such high skew. Thus it was decided to begin with examination of synthetic training sets with smaller number of non-GRS sources to explore the effect of skew on the quality of resulting decision trees. Tests were run with modified training sets with 48, 1000, 5000, 17000, and 34000 non-target source pairs, as well as the full training set. The AUC, and average probability estimate and maximum probability estimate of the training set GRS pairs were determined. Results are shown in Table 3. Calculation times for training set construction (with an ensemble of 50 decision trees) ranged from a few minutes for the smaller sets to a few hours for the full training set on an x86_64 processor.

Figure 1(a) shows ROC curve comparisons between the highest-skew and lowest-skew training sets and Figure 1(b) shows the comparison between the two highest skew sets examined. For intermediate skew ratios, not shown, various amounts of interweaving of curves occurred. However, it is noted that there appears to be a monotonic trend of improving AUC as skew ratios approach the inherent skew ratio for the population.

Application of the Hanley & McNeil (1983) formula, Eqn. 6 above, for the standard error of the difference in areas of two ROC curves, for the 48:48 skew curve and 51147:48 skew curve resulted in $z=1.22$. Thus the hypothesis of no significant difference is accepted at the 5% level. However, the 48:48 skew curve being entirely below the 51147:48 curve, whereas the others interweave suggests it is more marginal. A clear trend of degrading mean probability estimate and maximum probability estimate for the training set GRG as the skew ratio increases is apparent. Provost & Domingos (2003) in a study of tree induction for probability based rankings noted decision trees have been found to provide poor probability estimates, but discussed why decision tree representation is not intrinsically inadequate and recommended they be considered when rankings based on class membership is desired. Margineantu & Dietterich (2001) show probability estimation trees produce surprisingly good rankings, even when the probability estimates themselves are questionable. In Section 4 below, vote curve comparisons will be made that provide greater distinction between the various skew-ratio cases.

3.3. Impurity Measure Comparisons

Table 4 shows the results of utilization of the various impurity measures using the full training set. Figure 2(a) shows ROC curve comparisons for the Flach and Information-Gain impurity measures, the two extreme AUC values for impurity measure tests and Figure 2(b) shows the comparison between Flach and Hellinger impurity measures. The remaining impurity measures compared with Flach also show considerable interweaving of the ROC curves, suggesting that while there is little to distinguish Flach from Gini, Variance, Twoing Rule or Hellinger, the difference between Flach and Information-Gain impurity measures may be more marginal.

Applying the formula for the standard error of the difference in areas of two ROC curves, for $\hat{AUC}(Flach) - \hat{AUC}(information\ gain)$, the highest and lowest AUC in the table, results in $z=1.60$. This is less than the critical $z_c = 1.96$ for two sided value for normal distribution, for 5% probability of Type I error. Thus the hypothesis of no significant difference between these two ROC areas is accepted.

3.4. Order of Coefficient Perturbation Comparison

The order-of-coefficient-perturbation option determines the order in which the hyperplane coefficients are adjusted in searching for impurity measure improvement. Table 5 shows an order-of-coefficient-perturbation comparison. A difference of AUC comparison of best-first and random order gave $z = 1.40 < z_c$ indicating the hypothesis of no significant difference is accepted at the 5% level. It is noted however, -R12 option (loop 12 times picking a random coefficient and attempt to perturb it) also resulted in significantly longer computation times.

3.5. Number of Hyperplanes per Node Comparison

The number of restarts, i , is the number of hyperplanes searched for the best position at every node of the decision tree. The best axis parallel hyperplane and $(i-1)$ random hyperplanes are used. A difference of AUC calculation between using 20 and 60 hyperplanes/node resulted in accepting the hypothesis of no significant difference. Of course the more hyperplanes per node used the longer the decision tree generation run times.

3.6. *Number of Jumps attempted to escape local minimum*

Doubling the number of jumps, from 20 to 40 did not provide significantly different results.

3.7. *Feature Set Comparisons*

An eight feature set was compared to the six feature set using the Flach impurity measure. The two additional features used were the ratio of minor to major axes for each component. There was no significant difference in AUC at the 5% level.

3.8. *Outliers*

The sources J0317+769, J0342+636, J0508+609, J1036+677, J1847+707, J1951+706 showed votes consistently near zero over the various decision tree runs. Further examination of these sources suggested J0317+769 and J0508+609 may have had possible wrong choices for training set pairs for the lobes and J1847+707 has possible contamination from a unresolved chance projections. The poor results for J0342+636, J1036+677, and J1951+706 remain unclear, though contamination by chance projections remain a possible problem. Also, it is noted that while training set GRS have been lumped into one class, different types of GRS are expected to occupy different regions of feature space, a la FRI and FRII types. The small number of training set GRS may well be insufficient to distinguish these regions.

4. APPLICATION SET RESULTS

The candidate pairlist for the entire catalog was constructed with the same size constraints used for the training set and consisted of 870,370 pairs, of which 817,603 were below +60° declination. This section discusses types of contamination found, makes a comparison of training set and application set vote curves and finally presents a table and mosaic of higher-ranked sources.

4.1. Contamination

Initial examination of images and contour plots of higher-ranked sources showed significant contamination from galactic plane sources, including high probability HII regions, about four dozen ring and shell type sources, most previously identified as Super Nova Remnant (SNR), and some apparent chance projections. Also found were side-lobe artifacts from certain bright sources.

A plot of the galactic coordinates of sources with silhouette sizes greater than 2.438 square arc minutes (the minimum size of at least one component for pairlist inclusion) for the entire catalog are shown in Figure 3. It is noted that the ratio of large silhouette size sources to total number of sources for the training set and application set differed by about five percent. That difference may help account for excess galactic plane sources in the higher-ranked results. Additionally, the current feature set may not be sufficient for discriminating against these galactic plane sources.

Figure 4 shows example cutouts of some of this contamination. A random selection of example higher-ranked source pairs from the galactic plane are shown in the first row of Figure 4. The second row of Figure 4 shows a selection of NVSS cutouts of larger angular size entries from the Givon et al. (2005) catalog of Galactic radio compact HII regions at 1.4 GHz. These were chosen to demonstrate the various morphologies of these sources in NVSS. It is noted that it is possible for pairs of HII regions to mimic GRS. NVSS images of 890 high probability planetary nebula from Acker et al. (1992) were separately examined. While perhaps 8% showed no NVSS signature and most were point-like sources, a few had resolved morphology. The third row of Figure 4 shows a selection of NVSS images of larger angular diameter Acker sources for comparison. We note that only one planetary nebula (NGC 1514) was identified in the higher-ranked source pair list. Finally the fourth row of the figure shows NVSS images of known SNR with Green (2009) designations recovered from the higher-ranked sources. Of the ring and shell type sources, it is noted that 15 were subsequently identified as known SNR. The remainder were an assortment identified in SIMBAD with HII regions, young stellar objects, dark nebulae, and bubbles. More features may be needed to discriminate against these sources.

4.2. Vote Curve Comparison of the Training and Application Sets

Initial evaluation of the area under the vote curve for all pairs below $+60^\circ$ declination resulted in a determination of 0.00227 ± 0.00005 . Excluding galactic plane sources and sources around certain bright sources resulted in an improved area determination of 0.00196 ± 0.0003 ,

an improvement of over five standard deviations. Thus for further application set vote curve comparisons, pairs above $+60^\circ$ declination (the training set) and pairs within 5° of the galactic plane were excluded. Also excluded were pairs within $90'$ of Cyg A, M1, M42 and M87, areas with significant artifacts, leaving 438,350 pairs. Vote curves for this restricted application set, using decision trees from the training sets with 48, 1000, 5000, 34000, and 51147 nonGRS, are shown in Figure 5(a). As the training set skew increases, they show a clear trend of improvement towards the target area under the curve of $48/51195 \simeq .00094$. Note that for a perfect classifier, the vote curve would have a constant value of 1.0 at $x = 0$, then drop to 0.0 at the normalized index of ~ 0.00094 . For greater detail, comparison of the 51147:48 skew restricted application-set vote curve with the 51147:48 skew training-set vote curve over the initial index range is shown in Figure 5(b). Figure 5 clearly demonstrates the need to compare vote curves for the training set and application set when the population's underlying skew ratio is unknown. Of course, for sufficiently strong features, this skew effect may be obviated.

A possible explanation for the difference between the the training set vote-curve area and the ideal-classifier area may be found in those resolved GRS sources that were excluded from the training set. As will be discussed below, 19 were found to be higher-ranked sources when the decision trees were applied to their largest components. If those are included, the target area would become $(48+19)/(51195+19)=.00131$, a near match to the training set vote-curve area.

There are still some differences between the 51147:48 application set and training set vote-curve areas. Possible reasons include false positives in the training set due to, for example, unresolved chance projections in the GRS class. Also possible are unidentified true GRS labeled nonGRS. There do not appear to be any GRS source pairs in the Lara et al. (2001) sample for which both lobes are represented by approximate point-spread functions. It may be that such pairs, though possible GRS, were not included in the original pre-selection.

4.3. *Higher-Ranked Candidates - Table and Mosaic*

Images and contour plots of over four thousand higher-ranked sources were examined. These sources were produced using several of the better decision-tree ensembles. For the pairs above $+60^\circ$, the rankings from the training-set decision trees were used. Preliminary work was done to eliminate known HII regions, SNR, ambiguous galactic plane sources, and sources for which FIRST counterparts indicated chance projections. Table 6 presents the coordinates of 1616 GRS and candidate GRS from the higher-ranked sources, along with size and preliminary morphological annotations. The key to the annotations are given in

Table A-I. Typically, GRS/GRG are assigned an FR type. For this work, FR classification was not attempted. Though many of the larger 'd' and 't' annotated sources are clearly FR II, the smaller, less resolved sources were difficult to evaluate. It was felt such distinctions deserved more uniform analytic attention. In Table 6, other morphological types were retained. The annotations can be used to select promising candidates of morphologies of interest. A few incidental identifications, not from a candidate pair list are also included, and annotated as such.

While well over a thousand of the sources in Table 6 appear to be viable GRS candidates, approximately one quarter appear to be resolved cores, chance projections, or ambiguous cases. They are included for possible future investigations. Note a few of the systems in this table are less than $4''$, but also are included for documentation purposes. Also, it should be noted that not all sources in this table have both source pair members as components. This particularly applies to the resolved-core type morphology systems.

The significance of the system coordinate needs to be considered in the context of the system morphology and the estimated size, i.e. only the tabulated coordinates that are associated with an NVSS catalog source have the significant figures of the NVSS catalog itself. The remainder are given to the same number of digits as the NVSS catalog for tabular consistency and convenience. In this table, coordinate type 'c' indicates a NVSS catalog source presumed to be the core, and coordinate type 'v' is an estimate for the core position using lobe coordinates. These latter, in general, were the average of the component positions, though some may have had visual adjustments. Figure 6 is a sample of those candidates, including those judged to have chance projection or otherwise ambiguous appearance, for which future observations could be made. The complete mosaic is available in the online version of the journal.

While no comprehensive attempt was made at cross-identifications, those Lara et al. (2001), Solovyov & Verkhodanov (2014), and Machalski et al. (2001) GRS recovered were retained in the table, but not given another IAU designation. The coordinates listed for these systems are as constructed for the other candidates in the table. It is noted that of the 50 reported Solovyov & Verkhodanov (2014) GRS, 43 are in this candidate list. (J185618.6+013120, though in their table of 51 entries, is a galactic plane source and was dropped from their further discussion.) Of their remaining 7 GRS, four were resolved sources not expected to be found by this method, two (J172331.0-352542, J182708.3-124020) were perhaps more questionable galactic plane identifications, and one (J122045.0+055204) appeared to be a sidelobe of J121925.2+054945.9. It is of interest that 19 of the 38 GRS of Lara et al. (2001) sample that were excluded from the training set due to their resolved lobes, never-the-less, were recovered. The overall GRS recovery rate from the Lara sample

using a single ensemble is approximately half of such systems. The recovery rate for the FRII and FRI/II systems of Machalski was only 40%, perhaps reflecting the differing flux density limits for the Lara and Machalski samples.

5. SUMMARY and CONCLUSIONS

Pattern recognition procedures have been developed for recovering candidates for a significant fraction of the expected number GRS from the NVSS catalog data. An overview of the classification process was given. Training set construction, in particular the effect of skew was examined. ROC curve analysis and AUC comparisons, along with vote-curve comparisons were used to evaluate the classifiers.

The results of a selection of classifier options were compared. It was found necessary to avoid pruning. This is consistent with the observations of Provost & Domingos (2003) that larger trees may provide better classifications. The impurity measure comparisons showed that choice of impurity measure did not have a strong effect on classifier quality.

The need for vote curve comparison of the training set and application set when population skew ratios are unknown was demonstrated. The best consistency between the training set and application set was for training-set skew ratios that more closely approached the inherent ratio. The AUC and vote curve analysis suggested that for the given feature set, OC1 was quite adequate for the higher-skew training sets.

Cutouts of the higher-ranked source pairs were examined and preliminarily screened to eliminate known HII regions, SNR and chance projections, as well as ambiguous galactic plane sources. A table of GRS and GRS candidates was constructed. These include doubles and triples, and WAT, NAT, W, X, DD and S/Z-shaped systems. Some GRS candidates with more PSF-like lobes were found in the candidate list, but since they were not represented in the training set, they are not expected to be well represented in the high probability sources.

While classifiers have been generated that produce greater than 95% probability that a randomly selected (GRS,non-GRS) pair will be properly ranked, probability estimates show considerable room for improvement and thus, are an area for future study. Perhaps such low average probability estimates for training-set GRS are not surprising, given the relatively small number of training set GRS, the possibility of false positives and false negatives in the training set, and the grouping of GRS types into a single class. However, a baseline for future work has been established. A more profitable line of inquiry may be to apply pattern recognition to separating the morphological types using the enlarged database and then revisiting the GRS vs. nonGRS classification. A further area for study involves investigation

of more powerful statistical tests for AUC comparison.

With a larger database, there is promise of being able to sort types of radio galaxies and identify their regions in parameter space. Future work involves exploring options for including source shape as well as core and redshift information and examination of the decision tree ensembles to determine what they reveal about the locations of various morphologies in feature space. While it is not impossible to visually search the entire set of NVSS maps for particular morphologies, it is believed pattern recognition techniques will be needed to fully exploit information contained in the catalog and images. Observationally verified training sets will be needed and the importance of a representative training set should be emphasized.

The author is grateful to the anonymous referees for comments, suggestions, and a reference, which led to significant improvements of the paper. Thanks are due to NRAO for page charge support. The author appreciates office space and computing facilities provided by Lawrence Livermore National Laboratory’s Institute of Geophysics and Planetary Physics (IGPP), John Bradley, director, during first phases of this work and Kem Cook, during later stages, as well as office space and computing facilities provided by B Division and host Tom McAbee, also of LLNL. The author also appreciates further support provided by James Trebes, Physics Division, LLNL. Thanks are also due Bill Cotton for software to access the NVSS images and Robert Becker for computer resources.

Extensive use was made of the IDL astro package (<http://idlastro.gsfc.nasa.gov>), GDL (<http://gnudatalanguage.sourceforge.net>), and IDL (a registered trademark of Exelis Visual Information Solutions). The free availability of the OC1 decision tree software (<http://ccb.jhu.edu/software/oc1/oc1.tar.gz>) was greatly appreciated. The NVSS data was taken by the NRAO Very Large Array. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. This research has also made use of NASA’s Astrophysics Data System (ADS), the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration, and the Sloan Digital Sky Survey (SDSS), as well as the VizieR catalogue access tool, CDS, Strasbourg, France and the SIMBAD database, operated at CDS, Strasbourg, France.

This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory in part under Contract W-7405-Eng-48 and in part under Contract DE-AC52-07NA27344.

A. APPENDIX MATERIAL

Table A-I provides the key for annotations in Table 6, the table of GRS and candidates. It should be noted the annotations are not necessarily mutually exclusive, for example an S or Z-shaped source may also be a double-double candidate, thus groups may have multiple morphological classification types listed. If multiple morphological types are given, they generally are given in order of estimated probabilities.

While most keyed items in Table A-I are self-explanatory, a few comments are in order.

The 'cj' annotation refers to two components, one larger and brighter, the other smaller size and possible curved morphology. The 'd' annotation consists of two components of similar silhouette size and peak flux. The distinction between the two groups and becomes difficult for smaller and fainter systems.

The 'dd' annotation refers to a pair of double radio sources with a common center. Some of these are candidate double double radio galaxies.

A few systems were noted for which the lobes appeared to have different FR morphologies, one diffuse with brightest region within the inner half of the radio source, the other with brightest regions near the outer edge. See Gopal-Krishna & Wiita (2000) for detailed discussion of HYMORS.

The annotation 'irr' denotes a group whose members appear to be physically associated but are not well described by another morphology class. Such groups may provide candidate merger or relic type sources.

The annotation 'mg' (multiple possible groupings) was used to refer to groups for which multiple possibilities for physical associations appeared reasonable. This annotation was followed by possible likely groupings. These groups require higher resolution, sensitivity, and/or spectral data for definitive classification.

The annotation 'rc' (resolved core) was used to describe a single contiguous non-pointlike, extended source that may or may not have associated jets or lobes apparent and may or may not have been resolved into multiple catalog components. Their silhouette may be irregular. Optical counterparts for some of these sources were found to be the centers of large galaxies.

The distinction between double, resolved core, and core-jet morphology becomes rather arbitrary for smaller angular diameter groups.

The annotation 'unu' was generally applied to unusual configurations that could be explained by chance projection. If they are not chance projections, they would be deemed interesting.

For the purposes of this paper, the annotation 'dl' (dogleg) is used to describe jets or lobe portions for which there is an apparent abrupt change in direction.

The annotation 'int' (interesting) is a subjective evaluation, which may well have varied over the course of the classifications.

The comment 'tail' indicates a protrusion (normally low level) from lobe in direction away from core, whereas 'jet' indicates a protrusion from a presumed core.

An 's' annotation for a source indicates the source appeared to be a source associated with the group by chance projection. For some source components, the assignment of jet or lobe annotation was difficult and even arbitrary.

Table A-I. Key for Table 6

Code	Classification or Description
amb	ambiguous
arc	arc or C-shape, possible edge brightened lobe, bent jets, or ring fragment
cj	core-jet
d	double lobe radio galaxy - may include core-jet sources
dd	double double morphology (DD), may be candidate DDRG
dl	dogleg
fishtail	multiple (usually two) lower level tails
hymor	hybrid morphology radio source (HYMOR)
int	interesting
irr	irregular, distorted
incidental	non-source pair systems found during cutout examinations
jet	protrusion away from presumed core
lobe	lobe, resolved lobe
mg	multiple groupings possible
nat	narrow angle tail (NAT)
point	non-extended source for which the major and minor axes appear to be the same size
rc	resolved compact source, non-pointlike
ring	ring
ring-lobe	edge or rim brightened lobe or embedded ring
s	single source (sometimes with sidelobes) or probable chance projection of point-like source into group
sz	S or Z-shaped
t	triple, no bend, little extended structure
tail	protrusion (normally lower level) from lobe in direction away from core
unu	unusual, uncommon - may be due to chance projection
w	W-shape, wiggles
wat	wide angle tail (WAT)
x	X-shape

REFERENCES

- Acker, A., Marcout J., Ochsenbein F., Stenholm B., Tylanda R., Schohn C. S Strasbourg-ESO Catalogue of Galactic Planetary Nebulae (ISBN 3-923524-41-2), 1, 1 (1992) Garching, Germany: European Southern Observatory
- Andernach, H., Jiménez Andrade, E. F., Maldonado Sánchez, R. F., Vázquez Báez, I. R. 2012, in *Science from the Next Generation Imaging and Spectroscopic Surveys, 2012* P1, <http://www.eso.org/sci/meetings/2012/surveys2012/posters.html>
- Batista, G. E. A. P. A., Prati, R. C., Monard, M. C., 2004, *SIGKDD Explorations*, 6(1), 20
- Murthy, S. K., Kasif, S., & Salzberg, J. 1994, *J. Artif. Intell. Res.*, 2, 1
- Chawla, N. V., Bowyer, K. W., Hall, L. O., & Kegelmeyer, W. P., 2002, *J. Artif. Intell. Research*, 16, 321
- Chawla, N. V., Cieslak, D. A., Hall, L. O., Joshi, A. 2008, *Data Mining and Knowledge Discovery*, 17, 225
- Cieslak, D. A., Chawla, N. V., 2008, *Lecture Notes in Computer Science* No. 5211, 241 Springer-Verlag
- Condon, J. J., Cotton, W. D., Greisen, E. W., et al. 1998, *AJ*, 115, 1693
- Demšar, J., 2006, *Journal of Machine Learning Research* 7, 1
- Fanaroff, B. L., Riley, J. M. 1974, *MNRAS*, 167, 31
- Ferri, C., Flach, P., Hernandez-Orallo, J., 2002, *Proceedings of the 19th International Conference on Machine Learning (ICML-02)*, (San Francisco, CA: Morgan Kaufmann) 139
- Flach, P. A. 2003, *Proceedings of the Twentieth International Conference on Machine Learning (ICML-2003)*, (Palo Alto, CA: AAAI Press), 194
- Flach, P. A. 2003, *Proceedings of the Twentieth International Conference on Machine Learning (ICML-2003)*, Washington DC, 2004.
- Green, D. A., 2016, Private communication.
- Green, D. A., 2014, *Bull. Astr. Soc. India*, 42, 47
- Green, D. A., 2009, *Bull. Astr. Soc. India*, 37, 45

- Giveon, U., Becker, R. H., Helfand, D. J., White, R. L. 2005, *AJ* 130, 156
- Gopal-Krishna, Wiita, P. J. 2000, *A&A*, 363, 507
- Hanley, J. A., McNeil, B. J. 1982, *Radiology* 143, 29
- Hanley, J. A., McNeil, B. J. 1983, *Radiology* 148, 839
- Ishwara-Chandra, C. H., Saikia, D. J. 1999, *MNRAS*, 309, 100
- Lara, L., Cotton, W. D., Feretti, L., Giovannini, G., Marcaide, J. M., Márquez, I., Venturi, T. 2001, *A&A*, 370, 409
- Machalski, J., Jamroz, M., Zola, S. 2001, *A&A*, 371, 445
- Malarecki, J. M., Staveley-Smith, L., Saripalli, L., Subrahmanyan, R., Jones, D. H., Duffy, A. R., Rioja, M., 2013, *MNRAS*, 432, 200
- Margineantu, D. D., Dietterich, T. G. (2001) In C. Holmes (Ed.), *Nonlinear Estimation and Classification*. The Mathematical Sciences Research Institute, University of California, Berkeley.
- Press, W. H., Teukolsky, A. A., Vetterling, W. T., Flannery, B.P., *Numerical Recipes in C* - 2nd ed., 1992 Cambridge University Press, Cambridge
- Proctor, D. D. 2003, *J. Electron. Imaging*, 12, 398
- Proctor, D. D. 2006, *ApJS*, 165, 95
- Proctor, D. D. 2011, *ApJS*, 194, 31
- Provost, F., Fawcett, T. 2001, *Machine Learning*, 42, 203
- Provost, F., Domingos, P. 2003, *Machine Learning*, 52, 199
- Santiago-Bautista I., Andernach H., Rodríguez-Rico C.A. 2013, *The Modern Radio Universe* 2013, 178, <https://indico.mpifr-bonn.mpg.de/indico/getFile.py/access?rsId=19>
- Solov'yov, D. I., Verkhodanov, O. V. 2011, *AstBu*, 66, 4, 416
- Solov'yov, D. I., Verkhodanov, O. V. 2014, *AstBu*, 69, 2, 141
- White, R. L. et al., 2000, *ApJS*, 126, 133
- White, R. L. 2008, in C. A. L. Bailer-Jones (Ed.), *Classification and Discovery in Large Astronomical Surveys*, 2008 AIP Conference Proceedings, 1082, 37




Fig. 1.— (a) A ROC curve comparison of training sets with skew 51147:48 and skew 48:48. (b) A ROC curve comparison of training sets with skew 51147:48 and skew 34000:48.




Fig. 2.— (a) ROC curve comparison of classifiers using Flach and Information-Gain impurity measures. (b) ROC curve comparison of classifiers using Flach and Hellinger impurity measures.




Fig. 3.— Galactic coordinates of catalog sources with silhouette size ≥ 2.438 square arc minutes. A random selection of 10% of points are shown for improved plot clarity.




Fig. 4.— Examples of ‘contamination’ and possible contamination in the training and application sets. First row: Random selection of NVSS cutouts of galactic plane sources from higher-ranked pairlist. Second row: Example NVSS cutouts of larger silhouette size Givon et al. (2005) catalog high reliability compact HII sources with NVSS coordinate and the MSX6C designation listed by Givon et al. (2005). Third row: Example NVSS cutouts of larger silhouette size Acker et al. (1992) catalog high probability planetary nebulae with NVSS coordinate and PNG designation. Fourth row: NVSS cutouts of known SNR from higher-ranked pairlist with Green (2009) designations. Image cutout size is $20'$ square.




Fig. 5.— (a) Application Set Vote Curves from Training Sets with Various Skew. Galactic plane sources and artifacts around certain bright sources have been excluded. (b) Vote curve comparison of skew 51147:48 training set and its corresponding application set. Again galactic plane sources and artifacts around certain bright sources have been excluded from the application set. The areas shown are for the total area under the respective vote curve. The expected ideal vote curve for the population is also shown.




Fig. 6.— Giant Radio Sources and Candidates with IAU designation and Table 6 index. These include some higher-ranked systems with ambiguous morphology as well as resolved core, core-jets and presumed chance projections, retained for documentation purposes. Un-

Table 1. List of Features for GRS Pattern Recognition

Feature	Description
$silsz_l$	silhouette size, component with larger silhouette size
$silsz_s$	silhouette size, component with smaller silhouette size
flx_l	peak flux, component with larger silhouette size
flx_s	peak flux, component with smaller silhouette size
d	source pair separation
$N_{neighbors}$	Number of sources within 20' of larger silhouette-size source having silhouette size larger than 1.871 square arc minutes

Note. — The silhouette size is the area enclosed by the fitted model at the 1.4 mJy level.

Table 2. Contingency Table - Two Class Problem

Positive Prediction	Negative Prediction	Totals
True Positive (TP)	False Negative (FN)	Positive Class (POS)
False Positive (FP)	True Negative (TN)	Negative Class (NEG)
Predicted Positive (PP)	Predicted Negative (PN)	N_{ts}

Table 3. Synthetic Training Sets Compared with Complete Training Set

Training Set Skew	AUC ^a	mean ^b	maximum ^c
48:48	0.920 ± 0.030	0.862 ± 0.038	0.9760 ± 0.0002
1000:48	0.920 ± 0.027	0.682 ± 0.055	0.9688 ± 0.0004
5000:48	0.928 ± 0.026	0.474 ± 0.058	0.9601 ± 0.0006
17000:48	0.957 ± 0.020	0.330 ± 0.053	0.9588 ± 0.0010
34000:48	0.963 ± 0.019	0.296 ± 0.045	0.9484 ± 0.0007
51147:48	0.963 ± 0.019	0.230 ± 0.038	0.9269 ± 0.0057

Note. — Options: best first order of coefficient perturbation, 40 hyper-planes searched per node, no pruning, variance impurity measure

^aArea under ROC curve with standard error

^bUnweighted mean and standard deviation of mean for training set GRS

^cMean of maximum vote with standard deviation of mean

Table 4. Impurity Measure Comparisons

impurity measure	AUC ^a	mean ^b	maximum ^c
Flach	0.978 $\pm$ 0.015	0.191 $\pm$ 0.032	0.786 $\pm$ 0.083
Gini Index	0.965 $\pm$ 0.019	0.203 $\pm$ 0.032	0.898 $\pm$ 0.020
Variance	0.963 $\pm$ 0.019	0.230 $\pm$ 0.038	0.927 $\pm$ 0.006
Twoing Rule	0.962 $\pm$ 0.019	0.228 $\pm$ 0.042	0.929 $\pm$ 0.008
Hellinger	0.952 $\pm$ 0.021	0.313 $\pm$ 0.045	0.908 $\pm$ 0.021
Information Gain	0.935 $\pm$ 0.025	0.132 $\pm$ 0.022	0.563 $\pm$ 0.144

Note. — Options: best first order of coefficient perturbation, 40 hyperplanes searched per node, no pruning

^aArea under ROC curve with standard error

^bUnweighted mean and standard deviation of mean for training set GRS

^cMean of maximum vote with standard deviation of mean

Table 5. Order of Coefficient Perturbation Comparisons

Order of Coefficient Perturbations	AUC ^a	mean ^b	maximum ^c
best first	0.978 $\pm$ 0.015	0.191 $\pm$ 0.032	0.786 $\pm$ 0.083
sequential	0.966 $\pm$ 0.018	0.140 $\pm$ 0.025	0.602 $\pm$ 0.108
random (-R12)	0.942 $\pm$ 0.024	0.161 $\pm$ 0.026	0.606 $\pm$ 0.109

Note. — Options: 40 hyperplanes searched per node, no pruning, Flach impurity measure

^aArea under ROC curve with standard error

^bUnweighted mean and standard deviation of mean for training set GRS

^cMean of maximum vote with standard deviation of mean

Table 6. Giant Radio Sources and Candidates

Index ^a	Source Designation	R.A.(J2000)	Dec.(J2000)	coordinate type ^b	size (')	Annotations ^c
1	NVGRC J000042.9+121542	00 00 42.93	+12 15 42.4	v	10.0	d dd?
2	NVGRC J000106.4+340303	00 01 06.40	+34 03 03.0	c	5.0	t?
3	NVGRC J000333.3+514628	00 03 33.34	+51 46 28.9	c	7.0	t? asym core not apparent
4	NVGRC J000349.1-151734	00 03 49.19	-15 17 34.6	c	11.0	t
5	NVGRC J000449.3+124921	00 04 49.31	+12 49 21.5	v	7.0	mg d? 4' d with adj 4' d or irr?

Note. — Table 6 is published in its entirety in the electronic edition of this journal. A portion is shown here for guidance regarding its form and content.

Note. — Approximately two dozen candidates less than 4.0' were retained in the table for documentation purposes.

Note. — Coordinate based source designations are truncated, not rounded. Format=JHHMMSS.s+DDMMSS

^aFor easy reference, the index corresponds to numbered cutouts in Figure 6

^bCoordinate type c indicates system coordinate is a NVSS fitted catalog entry, v indicates visual estimate from symmetry considerations. It should be noted that only the coordinates associated with a NVSS catalog entry have the significance of the NVSS entry.

^cThe key to the annotations is given in Table A-I. Although the training set consisted of primarily FRI and FRII type sources, FR classification was not attempted. Though many of the larger 'd' and 't' annotated systems are clearly FRII, the smaller, less resolved sources were difficult to evaluate. It was felt such distinctions deserved more uniform analytic attention. Also note that other morphological types were retained for documentation purposes. GRS previously identified by Lara et al. (2001) [2001A&A...370..409L], Machalski et al. (2001) [2001A&A...371..445M], and Solovoyov & Verkhodanov (2014) [2014AstBu..69..141S], were retained but not given an NVGRC designation. Aside from the GRSs/GRGs of these three references and SNR identified by Green (2014,2016), no comprehensive attempt was made at cross-identifications. Coordinates are given as for other entries.

Title: A Selection of Giant Radio Sources from NVSS
 Author: Proctor D.D.
 Table: Giant Radio Sources and Candidates

Byte-by-byte Description of file: datafile6.txt

Bytes	Format	Units	Label	Explanations
1- 4	I4	---	Index	Index
8- 29	A22	---	Name	IAU Name (1)
33- 34	I2	h	RAh	Hour of Right Ascension (J2000)
36- 37	I2	min	RAm	Minute of Right Ascension (J2000)
39- 43	F5.2	s	RAs	Second of Right Ascension (J2000)
46- 46	A1	---	DE-	Sign of the Declination (J2000)
47- 48	I2	deg	DEd	Degree of Declination (J2000)
50- 51	I2	arcmin	DEm	Arcminute of Declination (J2000)
53- 56	F4.1	arcsec	DEs	Arcsecond of Declination (J2000)
58- 58	A1	---	Flag	System coordinate code (2)
60- 64	F5.1	arcmin	diam	System diameter, 1.4 MJy level (3)
71-188	A118	---	text	Annotations (4)

Note (1): Coordinate based source designations are truncated, not rounded.
 Format=JHHMMSS.s+DDMMSS.

Note (2): It should be noted that only the coordinates associated with a NVSS catalog entry have the significance of the NVSS entry.
 c = system coordinate is a NVSS fitted catalog entry;
 v = coordinate is visual estimate from symmetry considerations;

Note (3): Approximately two dozen candidates less than 4.0' were retained in the table for documentation purposes.

Note (4): Although the training set consisted of primarily FRI and FRII type sources, FR classification was not attempted. Though many of the larger 'd' and 't' annotated systems are clearly FRII, the smaller, less resolved sources were difficult to evaluate. It was felt such distinctions deserved more uniform analytic attention. Also note that other morphological types were retained for documentation purposes. GRS previously identified by Lara et al. (2001) [2001A&A...370..409L], Machalski et al. (2001) [2001A&A...371..445M], and Solov'yov & Verkhodanov (2014) [2014AstBu..69..141S], were retained but not given an NVGRC designation. Aside from the GRSs/GRGs of these three references and SNR identified by Green (2014,2016), no comprehensive attempt was made at cross-identifications. Coordinates are given as for other entries.

1	NVGR	J000042.9+121542	00 00 42.93	+12 15 42.4	v	10.0	d	dd?
2	NVGR	J000106.4+340303	00 01 06.40	+34 03 03.0	c	5.0	t?	
3	NVGR	J000333.3+514628	00 03 33.34	+51 46 28.9	c	7.0	t?	asym core not apparent
4	NVGR	J000349.1-151734	00 03 49.19	-15 17 34.6	c	11.0	t	
5	NVGR	J000449.3+124921	00 04 49.31	+12 49 21.5	v	7.0	mg	d? 4' d with adj 4' d or irr?
6	NVGR	J000622.1+263549	00 06 22.14	+26 35 49.8	v	5.0	d	dd?
7	NVGR	J000630.8+155000	00 06 30.89	+15 50 00.8	v	7.0	cp	elliptic with cj?
8	NVGR	J000730.5+175127	00 07 30.55	+17 51 27.4	v	5.0	mg	d?
9	NVGR	J000814.3-193947	00 08 14.36	-19 39 47.9	v	6.0	wat/nat?	sz?
10	NVGR	J001117.5+523204	00 11 17.53	+52 32 04.0	v	4.0	cp?	d? cj?
11	NVGR	J001118.9+321716	00 11 18.94	+32 17 16.9	c	9.0	wat	d
12	NVGR	J001258.3-232628	00 12 58.38	-23 26 28.9	v	5.5	d?	points
13	NVGR	J001458.8-391227	00 14 58.83	-39 12 27.8	c	10.0	rc	
14	NVGR	J001502.2+101054	00 15 02.23	+10 10 54.1	v	5.0	d	
15	NVGR	J001613.6-312312	00 16 13.65	-31 23 12.2	c	5.5	cj?	cp?
16	NVGR	J001636.8-382649	00 16 36.81	-38 26 49.0	v	5.0	sz?	wat?
17	NVGR	J001741.2-351539	00 17 41.26	-35 15 39.7	v	7.5	cp	
18	NVGR	J001742.3+082757	00 17 42.36	+08 27 57.8	c	4.5	d/t	
19			00 17 47.50	-22 23 11.3	v	9.5	d	dd? x? (Solov'yov & Verkhodanov(2014): J001748.5-222256)
20	NVGR	J001814.9+214134	00 18 14.91	+21 41 34.3	v	5.0	d	
21	NVGR	J001817.7-161233	00 18 17.77	-16 12 33.7	v	4.5	d?	
22	NVGR	J002011.3+000443	00 20 11.32	+00 04 43.7	v	4.5	sz?	cp?
23	NVGR	J002035.5-201649	00 20 35.52	-20 16 49.6	v	10.5	mg	w dd?
24	NVGR	J002045.0-375639	00 20 45.03	-37 56 39.2	v	6.5	sz?	shell fragment?
25	NVGR	J002104.4-190948	00 21 04.41	-19 09 48.5	v	6.5	sz?	d?
26	NVGR	J002221.8+234412	00 22 21.81	+23 44 12.4	c	7.0	wat?	core id amb
27	NVGR	J002523.3+072422	00 25 23.33	+07 24 22.6	v	5.0	d?	cp? points
28	NVGR	J002613.8+114352	00 26 13.83	+11 43 52.9	c	10.0	t	
29	NVGR	J002700.1-121829	00 27 00.19	-12 18 29.0	c	5.0	t	sz?
30	NVGR	J002728.9-301236	00 27 28.98	-30 12 36.8	c	5.5	t	sz? wat?
31	NVGR	J002824.7-341919	00 28 24.75	-34 19 19.8	v	7.5	cp	wat??
32	NVGR	J002828.1-002633	00 28 28.19	-00 26 33.8	v	4.5	d?	
33	NVGR	J002849.5+145853	00 28 49.57	+14 58 53.8	v	4.5	d?	

34	NVGR	J002956.9+140137	00 29 56.97	+14 01 37.9	v	5.5	cp point with elliptic
35	NVGR	J003014.1+112313	00 30 14.16	+11 23 13.1	v	5.0	wat?
36	NVGR	J003031.0+300202	00 30 31.08	+30 02 02.2	c	4.0	mg rc? may be larger
37	NVGR	J003049.2-030728	00 30 49.23	-03 07 28.2	v	4.0	d?
38	NVGR	J003130.0-112350	00 31 30.00	-11 23 50.0	v	4.0	t? (with cp?)
39	NVGR	J003147.6-151401	00 31 47.63	-15 14 01.7	v	4.5	mg d? wat/nat? cp?
40	NVGR	J003419.3+011857	00 34 19.34	+01 18 57.5	v	4.0	d
41	NVGR	J003525.6-053110	00 35 25.64	-05 31 10.8	v	7.0	mg wat?
42	NVGR	J003558.8-302237	00 35 58.89	-30 22 37.9	v	5.0	d dd?
43	NVGR	J003615.4+060918	00 36 15.47	+06 09 18.5	c	6.5	mg d? t? x? may be larger
44	NVGR	J003622.0-025825	00 36 22.02	-02 58 25.3	v	4.5	mg
45	NVGR	J003706.9+254102	00 37 06.99	+25 41 02.7	v	4.5	nat/wat?
46	NVGR	J003708.0+103930	00 37 08.04	+10 39 30.4	v	4.5	wat
47	NVGR	J003718.6+195841	00 37 18.69	+19 58 41.1	c	4.5	d t? no peak seen for core
48	NVGR	J003809.1+093606	00 38 09.11	+09 36 06.0	c	6.0	x? d
49	NVGR	J003821.5-185159	00 38 21.53	-18 51 59.8	v	4.0	d sz?
50	NVGR	J003939.8-130102	00 39 39.83	-13 01 02.1	c	13.0	mg unu complex region
51	NVGR	J004017.0+595948	00 40 17.04	+59 59 48.8	v	5.0	nat ring? int core id amb
52	NVGR	J004150.8+401110	00 41 50.88	+40 11 10.1	v	13.0	cp point and elliptic
53	NVGR	J004246.5+411610	00 42 46.56	+41 16 10.8	c	8.0	rc
54	NVGR	J004247.7-061457	00 42 47.72	-06 14 57.8	c	9.0	w dd? int core id amb
55	NVGR	J004340.1-295931	00 43 40.16	-29 59 31.6	v	6.5	d? elliptic
56	NVGR	J004516.3+535612	00 45 16.38	+53 56 12.5	v	4.0	wat?
57	NVGR	J004517.3+382512	00 45 17.39	+38 25 12.2	v	4.0	d? (with cp?) may be larger
58	NVGR	J004653.2+125528	00 46 53.25	+12 55 28.1	v	5.0	nat/wat? sz?
59	NVGR	J004733.1-251717	00 47 33.13	-25 17 17.1	c	20.0	rc
60	NVGR	J004830.4+432235	00 48 30.48	+43 22 35.1	v	9.0	cp
61			00 51 05.36	-20 28 47.6	c	9.5	t (Soloviyov & Verkhodanov(2014): J005108.0-202818)
62	NVGR	J005116.0-283144	00 51 16.04	-28 31 44.8	c	3.0	mg cj? (with cp?)
63	NVGR	J005130.0-282704	00 51 30.03	-28 27 04.6	v	8.0	cp mg
64			00 53 31.62	+40 31 22.1	c	9.5	t dd? 5 components with apparent core (Soloviyov & Verkhodanov(2014): J005331.6+403059)
65	NVGR	J005451.5+564842	00 54 51.53	+56 48 42.0	v	5.5	d? cp
66	NVGR	J005553.0+262650	00 55 53.07	+26 26 50.2	c	7.0	mg d? elliptic wat/nat? asym core id amb
67	NVGR	J005601.5-012034	00 56 01.57	-01 20 34.4	v	6.0	nat? sz? int similar adjacent system
68	NVGR	J005606.6+273314	00 56 06.65	+27 33 14.2	v	7.0	cp?
69	NVGR	J005615.3+490757	00 56 15.36	+49 07 57.5	v	4.0	d
70	NVGR	J005620.3-093623	00 56 20.37	-09 36 23.5	c	4.5	cj? rc?
71	NVGR	J005626.5-011558	00 56 26.51	-01 15 58.5	v	9.0	nat? sz? int similar adjacent system
72	NVGR	J005644.9-173840	00 56 44.94	-17 38 40.5	v	4.0	d (with cp?)
73	NVGR	J005655.9-115529	00 56 55.98	-11 55 29.9	v	4.5	nat/wat?
74	NVGR	J005700.9-201841	00 57 00.96	-20 18 41.5	v	5.0	d? points
75	NVGR	J005734.8-012332	00 57 34.86	-01 23 32.5	v	4.0	rc d SN
76			00 57 48.34	+30 21 14.4	c	34.0	nat? dl int (Soloviyov & Verkhodanov(2014): J005744.4+302156)
77	NVGR	J005824.7+265200	00 58 24.77	+26 52 00.5	v	6.0	w? sz?
78	NVGR	J005901.2-152116	00 59 01.29	-15 21 16.2	v	5.5	d? points
79	NVGR	J005938.1-124712	00 59 38.11	-12 47 12.4	v	4.0	wat/nat? sz? dl? cp?
80	NVGR	J005956.2-224747	00 59 56.21	-22 47 47.5	v	4.5	mg wat?
81	NVGR	J010001.3+300249	01 00 01.36	+30 02 49.9	v	8.0	rc? lobe of large nat/wat to west?
82	NVGR	J010051.6+061654	01 00 51.66	+06 16 54.3	c	6.0	sz
83	NVGR	J010144.4-025246	01 01 44.45	-02 52 46.4	c	9.5	wat/nat w? asym lobes?
84	NVGR	J010150.7-064803	01 01 50.76	-06 48 03.3	c	17.5	wat
85	NVGR	J010155.7+505259	01 01 55.72	+50 52 59.0	v	7.5	sz? (with cp?) w? int
86	NVGR	J010236.7+042049	01 02 36.77	+04 20 49.2	c	4.5	d (with cp?)
87	NVGR	J010326.6+084950	01 03 26.69	+08 49 50.5	v	7.0	mg wat?
88	NVGR	J010340.2+423912	01 03 40.29	+42 39 12.2	v	7.0	mg sz? cp? core id amb
89	NVGR	J010356.7+320627	01 03 56.72	+32 06 27.4	v	6.0	nat/wat? sz?
90	NVGR	J010535.5+283701	01 05 35.59	+28 37 01.8	v	4.0	d wat?
91			01 07 24.75	+32 25 12.6	v	18.0	w (Soloviyov & Verkhodanov(2014): J010725.4+322439)
92	NVGR	J010803.5+270001	01 08 03.50	+27 00 01.5	c	5.5	d t sz? core fit but not apparent
93	NVGR	J010854.6+132016	01 08 54.67	+13 20 16.0	v	6.5	wat? rc x? dd?
94	NVGR	J010936.1+252346	01 09 36.19	+25 23 46.0	c	5.0	d/t
95			01 09 39.85	+73 11 45.8	v	5.5	d (Lara et al.(2001): J0109+731)
96	NVGR	J011105.6-142245	01 11 05.61	-14 22 45.9	c	6.0	rc? t?
97			01 12 02.54	+49 28 41.9	c	13.0	t int gap around core (Soloviyov & Verkhodanov(2014): J011203.6+493004)
98	NVGR	J011258.6+152932	01 12 58.68	+15 29 32.9	v	10.0	wat/nat? w? 7 components
99			01 13 08.25	-17 19 32.2	c	7.5	sz t (Soloviyov & Verkhodanov(2014): J011306.9-1720280)
100	NVGR	J011341.8+010555	01 13 41.82	+01 05 55.9	v	4.5	mg x? cp?
101	NVGR	J011352.3+622434	01 13 52.32	+62 24 34.6	v	5.0	d
102	NVGR	J011429.7+050814	01 14 29.78	+05 08 14.2	v	7.0	d? (with tail?)
103	NVGR	J011453.1-223052	01 14 53.11	-22 30 52.9	v	5.0	cp d?
104	NVGR	J011544.9-241712	01 15 44.98	-24 17 12.4	c	3.5	cj? d?
105	NVGR	J011557.3+250720	01 15 57.32	+25 07 20.9	c	8.0	t
106	NVGR	J011800.8-364011	01 18 00.82	-36 40 11.1	v	5.5	mg int quad
107	NVGR	J011822.4+471213	01 18 22.48	+47 12 13.9	v	4.0	d (dl or with cp?)
108	NVGR	J012024.7+430950	01 20 24.75	+43 09 50.2	v	6.5	cp d? asym. one is arc/crescent
109	NVGR	J012041.6-170121	01 20 41.64	-17 01 21.8	v	11.0	wat d dd ring lobe
110	NVGR	J012317.0+485812	01 23 17.09	+48 58 12.2	v	6.0	wat/nat?
111	NVGR	J012322.5+014559	01 23 22.56	+01 45 59.1	v	10.0	rc?

112	NVGR	J012359.4+431259	01 23 59.40	+43 12 59.8	v	6.0	d (with cp?)
113	NVGR	J012400.4+805514	01 24 00.42	+80 55 14.5	c	4.5	wat? sz? core not apparent
114	NVGR	J012512.3+581040	01 25 12.39	+58 10 40.5	v	11.0	d
115	NVGR	J012532.2+070338	01 25 32.26	+07 03 38.3	c	11.0	nat?
116	NVGR	J012600.6-012052	01 26 00.61	-01 20 52.5	c	18.5	w? wat? may be larger
117	NVGR	J012718.4-041406	01 27 18.48	-04 14 06.5	c	4.5	d t
118	NVGR	J012848.8+243143	01 28 48.88	+24 31 43.8	c	5.5	t
119	NVGR	J013327.2-082420	01 33 27.28	-08 24 20.9	c	6.0	wat/nat? t?
120	NVGR	J013354.4-362914	01 33 54.42	-36 29 14.6	v	14.0	sz NGC 612
121	NVGR	J013413.9-010606	01 34 13.91	-01 06 06.8	v	12.0	wat? w?
122	NVGR	J013419.2+143014	01 34 19.20	+14 30 14.5	v	5.0	mg cp d?
123	NVGR	J013448.8+080047	01 34 48.85	+08 00 47.0	c	4.5	cj may be larger
124	NVGR	J013523.8-051640	01 35 23.86	-05 16 40.1	v	4.0	rc
125	NVGR	J013525.5-004436	01 35 25.59	-00 44 36.0	v	8.0	t?
126	NVGR	J013536.4+504036	01 35 36.40	+50 40 36.9	c	5.0	wat?
127	NVGR	J013615.5-381433	01 36 15.56	-38 14 33.7	v	7.0	mg cp two non-aligned elliptic
128	NVGR	J013847.4+472315	01 38 47.48	+47 23 15.1	v	6.0	d? wat?
129	NVGR	J013927.2+395654	01 39 27.25	+39 56 54.6	c	7.0	mg t
130	NVGR	J013956.3+230716	01 39 56.33	+23 07 16.4	c	9.0	wat?
131	NVGR	J014207.6-064148	01 42 07.66	-06 41 48.2	v	6.5	sz? wat?
132	NVGR	J014259.4+313833	01 42 59.44	+31 38 33.6	v	4.0	d
133	NVGR	J014356.0+062443	01 43 56.00	+06 24 43.9	c	7.0	wat t
134	NVGR	J014442.6+424417	01 44 42.61	+42 44 17.1	v	4.5	d sz?
135	NVGR	J014715.6+173725	01 47 15.68	+17 37 25.3	v	5.5	d? cp? int cluster of 'point' sources
136	NVGR	J014849.5+062258	01 48 49.58	+06 22 58.0	v	5.0	d
137	NVGR	J014942.8-133220	01 49 42.81	-13 32 20.6	v	4.0	d? points
138	NVGR	J014943.7-074226	01 49 43.72	-07 42 26.9	c	6.5	wat t?
139	NVGR	J015006.1-164043	01 50 06.11	-16 40 43.4	c	7.0	mg t? d+s?
140	NVGR	J015012.4-304151	01 50 12.46	-30 41 51.0	v	4.5	d?
141	NVGR	J015124.9-111202	01 51 24.95	-11 12 02.5	c	13.5	w?
142	NVGR	J015151.5-254704	01 51 51.52	-25 47 04.1	v	4.5	mg
143	NVGR	J015200.8-303317	01 52 00.84	-30 33 17.1	c	5.5	t? mg
144			01 53 02.45	+71 15 59.2	v	7.5	w? fades (Lara et al.(2001): J0153+712, Soloviyov & Verkhodanov(2014): J015302.5+711559)
145	NVGR	J015622.1+053739	01 56 22.14	+05 37 39.8	c	9.0	w? wat?
146			01 57 56.16	+02 10 17.5	c	9.0	x core id amb (Soloviyov & Verkhodanov(2014): J015756.3+020950)
147	NVGR	J015825.8+245140	01 58 25.88	+24 51 40.6	c	5.0	t sz?
148	NVGR	J015837.5+100452	01 58 37.54	+10 04 52.1	v	5.0	d?
149	NVGR	J020030.0+404900	02 00 30.00	+40 49 00.0	v	14.0	d?
150	NVGR	J020052.5+393439	02 00 52.57	+39 34 39.5	v	5.5	nat/wat arc
151	NVGR	J020211.1+152603	02 02 11.16	+15 26 03.1	v	5.5	mg wat/nat? (with rc?)
152	NVGR	J020320.9+431647	02 03 20.94	+43 16 47.8	v	4.0	d? points
153	NVGR	J020504.6+230517	02 05 04.61	+23 05 17.6	c	5.0	cj? rc?
154	NVGR	J020710.1+533901	02 07 10.14	+53 39 01.9	v	10.0	d
155	NVGR	J020813.3+273152	02 08 13.39	+27 31 52.6	c	6.0	t wat?
156	NVGR	J020822.3-320219	02 08 22.32	-32 02 19.9	c	10.0	sz t
157	NVGR	J020851.4-151354	02 08 51.47	-15 13 54.0	v	6.0	d wat? cp?
158	NVGR	J020942.4+150027	02 09 42.45	+15 00 27.3	c	6.5	mg wat? core id amb
159	NVGR	J021201.3-241152	02 12 01.34	-24 11 52.5	c	7.0	wat
160	NVGR	J021329.0+292139	02 13 29.05	+29 21 39.0	c	4.5	wat mg
161	NVGR	J021413.0+325053	02 14 13.04	+32 50 53.6	c	7.0	t
162	NVGR	J021427.6-140547	02 14 27.62	-14 05 47.9	v	8.0	d dd? resolved lobes
163	NVGR	J021625.1-024339	02 16 25.19	-02 43 39.0	v	7.0	nat/wat?
164	NVGR	J021717.3-145640	02 17 17.34	-14 56 40.7	v	6.0	d? cp? wat??
165	NVGR	J021837.3+530405	02 18 37.37	+53 04 05.3	v	7.5	d
166	NVGR	J021924.9+124620	02 19 24.92	+12 46 20.1	v	6.0	mg d? sz? w? (with cp's?)
167			02 20 22.95	+36 59 40.1	c	14.0	w t? dd? irr core id amb (Soloviyov & Verkhodanov(2014): J022033.5+365943)
168	NVGR	J022251.6+425744	02 22 51.60	+42 57 44.9	c	7.0	t ring-lobe?
169	NVGR	J022304.2-211400	02 23 04.23	-21 14 00.4	c	5.0	cj
170	NVGR	J022318.0+425939	02 23 18.00	+42 59 39.0	v	7.0	wat? rc?
171	NVGR	J022409.6+500700	02 24 09.65	+50 07 00.0	v	7.0	cp point with 3.5' arc/crescent
172	NVGR	J022526.9-313740	02 25 26.98	-31 37 40.1	c	5.5	t? cp?
173	NVGR	J022617.3-160108	02 26 17.39	-16 01 08.1	c	8.0	wat
174	NVGR	J022821.0-243351	02 28 21.06	-24 33 51.9	v	5.0	cp d? point and elliptic
175	NVGR	J022825.5-281405	02 28 25.52	-28 14 05.5	v	4.0	cp? nat?
176	NVGR	J023002.5+423415	02 30 02.50	+42 34 15.5	v	6.0	d? cp? mg?
177	NVGR	J023125.9-173914	02 31 25.98	-17 39 14.5	c	6.5	t wat/nat?
178	NVGR	J023137.1-204020	02 31 37.17	-20 40 20.4	c	5.5	nat/wat? t PKS 0229-208
179	NVGR	J023300.0-152100	02 33 00.00	-15 21 00.0	v	6.5	d? points
180	NVGR	J023413.9+312928	02 34 13.91	+31 29 28.1	v	5.5	d (teardrop shape lobes) 4 components
181	NVGR	J023432.1+471211	02 34 32.19	+47 12 11.2	c	5.0	rc w?
182	NVGR	J023830.5+023325	02 38 30.59	+02 33 25.1	v	8.5	w? nat? x? cp?
183	NVGR	J024024.4-162848	02 40 24.41	-16 28 48.0	v	6.0	cp d? point and elliptic
184	NVGR	J024104.9+084439	02 41 04.95	+08 44 39.3	v	13.0	w
185	NVGR	J024146.4-283954	02 41 46.46	-28 39 54.5	c	6.5	wat/nat cp?
186	NVGR	J024148.5-312214	02 41 48.55	-31 22 14.1	c	6.0	mg nat/wat? d?
187	NVGR	J024155.0+593613	02 41 55.03	+59 36 13.9	c	9.0	rc sz? unu wispy/clumpy
188	NVGR	J024429.2-254751	02 44 29.28	-25 47 51.6	v	5.0	cp d? point and elliptic
189	NVGR	J024432.8+393412	02 44 32.88	+39 34 12.5	v	4.5	d rc

190	NVGR	J024435.3-052610	02 44 35.30	-05 26 10.0	v	5.0	mg
191	NVGR	J024618.9-301629	02 46 18.93	-30 16 29.3	c	7.0	rc
192	NVGR	J024636.3+294306	02 46 36.39	+29 43 06.1	v	3.5	d
193	NVGR	J024733.6+615632	02 47 33.60	+61 56 32.5	v	7.0	cp? unu trough/cleared area, ring?
194	NVGR	J024738.1-383806	02 47 38.10	-38 38 06.2	v	5.0	d? points
195	NVGR	J024743.0-362400	02 47 43.00	-36 24 00.0	v	7.0	d? points
196	NVGR	J024811.4-041611	02 48 11.40	-04 16 11.3	v	4.5	rc? lobe? fades
197	NVGR	J024853.7-313231	02 48 53.75	-31 32 31.5	v	5.0	mg point and elliptic
198	NVGR	J024909.7+182003	02 49 09.78	+18 20 03.5	c	11.0	wat/nat beautiful 7 or more components
199	NVGR	J024930.4+430525	02 49 30.48	+43 05 25.6	v	5.0	d
200	NVGR	J024934.1-203047	02 49 34.19	-20 30 47.6	v	11.0	w? wat? sz?
201	NVGR	J025048.9+084313	02 50 48.97	+08 43 13.1	v	4.0	d
202	NVGR	J025100.5+465724	02 51 00.55	+46 57 24.0	c	15.0	w? nat/wat? sz? irr int core id amb
203	NVGR	J025102.8+261412	02 51 02.85	+26 14 12.9	v	3.5	d
204	NVGR	J025135.9+373146	02 51 35.92	+37 31 46.7	c	6.0	t fades
205	NVGR	J025204.1+381122	02 52 04.18	+38 11 22.1	c	4.5	cj? rc?
206	NVGR	J025209.4+025418	02 52 09.43	+02 54 18.1	c	5.0	t
207	NVGR	J025347.1-200007	02 53 47.15	-20 00 07.6	v	4.0	d? cp?
208	NVGR	J025739.9+060317	02 57 39.91	+06 03 17.3	v	12.5	w? nat/wat?
209	NVGR	J025831.8+133417	02 58 31.89	+13 34 17.4	c	5.5	cj fades
210	NVGR	J025847.9-263432	02 58 47.90	-26 34 32.6	v	11.0	cp d with elliptic
211	NVGR	J025926.4+355632	02 59 26.46	+35 56 32.7	v	6.5	wat
212	NVGR	J030039.8+362145	03 00 39.85	+36 21 45.8	v	5.0	d? points
213	NVGR	J030209.9+434229	03 02 09.93	+43 42 29.5	c	7.0	cj arc
214	NVGR	J030227.8+423300	03 02 27.88	+42 33 00.5	v	5.0	d dd?
215	NVGR	J030251.7+015456	03 02 51.73	+01 54 56.8	v	7.5	d?
216	NVGR	J030304.1-341413	03 03 04.18	-34 14 13.4	v	8.5	cp two elliptics
217	NVGR	J030314.5+162615	03 03 14.59	+16 26 15.3	v	5.0	rc d?
218	NVGR	J030826.2+040642	03 08 26.26	+04 06 42.8	c	5.0	rc NGC 1218
219	NVGR	J030954.7+191410	03 09 54.71	+19 14 10.1	v	7.5	d? asym one lobe wispy? or ring? int
220	NVGR	J031013.6+425002	03 10 13.64	+42 50 02.6	v	7.0	wat? t?
221	NVGR	J031242.8-363314	03 12 42.81	-36 33 14.8	c	5.0	rc sz?
222	NVGR	J031334.9-063204	03 13 34.97	-06 32 04.5	c	5.0	d x?
223	NVGR	J031400.3+202106	03 14 00.38	+20 21 06.8	v	6.0	d 4 components sz? x??
224	NVGR	J031447.9+150830	03 14 47.91	+15 08 30.4	c	8.0	wat/nat
225	NVGR	J031602.3-265738	03 16 02.38	-26 57 38.7	c	6.5	sz? d? t?
226	NVGR	J031633.0+411710	03 16 33.01	+41 17 10.1	v	10.0	w? d with cp? fades int
227			03 17 47.74	+76 57 22.7	v	5.0	mg cp? (Lara et al.(2001): J0317+769)
228	NVGR	J031817.7+415211	03 18 17.74	+41 52 11.5	c	11.5	nat? unu core id amb
229			03 18 18.87	+68 29 27.3	c	16.5	t x? (Lara et al.(2001): J0318+684, Solov'yov & Verkhodanov(2014): J031821.9+682932)
230	NVGR	J031931.6-211446	03 19 31.64	-21 14 46.2	v	4.0	wat? cp?
231	NVGR	J031948.1+413042	03 19 48.16	+41 30 42.1	c	9.0	rc 3C 84
232	NVGR	J032009.8-105539	03 20 09.84	-10 55 39.2	c	4.5	d t
233	NVGR	J032031.0+304248	03 20 31.07	+30 42 48.4	v	4.0	wat? cp?
234	NVGR	J032040.9+272748	03 20 40.97	+27 27 48.8	v	4.0	mg d with cp? w? sz? may be larger
235	NVGR	J032101.8-370927	03 21 01.85	-37 09 27.0	v	14.0	wat? artifact? apx. center of ~0.75 degree disturbance SN 1965J, NGC 1310 within 3.5'
236	NVGR	J032126.1+180559	03 21 26.13	+18 05 59.4	c	12.0	w wat/nat? int
237	NVGR	J032145.1+514855	03 21 45.11	+51 48 55.9	v	5.0	d
238	NVGR	J032156.0+434635	03 21 56.09	+43 46 35.1	v	5.0	d
239	NVGR	J032247.3+210749	03 22 47.35	+21 07 49.7	c	5.0	t
240	NVGR	J032252.5+233414	03 22 52.59	+23 34 14.8	v	4.0	d
241	NVGR	J032313.1+554226	03 23 13.16	+55 42 26.5	v	7.0	mg cj? t? asym hymors?
242	NVGR	J032533.6-392728	03 25 33.62	-39 27 28.2	c	13.0	wat
243	NVGR	J032625.3+094018	03 26 25.30	+09 40 18.4	c	8.0	wat/nat? unu core id amb
244	NVGR	J032636.9+023254	03 26 36.98	+02 32 54.4	v	6.5	d? cj? wat? w??
245	NVGR	J032638.5+415209	03 26 38.58	+41 52 09.8	v	4.0	d? cj?
246	NVGR	J032754.4+023341	03 27 54.44	+02 33 41.8	v	6.0	d sz? rc?
247	NVGR	J032758.0+382209	03 27 58.04	+38 22 09.1	c	6.0	sz?
248	NVGR	J032923.8+394732	03 29 23.89	+39 47 32.6	c	8.0	t sz?
249	NVGR	J033014.1-365309	03 30 14.19	-36 53 09.4	v	4.5	d
250	NVGR	J033018.5+573528	03 30 18.55	+57 35 28.7	v	6.0	mg cp? d?
251	NVGR	J033406.1-390058	03 34 06.16	-39 00 58.4	c	8.5	wat/nat core id amb
252	NVGR	J033418.4+392122	03 34 18.46	+39 21 22.0	c	6.5	rc?
253	NVGR	J033441.6-104328	03 34 41.61	-10 43 28.0	v	5.0	cp? d?
254	NVGR	J033524.1+550328	03 35 24.13	+55 03 28.0	c	7.0	wat? sz? t dl
255	NVGR	J033600.3+473536	03 36 00.35	+47 35 36.3	v	7.5	w? sz? 4 or more components
256	NVGR	J033737.8+330928	03 37 37.83	+33 09 28.7	c	8.0	t
257	NVGR	J033829.0-352700	03 38 29.02	-35 27 00.7	c	6.0	t
258	NVGR	J033845.9-352255	03 38 45.96	-35 22 55.0	v	4.0	d - incidental
259	NVGR	J033919.1+011548	03 39 19.16	+01 15 48.9	c	5.0	d
260	NVGR	J033931.3-181646	03 39 31.33	-18 16 46.2	v	6.5	cp elliptic with cj?
261	NVGR	J034015.7-015338	03 40 15.70	-01 53 38.0	v	4.0	sz d
262	NVGR	J034141.4-310427	03 41 41.49	-31 04 27.6	v	6.0	cp
263			03 42 09.26	+63 39 28.3	c	6.0	t core asymmetric position (Lara et al.(2001): J0342+636)
264	NVGR	J034253.9-065225	03 42 53.91	-06 52 25.3	c	4.5	d t
265	NVGR	J034346.8+231852	03 43 46.83	+23 18 52.7	c	5.0	wat? sz? core id amb
266	NVGR	J034419.8+510224	03 44 19.88	+51 02 24.8	v	12.0	wat cp? GRS or two
267	NVGR	J034629.7-342232	03 46 29.77	-34 22 32.0	v	7.5	rc d? x?

268	NVGR	J034901.6-385427	03 49 01.64	-38 54 27.8 v	7.0	wat?
269	NVGR	J034909.7+053418	03 49 09.74	+05 34 18.6 v	5.5	wat/nat? cp?
270	NVGR	J035052.1-014531	03 50 52.14	-01 45 31.4 c	9.0	t
271	NVGR	J035129.2-354258	03 51 29.20	-35 42 58.0 v	5.5	d sp?
272	NVGR	J035133.3-274500	03 51 33.33	-27 45 00.2 v	8.0	w? x?
273	NVGR	J035140.7-045620	03 51 40.73	-04 56 20.9 c	6.0	t? cp?
274	NVGR	J035322.1+355212	03 53 22.16	+35 52 12.0 v	6.5	mg d/sz? dl? (with cp?)
275	NVGR	J035339.2-011319	03 53 39.23	-01 13 19.8 c	6.0	wat?
276	NVGR	J035402.7-083431	03 54 02.76	-08 34 31.6 c	8.0	wat? ring? cp? core id amb
277	NVGR	J035422.2-383228	03 54 22.26	-38 32 28.5 v	7.5	wat? sz?
278	NVGR	J035434.7-151405	03 54 34.78	-15 14 05.2 v	5.0	d asym
279	NVGR	J035648.8-342654	03 56 48.83	-34 26 54.0 v	8.0	cp wat?
280	NVGR	J035650.9+001343	03 56 50.92	+00 13 43.1 v	4.0	d (with cp?)
281	NVGR	J035715.2+375653	03 57 15.23	+37 56 53.9 v	5.0	sz? rc?
282	NVGR	J035742.3+455758	03 57 42.33	+45 57 58.7 v	7.5	wat possible core at J000 03 57 38.59 +45 56 59.4
283	NVGR	J035743.2-305225	03 57 43.27	-30 52 25.5 v	4.0	d
284	NVGR	J035800.3-393629	03 58 00.39	-39 36 29.3 v	8.0	cp
285	NVGR	J035834.5+405544	03 58 34.56	+40 55 44.2 v	5.0	d? cp? both comp near pointlike
286	NVGR	J040023.2+314725	04 00 23.27	+31 47 25.6 v	6.0	d
287	NVGR	J040045.5+261910	04 00 45.54	+26 19 10.1 v	4.5	d sz? rc?
288	NVGR	J040052.5+344247	04 00 52.53	+34 42 47.6 v	5.0	wat/nat dl?
289	NVGR	J040108.5-253104	04 01 08.52	-25 31 04.2 v	5.0	mg d wat/nat? cp?
290	NVGR	J040310.8+342907	04 03 10.86	+34 29 07.5 v	6.0	d? (with cp?) asym
291	NVGR	J040409.8+151635	04 04 09.88	+15 16 35.2 c	5.0	t
292	NVGR	J040621.0+205622	04 06 21.07	+20 56 22.6 v	5.0	d
293	NVGR	J040650.7-064704	04 06 50.70	-06 47 04.9 v	5.0	cp? cj?
294	NVGR	J040651.0+094700	04 06 51.03	+09 47 00.5 c	6.0	sz
295	NVGR	J040715.8+034239	04 07 15.86	+03 42 39.4 c	8.0	wat/nat? dd? core id amb
296	NVGR	J040742.5+491050	04 07 42.59	+49 10 50.8 v	5.0	d
297	NVGR	J040905.8-052934	04 09 05.81	-05 29 34.8 v	5.5	cp
298	NVGR	J041203.9-325221	04 12 03.98	-32 52 21.0 v	5.5	rc? sz?
299	NVGR	J041524.2+422020	04 15 24.20	+42 20 20.5 v	4.0	d
300	NVGR	J041543.7+331413	04 15 43.77	+33 14 13.1 v	5.0	cp
301	NVGR	J041820.9+380148	04 18 20.96	+38 01 48.6 c	6.0	rc? t?
302	NVGR	J042109.0+260038	04 21 09.02	+26 00 38.1 v	6.0	d with tail
303	NVGR	J042115.9-295350	04 21 15.91	-29 53 50.6 v	4.5	cp
304	NVGR	J042220.9+151101	04 22 20.97	+15 11 01.4 c	14.5	d dd? dumbell int 'spherical' lobes at end of jets int grayscale wispy ring-lobes
305	NVGR	J042228.9+140748	04 22 28.95	+14 07 48.7 c	8.5	sz t core
306	NVGR	J042234.6-261606	04 22 34.62	-26 16 06.5 v	6.0	rc? t? irr (with cp?)
307	NVGR	J042340.7+324831	04 23 40.73	+32 48 31.4 v	5.5	cp d?
308	NVGR	J042418.7-274352	04 24 18.73	-27 43 52.5 v	5.0	d sp?
309	NVGR	J042538.6+194303	04 25 38.64	+19 43 03.1 c	5.0	nat?
310	NVGR	J042717.6+463415	04 27 17.64	+46 34 15.5 v	5.5	sz? cp?
311	NVGR	J042828.9+114627	04 28 28.90	+11 46 27.5 v	4.0	wat? (with cp?)
312	NVGR	J042857.2-031422	04 28 57.22	-03 14 22.1 v	5.0	cp
313	NVGR	J043025.3+242053	04 30 25.35	+24 20 53.1 c	4.0	rc? cj? d? asym
314	NVGR	J043145.5+082950	04 31 45.55	+08 29 50.6 v	5.5	d? point and elliptic
315	NVGR	J043308.6+060202	04 33 08.68	+06 02 02.3 v	5.0	dd? with cp? pair of doubles? sz?
316	NVGR	J043311.0+052115	04 33 11.03	+05 21 15.4 c	9.0	rc? t? sz? cp? irr int
317	NVGR	J043406.0-245534	04 34 06.02	-24 55 34.1 v	5.5	d? points
318	NVGR	J043409.1-132219	04 34 09.15	-13 22 19.5 c	9.0	nat? rc? with tail, 6 components core id amb
319	NVGR	J043503.2+215527	04 35 03.26	+21 55 27.8 c	5.0	d t?
320	NVGR	J043526.4-192503	04 35 26.44	-19 25 03.1 v	4.0	d t? - incidental
321	NVGR	J043558.1-233319	04 35 58.13	-23 33 19.8 c	8.5	t dumbell irr lobes hymors? ring-lobe?
322	NVGR	J043637.2-222629	04 36 37.22	-22 26 29.4 v	4.5	rc? d? with low-level 'hook' (or cp?)
323	NVGR	J043702.4+160459	04 37 02.41	+16 04 59.4 v	6.5	nat/wat?
324	NVGR	J043755.6-374050	04 37 55.62	-37 40 50.7 v	5.5	d rc? x?
325	NVGR	J043838.3+142422	04 38 38.38	+14 24 22.5 v	6.0	d? points - int cluster pattern of points
326	NVGR	J043857.8+095336	04 38 57.83	+09 53 36.7 c	9.5	t? wat? core id amb hymors? int
327	NVGR	J043927.3-233029	04 39 27.33	-23 30 29.2 v	10.0	d?
328	NVGR	J043956.9-112205	04 39 56.97	-11 22 05.9 v	5.5	d dd?
329	NVGR	J044024.6-080349	04 40 24.60	-08 03 49.3 v	8.0	cp d with elliptic
330	NVGR	J044111.2+251841	04 41 11.22	+25 18 41.6 c	11.0	rc sz
331	NVGR	J044152.7+082738	04 41 52.75	+08 27 38.0 v	4.5	d
332	NVGR	J044646.2+450144	04 46 46.22	+45 01 44.6 c	13.0	wat? artifact?
333	NVGR	J044805.1-202609	04 48 05.11	-20 26 09.9 v	5.5	cp?
334	NVGR	J044821.2+450201	04 48 21.25	+45 02 01.2 c	16.0	wat? w? ring fragment? with possible sidelobes int, core id amb, GRS or two?
335	NVGR	J045006.9+450315	04 50 06.91	+45 03 15.8 c	13.0	w? artifact?
336	NVGR	J045035.0+201055	04 50 35.00	+20 10 55.0 v	3.5	d? rc? (not original pair)
337	NVGR	J045058.9+224931	04 50 58.97	+22 49 31.0 v	5.5	cp? point and elliptic
338	NVGR	J045122.7-173047	04 51 22.71	-17 30 47.1 c	6.5	wat? rc? 5-components core id amb
339	NVGR	J045236.5+025212	04 52 36.59	+02 52 12.0 v	5.0	d? point-like
340	NVGR	J045250.8+520444	04 52 50.84	+52 04 44.1 v	10.0	w
341	NVGR	J045334.0-182218	04 53 34.05	-18 22 18.9 v	4.0	d t?
342	NVGR	J045346.5-182250	04 53 46.50	-18 22 50.0 v	3.5	d?
343	NVGR	J045359.4+120319	04 53 59.42	+12 03 19.1 v	5.0	d? points
344	NVGR	J045454.9+130208	04 54 54.94	+13 02 08.9 v	6.0	d 4 components inner pair sources bridge
345	NVGR	J045616.3-260137	04 56 16.34	-26 01 37.6 v	5.0	d asym

346	NVGR	J045742.1+062144	04 57 42.19	+06 21 44.1	v	5.0	d? cp? sz?
347	NVGR	J045935.0+283730	04 59 35.01	+28 37 30.5	v	7.0	cp
348	NVGR	J050022.1-244926	05 00 22.13	-24 49 26.7	c	4.0	t
349	NVGR	J050208.0+244229	05 02 08.01	+24 42 29.5	v	7.0	d? points
350	NVGR	J050222.5+244139	05 02 22.54	+24 41 39.5	v	5.5	d? points
351			05 02 53.32	+67 02 37.0	c	6.0	wat? w? (Lara et al.(2001): J0502+670)
352	NVGR	J050323.6+422804	05 03 23.60	+42 28 04.2	v	5.0	mg d (with cp?) all points
353	NVGR	J050341.2-191142	05 03 41.26	-19 11 42.8	v	4.0	d
354	NVGR	J050533.7-285707	05 05 33.75	-28 57 07.3	c	11.0	rc
355	NVGR	J050540.8-282445	05 05 40.86	-28 24 45.2	v	8.0	rc int ring? nat? in processing depression?
356	NVGR	J050717.5+305759	05 07 17.55	+30 57 59.0	c	8.0	mg w? wat? core id amb, possibly at 05 07 14.09 +30 58 17.0
357			05 08 27.45	+60 56 25.5	c	12.0	sz? cp? (Lara et al.(2001): J0508+609)
358	NVGR	J051103.2-131642	05 11 03.20	-13 16 42.8	v	5.0	rc? d?
359	NVGR	J051219.4+131945	05 12 19.48	+13 19 45.0	v	7.0	mg cp wat?
360	NVGR	J051336.6-302741	05 13 36.60	-30 27 41.2	v	13.0	w? butterfly? cp? irr
361	NVGR	J051354.8+015556	05 13 54.83	+01 55 56.6	v	4.5	cp?
362	NVGR	J051422.5-165230	05 14 22.55	-16 52 30.9	v	19.5	d?
363	NVGR	J051513.5-125451	05 15 13.51	-12 54 51.7	v	5.0	sz? nat? cp?
364	NVGR	J051535.8+151825	05 15 35.83	+15 18 25.8	v	5.5	d
365	NVGR	J051601.7+245826	05 16 01.74	+24 58 26.8	v	11.0	x? int cp? this coord center rc N-S system, E-W system listed separately
366	NVGR	J051605.7+245833	05 16 05.73	+24 58 33.7	v	10.5	x? int cp? this coord approximate center E-W system, N-S system listed separately
367	NVGR	J051845.2+282635	05 18 45.28	+28 26 35.7	v	6.5	d? cp?
368	NVGR	J051858.4-162800	05 18 58.42	-16 28 00.1	v	11.0	d? dd? 5 components core id amb
369			05 19 18.35	+70 13 46.8	c	6.0	nat (Lara et al.(2001): J0519+702)
370	NVGR	J051933.5+274400	05 19 33.50	+27 44 00.2	c	5.0	rc
371	NVGR	J052106.2+124606	05 21 06.27	+12 46 06.3	v	4.5	d? cp?
372	NVGR	J052258.3-293555	05 22 58.32	-29 35 55.1	v	5.0	d? points
373	NVGR	J052308.4+595057	05 23 08.40	+59 50 57.9	v	6.5	mg cp?
374	NVGR	J052322.9+594847	05 23 22.99	+59 48 47.2	v	5.5	mg cp?
375	NVGR	J052527.7-201048	05 25 27.78	-20 10 48.7	c	8.0	wat/nat? with cp? irr int core id amb
376	NVGR	J052528.4-324235	05 25 28.42	-32 42 35.5	c	7.5	sz
377	NVGR	J052823.7-173553	05 28 23.72	-17 35 53.1	c	5.5	t? core id amb asym
378	NVGR	J052829.2-185814	05 28 29.28	-18 58 14.5	v	5.5	sz? d tail?
379	NVGR	J052859.2+124644	05 28 59.26	+12 46 44.3	v	7.5	wat? core id amb
380	NVGR	J052949.6-314929	05 29 49.67	-31 49 29.3	v	5.0	d
381	NVGR	J053013.5-270343	05 30 13.51	-27 03 43.3	v	5.0	wat? d? cp?
382	NVGR	J053023.7+155023	05 30 23.70	+15 50 23.5	c	4.0	wat
383	NVGR	J053554.1+220300	05 35 54.16	+22 03 00.0	v	8.0	d? artifact? incidental
384	NVGR	J053559.1-045532	05 35 59.14	-04 55 32.0	v	4.0	d? artifact?
385	NVGR	J053617.3+271504	05 36 17.32	+27 15 04.3	v	8.0	mg cp d?
386	NVGR	J053901.8-094009	05 39 01.88	-09 40 09.8	v	6.0	d? cp?
387	NVGR	J054005.4+195233	05 40 05.44	+19 52 33.7	v	8.0	d? points not orig pair
388	NVGR	J054055.6+044211	05 40 55.61	+04 42 11.1	v	4.5	cp
389	NVGR	J054251.7-212715	05 42 51.76	-21 27 15.7	v	7.0	wat? dd? cp? core id amb
390	NVGR	J054359.8+304757	05 43 59.87	+30 47 57.8	c	4.5	t
391	NVGR	J054408.3+182609	05 44 08.30	+18 26 09.9	v	6.5	cp? points
392	NVGR	J054428.0+122314	05 44 28.00	+12 23 14.1	v	4.0	d
393	NVGR	J054444.1+164848	05 44 44.18	+16 48 48.9	c	10.0	w? sz? dl
394			05 46 15.74	+63 20 01.8	v	11.5	wat dd? core id amb (Lara et al.(2001): J0546+633)
395	NVGR	J054618.7-234653	05 46 18.74	-23 46 53.5	v	5.0	cp? wat??
396	NVGR	J054711.3+010930	05 47 11.35	+01 09 30.2	v	8.5	w? wat/nat?
397	NVGR	J054728.7-195625	05 47 28.72	-19 56 25.4	v	15.5	cp but similar shape orientation
398	NVGR	J054730.9-251506	05 47 30.93	-25 15 06.8	v	5.0	nat?
399	NVGR	J054733.0-251900	05 47 33.00	-25 19 00.0	v	4.5	t? cp?
400	NVGR	J054758.5+440256	05 47 58.59	+44 02 56.7	c	6.0	t? if d 4'
401	NVGR	J054808.6+134418	05 48 08.60	+13 44 18.4	v	6.0	d
402	NVGR	J054824.6+414544	05 48 24.62	+41 45 44.1	c	9.0	sz w?
403	NVGR	J054827.3-325850	05 48 27.35	-32 58 50.8	c	11.0	nat ']' appearance
404	NVGR	J054927.7-124917	05 49 27.71	-12 49 17.2	v	6.0	nat/wat? sz? cp?
405	NVGR	J055054.1+190412	05 50 54.13	+19 04 12.9	v	12.0	wat
406	NVGR	J055107.1+284726	05 51 07.15	+28 47 26.7	v	8.0	wat? w? dd?
407	NVGR	J055133.5+372531	05 51 33.59	+37 25 31.4	v	4.0	d
408	NVGR	J055231.1-070744	05 52 31.15	-07 07 44.2	v	6.0	cp? d?
409	NVGR	J055312.9+323411	05 53 12.93	+32 34 11.4	c	6.0	t? d+cp? asym
410	NVGR	J055443.3+024313	05 54 43.32	+02 43 13.6	v	5.5	d with cp?
411	NVGR	J055611.9+135837	05 56 11.91	+13 58 37.5	c	7.0	d?
412	NVGR	J055807.5+431653	05 58 07.55	+43 16 53.8	v	4.5	d? sz?
413			05 59 38.09	+60 44 07.8	c	7.5	wat/nat (Lara et al.(2001): J0559+607)
414	NVGR	J060125.3-323805	06 01 25.39	-32 38 05.8	v	4.5	nat/wat?
415	NVGR	J060210.5-395709	06 02 10.50	-39 57 09.1	c	4.0	nat
416	NVGR	J060238.1+592702	06 02 38.16	+59 27 02.0	c	4.5	d
417	NVGR	J060242.7+323012	06 02 42.77	+32 30 12.7	c	7.0	mg int point with d and loop, this coord of pt
418	NVGR	J060404.3-102611	06 04 04.36	-10 26 11.9	c	5.5	mg wat? sz?
419	NVGR	J060423.1+535151	06 04 23.19	+53 51 51.3	v	6.0	wat/nat?
420			06 07 34.96	+61 14 52.0	v	7.0	d t? core not apparent (Lara et al.(2001): J0607+612)
421	NVGR	J060854.0+072907	06 08 54.09	+07 29 07.1	c	4.5	nat
422	NVGR	J060914.4-142851	06 09 14.41	-14 28 51.2	v	5.5	d
423	NVGR	J060936.0+072914	06 09 36.09	+07 29 14.1	c	6.5	nat partial ring, int

424	NVGR	J061204.2-325744	06 12 04.22	-32 57 44.1	c	12.0	mg dd? wat? GRS or two?
425	NVGR	J061435.3-290909	06 14 35.39	-29 09 09.0	v	8.0	cp cj+d?
426	NVGR	J061508.4-033144	06 15 08.43	-03 31 44.7	v	5.5	d dd?
427	NVGR	J061621.6-212217	06 16 21.61	-21 22 17.7	v	5.0	ring
428	NVGR	J061640.7-042752	06 16 40.78	-04 27 52.0	c	7.5	w? x?
429	NVGR	J061648.6-171702	06 16 48.69	-17 17 02.5	v	4.5	wat?
430	NVGR	J061728.8-171354	06 17 28.83	-17 13 54.7	v	9.0	wat? irr int, may be larger (~16')
431	NVGR	J061935.2+325707	06 19 35.23	+32 57 07.2	v	5.5	cp? elliptic
432	NVGR	J062117.0+335034	06 21 17.00	+33 50 34.9	v	4.0	cp
433	NVGR	J062211.0-155919	06 22 11.06	-15 59 19.7	c	5.5	t
434	NVGR	J062212.0+014153	06 22 12.03	+01 41 53.6	c	5.0	d/t
435	NVGR	J062214.7-120308	06 22 14.73	-12 03 08.0	v	7.0	t? d? dd?
436			06 24 23.43	+63 03 39.4	v	5.0	sz? (Lara et al.(2001): J0624+630)
437	NVGR	J062426.2+503945	06 24 26.28	+50 39 45.0	v	5.5	d dd?
438	NVGR	J062433.8-012222	06 24 33.85	-01 22 22.7	c	4.5	sz? (with cp?) int may be larger
439	NVGR	J062449.9+094013	06 24 49.90	+09 40 13.9	v	4.5	t?
440	NVGR	J062707.1-352917	06 27 07.12	-35 29 17.1	c	5.0	rc t?
441	NVGR	J062710.1+365032	06 27 10.10	+36 50 32.5	v	5.0	cp two elliptic
442	NVGR	J062916.0+080540	06 29 16.02	+08 05 40.5	v	4.0	cp d?
443	NVGR	J062955.3+471713	06 29 55.37	+47 17 13.2	c	6.0	d sz? core fit but not apparent
444	NVGR	J063035.5-192424	06 30 35.50	-19 24 24.9	c	5.5	wat? sz?
445	NVGR	J063044.3+014552	06 30 44.35	+01 45 52.0	v	5.0	d
446	NVGR	J063055.2+062320	06 30 55.24	+06 23 20.0	v	5.5	d? two points
447	NVGR	J063112.9+425300	06 31 12.93	+42 53 00.9	v	8.0	cp point + elliptic
448			06 33 38.54	+72 09 28.3	c	6.5	nat almost ring (Lara et al.(2001): J0633+721)
449	NVGR	J063341.9+483018	06 33 41.90	+48 30 18.4	v	4.5	d
450	NVGR	J063352.1+393945	06 33 52.11	+39 39 45.9	c	6.5	t
451	NVGR	J063547.7+433410	06 35 47.77	+43 34 10.6	v	5.5	cp
452	NVGR	J063631.3-203518	06 36 31.37	-20 35 18.0	v	16.0	w? dd? td? int
453	NVGR	J063850.6+333357	06 38 50.62	+33 33 57.8	v	6.0	wat? cp? possible core 06 38 53.38 +33 34 51.8
454	NVGR	J063855.6+092732	06 38 55.66	+09 27 32.8	v	5.0	cp asym
455	NVGR	J063905.1+243308	06 39 05.11	+24 33 08.1	v	5.5	cp point and elliptic
456	NVGR	J064011.9+333221	06 40 11.97	+33 32 21.2	v	4.5	d
457	NVGR	J064319.9+421032	06 43 19.90	+42 10 32.8	v	4.5	d
458	NVGR	J064407.8+104348	06 44 07.83	+10 43 48.2	c	12.0	d? cp? irr GRS or two?
459	NVGR	J064449.3-273822	06 44 49.33	-27 38 22.0	c	5.0	rc
460	NVGR	J064521.4+173706	06 45 21.40	+17 37 06.9	v	4.0	d
461	NVGR	J064702.0-083941	06 47 02.08	-08 39 41.4	v	6.0	d
462	NVGR	J064832.7+403730	06 48 32.74	+40 37 30.9	v	5.0	d core not apparent
463	NVGR	J064922.1-330305	06 49 22.15	-33 03 05.4	v	7.5	mg cp?
464	NVGR	J065030.7+283332	06 50 30.77	+28 33 32.5	c	7.0	sz
465	NVGR	J065050.7+332908	06 50 50.71	+33 29 08.4	v	4.5	rc
466	NVGR	J065111.0+192744	06 51 11.00	+19 27 44.0	v	6.0	nat
467			06 51 36.69	+19 35 49.6	c	8.0	nat (Soloviyov & Verkhodanov(2014): J065122.5+193713)
468	NVGR	J065204.6-210351	06 52 04.65	-21 03 51.6	v	5.5	mg d?
469	NVGR	J065215.4+430449	06 52 15.44	+43 04 49.9	v	10.0	nat/wat? core id amb
470	NVGR	J065426.6+091148	06 54 26.69	+09 11 48.0	v	5.5	cp d? asym point and elliptic
471			06 54 27.26	+73 19 52.2	c	14.0	d dd?? t? int (Lara et al.(2001): J0654+733)
472	NVGR	J065516.6+103505	06 55 16.63	+10 35 05.5	c	6.0	wat? rc? sz?
473	NVGR	J065600.8+520623	06 56 00.82	+52 06 23.8	v	5.0	cp elliptics
474	NVGR	J065759.7-382951	06 57 59.78	-38 29 51.8	v	4.0	mg d? irr, may be larger system with five or more catalog sources
475	NVGR	J070025.6+222929	07 00 25.61	+22 29 29.9	v	6.5	mg nat+? int
476	NVGR	J070131.5+231317	07 01 31.50	+23 13 17.8	v	5.0	d (with fingers), cp?
477	NVGR	J070207.0+485924	07 02 07.06	+48 59 24.5	v	20.0	mg nat? w? may be smaller
478	NVGR	J070241.2+111115	07 02 41.27	+11 11 15.2	c	4.0	d t
479	NVGR	J070256.1-151930	07 02 56.15	-15 19 30.7	v	6.0	cp? d?
480	NVGR	J070403.0+572247	07 04 03.06	+57 22 47.8	v	5.5	sz? cp?
481	NVGR	J070427.8+115036	07 04 27.81	+11 50 36.0	v	5.0	rc d?
482	NVGR	J070500.3+271152	07 05 00.33	+27 11 52.9	v	5.0	cp
483	NVGR	J070635.0+365602	07 06 35.07	+36 56 02.0	c	7.0	nat partial ring?
484	NVGR	J070643.0+423203	07 06 43.01	+42 32 03.8	c	9.0	w? wat? sz? irr
485	NVGR	J070653.0+331835	07 06 53.06	+33 18 35.4	c	5.0	t
486	NVGR	J070703.0+423249	07 07 03.01	+42 32 49.7	v	4.0	d
487	NVGR	J070729.0+592814	07 07 29.03	+59 28 14.4	v	6.0	nat? cp?
488	NVGR	J070737.5+022616	07 07 37.57	+02 26 16.3	c	7.0	w
489	NVGR	J070746.1-400903	07 07 46.15	-40 09 03.7	v	8.0	dd?
490	NVGR	J070842.7+303041	07 08 42.79	+30 30 41.0	v	5.5	d
491	NVGR	J070907.8+483653	07 09 07.83	+48 36 53.2	c	8.0	wat? also dd? may be smaller
492	NVGR	J070912.5-152701	07 09 12.51	-15 27 01.2	c	5.0	rc? cj? sz?
493	NVGR	J070914.0-360119	07 09 14.04	-36 01 19.0	v	10.0	w (with cp?) dd?
494	NVGR	J071039.2-015913	07 10 39.24	-01 59 13.9	v	4.5	mg d? cj with point cp?
495	NVGR	J071044.5+023839	07 10 44.50	+02 38 39.6	v	5.5	wat?
496	NVGR	J071049.3-382431	07 10 49.37	-38 24 31.3	v	7.0	cp
497	NVGR	J071130.6+471030	07 11 30.68	+47 10 30.8	v	6.5	sz? wat? 4 components
498	NVGR	J071212.4+033307	07 12 12.43	+03 33 07.0	c	4.0	cj? d?
499	NVGR	J071234.0+005933	07 12 34.01	+00 59 33.9	v	4.0	mg cp?
500			07 12 43.96	-08 50 18.7	v	7.0	d dd? 4 components (Soloviyov & Verkhodanov(2014): J071244.0-085019)
501	NVGR	J071416.2+202057	07 14 16.28	+20 20 57.8	v	7.0	cj

502	NVGR	J071450.2-035005	07 14 50.20	-03 50 05.7	v	4.5	wat? d? cp? asym
503	NVGR	J071701.9-362120	07 17 01.93	-36 21 20.0	v	6.5	nat/wat
504	NVGR	J071716.6+430851	07 17 16.68	+43 08 51.6	c	8.0	t sz?
505	NVGR	J071745.5-345315	07 17 45.51	-34 53 15.9	v	7.0	mg dd? int unu beads - possible 5th
506	NVGR	J072223.0+375106	07 22 23.02	+37 51 06.9	c	5.0	t
507	NVGR	J072239.0+235532	07 22 39.01	+23 55 32.9	v	4.5	d?
508	NVGR	J072301.5-211715	07 23 01.57	-21 17 15.5	v	11.0	cp d?
509	NVGR	J072328.4-345300	07 23 28.43	-34 53 00.6	v	4.5	d
510	NVGR	J072405.8+095719	07 24 05.84	+09 57 19.7	v	5.0	d? t?
511	NVGR	J072427.7+151249	07 24 27.73	+15 12 49.5	c	5.0	t d
512	NVGR	J072505.4+031350	07 25 05.48	+03 13 50.1	v	5.0	d? points
513	NVGR	J072538.1-070949	07 25 38.16	-07 09 49.8	v	4.0	wat/nat? may be larger
514	NVGR	J072538.9+400414	07 25 38.95	+40 04 14.1	c	5.0	t
515	NVGR	J072621.0-053416	07 26 21.03	-05 34 16.9	v	5.5	wat/nat? d? cp?
516	NVGR	J072629.3+003347	07 26 29.37	+00 33 47.4	c	5.5	t
517	NVGR	J072802.4+493511	07 28 02.43	+49 35 11.4	v	5.0	rc? d?
518	NVGR	J072806.0+503442	07 28 06.07	+50 34 42.7	v	4.0	d
519	NVGR	J072850.1+543703	07 28 50.18	+54 37 03.5	c	6.0	d core id amb
520	NVGR	J072859.0+335023	07 28 59.01	+33 50 23.0	v	5.5	d? two elliptic
521	NVGR	J072947.4-153948	07 29 47.45	-15 39 48.0	c	6.0	t asym brightness, lobes similar size
522	NVGR	J073001.1+182017	07 30 01.12	+18 20 17.4	v	17.0	cp may be larger
523	NVGR	J073026.5-340715	07 30 26.54	-34 07 15.0	c	5.0	d/t
524	NVGR	J073204.7+155839	07 32 04.73	+15 58 39.0	v	6.5	d? (with cp?) w? wat?
525	NVGR	J073217.5-375759	07 32 17.52	-37 57 59.0	c	5.0	mg complex t?
526	NVGR	J073318.2-365428	07 33 18.23	-36 54 28.6	v	14.0	d?
527			07 33 30.92	+57 41 55.8	c	6.0	sz wat? rc? ring-lobes (Solovoyov & Verkhodanov(2014): J073331.1+574133)
528	NVGR	J073427.0-243022	07 34 27.06	-24 30 22.1	c	6.0	wat?
529	NVGR	J073557.3+145632	07 35 57.31	+14 56 32.6	v	5.5	wat/nat?
530	NVGR	J073610.1-345206	07 36 10.19	-34 52 06.9	v	4.0	d? points
531	NVGR	J073617.1+185705	07 36 17.13	+18 57 05.1	c	5.5	sz? cp?
532	NVGR	J073723.4-330939	07 37 23.43	-33 09 39.6	v	6.0	d
533	NVGR	J074655.3+171806	07 46 55.33	+17 18 06.1	v	5.0	d t?
534	NVGR	J074836.9+554858	07 48 36.93	+55 48 58.3	c	27.0	t
535	NVGR	J074930.1+062216	07 49 30.17	+06 22 16.8	v	5.0	d dd?
536	NVGR	J074947.3-381001	07 49 47.32	-38 10 01.8	v	5.0	d? points
537	NVGR	J075020.5+163258	07 50 20.58	+16 32 58.5	c	4.5	d/t
538	NVGR	J075026.7+541302	07 50 26.77	+54 13 02.9	v	5.5	d
539	NVGR	J075111.3+423109	07 51 11.31	+42 31 09.8	v	8.0	wat? sz?
540	NVGR	J075412.7-120014	07 54 12.70	-12 00 14.4	c	5.0	d/t
541	NVGR	J075441.2+232449	07 54 41.21	+23 24 49.8	c	7.5	w? t?
542	NVGR	J075519.2+010221	07 55 19.27	+01 02 21.9	c	5.0	t nat?
543	NVGR	J075633.4-273706	07 56 33.45	-27 37 06.3	c	5.5	wat 'starwars'
544	NVGR	J075706.6+095634	07 57 06.60	+09 56 34.9	c	3.5	rc? cj? cp?
545			07 57 32.90	+82 39 03.4	v	3.5	wat? (Lara et al.(2001): J0757+826)
546	NVGR	J075835.0-261513	07 58 35.08	-26 15 13.3	c	11.5	wat/nat? sz?
547	NVGR	J075903.7+475458	07 59 03.75	+47 54 58.8	c	4.0	d (with cp)
548	NVGR	J075916.1+270908	07 59 16.17	+27 09 08.7	v	4.0	d (with cp?)
549	NVGR	J075931.0+082547	07 59 31.02	+08 25 47.6	v	5.0	nat/wat? sz?
550	NVGR	J080053.5+392440	08 00 53.56	+39 24 40.8	c	5.0	sz? wat?
551	NVGR	J080135.7-152103	08 01 35.74	-15 21 03.4	v	7.0	mg t? cp?
552			08 02 37.10	-09 58 52.0	v	13.0	wat/nat? w? sz? GRS or two? (Solovoyov & Verkhodanov(2014): J080244.1-095757)
553	NVGR	J080321.6+515525	08 03 21.61	+51 55 25.6	v	5.0	cp? d?
554	NVGR	J080438.7+391906	08 04 38.78	+39 19 06.6	v	5.0	cp? nat?
555	NVGR	J080501.2+031422	08 05 01.24	+03 14 22.4	v	4.0	d? (with cp?) may be 6' wat
556	NVGR	J080557.7-372806	08 05 57.75	-37 28 06.2	v	6.0	sz? dd?
557	NVGR	J080633.3-393032	08 06 33.31	-39 30 32.3	c	6.0	wat/nat? dl core id amb
558			08 07 10.33	+74 00 34.0	c	9.5	t? rc with jets? dd? int a core-gap system? possibly larger (Lara et al.(2001): J0807+740)
559	NVGR	J080829.9+390148	08 08 29.99	+39 01 48.3	v	5.0	d? with tail? cp?
560	NVGR	J081208.2-394654	08 12 08.25	-39 46 54.5	v	7.0	d?
561	NVGR	J081252.9-270820	08 12 52.96	-27 08 20.0	v	4.0	d
562	NVGR	J081327.1-043043	08 13 27.10	-04 30 43.3	v	4.5	d
563	NVGR	J081346.2-113048	08 13 46.20	-11 30 48.8	v	7.0	wat/nat? distorted ring
564	NVGR	J081707.1-211222	08 17 07.18	-21 12 22.1	v	4.5	sz? d
565			08 19 50.42	+75 38 38.9	c	10.0	t asym (Lara et al.(2001): J0819+756)
566	NVGR	J082007.9-135008	08 20 07.94	-13 50 08.3	v	4.5	cp? nat? two adjacent elliptic
567	NVGR	J082015.8+094729	08 20 15.82	+09 47 29.6	c	4.0	rc?
568	NVGR	J082043.5-241438	08 20 43.52	-24 14 38.2	c	7.5	t? cp?
569			08 21 28.36	-30 11 24.0	v	8.0	w? dd? looks like 4 components but only 3 fit (Solovoyov & Verkhodanov(2014): J082128.4-301124)
570	NVGR	J082234.0+055624	08 22 34.03	+05 56 24.8	c	9.0	rc
571	NVGR	J082242.0-102059	08 22 42.07	-10 20 59.3	v	6.0	d two elliptic
572	NVGR	J082313.6+033257	08 23 13.62	+03 32 57.4	v	7.5	mg wat dd? wat dd? td? core id amb unu, may be larger
573	NVGR	J082333.8+113019	08 23 33.85	+11 30 19.2	v	5.5	d
574	NVGR	J082339.9-173525	08 23 39.97	-17 35 25.4	v	4.0	d
575	NVGR	J082556.1+175143	08 25 56.10	+17 51 43.8	v	4.0	d
576			08 26 02.05	+69 20 57.7	c	8.5	t w? sz? (Lara et al.(2001): J0825+693)
577	NVGR	J082621.7+105930	08 26 21.70	+10 59 30.6	v	5.0	d? points
578	NVGR	J082841.2+243637	08 28 41.26	+24 36 37.8	v	4.0	wat
579			08 28 54.99	+63 14 28.8	v	5.0	d sz? (Lara et al.(2001): J0828+632)

580	NVGR	J082942.9+224801	08 29 42.95	+22 48 01.3	v	6.0	d? asym sz? wat/nat?
581	NVGR	J083018.7+182640	08 30 18.70	+18 26 40.9	v	4.5	wat?
582	NVGR	J083040.7-191409	08 30 40.73	-19 14 09.2	v	4.5	d? cj?
583	NVGR	J083106.0+414745	08 31 06.09	+41 47 45.3	v	5.0	d (with cp?)
584	NVGR	J083127.8+321933	08 31 27.83	+32 19 33.4	v	13.0	x? or cp of strong 7'd and weaker 13' wat (with ring-lobes)?
585	NVGR	J083139.1+223423	08 31 39.10	+22 34 23.3	c	4.0	wat?
586	NVGR	J083148.2+223912	08 31 48.27	+22 39 12.5	v	4.0	sz fit missed core
587			08 35 34.01	-01 51 00.6	c	8.0	t rc pretty (Solov'yov & Verkhodanov(2014): J083533.9-015100)
588	NVGR	J083536.6-201030	08 35 36.62	-20 10 30.3	v	5.5	cp
589	NVGR	J083637.8-110531	08 36 37.81	-11 05 31.2	v	5.5	wat?
590	NVGR	J083915.7+285045	08 39 15.78	+28 50 45.1	c	8.0	wat? sz?
591	NVGR	J084001.7+294853	08 40 01.79	+29 48 53.2	c	7.5	rc sz?
592	NVGR	J084004.8-215019	08 40 04.81	-21 50 19.8	c	4.5	t may be larger
593	NVGR	J084010.6-201915	08 40 10.62	-20 19 15.9	v	6.0	wat? core id amb
594	NVGR	J084017.1+175720	08 40 17.13	+17 57 20.2	c	7.5	rc
595	NVGR	J084046.2-181635	08 40 46.24	-18 16 35.8	v	6.0	d dd? 4 components
596	NVGR	J084109.0+322348	08 41 09.02	+32 23 48.0	v	6.0	mg wat/nat? (with cp?) wisps
597	NVGR	J084202.3+385628	08 42 02.31	+38 56 28.6	v	5.0	wat?
598	NVGR	J084210.2+415634	08 42 10.27	+41 56 34.2	c	4.5	sz? wat?
599	NVGR	J084450.3-263222	08 44 50.36	-26 32 22.2	v	7.0	d
600	NVGR	J084451.6+004008	08 44 51.68	+00 40 08.5	c	5.0	d/t
601	NVGR	J084457.0-093817	08 44 57.08	-09 38 17.1	v	3.5	d with cp?
602	NVGR	J084504.2+510513	08 45 04.23	+51 05 13.5	v	7.5	cp 3' rc? and 3' cj?
603			08 45 08.06	-33 47 36.7	c	34.5	rc with jets no apparent lobes (Solov'yov & Verkhodanov(2014): J084507.2-334711)
604	NVGR	J084718.3+422329	08 47 18.37	+42 23 29.7	v	4.0	d
605	NVGR	J084728.0-331849	08 47 28.07	-33 18 49.5	v	8.0	w? wat? cp? core id amb
606	NVGR	J084745.9+383205	08 47 45.95	+38 32 05.7	v	5.0	d core not apparent
607	NVGR	J084759.6+314710	08 47 59.69	+31 47 10.7	c	7.0	sz? wat/nat?
608	NVGR	J084843.9+170105	08 48 43.92	+17 01 05.8	v	5.5	d rc?
609	NVGR	J084937.9-304945	08 49 37.98	-30 49 45.8	c	5.0	sz?
610	NVGR	J085250.3-192705	08 52 50.34	-19 27 05.6	v	4.5	d t? if t coord 08 52 49.23 -19 26 33.2 c
611	NVGR	J085316.0-313942	08 53 16.09	-31 39 42.0	v	5.5	d? two points
612	NVGR	J085350.0+145224	08 53 50.01	+14 52 24.3	c	9.5	t? wat?
613	NVGR	J085354.3-183910	08 53 54.36	-18 39 10.9	v	5.0	d? elliptic
614	NVGR	J085520.3+055454	08 55 20.31	+05 54 54.8	v	4.5	cj?
615			08 56 19.75	+66 21 28.0	v	5.5	d 4 components (Lara et al.(2001): J0856+663)
616	NVGR	J085634.0+595744	08 56 34.03	+59 57 44.3	v	4.5	d
617	NVGR	J085832.6+562011	08 58 32.69	+56 20 11.2	v	5.5	mg d? possible 9' system many neighboring sources
618	NVGR	J090017.4+074532	09 00 17.47	+07 45 32.2	c	8.0	wat?
619	NVGR	J090049.7-022559	09 00 49.72	-02 25 59.4	v	5.5	d x?
620	NVGR	J090112.1+294340	09 01 12.13	+29 43 40.7	v	4.5	d asym
621	NVGR	J090123.2+191422	09 01 23.24	+19 14 22.7	v	4.5	d
622	NVGR	J090212.2-225111	09 02 12.26	-22 51 11.0	c	5.0	d/t
623	NVGR	J090230.6+204434	09 02 30.64	+20 44 34.3	v	5.5	mg cp? with nat?
624	NVGR	J090238.4+173752	09 02 38.49	+17 37 52.1	c	9.0	t? wat?? int structure one lobe anchor-like
625	NVGR	J090302.0+120901	09 03 02.06	+12 09 01.7	v	7.0	d two elliptic
626	NVGR	J090438.0-243553	09 04 38.09	-24 35 53.6	v	9.0	t with extended sz tails dd?
627	NVGR	J090640.7+142518	09 06 40.79	+14 25 18.0	c	8.0	t?
628	NVGR	J090715.5-120912	09 07 15.52	-12 09 12.6	v	4.0	d sz?
629	NVGR	J090746.3-353553	09 07 46.33	-35 35 53.9	v	11.0	wat? dl
630	NVGR	J090806.2-100128	09 08 06.21	-10 01 28.5	c	6.5	wat/nat?
631	NVGR	J090841.5+594542	09 08 41.56	+59 45 42.0	c	6.0	t
632	NVGR	J091016.2-093545	09 10 16.29	-09 35 45.8	v	5.0	mg ring structure
633	NVGR	J091031.5+035310	09 10 31.51	+03 53 10.2	v	4.5	d wat?
634	NVGR	J091058.2-103342	09 10 58.25	-10 33 42.6	c	5.5	wat
635	NVGR	J091121.5-061539	09 11 21.55	-06 15 39.7	c	6.0	t
636	NVGR	J091130.3-364052	09 11 30.30	-36 40 52.1	c	8.0	nat/wat t
637	NVGR	J091234.1-095758	09 12 34.17	-09 57 58.0	v	5.0	mg point and elliptic
638			09 12 51.57	+35 09 28.8	v	8.0	d (Machalski et al.(2001): J0912+3510)
639	NVGR	J091350.3-143220	09 13 50.34	-14 32 20.0	c	8.0	nat int
640	NVGR	J091419.3+100638	09 14 19.36	+10 06 38.4	c	8.5	t nat? ring? dd? int
641	NVGR	J091731.0-081351	09 17 31.09	-08 13 51.9	v	7.0	d? w?
642	NVGR	J091856.9+315051	09 18 56.96	+31 50 51.8	c	8.0	w? sz?
643	NVGR	J091942.8+260928	09 19 42.85	+26 09 28.5	v	5.0	d
644	NVGR	J091950.4-150641	09 19 50.49	-15 06 41.4	v	4.0	d
645	NVGR	J092014.4-383439	09 20 14.48	-38 34 39.5	c	5.0	rc with cp?
646	NVGR	J092110.5-003342	09 21 10.50	-00 33 42.8	v	4.0	mg
647	NVGR	J092223.6+091914	09 22 23.61	+09 19 14.9	v	9.0	d?
648	NVGR	J092255.7+245259	09 22 55.76	+24 52 59.6	v	4.0	d? cj?
649	NVGR	J092328.8-213649	09 23 28.82	-21 36 49.6	v	8.0	d? with cp? may be smaller
650	NVGR	J092330.9+242758	09 23 30.92	+24 27 58.4	c	20.0	sz? t?
651	NVGR	J092402.3-170327	09 24 02.38	-17 03 27.1	v	5.0	d? points
652	NVGR	J092439.1+302839	09 24 39.15	+30 28 39.1	c	5.0	d/t
653	NVGR	J092545.8+404735	09 25 45.85	+40 47 35.6	v	5.5	mg d (with cp?)
654			09 26 00.89	+65 19 21.5	c	6.5	wat? dl (Lara et al.(2001): J0926+653)
655			09 26 53.34	+61 00 11.4	v	5.5	d core id amb (Lara et al.(2001): J0926+610)
656	NVGR	J092751.9+295902	09 27 51.92	+29 59 02.3	c	12.5	t
657	NVGR	J092910.7+414659	09 29 10.74	+41 46 59.7	v	8.0	dd 4 components

658	NVGR	J093058.9+034831	09 30 58.97	+03 48 31.4	c	7.0	wat
659	NVGR	J093237.0+161213	09 32 37.04	+16 12 13.4	v	5.5	sz? int complex
660	NVGR	J093241.8-401019	09 32 41.87	-40 10 19.4	c	8.0	d t? 5 components
661	NVGR	J093421.0+382327	09 34 21.03	+38 23 27.5	v	6.0	d
662	NVGR	J093800.8-291241	09 38 00.88	-29 12 41.5	v	6.0	rc? d?
663	NVGR	J093831.6-042049	09 38 31.67	-04 20 49.5	c	6.5	d core not apparent
664	NVGR	J094004.0-061859	09 40 04.07	-06 18 59.2	v	4.0	d
665	NVGR	J094005.7+510408	09 40 05.78	+51 04 08.2	c	6.0	mg t wat? (with cp?)
666	NVGR	J094120.1-065008	09 41 20.11	-06 50 08.0	v	4.5	d? cj?
667	NVGR	J094538.9-311127	09 45 38.90	-31 11 27.8	c	6.5	rc face on sp int NGC 2997
668	NVGR	J094645.0-080259	09 46 45.01	-08 02 59.1	v	4.0	wat?
669	NVGR	J094708.1-133828	09 47 08.16	-13 38 28.4	v	27.0	t
670	NVGR	J094745.7+072511	09 47 45.70	+07 25 11.3	v	6.0	rc t
671	NVGR	J094816.1-161957	09 48 16.16	-16 19 57.0	v	8.0	wat? with partial ring? int
672	NVGR	J094834.0-104358	09 48 34.05	-10 43 58.1	c	4.0	d core not apparent
673	NVGR	J094959.1-334217	09 49 59.11	-33 42 17.5	v	4.0	d
674			09 50 08.18	+73 15 14.4	v	16.0	x? int unusual lobe shape GRS or two? (Lara et al.(2001): J0949+732)
675	NVGR	J095136.3-392618	09 51 36.30	-39 26 18.0	c	7.0	w?
676	NVGR	J095150.9-265312	09 51 50.97	-26 53 12.5	v	4.0	d? points
677	NVGR	J095202.6-124116	09 52 02.69	-12 41 16.6	v	4.0	d? point and elliptic
678	NVGR	J095331.6-185518	09 53 31.67	-18 55 18.8	c	4.0	rc?
679	NVGR	J095419.2+271558	09 54 19.25	+27 15 58.3	c	5.5	t
680	NVGR	J095454.8+405913	09 54 54.81	+40 59 13.6	c	4.5	d t
681	NVGR	J095640.5-000128	09 56 40.53	-00 01 28.4	v	4.0	wat d
682	NVGR	J095653.8-134512	09 56 53.86	-13 45 12.7	v	4.0	d (with cp?)
683	NVGR	J095716.3+190645	09 57 16.36	+19 06 45.4	v	4.0	d
684	NVGR	J095809.2-371018	09 58 09.24	-37 10 18.5	c	5.0	sz? t
685	NVGR	J095813.5+265059	09 58 13.54	+26 50 59.1	v	5.5	d? lensing? int
686	NVGR	J095940.4+172533	09 59 40.43	+17 25 33.7	c	11.0	t sz?
687	NVGR	J100025.9-075423	10 00 25.90	-07 54 23.8	v	6.0	wat?
688	NVGR	J100601.7+345410	10 06 01.74	+34 54 10.4	c	42.0	mg t? asym jets, may be larger, incidental
689	NVGR	J100623.2-363645	10 06 23.28	-36 36 45.4	v	6.0	mg cp? 4 component group?
690	NVGR	J100645.9-062118	10 06 45.97	-06 21 18.8	v	5.0	d? t?
691	NVGR	J100651.8+142226	10 06 51.80	+14 22 26.4	c	9.0	sz? nat? w? core id amb, may be larger
692	NVGR	J100728.8+344217	10 07 28.87	+34 42 17.5	v	7.0	wat? part of jet of larger system?
693	NVGR	J100827.4-031726	10 08 27.49	-03 17 26.9	v	7.0	wat? 4 components
694	NVGR	J100834.4-213912	10 08 34.40	-21 39 12.5	c	5.0	rc?
695	NVGR	J100932.5-365708	10 09 32.51	-36 57 08.1	c	6.0	t
696	NVGR	J100943.6+033657	10 09 43.61	+03 36 57.8	v	6.0	w? wat?
697	NVGR	J100944.9+162555	10 09 44.92	+16 25 55.8	v	4.0	d rc?
698	NVGR	J101005.8-170740	10 10 05.80	-17 07 40.6	v	12.0	cp
699			10 11 11.00	+31 11 07.1	v	6.0	d dl (Machalski et al.(2001): J1011+3111)
700	NVGR	J101133.8+032757	10 11 33.83	+03 27 57.5	v	21.5	cp
701	NVGR	J101350.9-321954	10 13 50.99	-32 19 54.4	c	5.0	d t? core not apparent
702	NVGR	J101351.4+855400	10 13 51.48	+85 54 00.7	v	3.5	rc? (with cp?)
703	NVGR	J101442.3-014559	10 14 42.38	-01 45 59.0	v	7.0	mg irr? 4' d and 3.5' d
704	NVGR	J101453.0-045101	10 14 53.01	-04 51 01.0	c	5.5	t
705	NVGR	J101557.4+404649	10 15 57.49	+40 46 49.3	v	4.5	d
706	NVGR	J101815.1+032311	10 18 15.15	+03 23 11.2	v	5.0	d? point and elliptic
707			10 18 49.77	-12 40 56.3	c	12.0	sz? unu (Machalski et al.(2001): J1018-1240)
708			10 20 53.69	+48 31 27.2	c	9.0	w butterfly int (Solovoyov & Verkhodanov(2014): J102054.5+483044)
709	NVGR	J102054.7-165809	10 20 54.70	-16 58 09.0	c	5.0	d core not apparent
710	NVGR	J102103.4-023634	10 21 03.47	-02 36 34.2	c	7.5	d? x? core id amb
711	NVGR	J102131.6+051858	10 21 31.65	+05 18 58.2	c	16.0	t
712	NVGR	J102135.6-334339	10 21 35.65	-33 43 39.6	c	5.0	nat
713	NVGR	J102157.0+144334	10 21 57.01	+14 43 34.0	v	4.5	d
714	NVGR	J102407.4-020344	10 24 07.48	-02 03 44.9	c	8.0	wat? with cp?
715	NVGR	J102631.3-395617	10 26 31.33	-39 56 17.0	v	8.0	w? just jets?
716	NVGR	J102657.4-252220	10 26 57.48	-25 22 20.2	v	9.0	cp
717	NVGR	J103010.5-030950	10 30 10.54	-03 09 50.5	c	15.0	wat
718	NVGR	J103129.7+165018	10 31 29.75	+16 50 18.3	v	5.0	d
719	NVGR	J103147.1+224943	10 31 47.12	+22 49 43.8	v	5.5	d? sz?
720	NVGR	J103213.9+275610	10 32 13.98	+27 56 10.7	c	14.0	w?
721	NVGR	J103259.5+564451	10 32 59.33	+56 44 41.5	c	17.0	w?
722	NVGR	J103404.0+184044	10 34 04.08	+18 40 44.2	c	11.5	t? irr
723	NVGR	J103525.8+065007	10 35 25.85	+06 50 07.4	v	5.0	cp point and elliptic
724	NVGR	J103607.1+000618	10 36 07.12	+00 06 18.5	v	4.5	sz?
725	NVGR	J103622.9+383136	10 36 22.95	+38 31 36.8	c	5.0	t core fit, not apparent
726	NVGR	J103635.4+383537	10 36 35.46	+38 35 37.1	c	5.5	wat? sz?
727			10 36 37.92	+67 47 59.3	c	5.5	d (Lara et al.(2001): J1036+677)
728	NVGR	J103835.8-394138	10 38 35.89	-39 41 38.9	c	5.0	sz?
729	NVGR	J103938.6+051031	10 39 38.62	+05 10 31.3	c	5.0	cj
730	NVGR	J104013.4-261659	10 40 13.48	-26 16 59.8	v	4.0	d
731	NVGR	J104057.5-132915	10 40 57.55	-13 29 15.4	v	4.0	cp point with rc
732	NVGR	J104153.1-334029	10 41 53.13	-33 40 29.5	v	5.5	d? two elliptic
733	NVGR	J104332.9+245506	10 43 32.95	+24 55 06.9	v	5.5	rc
734	NVGR	J104631.1+543615	10 46 31.15	+54 36 15.4	v	6.0	dd
735	NVGR	J104633.2+134706	10 46 33.23	+13 47 06.2	v	6.5	cp? (two dissimilar elliptic, not aligned)

736	NVGR	J104751.1+195339	10 47 51.12	+19 53 39.0	c	10.0	wat/nat core id amb
737	NVGR	J104925.2-395854	10 49 25.24	-39 58 54.6	v	5.5	cp? d?
738	NVGR	J104945.9-130811	10 49 45.99	-13 08 11.7	v	10.0	t sz?, cat missed core?
739	NVGR	J105052.9+400055	10 50 52.99	+40 00 55.3	c	5.0	t asym
740	NVGR	J105112.3+464550	10 51 12.39	+46 45 50.3	v	6.0	cp? (d? with cp?)
741	NVGR	J105147.8+552316	10 51 47.82	+55 23 16.5	c	6.0	mg wat? or 3.5' d and elliptic
742	NVGR	J105225.0+373013	10 52 25.00	+37 30 13.4	v	4.0	rc? d?
743	NVGR	J105308.1+260200	10 53 08.16	+26 02 00.2	c	5.5	wat/nat? sz? core id amb
744	NVGR	J105349.1+402352	10 53 49.14	+40 23 52.0	c	4.5	rc? t?
745	NVGR	J105421.1+022754	10 54 21.16	+02 27 54.7	v	4.5	d?
746	NVGR	J105514.4+462105	10 55 14.45	+46 21 05.1	v	3.5	d
747	NVGR	J105538.3+143410	10 55 38.38	+14 34 10.3	c	5.0	t sz?
748	NVGR	J105553.0-021118	10 55 53.00	-02 11 18.4	c	3.5	rc? - incidental
749	NVGR	J105744.3-150741	10 57 44.39	-15 07 41.5	c	7.0	w? sz?
750	NVGR	J105850.5-361802	10 58 50.51	-36 18 02.3	c	8.0	nat/wat? cp?
751	NVGR	J105948.0-044623	10 59 48.08	-04 46 23.6	c	5.0	cj?
752	NVGR	J110152.8+164037	11 01 52.81	+16 40 37.5	v	7.0	x butterfly int
753	NVGR	J110423.0-270915	11 04 23.09	-27 09 15.5	c	6.5	t?
754	NVGR	J110522.0-220933	11 05 22.06	-22 09 33.5	v	4.0	wat? rc?
755	NVGR	J110548.6-000209	11 05 48.67	-00 02 09.9	v	7.0	mg rc (with cp d?)
756	NVGR	J110557.0-150422	11 05 57.08	-15 04 22.5	v	5.5	ring
757	NVGR	J110651.1+141212	11 06 51.10	+14 12 12.7	c	7.5	nat? core id amb
758	NVGR	J110712.5-311433	11 07 12.52	-31 14 33.4	c	6.0	t
759	NVGR	J111130.8+554027	11 11 30.84	+55 40 27.0	c	7.0	rc
760			11 11 41.31	-13 24 29.6	c	8.0	t rc (Soloviyov & Verkhodanov(2014): J111141.9-132417)
761	NVGR	J111254.2-122001	11 12 54.26	-12 20 01.0	v	4.0	mg wat? may be larger
762			11 13 05.93	+40 17 32.8	c	12.0	w wat? (Machalski et al.(2001): J1113+4017)
763	NVGR	J111318.6+412424	11 13 18.62	+41 24 24.0	v	5.0	d? wat? (with cp?)
764	NVGR	J111338.1-380146	11 13 38.17	-38 01 46.7	v	7.5	d wat?
765	NVGR	J111432.3-040102	11 14 32.35	-04 01 02.8	v	4.0	rc d
766	NVGR	J111435.2-033111	11 14 35.22	-03 31 11.2	v	5.5	d? (with cp?) points
767	NVGR	J111511.8-175049	11 15 11.84	-17 50 49.4	v	9.5	wat? sz? rc and rc?
768	NVGR	J111733.0-152908	11 17 33.06	-15 29 08.7	v	4.0	d t core fit, not apparent
769	NVGR	J111817.3-324844	11 18 17.34	-32 48 44.3	v	7.0	rc NGC 3621
770	NVGR	J111859.2+275409	11 18 59.20	+27 54 09.0	c	6.0	wat
771	NVGR	J111944.5+135747	11 19 44.56	+13 57 47.9	c	8.5	wat/nat? t?
772	NVGR	J112015.6+125909	11 20 15.64	+12 59 09.4	v	5.0	rc sz? int
773	NVGR	J112228.1-360512	11 22 28.14	-36 05 12.0	c	6.5	t fishtails
774	NVGR	J112253.9+010641	11 22 53.94	+01 06 41.5	v	4.5	sz?
775	NVGR	J112422.6+150958	11 24 22.67	+15 09 58.5	c	5.0	mg wat/nat?
776	NVGR	J112446.0-040229	11 24 46.01	-04 02 29.2	c	6.0	rc (with cp?) arc? part of ring? may be larger
777	NVGR	J112500.3+372223	11 25 00.36	+37 22 23.9	c	12.0	mg cp?
778	NVGR	J112539.9+082014	11 25 39.99	+08 20 14.9	v	5.0	rc? unu grayscale
779	NVGR	J112556.8+200529	11 25 56.85	+20 05 29.9	v	3.5	d (with cp?) may be larger
780	NVGR	J112605.5-004216	11 26 05.53	-00 42 16.8	v	5.5	sz? cp?
781	NVGR	J112725.1-160005	11 27 25.10	-16 00 05.3	v	6.0	d
782	NVGR	J112801.2-393408	11 28 01.24	-39 34 08.8	v	5.0	mg d? may be larger
783	NVGR	J112824.1-112216	11 28 24.15	-11 22 16.5	c	5.5	t? rc?
784	NVGR	J112912.0+273311	11 29 12.00	+27 33 11.8	c	6.0	rc (with cp?)
785	NVGR	J112936.5-285313	11 29 36.53	-28 53 13.1	v	4.5	wat? d? elliptic
786	NVGR	J113020.5-132053	11 30 20.55	-13 20 53.7	c	6.5	t 5 components
787	NVGR	J113022.5-345551	11 30 22.56	-34 55 51.4	v	6.5	w? nat? butterfly cp?
788	NVGR	J113051.7+062950	11 30 51.75	+06 29 50.9	v	6.5	d? asym
789	NVGR	J113153.2-262314	11 31 53.23	-26 23 14.9	v	4.5	cp
790	NVGR	J113305.0-040047	11 33 05.05	-04 00 47.6	c	9.5	sz nat/wat? hook
791	NVGR	J113315.6-230634	11 33 15.61	-23 06 34.8	v	5.0	wat? cj? cp?
792	NVGR	J113356.3+490334	11 33 56.35	+49 03 34.8	v	4.0	rc? quad
793	NVGR	J113439.8+485816	11 34 39.83	+48 58 16.5	v	7.0	rc?
794	NVGR	J113501.0+344402	11 35 01.06	+34 44 02.7	v	4.0	mg d?
795			11 35 37.99	+39 01 54.0	v	6.0	d? (Soloviyov & Verkhodanov(2014): J113538.0+390154)
796	NVGR	J113553.4+400013	11 35 53.40	+40 00 13.7	v	6.0	cp (two elliptic, not aligned)
797	NVGR	J113639.6+325310	11 36 39.67	+32 53 10.3	v	4.5	wat?
798			11 37 22.74	+61 19 56.3	v	5.0	d? x? wat? butterfly (Lara et al.(2001): J1137+613)
799	NVGR	J113724.3-005132	11 37 24.32	-00 51 32.0	v	6.5	rc? shell?
800	NVGR	J114000.4-033419	11 40 00.41	-03 34 19.8	v	6.0	wat/nat? cp?
801	NVGR	J114054.4+252557	11 40 54.48	+25 25 57.5	v	4.0	d
802	NVGR	J114105.0-212347	11 41 05.08	-21 23 47.5	v	7.0	d t?
803	NVGR	J114301.1+130447	11 43 01.17	+13 04 47.7	v	6.0	d x??
804	NVGR	J114344.5+222903	11 43 44.58	+22 29 03.6	v	4.0	t d
805	NVGR	J114410.7-392217	11 44 10.78	-39 22 17.9	v	5.5	wat
806	NVGR	J114428.7+370846	11 44 28.73	+37 08 46.0	c	7.0	nat/wat? irr
807	NVGR	J114505.2+193637	11 45 05.23	+19 36 37.8	c	7.5	nat? rc? lobe of FRII? int unu
808	NVGR	J114528.7-022517	11 45 28.76	-02 25 17.2	v	6.0	cp
809	NVGR	J114704.6+350302	11 47 04.62	+35 03 02.1	c	6.0	t (with tail?)
810	NVGR	J114740.5-381549	11 47 40.57	-38 15 49.6	c	7.5	t
811	NVGR	J114746.1+345932	11 47 46.10	+34 59 32.9	v	7.0	d? may be associated with brt source at J11 47 22.14 +35 01 07.3 - incidental
812	NVGR	J114746.6+263412	11 47 46.61	+26 34 12.1	v	8.0	mg group of 3 bright point sources + elliptic
813	NVGR	J114908.7-120431	11 49 08.70	-12 04 31.3	c	7.0	d/t

814	NVGR	J114931.8-074141	11 49 31.81	-07 41 41.5 v	5.0	d? two elliptic
815	NVGR	J115054.0-280339	11 50 54.01	-28 03 39.1 v	6.0	mg rc? int cluster, lots of pairs in neighborhood
816	NVGR	J115212.3+395012	11 52 12.31	+39 50 12.5 v	4.5	d? t? wat?
817	NVGR	J115252.3+042150	11 52 52.32	+04 21 50.5 v	6.5	cp? wat?
818	NVGR	J115400.4+093918	11 54 00.46	+09 39 18.9 v	6.5	nat/wat (with cp's)
819	NVGR	J115512.0+545250	11 55 12.00	+54 52 50.0 v	3.5	d
820	NVGR	J115530.6-121205	11 55 30.64	-12 12 05.8 v	6.0	d? two elliptic, long axes parallel
821	NVGR	J115530.6+545357	11 55 30.66	+54 53 57.0 v	4.0	d
822	NVGR	J115648.2-040542	11 56 48.27	-04 05 42.4 c	10.5	t sz? nat? ring-lobe hook, pretty
823	NVGR	J115820.0+262116	11 58 20.07	+26 21 16.3 v	6.0	d dd?
824			11 59 05.56	+58 20 26.3 v	8.5	w? sz? irr (with cp?) (Soloviyov & Verkhodanov(2014): J115909.1+582041)
825	NVGR	J115920.8-191536	11 59 20.87	-19 15 36.8 v	6.5	cp
826	NVGR	J115925.5+210659	11 59 25.59	+21 06 59.9 c	4.5	t
827	NVGR	J120121.4+492534	12 01 21.40	+49 25 34.0 v	6.5	d? x?
828	NVGR	J120227.1+584141	12 02 27.19	+58 41 41.5 v	4.5	d
829	NVGR	J120303.2+513931	12 03 03.20	+51 39 31.1 v	4.5	nat?
830	NVGR	J120337.9+275646	12 03 37.92	+27 56 46.2 v	5.5	mg d
831	NVGR	J120342.2+234315	12 03 42.26	+23 43 15.6 v	5.5	d
832	NVGR	J120400.7+201348	12 04 00.79	+20 13 48.7 c	5.0	nat
833	NVGR	J120445.9+483429	12 04 45.92	+48 34 29.0 v	7.0	mg cp wat with rc
834	NVGR	J120705.1-274154	12 07 05.18	-27 41 54.0 c	34.0	wat
835	NVGR	J120848.2-090204	12 08 48.22	-09 02 04.3 v	5.0	cp point and elliptic
836	NVGR	J120855.5+464113	12 08 55.57	+46 41 13.7 c	6.5	t sz? wat?
837	NVGR	J121129.3-035009	12 11 29.34	-03 50 09.2 v	4.5	mg sz?
838	NVGR	J121135.0+145131	12 11 35.07	+14 51 31.7 v	4.0	mg d with point
839			12 11 53.89	+74 19 45.8 c	9.0	t sz? (Lara et al.(2001): J1211+743)
840	NVGR	J121348.4+145403	12 13 48.40	+14 54 03.5 c	5.5	t
841	NVGR	J121350.3-394127	12 13 50.32	-39 41 27.5 v	5.0	cp
842	NVGR	J121605.2+202107	12 16 05.22	+20 21 07.2 v	4.5	d?
843	NVGR	J121610.0+415923	12 16 10.04	+41 59 23.9 c	7.0	t
844	NVGR	J121614.5+162446	12 16 14.57	+16 24 46.1 v	5.0	d sz?
845			12 16 41.40	+67 24 52.9 v	7.0	d? (Lara et al.(2001): J1216+674)
846	NVGR	J121705.0-031132	12 17 05.09	-03 11 32.4 c	4.0	rc? (with cp?)
847	NVGR	J121842.0-045847	12 18 42.04	-04 58 47.5 v	4.5	wat? cp?
848			12 18 47.23	+50 26 07.0 c	5.0	t x?? core not apparent (Machalski et al.(2001): J1218+5026)
849	NVGR	J121925.2+054945	12 19 25.20	+05 49 45.9 v	9.5	d may be larger
850			12 20 39.44	+63 41 39.0 v	6.5	t may be larger (Lara et al.(2001): J1220+636)
851	NVGR	J122044.3+055205	12 20 44.33	+05 52 05.6 c	8.5	mg t?
852	NVGR	J122112.6+075350	12 21 12.61	+07 53 50.3 v	5.0	mg
853	NVGR	J122154.9+042825	12 21 54.95	+04 28 25.3 c	5.5	rc? M 61
854	NVGR	J122255.0+154921	12 22 55.06	+15 49 21.5 c	6.0	rc M 100
855	NVGR	J122315.1+322321	12 23 15.12	+32 23 21.8 v	4.0	d? points
856	NVGR	J122444.5-254439	12 24 44.54	-25 44 39.7 c	5.0	wat
857	NVGR	J123037.7-234441	12 30 37.78	-23 44 41.3 c	5.5	t
858	NVGR	J123049.4+122321	12 30 49.46	+12 23 21.6 c	16.0	rc? size ambiguous M 87
859	NVGR	J123056.8+310135	12 30 56.87	+31 01 35.1 v	5.5	cp point and elliptic
860	NVGR	J123407.0-333845	12 34 07.06	-33 38 45.6 v	6.0	d
861	NVGR	J123419.9+081150	12 34 19.90	+08 11 50.7 c	5.0	rc core NGC 4535
862	NVGR	J123427.5-160740	12 34 27.53	-16 07 40.7 v	5.5	d (with cp's) 4 components
863	NVGR	J123527.0+212034	12 35 27.03	+21 20 34.8 v	5.0	d (int corduroy effect '1' diameter, artifact?)
864	NVGR	J123621.5+255904	12 36 21.58	+25 59 04.3 c	10.0	rc
865	NVGR	J123631.5+163229	12 36 31.50	+16 32 29.0 v	6.0	wat? sz?
866	NVGR	J123725.4+045729	12 37 25.48	+04 57 29.4 v	3.5	rc? wat?
867	NVGR	J123748.0-155807	12 37 48.03	-15 58 07.9 v	8.5	wat/nat
868			12 38 47.82	+44 09 35.3 c	6.5	t (Soloviyov & Verkhodanov(2014): J123846.9+440949)
869	NVGR	J123854.5-172700	12 38 54.57	-17 27 00.0 v	6.0	w? sz?
870	NVGR	J123908.8+001838	12 39 08.80	+00 18 38.1 v	4.5	d t?
871	NVGR	J123944.2+195436	12 39 44.25	+19 54 36.2 v	4.5	d? rc?
872	NVGR	J124013.7+533433	12 40 13.71	+53 34 33.3 c	4.5	d t
873	NVGR	J124056.7+183058	12 40 56.78	+18 30 58.2 v	6.5	cj? or lobe of possible 60' d?
874			12 42 04.72	+32 32 40.5 v	12.0	rc t? core id amb (Soloviyov & Verkhodanov(2014): J124159.1+323217)
875	NVGR	J124355.7+130745	12 43 55.76	+13 07 45.1 c	4.0	rc - incidental
876	NVGR	J124619.1+262650	12 46 19.16	+26 26 50.5 v	7.0	wat (with cp?)
877			12 47 33.31	+67 23 16.8 c	14.0	t (Lara et al.(2001): J1247+673)
878	NVGR	J124736.6-155947	12 47 36.62	-15 59 47.7 c	5.0	t
879	NVGR	J124736.7-290319	12 47 36.78	-29 03 19.5 v	4.0	d?
880	NVGR	J124819.9-214743	12 48 19.90	-21 47 43.3 v	6.5	w?
881	NVGR	J125053.4+410708	12 50 53.41	+41 07 08.6 c	5.0	rc - incidental
882			12 50 58.34	+75 38 02.0 v	5.0	nat? (Lara et al.(2001): J1251+756)
883	NVGR	J125153.4+351924	12 51 53.46	+35 19 24.7 c	5.0	d
884	NVGR	J125204.4-222601	12 52 04.47	-22 26 01.8 v	5.0	sz? core id amb
885	NVGR	J125258.4-152014	12 52 58.47	-15 20 14.9 v	5.5	wat/nat? cp? irr
886	NVGR	J125305.9+450025	12 53 05.91	+45 00 25.7 v	10.0	w?
887			12 53 06.31	-10 29 05.6 c	16.0	w? nat? (Soloviyov & Verkhodanov(2014): J125310.7-103115)
888	NVGR	J125309.8-013930	12 53 09.83	-01 39 30.0 v	5.5	d?
889			12 53 10.92	+40 41 22.5 v	6.5	d (Machalski et al.(2001): J1253+4041)
890	NVGR	J125321.1-154737	12 53 21.14	-15 47 37.6 c	6.0	t
891	NVGR	J125421.9-290247	12 54 21.92	-29 02 47.6 v	7.5	d?

892	NVGR	J125457.7-311025	12 54 57.76	-31 10 25.2 v	4.0	cp? cj? d?
893	NVGR	J125530.6-133822	12 55 30.61	-13 38 22.1 c	5.5	wat?
894	NVGR	J125604.0+225505	12 56 04.03	+22 55 05.3 v	4.5	cp point with elliptic
895	NVGR	J125607.6-271700	12 56 07.67	-27 17 00.5 v	4.0	d? points
896	NVGR	J125659.5+061858	12 56 59.57	+06 18 58.7 c	6.5	wat
897	NVGR	J125734.1-221607	12 57 34.17	-22 16 07.3 c	8.0	t int ring-lobes hooks
898	NVGR	J125808.6+421110	12 58 08.64	+42 11 10.5 v	5.0	d
899	NVGR	J125914.3+275441	12 59 14.33	+27 54 41.5 v	6.0	cj d?
900	NVGR	J130025.3-033704	13 00 25.35	-03 37 04.0 v	5.0	cp two elliptic not aligned
901	NVGR	J130026.6-092930	13 00 26.68	-09 29 30.1 v	4.0	d
902	NVGR	J130153.0+144907	13 01 53.04	+14 49 07.6 v	5.0	wat? d?
903	NVGR	J130202.8-005101	13 02 02.88	-00 51 01.5 v	6.5	wat? core id amb
904	NVGR	J130422.2-085254	13 04 22.20	-08 52 54.7 v	6.5	mg irr cp int cluster
905	NVGR	J130458.8-324920	13 04 58.85	-32 49 20.5 v	8.5	sz? butterfly
906	NVGR	J130512.2-304315	13 05 12.28	-30 43 15.2 v	4.0	d? rc?
907	NVGR	J130557.2-123545	13 05 57.28	-12 35 45.2 v	10.5	d t?
908	NVGR	J130731.8+520653	13 07 31.80	+52 06 53.3 v	4.0	d
909	NVGR	J130949.4+193907	13 09 49.49	+19 39 07.4 v	5.5	w? sz?
910	NVGR	J131143.3+405855	13 11 43.35	+40 58 55.5 v	8.0	d sz?
911	NVGR	J131231.2+211537	13 12 31.22	+21 15 37.8 v	5.0	d
912			13 14 05.52	+69 36 57.5 v	8.5	w? dd?? 4 components (Lara et al.(2001): J1313+696)
913	NVGR	J131536.0-194227	13 15 36.09	-19 42 27.0 c	5.5	t?
914	NVGR	J131548.4+420147	13 15 48.48	+42 01 47.3 c	7.5	rc M 63
915	NVGR	J131604.6-151043	13 16 04.63	-15 10 43.9 v	5.0	cp ~point and teardrop
916	NVGR	J131617.5+070237	13 16 17.53	+07 02 37.1 v	9.0	nat/wat
917	NVGR	J131824.7+094550	13 18 24.78	+09 45 50.3 c	5.0	d/t
918	NVGR	J131903.6+293829	13 19 03.64	+29 38 29.9 v	4.5	rc sz 4C 29.47
919	NVGR	J131920.8-005526	13 19 20.87	-00 55 26.8 v	5.0	nat
920			13 23 47.51	+31 33 41.7 v	12.0	w? rc? jets wisps int (Solovoyov & Verkhodanov(2014):? J132345.8+313402)
921	NVGR	J132403.3+433355	13 24 03.39	+43 33 55.2 v	4.0	d no core fit
922	NVGR	J132615.4-272551	13 26 15.41	-27 25 51.4 v	5.5	nat? cp?
923	NVGR	J132644.1-272621	13 26 44.11	-27 26 21.5 c	5.5	t?
924	NVGR	J132731.6+185418	13 27 31.66	+18 54 18.2 v	4.5	d
925	NVGR	J132736.3+350641	13 27 36.34	+35 06 41.6 v	5.5	d
926	NVGR	J132834.5-012915	13 28 34.53	-01 29 15.7 v	7.0	w
927	NVGR	J132846.9+275249	13 28 46.98	+27 52 49.2 v	5.0	t?
928	NVGR	J132853.3-395919	13 28 53.38	-39 59 19.9 v	4.5	d asym
929	NVGR	J132952.6+471145	13 29 52.65	+47 11 45.7 c	10.0	rc
930	NVGR	J133131.6+453918	13 31 31.60	+45 39 18.9 v	3.5	cp 3.5' t + 3.5' d
931	NVGR	J133242.4+071937	13 32 42.41	+07 19 37.8 c	8.5	wat/nat core id amb
932	NVGR	J133418.5-100929	13 34 18.56	-10 09 29.3 c	17.0	t
933	NVGR	J133447.8-270738	13 34 47.82	-27 07 38.1 c	8.0	wat? t? d? core id amb
934	NVGR	J133552.4-285519	13 35 52.48	-28 55 19.6 c	4.0	wat
935	NVGR	J133608.9-083054	13 36 08.94	-08 30 54.5 v	8.0	mg dd? int unusual config
936	NVGR	J133632.9+403725	13 36 32.97	+40 37 25.2 v	5.5	d? elliptics
937	NVGR	J133639.3-335759	13 36 39.37	-33 57 59.6 v	34.5	wat separate jets and lobes
938	NVGR	J133700.5-295155	13 37 00.52	-29 51 55.1 c	8.5	rc int spiral arms, also trough
939	NVGR	J133738.3+294216	13 37 38.31	+29 42 16.4 v	6.0	mg wat?
940	NVGR	J133855.0+241735	13 38 55.06	+24 17 35.0 c	4.5	cj? d?
941	NVGR	J134245.6+064256	13 42 45.62	+06 42 56.4 v	5.0	d? (two elliptic, not aligned)
942	NVGR	J134348.8-194930	13 43 48.85	-19 49 30.7 c	4.5	rc
943	NVGR	J134415.6+331716	13 44 15.66	+33 17 16.0 v	4.0	t d
944			13 44 46.23	+40 26 39.4 v	9.0	mg 5' wat? and 3.5' t? (Machalski et al.(2001): J1344+4028)
945			13 45 03.90	+39 52 37.8 v	5.0	d (Machalski et al.(2001): J1345+3952)
946	NVGR	J134554.6+540344	13 45 54.60	+54 03 44.6 c	6.5	t
947	NVGR	J134601.9-161656	13 46 01.92	-16 16 56.5 v	5.5	cp?
948	NVGR	J134642.1-253745	13 46 42.13	-25 37 45.9 v	5.0	d
949	NVGR	J134713.7-242223	13 47 13.76	-24 22 23.6 v	6.5	rc dd??
950	NVGR	J134729.4-134645	13 47 29.44	-13 46 45.2 c	4.5	mg nat? cp?
951	NVGR	J134759.8-131333	13 47 59.87	-13 13 33.7 v	4.5	d
952	NVGR	J135036.1-163449	13 50 36.16	-16 34 49.6 c	13.0	mg t hymors?
953	NVGR	J135106.6+072900	13 51 06.63	+07 29 00.7 v	5.0	mg
954	NVGR	J135144.9-054626	13 51 44.97	-05 46 26.9 v	17.0	cp
955	NVGR	J135217.8+312646	13 52 17.81	+31 26 46.7 c	6.0	rc?
956	NVGR	J135229.0+093543	13 52 29.00	+09 35 43.3 c	6.0	rc? d?
957	NVGR	J135306.2+330315	13 53 06.27	+33 03 15.8 v	7.0	sz? or perhaps wat/nat at N end
958	NVGR	J135420.7-070520	13 54 20.75	-07 05 20.7 v	5.5	d
959			13 55 17.92	+29 23 41.2 v	6.0	d (Machalski et al.(2001): J1355+2923)
960	NVGR	J135618.5+181726	13 56 18.52	+18 17 26.1 c	7.5	sz?
961	NVGR	J135956.5-105623	13 59 56.56	-10 56 23.1 v	4.5	wat/nat?
962	NVGR	J140000.2+282953	14 00 00.26	+28 29 53.8 v	4.0	d (with cp?) wat/nat?
963	NVGR	J140043.0+301800	14 00 43.00	+30 18 00.0 v	12.0	d?
964	NVGR	J140058.8+591929	14 00 58.82	+59 19 29.6 v	5.5	cp point and elliptic
965	NVGR	J140112.2-131433	14 01 12.21	-13 14 33.8 v	4.5	wat?
966	NVGR	J140140.9-113614	14 01 40.97	-11 36 14.2 c	9.0	t sz? hymors?
967	NVGR	J140305.6+004822	14 03 05.64	+00 48 22.6 v	4.0	d? points
968	NVGR	J140313.9+514447	14 03 13.91	+51 44 47.1 v	5.5	d dd?
969	NVGR	J140314.7+061126	14 03 14.70	+06 11 26.2 v	5.0	nat/wat? lobe? with hook, cp?

970	NVGR	J140319.8+252545	14 03 19.87	+25 25 45.2 v	5.5	d? point and irr
971	NVGR	J140326.5-060149	14 03 26.56	-06 01 49.7 c	7.5	mg wat?, may be smaller wat
972	NVGR	J140518.0+043545	14 05 18.01	+04 35 45.8 v	5.0	cp
973	NVGR	J140519.5-275509	14 05 19.56	-27 55 09.6 v	5.0	nat? sz?
974	NVGR	J140700.4+100836	14 07 00.47	+10 08 36.8 v	5.5	d? butterfly? nat??
975	NVGR	J140704.9+360117	14 07 04.97	+36 01 17.5 v	8.0	d? sz?
976	NVGR	J140719.1+513211	14 07 19.17	+51 32 11.1 c	4.5	d t core fit but not apparent
977			14 09 57.38	+17 32 44.1 c	8.0	t? (with cp?) w? (Solovyov & Verkhodanov(2014): J140955.9+173243)
978			14 10 30.25	+63 19 14.8 v	4.5	d (Lara et al.(2001): J1410+633)
979	NVGR	J141634.0-364036	14 16 34.00	-36 40 36.8 v	5.5	wat? rc? sz?
980	NVGR	J141653.5+104840	14 16 53.50	+10 48 40.2 c	8.5	nat?
981	NVGR	J141838.9-332800	14 18 38.97	-33 28 00.5 v	4.5	d asym
982	NVGR	J141913.6-310153	14 19 13.61	-31 01 53.1 v	3.5	d
983	NVGR	J141946.8+081421	14 19 46.88	+08 14 21.7 v	4.0	d sz?
984	NVGR	J142008.7+450055	14 20 08.74	+45 00 55.1 v	8.0	mg int
985	NVGR	J142251.7-092456	14 22 51.70	-09 24 56.0 v	4.5	cp? d?
986	NVGR	J142301.2-051428	14 23 01.22	-05 14 28.6 v	6.0	d (with cp?) bridge apparent
987	NVGR	J142326.3+244911	14 23 26.34	+24 49 11.7 v	5.5	cp d with teardrop
988	NVGR	J142420.3-164434	14 24 20.32	-16 44 34.9 v	6.5	d? two elliptic
989	NVGR	J142452.7-192248	14 24 52.74	-19 22 48.8 v	13.5	t
990			14 25 54.59	-08 04 00.9 v	7.5	d (Solovyov & Verkhodanov(2014): J142554.6-080401)
991	NVGR	J142622.6+585449	14 26 22.67	+58 54 49.6 v	6.0	cp? d with 'point at west end
992	NVGR	J142720.4+163438	14 27 20.49	+16 34 38.3 v	4.5	d? cp?
993	NVGR	J142727.2-240722	14 27 27.29	-24 07 22.4 c	4.0	rc? (with cp?)
994	NVGR	J142735.5+263210	14 27 35.51	+26 32 10.3 c	6.0	t?
995			14 28 19.46	+29 18 45.3 c	17.5	wat (Machalski et al.(2001): J1428+2918)
996			14 28 45.60	+39 38 44.3 c	5.5	t (Machalski et al.(2001): J1428+3938)
997	NVGR	J142926.6+544341	14 29 26.63	+54 43 41.2 v	6.0	sz? wat?
998	NVGR	J142950.9+202223	14 29 50.96	+20 22 23.6 v	10.0	d? hymors? N component ringlike
999	NVGR	J142954.7+071919	14 29 54.75	+07 19 19.2 c	13.5	w? nat? core id amb core
1000	NVGR	J143308.4-183405	14 33 08.43	-18 34 05.7 v	7.5	mg
1001	NVGR	J143406.5+013557	14 34 06.59	+01 35 57.4 v	5.0	wat/nat
1002	NVGR	J143506.1-161427	14 35 06.11	-16 14 27.3 c	5.5	t sz?
1003	NVGR	J143549.9+201339	14 35 49.96	+20 13 39.9 v	4.5	d
1004	NVGR	J143639.3-202354	14 36 39.39	-20 23 54.5 v	5.0	d? points
1005	NVGR	J143642.0+080713	14 36 42.08	+08 07 13.2 v	6.5	d
1006	NVGR	J143850.4-140842	14 38 50.46	-14 08 42.2 v	6.0	d
1007	NVGR	J144020.1+055411	14 40 20.15	+05 54 11.3 v	6.0	nat? sz?
1008	NVGR	J144152.6+340117	14 41 52.61	+34 01 17.8 v	5.0	rc? wat? w?
1009	NVGR	J144255.4+520934	14 42 55.44	+52 09 34.5 v	6.0	d? mg
1010	NVGR	J144406.2+260127	14 44 06.20	+26 01 27.6 v	5.0	d
1011	NVGR	J144429.1-400147	14 44 29.15	-40 01 47.3 c	5.0	cj? - incidental
1012	NVGR	J144433.0+275259	14 44 33.02	+27 52 59.1 v	4.5	d
1013	NVGR	J144626.0+214151	14 46 26.01	+21 41 51.5 v	4.0	wat d
1014	NVGR	J144722.4+424225	14 47 22.44	+42 42 25.7 v	6.0	d?
1015	NVGR	J144823.8+442339	14 48 23.88	+44 23 39.5 v	5.5	d?
1016	NVGR	J144853.7-400611	14 48 53.77	-40 06 11.0 v	9.0	mg cp?
1017			14 51 32.16	+33 57 55.4 c	5.5	t (Machalski et al.(2001): J1451+3357)
1018	NVGR	J145204.7+385810	14 52 04.74	+38 58 10.1 v	8.0	rc? cj? sz? unu
1019	NVGR	J145228.7-242946	14 52 28.74	-24 29 46.7 v	5.5	mg cp? d? elliptic with point(s)
1020	NVGR	J145235.6-131119	14 52 35.62	-13 11 19.6 v	5.5	d
1021			14 53 02.99	+33 08 55.8 c	7.5	t (Machalski et al.(2001): J1453+3309)
1022	NVGR	J145320.3+015217	14 53 20.39	+01 52 17.8 v	5.5	d? cp?
1023	NVGR	J145345.3+224313	14 53 45.38	+22 43 13.9 v	5.0	d core fit but not apparent
1024	NVGR	J145411.7-304528	14 54 11.78	-30 45 28.6 c	4.5	cj?
1025	NVGR	J145421.9+162127	14 54 21.91	+16 21 27.4 v	8.0	d t? dd? 4 or more components
1026	NVGR	J145448.7+122456	14 54 48.70	+12 24 56.4 c	4.5	t (with adjacent t?)
1027	NVGR	J145534.9+233534	14 55 34.94	+23 35 34.8 v	9.0	cp d? elliptic and irr
1028	NVGR	J145614.7+181830	14 56 14.72	+18 18 30.1 v	4.5	cp elliptic with d?
1029	NVGR	J145631.3-241327	14 56 31.32	-24 13 27.6 v	5.0	d? points
1030	NVGR	J145736.0+250256	14 57 36.03	+25 02 56.2 v	12.0	cp wat? elliptic
1031			14 57 46.96	-06 12 58.4 c	6.0	d t? 4 components (Machalski et al.(2001): J1457-0613)
1032	NVGR	J145753.6+283213	14 57 53.62	+28 32 13.2 v	5.5	d
1033	NVGR	J145753.7-212633	14 57 53.77	-21 26 33.7 v	5.5	cp
1034	NVGR	J145950.2+185240	14 59 50.21	+18 52 40.8 v	5.0	sz?
1035	NVGR	J150023.2+201235	15 00 23.25	+20 12 35.8 v	5.0	d sz?
1036	NVGR	J150147.3+434611	15 01 47.39	+43 46 11.0 c	5.0	d t sz?
1037	NVGR	J150336.8-295621	15 03 36.83	-29 56 21.5 c	5.0	d t?
1038	NVGR	J150339.5+101602	15 03 39.51	+10 16 02.8 c	5.5	sz
1039	NVGR	J150354.6-281119	15 03 54.61	-28 11 19.0 c	11.5	wat?
1040			15 04 12.51	+68 56 14.8 c	5.5	mg 4.5' d and 3.5' d (Lara et al.(2001): J1504+689)
1041	NVGR	J150419.5+283534	15 04 19.50	+28 35 34.3 v	7.5	nat? rc? arc
1042	NVGR	J150458.4+260132	15 04 58.43	+26 01 32.9 c	7.0	rc
1043	NVGR	J150505.5+261030	15 05 05.50	+26 10 30.0 v	3.5	wat? faint - incidental
1044	NVGR	J150725.7+082915	15 07 25.77	+08 29 15.9 v	8.0	sz? dl
1045	NVGR	J150831.9+355432	15 08 31.94	+35 54 32.4 v	4.0	d?
1046	NVGR	J150911.0-353224	15 09 11.00	-35 32 24.6 v	4.5	wat?
1047	NVGR	J150912.0-064020	15 09 12.01	-06 40 20.5 v	7.0	cp point with elliptic

1048	NVGR	J150916.3+094846	15 09 16.32	+09 48 46.1 v	4.0	t cat missed core?
1049	NVGR	J150959.5+030004	15 09 59.53	+03 00 04.4 c	7.0	sz? nat?
1050	NVGR	J151048.7-042535	15 10 48.72	-04 25 35.3 v	4.5	wat?
1051	NVGR	J151128.2+071110	15 11 28.23	+07 11 10.7 v	11.5	d? with bridge cj?
1052			15 12 05.80	+30 50 22.1 c	6.0	t (Machalski et al.(2001): J1512+3050)
1053	NVGR	J151304.1-225812	15 13 04.11	-22 58 12.2 c	4.0	d t
1054	NVGR	J151308.9-252315	15 13 08.93	-25 23 15.9 v	6.0	mg t? wat? GRS or two?
1055	NVGR	J151329.6-381635	15 13 29.68	-38 16 35.9 v	6.5	t cat missed core?
1056	NVGR	J151341.0-323349	15 13 41.07	-32 33 49.8 v	4.0	d?
1057	NVGR	J151356.7+241255	15 13 56.72	+24 12 55.0 v	8.0	sz? t? cp? cat missed core?
1058	NVGR	J151454.4+420057	15 14 54.46	+42 00 57.0 v	3.5	d
1059	NVGR	J151540.0-193931	15 15 40.03	-19 39 31.2 v	4.0	d
1060	NVGR	J151551.4+562006	15 15 51.48	+56 20 06.8 v	8.0	rc? (with cp?) asym
1061	NVGR	J151604.5-381858	15 16 04.54	-38 18 58.0 v	19.0	dd? sz? int or 5' arc and 5.5' cj?
1062	NVGR	J151640.2+001502	15 16 40.21	+00 15 02.4 c	6.5	t sz? rc? wisps
1063	NVGR	J151657.5+051838	15 16 57.57	+05 18 38.0 v	6.0	d? (with cp?) unu
1064	NVGR	J151743.4+285343	15 17 43.42	+28 53 43.6 v	4.5	mg cp wat?
1065	NVGR	J151836.3-240545	15 18 36.32	-24 05 45.6 v	6.5	rc? lobe?
1066	NVGR	J151853.6-101406	15 18 53.61	-10 14 06.8 v	4.0	d? points
1067	NVGR	J152116.2+510458	15 21 16.25	+51 04 58.4 c	6.5	t dd?? x?
1068	NVGR	J152139.4+222727	15 21 39.47	+22 27 27.2 v	5.0	d? points
1069	NVGR	J152309.6-172039	15 23 09.65	-17 20 39.2 v	4.5	d
1070	NVGR	J152325.5-285128	15 23 25.58	-28 51 28.0 v	5.5	d 4 components
1071	NVGR	J152329.3+283651	15 23 29.36	+28 36 51.5 v	8.0	wat? (with cp?)
1072			15 23 45.74	+63 39 18.7 v	5.0	rc? (Lara et al.(2001): J1523+636)
1073	NVGR	J152613.7+170109	15 26 13.77	+17 01 09.1 v	5.5	cp (two elliptic not aligned)
1074	NVGR	J152630.2+272452	15 26 30.26	+27 24 52.4 v	5.0	d
1075	NVGR	J152805.0+054428	15 28 05.02	+05 44 28.5 c	16.5	t? sz? wat? wispy lobes asym
1076	NVGR	J153009.9+290040	15 30 09.90	+29 00 40.6 v	5.5	d with bridge
1077	NVGR	J153035.4-353534	15 30 35.42	-35 35 34.0 v	5.0	mg d? points
1078	NVGR	J153228.9-381834	15 32 28.92	-38 18 34.0 v	4.5	d/t
1079	NVGR	J153429.3-221756	15 34 29.31	-22 17 56.7 v	5.5	d? points
1080	NVGR	J153636.7-020315	15 36 36.71	-02 03 15.6 v	7.5	cp elliptic with irr/arc
1081			15 37 03.88	+84 23 17.0 c	9.5	w? dd? x? (Lara et al.(2001): J1536+843)
1082	NVGR	J153850.0-280500	15 38 50.08	-28 05 00.2 v	4.5	mg wat?
1083	NVGR	J153853.8+170138	15 38 53.87	+17 01 38.1 c	13.0	t? sz? int ring-lobe and two ring-lobe pretzel-lobe?
1084	NVGR	J153958.8+483539	15 39 58.88	+48 35 39.7 v	4.0	d
1085	NVGR	J154011.9-232435	15 40 11.93	-23 24 35.3 v	5.5	d
1086	NVGR	J154025.0-052840	15 40 25.00	-05 28 40.7 v	6.0	d? cp? fuzzy point and elliptic
1087	NVGR	J154057.4-012636	15 40 57.49	-01 26 36.3 v	7.0	mg sz? nat? dl
1088	NVGR	J154130.3-253440	15 41 30.38	-25 34 40.2 v	5.5	d? wat/nat?
1089	NVGR	J154132.1-270744	15 41 32.13	-27 07 44.8 v	4.5	wat/nat? may be larger
1090	NVGR	J154149.7-365943	15 41 49.76	-36 59 43.5 v	9.0	d? cp?
1091	NVGR	J154640.1+012358	15 46 40.12	+01 23 58.9 v	4.5	wat/nat?
1092	NVGR	J154818.4+072558	15 48 18.48	+07 25 58.5 c	6.5	w? sz? t?
1093			15 48 57.72	-32 16 47.0 v	10.0	x? w? (Solov'yov & Verkhodanov(2014): J154901.7-321747)
1094	NVGR	J154858.3+085431	15 48 58.34	+08 54 31.9 c	5.0	rc? cj?
1095	NVGR	J155113.1-221304	15 51 13.14	-22 13 04.8 v	4.0	d points
1096	NVGR	J155135.6+200406	15 51 35.64	+20 04 06.0 c	10.5	mg t? w?
1097	NVGR	J155139.8+103553	15 51 39.86	+10 35 53.9 v	4.0	d
1098	NVGR	J155141.3-210948	15 51 41.38	-21 09 48.1 v	5.0	d
1099	NVGR	J155209.0+200530	15 52 09.00	+20 05 30.0 v	13.5	mg d
1100	NVGR	J155210.9+224508	15 52 10.97	+22 45 08.0 c	13.5	t asym int
1101	NVGR	J155216.2-060219	15 52 16.27	-06 02 19.2 v	4.0	d? points
1102	NVGR	J155220.1+361404	15 52 20.14	+36 14 04.2 v	4.0	sz?
1103	NVGR	J155235.1+223418	15 52 35.17	+22 34 18.8 v	13.0	t dd? asym
1104	NVGR	J155727.3+544132	15 57 27.30	+54 41 32.8 v	10.0	mg wat/nat w? interacting systems? very twisted int
1105			15 57 29.63	+70 41 35.8 c	12.0	sz? nat? int hook (Lara et al.(2001): J1557+706)
1106	NVGR	J155902.5-213857	15 59 02.53	-21 38 57.1 v	10.0	wat? GRS or two?
1107	NVGR	J160028.7+083752	16 00 28.72	+08 37 52.9 v	4.5	rc? w?
1108	NVGR	J160121.7+460042	16 01 21.79	+46 00 42.2 c	5.0	t
1109	NVGR	J160303.9-184455	16 03 03.97	-18 44 55.1 v	5.0	d? elliptic
1110	NVGR	J160312.5-131945	16 03 12.54	-13 19 45.9 v	5.5	d asym
1111	NVGR	J160332.1+171158	16 03 32.13	+17 11 58.9 v	8.0	mg 8' nat/wat or 6' wat/nat with fainter 6' wat/nat?
1112	NVGR	J160337.9+524228	16 03 37.93	+52 42 28.8 v	4.0	cp
1113	NVGR	J160349.2+251038	16 03 49.28	+25 10 38.4 v	5.5	cp two elliptic non-aligned
1114	NVGR	J160417.6-015428	16 04 17.66	-01 54 28.6 c	7.0	rc
1115	NVGR	J160435.8+252028	16 04 35.83	+25 20 28.8 v	4.5	d? cj? rc? asym
1116			16 04 46.27	+34 38 12.1 v	5.0	d (Machalski et al.(2001): J1604+3438)
1117	NVGR	J160500.0+183602	16 05 00.03	+18 36 02.1 v	4.5	d?
1118	NVGR	J160509.8-334405	16 05 09.83	-33 44 05.4 c	4.5	d/t
1119	NVGR	J160513.8+071143	16 05 13.82	+07 11 43.4 v	4.5	rc? d?
1120	NVGR	J160531.3-092738	16 05 31.35	-09 27 38.8 v	7.0	w? sz? 4 components
1121	NVGR	J160820.7-233356	16 08 20.70	-23 33 56.1 v	5.0	d sz?
1122	NVGR	J161034.6-051116	16 10 34.66	-05 11 16.0 v	6.0	nat? sz? d?
1123	NVGR	J161051.4-053602	16 10 51.45	-05 36 02.1 v	6.0	d
1124	NVGR	J161157.4+073855	16 11 57.45	+07 38 55.1 c	7.0	t asym
1125	NVGR	J161241.3+431320	16 12 41.34	+43 13 20.6 c	6.5	d sz? core id amb

1126	NVGR	J161419.4-285503	16 14 19.44	-28 55 03.6 v	6.0	d four component
1127	NVGR	J161521.2-170845	16 15 21.23	-17 08 45.1 v	6.0	d? elliptic
1128	NVGR	J161534.8+095715	16 15 34.80	+09 57 15.4 c	5.0	sz? core id amb
1129	NVGR	J161713.0+063729	16 17 13.03	+06 37 29.1 v	3.0	d rc
1130	NVGR	J161724.9+350453	16 17 24.92	+35 04 53.6 v	12.0	sz w? may be larger
1131	NVGR	J161829.7+295902	16 18 29.76	+29 59 02.0 v	4.5	d wat? sz? not original pair
1132	NVGR	J161830.9-041033	16 18 30.98	-04 10 33.1 v	5.5	d? point and elliptic
1133	NVGR	J161839.5+443607	16 18 39.50	+44 36 07.9 v	6.5	d? two aligned elliptic
1134	NVGR	J162006.7-253002	16 20 06.74	-25 30 02.1 v	13.0	sz 4 or 5 components
1135	NVGR	J162344.4+252301	16 23 44.47	+25 23 01.1 c	7.0	t core id amb
1136	NVGR	J162549.9+402921	16 25 49.96	+40 29 21.0 c	7.0	t? asym lobes, may be smaller d
1137	NVGR	J162610.8+052802	16 26 10.89	+05 28 02.5 c	5.0	d/t
1138	NVGR	J162730.5+211318	16 27 30.56	+21 13 18.2 v	4.5	cp fainter 4.5' d + point?
1139	NVGR	J162748.8+514226	16 27 48.82	+51 42 26.6 v	7.0	t? core id amb, may be larger, fades
1140			16 29 01.72	+82 36 05.8 v	18.0	mg sz? (Lara et al.(2001): J1632+825)
1141	NVGR	J163134.2-064642	16 31 34.21	-06 46 42.9 c	3.0	rc? cj? - incidental
1142	NVGR	J163145.0-065051	16 31 45.09	-06 50 51.3 v	5.0	wat? d? core not apparent
1143	NVGR	J163534.6+150258	16 35 34.65	+15 02 58.0 v	5.0	d?
1144	NVGR	J163621.6+491325	16 36 21.68	+49 13 25.5 c	6.0	wat? sz?
1145	NVGR	J163622.6-032216	16 36 22.69	-03 22 16.5 v	5.0	cp nat?
1146	NVGR	J163636.0+333504	16 36 36.03	+33 35 04.8 v	4.0	wat?
1147	NVGR	J163706.7+022435	16 37 06.79	+02 24 35.8 v	4.5	d?
1148	NVGR	J163747.0-034500	16 37 47.08	-03 45 00.4 v	13.5	cp
1149	NVGR	J163801.9+375251	16 38 01.90	+37 52 51.9 v	4.0	rc d
1150	NVGR	J163923.4+295159	16 39 23.40	+29 51 59.6 v	4.5	nat? shell? wispy fingers
1151	NVGR	J164042.1-264030	16 40 42.10	-26 40 30.1 v	5.5	cp wat?
1152	NVGR	J164610.9-130927	16 46 10.96	-13 09 27.2 v	6.0	d? cp? irr
1153	NVGR	J164626.3+281227	16 46 26.34	+28 12 27.9 v	4.0	d
1154	NVGR	J164747.3+190251	16 47 47.39	+19 02 51.4 v	5.0	cp point with elliptic
1155	NVGR	J164852.5-280459	16 48 52.55	-28 04 59.9 v	5.0	d? elliptic
1156	NVGR	J164938.7+013833	16 49 38.70	+01 38 33.9 c	5.0	t?
1157			16 50 49.60	+81 33 37.4 v	6.0	d? asym, low level ring? (Lara et al.(2001): J1650+815)
1158	NVGR	J165346.2+010015	16 53 46.27	+01 00 15.3 v	5.5	cp? wat/nat? cj with cj?
1159	NVGR	J165358.1-051017	16 53 58.17	-05 10 17.9 c	6.5	sz? nat?
1160	NVGR	J165638.1-090452	16 56 38.12	-09 04 52.9 v	4.0	wat/nat? rc?
1161	NVGR	J165645.8-331248	16 56 45.85	-33 12 48.6 v	7.5	mg w? cp? GRS or 2
1162	NVGR	J165719.3+534012	16 57 19.37	+53 40 12.4 v	4.5	d? (with cp?)
1163	NVGR	J165730.1-224419	16 57 30.18	-22 44 19.6 c	9.5	nat? t? with bridge/jets
1164	NVGR	J170202.5+531234	17 02 02.50	+53 12 34.0 c	5.0	t core fit but not apparent
1165	NVGR	J170255.9+421721	17 02 55.97	+42 17 21.6 v	5.0	d 4 components
1166	NVGR	J170312.9+313224	17 03 12.95	+31 32 24.2 v	4.0	d?
1167	NVGR	J170326.1-094127	17 03 26.15	-09 41 27.2 v	5.5	t? asym cp?
1168			17 05 00.17	-24 45 25.8 c	8.0	t sz? (Solovyov & Verkhodanov(2014): J170502.0-24460)
1169	NVGR	J170548.5-230716	17 05 48.58	-23 07 16.5 v	5.5	d points, W source does have finger toward other source
1170	NVGR	J170550.3-033112	17 05 50.38	-03 31 12.0 v	7.0	nat/wat?
1171	NVGR	J170628.0+102450	17 06 28.07	+10 24 50.7 v	5.5	d/t
1172	NVGR	J170727.8+223817	17 07 27.80	+22 38 17.8 v	4.5	d
1173	NVGR	J170825.7-123311	17 08 25.77	-12 33 11.7 v	5.0	d/t
1174	NVGR	J170833.4-113816	17 08 33.45	-11 38 16.8 c	6.5	t
1175	NVGR	J170841.4+415022	17 08 41.42	+41 50 22.5 v	5.0	d? two points, int alignment of 4 or more sources
1176	NVGR	J170927.5-222640	17 09 27.58	-22 26 40.9 c	12.5	nat int
1177	NVGR	J170931.9+225211	17 09 31.99	+22 52 11.4 v	5.5	mg cp? d?
1178	NVGR	J171008.6-353004	17 10 08.64	-35 30 04.2 c	5.0	d? sz? nat? dl? cj?
1179	NVGR	J171057.3+462416	17 10 57.32	+46 24 16.6 c	11.5	t int lobe structure hymors? ring-lobe with fingers
1180	NVGR	J171057.6+394135	17 10 57.62	+39 41 35.3 c	7.0	cj? fishtail int
1181	NVGR	J171136.5-225900	17 11 36.50	-22 59 00.0 v	4.0	cj? N lobe of sz? - incidental
1182	NVGR	J171145.0-230430	17 11 45.00	-23 04 30.0 v	13.0	sz? cp?
1183	NVGR	J171150.4-230821	17 11 50.41	-23 08 21.8 v	6.0	cj? rc? or S lobe of sz?
1184	NVGR	J171229.9-231319	17 12 29.96	-23 13 19.7 c	5.5	rc t
1185	NVGR	J171240.0-090236	17 12 40.06	-09 02 36.7 v	5.0	cp
1186	NVGR	J171301.1+205144	17 13 01.11	+20 51 44.9 v	5.5	cp nat?
1187	NVGR	J171308.2+043556	17 13 08.27	+04 35 56.2 v	4.5	cp point with teardrop
1188	NVGR	J171524.9-042546	17 15 24.91	-04 25 46.5 v	5.0	wat? rc? cp?
1189	NVGR	J171541.1-120621	17 15 41.19	-12 06 21.7 c	6.5	sz
1190	NVGR	J171813.9+153907	17 18 13.90	+15 39 07.0 v	4.0	d
1191	NVGR	J171819.5+062731	17 18 19.50	+06 27 31.8 v	5.0	wat/nat? cp? possibly larger system
1192	NVGR	J172104.4+262417	17 21 04.45	+26 24 17.9 c	5.5	t
1193	NVGR	J172206.2+330659	17 22 06.23	+33 06 59.4 v	4.5	cp? asym
1194	NVGR	J172209.9+092036	17 22 09.91	+09 20 36.5 v	4.5	d
1195	NVGR	J172320.8+341757	17 23 20.85	+34 17 57.3 c	6.0	t sz?
1196	NVGR	J172508.0-332518	17 25 08.00	-33 25 18.4 v	45.0	rc? jets? irr linear structure, artifact?
1197	NVGR	J172749.9+534647	17 27 49.93	+53 46 47.0 v	4.5	d cj?
1198	NVGR	J172854.7-225139	17 28 54.71	-22 51 39.0 v	6.0	d?
1199	NVGR	J172858.2+084858	17 28 58.21	+08 48 58.1 c	4.0	cj
1200	NVGR	J172958.2+450621	17 29 58.26	+45 06 21.3 c	4.0	t rc
1201	NVGR	J173106.8+242859	17 31 06.82	+24 28 59.9 c	4.5	d/t
1202	NVGR	J173112.1+225015	17 31 12.10	+22 50 15.9 v	4.5	mg cp?
1203	NVGR	J173215.3+170605	17 32 15.30	+17 06 05.3 c	6.0	sz

1204		17 32 32.75	+71 24 03.9 c	5.0	t (Lara et al.(2001): J1732+714)
1205	NVGR	17 32 59.05	+56 34 16.2 v	5.0	cp
1206		17 33 12.89	+70 46 32.5 c	6.0	t sz? (Lara et al.(2001): J1733+707)
1207	NVGR	17 33 27.32	-04 45 40.0 c	4.5	wat?
1208	NVGR	17 35 21.80	-08 02 39.2 v	5.5	d
1209	NVGR	17 36 41.94	-20 48 43.3 c	5.5	wat
1210	NVGR	17 37 42.13	-18 32 40.9 c	8.0	t? d? irr hymors? cp?
1211	NVGR	17 38 57.42	+13 00 34.1 v	5.0	cp point with teardrop
1212	NVGR	17 40 24.64	+19 41 52.9 c	5.0	wat? d?
1213	NVGR	17 40 30.26	-14 15 30.9 v	4.0	cp? rc with d?
1214	NVGR	17 41 32.50	-05 24 33.5 c	10.0	t w?
1215	NVGR	17 41 42.67	-12 58 29.9 v	6.0	t (only two components fit)
1216	NVGR	17 42 06.85	+18 27 29.9 c	5.5	t rc 'long' core
1217		17 43 18.67	+71 13 11.6 c	4.5	t sz? dl? (Lara et al.(2001): J1743+712)
1218	NVGR	17 43 57.96	+19 35 08.6 c	4.5	rc? sz?
1219	NVGR	17 44 56.50	+55 42 16.6 c	9.0	sz rc?
1220		17 45 37.72	+71 15 41.2 v	6.0	d? asym x? (Lara et al.(2001): J1745+712)
1221	NVGR	17 46 31.28	+23 43 26.0 v	5.0	d? wat?
1222	NVGR	17 47 55.65	-11 01 07.6 c	5.0	sz? rc?
1223	NVGR	17 48 39.25	-23 35 17.6 c	8.0	t
1224	NVGR	17 49 11.39	-15 21 51.1 v	5.0	d
1225	NVGR	17 49 23.62	-07 07 40.9 v	6.0	cp point with teardrop
1226	NVGR	17 49 33.58	+18 24 03.5 c	4.5	cj? rc? d?
1227		17 54 28.73	+62 35 16.1 v	10.0	mg irr interacting systems? (Lara et al.(2001): J1754+626)
1228	NVGR	17 56 45.96	+45 49 33.8 v	4.0	d
1229	NVGR	17 57 44.35	-08 25 45.3 c	5.0	d/t
1230	NVGR	17 57 58.11	-08 45 27.0 v	4.5	w? d?
1231	NVGR	17 58 14.15	+15 44 10.6 v	10.0	d? t? both 'lobes' fuzzy
1232	NVGR	17 58 30.12	+65 53 58.1 v	4.5	d
1233	NVGR	17 59 33.04	+06 16 15.6 v	4.5	d
1234	NVGR	17 59 48.56	-34 49 41.3 v	5.5	d
1235		18 00 39.34	+71 44 56.2 v	6.0	d (Lara et al.(2001): J1800+717)
1236	NVGR	18 02 00.65	+12 04 34.5 v	5.5	nat?
1237	NVGR	18 04 49.97	+10 42 11.6 v	5.5	d
1238	NVGR	18 07 26.66	-09 05 20.3 v	7.0	wat/nat
1239	NVGR	18 08 23.16	+08 41 02.6 c	13.0	rc? lobe?
1240	NVGR	18 09 06.48	-15 01 13.6 v	6.5	w?
1241	NVGR	18 10 37.74	+37 36 01.8 v	5.5	d? cj?
1242	NVGR	18 11 26.69	+10 08 10.6 c	6.5	sz cp?
1243	NVGR	18 12 03.99	+21 24 02.1 v	10.0	w? irr or 5' wat with 6' wat
1244	NVGR	18 13 58.35	+35 03 50.7 c	4.0	d t
1245	NVGR	18 16 53.36	+03 39 52.5 v	5.5	d? sz?
1246	NVGR	18 20 06.29	+08 56 11.0 v	8.0	wat? sz?
1247	NVGR	18 20 08.08	+22 51 25.1 c	5.0	d t?
1248	NVGR	18 21 05.01	-05 02 21.2 v	5.5	d? points, many pairs in neighborhood
1249	NVGR	18 21 28.08	+11 38 28.9 v	5.0	d? points
1250	NVGR	18 21 54.32	-02 38 42.6 v	9.0	sz wat? suggestion of core in image, but none fit
1251	NVGR	18 22 41.20	+12 15 50.8 v	7.5	t
1252	NVGR	18 22 53.93	+36 09 03.4 v	4.0	d
1253	NVGR	18 24 46.07	+36 04 29.6 c	6.0	t dd? asym?
1254	NVGR	18 28 33.50	+08 19 39.3 v	6.0	sz
1255	NVGR	18 28 43.52	-30 44 14.6 c	5.0	t
1256	NVGR	18 29 00.38	+32 21 16.5 v	8.0	wat? sz? core id amb
1257	NVGR	18 29 18.65	-10 50 49.9 v	7.0	shell? int twisted (see a contour map) irr disturbed area
1258	NVGR	18 29 37.53	+22 30 11.2 v	5.5	sz?
1259	NVGR	18 30 22.63	+04 23 03.0 v	4.5	rc? with low level dl
1260	NVGR	18 31 21.45	-22 57 15.0 c	4.5	d t core not apparent
1261		18 32 59.35	-10 12 58.9 v	16.0	wat? artifact? disturbed area (Green(2014, 2016): SNR Kes 69 (G21.8-0.6))
1262	NVGR	18 33 36.49	-23 17 50.4 c	6.0	cj
1263	NVGR	18 34 40.73	-23 41 53.0 c	5.0	cj
1264	NVGR	18 34 56.71	+45 02 37.0 v	4.5	d?
1265	NVGR	18 35 03.49	+32 41 33.9 v	4.5	rc nat/wat?
1266		18 35 06.55	+66 35 09.4 c	5.5	t (Lara et al.(2001): J1835+665)
1267	NVGR	18 35 22.14	-25 46 13.0 v	7.0	sz? wat?
1268	NVGR	18 36 32.15	+19 41 28.6 c	11.0	w int core id ambiguous
1269	NVGR	18 38 05.72	-12 46 16.3 v	6.0	mg unu int cp of 4.5' d with ? GRS or two?
1270	NVGR	18 38 25.85	+17 11 42.0 v	6.0	rc
1271	NVGR	18 41 07.61	-35 57 20.3 v	7.0	d? cp?
1272	NVGR	18 41 12.79	+09 29 18.5 v	7.0	d? cp? point and pointlike
1273	NVGR	18 43 30.58	-03 48 47.2 v	12.0	mg w? int disturbed area
1274		18 45 06.80	+81 49 33.1 v	5.5	d? butterfly asym (Lara et al.(2001): J1845+818)
1275	NVGR	18 45 24.68	-19 24 51.7 v	5.0	wat? d?
1276	NVGR	18 46 30.34	+49 16 58.5 v	6.0	nat? (with cp?)
1277	NVGR	18 47 13.09	-26 44 52.2 v	4.5	cp? point with arc/rc
1278		18 47 36.08	+70 43 57.3 c	5.0	t (with cp?) (Lara et al.(2001): J1847+707)
1279	NVGR	18 47 36.22	+09 24 49.5 c	5.5	rc? HII? fuzzy
1280	NVGR	18 48 42.04	+19 19 56.9 c	8.0	nat dumbell dl
1281	NVGR	18 49 52.83	-07 15 49.3 v	5.0	d

1282	NVGR	J185036.1-290517	18 50 36.17	-29 05 17.2	v	6.0	wat? sz? x?
1283			18 50 45.80	+64 30 33.0	c	7.0	w? rc? d? (Lara et al.(2001): J1850+645)
1284	NVGR	J185050.2-165554	18 50 50.23	-16 55 54.1	v	4.5	d
1285			18 52 29.98	+00 40 14.7	v	9.5	mg irr disturbance area (Green(2014, 2016): SNR Kes 79 (G33.6+0.1))
1286	NVGR	J185234.7-101347	18 52 34.78	-10 13 47.7	v	6.0	wat/nat?
1287	NVGR	J185332.2+504612	18 53 32.26	+50 46 12.3	c	8.0	wat? ring-lobe hymors?
1288			18 54 00.07	+80 02 50.3	v	7.0	d 4 components (Lara et al.(2001): J1853+800)
1289	NVGR	J185419.3+050213	18 54 19.32	+05 02 13.9	v	5.5	cp
1290			18 55 28.61	-07 16 12.6	v	7.0	sz? (Solovyov & Verkhodanov(2014): J185528.6-071613)
1291	NVGR	J185621.3-313507	18 56 21.39	-31 35 07.4	v	5.5	d (with cp?)
1292	NVGR	J185722.6-191308	18 57 22.63	-19 13 08.8	v	4.0	d
1293	NVGR	J185739.6+354317	18 57 39.63	+35 43 17.7	c	4.5	t
1294	NVGR	J185748.5+613535	18 57 48.57	+61 35 35.7	c	6.0	mg nat/wat?
1295	NVGR	J185752.2+243428	18 57 52.26	+24 34 28.8	c	5.0	t
1296	NVGR	J185832.3-155702	18 58 32.37	-15 57 02.7	c	4.0	wat/nat (with cp?) may be larger
1297	NVGR	J190241.5+145100	19 02 41.53	+14 51 00.9	v	5.5	cp point and elliptic
1298	NVGR	J190304.6+361647	19 03 04.62	+36 16 47.9	c	9.0	w
1299	NVGR	J190521.4+391735	19 05 21.42	+39 17 35.1	v	4.5	d cj?
1300	NVGR	J190927.4-242833	19 09 27.41	-24 28 33.9	v	10.0	t dd?
1301	NVGR	J191029.7+135909	19 10 29.74	+13 59 09.1	v	7.5	d wat? four components
1302	NVGR	J191149.6+780045	19 11 49.67	+78 00 45.9	v	11.5	d? cp? points fuzzy
1303	NVGR	J191315.7+470656	19 13 15.72	+47 06 56.7	c	8.0	sz? wat?
1304	NVGR	J191327.9+420139	19 13 27.96	+42 01 39.7	c	6.5	wat
1305	NVGR	J191407.4+471829	19 14 07.47	+47 18 29.7	c	9.0	wat irr int hook, core id amb
1306	NVGR	J191442.8+380038	19 14 42.83	+38 00 38.0	c	5.5	wat/nat
1307	NVGR	J191452.0+271941	19 14 52.07	+27 19 41.8	c	5.0	d t?
1308	NVGR	J191456.2-255145	19 14 56.20	-25 51 45.5	c	8.5	d? rc?
1309	NVGR	J191742.3-025654	19 17 42.32	-02 56 54.3	v	4.5	d
1310	NVGR	J191756.4-243923	19 17 56.48	-24 39 23.6	c	8.0	wat? sz?
1311	NVGR	J191805.7+554237	19 18 05.70	+55 42 37.5	v	4.5	d
1312			19 18 14.64	+74 15 29.5	v	8.0	d? (Lara et al.(2001): J1918+742)
1313	NVGR	J191820.4-204044	19 18 20.41	-20 40 44.8	v	6.0	d? t?
1314	NVGR	J191820.7-064441	19 18 20.72	-06 44 41.8	c	5.5	wat/nat? sz?
1315			19 19 20.22	+51 42 53.0	v	9.0	x? dd? appears to be at least 5 components unu (Solovyov & Verkhodanov(2014): J191918.2+514208)
1316	NVGR	J192001.9+452653	19 20 01.99	+45 26 53.5	c	19.0	wat? t? d?
1317	NVGR	J192016.5-055155	19 20 16.59	-05 51 55.4	v	4.0	rc? d?
1318	NVGR	J192048.3-201222	19 20 48.39	-20 12 22.3	c	6.5	mg t? ring lobe?
1319	NVGR	J192115.5+480416	19 21 15.59	+48 04 16.2	v	11.0	mg 11' nat/wat? or 8'wat? with 7'wat? GRS or two?
1320	NVGR	J192157.4+530638	19 21 57.47	+53 06 38.8	v	4.0	rc? cj?
1321	NVGR	J192326.4-334707	19 23 26.41	-33 47 07.2	c	5.0	d t? this core coord, central source looks like bridge not core
1322	NVGR	J192432.6+352241	19 24 32.64	+35 22 41.2	c	6.5	t x? sz? rc? dd??
1323	NVGR	J192530.3+161429	19 25 30.39	+16 14 29.9	v	20.0	shell fragments? artifact? HII? disturbance area int structures
1324	NVGR	J192604.1+224259	19 26 04.14	+22 42 59.5	v	6.0	wat? sz? cp? irr 4 components
1325	NVGR	J192638.1-323855	19 26 38.12	-32 38 55.4	v	11.5	cp two elliptic not aligned
1326	NVGR	J192655.1-391743	19 26 55.19	-39 17 43.9	c	8.0	t (with artifact?) size ambiguous
1327	NVGR	J192821.1-293137	19 28 21.17	-29 31 37.5	v	6.0	w? nat/wat? core id amb
1328	NVGR	J192826.1+033244	19 28 26.17	+03 32 44.2	c	5.0	mg wat/nat? cp?
1329	NVGR	J192933.4-224421	19 29 33.45	-22 44 21.3	c	6.0	mg t+rc? t fades
1330	NVGR	J192954.0-312615	19 29 54.07	-31 26 15.0	c	5.5	t? rc?
1331	NVGR	J192959.4-001538	19 29 59.48	-00 15 38.1	c	8.0	wat?
1332	NVGR	J193009.7-051647	19 30 09.73	-05 16 47.3	v	5.5	d
1333	NVGR	J193016.2+462752	19 30 16.28	+46 27 52.3	v	6.0	mg d? wat? cp? could be larger, GRS or two?
1334	NVGR	J193045.9-393829	19 30 45.99	-39 38 29.8	v	4.0	d
1335	NVGR	J193138.5-094823	19 31 38.59	-09 48 23.8	v	5.5	d
1336	NVGR	J193310.3-022844	19 33 10.30	-02 28 44.8	v	4.0	d
1337	NVGR	J193326.4-394034	19 33 26.42	-39 40 34.5	v	4.0	d
1338	NVGR	J193457.2+432834	19 34 57.21	+43 28 34.2	v	7.0	d
1339	NVGR	J193503.9+215647	19 35 03.92	+21 56 47.8	v	8.0	wat? arc wispy irr
1340	NVGR	J193515.0-342507	19 35 15.03	-34 25 07.2	v	5.5	d
1341	NVGR	J193601.3+470127	19 36 01.39	+47 01 27.4	v	6.5	cp nat?
1342	NVGR	J193613.2-171125	19 36 13.25	-17 11 25.1	c	5.0	t
1343	NVGR	J193925.6+074359	19 39 25.61	+07 43 59.8	v	4.0	d? dl
1344	NVGR	J194019.5-155502	19 40 19.56	-15 55 02.6	v	8.0	cp
1345	NVGR	J194142.1+503750	19 41 42.12	+50 37 50.0	c	9.5	wat/nat
1346	NVGR	J194219.2+525206	19 42 19.20	+52 52 06.1	v	5.0	mg d? cp?
1347	NVGR	J194319.7+072733	19 43 19.73	+07 27 33.7	v	5.0	d t?
1348			19 43 49.27	-35 46 46.4	c	8.0	mg wat? t? dd?? multiple choices for core, may be larger (Solovyov & Verkhodanov(2014): J194348.3-354646)
1349	NVGR	J194436.7+552843	19 44 36.71	+55 28 43.6	v	6.0	mg cp nat?
1350	NVGR	J194530.0-005054	19 45 30.08	-00 50 54.5	v	5.0	wat? d?
1351	NVGR	J194556.7+251700	19 45 56.79	+25 17 00.4	v	5.5	rc? HII? not original pair, may be larger
1352	NVGR	J194625.5+243638	19 46 25.59	+24 36 38.3	v	8.0	cp? rc? irr
1353	NVGR	J194647.9+251245	19 46 47.91	+25 12 45.0	c	5.5	cj? HII?
1354	NVGR	J194740.7+390558	19 47 40.79	+39 05 58.3	v	6.0	cp d? faint
1355	NVGR	J194808.8-243132	19 48 08.89	-24 31 32.1	c	6.0	t nat?
1356	NVGR	J194837.3-291608	19 48 37.37	-29 16 08.2	c	6.5	mg t+cp?
1357	NVGR	J194906.1-155641	19 49 06.18	-15 56 41.9	v	4.5	w? rc? sz?
1358	NVGR	J194913.3+485045	19 49 13.33	+48 50 45.3	v	4.5	d
1359			19 51 40.40	+70 37 44.2	c	7.0	d t? core id amb (Lara et al.(2001): J1951+706)

			19 52 32.3	-01 17 34.3	c	9.0	sz? w? (Solovyov & Verkhodanov(2014): J195230.9-011711)
1361	NVGR	J195232.3+400617	19 52 32.38	+40 06 17.4	v	5.5	cp? rc with rc? HII?
1362	NVGR	J195440.6+401106	19 54 40.61	+40 11 06.1	v	5.5	d? points - incidental
1363	NVGR	J195637.7-123518	19 56 37.70	-12 35 18.1	v	3.5	d with cp?
1364	NVGR	J195825.3+545835	19 58 25.35	+54 58 35.6	v	4.0	d
1365	NVGR	J195830.4-373830	19 58 30.45	-37 38 30.2	c	13.0	t sz?
1366	NVGR	J195928.0+404403	19 59 28.05	+40 44 03.5	v	4.0	d may be larger
1367	NVGR	J200101.7+333111	20 01 01.75	+33 31 11.1	v	8.0	mg complex cluster
1368	NVGR	J200140.4-383706	20 01 40.49	-38 37 06.8	c	7.5	t
1369	NVGR	J200206.1+413243	20 02 06.19	+41 32 43.5	c	4.0	rc? may be larger - incidental
1370	NVGR	J200248.1-032610	20 02 48.13	-03 26 10.9	v	5.0	wat/nat? sz?
1371	NVGR	J200650.0+475047	20 06 50.00	+47 50 47.3	c	5.5	t
1372	NVGR	J200817.3+465723	20 08 17.35	+46 57 23.6	c	4.0	t
1373	NVGR	J200951.7+031448	20 09 51.78	+03 14 48.5	v	6.0	wat?
1374	NVGR	J201015.2-301230	20 10 15.22	-30 12 30.6	v	4.0	rc butterfly
1375	NVGR	J201030.5-194851	20 10 30.58	-19 48 51.1	c	5.5	t
1376	NVGR	J201128.8+211241	20 11 28.88	+21 12 41.5	v	5.5	d 4 components
1377	NVGR	J201329.3+012904	20 13 29.38	+01 29 04.0	c	6.0	t sz?
1378	NVGR	J201554.1+592247	20 15 54.13	+59 22 47.0	v	5.0	mg cp? elliptic may be larger
1379			20 16 13.60	+60 54 13.7	c	4.5	wat (with cp?) core id amb (Lara et al.(2001): J2016+608)
1380	NVGR	J201615.8-304124	20 16 15.86	-30 41 24.9	v	6.0	t dumbell 5 components
1381	NVGR	J201706.5-310306	20 17 06.50	-31 03 06.1	v	4.0	mg d? at least 4 components, this coord for W pair (not original pair), may be larger GRS or two?
1382	NVGR	J201719.6-310306	20 17 19.69	-31 03 06.1	v	3.5	d?
1383	NVGR	J201720.8-214023	20 17 20.89	-21 40 23.4	c	5.5	t
1384	NVGR	J201812.6-043022	20 18 12.61	-04 30 22.5	v	5.5	cp? nat/wat? fairly symmetrical
1385	NVGR	J201938.8+141653	20 19 38.83	+14 16 53.9	v	5.5	sz? dd? cp? int
1386	NVGR	J202003.3+040828	20 20 03.34	+04 08 28.8	v	5.0	d
1387	NVGR	J202147.5+161606	20 21 47.51	+16 16 06.7	v	5.0	d
1388	NVGR	J202235.2+024054	20 22 35.22	+02 40 54.4	v	4.5	d?
1389	NVGR	J202320.4-181107	20 23 20.40	-18 11 07.1	v	4.0	d? rc?
1390			20 23 40.09	+17 03 56.7	c	8.0	t asym (Solovyov & Verkhodanov(2014): J202339.8+170350)
1391	NVGR	J202355.8-095247	20 23 55.80	-09 52 47.3	v	4.5	wat?
1392	NVGR	J202442.4+025603	20 24 42.46	+02 56 03.0	v	5.5	d? cp? irr
1393	NVGR	J202805.1+563113	20 28 05.19	+56 31 13.7	v	5.5	d
1394	NVGR	J203059.1+101323	20 30 59.10	+10 13 23.5	v	5.0	d
1395	NVGR	J203121.3+082220	20 31 21.34	+08 22 20.5	v	6.5	mg d? non-aligned elliptics
1396	NVGR	J203242.4+591747	20 32 42.47	+59 17 47.6	v	5.0	wat? non-aligned elliptics
1397	NVGR	J203506.6+005246	20 35 06.66	+00 52 46.8	v	6.0	cp elliptic and point
1398	NVGR	J203616.7+063159	20 36 16.76	+06 31 59.4	c	5.5	w? butterfly d? t?
1399	NVGR	J203901.5-300555	20 39 01.58	-30 05 55.4	v	4.5	d
1400	NVGR	J203931.2-351219	20 39 31.21	-35 12 19.5	c	6.0	d? t? core id amb
1401	NVGR	J203942.3+241920	20 39 42.33	+24 19 20.0	v	7.0	d? cp? one component with fishtail
1402	NVGR	J203943.1+321145	20 39 43.17	+32 11 45.9	c	5.0	rc (with cp?) may be larger GRS or two?
1403	NVGR	J204018.7-065915	20 40 18.75	-06 59 15.4	c	5.0	d three components
1404	NVGR	J204113.7+220425	20 41 13.79	+22 04 25.9	v	4.0	d
1405	NVGR	J204153.3+453023	20 41 53.35	+45 30 23.0	v	7.0	cj? rc? cp? wat? arc
1406			20 42 37.93	+75 08 02.1	c	12.5	t beautiful (Lara et al.(2001): J2042+751)
1407	NVGR	J204251.6-350921	20 42 51.60	-35 09 21.8	v	7.0	wat
1408	NVGR	J204345.3-263256	20 43 45.36	-26 32 56.1	c	7.5	sz t
1409	NVGR	J204403.0+071220	20 44 03.01	+07 12 20.5	v	5.5	mg cj? wat?
1410			20 48 05.26	+30 42 47.8	c	15.0	wat? artifact? region of wrinkles (Green(2014, 2016): Cygnus Loop SNR (G74.0-8.5))
1411	NVGR	J204811.6-284703	20 48 11.64	-28 47 03.3	v	5.0	mg points and elliptic
1412	NVGR	J204900.7+193007	20 49 00.74	+19 30 07.3	c	5.0	t wat?
1413	NVGR	J204941.0-302946	20 49 41.09	-30 29 46.9	v	5.0	d? points, int configuration
1414			20 50 51.15	+31 27 28.0	v	105.0	sz? wrinkle? (Green(2014, 2016): Cygnus Loop SNR (G74.0-8.5))
1415	NVGR	J205508.5-011226	20 55 08.55	-01 12 26.9	v	5.5	cp d?
1416	NVGR	J205631.6-190612	20 56 31.68	-19 06 12.3	v	6.0	cp?
1417			20 57 19.49	+31 12 00.1	c	8.0	rc? artifact? (Green(2014, 2016): Cygnus Loop SNR (G74.0-8.5))
1418			20 59 08.54	+62 47 37.7	v	6.0	d (Lara et al.(2001): J2059+627)
1419	NVGR	J205949.4-250626	20 59 49.47	-25 06 26.8	v	4.0	wat?
1420			21 01 38.31	-28 01 45.0	v	14.0	w twisted int (Solovyov & Verkhodanov(2014): J210138.4-280158)
1421	NVGR	J210145.4-131829	21 01 45.43	-13 18 29.1	v	9.0	sz?
1422	NVGR	J210239.7-205810	21 02 39.72	-20 58 10.9	c	11.0	t
1423	NVGR	J210245.4+294130	21 02 45.48	+29 41 30.0	v	5.5	d? point and near point
1424	NVGR	J210300.2-305840	21 03 00.26	-30 58 40.8	v	5.0	d (with cp?)
1425			21 03 10.47	+64 56 31.6	v	6.0	t? sz? core id amb (Lara et al.(2001): J2103+649)
1426	NVGR	J210428.3+323549	21 04 28.38	+32 35 49.1	v	6.0	d? cp?
1427	NVGR	J210611.1+123957	21 06 11.16	+12 39 57.3	v	5.0	d
1428	NVGR	J210632.3-170324	21 06 32.38	-17 03 24.6	v	5.0	d
1429	NVGR	J210720.2-252844	21 07 20.27	-25 28 44.2	v	10.0	wat with adjacent nat/wat
1430	NVGR	J210724.1-252545	21 07 24.15	-25 25 45.8	v	5.5	nat/wat with adjacent wat
1431	NVGR	J210911.6-125243	21 09 11.69	-12 52 43.5	v	6.0	mg d cp? dd?? core amb
1432	NVGR	J211005.1+081022	21 10 05.11	+08 10 22.4	v	4.0	d? point and elliptic
1433			21 11 31.09	+63 00 07.6	c	8.0	t? nat? with extended emission irr (Lara et al.(2001): J2111+630)
1434	NVGR	J211155.9-132519	21 11 55.92	-13 25 19.3	v	5.5	mg points+elliptic
1435	NVGR	J211306.6-394059	21 13 06.61	-39 40 59.9	v	6.5	cp point with elliptic
1436	NVGR	J211314.2+023022	21 13 14.27	+02 30 22.5	v	8.0	mg irr int tails or cp?
1437	NVGR	J211325.7+562913	21 13 25.76	+56 29 13.3	v	4.5	d?

			21 14 00.33	+82 04 52.0 c	6.5	t sz? (Lara et al.(2001): J2114+820)
1439	NVGR	J211514.9+225344	21 15 14.91	+22 53 44.1 c	4.5	t sz?
1440	NVGR	J211727.0+512130	21 17 27.00	+51 21 30.0 v	5.0	mg irr d+cp? may be larger system
1441	NVGR	J211735.0+512150	21 17 35.00	+51 21 50.0 v	10.0	mg t?
1442	NVGR	J211821.5-244952	21 18 21.52	-24 49 52.4 v	21.0	cp
1443	NVGR	J211832.9+262643	21 18 32.90	+26 26 43.7 c	8.5	rc
1444	NVGR	J211858.6+003340	21 18 58.68	+00 33 40.5 c	6.5	mg cp? d? t?, core if t
1445	NVGR	J211941.9-232056	21 19 41.94	-23 20 56.0 v	6.0	d? cp?
1446	NVGR	J211944.3-251539	21 19 44.38	-25 15 39.0 v	4.5	d
1447	NVGR	J212155.4+293602	21 21 55.43	+29 36 02.8 v	5.0	d
1448	NVGR	J212311.2-210500	21 23 11.22	-21 05 00.6 c	7.0	t wat?
1449	NVGR	J212439.2+505823	21 24 39.25	+50 58 23.8 c	10.5	w? t
1450	NVGR	J212544.3+005550	21 25 44.35	+00 55 50.3 v	4.0	rc (with cp?)
1451	NVGR	J212619.8+132116	21 26 19.89	+13 21 16.0 v	4.0	d? points
1452			21 28 07.45	+60 20 38.3 v	7.0	w? x? 4 components (Lara et al.(2001): J2128+603)
1453	NVGR	J212840.5-393342	21 28 40.52	-39 33 42.5 v	5.0	d? with dl sz?
1454	NVGR	J212850.9+011525	21 28 50.91	+01 15 25.1 c	4.5	mg tri int
1455	NVGR	J212859.1+510446	21 28 59.16	+51 04 46.5 v	6.0	d
1456	NVGR	J212937.4-211145	21 29 37.48	-21 11 45.2 c	15.0	rc sz
1457	NVGR	J213006.9+554806	21 30 06.99	+55 48 06.7 v	4.0	shell? rc? artifact? int grayscale
1458	NVGR	J213039.2+073542	21 30 39.26	+07 35 42.4 v	4.5	d/t
1459			21 32 08.23	+34 21 02.9 c	8.0	t (with cp?) (Solovyov & Verkhodanov(2014): J213208.6+342057)
1460	NVGR	J213446.8+144333	21 34 46.81	+14 43 33.5 v	4.5	d
1461	NVGR	J213526.0+534903	21 35 26.02	+53 49 03.8 v	10.0	mg d/t? shell?
1462	NVGR	J213535.1-182219	21 35 35.13	-18 22 19.5 c	5.0	t
1463	NVGR	J213745.3-143257	21 37 45.39	-14 32 57.1 v	4.5	d
1464	NVGR	J213959.6+092905	21 39 59.60	+09 29 05.1 v	6.5	cp point and elliptic
1465	NVGR	J214002.0-142122	21 40 02.08	-14 21 22.2 v	4.5	d/t
1466	NVGR	J214045.5+370419	21 40 45.59	+37 04 19.9 c	4.5	t (a component of original pair)
1467	NVGR	J214048.1-274431	21 40 48.15	-27 44 31.9 v	18.0	cp
1468	NVGR	J214103.5-164137	21 41 03.59	-16 41 37.4 c	9.0	wat
1469	NVGR	J214209.4+231104	21 42 09.41	+23 11 04.3 v	4.0	cp point with teardrop
1470	NVGR	J214231.0+401758	21 42 31.00	+40 17 58.9 c	4.5	t with dl
1471	NVGR	J214515.9-195913	21 45 15.96	-19 59 13.0 v	11.0	mg w wat?
1472			21 45 30.83	+81 54 55.5 c	21.0	t x?? (11" d/t?, diagonal with same core?) int (Lara et al.(2001): J2145+819)
1473	NVGR	J214536.6-042642	21 45 36.66	-04 26 42.2 v	5.0	d/t
1474	NVGR	J214623.3+333213	21 46 23.33	+33 32 13.6 v	7.0	nat/wat sz?
1475	NVGR	J214640.8-370153	21 46 40.82	-37 01 53.7 v	4.0	d t? may be larger
1476	NVGR	J214732.3-274255	21 47 32.36	-27 42 55.1 v	4.0	d cj? cp?
1477	NVGR	J214920.3+194037	21 49 20.38	+19 40 37.3 c	5.0	d t? core fit but not apparent
1478	NVGR	J214936.7+205148	21 49 36.75	+20 51 48.3 v	5.0	cp
1479	NVGR	J215157.7-153745	21 51 57.71	-15 37 45.8 v	6.5	sz? twisted int
1480	NVGR	J215305.0-071107	21 53 05.08	-07 11 07.6 c	9.5	nat/wat?
1481	NVGR	J215337.7+474854	21 53 37.79	+47 48 54.1 v	5.5	d?
1482	NVGR	J215356.6+125211	21 53 56.65	+12 52 11.5 v	5.5	d?
1483	NVGR	J215431.8+030529	21 54 31.80	+03 05 29.1 v	5.0	d? cp?
1484	NVGR	J215657.7+325908	21 56 57.76	+32 59 08.4 c	6.0	wat? core id ambiguous
1485	NVGR	J215659.6+523356	21 56 59.61	+52 33 56.3 v	5.0	d? wat? wat? points
1486			21 57 03.01	+66 26 10.8 c	5.0	d? cj? int tail, core assumes cj (Lara et al.(2001): J2157+664)
1487	NVGR	J215832.0+595045	21 58 32.06	+59 50 45.5 v	5.0	rc? d? wat?
1488	NVGR	J215935.5+565621	21 59 35.50	+56 56 21.9 v	3.0	rc? HII? arc
1489	NVGR	J220136.2-192344	22 01 36.21	-19 23 44.9 c	8.0	t? nat?
1490	NVGR	J220319.4+541238	22 03 19.42	+54 12 38.2 v	5.0	d asym
1491			22 04 13.70	+78 22 47.3 v	5.5	t (Lara et al.(2001): J2204+783)
1492	NVGR	J220515.8-303305	22 05 15.85	-30 33 05.2 v	6.0	d sz?
1493	NVGR	J220544.8+780819	22 05 44.85	+78 08 19.6 v	19.5	d? points, both fuzzy, also at least two other similar nearby
1494	NVGR	J220612.6+153209	22 06 12.64	+15 32 09.4 v	4.0	d cj?
1495			22 09 28.08	+72 46 13.8 c	5.5	t (with cp?) (Lara et al.(2001): J2209+727)
1496	NVGR	J221114.8+385546	22 11 14.89	+38 55 46.7 c	12.0	w? sz? irr
1497	NVGR	J221242.4+451941	22 12 42.45	+45 19 41.8 v	6.5	cp d?
1498	NVGR	J221402.4-290713	22 14 02.45	-29 07 13.3 v	5.5	d dd? 4 components
1499	NVGR	J221443.7+135019	22 14 43.79	+13 50 19.0 v	10.0	wat/nat? cp? int
1500	NVGR	J221536.9+290230	22 15 36.98	+29 02 30.3 c	5.0	t
1501	NVGR	J221614.6-163144	22 16 14.65	-16 31 44.5 v	5.0	t? d? sz?
1502	NVGR	J221743.3-242030	22 17 43.39	-24 20 30.1 v	4.0	mg d? points
1503	NVGR	J221815.7+193138	22 18 15.72	+19 31 38.0 v	6.0	t
1504	NVGR	J221837.8-082307	22 18 37.86	-08 23 07.1 v	8.0	d? mg
1505	NVGR	J222001.2+573108	22 20 01.20	+57 31 08.3 v	7.0	rc? hazy
1506	NVGR	J222002.5-040707	22 20 02.50	-04 07 07.7 v	6.0	sz rc?
1507	NVGR	J222150.8+421521	22 21 50.87	+42 15 21.6 v	5.0	mg t? d? cp? wat?
1508	NVGR	J222349.2-020556	22 23 49.27	-02 05 56.1 c	12.0	w
1509	NVGR	J222511.5+042112	22 25 11.58	+04 21 12.5 c	8.0	mg nat/wat? irr (with cp?), core id amb
1510	NVGR	J222521.9-161638	22 25 21.95	-16 16 38.4 c	12.5	wat? w? irr cp? core id amb
1511	NVGR	J222525.0+581200	22 25 25.00	+58 12 00.0 v	6.0	rc? HII? SNR? lobe? very mottled/wispy
1512	NVGR	J222805.0+493435	22 28 05.04	+49 34 35.2 v	5.0	d
1513	NVGR	J222845.6+395741	22 28 45.60	+39 57 41.7 v	4.5	rc? sz? dl? d asym
1514	NVGR	J222932.4-292423	22 29 32.43	-29 24 23.5 v	4.5	d/t
1515	NVGR	J223036.6+135854	22 30 36.69	+13 58 54.8 v	5.0	w? cp?

1516	NVGR	J223040.3-394249	22 30 40.36	-39 42 49.5	c	5.5	t? somewhat asymmetric, d with s?
1517	NVGR	J223111.1+010114	22 31 11.16	+01 01 14.7	v	9.0	mg cp? w?
1518			22 31 19.78	+39 22 47.6	c	52.0	w? jets? dd? int, may be larger, core id amb (Soloviyov & Verkhodanov(2014): J223115.9+392501)
1519	NVGR	J223128.8-082907	22 31 28.83	-08 29 07.6	c	6.5	w? wat/nat? - incidental
1520	NVGR	J223129.6-203549	22 31 29.60	-20 35 49.0	v	4.5	cp
1521	NVGR	J223143.7-082433	22 31 43.76	-08 24 33.5	c	7.5	t dumbell with core
1522	NVGR	J223241.5+213452	22 32 41.58	+21 34 52.6	v	5.0	cp
1523	NVGR	J223251.6-053015	22 32 51.62	-05 30 15.3	v	4.5	cj? d? asymmetric size/peaks
1524			22 35 35.97	+37 35 22.9	v	8.0	d (Soloviyov & Verkhodanov(2014): J223536.0+373523)
1525	NVGR	J223704.6+342427	22 37 04.66	+34 24 27.9	c	7.5	rc int two pairs of 'satellites' plus elliptic
1526	NVGR	J223803.0+410821	22 38 03.03	+41 08 21.5	v	7.0	wat? w? dd?
1527	NVGR	J223808.4+595208	22 38 08.42	+59 52 08.2	v	7.0	sz
1528	NVGR	J223835.9-070511	22 38 35.97	-07 05 11.3	v	5.5	d
1529	NVGR	J223911.3-172047	22 39 11.30	-17 20 47.7	v	4.5	wat/nat?
1530	NVGR	J224140.1+473709	22 41 40.17	+47 37 09.7	v	6.5	d
1531			22 42 35.20	+62 12 24.8	v	6.0	sz? d? 4 components (Lara et al.(2001): J2242+622)
1532	NVGR	J224250.1+332153	22 42 50.15	+33 21 53.7	c	4.0	cj? d? asym
1533	NVGR	J224319.4+440049	22 43 19.44	+44 00 49.6	v	5.0	d
1534			22 45 49.29	+39 41 22.1	c	6.5	t rc? (Soloviyov & Verkhodanov(2014): J224548.5+394113)
1535	NVGR	J224620.5+314211	22 46 20.59	+31 42 11.3	v	5.0	d sz?
1536	NVGR	J224644.6-131947	22 46 44.64	-13 19 47.6	v	4.0	cp
1537	NVGR	J224710.4+441955	22 47 10.44	+44 19 55.1	v	5.5	d rc?
1538			22 47 27.83	+63 21 01.3	v	5.0	wat? dl (Lara et al.(2001): J2247+633)
1539	NVGR	J224817.4+390415	22 48 17.41	+39 04 15.5	c	6.0	mg wat? w? irr core id amb
1540	NVGR	J224943.3+113330	22 49 43.34	+11 33 30.3	c	10.5	rc? with ?, irr, may be larger, fades
1541	NVGR	J225102.7+061931	22 51 02.79	+06 19 31.0	v	5.0	d
1542	NVGR	J225123.3+381250	22 51 23.36	+38 12 50.0	v	4.5	rc? ring? nat? int
1543	NVGR	J225123.6+160817	22 51 23.61	+16 08 17.4	v	5.5	d/t
1544	NVGR	J225309.6+084956	22 53 09.67	+08 49 56.1	c	4.0	cj nat? fishtail
1545			22 53 20.99	+16 20 21.5	c	10.5	t (Soloviyov & Verkhodanov(2014): J225321.7+162014)
1546	NVGR	J225336.2-345529	22 53 36.22	-34 55 29.6	c	7.0	t
1547	NVGR	J225356.2-070638	22 53 56.20	-07 06 38.9	c	9.0	sz? w? cp??
1548			22 55 29.53	+64 30 14.7	c	5.5	d t? core fit but not apparent (Lara et al.(2001): J2255+645)
1549	NVGR	J225600.3+212725	22 56 00.36	+21 27 25.9	c	5.0	nat/wat 3 component
1550	NVGR	J225631.8+412954	22 56 31.80	+41 29 54.8	v	7.0	cp
1551	NVGR	J225933.6+082037	22 59 33.60	+08 20 37.3	v	7.0	sz? irr sz? cp?
1552	NVGR	J230022.1+092150	23 00 22.12	+09 21 50.1	c	5.0	wat? sz? sz?
1553	NVGR	J230028.8+313759	23 00 28.88	+31 37 59.4	v	5.0	d? points
1554	NVGR	J230034.1+505017	23 00 34.14	+50 50 17.9	c	5.0	t
1555	NVGR	J230057.5+282517	23 00 57.55	+28 25 17.4	v	4.5	cp?
1556	NVGR	J230157.7-020801	23 01 57.75	-02 08 01.6	v	7.5	cp d?
1557	NVGR	J230235.2+311805	23 02 35.26	+31 18 05.3	v	4.5	sz
1558	NVGR	J230302.9-184118	23 03 02.97	-18 41 18.9	c	5.5	rc sz?
1559	NVGR	J230317.7+014325	23 03 17.72	+01 43 25.9	v	4.5	rc? sz?
1560	NVGR	J230356.6-253223	23 03 56.60	-25 32 23.2	v	7.0	d? wat/nat? x? cp?
1561	NVGR	J230511.7+283428	23 05 11.71	+28 34 28.8	c	4.0	sz? rc?
1562	NVGR	J230642.1-282610	23 06 42.12	-28 26 10.4	c	5.5	d
1563			23 08 01.71	+64 02 07.1	c	6.0	t (Lara et al.(2001): J2307+640)
1564	NVGR	J230821.7-061322	23 08 21.71	-06 13 22.4	v	5.5	d? similar shapes
1565	NVGR	J230822.7+501901	23 08 22.75	+50 19 01.8	v	4.0	d (with cp west end?)
1566	NVGR	J230828.5+325152	23 08 28.51	+32 51 52.3	v	5.0	cp
1567	NVGR	J230848.9+532244	23 08 48.96	+53 22 44.5	v	6.5	mg cp? sz?
1568	NVGR	J230945.1-195705	23 09 45.19	-19 57 05.4	v	7.0	wat? w?
1569	NVGR	J231046.3-210828	23 10 46.36	-21 08 28.8	v	3.5	wat/nat? with cp? may be larger
1570	NVGR	J231136.1-285921	23 11 36.16	-28 59 21.8	v	6.0	d cp?
1571	NVGR	J231333.5+101905	23 13 33.52	+10 19 05.8	v	5.5	mg d? cp?
1572	NVGR	J231412.5+192324	23 14 12.54	+19 23 24.1	v	8.5	w? possible core J231411.88+192411.3
1573	NVGR	J231524.0-371701	23 15 24.06	-37 17 01.9	v	7.5	d? t? cp?
1574			23 16 00.60	-28 23 52.7	c	10.0	t (Soloviyov & Verkhodanov(2014): J231600.3-282359)
1575	NVGR	J231622.4+224712	23 16 22.42	+22 47 12.0	c	8.5	d 5 components core id amb
1576	NVGR	J231627.2+205559	23 16 27.29	+20 55 59.6	v	4.5	d? points
1577	NVGR	J231628.5+205844	23 16 28.59	+20 58 44.3	v	5.0	mg d? points
1578	NVGR	J231631.2+210100	23 16 31.20	+21 01 00.0	v	3.5	d? points
1579	NVGR	J231757.0-011741	23 17 57.04	-01 17 41.7	c	5.5	t
1580	NVGR	J231956.4-272837	23 19 56.46	-27 28 37.7	v	6.5	wat?
1581	NVGR	J232002.6-151650	23 20 02.63	-15 16 50.4	v	4.0	cp point and elliptic
1582	NVGR	J232013.9-132104	23 20 13.93	-13 21 04.4	v	5.5	d
1583	NVGR	J232025.5+163213	23 20 25.55	+16 32 13.3	v	5.0	d cp? elliptic
1584	NVGR	J232042.2+081256	23 20 42.21	+08 12 56.3	c	14.0	wat/nat core id amb
1585	NVGR	J232320.4+203525	23 23 20.45	+20 35 25.9	c	6.0	wat
1586	NVGR	J232344.2+145753	23 23 44.27	+14 57 53.3	c	7.0	w? nat/wat? dd?
1587	NVGR	J232512.7-323653	23 25 12.74	-32 36 53.0	v	4.5	d?
1588	NVGR	J232710.6-004158	23 27 10.68	-00 41 58.4	c	4.5	t
1589	NVGR	J232715.4-050350	23 27 15.49	-05 03 50.8	v	6.5	d? cp? hymors?
1590	NVGR	J232823.3+214557	23 28 23.33	+21 45 57.6	c	5.5	rc? t? fades core
1591	NVGR	J232944.4+431636	23 29 44.42	+43 16 36.1	v	4.0	nat
1592	NVGR	J233205.7+235107	23 32 05.76	+23 51 07.9	c	5.5	nat
1593	NVGR	J233318.7-002703	23 33 18.70	-00 27 03.2	c	5.0	t

1594	NVGR	J233355.2-234340	23 33 55.24	-23 43 40.0	c	22.0	t
1595	NVGR	J233402.3-395149	23 34 02.37	-39 51 49.8	v	5.5	d
1596	NVGR	J233513.3+174134	23 35 13.32	+17 41 34.9	v	6.0	d
1597	NVGR	J233717.7+163245	23 37 17.70	+16 32 45.1	v	5.5	d? points
1598	NVGR	J233832.4+270136	23 38 32.42	+27 01 36.2	c	11.5	wat/nat nice, core id amb
1599	NVGR	J234000.6+270800	23 40 00.65	+27 08 00.7	c	17.0	wat? cp?
1600			23 40 57.69	+62 10 38.4	v	5.0	sz d (Lara et al.(2001): J2340+621)
1601	NVGR	J234058.3-283617	23 40 58.36	-28 36 17.3	c	5.0	d
1602	NVGR	J234137.0+082855	23 41 37.09	+08 28 55.8	v	8.0	d sz?
1603	NVGR	J234622.6+361728	23 46 22.63	+36 17 28.7	c	5.0	d core fit, but not apparent
1604	NVGR	J234839.4+230218	23 48 39.49	+23 02 18.2	v	4.5	cp
1605	NVGR	J234847.5+160000	23 48 47.53	+16 00 00.3	v	4.5	d
1606	NVGR	J235305.3-172639	23 53 05.39	-17 26 39.8	v	4.0	d
1607	NVGR	J235346.6-400134	23 53 46.67	-40 01 34.2	c	9.5	wat t
1608	NVGR	J235433.9-013958	23 54 33.95	-01 39 58.6	v	4.0	t d
1609	NVGR	J235435.3+535408	23 54 35.38	+53 54 08.5	v	4.0	d? points
1610	NVGR	J235506.1+125250	23 55 06.11	+12 52 50.6	v	6.0	d t? sz? int configuration of six sources
1611	NVGR	J235525.2+262401	23 55 25.25	+26 24 01.7	v	5.5	d 4 components
1612			23 57 27.74	+47 52 28.2	v	10.0	w? sz? (Solovyov & Verkhodanov(2014): J235721.0+475238)
1613	NVGR	J235730.4+472624	23 57 30.42	+47 26 24.9	v	5.5	d
1614	NVGR	J235910.9+170613	23 59 10.94	+17 06 13.6	c	10.0	w? x? sz?
1615	NVGR	J235926.6-242530	23 59 26.62	-24 25 30.8	c	6.5	rc sz?
1616	NVGR	J235959.3+004200	23 59 59.35	+00 42 00.6	v	7.0	t