

TOWARD A CONJECTURE OF TAN AND TU ON FIBERED GENERAL TYPE SURFACES

A. HUITRADO-MORA, M. CASTAÑEDA-SALAZAR AND A. G. ZAMORA

ABSTRACT. Given a semistable non-isotrivial fibered surface $f : X \rightarrow \mathbb{P}^1$ it was conjectured by Tan and Tu that if X is of general type, then f admits at least 7 singular fibers. In this paper we prove this conjecture in several particular cases, i.e. assuming f is obtained from blowing-up the base locus of a transversal pencil on an exceptional minimal surface S or assuming that f is obtained as the blow-up of the base locus of a transversal and adjoint pencil on a minimal surface.

1. NOTATION

We work on the complex field number \mathbb{C} . All the considered varieties will be assumed irreducible and projective. Through the paper we shall use the following notation:

- . X will be a general type surface and S its minimal model. $\pi : X \rightarrow S$ will be the associated chain of blowing-downs.
- . $f : X \rightarrow \mathbb{P}^1$ will be a semi-stable, non-isotrivial fibration, and F the general fibre. We set $g =$ the genus of F . By C we will denote the image of F under π and

$$\Lambda : S \dashrightarrow \mathbb{P}^1,$$

the pencil induced by f . We denote by s the number of singular fibers.

- . We shall say that Λ is transversal if its general member $C \in \Lambda$ is non-singular and intersects transversally any other general member $C' \in \Lambda$.
- . We'll freely use the standard notation in surfaces' theory. In particular $q = h^1(X, \mathcal{O}_X)$ will be the irregularity of X and $p_g = h^0(X, K_X)$ its geometric genus. By $e(X)$ will be denote the topological Euler characteristic.
- . Given divisors D_1 and D_2 in an algebraic surface we denote as usual $D_1 \equiv D_2$ for the numeric equivalence and $D_1 \sim D_2$ for the linear one. Most of the time we will be working on regular surfaces and in this case we indistinctly use both symbols.
- . The number m will be:

$$m := K_S^2 - K_X^2 = e(X) - e(S).$$

Date: 03/25/16.

2000 Mathematics Subject Classification. 14D06, 14J25.

Key words and phrases. Algebraic Surfaces, Fibrations, Singular Fibres.

This paper was partially supported by CONACyT Grant 257079. First and second authors were supported by Conacyt Doctoral Scholarships. Third author was partially supported by Conacyt Grant 265621 for Academic Sabbatical Year.

Note that in general $m \leq C^2$. Adjunction Formula gives the important inequality $C.K_S + m \leq 2(g-1)$ with equality holding if Λ is transversal.

2. INTRODUCTION

Let $f : X \rightarrow \mathbb{P}^1$ be a non-isotrivial semistable fibered surface. It is a classical result that such a fibration admits a certain number of singular fibers (in contrast to the case when the base of the fibration is not rational or elliptic [1]). In the seminal paper [2] it was proved that this number s must be at least 4. Subsequently the bound have been sharpened to $s \geq 5$ if $g \geq 2$ and $s \geq 6$ if the surface X is not birationally ruled ([11] [12], [13]). It was conjectured by Tan and Tu, in a preprint previous to [13] that this bound must raise to 7 if X is of general type (Tan-Tu conjecture for what follows). They also proved the conjecture for genus $2 \leq g \leq 4$ and characterized fibrations of genus 5 with $s = 6$ on a general type surface as those obtained from blowing up the base locus of a transversal pencil on a Horikawa surface. Using this characterization the proof of the conjecture for $g = 5$ was completed in [15].

Roughly speaking the proof of these bounds are based, in case $g \geq 2$ on the canonical class inequality:

$$K_f^2 < 2(g-1)(2g_B - 2 + s),$$

for any non iso-trivial semi-stable fibration $f : X \rightarrow B$ of genus $g \geq 2$. Here K_f is the relatively canonical divisor, $K_f := K_X - f^*K_B$, which turns out to be $K_X(-2F)$ if $B = \mathbb{P}^1$. The bounds for s are obtained from the positivity properties of K_f and $K_f(-F)$.

Unfortunately, this approach is useless for proving Tan-Tu conjecture, since in this case the only relevant information the inequality provides is that $K_X^2 < 0$ if $s = 6$. However for most of the cases a fibered surface (of general type or not) satisfies $K_X^2 < 0$. Indeed, such a surface is obtained by blowing up the base locus of some pencil Λ on a minimal surface S .

In this paper we deal with Tan-Tu conjecture in some particular cases. We first impose on the minimal model S of X the conditions of being exceptional in the sense that either $K_S^2 = 2, p_g = 3$ or $K_S^2 = 1$ and $p_g = 2$ (see [1], VII.8) and assuming that f is obtained as the blowing-up of the base locus of a transversal pencil Λ in S . In these cases we are able to prove the conjecture using explicit descriptions of these surfaces as double coverings of rational surfaces, by means of the canonical or bi-canonical map. This is the content of Section 3, Theorems 4.1 and 4.2. In this sense there is some hope of extending the result to a wider class of surfaces.

Next, in section 5 we prove the conjecture assuming that the pencil Λ is adjoint, i.e. $C = B + K_S$ with B a big and nef divisor in S and $K_S^2 \geq 3$ and for $K_S^2 \leq 2$ assuming not only that Λ is adjoint, but also transversal (Theorem 5.3). The case $K_S^2 = p_g = 1$ is the subtler and is stated and proved in Proposition 5.2.

We list below the cases in which Tan-Tu conjecture have been proved in this article:

- If $\chi(\mathcal{O}_S) = 1$ with no extra assumption on S or Λ .
- If either $K_S^2 = 1, p_g = 2$ or $K_S^2 = 2, p_g = 3$ assuming Λ is transversal.
- If Λ is transversal and adjoint and $K_S^2 \leq 2$ or if Λ is merely adjoint and $K_S^2 \geq 3$.

3. SOME GENERAL FACTS AND RESULTS

The following inequality will be systematically used: given a semi-stable non-isotrivial fibration of genus $g \geq 2$, $f : X \rightarrow B$, for any integer $e \geq 2$:

$$\frac{1}{3}e^2(K_X^2 - 2(g-1)(6(g_B - 1) + s - s/e)) \leq e_f.$$

Original formulation involves the number of (-2) vertical curves in X , but for our purposes this version will be sufficient. The proof can be founded in [12] and is based on successive changes of the base B of the fibration. We call this Tan's inequality. In particular if $B = \mathbb{P}^1$ (our interest's case) and $s = 6$ we obtain:

$$(1) \quad \frac{1}{3}e(K_X^2 e + 12(g-1)) \leq e_f.$$

Useful forms of this inequality are collected in the following:

Lemma 3.1. *Let $f : X \rightarrow \mathbb{P}^1$ be semistable, non-isotrivial of genus $g \geq 2$. If $s = 6$, then evaluating (1) we obtain:*

i)

$$K_S^2 + C.K_S \leq 3\chi(\mathcal{O}_S) \text{ if } e = 3,$$

ii)

$$19K_S^2 + 18C.K_S \leq m + 36\chi(\mathcal{O}_S) \text{ if } e = 4,$$

iii)

$$7K_S^2 + 6C.K_S \leq m + 9\chi(\mathcal{O}_S) \text{ if } e = 5.$$

Proof. Evaluate (1) at the indicated value of e and substitute:

$$m + C.K_S \leq 2(g-1),$$

$$e(X) = 12\chi(\mathcal{O}_X) - K_X^2 \text{ (Noether's Formula),}$$

$$K_X^2 = K_S^2 - m,$$

and

$$e_f = 4(g-1) + e(X) \text{ (because } f \text{ is semistable) .}$$

□

We start by sharpening the bound for m obtained in [13], inequality (3.2) (compare with the proof of Theorem 2.1(4) in [13]).

Lemma 3.2. *Assume $f : X \rightarrow \mathbb{P}^1$ is a semistable fibration and the minimal model S of X is a general type surface, then:*

$$m \leq C^2 \leq \frac{4(g-1) + K_S^2 - \sqrt{8(g-1)K_S^2 + (K_S^2)^2}}{2}.$$

Proof. From Index Hodge Theorem applied to K_S and C we get:

$$mK_S^2 \leq C^2 K_S^2 \leq (C.K_S)^2.$$

Adjoint formula gives $C.K_S \leq 2(g-1) - m$, therefore

$$0 \leq m^2 - (4(g-1) + K_S^2)m + 4(g-1)^2.$$

Consider the right hand term of the previous inequality as a polynomial in m . Its discriminant turns out to be

$$\Delta = 8(g-1)K_S^2 + (K_S^2)^2.$$

Thus its roots are:

$$m_{\pm} = \frac{4(g-1) + K_S^2 \pm \sqrt{\Delta}}{2}.$$

It follows that either $m \leq m_-$ or $m \geq m_+$. We claim that $m \leq m_-$ is the only possible case. Indeed, if

$$\frac{4(g-1) + K_S^2 + \sqrt{\Delta}}{2} \leq m,$$

then $C.K_S \leq 2(g-1) - m \leq 0$ that is impossible. \square

As a consequence of Lemma 3.2 we obtain our first general fact concerning the number s :

Proposition 3.3. *Let S be of general type. If $s = 6$, then*

$$K_S^2 + \sqrt{8(g-1)K_S^2 + (K_S^2)^2} \leq 6\chi(\mathcal{O}_S).$$

In particular, if $g \geq 6$ and $\chi(\mathcal{O}_S) = 1$, then $s \geq 7$.

Proof. Assume $s = 6$, by Tan's inequality:

$$\frac{1}{3}e(K_X^2 e + 12(g-1)) \leq e_f,$$

for any natural number $e \geq 2$. Evaluating in $e = 3$ we obtain:

$$(2) \quad 3K_X^2 + 12(g-1) \leq e_f.$$

Since f is semistable $e(X) = -4(g-1) + e_f$ and (2) becomes:

$$(3) \quad 3K_X^2 + 12(g-1) \leq e_f = e(X) + 4(g-1).$$

By definition $m = K_S^2 - K_X^2 = e(X) - e(S)$. This, combined with Noether Formula $e(S) = 12\chi(\mathcal{O}_S) - K_S^2$ leads to:

$$K_S^2 + 2(g-1) \leq 3\chi(\mathcal{O}_S) + m.$$

The desired inequality follows after applying Lemma 3.2. \square

The importance of this Proposition is that given a family of general type surfaces with given invariants K_S^2 and $\chi(\mathcal{O}_S)$ only a finite numbers of values of g must be discharged in order to conclude that a fibered surface birational to S has at least 7 singular fibers. This principle will be illustrated in the next section.

Note that in Proposition 3.3 the hypothesis of being S of general type is essential ([2], Example 2).

4. FIBRATIONS OBTAINED FROM EXCEPTIONAL SURFACES

Surfaces satisfying either $K_S^2 = 2$ and $p_g = 3$ or $K_S^2 = 1$ and $p_g = 2$ are called exceptional because of the behavior of the tri-canonical map (Theorem VII 8.3 in [1]). In this section we study fibrations in these surfaces by means of the canonical and bi-canonical map, respectively.

Theorem 4.1. *Assume S satisfies $K_S^2 = 2$ and $p_g = 3$. Let $f : X \rightarrow \mathbb{P}^1$ be obtained as the blowing-up of a transversal pencil Λ in S . Then, $s \geq 7$.*

Proof. By Debarre's Inequality ([6]) we know that $q = 0$ and therefore $\chi(\mathcal{O}_S) = 4$. Assume $s = 6$, by Proposition 3.3 it is sufficient to consider:

$$6 \leq g \leq 31.$$

In this case the canonical map ϕ_{K_S} defines a $2 : 1$ covering:

$$\phi_{K_S} : S \rightarrow \mathbb{P}^2,$$

ramified along $R \equiv 4K_S$ ([8]). Consider the restriction $\phi := \phi_{K_S}|_C$.

First, we consider the case ϕ is a $2 : 1$ covering. Denote by $G \subset \mathbb{P}^2$ the image of C under ϕ and d for its degree. We have $C = \phi^*G$ and therefore:

$$m = C^2 = (\phi^*G)^2 = 2G^2 = 2d^2.$$

On the other hand, taking into account that $\phi^*H = K_S$, with H a hyperplane section (i.e. the divisor associated with $\mathcal{O}_{\mathbb{P}^2}(1)$) we obtain:

$$d = \frac{C.K_S}{2} \text{ and } m = \frac{(C.K_S)^2}{2}.$$

From this we get $2m = (2(g-1) - m)^2$. The possible values of g satisfying such a relation in the range $6 \leq g \leq 31$ are: $g = 7, 13, 21, 31$. The corresponding values of m and $C.K_S$ are listed below:

$g - 1$	$C.K_S$	m
6	4	8
12	6	18
20	8	32
30	10	50

If $g - 1 = 20$ or 30 we use the inequalities of Lemma 3.1 ii), in order to obtain a contradiction.

Assume $g - 1 = 6$. In this case the number of singular points in the fibers of f will be:

$$\begin{aligned} e_f &= e(X) + 4(g - 1) \\ &= -K_X^2 + 12\chi(\mathcal{O}_X) + 4(g - 1) \\ &= 6 + 48 + 24 = 78. \end{aligned}$$

Since $s = 6$ there exists at least a singular fiber F_0 of f containing $\sigma_0 = 13 = 78/6$ singular points. But then, denoting $F_0 = F_1 + \dots + F_l$ for the decomposition into irreducible components:

$$\begin{aligned} 6 = (g - 1) &= \sum_{i=1}^l (g_i - 1) + \sigma_0 \\ &\geq \sum (g_i - 1) + 13, \end{aligned}$$

with g_i standing for the geometric genus of F_i and σ_0 for the number of singular points of F_0 . We have:

$$\sum_{i=1}^l (g_i - 1) \leq -7.$$

In particular, F_0 have at least 7 irreducible rational components. Being ϕ a $2 : 1$ covering and C be applied under ϕ to a plane curve G of degree $d = C.K_S/2 = 2$ we have that the number l of irreducible components of F_0 is at most 4. In this way we get a contradiction with the assumption $s = 6$.

The case $g - 1 = 12$ follows after similar considerations, this time tanking into account that the covering ϕ sends C onto a curve G of degree 3.

Consider now the case ϕ restricted to C is $1 : 1$. In this case there exists a curve C' such that

$$C + C' = \phi^*G.$$

Moreover, since the ramifications of ϕ over C occurs exactly on the intersections of C and C' we have $C.C' = C.R = C.4K_S$. Similarly, we conclude that $C.K_S = C'.K_S$. Also, we have that:

$$2C.K_S = \phi^*G.K_S = 2d,$$

i.e. $d = C.K_S$.

It follows that $\phi^*G = \phi^*((C.K_S)H) = (C.K_S)K_S$. From

$$C + C' = (C.K_S)K_S,$$

we obtain, after intersecting with C :

$$4K_S.C = C.C' = (C.K_S)^2 - m.$$

The possibilities for such a relationship are listed below:

$(g - 1)$	$C.K_S$	m
5	5	5
9	6	12
14	7	21
20	8	32
27	9	45

Using Lemma 3.1 we obtain a contradiction in the following cases: $g - 1 = 20$ and 27 when evaluating at $e = 4$, and $g - 1 = 5, 9$ when evaluating at $e = 5$.

Finally, we must analyze the case $g - 1 = 14$. In this case,

$$\begin{aligned} e_f &= 4(g - 1) - K_X^2 + 12\chi(\mathcal{O}_X) \\ &= 123. \end{aligned}$$

Assuming $s = 6$ there must to exist a singular fiber F_0 with its number of nodes $\sigma_0 \geq 21$. Assume $F_0 = F_1 + \dots + F_l$ is its decomposition into irreducible components. Denoting by g_i the geometric genus of F_i we have:

$$14 = \sum_{i=1}^l (g_i - 1) + \sigma_0 \geq \sum_{i=1}^l (g_i - 1) + 21.$$

From this we get $l \geq 7$. Now, the image of F_0 in \mathbb{P}^2 (under the composition $\phi \circ \pi$) is a degree 7 curve G_0 . It follows that $l = 7$, $G_0 = L_1 + \dots + L_7$ is the sum of seven lines L_i , and $C_0 := \pi(F_0)$ must be the sum of seven irreducible components $C_0 = C_1 + \dots + C_7$ ($C_i = \pi(F_i)$). We have, moreover, $\phi^* L_i = C_i + C'_i \equiv K_S$, $C_i^2 = C_i'^2 = -3$ and $C_i C'_i = 4$.

A simple cohomological computation shows that $h^0(\mathcal{O}_S(C_{i_1} + C_{i_2} + C_{i_3})) = 1$ for any indexes $i_1, i_2, i_3 \in \{1, \dots, 7\}$ and $h^0(\mathcal{O}_S(C_1 + \dots + C_4)) = 2$. This, together with $(C_1 + \dots + C_4)^2 = 0$ means that $|C_1 + \dots + C_4|$ is a base point free pencil. Call $\Delta := C_1 + \dots + C_4$. We have, $C'_i \cdot \Delta = 0$ for $i = 5, 6, 7$, thus $C'_5 + C'_6 + C'_7$, being connected, must be a vertical divisor with respect to $|\Delta|$. We conclude the existence of an effective divisor D' such that $\Delta \sim C'_5 + C'_6 + C'_7 + D'$. It is easy to deduce that D' is a rational (-3) -curve and $\phi(D')$ is a line in \mathbb{P}^2 .

From this relation and

$$(C_1 + C'_1) + \dots + (C_7 + C'_7) \sim 7K_S,$$

it follows that:

$$C \sim C_1 + \dots + C_7 \sim 3K_S + D'.$$

This gives a contradiction, as the image $\phi(C)$ is a degree 7 curve and the image $\phi(3K_S + D')$ is a degree 4 curve. \square

Theorem 4.2. *Assume $f : X \rightarrow \mathbb{P}^1$ is obtained as a blow up of a transversal pencil Λ on a minimal surface S with $K_S^2 = 1$ and $p_g = 2$. Then $s \geq 7$.*

Proof. By Proposition 3.3 we must consider only the values $6 \leq g \leq 37$.

Considering the classical Horikawa's construction and notation ([9]), let $\Pi : \bar{S} \rightarrow S$ be the blowing up centered in the base point p of $|K_S|$, denote by E its exceptional divisor and consider the ramified double covering:

$$\phi_2 : \bar{S} \rightarrow \mathbb{F}_2.$$

The map ϕ_2 is given as follow: the bicanonical map of S determines a double cover on the singular quadric $Q \subset \mathbb{P}^3$, the singular point being the image of p . ϕ_2 is the induced map on \bar{S} after considering the desingularization \mathbb{F}_2 of the quadric. The locus branch of ϕ_2 is the divisor $B = 6\Delta + 10\Gamma$, with Δ and Γ denoting respectively the class of the (-2) -section and the class of the fiber in \mathbb{F}_2 of the structural morphism and the ramification divisor R is $5K_S + E$. Here and in what follows given D any divisor in S we just write D for the divisor Π^*D in \bar{S} .

Denote by $\bar{\Lambda}$ the induced pencil on \bar{S} . Depending on whether p is a base point of Λ or not we have

$$(4) \quad \bar{C} = \begin{cases} \Pi^*C & \text{if } p \notin \Lambda \\ \Pi^*C - E & \text{if } p \in \Lambda. \end{cases}$$

Let G be the image of \bar{C} under ϕ_2 . If we denote $G = a\Delta + b\Gamma$ and considering that $\phi^*\Delta = 2E$, $\phi^*\Gamma = K_S - E$ we have

$$(5) \quad \phi^*G = bK_S + (2a - b)E.$$

Let be $\phi := \phi_2|_{\bar{C}} : \bar{C} \rightarrow G$ with $\deg \phi = n = 1$ or 2 .

We analyze the two cases $n = 1$ or 2 and within each one the subcases $p \in \Lambda$ or not.

Case 1 ϕ is 2:1. In this case $\phi^*G = \bar{C}$.

Assume first that $p \notin \Lambda$.

By (4) and (5) we have that $2a - b = 0$, using this $\bar{C}^2 = m = b^2$. On the other hand $m = bC.K_S$, therefore $b = C.K_S$. The next table shows the possible values of m , $C.K_S$ and $g - 1$.

$C.K_S$	a	m	$g - 1$
4	2	16	10
6	3	36	21
8	4	64	36

Assume $g - 1 = 10$. In this case the number of singular points in the fibers of f will be:

$$\begin{aligned} e_f &= 12\chi(\mathcal{O}_X) - K_X^2 + 4(g - 1) \\ &= 36 + 15 + 4(10) = 91 \end{aligned}$$

Since $s = 6$ there exists at least a singular fiber F_0 of f containing 16 singular points. Let $F_0 = F_1 + \dots + F_l$ be the decomposition into irreducible components, then:

$$(6) \quad \begin{aligned} 10 = g - 1 &\geq \sum_{i=1}^l (g_i - 1) + \sigma_0 \\ &\geq \sum (g_i - 1) + 16 \end{aligned}$$

where g_i denotes the geometric genus of F_i . This imply that $l \geq 6$. Moreover there are at least 6 of this components that are rational curves and are mapping onto rational components of G_0 .

From [7] (Corollary V.5.18) we know that the possible irreducible curves in \mathbb{F}_2 are: Γ , Δ and $\alpha\Delta + \beta\Gamma$ with $\alpha > 0$, $\beta \geq 2a$. Let $G_0 = G_1 + \dots + G_s$ be the decomposition into irreducible components.

Note that even being Δ a rational curve, is not a possibility for any of the G_i 's, that because if $G_1 = \Delta$ then

$$\bar{C}_1 = \phi^*G_0 = 2E$$

that contradicts the semistability of f .

With respect to the components $G_i = a_i\Delta + b_i\Gamma$, $a_i > 0$, $b_i \geq 2a$, G_i is rational if and only if $a_i = 1$ and $b_i \geq 2$. Since $a = \sum_{i=1}^s a_i$ and $b = \sum_{i=1}^s b_i$ the only possible

decomposition is $G_0 = G_1 + G_2$ with $G_i = \Delta + 2\Gamma$, $i = 1, 2$, i.e. we can't have the 6 needed rational components.

If $g - 1 = 21, 36$ we use similar arguments with $\sigma_0 = 26, 41$ respectively. In both cases there exist at least 5 rational components and this is not possible because $a < 5$.

Now consider the case $p \in \Lambda$.

By (4) and (5) $b = 2a + 1$, so $m - 1 = \bar{C}^2 = 4a(a + 1)$ and $m = b^2$. Moreover,

$$\bar{C}.K_{\bar{S}} = C.K_S + 1 = b + 1$$

Therefore $m = (C.K_S)^2$. Keeping in mind the previous notation, we get the next possible values

$C.K_S$	a	m	$g - 1$	e_f	σ_0	$l \geq$
3	1	9	6	68	12	6
5	2	25	15	120	20	5
7	3	49	28	196	33	5

Observe that l is, as before, the minimal number of rational components in G_0 , so analogous to the case $p \notin \Lambda$, all possibilities in the table can't occur because of $a < l$.

Case 2 ϕ is 1:1. In this case there exists a divisor \bar{C}' such that $\phi^*G = \bar{C} + \bar{C}'$.

Denote as before by F_0 a singular fiber of f , C_0 its image under π . If $F_0 = F_1 + \dots + F_k$ is the decomposition of F_0 into irreducible components, we denote by $C_0 = C_1 + \dots + C_k$ the corresponding decomposition for C_0 and by $\bar{C}_0 = \bar{C}_1 + \dots + \bar{C}_k$ the corresponding curves and decomposition in \bar{S} and by $G_0 = G_1 + \dots + G_k$ their images in \mathbb{F}_2 .

We begin by stating the following:

Lemma 4.3. *In the previous situation, let G_1 be any irreducible component of G_0 , then neither $G_1 \sim \Delta$, nor $G_1 \sim \Gamma$ nor $G_1 \sim \Delta + 2\Gamma$. In particular, if $G_0 \sim a\Delta + b\Gamma$, then the number of rational irreducible components of G_0 is least or equal than a .*

Proof. Let G_1 be equivalent to Δ . Then $\phi_2^*(G_1) \sim 2E$. Note that there exists a divisor \bar{C}'_1 such that $\bar{C}_1 + \bar{C}'_1 = \phi_2^*(G_1)$. This implies $\bar{C}_1 = E$, which is impossible by the definition of \bar{C} .

Now, assume $G_1 \equiv \Gamma$, then $\phi^*G_1 \equiv K_S - E = C_1 + C'_1$. From this, intersecting with K_S , and using that K_S is nef we obtain that either $C_1.K_S = 0$ or $K_S.C'_1 = 0$. It follows that there exists a curve D on S with $K_S.D = 0$, which contradicts that ampleness of $|K_S|$.

Finally, suppose that $G_1 = \Delta + 2\Gamma$, in this case $\phi^*G_1 \equiv 2K_S$.

Note that for any decomposition $2K_S = A + A'$ we must have that both, A and A' must be irreducible and equivalent to K_S and $A.A' = 1$. Indeed, assume A irreducible, that from $2K_S = A + A'$ with easy it follows that $A.K_S = A'.K_S = 1$. Therefore, $A^2 = 1$, because $2K_S$ is 1-connected, $A \equiv K_S$ follows from HIT.

One established this fact, just note that if $2K_S = C_1 + C'_1$, then $\phi : C_1 \rightarrow G_1$ can not be 1 : 1, because C_1 is a genus 2 curve.

The last assertion follows from the fact that the only irreducible rational curves on \mathbb{F}_2 are equivalent to either Δ , Γ or $\Delta + b\Gamma$ with $b \geq 2$. \square

Continuing the proof of the Theorem, assume first that $p \notin \Lambda$. We have the commutative diagram:

$$\begin{array}{ccc} \bar{S} & \xrightarrow{\phi_2} & \mathbb{F}_2, \\ \Pi \downarrow & & \downarrow \\ S & \xrightarrow{\phi_{2K}} & Q \end{array}$$

therefore $\phi_{2K}(\Pi(\bar{C}'))$ contains the singular point of Q and from this it follows that $\bar{C}' = \Pi^*C'$, for some effective divisor C' in S . By (5) $b = 2a$ and $\bar{C} + \bar{C}' \sim bK_S$. Therefore,

$$C.K_S + C'.K_S = (\bar{C} + \bar{C}')K_{\bar{S}} = (bK_S)(K_S + E) = b.$$

The ramifications of ϕ_2 occurring on \bar{C} are given by the intersections of \bar{C} and \bar{C}' , so we have $\bar{C}.\bar{C}' = \bar{C}.R = \bar{C}'.R$. In particular, the right hand term of the previous equation implies that $C.K_S = C'.K_S$ and $b = 2C.K_S$.

Moreover,

$$bC.K_S = \phi^*G.\bar{C} = (\bar{C} + \bar{C}').\bar{C} = m + \bar{C}.\bar{C}' = m + 5C.K_S.$$

We conclude that $m = C.K_S(2C.K_S - 5)$ and we get the possible values (keeping in mind the previous notation introduced for σ_0 and l):

b	$C.K_S$	m	$g - 1$
8	4	12	8
10	5	25	15
12	6	42	24
14	7	63	35

The values $g - 1 = 24, 35$ are impossible because of the Hodge Index Theorem.

If $g - 1 = 8$ we have that $e_f = 79$ and then there must exist a singular fiber F_0 with at least 14 singular points. It follows that G_0 has 6 or more rational components. Using that $a = 4$ and Lemma 4.3, we get a contradiction. The case $g - 1 = 15$ follows after similar considerations.

It remains to analyze the case $p \in \Lambda$.

As in the previous case, like $\bar{C} = C - E$ also $\bar{C}' = C' - E$. From this we get $2a - b = -2$. Therefore we have:

$$\phi_2^*G = C + C' - 2E = bK_S - 2E.$$

Moreover, $\bar{C}.R = \bar{C}'.R$ and then $C.K_S + 1 = \bar{C}.K_{\bar{S}} = \bar{C}'.K_{\bar{S}}$

We get the next formulas

$$\phi_2^*G.K_{\bar{S}} = b + 2 = 2C.K_S + 2$$

and therefore

$$m + 5C.K_S = \phi_2^*G.\bar{C} = (bK_S - 2E).(C - E) = bC.K_S - 2$$

We conclude that $b = 2C.K_S$ and $m = 2(C.K_S)^2 - 5C.K_S - 2$. The table of possible values is:

$C.K_S$	b	a	m	$g-1$
4	8	3	10	7
5	10	4	23	14
6	12	5	40	23
7	14	6	61	34

If $g-1 = 23, 34$ we get a contradiction by Hodge Index Theorem. If $g-1 = 7$, then $e_f = 73$, therefore there exists a singular fiber F_0 with at least 13 singular points and at least 6 rational components. Taking in consideration that $a = 3$ and Lemma 4.3 we obtain a contradiction. The case $g-1 = 14$ is similar. \square

5. THE ADJOINT CASE

In this section we consider fibrations $f : X \rightarrow \mathbb{P}^1$ satisfying the property that C is an adjoint linear system, i.e., $C \equiv B + K_S$ with B a big and nef divisor. The typical example for bearing in mind is $C \equiv nK_S$, i.e. the fibration f is obtained after blowing up the base locus of a generic pencil of curves $\Lambda \subset |nK_S|$.

We collect, for further use, some general elemental facts in the following:

Lemma 5.1. *Assume $C \equiv B + K_S$ with B a big and nef divisor, denoting by g_B the arithmetic genus of B , we have:*

i)

$$2(g_B - 1) = B^2 + B.K_S.$$

ii)

$$m = (g_B - 1) + (g - 1).$$

iii)

$$(g - 1) = (g_B - 1) + B.K_S + K_S^2.$$

iv) $2 \leq B.K_S$, and if $g \geq 5$, then

$$g + 1 \leq m.$$

Proof. Assertions i)-iii) follow immediately from adjunction formula. As for iv), note that it is enough to prove that $2 \leq B.K_S$, because then using i) $g_B - 1 \geq 2$ and the desired inequality follows from ii). Now, if $B.K_S = 1$, by Index Hodge Theorem $B^2 = K_S^2 = 1$ and applying from $m = B^2 + 2B.K_S + K_S^2$ we have $m = 4$. On the other hand, from i) and ii) $m = 1 + (g - 1) = g \geq 5$. \square

We start by studying the case $K_S^2 = 1$, which is similar in nature to Theorems 4.1 and 4.2:

Proposition 5.2. *Let $f : X \rightarrow \mathbb{P}^1$ be a fibration obtained as the blowing up of the base locus of a transversal and adjoint pencil Λ on a minimal surface with $K_S^2 = 1$. Then, $s \geq 7$.*

Proof. By Noether Inequality $p_g \leq 2$, and by Proposition 3.3 and Theorem 4.2, we can assume $p_g = 1$. It is well known that for a surface with such invariants the bicanonical map ϕ_{2K_S} defines a 4 : 1 morphism onto \mathbb{P}^2 ([5], [10], [14]):

$$\phi_{2K_S} : S \rightarrow \mathbb{P}^2,$$

ramified along a divisor $R \equiv 7K_S$. We consider, as before, the restriction of this map to C :

$$\begin{array}{ccc} C & \xrightarrow{n:1} & \tilde{G} \\ \phi \downarrow & & \nearrow j \\ & & G \end{array}$$

In this case n is a divisor of 4, $G \subset \mathbb{P}^2$ is the image of C and j denotes its normalization. Denote by d the degree of G .

If we assume $s = 6$ we only need, according to Proposition 3.3 to consider $6 \leq g \leq 16$.

We start by analyzing the case $n = 4$: in this case we have $C = \phi^*G$ and therefore:

$$m = C^2 = (\phi^*G)^2 = 4G^2 = 4d^2.$$

On the other hand, taking in account that $\phi^*H \equiv 2K_S$, with H hyperplane section (i.e the divisor associated with $\mathcal{O}_{\mathbb{P}^2}(1)$) we obtain:

$$C.K_S = \phi^*G.K_S = \phi^*(dH).K_S = 2d.$$

Adjunction formula gives $2(g-1) = 2d(2d+1)$. The only value of g that satisfies the relation in the range $6 \leq g \leq 16$ is $g-1 = 10$ with $d = 2$ and $m = 16$. Evaluating in Tan's inequality for $e = 4$ (Lemma 3.1 ii)) we obtain a contradiction.

If $n < 4$, then there exists an effective divisor $C' > 0$ such that:

$$C + C' \equiv \phi_{2K_S}^*G \equiv 2dK_S.$$

Note that:

$$C'.K_S = 2d - C.K_S,$$

and

$$C'^2 = 4d^2 - 4dC.K_S + C^2.$$

Moreover, since $h^1(C) = h^2(C) = 0$,

$$h^0(C) = \frac{C^2 - C.K_S}{2} + 2.$$

Next, the only possibility for $h^2(C') = h^0(K_S - C') \neq 0$ is $C' \equiv K_S$, because $p_g = 1$. This would imply $C \equiv K_S$, which is impossible.

It is easy to prove that $H^0(C) \simeq H^0(C')$ and by Riemann-Roch we get:

$$h^0(C) = h^0(C') \geq \frac{C'^2 - C'.K_S}{2} + 2,$$

and substituting the values of C'^2 and $C'.K_S$:

$$h^0(C) \geq h^0(C') + \frac{4d^2 - (4d-2)C.K_S - 2d}{2}.$$

This implies,

$$2d^2 - (2d-1)C.K_S - d \leq 0,$$

that is equivalent to $d \leq C.K_S$.

Now, assume $n = 1$. The linear system $|2K_S|_C$ defines a base point free linear system on C and the associated map a $1 : 1$ cover onto a plane degree d curve.

Thus, we have $d = 2C.K_S$ and we obtain a contradiction with the just obtained bound $d \leq C.K_S$.

The case $n = 2$ remains to be analyzed: in this case we have $C.K_S = d$. Note that the intersections of C and C' gives place to ramifications points of ϕ_{2K_S} . Therefore:

$$2dC.K_S - m = C.C' \leq C.R = 7C.K_S.$$

From this we get:

$$d(2d - 7) \leq m \leq 2d^2 - 2.$$

It follows that $d \leq 5$. In general we have that

$$(7) \quad \begin{aligned} e_f &= e(X) + 4(g - 1) \\ &= 24 - (1 - m) + 4(g - 1), \\ &= 23 - d + 6(g - 1). \end{aligned}$$

Thus, assuming $s = 6$ and $d \leq 3$, there must exist a singular fiber F_0 of f having at least $(g - 1) + 4$ nodes. Call σ_0 the number of nodes of F_0 . Note that the number of nodes of $C_0 = \pi(F_0)$ is also σ_0 . Then, $\sigma_0 \geq 9$. On the other hand, the plane curve $G_0 = \phi_{2K_S}(C_0)$, being of degree $d \leq 3$ admits at most 3 nodes. In this way we get the contradiction $\sigma_0 \leq 6$.

Similar argumentations lead to contradictions for the cases $d = 4, 5$. Indeed, if $d = 4$, then $\sigma_0 \geq g + 3$. We have:

$$g - 1 = \sum_{i=1}^l (g_i - 1) + \sigma_0 \geq -l + g + 3,$$

with g_i standing for the geometric genus of the components F_i of F_0 . It follows that $l \geq 4$ and therefore F_0 (and in consequence C_0) has at least 4 rational components. From this it follows that G_0 has at least 2 irreducible rational components. Taking in account that G_0 is a degree 4 curve we have that, either G_0 contains a line as an irreducible component or it is the product of two irreducible conics. If G_0 is the product of two irreducible conics then it has only 4 nodes and we get, as before a contradiction, in any other case, if L is an irreducible component of G_0 and $C_0 = C_1 + \dots + C_l$, then

$$\phi_{2K_S}^* L = C_i + C'_i,$$

for C_i some rational components of C_0 . But then:

$$2K_S \equiv C_i + C'_i,$$

and it follows ([4] Lemma 1, page 181) that $C_i = \Delta$, the only effective divisor in $|K_S|$. This gives a contradiction, since Δ is a curve of geometric genus 2.

Finally the case $d = 5$ follows after similar considerations. In this case C_0 admits at least 3 irreducible rational components and G_0 at least 2 irreducible rational components. The only subtle case to be treated carefully being the possibility that $G_0 = Q + E$, with Q an irreducible conic and E a singular irreducible cubic. But in this case either Q or E must satisfy that it pull back under ϕ_{2K_S} is the sum of two irreducible components $C_i + C_j$ of C_0 .

Note that $\phi_{2K_S}^* E = C_i + C_j$ is impossible, because then

$$6K_S \equiv C_i + C_j$$

and then $K_S.(C_i + C_j) = 6$, that contradicts $K_S.C_0 = 5$. On the other hand

$$4K_S = \phi_{2K_S}^* Q = C_i + C_j$$

implies that C_0 has exactly 3 irreducible components: $C_0 = C_1 + C_2 + C_3$. Suppose $i = 1$ and $j = 2$, then $C_3.K_S = 1$ and

$$\phi_{2K_S} : C_3 \rightarrow E,$$

must be a 2 : 1 map onto a degree cubic and we obtain the contradiction

$$2K_S.C_3 = 6.$$

□

Finally we have:

Theorem 5.3. *Let $f : X \rightarrow \mathbb{P}^1$ a semi-stable non-isotrivial fibration obtained as the blow-up of the base locus of an adjoint pencil Λ on the minimal surface S . Then if $K_S^2 \geq 3$ the number s of singular fibers of f is at least 7. If $K_S^2 \leq 2$ and Λ is also transversal, then $s \geq 7$.*

Proof. We assume $s = 6$ and $2 \leq K_S^2$. From Noether's inequality:

$$p_g \leq \frac{K_S^2}{2} + 2,$$

we have:

$$\chi(\mathcal{O}_S) \leq \frac{K_S^2}{2} + 3.$$

Applying Lemma 3.3:

$$K_S^2 + \sqrt{8K_S^2(g-1) + (K_S^2)^2} \leq 3K_S^2 + 18,$$

which implies

$$8K_S^2(g-1) + (K_S^2)^2 \leq 4(K_S^2)^2 + 72K_S^2 + 18^2.$$

Substituting $2(g-1) \geq C.K_S + m = B.K_S + K_S^2 + m$, we get:

$$m \leq \frac{72 - 4B.K_S}{4} + \frac{18^2 - (K_S^2)^2}{4K_S^2}.$$

Now, combine the previous bound for m with Lemma 3.1 ii), in order to deduce:

$$19K_S^2 + 19B.K_S \leq 18 + \frac{18^2 - (K_S^2)^2}{4K_S^2} + 108.$$

Using again the adjoint hypothesis, this amount to:

$$\frac{(K_S^2)}{4} + 19C.K_S \leq 166,$$

that is,

$$C.K_S \leq 8.$$

Now, use Hodge Index Theorem:

$$m \leq \frac{(C.K_S)^2}{K_S^2} \leq \frac{64}{K_S^2},$$

and apply one more time Lemma 3.1 ii):

$$K_S^2 + 18C.K_S \leq \frac{64}{K_S^2} + 108,$$

$$19K_S^2 + 18B.K_S \leq \frac{64}{K_S^2} + 108.$$

Finally, using $2 \leq B.K_S$, (Lemma 5.1 iv)) we arrive to:

$$19K_S^2 \leq \frac{64}{K_S^2} + 72.$$

This implies $K_S^2 \leq 4$. Thus, only the cases $K_S^2 = 2, 3, 4$ remains to be discharged. This is easy and essentially is a reproduction of the previous argument.

For instance, for case $K_S^2 = 2$ we have by Lemma 3.2 and Proposition 3.3 that $g \leq 16$ and $m \leq 26$, moreover, assuming Λ is transversal, we can apply Theorem 4.1 and assume that $\chi(\mathcal{O}_S) \leq 3$. Evaluating Tan's Inequality at $e = 5$ (Lemma 3.1 iii)) we get:

$$7K_S^2 + 6C.K_S \leq m + 9\chi(\mathcal{O}_S),$$

that, under our fixed values becomes $C.K_S \leq 6$. Using Hodge Index Theorem we obtain $m \leq 18$. Evaluating again Tan's Inequality at $e = 4$ we have $C.K_S \leq 4$ and $m \leq 8$. Finally, we evaluate once again Tan's Inequality at $e = 5$ and get the final contradiction $C.K_S \leq 3$ and $g + 1 \leq m \leq 4$.

Cases $K_S^2 = 3, 4$ are quite analogous, only that in these cases we don't need Theorem 4.1, and in consequence the transversality hypothesis can be avoided. \square

REFERENCES

- [1] Barth, W., Peters, C., Van de Ven, A. *Compact Complex Surfaces* Springer Verlag (1984).
- [2] Beauville, A.: *Le nombre minimum de fibres singulieres d'un courbe stable sur \mathbb{P}^1* . Asterisque, 86, (1981), 97-108.
- [3] Beauville A.: *Surfaces algébriques complexes*. Asterisque 54, (1978).
- [4] Bombieri, E. *Canonical models of surfaces of general type*. Publ. Math. IHES, 42, (1973), 171-219.
- [5] Catanese, F. *Surfaces with $K^2 = pg = 1$ and their period mapping*. In Algebraic geometry (Proc. Summer Meeting, Univ. Copenhagen, 1978), Springer LNM, 732 (1979), 1-29.
- [6] Debarre, O. *Inégalités numériques pour les surfaces de type général*. Bull. Soc. Math. France 110 (3), (1982), 319-346.
- [7] Hartshorne, R. *Algebraic Geometry*. Springer-Verlag GTM, (1997).
- [8] Horikawa, E. *Algebraic surfaces of general type with small c_1^2 . I*. Annals of Mathematics, 104, (1976), 357-387.
- [9] Horikawa, E. *Algebraic surfaces of general type with small c_1^2 II*, Inventiones Math. 37 (1976), 121-155.
- [10] Kunev, V. Thesis for Masters Degree. Sofia University, (1976).
- [11] Liu, K. *Remarks on the Geometry of Moduli Spaces* Proceedings of the AMS 124, (3), (1996), 689-695.
- [12] Tan S-L: *The minimal number of singular fibers of a semi stable curve over \mathbb{P}^1* . J. Algebraic Geom., 4, (1995), 591-596.

- [13] Tan S-L , Tu Y. , Zamora A. G. *On complex surfaces with 5 or 6 semistable singular fibers over \mathbb{P}^1* . Math. Zeitschrift 249 (2005), 427-438.
- [14] Todorov, A. N. *Surfaces of General type with $p_g = 1$ and $(K, K) = 1$. I* Annales scientifiques de l'E.N.S. 13 (1980), 1-21.
- [15] Zamora, A. G. *Genus 5 general type \mathbb{P}^1 -semistable curves have at least 7 singular points.* Note di Matematica 32 (2) (2012), 1-4.

CENTRO DE CIENCIAS MATEMÁTICAS,
UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, CAMPUS MORELIA.
APARTADO POSTAL 61-3, SANTA MARÍA, 58089.
MORELIA, MICHOACÁN, MÉXICO.

E-mail address: `huitrado at matmor.unam.mx`

CENTRO DE CIENCIAS MATEMÁTICAS,
UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, CAMPUS MORELIA.
APARTADO POSTAL 61-3, SANTA MARÍA, 58089.
MORELIA, MICHOACÁN, MÉXICO.

E-mail address: `mcastaneda at matmor.unam.mx`

U. A. MATEMÁTICAS, U.DE ZACATECAS
CAMINO A LA BUFA Y CALZADA SOLIDARIDAD, C.P. 98000
ZACATECAS, ZAC. MÉXICO

E-mail address: `alexiszamora06 at gmail.com`