

A CHERNOFF BOUND FOR BRANCHING RANDOM WALK

Changqing Liu
 ClinTFL Ltd.
 c.liu@ClinTFL.com

Concentration inequalities, which prove to be very useful in a variety of fields, provide fairly tight bounds for large deviation probability while central limit theorem (CLT) describes the asymptotic distribution around the mean (within scope of \sqrt{n} order). Harris (1963) conjectured that for a supercritical branching random walk (BRW) of i.i.d offspring and iid displacement, population's positions in n th generation approach to Gaussian distribution — central limit theorem. This conjecture was latter proved by Stam (1966) and Kaplan & Asmussen (1976). Refinements and extensions followed. Yet little effort is known on large deviation probability for BRW. In this note, we suggest and verify a more general and probably more formal setting of BRW. Benefiting from this framework, a Chernoff bound for BRW is immediately obtained. The relation between RW (random walk) and BRW is addressed.

1. Introduction.

1.1. *Concentration inequalities.* Let $(S_i)_{i=1,2,\dots}$ be a real-valued random process, one-dimensional random walk. S_n can be formulated as $S_n = X_1 + X_2 + \dots + X_n$; $X_i = S_i - S_{i-1}$. If some conditions are satisfied, among which independence is crucial, S_n converges to a Gaussian distribution in a \sqrt{n} neighborhood of its mean — central limit theorem (CLT). For S_n beyond the neighborhood, i.e., $|S_n - E(S_n)| \gg \sqrt{n}$, Chernoff inequality ([14] [21]), also referred as Chernoff bound,

$$(1.1) \quad Pr(|S_n - E(S_n)| \geq \lambda) \leq e^{-c \frac{\lambda^2}{n}}$$

tells us how unlikely this occurrence is. Chernoff inequality is extended to the more general context of bounded martingale difference (say, $E(X_n | S_{n-1}) = 0$), referred as Azuma-Hoeffding inequality [4] (see [15], [28] for the survey and references therein). Refinements and extensions followed (e.g. [33], [28], [6], [7], [18], [25]). Known as concentration inequalities, these prove very useful and have a wide variety of applications in computer science, combinatorics, information theory (see e.g. [17], [3], [31], [30]).

1.2. *Branching random walk (BRW).* In the literature, a branching random walk on the real line is described as follows (see, for example, [11] [8] [9]). In generation zero, an initial particle at the original place on the real line \mathbf{R} . It splits into a random number of child particles who form generation one. The children's displacements, relative to their parent, correspond to a point process on \mathbf{R} . The children in turn split too to form the second generation, and so on. If the average split number (branching factor) is greater than one, with positive probability the number of the descendants grows exponentially through generations. Current BRW studies typically address models where the offspring's behavior is independent of that of their previous generation (e.g. [8], [9] [10], [22], [19]). The law of large numbers and central limit theorem type results about the distribution of position are established under certain conditions of independence (for instance, i.i.d

Key words: Concentration inequality, Chernoff bound, Branching random walk

of branching and walking). Yet unlike random walk, Chernoff bound is not known so far even in the case of i.i.d. aforementioned, while the minimal (and maximal) is studied by many.

Considering i.i.d. offspring (and hence independent of the parent's position), and i.i.d. displacement, Harris [20] conjectured that the distribution of the descendants' position of the n th generation approaches Gaussian distribution. This conjecture was proved by [32], [23]. Its extended generation-dependent versions, where offspring and displacement distribution are determined by generation n , were proved by [12], [24], to mention a few. Problems where offspring and displacement are dependent of parents' positions are studied in adhoc approaches though (e.g. [34]). Concerning the deviation from the expectation, [9] proved that for any $\delta > 0$ the number of particles locating beyond $\mu - \delta$ is zero almost surely, μ denoting the scaled expectation. In another direction of estimating the rareness of the large deviation, extremum is well studied (e.g. [16] [29], [5], [13] [22], [1] [2]). [16] showed tightness for $M_n - EM_n$, where M_n is minimum of n th generation. [29] gave probability bound for the deviation of M_n , i.e. $Pr(M_n - Med_n > x) < e^{-\delta x}$, where Med_n is the median of M_n . So far probability bound for only deviation ϵ of appropriate scale ($\epsilon > 0$) is established. Little is known as to how sparse the population is at a distance of $O(\sqrt{n})$ far from the mean in terms of concentration inequalities.

2. Chernoff bound for BRW. Throughout we consider BRW on \mathbf{N} for notational simplicity, though the argument applies to \mathbf{R} . We start by suggesting a new setting for BRW, which is more general in three aspects than the classical in the literature. First, in our framework both subcritical and supercritical BRW are treated equally. While traditionally subcritical BRW is considered trivial because a branching random walk almost surely ends with zero population when n is large. We argue that the probability space for the survival BRW processes is well defined with large enough number of ancestors (say, as large as exponential in n). Second, the underlying random walks between siblings are not assumed to be independent. Third, branching factor (birth rate) is not assumed identical across generations and siblings. The only major requirement is the independence between birth-rate and birth-place. Specifically, we define a BRW as a sequence of pair

$$(m_i(u), p_i(x|u))_{i=1,2,\dots,n}$$

where $m_i(u)$ is the expectation of offsprings (branching factor) of a parent at position u , and $p_i(x|u)$ is offsprings' displacement (relative to their parent's position) p.d.f, the probability (or proportion) density function, or mass probability in discrete cases.

Starting from one initial ancestor (1st parent), a realization of branching random walk process constitutes a rooted tree. Each node in the tree is associated with a position which equals to the sum of the displacements of its previous parents and the displacement of itself, u . As a parent, this node in turn produces some number of children with u as the birth place. Infinite BRW processes with such initial ancestors form a forest, a probability space of the BRW. In other words, the probability space of the BRW can be interpreted as a forest grows from infinite such initial ancestors. Let $S_n = X_1 + X_2 + \dots + X_n$ be the position of a leaf in the n th generation, where X_i is displacement (step size relative from the birth place) of its i th parent. (X_1, X_2, \dots, X_n) is called a *spine* (a path from the original to S_n). The central limit theorem for BRW says that almost surely a randomly chosen tree has a Gaussian "canopy"; namely, S_n on the tree has Gaussian distribution. Let u_i be an individual's position, i.e. $X_1 + \dots + X_i$, at generation i . At position u_i , the mean of birth rate is $m_i(u)$ (generally the mean of birth rate may be dependent on the birth place).

Let M_0 be the number of initial ancestors which may approach to infinity, and $M_i(u)$ the total population of i th generation at position u , in the forest. The population, at $i + 1$ generation,

produced by those $M_i(u)$ particles can be written as

$$M_i(u) \sum_x m_i(u) p_{i+1}(x)$$

where

$$M_i(u) \cdot m_i(u) p_{i+1}(x)$$

is population at $u + x$, $m_i(u)$ is the mean of birth rate of a (parent) particle at u in i th generation, and $\sum_x p_{i+1}(x) = 1$. Note, $p_{i+1}(x)$ is dependent of u which we drop off for notational simplicity; though given u , $p_{i+1}(x)$ is a function of x only. With

$$M_0 \cdot m_1(u_1) p_2(x) \quad (u_1 = X_1)$$

being the population (of 2nd generation) at $u_1 + x$ produced by 1st generation, we have by induction the total population at generation n (note, $p_1(u_1) = 1$)

$$\sum_{x_1} \sum_{x_2} \dots \sum_{x_n} M_0 p_1(x_1) m_1(u_1) p_2(x_2) \dots m_{n-1}(u_{n-1}) p_n(x_n)$$

and the law of spine (x_1, x_2, \dots, x_n)

$$(2.1) \quad \frac{p_1(x_1) m_1(u_1) p_2(x_2) \dots m_{n-1}(u_{n-1}) p_n(x_n)}{\sum_{x_1} \sum_{x_2} \dots \sum_{x_n} p_1(x_1) m_1(u_1) p_2(x_2) \dots m_{n-1}(u_{n-1}) p_n(x_n)}$$

which can be interpreted as the proportion of (x_1, x_2, \dots, x_n) in the whole forest. As aforementioned earlier in this section, we are concerned with BRW where birth rate is independent of birth position. In this case, the law of spine (2.1) is reduced to

$$(2.2) \quad p_1(x_1) p_2(x_2) \dots p_n(x_n)$$

which is the probability distribution of the random walk $X_1 + X_2 + \dots + X_n$ where $Pr(X_i = x \mid u_{i-1}) = p_i(x)$. This observation turns questions about BRW into questions about random walk (without branching).

Remark.

- As far as the stochastic behavior at generation n is concerned, random walk can be viewed as a special BRW (with branching factor = 1), or random walk is a special BRW where birth-rate is independent of birth-place.

- A BRW of spatial homogeneity in branching (i.e. $m_n(u) = m_n$) can be treated as a random walk process (without branching).

In the following, we refer the random walk by $(p_i)_{i=1,2,\dots,n}$ (note this random walk is not necessarily the underlying random walk of the original BRW, though under the condition of independence between siblings' motion, the underlying random walk is (p_i)). The following theorem presents a BRW version of Chernoff bound.

THEOREM 2.1. *For BRW $(m_i, p_i)_{i=1,2,\dots,n}$, if (a) m_i is position-independent, and (b) Chernoff bound holds for random walk (p_i) , then*

$$Pr(|Q_n(\alpha) - na| \geq \lambda) \leq 2e^{-\frac{1}{2}c\lambda^2/n}$$

where na is the expected position¹ for an individual in n th generation, $Q_n(\alpha)$ is α quantile and $\alpha \in [e^{-\frac{1}{2}c\lambda^2/n}, 1 - e^{-\frac{1}{2}c\lambda^2/n}]$. Throughout, generation index n will be omitted when no confusion can arise.

The theorem conveys the same idea as Chernoff inequalities for random walk, stating that “almost surely” (with as small as about $e^{-O(1)\lambda^2/n}$ chance of exception) a BRW tree’s almost all leaves (about $1 - e^{-O(1)\lambda^2/n}$ proportion) are concentrating around the expected position within area of \sqrt{n} order (when λ is set $O(\sqrt{n})$).

PROOF. Let $z_1^{(n)}, z_2^{(n)}, \dots$ be an enumeration of the positions of the particles (leaves) in the n th generation and $Z^{(n)}$ be its population; i.e. $Z^{(n)} = |\{z_1^{(n)}, z_2^{(n)}, \dots\}|$. There should be an index variable τ for trees in the above notations which we omit. Define

$$F(t) = \frac{\sum_i \mathbf{1}_{\{z_i \leq t\}}}{Z^{(n)}}$$

namely, the cumulative distribution function of the data set $\{z_1^{(n)}, z_2^{(n)}, \dots\}$. The $1 - \alpha$ quantile is

$$Q_n(1 - \alpha) = \inf\{t : F(t) \geq 1 - \alpha\}$$

Let τ be a tree of Z leaves and $p_{(\tau)}(\lambda)$ be the proportion of positions which $\geq na + \lambda$ (in the n th generation), i.e.

$$p_{(\tau)}(\lambda) = \frac{\sum_{i=1}^Z \mathbf{1}_{\{z_i(\tau) \geq na + \lambda\}}}{Z}$$

By the definitions,

$$p_{(\tau)}(\lambda) \geq \alpha \text{ iff } Q_n(1 - \alpha) \geq na + \lambda$$

We have $E(p_{(\tau)}(\lambda)) = Pr(S_n - na \geq \lambda)$, because

$$(2.3) \quad E(p_{(\tau)}(\lambda)) = \frac{\sum_{i=1}^Z E(\mathbf{1}_{\{z_i - na \geq \lambda\}})}{Z} = \frac{\sum_1^Z Pr(S_n - na \geq \lambda)}{Z} = Pr(S_n - na \geq \lambda)$$

where τ is a tree of population Z . In other words, the expectation of $p_{(\tau)}(\lambda)$ over all the trees of the same population is $Pr(S_n - na \geq \lambda)$. The reason Z does not appear in $E(p_{(\tau)}(\lambda))$ is that for a randomly selected leaves z_i from a tree $E(\mathbf{1}_{\{z_i - na \geq \lambda\}})$ is independent of the size of the tree because branching is independent of walking. Branching is also independent from leaf indexing so that $E(\mathbf{1}_{\{z_i - na \geq \lambda\}})$ is the same for any i . Below we will use k instead of capital Z for readability in summation.

On the other hand, denoting the number of trees of size k by n_k , we have

$$(2.4) \quad \frac{\sum_{i=1}^k E(\mathbf{1}_{\{z_i - na \geq \lambda\}})}{k} = \frac{1}{k} \sum_{i=1}^k \frac{1}{n_k} \sum_{\tau_k} \mathbf{1}_{\{z_i(\tau_k) - na \geq \lambda\}}$$

¹“ na ” is used for mean in order to be consistent with the literature of BRW, not indicating a linear relationship with n

Note n_k is very large for we have infinity initial ancestors. Suppose there are N_t trees in the forest, n_1 trees of population 1, n_2 trees of population 2, ... and so on. In view of above equations, we have

$$\begin{aligned}
Pr(S_n - na \geq \lambda) &= \sum_{k=1,2,\dots} \frac{n_k}{N_t} \frac{1}{k} \sum_{i=1}^k \frac{\sum \tau_k \mathbf{1}_{\{z_i(\tau_k) - na \geq \lambda\}}}{n_k} \quad (2.3) \text{ and then } (2.4) \\
&= \frac{1}{N_t} \sum_{k=1,2,\dots} \sum_{\tau_k} \frac{\sum_{i=1}^k \mathbf{1}_{\{z_i(\tau_k) - na \geq \lambda\}}}{k} \\
&= \frac{1}{N_t} \sum_{\tau} \frac{\sum_{i=1}^k \mathbf{1}_{\{z_i(\tau) - na \geq \lambda\}}}{k} \\
&= \frac{1}{N_t} \sum_{\tau} p_{(\tau)}(\lambda) \\
&\geq \frac{1}{N_t} \sum_{\tau} p_{(\tau)}(\lambda) \mathbf{1}_{\{p_{(\tau)}(\lambda) \geq \alpha\}} \\
&\geq \frac{1}{N_t} \sum_{p_{(\tau)}(\lambda) \geq \alpha} \alpha \\
&= \alpha \cdot Pr(p_{(\tau)}(\lambda) \geq \alpha) \\
&= \alpha \cdot Pr(Q_n(1 - \alpha) \geq na + \lambda)
\end{aligned}$$

By chernoff bound on the left-hand side of the above, for a certain constant $c > 0$

$$\alpha \cdot Pr(Q_n(1 - \alpha) \geq na + \lambda) \leq e^{-c\lambda^2/n}$$

Choosing $\alpha = e^{-\frac{1}{2}c\lambda^2/n}$, we have

$$(2.5) \quad Pr(Q_n(1 - e^{-\frac{1}{2}c\lambda^2/n}) - na \geq \lambda) \leq e^{-\frac{1}{2}c\lambda^2/n}$$

and therefore

$$(2.6) \quad Pr(Q_n(\alpha) - na \geq \lambda) \leq e^{-\frac{1}{2}c\lambda^2/n}$$

for $\alpha \leq 1 - e^{-\frac{1}{2}c\lambda^2/n}$.

Similarly, for $\alpha \geq e^{-\frac{1}{2}c\lambda^2/n}$

$$(2.7) \quad Pr(Q_n(\alpha) - na \leq -\lambda) \leq e^{-\frac{1}{2}c\lambda^2/n}$$

The claim follows. □

Remark.

- The Chernoff bound holds for both supercritical and subcritical BRW.
- If displacements of siblings are independent of each other, with step X_i then, p_i and the law of X_i are identical (Note, generally they are not equal). In other words, if $S_n = X_1 + \dots + X_n$ is a random walk with (p_i) as the probability density of the increment, then the distribution of S_n has the same "shape" as the forest of any such a BRW (m_i, p_i) ; originally, probability space of random walk is interpreted as the forest of BRW of $(1, p_i)$.

• In CSP (constraints satisfaction problem) (e.g. K-SAT [26]), enumerating the whole problem instances can be formulated as a branching random walk. In this BRW the forest has only one tree because every tree is the same; m_i is not random, given birth place, and in addition branching factor is large (say $(2n)^k$). Because of dependence of branching and position, the BRW can not be reduced to random walk; (2.1) can not be reduced to (2.2). The concentration inequalities for this BRW are developed in a separate paper ([27]).

REFERENCES

- [1] ADDARIO-BERRY, L. AND REED, B. (2009). *Minima in branching random walks*. Ann. Probab. 37, 1044-1079
- [2] AÏDÉKON, E. (2013). *Convergence in law of the minimum of a branching random walk*. Ann. Probab. 41, 1362-1426.
- [3] N. ALON AND J. H. SPENCER (2000). *The Probabilistic Method*. Second Edition, Wiley, New York, 2000, page 162.
- [4] KAZUOKI AZUMA (1967). *Weighted sums of certain dependent random variables*. Tohoku Math. J. (2) 19 357-367
- [5] BACHMANN, M. (2000). *Limit theorems for the minimal position in a branching random walk with independent logconcave displacements*. Adv. in Appl. Probab. 32 159-176.
- [6] V. BENTKUS (2004). *On Hoeffdings inequalities*. Ann. Probab., 32(2), 1650-1673, 2004
- [7] V. BENTKUS (2008). *An extension of the Hoeffding inequality to unbounded random variables*. Lith. Math. J. 48,137-157
- [8] BIGGINS, J. D. (1977). *Martingale convergence in the branching random walk*. J. Appl. Prob. 14, 25-37.
- [9] BIGGINS, J.D. (1977). *Chernoff's Theorem in the branching random walk*. J. Appl. Probab. 14, 630-636.
- [10] BIGGINS, J. D. AND KYPRIANOU, A. E. (1997). *Seneta-Heyde norming in the branching random walk*. Ann. Probab. 25 337-360.
- [11] BIGGINS, J. D. (1976). *The First- and Last-Birth Problems for a Multitype Age-Dependent Branching Process*. Advances in Applied Probability. Vol. 8, No. 3, 446-459
- [12] J. D. BIGGINS (1990). *The central limit theorem for the supercritical branching random walk, and related results*. Stochastic Process. Appl. 34 (1990), no. 2, 255-274.
- [13] BRAMSON, M. D. AND ZEITOUNI, O. (2009). *Tightness for a family of recursion equations*. Ann. Probab. 37 615-653.
- [14] H. CHERNOFF (1952). *A Measure of Asymptotic Efficiency for Tests of a Hypothesis Based on the sum of Observations*. Annals of Mathematical Statistics 23 (4), 493-507
- [15] F. CHUNG AND L. LU (2006). *Concentration inequalities and martingale inequalities: A survey*. Internet Mathematics, 3(1):79-127, 2006.
- [16] DEKKING, F. M. AND HOST, B. (1991). *Limit distributions for minimal displacement of branching random walks*. Probab. Theory Related Fields 90 403-426
- [17] D. P. DUBASHI AND A. PANCONESI (2009). *Concentration of Measure for the Analysis of Randomized Algorithms*. Cambridge University Press, 2009.
- [18] X. FAN, I. GRAMA, AND Q. LIU (2012). *Hoeffding's inequality for supermartingales*. Stochastic Processes and their Applications, vol. 122, no. 10, pp. 3545-3559,
- [19] Z. GAO, Q. LIU, AND H. WANG. (2014). *Central limit theorems for a branching random walk with a random environment in time*. Acta Math. Sci. Ser. B Engl. Ed., 34(2): 501-512, 2014.
- [20] HARRIS, T.E. (1963). *The Theory of Branching Processes*. Springer, Berlin.
- [21] W. Hoeffding (1963). *Probability inequalities for sums of bounded random variables*. J. Amer. Statist. Assoc. 58 (1963) 13-30.
- [22] Y. HU AND Z. SHI (2009). *Minimal position and critical martingale convergence in branching random walks, and directed polymers on disordered trees*. Ann. Probab. 37 742-789.
- [23] N. KAPLAN AND S. ASMUSSEN (1976). *Branching random walks. II*. Stochastic Processes Appl., 4(1): 15-31
- [24] C. F. KLEBANER (1982). *Branching random walk in varying environments*. Adv. in Appl. Probab. 14 (1982), no. 2, 359-367.
- [25] A. KONTOROVICH (2014). *Concentration in unbounded metric spaces and algorithmic stability*. Proc. of ICML 2014
- [26] C. LIU (2016). *Concentration Inequalities for Branching Random Walk and Applications (II): On the threshold of K-SAT/q-COL*.

- [27] C. LIU (2016). *Concentration Inequalities for Branching Random Walk and Applications (I)*.
- [28] C. McDIARMID (1998). *Concentration*, In Probabilistic Methods for Algorithmic Discrete Mathematics, edited by M. Habib, C. McDiarmid, J. RamirezAlfonsin, and B. Reed, pp. 195-248, Algorithms and Combinatorics 16. Berlin: Springer, 1998.
- [29] C. McDIARMID (1995). *Minimal positions in a branching random walk*. Ann. Appl. Probab. Vol. 5, No. 1, 128–139
- [30] R. MOTWANI AND P. RAGHAVAN (1990). *Randomized Algorithms*. Cambridge University Press, Cambridge, MA, USA, 1995.
- [31] M. RAGINSKY AND I. SASON (2013). *Concentration of measure inequalities in information theory, communications, and coding*. Foundations and Trends in Communications and Information Theory 10 (2013), no. 1–2, 1-246. 24
- [32] A.J. STAM (1966). *On a conjecture of Harris*. Z. Wahrsch. Verw. Geb. 5. 202–206 (1966)
- [33] M. TALAGRAND (1995). *Concentration of measure and isoperimetric inequalities in product spaces*. Inst. Hautes Etudes Sci. Publ. Math. 81, 73-205
- [34] YOSHIDA, N. (2008). *Central limit theorem for branching random walks in random environment*. Ann. Appl. Probab. Volume 18, Number 4 (2008), 1619-1635.