

AN EXAMPLE OF A NON-BOREL LOCALLY-CONNECTED FINITE-DIMENSIONAL TOPOLOGICAL GROUP

I. BANAKH, T. BANAKH, M. VOVK

ABSTRACT. Answering a question posed by S.Maillot in MathOverflow, for every $n \in \mathbb{N}$ we construct a locally connected subgroup $G \subset \mathbb{R}^{n+1}$ of dimension $\dim(G) = n$, which is not locally compact.

By a classical result of Gleason [3] and Montgomery [6], every locally path-connected finite-dimensional topological group G is locally compact. Generalizing this result of Gleason and Montgomery, Banakh and Zdomskyy [1] proved that a topological group G is locally compact if G is compactly finite-dimensional and locally continuum-connected. In [5] Sylvain Maillot asked if the locally path-connectedness in the result of Gleason and Montgomery can be replaced by the local connectedness. In this paper we construct a counterexample to this question of S. Maillot. We recall that a subset B of a Polish space X is called a *Bernstein set* in X if both B and $X \setminus B$ meet every uncountable closed subset F of X . Bernstein sets in Polish space can be easily constructed by transfinite induction, see [4, 8.24].

Theorem 1. *For every $n \geq 1$ the Euclidean space \mathbb{R}^n contains a dense additive subgroup $G \subset \mathbb{R}^n$ such that*

- (1) G is a Bernstein set in \mathbb{R}^n ;
- (2) G is locally connected;
- (3) G has dimension $\dim(G) = n - 1$;
- (4) G is not Borel and hence not locally compact.

Proof. Let $(F_\alpha)_{\alpha < \mathfrak{c}}$ be an enumeration of all uncountable closed subsets of \mathbb{R}^n by ordinal $\alpha < \mathfrak{c}$. Fix any point $p \in \mathbb{R}^n \setminus \{0\}$. By transfinite induction, for every ordinal $\alpha < \mathfrak{c}$ we shall choose a point $z_\alpha \in F_\alpha$ such that the subgroup $G_\alpha \subset \mathbb{R}^n$ generated by the set $\{z_\beta\}_{\beta < \alpha}$ does not contain the point p . Assume that for some ordinal $\alpha < \mathfrak{c}$ we have chosen points z_β , $\beta < \alpha$, so that the subgroup $G_{<\alpha}$ generated by the set $\{z_\beta\}_{\beta < \alpha}$ does not contain p . Consider the set $Z = \{\frac{1}{n}(p - g) : n \in \mathbb{Z} \setminus \{0\}, g \in G_{<\alpha}\}$ and observe that it has cardinality $|Z| \leq \omega \cdot |G_{<\alpha}| \leq \omega + |\alpha| < \mathfrak{c}$. Since the uncountable closed subset F_α of \mathbb{R}^n has cardinality $|F_\alpha| = \mathfrak{c}$ (see [4, 6.5]), there is a point $z_\alpha \in F_\alpha \setminus Z$. For this point we get $p \neq nz_\alpha + g$ for any $n \in \mathbb{Z} \setminus \{0\}$, and $g \in G_{<\alpha}$. Consequently, the subgroup $G_\alpha = \{nz_\alpha + g : n \in \mathbb{Z}, g \in G_{<\alpha}\}$ generated by the set $\{z_\beta\}_{\beta \leq \alpha}$ does not contain the point p . This completes the inductive step.

After completing the inductive construction, consider the subgroup G generated by the set $\{a_\alpha\}_{\alpha < \mathfrak{c}}$ and observe that $p \notin G$ and G meets every uncountable closed subset F of \mathbb{R}^n . Moreover, since G meets the closed uncountable set $F - p$, the coset $p + G \subset \mathbb{R}^n \setminus G$ meets F . So, both the subgroup G and its complement $\mathbb{R}^n \setminus G$ meet each uncountable closed subset of \mathbb{R}^n , which means that G is a Bernstein set in \mathbb{R}^n . The following proposition implies that the group G has properties (2)–(4). \square

Proposition 1. *Let $n \geq 2$. Every Bernstein subset B of \mathbb{R}^n has the following properties:*

- (1) *B is not Borel;*
- (2) *B is connected and locally connected;*
- (3) *B has dimension $\dim(B) = n - 1$.*

Proof. 1. By [4, 8.24], the Bernstein set B is not Borel (more precisely, B does not have the Baire property in \mathbb{R}^n).

2. To prove that B is connected and locally connected, it suffices to prove that for every open subset $U \subset \mathbb{R}^n$ homeomorphic to \mathbb{R}^n the intersection $U \cap B$ is connected. Assuming the opposite, we could find two non-empty open disjoint sets $U_1, U_2 \subset U$ such that $U \cap B = (U_1 \cap B) \cup (U_2 \cap B)$. Consider the complement $F = U \setminus (U_1 \cup U_2) \subset U \setminus B$ and observe that F is closed in U and hence of type F_σ in \mathbb{R}^n . If F is uncountable, then F contains an uncountable closed subset of \mathbb{R}^n and hence meets the set B , which is not the case. So, the closed subset F of U is at most countable and separates the space $U \cong \mathbb{R}^n$, which contradicts Theorem 1.8.14 of [2].

3. Since the subset B has empty interior in \mathbb{R}^n , we can apply Theorem 1.8.11 of [2] and conclude that $\dim(B) < n$. On the other hand, Lemma 1.8.16 [2] guarantees that B has dimension $\dim(B) \geq n - 1$ (since B meets every non-trivial compact connected subset of \mathbb{R}^n). So, $\dim(B) = n - 1$. \square

REFERENCES

- [1] T. Banach, L. Zdomskyy, *Closed locally path-connected subspaces of finite-dimensional groups are locally compact*, Topology Proc. **36** (2010) 399–405.
- [2] R. Engelking, *Theory of Dimensions, Finite and Infinite*, Heldermann Verlag, 1995.
- [3] A. Gleason, *Arcs in locally compact groups*, Proc. Natl. Acad. Sci. USA. **38** (1950) 663–667.
- [4] A. Kechris, *Classical descriptive set theory*, Springer-Verlag, New York, 1995.
- [5] S. Maillot, *A non locally compact group of finite topological dimension?*, MathOverflow (<http://mathoverflow.net/questions/230878/a-non-locally-compact-group-of-finite-topological-dimension>)
- [6] D. Montgomery, *Theorems on the topological structure of locally compact groups*, Ann. of Math. **50** (1949) 570–580.

T.BANAKH: IVAN FRANKO NATIONAL UNIVERSITY OF LVIV
E-mail address: `t.o.banakh@gmail.com`

I.BANAKH: YA. PIDSTRYHACH INSTITUTE FOR APPLIED PROBLEMS OF MECHANICS AND MATHEMATICS, LVIV
E-mail address: `ibanakh@yahoo.com`

M.VOVK: LVIV POLYTECHNIC NATIONAL UNIVERSITY
E-mail address: `mira.i.kopych@gmail.com`