## NEW PROOFS OF BORWEIN-TYPE ALGORITHMS FOR PI

# JESÚS GUILLERA

ABSTRACT. We use a method of translation to recover Borweins' quadratic and quartic iterations. Then, by using the WZ-method, we obtain some initial values which lead to the limit  $1/\pi$ . We will not use the modular theory nor either the Gauss' formula that we used in [7]. Our proofs are short and self-contained.

# 1. Introduction

The complete elliptic integral K(x) is defined as follows:

$$K(x) = \int_0^{\frac{\pi}{2}} \frac{dt}{\sqrt{1 - x^2 \sin^2 t}},$$

and its expansion in power series is given by

$$\frac{2}{\pi}K(x) = {}_{2}F_{1}\left(\begin{vmatrix} \frac{1}{2}, & \frac{1}{2} \\ 1 & \end{vmatrix} x^{2}\right) = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_{k}^{2}}{\left(1\right)_{k}^{2}} x^{2k},$$

where  $(a)_k = a(a+1)\cdots(a+k-1)$  is the rising factorial, also known as Pochhammer symbol. Following the Indian mathematical genius Srinivasa Ramanujan, an identity of the form

$$\frac{K(\sqrt{1-v^2})}{K(v)} = n \frac{K(\sqrt{1-u^2})}{K(u)},$$

where n is any positive rational number, induces an algebraic transformation of degree n, which is of the following form [9] or [1, Ch. 29]:

$$K(u) = m(u, v) K(v), \quad P(u, v) = 0,$$

where m(u, v) (the multiplier) and P(u, v) (the modular equation) are algebraic functions in u and v. Many modular equations stated by Ramanujan in the variables  $\alpha = u^2$  and  $\beta = v^2$  are proved by Berndt in [2, Ch.19]. In addition, Ramanujan proved that if  $u_0$  is a solution of the equation  $P(u, \sqrt{1-u^2}) = 0$ , then there exist two algebraic numbers  $C_1$  and  $C_2$  such that

(1) 
$$\left( C_1 \frac{K^2}{\pi^2} + C_2 \frac{KK'}{\pi^2} \right) (u_0) = \frac{1}{\pi},$$

Observe that as K is a common factor, we can write (1) in the following way

$$\sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} u_0^{2k} \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} u_0^{2k} (A + Bk) = \frac{1}{\pi},$$

where A and B are algebraic numbers related to  $u_0$ ,  $C_1$  and  $C_2$ . Instead, in his proof Ramanujan relates (1) to a unique  ${}_3F_2$  hypergeometric sum of the following form:

$$\sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} \left[ 4u_0^2 (1 - u_0^2) \right]^k (a + bk) = \frac{1}{\pi},$$

where a, b are algebraic numbers. One example given by Ramanujan is

$$\sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} \frac{5+42k}{64^k} = \frac{16}{\pi}.$$

In the middle eighties, the brothers Jonathan and Peter Borwein observed that the modular equations given by Ramanujan could be connected to the number  $\pi$  via the *elliptic alpha function*, to construct extraordinary rapid algorithms to calculate  $\pi$  [3, p.170]. The point of view of the Borweins' is explained in [4], and their proofs require a good knowledge of the elliptic modular functions. In this paper we show a different point of view, and although the idea of our method looks very general, we only focus in a quadratic transformation, which will allow us to recover the quadratic and quartic algorithms for a few sets of initial values (got with the WZ-method). The main point is that we avoid completely the use of the modular stuff, and also the Gauss's formula used in [7].

# 2. Transformations and algorithms

Applying Zudilin's translation method [10] to a quadratic transformation we will derive Borweins' quadratic iteration.

# 2.1. A quadratic transformation. The following identity:

(2) 
$$\sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} x^{2k} = (1+t) \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} t^{2k}.$$

where

(3) 
$$t = \frac{1 - \sqrt{1 - x^2}}{1 + \sqrt{1 - x^2}},$$

and 0 < x < 1 is a known quadratic transformation [3, Thm 1.2 (b)]. Observe that the transformation is quadratic because  $t \sim 4^{-1} \cdot x^2$ . To prove (2) just expand in powers of x, observe that the coefficients up to  $x^2$  are equal and check that both sides of (2) satisfy the differential equation:

$$((x^2 - 1)\vartheta_x^2 + 2x^2\vartheta_x + x^2) f = 0, \qquad \vartheta_x = x\frac{d}{dx},$$

taking into account that the recurrence satisfied by the coefficient  $D_k = (1/2)_k^2/(1)_k^2$  of  $x^{2k}$  and  $t^{2k}$  is:

$$D_0 = 1,$$
  $D_{k+1} = \frac{(1+2k)^2}{4(1+k)^2}D_k,$ 

the following relations which come from (3):

$$\sqrt{1-x^2} = \frac{1-t}{1+t}$$
,  $x^2 = \frac{4t}{(1+t)^2}$ ,  $x\frac{dt}{dx} = \frac{x^2(1+t)^3}{2-2t} = \frac{2t(1+t)}{1-t}$ ,

and

$$\vartheta_x = \frac{x}{t} \frac{dt}{dx} \vartheta_t = \frac{2+2t}{1-t} \vartheta_t, \qquad \vartheta_x^2 = \frac{4(1+t)^2}{(1-t)^2} \vartheta_t^2 + \frac{8t(1+t)}{(1-t)^3} \vartheta_t.$$

2.2. A quadratic algorithm. We consider the following operator:

$$a + \frac{b}{2}\vartheta_x = a + b\frac{1+t}{1-t}\vartheta_t,$$

and apply it to both sides of (2):

$$\left(a + \frac{b}{2}\,\vartheta_x\right) \left\{ \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} x^{2k} \right\} = \left(a + b\,\frac{1+t}{1-t}\,\vartheta_t\right) \left\{ (1+t) \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} t^{2k} \right\}.$$

We have

(4) 
$$\sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} (a+bk) x^{2k} = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} \left(a(1+t) + b \frac{t(1+t)}{1-t} + 2b \frac{(1+t)^2}{1-t} k\right) t^{2k}$$

Multiplying (2) and (4), we obtain

(5) 
$$\sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_{k}^{2}}{(1)_{k}^{2}} x^{2k} \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_{k}^{2}}{(1)_{k}^{2}} (a+bk) x^{2k}$$

$$= \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_{k}^{2}}{(1)_{k}^{2}} t^{2k} \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_{k}^{2}}{(1)_{k}^{2}} \left(a(1+t)^{2} + b\frac{t(1+t)^{2}}{1-t} + 2b\frac{(1+t)^{3}}{1-t}k\right) t^{2k}$$

Let

$$A_n = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} d_n^{2k} \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} (a_n + b_n k) d_n^{2k}.$$

We define the sequences  $d_n$ ,  $a_n$ ,  $b_n$ ,  $A_n$  of initial values  $d_0$ ,  $a_0$ ,  $b_0$ ,  $A_0$ , implicitly by means of  $A_n = A_{n+1}$ , where  $A_{n+1}$  comes from  $A_n$  in the same way than the right hand side of (5) comes from its left side. Hence we have the following recurrences:

(6) 
$$d_{n+1} = \frac{1 - \sqrt{1 - d_n^2}}{1 + \sqrt{1 - d_n^2}}, \qquad b_{n+1} = 2b_n \frac{(1 + d_{n+1})^3}{1 - d_{n+1}},$$

(7) 
$$a_{n+1} = a_n (1 + d_{n+1})^2 + \frac{b_{n+1} d_{n+1}}{2(1 + d_{n+1})}.$$

As  $A_n$  is a constant sequence, it implies that  $\lim A_n = A_0$ , and as  $d_n^{2k} \to 0$  and  $b_n d_n^{2k} \to 0$  when  $n \to \infty$  and  $k \neq 0$ , we deduce that  $\lim A_n = \lim a_n$ . Hence  $\lim a_n = A_0$ .

**Remark.** Instead of dealing with algebraic transformations of products of two  ${}_{2}F_{1}$  hypergeometric sums, we can deal with transformations of single  ${}_{3}F_{2}$  hypergeometric series. In fact, there is an equivalence. For example, from the known transformation

$$\left(\sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} x^{2k}\right)^2 = \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^3}{(1)_k^3} \left[4x^2(1-x^2)\right]^k,$$

we can, via translation, rewrite our products of two  $_2F_1$  series as unique  $_3F_2$  series. When the sum is equal to  $1/\pi$  these single sums constitute one of the families of Ramanujan-type series for  $1/\pi$ .

#### 3. Other algorithms

Here we give some related algorithms:

3.1. A simplified version. From (6), we have

$$\frac{b_{n+1}}{1 - d_{n+1}^2} = 2b_n \frac{(1 + d_{n+1})^2}{(1 - d_{n+1})^2} = 2\frac{b_n}{1 - d_n^2}.$$

Then, writing  $c_n = b_n/(1-d_n^2)$ , we have  $c_{n+1} = 2c_n$ , which imply  $c_n = c_0 \cdot 2^n$ , and we get a simplified version of the algorithm of subsection 2.2. It is

(8) 
$$d_{n+1} = \frac{1 - \sqrt{1 - d_n^2}}{1 + \sqrt{1 - d_n^2}},$$

(9) 
$$a_{n+1} = a_n (1 + d_{n+1})^2 + c_0 2^n d_{n+1} (1 - d_{n+1}),$$

with  $\lim a_n = A_0$ .

3.2. An even more simplified version: Borweins' quadratic algorithm. As  $c_0 2^n d_{n+1} \sim a_{n+1} - a_n$ , we deduce that  $2^n d_{n+1}$  tends to 0 as  $n \to \infty$ . Then, writing  $a_n = r_n - c_0 2^{n-1} d_n^2$ , we arrive to the Borweins' quadratic iteration

(10) 
$$d_{n+1} = \frac{1 - \sqrt{1 - d_n^2}}{1 + \sqrt{1 - d_n^2}},$$

(11) 
$$r_{n+1} = r_n (1 + d_{n+1})^2 - c_0 2^n d_{n+1},$$

with  $\lim r_n = \lim a_n = A_0$ .

3.3. Borweins' quartic algorithm. Making  $s_n = \sqrt{d_{2n}}$ ,  $t_n = r_{2n}$ , we can arrive to the following Borweins' quartic algorithm (see [7] for the steps):

$$s_0 = \sqrt{d_0}, \quad t_0 = r_0,$$

$$s_{n+1} = \frac{1 - \sqrt[4]{1 - s_n^4}}{1 + \sqrt[4]{1 - s_n^4}},$$

$$t_{n+1} = (1 + s_{n+1})^4 t_n - c_0 2^{2n+1} s_{n+1} (1 + s_{n+1} + s_{n+1}^2),$$

with  $\lim t_n = A_0$ . Observe that it is a quartic algorithm because  $s_{n+1} \sim 8^{-1} \cdot s_n^4$ .

3.4. **Initial values.** The identities of this subsection provide initial values in order that the iterations tend to  $A_0 = 1/\pi$ . Below each identity we show the corresponding initial values in the algorithms. We will prove these evaluations in next subsection.

Identity 1.

$$\sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k^2}{(1)_k^2} (3 - 2\sqrt{2})^{2k} \sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k^2}{(1)_k^2} (3 - 2\sqrt{2})^{2k} \left[ 16(3\sqrt{2} - 4)k + 4(5\sqrt{2} - 7) \right] = \frac{1}{\pi},$$

and

$$d_0 = s_0^2 = 3 - 2\sqrt{2}, \quad b_0 = 48\sqrt{2} - 64, \quad a_0 = 20\sqrt{2} - 28, \quad c_0 = 4, \quad r_0 = t_0 = 6 - 4\sqrt{2}$$
Identity 2.

$$\sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} \frac{1}{2^k} \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} \frac{k}{2^k} = \frac{1}{\pi},$$

and

$$d_0 = s_0^2 = \frac{1}{\sqrt{2}}, \quad b_0 = 1, \quad a_0 = 0, \quad c_0 = 2, \quad r_0 = t_0 = \frac{1}{2}.$$

Identity 3.

$$\sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} (\sqrt{2} - 1)^{2k} \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} (\sqrt{2} - 1)^{2k} \left[ (8 - 4\sqrt{2})k + (3 - 2\sqrt{2}) \right] = \frac{1}{\pi},$$

and

$$d_0 = s_0^2 = \sqrt{2} - 1$$
,  $b_0 = 8 - 4\sqrt{2}$ ,  $a_0 = 3 - 2\sqrt{2}$ ,  $c_0 = 2\sqrt{2}$ ,  $r_0 = t_0 = \sqrt{2} - 1$ .

Identity 4.

$$\sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} \left(\frac{2-\sqrt{3}}{4}\right)^k \sum_{k=0}^{\infty} \frac{\left(\frac{1}{2}\right)_k^2}{(1)_k^2} \left(\frac{2-\sqrt{3}}{4}\right)^k \left[\left(\frac{3}{2}+\sqrt{3}\right)k+\frac{1}{4}\right] = \frac{1}{\pi},$$

and

$$d_0 = s_0^2 = \frac{\sqrt{6} - \sqrt{2}}{4}, \quad b_0 = \frac{3}{2} + \sqrt{3}, \quad a_0 = \frac{1}{4}, \quad c_0 = 2\sqrt{3}, \quad r_0 = t_0 = \frac{\sqrt{3} - 1}{2}.$$

Identity 5.

$$\sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k^2}{(1)_k^2} \left(2\sqrt{2} - 2\right)^k \sum_{k=0}^{\infty} \frac{(\frac{1}{2})_k^2}{(1)_k^2} \left(2\sqrt{2} - 2\right)^k \left[ (3\sqrt{2} - 4)k + \sqrt{2} - \frac{3}{2} \right] = \frac{1}{\pi},$$

and

$$d_0 = s_0^2 = \sqrt{2\sqrt{2} - 2}, \quad b_0 = 3\sqrt{2} - 4, \quad a_0 = \sqrt{2} - \frac{3}{2}, \quad c_0 = \sqrt{2}, \quad r_0 = t_0 = \frac{1}{2}.$$

3.5. Elementary proofs of identities 1 to 5. In this section we prove Identities 1 to 5. We avoid the modular theory and use only the WZ-method together with Carlson's Theorem [5]. In the right side of the formulas we use the general definition of the Pochhammer symbol, namely  $(v)_k = \Gamma(v+k)/\Gamma(v)$ .

WZ-proof of Identity 1. The following identity:

(12) 
$$\sum_{n=0}^{\infty} \frac{(\frac{1}{2} - 4k)_n (\frac{1}{2} - 2k)_n}{(1 + 2k)_n (1)_n} (3 - 2\sqrt{2})^{2n} = C_1 (3\sqrt{2} - 4)^{2k} \frac{2^{9k}}{3^{3k}} \frac{(1)_k (\frac{1}{2})_k}{(\frac{7}{12})_k (\frac{11}{12})_k},$$

valid for any complex number k is WZ-provable [8], due to the fact that it has a free parameter, and we can determine the constant  $C_1$  by taking k = 1/4. To discover other formulas like this, we need that the function H(n, k) inside the sum is such that the output of the Maple code

is 1. In this case we can easily guess the function at the right hand side of the equal, which is a function depending only in k. Many terminating versions of formulas of this kind have been discovered and proved automatically by a program written by D. Zeilberger [6]. It is good for our purposes that we found the following complement of (12):

$$\sum_{n=0}^{\infty} \frac{(\frac{1}{2} - 4k)_n (\frac{1}{2} - 2k)_n}{(1 + 2k)_n (1)_n} (3 - 2\sqrt{2})^{2n} \left[ 16(3\sqrt{2} - 4)n + 16(13 - 9\sqrt{2})k + 4(5\sqrt{2} - 7) \right]$$

$$= C_2 (3\sqrt{2} - 4)^{2k} \frac{2^{9k}}{3^{3k}} \frac{(1)_k (\frac{1}{2})_k}{(\frac{1}{12})_k (\frac{5}{12})_k},$$
(13)

which is WZ-provable as well. To discover this complement we have used the following Maple code:

```
with(SumTools[Hypergeometric]);
coK2:=coeff(Zeilberger(H(n,k)*(n+b*k+c),k,n,K)[1],K,2);
coes:=coeffs(coK2,k); solve({coes},{b,c});
```

where H(n, k) is the function inside the sum in (12) written in the notation of Maple. Again, we can determine  $C_2$  taking k = 1/4, and we have that  $C_1C_2 = 1/\pi$ . Multiplying the formulas (12) and (13), taking k = 0, and finally replacing n with k to agree with the notation of the other subsections, we arrive at Identity 1.

Below we give couples of identities which prove Identities 2 to 5.

WZ-proof of Identity 2.

$$\sum_{n=0}^{\infty} \frac{(\frac{1}{2} - 2k)_n (\frac{1}{2} + 2k)_n}{(1 + 4k)_n (1)_n} \left(\frac{1}{2}\right)^n = C_1 \left(\frac{16}{27}\right)^k \frac{(1)_k \left(\frac{1}{2}\right)_k}{\left(\frac{7}{12}\right)_k \left(\frac{11}{12}\right)_k},$$

$$\sum_{n=0}^{\infty} \frac{(\frac{1}{2} - 2k)_n (\frac{1}{2} + 2k)_n}{(1 + 4k)_n (1)_n} \left(\frac{1}{2}\right)^n (n + 4k) = C_2 \left(\frac{16}{27}\right)^k \frac{(1)_k \left(\frac{1}{2}\right)_k}{\left(\frac{1}{12}\right)_k \left(\frac{5}{12}\right)_k}.$$

WZ-proof of Identity 3.

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{2} - 2k\right)_n \left(\frac{1}{2}\right)_n}{(1+2k)_n (1)_n} (\sqrt{2} - 1)^{2n} = C_1 \frac{\left(1\right)_k \left(\frac{1}{2}\right)_k}{\left(\frac{5}{8}\right)_k \left(\frac{7}{8}\right)_k},$$

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{2} - 2k\right)_n \left(\frac{1}{2}\right)_n}{(1+2k)_n (1)_n} (\sqrt{2} - 1)^{2n} \left[ (8 - 4\sqrt{2})n + (8\sqrt{2} - 8)k + (3 - 2\sqrt{2}) \right] = C_2 \frac{\left(1\right)_k \left(\frac{1}{2}\right)_k}{\left(\frac{1}{8}\right)_k \left(\frac{3}{8}\right)_k}.$$

WZ-proof of Identity 4.

$$\sum_{n=0}^{\infty} \frac{(\frac{1}{2} - k)_n (\frac{1}{2} + 3k)_n}{(1+k)_n (1)_n} \left(\frac{2 - \sqrt{3}}{4}\right)^n = C_1 \left(\frac{4}{3\sqrt{3}}\right)^k \frac{(1)_k}{\left(\frac{5}{6}\right)_k},$$

$$\sum_{n=0}^{\infty} \frac{(\frac{1}{2} - k)_n (\frac{1}{2} + 3k)_n}{(1+k)_n (1)_n} \left(\frac{2 - \sqrt{3}}{4}\right)^n \left[\left(\frac{3}{2} + \sqrt{3}\right)n + \frac{3}{2}k + \frac{1}{4}\right] = C_2 \left(\frac{4}{3\sqrt{3}}\right)^k \frac{(1)_k}{\left(\frac{1}{6}\right)_k}.$$

WZ-proof of Identity 5.

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{2} + 4k\right)_n \left(\frac{1}{2} + 2k\right)_n}{(1+k)_n (1)_n} (2\sqrt{2} - 2)^n = C_1 \left(3 + 2\sqrt{2}\right)^{2k} \frac{\left(1\right)_k \left(\frac{1}{2}\right)_k}{\left(\frac{5}{8}\right)_k \left(\frac{7}{8}\right)_k},$$

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{2} + 4k\right)_n \left(\frac{1}{2} + 2k\right)_n}{(1+k)_n (1)_n} (2\sqrt{2} - 2)^n \left[ (3\sqrt{2} - 4)n + (8\sqrt{2} - 12)k + \left(\sqrt{2} - \frac{3}{2}\right) \right]$$

$$= C_2 \left(3 + 2\sqrt{2}\right)^{2k} \frac{\left(1\right)_k \left(\frac{1}{2}\right)_k}{\left(\frac{1}{8}\right)_k \left(\frac{3}{8}\right)_k}.$$

Although with the WZ-method we have only obtained a few sets of initial values, it supposes no disadvantage with respect to the modular theory because the interesting fact is to have one set of simple initial values.

### References

- [1] L. Berggren, J. Borwein and P. Borwein, *Pi: A Source Book*, (Springer-Verlag, New York, 1997, 2000).
- [2] B. Berndt, Ramanujan's Notebooks, Part III, (Springer-Verlag, New York, 1991).
- [3] J. Borwein and P. Borwein, Pi and the AGM: A Study in Analytic Number Theory and Computational Complexity, Canad. Math. Soc. Series Monographs Advanced Texts, (John Wiley, New York, 1987).
- [4] J. Borwein and P. Borwein, Ramanujan, Modular Equations, and Approximations to Pi or How to Compute One Billion Digits of Pi, Amer. Math. Monthly, 96 (1989), 201-219.
- [5] S. EKHAD AND D. ZEILBERGER, A WZ proof of Ramanujan's formula for  $\pi$ , Geometry, Analysis and Mechanics, J. M. Rassias (ed.), World Scientific, Singapore 1994.
- [6] S. EKHAD, Forty "Strange" Computer-Discovered and Computer Proved (of course!) Hypergeometric Series Evaluations. (available on Zeilberger's web-site).
- [7] J. GUILLERA, Easy Proofs of Some Borwein Algorithms for  $\pi$ , Amer. Math. Monthly, 115 (2008), 850-854.
- [8] M. Petkovšek, H. Wilf and D. Zeilberger, A=B, (A K Peters, Ltd., Wellesley, Massachusetts, 1996).

- [9] S. RAMANUJAN, Modular equations and approximations to  $\pi$ , Quarterly Journal of Mathematics 45 (1914), 350-372.
- [10] W. Zudilin, Lost in translation, in *Advances in Combinatorics*, Waterloo Workshop in Computer Algebra W80 (May 26–29, 2011), I. Kotsireas and E.V. Zima (eds.) (2013), Springer, 287–293.

Department of Mathematics, University of Zaragoza, 50009 Zaragoza, SPAIN  $E\text{-}mail\ address:}$  jguillera@gmail.com