

On Dirac's Conjecture

Zh.G. Nikoghosyan

February 12, 2022

Abstract

Let G be a 2-connected graph, l be the length of a longest path in G and c be the circumference - the length of a longest cycle in G . In 1952, Dirac proved that $c > \sqrt{2l}$ and conjectured that $c \geq 2\sqrt{l}$. In this paper we present more general sharp bounds in terms of l and the length m of a vine on a longest path in G including Dirac's conjecture as a corollary: if $c = m + y + 2$ (generally, $c \geq m + y + 2$) for some integer $y \geq 0$, then $c \geq \sqrt{4l + (y + 1)^2}$ if m is odd; and $c \geq \sqrt{4l + (y + 1)^2} - 1$ if m is even.

Key words: longest cycle, longest path, circumference.

1 Introduction

We consider only undirected graphs with no loops or multiple edges. Let G be a 2-connected graph. We use c and l to denote the circumference (the length of a longest cycle) and the length of a longest path of G . A good reference for any undefined terms is [1].

In 1952, Dirac [3] proved the following.

Theorem A [3]. In every 2-connected graph, $c > \sqrt{2l}$.

In the same paper [3], Dirac conjectured a sharp version of Theorem A.

Conjecture A [3]. In every 2-connected graph, $c \geq 2\sqrt{l}$.

In this paper we present more general sharp bounds in terms of l and the length of a vine on a longest path of G including Dirac's conjecture as a corollary. In order to formulate this result, we need some additional definitions and notations.

The set of vertices of a graph G is denoted by $V(G)$ and the set of edges by $E(G)$. If Q is a path or a cycle, then the length of Q , denoted by $l(Q)$, is $|E(Q)|$ - the number of edges in Q . We write a cycle Q with a given orientation by \vec{Q} . For $x, y \in V(Q)$, we denote by $x\vec{Q}y$ the subpath of Q in the chosen direction from x to y . We use $P = x\vec{P}y$ to denote a path with end vertices x and y in

the direction from x to y . We say that vertex z_1 precedes vertex z_2 on \vec{Q} if z_1, z_2 occur on \vec{Q} in this order, and indicate this relationship by $z_1 \prec z_2$. We will write $z_1 \preceq z_2$ when either $z_1 = z_2$ or $z_1 \prec z_2$.

Let $P = x\vec{P}y$ be a path. A vine of length m on P is a set

$$\{L_i = x_i\vec{L}_iy_i : 1 \leq i \leq m\}$$

of internally-disjoint paths such that

- (a) $V(L_i) \cap V(P) = \{x_i, y_i\}$ ($i = 1, \dots, m$),
- (b) $x = x_1 \prec x_2 \prec y_1 \preceq x_3 \prec y_2 \preceq x_4 \prec \dots \preceq x_m \prec y_{m-1} \prec y_m = y$ on P .

Theorem 1. Let G be a 2-connected graph. If $\{L_1, L_2, \dots, L_m\}$ is a vine on a longest path of G and $c = m + y + 2$ for some integer $y \geq 0$, then

$$c \geq \begin{cases} \sqrt{4l + (y+1)^2} & \text{when } m \equiv 1 \pmod{2}, \\ \sqrt{4l + (y+1)^2 - 1} & \text{when } m \equiv 0 \pmod{2}. \end{cases}$$

The following lemma guarantees the existence of at least one vine on a longest path in a 2-connected graph.

The Vine Lemma [2]. Let G be a k -connected graph and P a path in G . Then there are $k - 1$ pairwise-disjoint vines on P .

2 The proof of Theorem 1

Let $P = x\vec{P}y$ be a longest path in G and let

$$\{L_i = x_i\vec{L}_iy_i : 1 \leq i \leq m\}$$

be a vine of length m on P . Put

$$\begin{aligned} L_i &= x_i\vec{L}_iy_i \quad (i = 1, \dots, m), \quad A_1 = x_1\vec{P}x_2, \quad A_m = y_{m-1}\vec{P}y_m, \\ A_i &= y_{i-1}\vec{P}x_{i+1} \quad (i = 2, 3, \dots, m-1), \quad B_i = x_{i+1}\vec{P}y_i \quad (i = 1, \dots, m-1), \\ l(A_i) &= a_i \quad (i = 1, \dots, m), \quad l(B_i) = b_i \quad (i = 1, \dots, m-1). \end{aligned}$$

By combining appropriate L_i, A_i, B_i , we can form the following cycles:

$$Q_0 = \bigcup_{i=1}^m (A_i \cup L_i),$$

$$Q_j = \bigcup_{i=j+1}^{m-j} (A_i \cup L_i) \cup B_j \cup B_{m-j} \quad (j = 1, 2, \dots, \lfloor (m-1)/2 \rfloor).$$

Since $l(L_i) \geq 1$ ($i = 1, 2, \dots, m$) and $a_1 \geq 1, a_m \geq 1$, we have

$$c \geq l(Q_0) = \sum_{i=1}^m l(L_i) + a_1 + a_m + \sum_{i=2}^{m-1} a_i \geq m + 2,$$

Let $c = m + y + 2$, where $y \geq 0$. If $a_1 + a_m \geq y + 3 - \sum_{i=2}^{m-1} a_i$, then

$$\begin{aligned} l(Q_0) &= \sum_{i=1}^m l(L_i) + (a_1 + a_m) + \sum_{i=2}^{m-1} a_i \\ &\geq m + \left(y + 3 - \sum_{i=2}^{m-1} a_i \right) + \sum_{i=2}^{m-1} a_i = m + y + 3 > c, \end{aligned}$$

a contradiction. Hence,

$$a_1 + a_m \leq y + 2 - \sum_{i=2}^{m-1} a_i. \quad (1)$$

Next, if $b_1 + b_{m-1} \geq y + 5 - \sum_{i=2}^{m-1} a_i$, then

$$\begin{aligned} l(Q_1) &= \sum_{i=2}^{m-1} l(L_i) + (b_1 + b_{m-1}) + \sum_{i=2}^{m-1} a_i \\ &\geq (m-2) + \left(y + 5 - \sum_{i=2}^{m-1} a_i \right) + \sum_{i=2}^{m-1} a_i = m + y + 3 > c, \end{aligned}$$

a contradiction. Hence,

$$b_1 + b_{m-1} \leq y + 4 - \sum_{i=2}^{m-1} a_i \leq y + 4.$$

Furthermore, if $b_2 + b_{m-2} \geq y + 7 - \sum_{i=3}^{m-2} a_i$, then

$$\begin{aligned} l(Q_2) &= \sum_{i=3}^{m-2} l(L_i) + (b_2 + b_{m-2}) + \sum_{i=3}^{m-2} a_i \\ &\geq (m-4) + \left(y + 7 - \sum_{i=3}^{m-2} a_i \right) + \sum_{i=3}^{m-2} a_i = m + y + 3 > c, \end{aligned}$$

again a contradiction. Hence

$$b_2 + b_{m-2} \leq y + 6 - \sum_{i=3}^{m-2} a_i \leq y + 6.$$

Analogously,

$$\begin{aligned} b_j + b_{m-j} &\leq y + 2(j+1) - \sum_{i=j+1}^{m-j} a_i \\ &\leq y + 2(j+1) \quad (j = 1, 2, \dots, \lfloor (m-1)/2 \rfloor). \end{aligned} \quad (2)$$

Case 1. $m = 2k + 1$ for some integer $k \geq 0$.

By (2),

$$\begin{aligned} \sum_{i=1}^{m-1} b_i &= \sum_{j=1}^{(m-1)/2} (b_j + b_{m-j}) \leq \sum_{j=1}^{(m-1)/2} (y + 2(j+1)) \\ &= \frac{m-1}{2}(y+2) + 2 \sum_{j=1}^{(m-1)/2} (j+1) = \frac{m-1}{2}(y+2) + \frac{(m-1)(m+1)}{4}. \end{aligned}$$

Then

$$\begin{aligned} l &= (a_1 + a_m) + \sum_{i=2}^{m-1} a_i + \sum_{i=1}^{m-1} b_i \\ &\leq \left(y + 2 - \sum_{i=2}^{m-1} a_i \right) + \sum_{i=2}^{m-1} a_i + \frac{m-1}{2}(y+2) + \frac{(m-1)(m+1)}{4} \\ &= (y+2) \frac{m+1}{2} + \frac{(m-1)(m+1)}{4} \\ &= (y+2) \frac{c-y-1}{2} + \frac{(c-y-3)(c-y-1)}{4} = \frac{c^2 - (y+1)^2}{4}, \end{aligned}$$

implying that $c \geq \sqrt{4l + (y+1)^2}$.

Case 2. $m = 2k$ for some integer $k \geq 1$.

As in Case 1,

$$\begin{aligned} \sum_{i=1}^{m-1} b_i &= \sum_{j=1}^{(m-2)/2} (b_j + b_{m-j}) + b_{m/2} \\ &\leq (y+2) \frac{m-2}{2} + \frac{(m-2)m}{4} + b_{m/2}. \end{aligned}$$

For the cycle

$$Q^* = B_{m/2} \cup B_{(m-2)/2} \cup A_{m/2} \cup L_{m/2}$$

we have

$$l(Q^*) = b_{m/2} + b_{(m-2)/2} + a_{m/2} + l(L_{m/2}) \leq c = y + m + 2,$$

implying that

$$b_{m/2} + b_{(m-2)/2} \leq y + m + 1.$$

Then it easy to see that

$$\sum_{i=1}^{m-1} b_i \leq (y+2)\frac{m-2}{2} + \frac{(m-2)m}{4} + \frac{y}{2} + \frac{m+2}{2}.$$

Furthermore, we have

$$\begin{aligned} l &= (a_1 + a_m) + \sum_{i=2}^{m-1} a_i + \sum_{i=1}^{m-1} b_i \\ &\leq (y+2) + (y+2)\frac{m-2}{2} + \frac{(m-2)m}{4} + \frac{y}{2} + \frac{m+2}{2} \\ &= \frac{(m+2)^2 + 2y(m+1)}{4} = \frac{(c-y)^2 + 2y(c-y-1)}{4} = \frac{c^2 - y^2 - 2y}{4}, \end{aligned}$$

implying that $c \geq \sqrt{4l + (y+1)^2 - 1}$. Theorem is proved. \blacksquare

Let $P = x\vec{P}y$ be a path and let

$$\{L_i = x_i\vec{L}_iy_i : 1 \leq i \leq m\}$$

be a vine of length m on P . Put

$$\begin{aligned} L_i &= x_i\vec{L}_iy_i \quad (i = 1, \dots, m), \quad A_1 = x_1\vec{P}x_2, \quad A_m = y_{m-1}\vec{P}y_m, \\ A_i &= y_{i-1}\vec{P}x_{i+1} \quad (i = 2, 3, \dots, m-1), \quad B_i = x_{i+1}\vec{P}y_i \quad (i = 1, \dots, m-1), \\ l(A_i) &= a_i \quad (i = 1, \dots, m), \quad l(B_i) = b_i \quad (i = 1, \dots, m-1). \end{aligned}$$

Let $y \geq 0$ by an integer and

$$\begin{aligned} a_1 = a_m &= \frac{y}{2} + 1, \quad a_2 = a_3 = \dots = a_{m-1} = 0, \\ b_i = b_{m-i} &= \frac{y}{2} + i + 1 \quad (i = 1, 2, \dots, \lfloor (m-1)/2 \rfloor). \end{aligned}$$

If m is odd, then $c = m + y + 2$ and $c = \sqrt{4l + (y+1)^2}$.

If m is even, we put $b_{m/2} = \frac{y}{2} + \frac{m+2}{2}$, implying that $c = m + y + 2$ and $c = \sqrt{4l + (y+1)^2 - 1}$. Thus, the bounds in Theorem 1 are best possible.

References

- [1] J.A. Bondy and U.S.R. Murty, Graph Theory with Applications, Macmillan, London and Elsevier, New York (1976).
- [2] J.A. Bondy, Basic Graph Theory: Paths and circuits. Handbook of combinatorics, Vol. 1,2, Elsevier, Amsterdam (1990).

- [3] G.A. Dirac, Some theorems on abstract graphs, Proc. London, Math. Soc. 2 (1952) 69-81.

Institute for Informatics and Automation Problems
National Academy of Sciences
P. Sevak 1, Yerevan 0014, Armenia
E-mail: zhora@ipia.sci.am