# Some results on chromatic number as a function of triangle count

David G. Harris\*

February 21, 2022

### Abstract

A variety of powerful results have been shown for the chromatic number of triangle-free graphs. Three noteworthy bounds are in terms of the number of vertices, edges, and maximum degree given by Nilli (2000), Gimbel & Thomassen (2000), and Johansson. There have been comparatively fewer works extending these types of bounds to graphs with relatively few, but non-zero, number of triangles. One noteworthy exception is a result of Alon et. al (1999) bounding the chromatic number for graphs with low degree and few triangles per vertex; this bound is nearly the same as for triangle-free graphs. This type of parametrization is much less rigid, and has appeared in dozens of combinatorial constructions.

In this paper, we show a similar type of result for  $\chi(G)$  as a function of the number of vertices n, the number of edges m, as well as the triangle count (both local and global measures). Our results smoothly interpolate between the generic bounds true for all graphs and bounds for triangle-free graphs. Our results are tight for most of these cases.

### 1 Introduction

In this paper, we will show bounds on the chromatic number of undirected graphs G = (V, E) which have relatively few triangles (a triangle is a triple of vertices  $x, y, z \in V$  where all three edges (x, y), (x, z), (y, z)are present in G).

A variety of powerful results have been shown for the chromatic number of triangle-free graphs. Two noteworthy bounds, which we explore further in this paper, are in terms of the number of vertices and edges.

**Theorem 1.1** ([1]). Suppose G is triangle-free and has n vertices. Then  $\chi(G) \leq O\left(\sqrt{\frac{n}{\log n}}\right)$ .

**Theorem 1.2** ([8], [3]). Suppose G is triangle-free and has m edges. Then  $\chi(G) \leq O\left(\frac{m^{1/3}}{\log^{2/3} m}\right)$ .

Another powerful bound for triangle-free graphs can be given in terms of the maximum degree:<sup>1</sup>

**Theorem 1.3.** Suppose G is triangle-free and has maximum degree d. Then  $\chi(G) \leq O\left(\frac{d}{\log d}\right)$ 

These bounds are much smaller than would be possible for a generic graph (with no restriction on the number of triangles); in those cases one can only show the bounds

$$\chi(G) \le \min(n, \sqrt{m}, d)$$

The requirement that a graph has no triangles is quite rigid. There have been comparatively fewer works which relax the triangle-free condition to allow a small number of triangles. One noteworthy exception to this is the result of [2]:

**Theorem 1.4** ([2]). Suppose G is a graph with maximum degree d and each vertex incident on at most y triangles, where  $1 \le y \le d^2/2$ . Then

$$\chi(G) \le O(\frac{d}{\ln(d^2/y)})$$

<sup>\*</sup>Department of Computer Science, University of Maryland, College Park, MD 20742. Email: davidgharris290gmail.com

<sup>&</sup>lt;sup>1</sup>Theorem 1.3 is attributed to Johansson, as attributed by [6]. See [4] and [9] for more recent proofs, which also give bounds on the constant term.

Thus, for instance, if  $y < d^{2-\Omega(1)}$ , then  $\chi(G)$  is roughly the same as if G had no triangles at all. The result of [2] has been used in dozens of combinatorial constructions. Roughly speaking, if G is produced in a somewhat "random" or "generic" way, then G will have relatively few triangles; only a few extremal cases (such as a graph containing a d-clique) give the full triangle count.

In this paper, we show upper bounds for  $\chi(G)$  as a function of the number of vertices n, the number of edges m, as well as the triangle count. Our results smoothly interpolate between the generic bounds true for all graphs, and the triangle-free bounds on Theorems 1.1,1.2. We will see here as well that graphs with few triangles have nearly the same behavior as graphs with no triangles.

**Definition 1.5.** We say that a vertex  $v \in V$  has local triangle count y, if there are y pairs of vertices  $u, w \in V$  such that  $(v, u), (v, w), (u, w) \in E$ . We say that G has local triangle bound y if every vertex has local triangle count at most y.

We shall show the following bounds on the chromatic number:

**Theorem 1.6.** Suppose a graph G has m edges, n vertices, t triangles, and local triangle bound y. Then

$$\chi(G) \le \min(a_1, a_2, a_3, a_4, a_5, a_6)$$

where

$$a_{1} = O\left(\sqrt{\frac{n}{\log n}} + \frac{n^{1/3}y^{1/3}}{\log^{2/3}(n^{2}/y)}\right) \qquad a_{2} = O\left(\frac{m^{1/3}}{\log^{2/3}m} + \frac{m^{1/4}y^{1/4}}{\log^{3/4}(m/y)}\right)$$
$$a_{3} = O\left(\sqrt{\frac{n}{\log n}} + \frac{t^{1/3}\log\log(t^{2}/y^{3})}{\log^{2/3}(t^{2}/y^{3})}\right) \qquad a_{4} = O\left(\frac{m^{1/3}}{\log^{2/3}m} + \frac{t^{1/3}\log\log(t^{2}/y^{3})}{\log^{2/3}(t^{2}/y^{3})}\right)$$
$$a_{5} = \sqrt{\frac{n}{\log n}} + (6^{1/3} + o(1))t^{1/3} \qquad a_{6} = \frac{m^{1/3}}{\log^{2/3}m} + (6^{1/3} + o(1))t^{1/3}$$

Furthermore, colorings with the indicated bounds can be constructed by a randomized polynomial-time algorithm.

The bounds  $a_1, a_2, a_5, a_6$  are tight up to second-order terms for all admissible values of m, n, y, t; the bounds  $a_3, a_4$  are tight up to factors of  $\log \log(t^2/y^3)$ .

### 1.1 Note on constructive algorithms

All our bounds on chromatic number can be realized constructively; that is, there are randomized algorithms which construct such colorings in expected polynomial time. Our proofs will use a few previous results on chromatic number, which did not have constructive algorithms stated directly. For completeness, we state these here with a proof sketch.

**Theorem 1.7** ([2]). Suppose G has maximum degree d and local triangle bound y, where  $1 \le y \le d^2/2$ . Then there is a randomized polynomial-time algorithm producing a coloring with  $O(\frac{d}{\ln(d^2/y)})$  colors.

*Proof.* The original proof of Theorem 1.4 uses a result on constructing coloring with  $O(d/\log d)$  colors for triangle-free graphs. This has been made into a constructive algorithm in [9]. The other main probabilistic tool used in Theorem 1.4 is the Lovász Local Lemma; the algorithm of Moser & Tardos [7] makes this constructive in a fairly straightforward way.

Theorem 1.4 was extended to list-chromatic number by Vu:

**Theorem 1.8** ([10]). Suppose G has maximum degree d and local triangle bound y, where  $1 \le y \le d^2/2$ . If each vertex has a palette of size  $\frac{cd}{\log(d^2/y)}$ , where c is a universal constant, then there is a randomized polynomial-time algorithm to producing a list-coloring.

**Theorem 1.9** (Turán's Theorem). Suppose G is a graph with n vertices and m edges. Then G has an independent set of size at least  $\frac{n}{2m/n+1}$ ; furthermore, such an independent set can be constructed in randomized polynomial time.

#### 1.2Notation

We use ln to refer to the natural logarithm, while the expression log always refers to the truncated logarithm, i.e.

$$\log(x) = \begin{cases} \ln(x) & x > e \\ 1 & x \le e \end{cases}$$

We write  $f \leq g$  to mean that f = O(g). We write  $f \approx g$  if  $f = \Theta(g)$ .

Given a graph G on a vertex set V and a subset  $B \subseteq V$ , we use  $\chi(B)$  as a shorthand for  $\chi(G[B])$ , where G[B] denotes the subgraph induced on B. We let N(v) denote the neighborhood of v, i.e. the set of vertices w such that  $(w, v) \in E$ . The degree of v is |N(v)|.

#### $\mathbf{2}$ Bounds in terms of n

We begin by showing a bound in terms of the vertex count and local triangle bound. This generalizes a result of [1], which show a similar bound for chromatic number as a function of the vertex count for triangle-free graphs.

**Theorem 2.1.** Suppose G has n vertices and local triangle bound y. Then

$$\chi(G) \lesssim \sqrt{\frac{n}{\log n}} + \frac{n^{1/3}y^{1/3}}{\log^{2/3}(n^2/y)}$$

Further, such a coloring can be produced in expected polynomial time.<sup>2</sup>

*Proof.* Let  $f = \log(n^2/y)$ . Simple analysis shows that it suffices to show that  $\chi(G) \lesssim \sqrt{n/\log n}$  for  $y \leq \sqrt{n/\log n}$  $\sqrt{n\log n}$  and  $\chi(G) \lesssim \frac{(ny)^{1/3}}{f^{2/3}}$  for  $y \ge \sqrt{n\log n}$ . Our plan is to repeatedly remove independent sets from G, as long as its maximum degree exceeds d, where d is some parameter to be chosen.

Initially let  $G_0 = G$ . Now, repeat the following for i = 0, 1, 2, ..., k: if all the vertices of  $G_i$  have degree  $\leq d$ , then abort the process and color the residual graph  $G_i$  using Theorem 1.7; this requires  $O(\frac{d}{\log(d^2/y)})$ colors.

Otherwise, select some vertex  $v_i$  which has degree > d in  $G_i$ , and let  $I_i$  be a maximum independent set

of the neighborhood of v in  $G_i$ . Color all the vertices of  $I_i$  with a new color, and let  $G_{i+1} = G_i - I_i$ . The total number of colors used during this process is at most  $O(\frac{d}{\log(d^2/y)} + k)$ . To bound k, observe the following. Each vertex  $v_i$  has degree > d in  $G_i$ , so the neighborhood of  $v_i$  contains > d vertices. By hypothesis, the neighborhood of  $v_i$  contains  $\leq y$  edges. Hence, by Theorem 1.9, one can construct an independent set of size  $\geq \frac{d}{y/d+1}$ . So we have  $|I_i| \geq \frac{d}{y/d+1}$  for  $i = 1, \ldots, k$ . As  $\sum |I_i| \leq n$ , it follows that  $k \le \frac{n(y/d+1)}{d}.$ 

So, overall we have  $\chi \lesssim \frac{d}{\log(d^2/y)} + \frac{n(y/d+1)}{d}$ . It remains now to select the parameter d. First let us suppose that  $y \leq \sqrt{n \log n}$ . In this case, take  $d = \sqrt{n \log n}$ . Then

$$\chi(G) \lesssim \frac{\sqrt{n}(1 + \frac{y}{\sqrt{n\log n}})}{\sqrt{\log n}} + \frac{\sqrt{n\log n}}{\log \frac{n\log n}{y}} \lesssim \sqrt{\frac{n}{\log n}}$$

Next, suppose that  $y > \sqrt{n \log n}$ . In this case, take  $d = (nyf)^{1/3}$ . Then

$$\begin{split} \chi(G) &\lesssim \frac{n(1 + \frac{y}{(nyf)^{1/3}})}{(nyf)^{1/3}} + \frac{(nyf)^{1/3}}{\log\frac{(nyf)^{2/3}}{y}} \\ &\leq \frac{2n^{1/3}y^{1/3}}{f^{2/3}} + \frac{(nyf)^{1/3}}{\log(f^{1/3}e^f)} \quad \text{as } y > \sqrt{nf} \\ &\lesssim \frac{n^{1/3}y^{1/3}}{f^{2/3}} \end{split}$$

 $<sup>^{2}</sup>$ For the remainder of this paper, we will not explicitly state that the colorings can be produced in expected polynomial time, if it is clear from context.

Next, we will show bounds on  $\chi(G)$  given information both about the global triangle count and the local triangle bound. We begin with a useful lemma.

**Lemma 2.2.** Suppose that a graph G has local triangle bound y, and there is a partition of the vertices  $V = V_1 \sqcup V_2 \sqcup \cdots \sqcup V_k$  such that, for all  $1 \le i \le j \le k$ , every  $v \in A_i$  satisfies

$$|N(v) \cap A_i| \le dx^{x-j}$$

for some parameters  $d, x \ge 1$ .

Then

$$\chi(G) \lesssim \frac{d\lceil \log_x \log(d^2/y) \rceil}{\log(d^2/y)}$$

Furthermore, such a coloring can be found in polynomial time. In particular, if  $x \ge 1 + \Omega(1)$ , then

$$\chi(G) \lesssim \frac{d \log \log(d^2/y)}{\log(d^2/y)}$$

*Proof.* Define  $f = \log(d^2/y)$ . We will first show that this result hold for  $x \ge f$ . In this case, we want to show that  $\chi(G) \le d/f$ .

We will allocate a palette of size cd/f to each vertex v, where c is a constant to be determined. We proceed for j = k, k - 1, ..., 1, attempting to list-color  $G[A_j]$ .

When we reach a given value j, note that each  $v \in A_j$  began to cd/f colors in its palette. Only the neighbors of v in  $A_{j+1}, A_{j+2}, \ldots, A_k$  have been colored so far; the total number of such neighbors is at most

$$dx^{-1} + dx^{-2} + \dots \le d(\frac{1}{f} + \frac{1}{f^2} + \frac{1}{f^3}) \le 2d/f$$

Thus, the vertex v remains with cd/f - 2d/f = (c-2)d/f colors in its palette. If c is a sufficiently large constant, then we may apply Theorem 1.8 to extend the coloring to v.

Now that we have shown this result holds for x = f, let us show that it holds for an arbitrary value x < f. Set  $s = \lceil \log_x f \rceil$  and for each i = 1, ..., s define  $B_i = A_i \cup A_{i+s} \cup A_{i+2s} \cup ...$  Observe that each  $G[B_i]$  satisfies the hypotheses of this Lemma with the vertex-partition  $A_i, A_{i+s}, A_{i+2s}, ...$  and with parameters  $d' = d, x' = x^s \ge f$ . Thus  $\chi(B_i) \le d/f$  and so

$$\chi(G) \le \chi(B_1) + \dots + \chi(B_s) \lesssim sd/f = \lceil \log_x \log(d^2/y) \rceil \times \frac{d}{\log(d^2/y)}$$

**Proposition 2.3.** Suppose G has n vertices, t triangles, and local triangle bound y. Then

$$\chi(G) \lesssim \frac{t^{1/3} \log \log(t^2/y^3)}{\log^{2/3}(t^2/y^3)} + \sqrt{n}$$

*Proof.* Let  $A_i$  denote the set of vertices in G with between  $2^i$  and  $2^{i+1}$  triangles and let d be a parameter which we will specify shortly.

Let  $k = \log_2 d$ . Suppose that for  $i \ge k$  there is a vertex  $v \in A_i$  with more than  $2^{(i-j)/2}d$  neighbors in  $A_j$  where  $j \ge i$ . Then apply Theorem 1.9 to  $N(v) \cap A_j$ . This yields an independent set I of size  $|I| \ge \frac{2^{(i-j)/2}d}{1+\frac{2^{i+2}}{2^{(i-j)/2}d}} \gtrsim 2^{-j}d^2$ . Assign each vertex in I to one new color and remove I from the graph. Since each  $w \in I$  is incident on at least  $2^j$  triangles, this removes  $\Omega(d^2)$  triangles. Note that this process of removing

 $w \in I$  is incident on at least 2<sup>o</sup> triangles, this removes  $\Omega(a^{\circ})$  triangles. Note that this process of removing vertices may cause the membership of the sets  $A_l$  to change. Overall the total number of new colors that can be produced this way is  $O(t/d^2)$ .

Next, apply Lemma 2.2 to color  $G[A_{k+1} \cup A_{k+2} \cup \dots]$ ; this satisfies the requirements of that theorem

with the parameter  $x = \sqrt{2}$ . So it can be colored using  $O(\frac{d \log \log(d^2/y)}{\log(d^2/y)})$  colors. Next, apply Theorem 2.1 to color  $G[A_1 \cup \cdots \cup A_k]$ . This graph has maximum of  $2^{k+1} \le O(d)$  triangles per vertex, hence it can be colored using  $O(\sqrt{\frac{n}{\log n} + \frac{(nd)^{1/3}}{\log^{2/3}(n^2/d)}})$  colors.

Putting all these terms together,

$$\chi(G) \lesssim \frac{d \log \log(d^2/y)}{\log(d^2/y)} + \sqrt{\frac{n}{\log n}} + \frac{(nd)^{1/3}}{\log^{2/3}(n^2/d)} + \frac{t}{d^2}$$

Now set  $f = \log(t^2/y^3)$  and  $d = (ft)^{1/3} + \sqrt{n}$ . Observe that  $d^2/y \ge \frac{(t\log(t^2/y^3))^{2/3}}{y} \approx (t^{2/3}/y)\log(t^{2/3}/y)$ . Thus  $\log(d^2/y) \gtrsim f$ . Similarly,  $\log(n^2/d) \gtrsim \log n$ .

Then we compute these terms in turn.

$$\frac{t}{d^2} = \frac{t^{1/3}}{(\sqrt{n} + (t\log(t^2/y^3))^{1/3})^2} \lesssim \frac{t^{1/3}}{\log^{2/3}(t^2/y^3)}$$
$$\frac{(nd)^{1/3}}{\log^{2/3}(n^2/d)} \lesssim \frac{\sqrt{n}}{\log^{2/3}n} + \frac{n^{1/3}t^{1/9}f^{1/9}}{\log^{2/3}n}$$
$$\frac{d\log\log(d^2/y)}{\log(d^2/y)} \lesssim \frac{d\log f}{f} \lesssim \frac{t^{1/3}\log f}{f^{2/3}} + \frac{\sqrt{n}\log f}{f}$$

We see that

$$\chi(G) \lesssim \frac{t^{1/3} \log f}{f^{2/3}} + \sqrt{n} + \frac{n^{1/3} t^{1/9} f^{1/9}}{\log^{2/3} n}$$

Observe that when  $t \leq n^{3/2}\sqrt{\log n}$ , we have  $\frac{n^{1/3}t^{1/9}f^{1/9}}{\log^{2/3}n} \lesssim \sqrt{\frac{n}{\log n}}$ . When  $t > n^{3/2}\sqrt{\log n}$ , then

$$\frac{n^{1/3}t^{1/9}f^{1/9}}{\log^{2/3}n} \le \frac{t^{1/3}}{f^{2/3}} \times \frac{f^{7/9}n^{1/3}}{t^{2/9}\log^{2/3}n} \le \frac{t^{1/3}}{f^{2/3}} \times (f/\log n)^{7/9} \le \frac{t^{1/3}}{f^{2/3}}$$

Thus, either way,  $\frac{n^{1/3}t^{1/9}f^{1/9}}{\log^{2/3}n}$  is dominated by the other two terms.

**Theorem 2.4.** Suppose a graph G has n vertices, t triangles, and local triangle bound y. Then

$$\chi(G) \lesssim \sqrt{\frac{n}{\log n}} + \frac{t^{1/3} \log f}{\log^{2/3} (t^2/y^3)}$$

*Proof.* Let  $f = \log(t^2/y^3)$  and let z = t/n. Let A denote the vertices of G which are incident on at least z triangles, and let B denote the vertices incident on at most z triangles.

As  $|A| \leq t/z$ , by Proposition 6.4, we have

$$\chi(A) \lesssim \sqrt{\frac{t/z}{\log(t/z)}} + \frac{t^{1/3}}{\log^{2/3}(t^2/y^3)} = \sqrt{\frac{n}{\log n}} + \frac{t^{1/3}\log f}{f^{2/3}}$$

By Proposition 2.1, we have

$$\chi(B) \lesssim \sqrt{\frac{n}{\log n}} + \frac{n^{1/3} z^{1/3}}{\log^{2/3} (n^2/z)} = \sqrt{\frac{n}{\log n}} + \frac{t^{1/3}}{\log^{2/3} (n^3/t)}$$

Now observe that  $n^3/t \ge n^3/(ny) = n^2/y \ge t^{2/3}/y$ , and so  $\log^{2/3}(n^3/t) \approx f$ . So  $\chi(B) \lesssim \sqrt{\frac{n}{\log n}} + \frac{t^{1/3}}{f^{2/3}}$ . Putting these terms together, we have that

$$\chi(G) \le \chi(A) + \chi(B) \lesssim \sqrt{\frac{n}{\log n}} + \frac{t^{1/3}\log f}{f^{2/3}}$$

#### 3 Bounds in terms of m

In this section, we show some bounds in terms of the edge count m, as well as triangle count (local and global). We next show a bound in terms of the global edge count and local triangle bound. This generalizes results of [8] and [3], which show similar bounds for the chromatic number as a function of m for triangle-free graphs.

**Theorem 3.1.** Suppose G has m edges and local triangle bound y. Then

$$\chi(G) \lesssim \frac{m^{1/3}}{\log^{2/3} m} + \frac{m^{1/4} y^{1/4}}{\log^{3/4}(m/y)}$$

*Proof.* Let  $f = \log(m/y)$ .

Let A denote the vertices of degree greater than d, and let B denote the vertices of degree  $\leq d$ , where d is some parameter to be chosen. As  $|A| \leq m/d$ , Theorem 2.1 gives

$$\chi(A) \lesssim \sqrt{\frac{m/d}{\log(m/d)}} + \frac{(m/d)^{1/3}y^{1/3}}{\log^{2/3}((m/d)^2/y)}$$

By Theorem 1.7 we have

$$\chi(B) \lesssim \frac{d}{\log(d^2/y)}$$

Suppose first that  $y > (m \log m)^{1/3}$ . Then let  $d = (myf)^{1/4}$ . We have  $\chi(B) \lesssim (myf)^{1/4} / \log(\sqrt{fe^f}) \lesssim$  $m^{1/4}y^{1/4}f^{-3/4}$ . To compute  $\chi(A)$ :

$$\begin{split} \chi(A) \lesssim \sqrt{\frac{m/d}{\log(m/d)}} + \frac{(m/d)^{1/3}y^{1/3}}{\log^{2/3}((m/d)^2/y)} \\ \lesssim \frac{m^{3/8}f^{-1/8}y^{-1/8}}{\log^{1/2}(m^{1/2}g^{1/4}f^{-1/4})} + \frac{m^{3/8}m^{1/4}y^{1/4}f^{-1/2}}{\log^{2/3}(e^{3/2f}f^{-1/12})} \\ \lesssim \frac{m^{3/8}f^{-1/8}y^{-1/8}}{f^{1/2}} + \frac{m^{3/8}m^{1/4}y^{1/4}f^{-1/2}}{f^{2/3}} \\ \lesssim m^{1/3}f^{-2/3} + m^{1/4}y^{1/4}f^{-3/4} \quad \text{as } y > (mf)^{1/3} \end{split}$$

Next, suppose that  $y \leq (m \log m)^{1/3}$ . Then  $y' = (m \log m)^{1/3}$  is also a valid upper bound on the local triangle count, and so (as we have already shown) this gives

$$\chi(G) \lesssim \frac{m^{1/3}}{\log^{2/3} m} + \frac{m^{1/4} (m \log m)^{1/12}}{\log^{3/4} (m/(m \log m)^{1/3})} \lesssim \frac{m^{1/3}}{\log^{2/3} m} + \frac{m^{1/3}}{\log^{2/3} m}$$

as desired.

**Theorem 3.2.** Suppose a graph G has m edges, t triangles, and local triangle bound y. Then

$$\chi(G) \lesssim \frac{t^{1/3} \log \log(t^2/y^3)}{\log^{2/3}(t^2/y^3)} + \frac{m^{1/3}}{\log^{2/3} m}$$

*Proof.* Let  $f = \log(t^2/y^3)$ . Let A denote the set of vertices incident on at least  $z = \frac{f^{1/3}t^{2/3}}{m^{1/3}}$  triangles, and

let *B* denote the remaining vertices; here *z* is a parameter to be determined. Then  $|A| \leq t/z$ . By Theorem 2.4, we have  $\chi(A) \leq \frac{t^{1/3} \log f}{f} + \sqrt{\frac{t/z}{\log(t/z)}}$ . Here  $t/z = (mt/\log m)^{1/3}$  and hence  $\log(t/z) \gtrsim \log m$  so  $\sqrt{\frac{t/z}{\log(t/z)}} \lesssim m^{1/6} t^{1/6} / (\log m)^{2/3}$ .

By Theorem 3.1, we have  $\chi(B) \lesssim \frac{m^{1/3}}{\log^{2/3} m} + \frac{(mz)^{1/4}}{\log^{3/4}(m/z)}$ . As  $m/z = \frac{m^{4/3}}{f^{1/3}t^{2/3}} \geq \frac{m^{1/3}}{f^{1/3}}$  we have  $\log(m/z) \gtrsim \frac{m^{1/4}}{d(m/z)}$ . log *m* and hence  $\frac{(mz)^{1/4}}{\log^{3/4}(m/z)} \lesssim m^{1/6} t^{1/6} / (\log m)^{2/3}$ .

E	-	
L		
L		
L		

We now claim that  $m^{1/6}t^{1/6}/(\log m)^{2/3} \lesssim (\frac{t^{1/3}\log f}{f} + \frac{m^{1/3}}{\log^{2/3}m})$ , which will prove the desired claim. When  $t \le m$ , we have  $m^{1/6}t^{1/6}/(\log m)^{2/3} \le \frac{m^{1/3}}{\log^{2/3}m}$ . When t > m, we have that  $m^{1/6}t^{1/6}/(\log m)^{2/3} \le \frac{t^{1/3}}{\log^{2/3}m}$ . As  $f \lesssim \log m$ , this shows that  $m^{1/6}t^{1/6}/(\log m)^{2/3} \lesssim \frac{t^{1/3}}{f^{2/3}}$ .

### 4 Lower bounds

We next show matching lower bounds. We will show that Theorems 2.1,3.1 are asymptotically tight for all admissible values of m, n, y, while Theorems 2.4 is tight up to factors for  $\log \log(t^2/y^3)$  for all admissible values of m, n, y, t. The situation for Theorem 3.2 is slightly more complicated; in general, the bounds given by Theorem 3.2 and Theorem 3.1 are incomparable. For a given value of m, n, y, t, we show that either Theorem 3.2 or Theorem 3.1 is asymptotically tight (the latter up to a factor of  $\log \log(t^2/y^3)$ ).

The lower bounds apply even for the fractional chromatic number (which is always less than or equal to the ordinary chromatic number).

We begin by recalling a result of [5]:

**Theorem 4.1** ([5]). For any integer  $n \ge 1$ , there exists a graph  $H_n$  on n vertices with the following properties:

- (A1)  $H_n$  is triangle-free
- (A2) Each vertex has degree at most  $O(\sqrt{n \log n})$
- $(A3) \ \alpha(H_n) \leq O(\sqrt{n \log n})$  (where  $\alpha(G)$  denotes the size of the maximum independent set of G)
- (A4)  $H_n$  has chromatic number  $\chi(H_n) \ge \Omega(\sqrt{\frac{n}{\log n}}).$

(We note that item (A2) is not stated explicitly in the paper [5], but can be deduced in terms of "Property 1" in that paper.)

Following a strategy of [2], we may construct a blow-up graph, which is an version of  $H_n$  inflated to have a small number of triangles, using the following construction:

**Proposition 4.2.** For any integers  $k, i \geq 1$ , there is a graph  $H_{k,i}$  with the following properties:

- (B1)  $H_{k,i}$  contains ni vertices
- (B2) Each vertex has triangle count at most  $\leq O(i^2 \sqrt{k \log k})$ .
- (B3) Each vertex has degree at most  $O(i\sqrt{k\log k})$
- $(B4) \ \alpha(H_{k,i}) \le O(\sqrt{k \log k}).$
- (B5) The fractional chromatic number of  $H_{n,i}$  satisfies  $\chi_f(H_{k,i}) \ge \Omega(i\sqrt{\frac{k}{\log k}})$ .

*Proof.* We replace each vertex of  $H_k$  with a clique on *i* vertices. For every edge  $(x, y) \in H_n$ , we place an edge between all the corresponding copies of x, y in  $H_{k,i}$ , a total of  $i^2$  edges. Now (B1) follows immediately from the fact that  $H_k$  contains k vertices and (B3) follows immediately from (A2).

To show (B2). Suppose that x', y', z' form a triangle of  $H_{k,i}$ , where x' is some fixed vertex of  $H_{k,i}$ . These vertices correspond to vertices x, y, z of  $H_k$ . If x, y, z are distinct, then this would imply that there are edges (x, y), (y, z), (x, z) in  $H_k$  which is impossible as  $H_k$  is triangle-free. In the next case, suppose that y = z = x. The total number of such triangles is at most  $i^2$  (since x is fixed from x' and y', z' must lie in the same clique as x'). In the next case, suppose  $y = z \neq x$ . Then there must be an edge in  $H_k$  from x to y. There are at most  $O(\sqrt{k \log k})$  choices of y and once y is fixed, at most  $i^2$  choices for y', z'. In the final case, suppose  $y = x \neq z$ . Again there must be an edge in  $H_k$  from x to z. There are at most  $O(\sqrt{k \log k})$  choices for z and at most  $i^2$  choices for y', z'. Adding all these cases, we see that there are  $O(i^2\sqrt{k \log k})$  triangles.

To show (B4). Observe that if I is an independent set of  $H_{k,i}$ , then all of its vertices must correspond to distinct vertices of  $H_k$ , and it must correspond to an independent set of  $H_k$ . So  $|I| \leq O(\sqrt{k \log k})$ .

The bound (B5) follows from (B1), (B4) and the general bound on fractional chromatic number  $\chi_f(G) \geq \frac{|V(G)|}{\alpha(G)}$ .

When apply Proposition 4.2, it will be convenient to sidestep the condition that i, k are integers.

**Proposition 4.3.** Let *i*, *k* be real numbers with *i*,  $k \ge 1$ . There is a graph  $H_{k,i}$  satisfying the following properties:

(B1)  $H_{k,i}$  contains O(ni) vertices

(B2) Each vertex has triangle count at most  $\leq O(i^2 \sqrt{k \log k})$ .

(B3) Each vertex has degree at most  $O(i\sqrt{k\log k})$ 

 $(B4) \ \alpha(H_{k,i}) \le O(\sqrt{k \log k}).$ 

(B5) The fractional chromatic number of  $H_{n,i}$  satisfies  $\chi_f(H_{k,i}) \ge \Omega(i\sqrt{\frac{k}{\log k}})$ .

*Proof.* Apply Proposition 4.2 with to construct  $H_{[i],[k]}$ . The condition that  $i, k \geq 1$  ensures that  $[i] \approx i$ and  $[k] \approx k$ . 

**Proposition 4.4.** For any integers  $n, y \le n^2, t \le ny$ , there is a graph G with at most n vertices, at most t triangles, local triangle bound y, and such that

$$\chi(G) \ge \chi_f(G) \gtrsim \sqrt{\frac{n}{\log n}} + \frac{t^{1/3}}{\log^{2/3}(t^2/y^3)}$$

*Proof.* By rescaling, it suffices to show that a graph has this value of  $\chi_f(G)$  and O(n) vertices, O(t) triangles, and O(y) local triangle bound. We will break this into a number of cases.

Define  $f = \log(t^2/y^3)$ . If f = 1, then we need to construct a graph G with  $\chi_f(G) \gtrsim \sqrt{\frac{n}{\log n}} + t^{1/3}$  and O(n) vertices. If  $\sqrt{n/\log n} > t^{1/3}$ , then simply take  $G = H_n$ . If  $\sqrt{n/\log n} < t^{1/3}$ , take G to be the complete graph on  $t^{1/3}$  vertices. This clearly has O(t) triangles and O(n) vertices. Also, it has a local triangle bound of  $t^{2/3} < y$ .

So we suppose that  $f \ge 1$ . If  $\sqrt{n/\log n} > \frac{t^{1/3}}{t^{2/3}}$ , this is easily achieved by taking  $G = H_n$ . Thus we assume throughout that  $\sqrt{n/\log n} \leq \frac{t^{1/3}}{t^{2/3}}$ .

Note that as  $t \le ny$  we have  $\frac{t^{1/3}}{f^{2/3}} \le \frac{(ny)^{1/3}}{\log^{2/3}(n^2/y)}$ . We thus claim that  $y \ge \sqrt{n \log n}$ . For,  $\frac{(ny)^{1/3}}{\log^{2/3}(n^2/y)}$  is an increasing function of y, and so if  $y < \sqrt{n \log n}$  then we would have  $\frac{(ny)^{1/3}}{\log^{2/3}(n^2/y)} < \sqrt{\frac{n}{\log n}} \le \frac{t^{1/3}}{f^{2/3}}$ , which we have already ruled out.

Then apply Proposition 4.3 with

$$i = \frac{y}{f^{1/3}t^{1/3}}, \qquad k = \frac{f^{1/3}t^{4/3}}{y^2}$$

We must first show that  $i, k \geq \Omega(1)$ .

Observe that  $\frac{y}{f^{1/3}t^{1/3}}$  is a decreasing function of t, and as  $t \le ny$  this implies that  $\frac{y}{f^{1/3}t^{1/3}} \ge \frac{y^{2/3}}{n^{1/3}\log(n^2/y)^{1/3}}$ . Our asymption that  $y \ge \sqrt{n \log n}$  implies that this is at least 1.

Similarly, the assumption that  $t^2 \ge y^3$  implies that  $\frac{f^{1/3}t^{4/3}}{y^2} \ge 1$ .

With these observations, one can immediately observe that  $H_{k,i}$  has O(t) triangles, O(y) triangles per vertex, and O(ty) vertices, and  $\chi_{\rm f}(H_{k,i}) \geq \Omega(i\sqrt{\frac{k}{\log k}})$ . Observe that  $k = t^2/y^3 \times \frac{f^{1/3}y}{t^{2/3}} \leq f \log^{1/3} f$ . Thus  $\log k = \Theta(f)$ . So  $i\sqrt{\frac{k}{\log k}} = \Omega(t^{1/3}/f^{2/3})$  as desired. 

Finally, observe that  $H_{k,i}$  has  $O(ik) \leq ty \leq n$  vertices.

**Corollary 4.5.** For any integers  $n, y \leq n^2$ , there is a graph G with at most n vertices and local triangle bound y, such that

$$\chi(G) \ge \chi_f(G) \gtrsim \sqrt{\frac{n}{\log n}} + \frac{(ny)^{1/3}}{\log^{2/3}(n^2/y)}$$

*Proof.* Apply Proposition 4.5 with t = ny.

We next show a lower bound on  $\chi(G)$  as a function of t, m, y. We first observe a simple relationship that must hold between these quantities:

**Proposition 4.6.** Suppose G has m edges, t triangles, and local triangle bound y. Then  $t \leq m\sqrt{y}$ .

*Proof.* Let  $a_v$  denote the number of triangles incident on each vertex  $v \in V$ . Then  $t = \frac{\sum_v a_v}{3}$ . Also, we have  $a_v \leq y$  and  $a_v \leq {d_v \choose 2}$  for each vertex v, where  $d_v$  denotes the degree of v. Finally,  $\sum_v d_v = 2m$ . Thus, we have that

$$t \le \frac{\sum_{v} \min(d^2/2, y)}{3}$$

Note that  $\min(\binom{x}{2}, y) \leq \frac{xy}{\sqrt{2y}}$ , so  $t \leq \sum_{v} \frac{d_v y}{3\sqrt{2y}} \leq \frac{2my}{3\sqrt{2y}} \leq m\sqrt{y}$ .

**Proposition 4.7.** Given any integers  $m, y \leq m, t \leq m\sqrt{y}$ , there is a graph G with at most m edges, at most t triangles, local triangle bound y, and

$$\chi(G) \ge \chi_f(G) \gtrsim \min\left(\frac{(my)^{1/4}}{\log^{3/4}(m/y)}, \frac{t^{1/3}}{\log^{2/3}(t^2/y^3)}\right) + \frac{m^{1/3}}{\log^{2/3}m}$$

Proof. Let  $f = \log(m/y)$  and let  $g = \log(t^2/y^3)$ . **Case I:**  $t \ge \frac{g^2(my)^{3/4}}{f^{9/4}}$ . Then we need to construct G such that  $\chi_f(G) \gtrsim \frac{(my)^{1/4}}{f^{3/4}} + \frac{m^{1/3}}{\log^{2/3} m}$ . We can construct a triangle-free graph with  $\Omega(m^{1/3}/\log^{2/3} m)$  edges, so we may assume that  $y \ge \frac{f^3m^{1/3}}{\log^{8/3} m}$ .

Apply Proposition 4.3 with

$$i = \frac{y^{3/4}}{(mf)^{1/4}}$$
  $k = m/y$ 

We must show that  $i, k \gtrsim 1$ . Clearly  $k \ge 1$ . If  $y \ge \sqrt{m}$ , then  $i \ge m^{1/8}/f^{1/4} \gtrsim 1$ ; otherwise, then  $f \approx \log m$ . Using the bound  $y \ge \frac{f^3 m^{1/3}}{\log^{8/3} m}$ , we see that  $i \ge \frac{f^2}{\log^2 m} \gtrsim 1$ . One may easily verify that this graph has m edges and local triangle bound y. It has at most iky triangles. As  $t \ge \frac{g^2 (my)^{3/4}}{f^{9/4}}$ , it suffices to show that  $iky \lesssim \frac{g^2 (my)^{3/4}}{f^{9/4}}$ . This reduces to showing that  $f \lesssim g$ . To see this, observe that  $t^2/y^3 \ge \frac{g^4 (m/y)^{3/2}}{f^{9/2}} \ge \frac{g^4 \exp(f)}{f^{9/2}}$ . This implies that  $\log(t^2/y^3) \gtrsim \log(e^{f/2}) \gtrsim f$  as desired.

desired.

Case II:  $t \leq \frac{g^2(my)^{3/4}}{f^{9/4}}$ . Then we need to construct G such that  $\chi_{\rm f}(G) \gtrsim \frac{t^{1/3}}{g^{2/3}} + \frac{m^{1/3}}{\log^{2/3} m}$ . We can construct a triangle-free graph with  $\Omega(m^{1/3}/\log^{2/3}m)$  edges, so we may assume that  $t \geq \frac{mg^2}{\log^2 m}$ 

If  $y \ge t^{2/3}$ , then g = 1 and our goal is achieved by taking G to be a clique on  $\sqrt{y}$  vertices. So we assume  $y \le t^{2/3}$ .

Apply Proposition 4.3 with

$$i = \frac{y}{(gt)^{1/3}}, k = \frac{g^{1/3}t^{4/3}}{y^2}$$

The bound  $k \gtrsim 1$  follows immediately from the bound  $y \leq t^{2/3}$ . If  $y > m^{0.51}$ , then  $i \geq \frac{m^{0.51}}{(am^{3/2})^{1/3}} \gtrsim 1$ . Let us thus assume that  $y \leq m^{0.51}$ , in which case  $f = \log(m/y) \approx \log m$ .

Note that we have the two bounds

$$\frac{mg^2}{\log^2 m} \le t \le \frac{g^2 (my)^{3/4}}{f^{9/4}}$$

These together imply that  $y \ge \frac{f^3 m^{1/3}}{\log^{8/3} m} \gtrsim \frac{(\log m)^3 m^{1/3}}{\log^{8/3} m} = (m \log m)^{1/3}$ . So

$$i \gtrsim \frac{(m\log m)^{1/3}}{(gt)^{1/3}} \ge \frac{(m\log m)^{1/3}}{((\log m)m^{3/2})^{1/3}} = 1$$

We easily verify that this graph has O(t) triangles and local triangle bound O(y). Finally, we need to verify that G has O(m) edges. Property (B3) implies that G has at most  $ik \times O(i\sqrt{k\log k})$  edges. Note that

here  $k = g^{1/3} t^{4/3} / y^2 = t^2 / y^3 \times g \frac{y}{t^{2/3}} = \sqrt{\exp(g)} / g$ . So  $\log k \approx g$ . So G has  $O(\frac{g^{1/3} t^{4/3} / y}{\lesssim} (g/f)^3 m$  edges. So, we need to show that  $g \lesssim f$ ; we assume without loss of generality that g is large than any fixed constant. Now observe that  $t^2 / y^3 \le \frac{g^4 (m/y)^{3/2}}{f^{9/2}} \le 2g^4 \exp(f)$ . Taking logarithms (and noting that log agrees with ln for g sufficiently large), we see that  $g \leq \log 2 + 4 \log g + f$ . As  $g - 4 \log g - \log 2 \approx g$  for g sufficiently large, this implies that  $g \lesssim f$  as desired. 

#### Getting the correct coefficient of $t^{1/3}$ $\mathbf{5}$

Suppose we have no information on the local triangle counts; in this case, Theorem 2.3 would give us a bound (solely in terms of n and t) of the form

$$\chi(G) \leq O(t^{1/3} + \sqrt{\frac{n}{\log n}})$$

for a graph with n vertices and t triangles. This bound is clearly tight up to constant factors. In this section, we compute a more exact formula, which gives us the correct coefficient of the term  $t^{1/3}$ . The correct coefficient of the term  $\sqrt{\frac{n}{\log n}}$  is not currently known even for triangle-free graphs. We begin by showing a weaker bound:

**Proposition 5.1.** If G has at most n vertices and at most t triangles, then

$$\chi(G) \le 100\sqrt{n} + (6t)^{1/3}$$

*Proof.* Let us define  $f(x,y) = 100\sqrt{n} + (6t)^{1/3}$ , and let d = |f(n,t)|. Suppose that some vertex v of G has degree strictly less than d. In that case, apply the induction hypothesis to G - v, obtaining a coloring using at most  $|f(n-1,t)| \leq d$  colors. At least one of those colors does not appear in a neighbor of v, so we can extend the coloring to v as desired.

So we suppose that G has minimum degree d. Among all vertices, select one vertex v which participtes in the minimum number y of triangles. Select a set Y consisting of exactly d neighbors of v; let r denote the number of edges in G[Y]. It must be the case that  $r \leq y$  and every vertex in G[Y] is incident upon at least y triangles.

Now, by Theorem 1.9, G[Y] contains an independent set I of size at least  $\frac{d}{2r/d+1} \geq \frac{d}{2u/d+1}$ . Let us define

$$s = \frac{d}{2y/d + 1}$$

so that |I| > s.

Assign all vertices in I one new color, and recurse on G - I. As v is chosen to be incident on the minimum number of triangles, every vertex of I is itself incident on at least y triangles; furthermore as I is independent these triangles are all distinct. So I is incident on at least ys triangles, which are all removed in G-I. Thus G-I has at most n-s vertices and at most t-ys triangles. By induction hypothesis, then  $\chi(G-I) \leq f(n-s,t-ys)$  and so we have

$$\begin{split} \chi(G) &\leq 1 + f(n-s,t-ys) \\ &\leq 1 + 100\sqrt{n-s} + 6^{1/3}(t-ys)^{1/3} \\ &\leq 1 + 1 + 100\sqrt{n} - \frac{50s}{\sqrt{n}} + 6^{1/3}(t^{1/3} - \frac{ys}{3t^{2/3}}) \\ &= f(n,t) + 1 - \frac{50s}{\sqrt{n}} - \frac{6^{1/3}ys}{3t^{2/3}} \end{split}$$

We want to show that  $\chi(G) \leq f(n, t)$ ; thus, it suffices to show that

$$\frac{50s}{\sqrt{n}} + \frac{6^{1/3}ys}{3t^{2/3}} \ge 1 \tag{1}$$

Substituting in the value for s, simple algebraic manipulations show that this is equivalent to showing:

$$d^{2}\left(\frac{150}{\sqrt{n}} + \frac{6^{1/3}y}{t^{2/3}}\right) \ge 3(d+2y) \tag{2}$$

Now, observe that the LHS and RHS of (2) are both linear functions of y. As y is a non-negative real number, it suffices to show that (2) holds at the extreme values  $y = 0, y \to \infty$  respectively.

When y = 0, then (2) reduces to:

$$d\frac{150}{\sqrt{n}} \ge 3 \tag{3}$$

We have  $d \ge \lfloor 100\sqrt{n} \rfloor \ge 50\sqrt{n}$ . Thus (3) easily holds. When  $y \to \infty$ , then (2) reduces to:

$$d^2(\frac{6^{1/3}}{t^{2/3}}) \ge 6 \tag{4}$$

Note that  $d \ge (6t)^{1/3}$ , and so again this is easily seen to hold. This completes the induction.

**Theorem 5.2.** Suppose that G contains t triangles and n vertices. Then

$$\chi(G) \le O\Big(\sqrt{\frac{n}{\log n}} + \frac{t^{1/3} (\log \log n)^{3/2}}{\log n}\Big) + (6t)^{1/3} = O(\sqrt{\frac{n}{\log n}}) + (6^{1/3} + o(1))t^{1/3}$$

*Proof.* Let  $y = t^{1/3} \log^2 n$ . Let A denote the vertices of G which are incident on at least y triangles, and let a be the number of triangles contained in G[A]. As  $|A| \leq 3t/y$  vertice, by Proposition 5.1 we have  $\chi(A) \leq O(\sqrt{t/y}) + (6a)^{1/3}$ .

Let B = V - A, and suppose G[B] contains b triangles. By Theorem 2.4 we have

$$\chi(B) \lesssim \sqrt{\frac{n}{\log n}} + \frac{(t-a)^{1/3} \log \log((t-a)^2/y^3)}{\log^{2/3}((t-a)^2/y^3)}$$

Putting these terms together, we have

$$\chi(G) \le (6a)^{1/3} + O\left(\sqrt{\frac{n}{\log n}} + \frac{b^{1/3}\log\log(b^2/y^3)}{\log^{2/3}(b^2/y^3)} + \frac{t^{1/3}}{\log n}\right)$$

As  $a + b \leq t$ , we have

$$\chi(G) \le (6t)^{1/3} + O\left(\sqrt{\frac{n}{\log n}} - \frac{2^{1/3}b}{3^{2/3}t^{2/3}} + \frac{b^{1/3}\log\log(b^2/y^3)}{\log^{2/3}(b^2/y^3)} + \frac{t^{1/3}}{\log n}\right)$$
(5)

Our next task is to show an upper bound on the quantity

$$-\frac{2^{1/3}b}{3^{2/3}t^{2/3}} + \frac{b^{1/3}\log\log(b^2/y^3)}{\log^{2/3}(b^2/y^3)}$$
(6)

If  $b \leq t/\log^3 n$ , this this is clearly at most  $\frac{t^{1/3}}{\log n}$ . Otherwise, we have  $b^2/y^3 \geq t/\log^{12} n$ . Again, if  $t \leq n$  then Theorem 5.2 easily holds, so  $\log(b^2/y^3) \gtrsim \log n$ , and

$$-\frac{2^{1/3}b}{3^{2/3}t^{2/3}} + \frac{b^{1/3}\log\log(b^2/y^3)}{\log^{2/3}(b^2/y^3)} \le O(\frac{b}{t^{2/3}} + \frac{b^{1/3}\log\log n}{\log^{2/3}n})$$

and simple calculus shows that is bounded by  $O(\frac{t^{1/3}(\log \log n)^{3/2}}{\log n})$ .

Thus, substituting this bound into (5), we have

$$\chi(G) \le (6t)^{1/3} + O\left(\sqrt{\frac{n}{\log n}} + \frac{t^{1/3}(\log\log n)^{3/2}}{\log n}\right)$$

**Theorem 5.3.** Suppose that G contains t triangles and m edges. Then

$$\chi(G) \le O\Big(\frac{m^{1/3}}{\log^{2/3} m} + \frac{t^{1/3} (\log \log m)^{3/2}}{\log m}\Big) + (6t)^{1/3} = O\Big(\frac{m^{1/3}}{\log^{2/3} m}\Big) + (6^{1/3} + o(1))t^{1/3}$$

*Proof.* The proof is nearly identical to Theorem 5.2, except that we apply Theorem 3.2 instead of Theorem 2.4.  $\Box$ 

Theorems 5.2, 5.3 are easily seen to be tight, up to lower-order terms.

**Observation 5.4.** For any integers  $n \ge 1$  and  $1 \le t \le n^3$  there is a graph G with  $\le n$  vertices and  $\le t$  triangles and

$$\chi_f(G) \ge \Omega(\sqrt{\frac{n}{\log n}}) + (6^{1/3} - g(n))t^{1/3}$$

where  $g(n) \to 0$  as  $n \to \infty$ .

For any integers  $m \ge 1$  and  $1 \le t \le m^{3/2}$  there is a graph G with  $\le m$  vertices and  $\le t$  triangles and

$$\chi_f(G) \ge \Omega(\sqrt{m^{1/3} \log^{2/3} m} + (6^{1/3} - g(m))t^{1/3})$$

where  $g(m) \to 0$  as  $m \to \infty$ .

*Proof.* We show only the bound in terms of the vertex count n; the bound in terms of edge count is similar.

Suppose  $t^{1/3} > \sqrt{n/\log n}$ . Then create a clique on  $\lfloor (6t)^{1/3} \rfloor$  vertices. This satisfies the triangle count and has less t triangles and  $(6t)^{1/3} \leq n$  vertices. Its fractional chromatic number is at least  $\lfloor (6t)^{1/3} \rfloor \geq (6t)^{1/3} - 1$ . Thus we have

$$\begin{split} \chi(G) &\geq (6t)^{1/3} + \sqrt{\frac{n}{\log n}} - \sqrt{\frac{n}{\log n}} - 1\\ &\geq (6t)^{1/3} + \sqrt{\frac{n}{\log n}} - t^{1/3} (\sqrt{\frac{n}{2t^{1/3}\log n}})\\ &\geq (6t)^{1/3} + \sqrt{\frac{n}{\log n}} - O(t^{1/3}/\sqrt{\log n}) \end{split}$$

as desired.

If  $t^{1/3} \leq \sqrt{n/\log n}$ , then use the graph  $H_n$  as in Theorem 4.1, which has n vertices and fractional chromatic number  $\Omega(\sqrt{n/\log n})$  and is triangle-free.

## 6 Conjectured tight bounds

The following conjecture seems natural:

**Conjecture 6.1.** Suppose G has n vertices, t triangles, and local triangle bound y. Then

$$\chi(G) \lesssim \frac{t^{1/3}}{\log^{2/3}(t^2/y^3)} + \sqrt{\frac{n}{\log n}}$$

This conjecture would strengthen Theorem 2.4, would give Proposition 2.1 as a special case (as  $t \le ny$ ), and would match the lower bound Proposition 4.4.

As further evidence for Conjecture 6.1, we show that it would follow from a conjecture regarding fractional chromatic number.

**Conjecture 6.2.** Suppose that G is triangle-free and is d-degenerate. Then

$$\chi_f(G) \lesssim \frac{d}{\log d}$$

We observe that [2] showed that a triangle-free, d-degenerate graph may have  $\chi(G)$  as large as d; the graph G which achieves this indeed has  $\chi_{\rm f}(G) \lesssim \frac{d}{\log d}$ . Thus Conjecture 6.2 is itself a natural conjecture, which may have applications outside this paper. In the remainder of this section, we shall show that Conjecture 6.2 implies Conjecture 6.1. For the remainder of this section, we shall suppose that Conjecture 6.2 holds.

**Proposition 6.3.** Suppose that G is d-degenerate and has local triangle bound y. Then

$$\chi_f(G) \lesssim \frac{d}{\log(d^2/y)}$$

*Proof.* We need to show a random distribution on the independent sets I of G, such that for any vertex  $v \in G$  we have  $P(v \in I) \gtrsim \frac{\log(d^2/y)}{d}$ .

Since G is d-degenerate, let us fix an ordering of its edges such that each vertex has out-degree at most d. Consider the following random process: we mark each vertex of G independent with probability  $p = 0.01/\sqrt{y}$ . If some vertex w has more than  $d/\sqrt{y}$  marked out-neighbors, or has any triangles, then we unmark w. Next, we let H denote the subgraph induced on the marked vertices. H is d'-degenerate and triangle-free, where  $d' = d/\sqrt{y}$ , so by Conjecture 6.2 we select an independent set I wherein each  $v \in H$  survives to I with probability  $\Omega(\frac{\log d'}{d'})$ .

Observe that for any vertex  $v \in G$ , the expected number of marked triangles containing v is at most  $yp^2 \leq 0.01$ . Also, the expected number of marked out-neighbors of v is at most pd. Hence, by Markov's inequality, each vertex v is unmarked with probability at most 1/2. Thus, the overall probability that v survives to H is  $\Omega(p)$ . Overall, the probability that  $v \in I$  is  $\Omega(p) \times \Omega(\frac{\log d'}{d'}) \gtrsim \frac{d}{\log(d^2/y)}$ 

We may now prove Conjecture 6.1.

**Proposition 6.4.** Suppose a graph G has n vertices, t triangles, and local triangle bound y. Then for some universal constant C we have

$$\chi(G) \le C(\sqrt{n} + \frac{t^{1/3}}{\log^{2/3}(t^2/y^3)})$$

*Proof.* We prove this by induction on G. Let us set  $f = \log(t^2/y^3)$  and  $d = (tf)^{1/3} + \sqrt{n}$ . We may assume that n, t, and f are larger than any needed constant; otherwise, this follows from Theorem 2.4 if the constant C is sufficiently large.

We will first show that there is an independent set I of G which is incident upon either  $\Omega(\sqrt{n})$  vertices or  $\Omega((tf)^{2/3})$  triangles. In order to show this, we consider two cases: whether G is or is not d-degenerate.

**CASE I:** G is not d-degenerate. Then there is some set of vertices U such that G[U] has minimum degree at d. Let  $w \in U$  be the vertex of U which is incident upon the fewest triangles of G; suppose that w is incident upon k triangles and has degree at least d.

If  $k \leq d$ , then by Turán's Theorem we may select an independent set  $I \subseteq N(w) \cap U$  with  $|I| \geq \frac{d}{1+\frac{k}{d}} \gtrsim d \geq \sqrt{n}$ , as desired.

If  $k \leq d$ , then by Turán's Theorem we may select an independent set  $I \subseteq N(w) \cap U$  with  $|I| \geq \frac{d}{1+\frac{k}{d}} \gtrsim d^2/k$ . Since w is incident on the smallest number of triangles of U, every vertex in I must be incident upon at least k triangles. So I is incident upon at least  $|I|k \gtrsim d^2 \geq (tf)^{2/3}$  triangles, as desired.

**CASE II:** G is d-degenerate and  $\sqrt{n} < (tf)^{1/3}$ . One equivalent formulation of fractional chromatic number is the following: if  $\chi_f(G) \leq r$ , then given any weighting function  $w: V \to \mathbf{R}_{\geq 0}$ , there exists an independent set I with  $\sum_{v \in I} w(v) \geq \frac{\sum_{v \in V} w(v)}{r}$ .

Now use the weighting function w defined by setting w(v) to the number of triangles incident upon v. By Proposition 6.3, there is an independent set I with  $\sum_{v \in I} w(v) \gtrsim \frac{\sum_{v \in V} w(v)}{d/\log(d^2/y)} \gtrsim \frac{t \log(d^2/y)}{d}$ . As  $(tf)^{1/3} > \sqrt{n \log n}$ , observe that

$$\frac{\log(d^2/y)}{d} \gtrsim \frac{\log((tf)^{2/3}/y)}{(tf)^{1/3}} \gtrsim \frac{\log(e^f f^2)}{(tf)^{1/3}} \approx \frac{f^{2/3}}{t^{1/3}}$$

Thus, the number of triangles incident upon the vertices of I is at least  $\sum_{v \in I} w(v) \gtrsim (tf)^{2/3}$ .

**CASE III:** G is d-degenerate and  $\sqrt{n} \ge (tf)^{1/3}$ . Since G is d-degenerate, we have  $\chi(G) \le d$ . Hence G has an independent set of size at least  $n/d \ge \sqrt{n}$ .

Now suppose that we have found an independent set I which is incident upon at least  $\Omega(\sqrt{n})$  vertices or  $\Omega((tf)^{2/3})$  triangles. We assign all the vertices in I one new color, remove I from the graph, and apply our induction hypothesis to color G - I.

If  $|I| \ge c\sqrt{n}$ , this gives

$$\chi(G) \le 1 + C(\sqrt{n - c\sqrt{n}} + \frac{t^{1/3}}{\log^{2/3}(t^2/y^3)}) \le 1 + C(\sqrt{n} + \frac{t^{1/3}}{\log^{2/3}(t^2/y^3)}) - \frac{c\sqrt{n}}{2\sqrt{n}}$$

This is at most  $C(\sqrt{n} + \frac{t^{1/3}}{\log^{2/3}(t^2/y^3)})$  for C a sufficiently large constant, and the induction proceeds in this case.

Next, suppose that I is incident upon at least r triangles, where  $r = c(tf)^{2/3}$  for some constant c. These triangles are all removed in G - I, and so by induction hypothesis we have

$$\chi(G) \le 1 + C(\sqrt{n} + \frac{(t-r)^{1/3}}{\log^{2/3}((t-r)^2/y^3)})$$

Now consider the function  $F(x) = x^{1/3}/\log^{2/3}(x^2/y^3)$ . For  $x \ge e^2y^{3/2}$ , this function is increasing concave-down. We claim that  $t - r \ge e^2y^{3/2}$ . For, observe that  $r \le t^{0.9}$  and so  $t - r \ge t/2$  for t above a sufficiently large constant. Thus, it suffices to show that  $t \ge 2e^2y^{3/2}$ , which follows from our assumption that f exceeds any sufficiently large constant.

Thus, we may upper-bound the expression F(t-r) by its tangent line F(t) - rF'(t), yielding

$$\chi(G) \le 1 + C(\sqrt{n} + F(t)) \le 1 + C(\sqrt{n} + \frac{t^{1/3}}{f^{2/3}} - r\frac{f-4}{t^{2/3}f^2})$$

Tto show that  $\chi(G) \leq C(\sqrt{n} + \frac{t^{1/3}}{f^{2/3}})$  it thus suffices to show that

$$1 - Cc \frac{(tf)^{2/3}(f-4)}{3t^{2/3}f^{5/3}}) \le 0$$

By taking C to be sufficiently large, this follows immediately from our assumption that f is larger than any needed constant. Thus the induction again holds in this case.

From Proposition 6.4, we can tighten many of the bounds in our paper. We omit the proofs since they are essentially identical to proofs we have already encountered.

**Proposition 6.5.** Suppose a graph G has n vertices, t triangles, and local triangle bound y. Then

$$\chi(G) \lesssim \sqrt{\frac{n}{\log n}} + \frac{t^{1/3}}{\log^{2/3}(t^2/y^3)}$$

**Proposition 6.6.** Suppose a graph G has m edges, t triangles, and local triangle bound y. Then

$$\chi(G) \lesssim \frac{t^{1/3}}{\log^{2/3}(t^2/y^3)} + \frac{m^{1/3}}{\log^{2/3}m}$$

**Proposition 6.7.** Suppose that G contains t triangles and n vertices. Then

$$\chi(G) \le O\left(\sqrt{\frac{n}{\log n}} + \frac{t^{1/3}}{\log n}\right) + (6t)^{1/3} = O\left(\sqrt{\frac{n}{\log n}}\right) + (6^{1/3} + o(1))t^{1/3}$$

### 7 Acknowledgements

Thanks to Vance Faber, Paul Burkhardt, Louis Ibarra, and Aravind Srinivasan for helpful suggestions, discussions, and proofreading. Thanks to Seth Pettie for some clarifications about Theorem 1.3. Thanks to Hsin-Hao Su for some discussions about fractional chromatic number.

## References

- Ajtai, M., Komlós, J., Szemerédi, E.: A note on Ramsey numbers. Journal of Combinatorial Theory Series A 29, pp. 354-360 (1980)
- [2] Alon, N., Krivelevich, M., Sudakov, B.: Coloring graphs with sparse neighborhoods. Journal of Combinatorial Theory, Series B 77.1, pp. 73-82 (1999)
- [3] Gimbel, J., Thomassen, C.: Coloring triangle-free graphs with fixed size. Discrete Mathematics 219-1, pp. 275-277 (2000)
- [4] Jamall, M.: A Brooks' theorem for triangle-free graphs, arXiv:1106.1958 (2011)
- [5] Kim, J.H.: The Ramsey number R(3,t) has order of magnitude  $t^2/\log t$ . Random Structures & Algorithms 7-3, pp. 173-207 (1995).
- [6] Molloy, M., Reed, B.: Graph colouring and the probabilistic method. Algorithms and Combinatorics, Springer (2001)
- [7] Moser, R., Tardos, G.: A constructive proof of the general Lovász Local Lemma. Journal of the ACM 57.2, p. 11 (2010)
- [8] Nilli, A.: Triangle-free graphs with large chromatic number. Discrete Mathematics 211-1, pp. 261-262 (2000)
- [9] Pettie, S., Su, H.: Distributed coloring algorithms for triangle-free graphs. Information and Computation 243, pp, 263-280 (2015)
- [10] Vu, V.: A general upper bound on the listing chromatic number of locally sparse graphs. Combinatorics, Probability, and Computing 11, pp. 103-111 (2002).