

The higher sharp I: on $M_1^\#$

Yizheng Zhu

Institut für mathematische Logik und Grundlagenforschung
 Fachbereich Mathematik und Informatik
 Universität Münster
 Einsteinstr. 62
 48149 Münster, Germany

February 8, 2022

Abstract

We establish the descriptive set theoretic representation of the mouse $M_n^\#$, which is called $0^{(n+1)\#}$. This part deals with the case $n = 1$.

1 Introduction to this series of papers

This is an introduction to a series of four papers.

The collection of projective subsets of \mathbb{R} is the minimum one which contains all the Borel sets and is closed under both complements and continuous images. Despite its natural-looking definition, many fundamental problems about projective sets are undecidable in ZFC, for instance, if all projective sets are Lebesgue measurable. The axiom of Projective Determinacy (PD) is the most satisfactory axiom that settles these problems by producing a rich structural theory of the projective sets. PD implies certain regularity properties of projective sets: all projects of reals are Lebesgue measurable (Mycielski, Swierczkowski), have the Baire property (Banach, Mazur) and are either countable or have a perfect subset (Davis) (cf. [24]). The structural theory of the projective sets are centered at good Suslin representations of projective sets. Moschovakis [24] shows that PD implies the scale property of the pointclasses Π_{2n+1}^1 and Σ_{2n+2}^1 . It follows that there is a nicely behaved tree T_{2n+1} that projects to the good universal Σ_{2n+2}^1 set. So the

analysis of Σ_{2n+2}^1 sets is reduced to that of the tree T_{2n+1} , the canonical model $L[T_{2n+1}]$ and its relativizations. The canonicity of $L[T_{2n+1}]$ is justified by Becker-Kechris [3] in the sense that $L[T_{2n+1}]$ does not depend on the choice of T_{2n+1} . The model $L[T_{2n+1}]$ turns out to have many analogies with $L = L[T_1]$. These analogies support the generalizations of classical results on Σ_2^1 sets to Σ_{2n+2}^1 sets.

The validity of PD is further justified by Martin-Steel [23]. They show that PD is a consequence of large cardinals: if there are n Woodin cardinals below a measurable cardinal, then Π_{n+1}^1 sets are determined. Inner model quickly developed into the region of Woodin cardinals. $M_n^\#$, the least active mouse with n Woodin cardinals, turns out to have its particular meaning in descriptive set theory. Martin [22] (for $n = 0$) and Neeman [25, 26] (for $n \geq 1$) show that $M_n^\#$ is many-one equivalent to the good universal $\mathcal{O}^{n+1}(<\omega^2\text{-}\Pi_1^1)$ real. Steel [32] shows that $L[T_{2n+1}] = L[M_{2n,\infty}^\#|\delta_{2n+1}^1]$ where $M_{2n,\infty}^\#$ is the direct limit of all the countable iterates of $M_{2n}^\#$, and that δ_{2n+1}^1 is the least cardinal that is strong up to the least Woodin of $M_{2n,\infty}^\#$. This precisely explains the analogy between $L[T_{2n+1}]$ and L . The mechanism of inner model theory is therefore applicable towards understanding the structure $L[T_{2n+1}]$.

In this series of papers, we generalize the Silver indiscernibles for L to the level- $(2n + 1)$ indiscernibles of $L[T_{2n+1}]$. The theory of $L[T_{2n+1}]$ with the level- $(2n + 1)$ indiscernibles will be called $0^{(2n+1)\#}$, which is many-one equivalent to $M_{2n}^\#$. At the level of mice with an odd number of Woodins, $M_{2n-1}^\#$ is the optimal real with the basis result for Σ_{2n+1}^1 sets (cf. [34, Section 7.2]): Every nonempty Σ_{2n+1}^1 set has a member recursive in $M_{2n-1}^\#$. The basis result for Σ_{2n+1}^1 was originally investigated in [16], with the intention of generalizing Kleene's basis theorem: Every nonempty Σ_1^1 set of real has a member recursive in Kleene's \mathcal{O} . The real y_{2n+1} , defined [16], turns out Δ_{2n+1}^1 equivalent to $M_{2n-1}^\#$. In this series of papers, we define the canonical tree T_{2n} that projects to a good universal Π_{2n}^1 set. It is the natural generalization of the Martin-Solovay tree T_2 that projects a good universal Π_2^1 set. We show that $L_{\kappa_{2n+1}}[T_{2n}]$, the minimum admissible set over T_{2n} , shares most of the standard properties of $L_{\omega_1^{CK}}$, in particular, the higher level analog of the Kechris-Martin theorem [13, 14]. We define $0^{(2n)\#}$ as the set of truth values in $L_{\kappa_{2n+1}}[T_{2n}]$ for formulas of complexity slightly higher than Σ_1 . $0^{(2n)\#}$ is many-one equivalent to both $M_{2n+1}^\#$ and y_{2n+1} . Summing up, we have

$$0^{(n+1)\#} \equiv_m M_n^\#.$$

We start to give a detailed explanation of the influence of the higher sharp in the structural theory of projective sets and in inner model theory. The

set theoretic structures tied to Π_1^1 sets are $L_{\omega_1^{CK}}$ and its relativizations. The classical results on Π_1^1 sets and $L_{\omega_1^{CK}}$ include:

1. (Model theoretic representation of Π_1^1) $A \subseteq \mathbb{R}$ is Π_1^1 iff there is a Σ_1 formula φ such that $x \in A \leftrightarrow L_{\omega_1^x}[x] \models \varphi(x)$.
2. (Mouse set) $x \in \mathbb{R} \cap L_{\omega_1^{CK}}$ iff x is Δ_1^1 iff x is Δ_1^1 in a countable ordinal.
3. (The transcendental real over $L_{\omega_1^{CK}}$) \mathcal{O} is the Σ_1 -theory of $L_{\omega_1^{CK}}$.
4. (Π_1^1 -coding of ordinals below ω_1) $x \in \text{WO}$ iff x codes a wellordering of a subset of ω . Every ordinal below ω_1 is coded by a member of WO . WO is Π_1^1 .

Σ_2^1 sets are ω_1 -Suslin via the Shoenfield tree T_1 . The complexity of T_1 is essentially that of WO , or Π_1^1 . The set-theoretic structures in our attention are $L = L[T_1]$ and its relativizations. Assuming every real has a sharp, the classical results related to L include:

1. (Model theoretic representation of Σ_2^1) $A \subseteq \mathbb{R}$ is Σ_2^1 iff there is a Σ_1 formula φ such that $x \in A \leftrightarrow L[x] \models \varphi(x)$.
2. (Mouse set) $x \in \mathbb{R} \cap L$ iff x is Δ_2^1 in a countable ordinal.
3. (The transcendental real over L) $0^\#$ is the theory of L with Silver indiscernibles, or equivalently, the least active sound mouse projecting to ω .
4. (Δ_3^1 -coding of ordinals below u_ω) WO_ω is the set of sharp codes. Every ordinal $\alpha < u_\omega$ has a sharp code $\langle \ulcorner \tau \urcorner, x^\# \rangle$ so that $\alpha = \tau^{L[x]}(x, u_1, \dots, u_k)$. The comparison of sharp codes is Δ_3^1 .

Inner model theory start to participate at this level. Based on the theory of sharps for reals, the Martin-Solovay tree T_2 is defined. T_2 is essentially a tree on u_ω . The complexity of T_2 is Δ_3^1 via the sharp coding of ordinals.

Σ_3^1 sets are u_ω -Suslin via the Martin-Solovay tree T_2 . The structures tied to Π_3^1 sets are $L_{\kappa_3}[T_2]$ and its relativizations. The theory at this level is in parallel to Π_1^1 sets and $L_{\omega_1^{CK}}$:

1. (Model theoretic representation of Π_3^1 , [13, 14]) $A \subset \mathbb{R}$ is Π_3^1 iff there is a Σ_1 formula φ such that $x \in A \leftrightarrow L_{\kappa_3^x}[T_2, x] \models \varphi(T_2, x)$.
2. (Mouse set, [13, 14, 16, 30]) $x \in \mathbb{R} \cap L_{\kappa_3}[T_2]$ iff x is Δ_3^1 in a countable ordinal iff $x \in \mathbb{R} \cap M_1^\#$.

3. (The transcendental real over $L_{\kappa_3}[T_2]$, Theorem 3.8) $M_1^\# \equiv_m 0^{2\#}$.
4. (Π_3^1 -coding of ordinals below δ_3^1 , essentially by Kunen in [29]) $\text{WO}^{(3)}$ is the set of reals that naturally code a wellordering of u_ω . $\text{WO}^{(3)}$ is Π_3^1 .

In general, if Γ is a pointclass, α is an ordinal, and $f : \mathbb{R} \rightarrow \alpha$ is a surjection, then $\text{Code}(f) = \{(x, y) : f(x) \leq f(y)\}$ and f is in Γ iff $\text{Code}(f)$ is in Γ ; α is Γ -wellordered cardinal iff there is a surjection $f : \mathbb{R} \rightarrow \alpha$ such that f is in Γ but there is no $\beta < \alpha$ and surjections $g : \mathbb{R} \rightarrow \beta$, $h : \beta \rightarrow \alpha$ such that both g and $\{(x, y) : f(x) = h \circ g(y)\}$ is in Γ . The above list can be continued:

5. The uncountable Δ_3^1 wellordered cardinals are $(u_k : 1 \leq k \leq \omega)$.

The heart of the new knowledge at this level is the equality of pointclass in Theorem 3.5: $\mathcal{O}^2(<\omega^2\text{-}\Pi_1^1) = <u_\omega\text{-}\Pi_3^1$. This is the main objective of this paper. Philosophically speaking, as $\mathcal{O}^2\Pi_1^1 = \Pi_3^1$, this equality reduces the “non-linear” part \mathcal{O}^2 to the “linear” part $<u_\omega$. Based on this equality, $0^{2\#}$ is defined to be the set of truth of $L_{\kappa_3}[T_2]$ for formulas of complexity slightly larger than Σ_1 , cf. Definitions 3.6-3.7. $0^{2\#}$ is essentially y_3 , defined in [16]. It is a good universal $<u_\omega\text{-}\Pi_3^1$ subset of ω . The many-one equivalence $M_1^\# \equiv_m 0^{2\#}$ is thus obtained using Neeman [25, 26]. Under AD , we have $u_k = \aleph_k$, and [15] summarizes the further structural theory at this level. The expression of $0^{2\#}$ opens the possibility of running recursion-theoretic arguments in $L_{\kappa_3}[T_2]$ that generalize those in $L_{\omega_1^{CK}}$.

The Moschovakis tree T_{2n+1} projects to the good universal Σ_{2n+2}^1 set. The structures tied to Σ_{2n+2}^1 sets are $L[T_{2n+1}]$ and its relativizations. $L[T_{2n+1}]$ is the higher level analog of L :

1. (Model theoretic representation of Σ_{2n+2}^1) $A \subseteq \mathbb{R}$ is Σ_{2n+2}^1 iff there is a Σ_1 formula φ such that $x \in A \leftrightarrow L[T_{2n+1}, x] \models \varphi(T_{2n+1}, x)$.
2. (Mouse set, [30]) $x \in \mathbb{R} \cap L[T_{2n+1}]$ iff x is Δ_{2n+2}^1 in a countable ordinal iff $x \in \mathbb{R} \cap M_{2n}^\#$.
3. (The transcendental real over $L[T_{2n+1}]$, to be proved in this series of papers) $M_{2n}^\# \equiv_m 0^{(2n+1)\#}$.
4. (Δ_{2n+3}^1 -coding of ordinals below $u_{E(2n+1)}^{(2n+1)}$) $\text{WO}_{E(2n+1)}^{(2n+1)}$ is the set of level- $(2n+1)$ sharp codes for ordinals in $u_{E(2n+1)}^{(2n+1)}$. The comparison of level- $(2n+1)$ sharp codes is Δ_{2n+3}^1 .

$0^{(2n+1)\#}$ is the theory of $L[T_{2n+1}]$ with level- $(2n+1)$ indiscernibles. The structure of the level- $(2n+1)$ indiscernibles is more complicated than their order,

as opposed to the order indiscernibles for L . The level- $(2n+1)$ indiscernibles form a tree structure, and the type realized in $L[T_{2n+1}]$ by finitely many of them depends only on the finite tree structure that relates them. This tree structure resembles the structure of measures (under AD) witnessing the homogeneity of S_{2n+1} , a tree on $\omega \times \delta_3^1$ that projects to the good universal Π_{2n+1}^1 set. We give a purely syntactical definition of $0^{(2n+1)\#}$ as the unique iterable, remarkable, level $\leq 2n$ correct level- $(2n+1)$ EM blueprint. This is the higher level analog of $0^\#$ as the unique wellfounded remarkable EM blueprint. The “iterability” part takes the form $\forall^{\mathbb{R}}(\Pi_{2n+1}^1 \rightarrow \Pi_{2n+1}^1)$, making the complexity of the whole definition Π_{2n+2}^1 . The ordinal $u_{E(2n+1)}^{(2n+1)}$ is a level- $(2n+1)$ uniform indiscernible. It will be discussed in the next paragraph. When $n = 0$, $u_{E(1)}^{(1)} = u_\omega$.

The structure tied to arbitrary Π_{2n+1}^1 sets are defined. By induction, we have level- $(2n-1)$ indiscernibles for $L_{\delta_{2n-1}^1}[T_{2n-1}]$ and the real $0^{(2n-1)\#}$. Based on the EM blueprint formulation of $0^{(2n-1)\#}$, we define the level- $2n$ Martin-Solovay tree T_{2n} . It is the higher level analog of T_2 . This is the most canonical tree that enables the correct generalization of the structural theory related to Π_{2n+1}^1 sets. The structures in our attention are $L_{\kappa_{2n+1}}[T_{2n}]$, the least admissible set over T_{2n} , and its relativizations:

1. (Model theoretic representation of Π_{2n+1}^1 , to be proved in this series of papers) $A \subseteq \mathbb{R}$ is Π_{2n+1}^1 iff there is a Σ_1 formula φ such that $x \in A \leftrightarrow L_{\kappa_{2n+1}}^x[T_{2n}, x] \models \varphi(T_{2n}, x)$.
2. (Mouse set, [30]) $x \in \mathbb{R} \cap L_{\kappa_{2n+1}}[T_{2n}]$ iff x is Δ_{2n+1}^1 in a countable ordinal iff $x \in M_{2n-1}^\#$.
3. (The transcendental real over $L_{\kappa_{2n+1}}[T_{2n}]$, to be proved in this series of papers) $M_{2n-1}^\# \equiv_m 0^{(2n)\#}$.
4. (Π_{2n+1}^1 -coding of ordinals below δ_{2n+1}^1) $\text{WO}^{(2n+1)}$ is the set of reals that naturally code a wellordering of $u_{E(2n-1)}^{(2n-1)}$. $\text{WO}^{(2n+1)}$ is Π_{2n+1}^1 .
5. The uncountable Δ_{2n+1}^1 wellordered cardinals are $(u_k : 1 \leq k \leq \omega)$, $(u_\xi^{(3)} : 1 \leq \xi \leq E(3))$, \dots , $(u_\xi^{(2n-1)} : 1 \leq \xi \leq E(2n-1))$, where $E(0) = 1$, $E(i+1) = \omega^{E(i)}$ via ordinal exponentiation.

The equivalence $M_{2n-1}^\# \equiv_m 0^{(2n)\#}$ will be based on the equality of point-classes: $\mathfrak{D}^{2n}(<\omega^2\text{-}\Pi_1^1) = <u_{E(2n-1)}^{(2n-1)}\text{-}\Pi_{2n+1}^1$. $\{u_\xi^{(2n-1)} : 1 \leq \xi \leq E(2n-1)\}$ is the set of level- $(2n-1)$ uniform indiscernibles. It is the higher level analog of the first $\omega+1$ uniform indiscernibles $\{u_n : 1 \leq n \leq \omega\}$. Under full

AD, the uncountable Δ_{2n+1}^1 wellordered cardinals enumerate all the uncountable cardinals below δ_{2n+1}^1 : $u_k = \aleph_k$ for $1 \leq k < \omega$, $u_\xi^{(2i+1)} = \aleph_{E(2i-1)+\xi}$ for $1 \leq \xi \leq E(2i+1)$. Assume AD for the moment. The equation $\delta_{2n+1}^1 = \aleph_{E(2n-1)+1}$ is originally proved by Jackson in [8, 10]. Jackson shows that every successor cardinal in the interval $[\delta_{2n-1}^1, \aleph_{E(2n-1)})$ is the image of δ_{2n-1}^1 via an ultrapower map induced by a measure on δ_{2n-1}^1 . [11] goes on to show that for a certain collection of measures μ on δ_3^1 , every description leads to a canonical function representing a cardinal modulo μ . [11, 12] compute the cofinality of the cardinals below δ_ω^1 . In this series of papers, we demonstrate the greater importance of the set theoretic *structures* tied to these cardinals over their *order type*. It is the inner model $L[T_{2n-1}]$ and its images via different ultrapower maps that give birth to the uncanny order type $E(2n-1)+1$. The level- $(2n-1)$ uniform indiscernibles $(u_\xi^{(2n-1)} : 1 \leq \xi \leq E(2n-1))$ are defined under this circumstance. Recall that the first ω uniform indiscernibles can be generated by $j^{\mu^n}(L_{\omega_1}) = L_{u_{n+1}}$, where μ^n is the n -fold product of the club measure on ω_1 ; if $1 \leq i \leq n+1$, then u_i is represented modulo μ^n by a projection map; every ordinal below u_{n+1} is in the Skolem hull of $\{x, u_1, \dots, u_n\}$ over $L[x]$ for some $x \in \mathbb{R}$. This scenario is generalized by the level- $(2n-1)$ uniform indiscernibles. As a by-product, we simplify the arguments in [8, 10–12], show in full generality that any description represents a cardinal modulo any measure on δ_{2n-1}^1 , and establish the effective version of the cofinality computations.

The whole argument is inductive. Assume AD for simplicity. In the computation of δ_{2n+1}^1 in [8, 10], the strong partition property of δ_{2n+1}^1 is proved and used inductively in the process. Our argument reproves the strong partition property of δ_{2n+1}^1 using the EM blueprint formulation of $0^{(2n+1)\#}$. The definition of $0^{(2n+1)\#}$ is based on the analysis of level- $(2n+1)$ indiscernibles, whose existence depend on the homogeneous Suslin representations of Π_{2n}^1 sets, which in turn follow from the strong partition property of δ_{2n-1}^1 . Just as the main ideas of the computation of δ_{2n+1}^1 boil down to that of δ_5^1 , all the non-trivialities are contained in the first few levels. The general inductive step is merely a technical manifestation.

A deeper insight into the interaction between inner model theory and Jackson's computation of projective ordinals in [8, 10] is the concrete information on the direct system of countable iterates of $M_{2n}^\#$. Put $n = 1$ and assume AD for simplicity sake. Put $M_{2,\infty}^- = L_{\delta_3^1}[T_3]$. We define $(c_\xi^{(3)} : \xi < \delta_3^1)$, a continuous sequence in δ_3^1 that generates the set of level-3 indiscernibles for $M_{2,\infty}^-$. Each $M_{2,\infty}^-|c_\xi^{(3)}$ is the direct limit of Π_3^1 -iterable mice whose Dodd-Jensen order is $c_\xi^{(3)}$. We define an alternative direct limit system indexed by

ordinals in u_ω which is dense in the system leading to $M_{2,\infty}^-|c_\xi^{(3)}$. The advantage of this dense subsystem is that it leads to a good coding of $M_{2,\infty}^-|c_\xi^{(3)}$ by a subset of u_ω . The indexing ordinals are represented by wellorderings on ω_1 of order type $\omega_1 + 1$ modulo measures on ω_1 arising from the strong partition property on ω_1 . Any order-preserving injection between two such wellorderings corresponds to an elementary embedding between models of this new direct limit. This injection is an isomorphism just in case its corresponding elementary embedding is essentially an iteration map, i.e., commutes with the comparison maps. The new direct system is then guided by isomorphisms between wellorderings on ω_1 of order type $\omega_1 + 1$. In this regard, the Dodd-Jensen property of mice corresponds to the simple fact that if f is an order preserving map between ordinals, then $\alpha \leq f(\alpha)$ pointwise. This observation is not surprising at all, as the Dodd-Jensen property on iterates of $0^\#$ is originated from this simple fact. This viewpoint might be a prelude to understanding the combinatorial nature of iteration trees on mice with finitely many Woodin cardinals.

A key step in computing the upper bound of δ_5^1 in [8] is the (level-3) Martin tree. For the reader familiar with the Martin tree and the purely descriptive set theoretical proof of the Kechris-Martin theorem in [9, Section 4.4], the level-1 version of the Martin tree is essentially an analysis of partially iterable sharps. The level-3 Martin tree is therefore replaced by an analysis of partially iterable level-3 sharps in this series of papers. The aforementioned new direct limit system indexed by ordinals in u_ω applies to any partially iterable mouse, so that its possibly illfounded direct limit is naturally coded by a subset of u_ω . This is yet another incidence that descriptive set theory and inner model theory are two sides of the same coin.

Apart from inner model theory, the pure computational component in [8, 10] has a major simplification. Under AD, a successor cardinal in the interval $[\delta_3^1, \aleph_{\omega^\omega})$ is represented by a measure μ on δ_3^1 and a description. The original definition of description involves a finite iteration of ultrapowers on u_ω . The “finite iteration of ultrapowers” part is now simplified to a single ultrapower. The analysis will be in the third paper of this series.

As $L_{\kappa_{2n+1}}[T_{2n}]$ is the correct structure tied to Π_{2n+1}^1 sets, it is natural to investigate its intrinsic structure. However, little is known at this very step. The closest result is on the full model $L[T_{2n}]$. The uniqueness of $L[T_{2n}]$ is proved by Hjorth [6] for $n = 1$ and Atmai [2] for general n . Here, uniqueness means that if T' is the tree of another Δ_{2n+1}^1 -scale on a good universal Π_{2n}^1 set, then $L[T_{2n}] = L[T']$. Atmai-Sargsyan [2] goes on to show that the full model $L[T_{2n}]$ is just $L[M_{2n-1,\infty}^\#]$, where $M_{2n-1,\infty}^\#$ is the direct limit of all the countable iterates of $M_{2n-1}^\#$. A test question that separates

$L_{\kappa_3}[T_2]$ from $L[T_2]$ is the inner model theoretic characterization of C_3 , the largest countable Π_3^1 set: if $x \in C_3$, must x be Δ_3^1 -equivalent to a master code in M_2 ? (cf. [33, p.13]) We will set up a good preparation for tackling this problem.

Looking higher up, the technique in this series of papers should generalize to arbitrary projective-like pointclasses in $L(\mathbb{R})$ and beyond. The descriptive set theory counterpart of larger mice should enhance our understanding of large cardinals. Typical open questions in the higher level include:

1. (cf. [1, Problem 19]) Assume AD. Let Γ be a Π_1^1 -like scaled pointclass (i.e., closed under $\forall^{\mathbb{R}}$, continuous preimages and non-self-dual) and Let $\Delta = \Gamma \cap \Gamma^\sim$, $\delta = \sup\{|\alpha| : \alpha \text{ is a prewellordering in } \Delta\}$. Is Γ closed under unions of length $< \delta$?
2. Assume AD. Let Γ , δ be as in 1. Must δ have the strong partition property?
3. Assume AD. If $\kappa \leq \lambda$ are cardinals, must $\text{cf}(\kappa^{++}) \leq \text{cf}(\lambda^{++})$?

We now switch to some immediate applications on the theory of higher level indiscernibles. Our belief is that any result in set theory that involves sharp and Silver indiscernibles should generalize to arbitrary projective levels.

Woodin [28] proves that boldface Π_{2n+1}^1 -determinacy is equivalent to “for any real x , there is an (ω, ω_1) -iterable $M_{2n}^\#(x)$ ”. The lightface scenario is tricky however. Woodin (in unpublished work for odd n) and Neeman [25, 26] proves that the existence of an ω_1 -iterable $M_n^\#$ implies boldface Π_n^1 -determinacy and lightface Π_{n+1}^1 -determinacy.

Question 1.1 (cf. [4, #9]). Assume Π_n^1 -determinacy and Π_{n+1}^1 -determinacy. Must there exist an ω_1 -iterable $M_n^\#$?

Note that the assumption of boldface Π_n^1 -determinacy in Question 1.1 is necessary, as Δ_2^1 -determinacy alone is enough to imply that there is a model of OD-determinacy (Kechris-Solovay [17]). The cases $n \in \{0, 1\}$ in Question 1.1 are solved positively by Harrington in [5] and by Woodin in [31]. The proof of the $n = 1$ case heavily relies on the theory of Silver indiscernibles for L . The theory of level-3 indiscernibles for $L_{\delta_3^1}[T_3]$ is thus involved in proving the general case when n is odd.

Theorem 1.2. *Assume Π_{2n+1}^1 -determinacy and Π_{2n+2}^1 -determinacy. Then there exists an (ω, ω_1) -iterable $M_{2n+1}^\#$.*

The proof of Theorem 1.2 will appear in further publications. The case $n \geq 2$ even in Question 1.1 remains open.

Another application is the δ -ordinal of intermediate pointclasses between Π_m^1 and Δ_{m+1}^1 . If Γ is a pointclass, $\delta(\Gamma)$ is the supremum of the lengths of Γ -prewellorderings on \mathbb{R} . $A \subseteq \mathbb{R}$ is $\Gamma_{m,n}(z)$ iff for some formula ψ we have $x \in A \leftrightarrow M_{m-1}[x, z] \models \psi(x, z, \aleph_1, \dots, \aleph_n)$. A is $\mathbf{\Gamma}_{m,n}$ iff A is $\Gamma_{m,n}(z)$ for some real z . Hjorth [7] proves that $\delta(\mathbf{\Gamma}_{1,n}) = u_{n+2}$ under Δ_2^1 -determinacy. Sargsyan [27] proves that under AD, $\sup_{n < \omega} \delta(\mathbf{\Gamma}_{2k+1,n})$ is the cardinal predecessor of δ_{2k+3}^1 . The exact value of $\delta(\mathbf{\Gamma}_{2k+1,n})$ remains unknown. Based on the theory of higher level indiscernibles, we can define the pointclasses $\Lambda_{2k+1,\xi}$ for $0 < \xi \leq E(2k+1)$. For the moment we need the notations in this series of papers. $A \subseteq \mathbb{R}$ is $\Lambda_{3,\xi+1}(z)$ iff for some level-3 tree R such that $\llbracket \emptyset \rrbracket_R = \widehat{\xi}$, for some $\mathcal{L}^{\mathbb{Q},R}$ -formula ψ we have $x \in A \leftrightarrow \ulcorner \psi \urcorner \in (x, z)^{3\#}(R)$. When ξ is a limit, $\Lambda_{3,\xi}(z) = \bigcup_{\eta < \xi} \Lambda_{3,\eta}(z)$. A is $\Lambda_{3,\xi}$ iff A is $\Lambda_{3,\xi}(z)$ for some real z .

Theorem 1.3. *Assume Δ_4^1 -determinacy and $0 < \xi < \omega^{\omega^\omega}$. If ξ is a successor ordinal, then $\delta(\Lambda_{3,\xi}) = u_{\xi+1}^{(3)}$. If ξ is a limit ordinal, then $\delta(\Lambda_{3,\xi}) = u_\xi^{(3)}$.*

The proof of Theorem 1.3 and its higher level analog will appear in further publications. The question on the value of $\delta(\mathbf{\Gamma}_{3,n})$ is then reduced to the relative position of $\mathbf{\Gamma}_{3,n}$ in the hierarchy $(\Lambda_{3,\xi} : 0 < \xi < \omega^{\omega^\omega})$. The results of this series of papers combined with Neeman [25, 26] yield the following estimate:

$$\Lambda_{3,\omega^{\omega^n}} \subseteq \mathbf{\Gamma}_{3,n} \subseteq \Lambda_{3,\omega^{\omega^{n+1}}+1}.$$

We conjecture that $\Lambda_{3,\omega^{\omega^{n+1}}} \subsetneq \mathbf{\Gamma}_{3,n} \subsetneq \Lambda_{3,\omega^{\omega^{n+1}}+1}$ and $\delta(\mathbf{\Gamma}_{3,n}) = u_{\omega^{\omega^{n+1}}+1}^{(3)}$.

We try to make this series of papers as self-contained as possible. The reader is assumed to have some minimum background knowledge in descriptive set theory and inner model theory. On the descriptive set theory side, we assume basic knowledge of determinacy, scale and its tree representation, homogeneous tree and its ultrapower representation, and at least the results of Moschovakis periodicity theorems. We will briefly recall them in Section 2. Theorem 2.1 by Becker-Kechris [3] and Kechris-Martin [13, 14] will basically be treated as a black box. Knowing its proof would help, though not necessary. On the inner model theory side, we assume basic knowledge of mice and iteration trees in the region of finitely many Woodin cardinals, especially Theorem 6.10 in [34]. The level-wise projective complexity associated to mice will be recalled in the second paper of this series. Steel's computation of $L[T_{2n+1}]$ in [32] will be treated as a black box. In particular, we require absolutely no knowledge of Jackson's analysis in [8, 10].

This series of papers is organized as follows. This paper is the first one of this series, establishing the many-one equivalence of $0^{2\#}$ and $M_1^\#$. The second

paper will define $0^{3\#}$ as the theory of $L_{\delta_3^1}[T_3]$ with its “level-3 indiscernibles” and prove the many-one equivalence of $0^{3\#}$ and $M_2^\#$. The third paper will give a Π_4^1 -axiomatization of the real $0^{3\#}$, define the level-4 Martin-Solovay tree and prove the level-4 Kechris-Martin theorem, which prepares for the induction into the next level. The fourth paper will deal with the general inductive step in the projective hierarchy.

We arrange these papers in such a way in order to minimize the background knowledge of the first and second papers. The first paper uses only sharps for reals and blackboxed Kechris-Martin. The second paper will introduce homogeneous trees with restricted complexity without too much technicalities that are enough to define $0^{3\#}$. The fine analysis associated to homogeneous trees, especially to the generalized Jackson’s analysis, will only show up in the third paper. The fourth paper will be pretty much a routine generalization of the first three.

2 Backgrounds and preliminaries

2.1 Basic descriptive set theory

Following the usual treatment in descriptive set theory, $\mathbb{R} = \omega^\omega$ is the Baire space, which is homeomorphic to the irrationals of the real line. If $A \subseteq \mathbb{R} \times X$, then $y \in \exists^\mathbb{R} A$ iff $\exists x \in \mathbb{R} (x, y) \in A$, $y \in \forall^\mathbb{R} A$ iff $\forall x \in \mathbb{R} (x, y) \in A$, $y \in \exists A$ iff Player I has a winning strategy in the game with output $A_y =_{\text{DEF}} \{x : (x, y) \in A\}$. $\exists^{n+1} A = \exists(\exists^n(A))$ when A is a subset of an appropriate product space. A pointclass is a collection of subsets of Polish spaces (typically finite products of ω and \mathbb{R}). If Γ is a pointclass, then $\exists^\mathbb{R} \Gamma = \{\exists^\mathbb{R} A : A \in \Gamma\}$, and similarly for $\forall^\mathbb{R} \Gamma$, $\exists \Gamma$, $\exists^n \Gamma$. $\Sigma_1^0 = \Sigma_0^1$ is the pointclass of open sets. $\Sigma_1^0 = \Sigma_0^1$ is the pointclass of effectively open sets. $\Pi_{n+1}^1 = \forall^\mathbb{R} \Sigma_n^1$, $\Sigma_{n+1}^1 = \exists^\mathbb{R} \Pi_n^1$, $\Pi_{n+1}^1 = \forall^\mathbb{R} \Sigma_n^1$, $\Sigma_{n+1}^1 = \exists^\mathbb{R} \Pi_n^1$.

If α is an ordinal and $A \subseteq \alpha \times X$, then

$$x \in \text{Diff } A \leftrightarrow \exists i < \alpha (i \text{ is odd} \wedge \forall j < i ((j, x) \in A) \wedge (i, x) \notin A).$$

If $\alpha < \omega_1^{CK}$ then $A \subseteq X$ is α - Π_1^1 iff $A = \text{Diff } B$ for some Π_1^1 $B \subseteq \alpha \times X$. A is $<\alpha$ - Π_1^1 iff A is β - Π_1^1 for some $\beta < \alpha$. Martin [22] proves that Π_1^1 -determinacy implies $<\omega^2$ - Π_1^1 -determinacy.

A tree on X is a subset of $X^{<\omega}$ closed under initial segments. If T is a tree on X , $[T]$ is the set of infinite branches of T , i.e., $x \in T$ iff $\forall n (x \upharpoonright n) \in T$. If T is a tree on λ , λ is an ordinal, $[T] \neq \emptyset$, the leftmost branch is $x \in [T]$ such that for any $y \in [T]$, $(x(0), x(1), \dots)$ is lexicographically smaller than or equal to $(y(0), y(1), \dots)$. In addition, if $x \in [T]$ and for any $y \in [T]$ we have

$\forall n \ x(n) \leq y(n)$, then x is the honest leftmost branch of T . A tree T on $\omega \times X$ is identified with a subset of $\omega^{<\omega} \times X^{<\omega}$ consisting of (s, t) so that $\text{lh}(s) = \text{lh}(t)$ and $((s(i), t(i)))_{i < \text{lh}(s)} \in T$. If T is a tree on $\omega \times X$, $[T] \subseteq \omega^{<\omega} \times X^{<\omega}$ is the set of infinite branches of T . $p[T] = \{x : \exists y \ (x, y) \in [T]\}$ is the projection of T . If T is a tree on $\omega \times \lambda$ and $p[T] \neq \emptyset$, then x is the leftmost real of T iff $\exists \vec{\alpha} \ (x, \vec{\alpha})$ is the leftmost branch of T .

2.2 The Martin-Solovay tree

We assume Π_1^1 -determinacy. This is equivalent to $\forall x \in \mathbb{R} \ (x^\# \text{ exists})$ by Martin [21] and Harrington [5].

γ is a uniform indiscernible iff for every $x \in \mathbb{R}$, γ is an x -indiscernible. The uniform indiscernibles form a club in Ord , which are listed u_1, u_2, \dots in the increasing order. In particular, $u_1 = \omega_1$ and $u_\omega = \sup_{n < \omega} u_n$.

The set $\{x^\# : x \in \mathbb{R}\}$ is Π_2^1 . $\text{WO} = \text{WO}_1$ is the set of codes for countable ordinals. For $1 \leq m < \omega$, WO_{m+1} is the set of $\langle \ulcorner \tau \urcorner, x^\# \rangle$ where τ is an $(m+1)$ -ary Skolem term for an ordinal in the language of set theory and $x \in \mathbb{R}$. The ordinal coded by $w = \langle \ulcorner \tau \urcorner, x^\# \rangle \in \text{WO}_{m+1}$ is

$$|w| = \tau^{L[x]}(x, u_1, \dots, u_m).$$

Every ordinal in u_{m+1} is of the form $|w|$ for some $w \in \text{WO}_{m+1}$. For each $1 \leq m < \omega$,

$$\{\tau^{L[x]}(x, u_m) : \langle \ulcorner \tau \urcorner, x^\# \rangle \in \text{WO}_2\}$$

is a cofinal subset of u_{m+1} . $\text{WO}_\omega = \bigcup_{1 \leq m < \omega} \text{WO}_m$. WO is Π_1^1 , and WO_{m+1} is Π_2^1 for $1 \leq m < \omega$. If $\sigma : \{1, \dots, m\} \rightarrow \{1, \dots, n\}$ is order preserving, then define

$$j^\sigma(\tau^{L[x]}(x, u_1, \dots, u_m)) = \tau^{L[x]}(x, u_{\sigma(1)}, \dots, u_{\sigma(m)}).$$

If \mathcal{X} is a Polish space, $A \subseteq \mathcal{X} \times u_\omega$ and Γ is a pointclass, say that A is in Γ iff

$$A^* = \{(x, w) : x \in \text{WO}_\omega \wedge (x, |w|) \in A\}$$

is in Γ . Γ acting on product spaces are similarly defined.

T_2 , defined in [3, 14, 20], refers to the Martin-Solovay tree on $\omega \times u_\omega$ that projects to $\{x^\# : x \in \mathbb{R}\}$, giving the scale

$$\varphi_{\ulcorner \tau \urcorner}(x^\#) = \tau^{L[x]}(x, u_1, \dots, u_{k_\tau}),$$

where $\ulcorner \tau \urcorner$ is the Gödel number of τ , τ is $k_\tau + 1$ -ary. T_2 is a Δ_3^1 subset of $(\omega \times u_\omega)^{<\omega}$. From T_2 one can compute a tree \widehat{T}_2 on $\omega \times u_\omega$ that projects to a good universal Π_2^1 set.

To conclude this section, we define the Martin-Solovay tree T_2 projecting to $\{x^\# : x \in \mathbb{R}\}$ and its variant \widehat{T}_2 projecting to a good universal Π_2^1 set. This formulation of T_2 and \widehat{T}_2 will generalize to the higher levels in this series of papers. Let $T \subseteq 2^{<\omega}$ be a recursive tree such that $[T]$ is the set of remarkable EM blueprints over some real. Here we have fixed in advance an effective Gödel coding of first order formulas in the language $\{\underline{\in}, \underline{x}, \underline{c}_n : n < \omega\}$, so that an infinite string $x \in 2^\omega$ represents the theory $\{\varphi : x_{\ulcorner \varphi \urcorner} = 0\}$. Fix an effective list of Skolem terms $(\tau_k)_{k < \omega}$ in the language of set theory, where τ_k is $f(k) + 1$ -ary, f is effective. T_2 is defined as a tree on $2 \times u_\omega$ where

$$(s, (\alpha_0, \dots, \alpha_{n-1})) \in T_2$$

iff $s \in T$, $\text{lh}(s) = n$, and for any $k, l < n$, for any order preserving $\sigma : \{1, \dots, f(k)\} \rightarrow \{1, \dots, f(l)\}$,

1. if “ $\tau_k(\underline{x}, \underline{c}_{\sigma(1)}, \dots, \underline{c}_{\sigma(f(k))}) = \tau_l(\underline{x}, \underline{c}_1, \dots, \underline{c}_{f(l)})$ ” is true in s , then $j^\sigma(\alpha_k) = \alpha_l$;
2. if “ $\tau_k(\underline{x}, \underline{c}_{\sigma(1)}, \dots, \underline{c}_{\sigma(f(k))}) < \tau_l(\underline{x}, \underline{c}_1, \dots, \underline{c}_{f(l)})$ ” is true in s , then $j^\sigma(\alpha_k) < \alpha_l$;

In essence, the second coordinate of T_2 attempts to verify the wellfoundedness of the EM blueprint coded in the first coordinate. From T_2 we compute \widehat{T}_2 , a tree on $\omega \times (\omega \times u_\omega)$ that projects to a good universal Π_2^1 set. By Shoenfield absoluteness, if $\varphi(v)$ is a Π_2^1 formula, effectively from $\ulcorner \varphi \urcorner$ we can compute a unary Skolem term $\tau_{\ulcorner \varphi \urcorner}$ such that $\tau_{\ulcorner \varphi \urcorner}^{L[x]}(x) = 0$ iff $\varphi(x)$ holds. Define $(\ulcorner \varphi \urcorner^\frown(v), (s, \vec{\alpha})) \in \widehat{T}_2$ iff $(s, \vec{\alpha}) \in T_2$ and

1. if “ $\underline{x}(m) = n$ ” is true in s , then $v(m) = n$;
2. “ $\tau_{\ulcorner \varphi \urcorner}(\underline{x}) \neq 0$ ” is not true in s .

So $p[\widehat{T}_2] = \{\ulcorner \varphi \urcorner^\frown(x) : \varphi(x)\}$.

2.3 Q-theory

From now on until the end of this paper, we assume Δ_2^1 -determinacy. By Kechris-Woodin [18], $\mathfrak{D}(<\omega^2\text{-}\Pi_1^1)$ -determinacy follows. By Neeman [25, 26] and Woodin [19, 31], this is also equivalent to “for every $x \in \mathbb{R}$, there is an (ω, ω_1) -iterable $M_1^\#(x)$ ”.

For $x \in \mathbb{R}$, $L_{\kappa_3^x}[T_2, x]$ is the minimum admissible set containing (T_2, x) . κ_3^x is the higher level analog of ω_1^x , the least x -admissible. The fact that the \widehat{T}_2 projects to a good universal Π_2^1 set implies for every Π_3^1 set of reals A , there is

a Σ_1 -formula φ such that $x \in A$ iff $L_{\kappa_3^x}[T_2, x] \models \varphi(T_2, x)$; φ can be effectively computed from the definition of A . Becker-Kechris in [3] strengthens this fact by allowing a parameter in u_ω . The converse direction is shown by Kechris-Martin in [13, 14]. The back-and-forth conversion is concluded in [3].

Theorem 2.1 (Becker-Kechris-Martin). *Assume Δ_2^1 -determinacy. Then for each $A \subseteq u_\omega \times \mathbb{R}$, the following are equivalent.*

1. A is Π_3^1 .
2. There is a Σ_1 formula φ such that $(\alpha, x) \in A$ iff $L_{\kappa_3^x}[T_2, x] \models \varphi(T_2, \alpha, x)$.

The conversions between the Π_3^1 definition of A and the Σ_1 -formula φ are effective.

3 The equivalence of $x^{2\#}$ and $M_1^\#(x)$

Definition 3.1. Suppose $\mathcal{X} = \omega^k \times \mathbb{R}^l$ is a product space. Suppose x is a real and $\beta \leq u_\omega$. A subset $A \subseteq \mathcal{X}$ is β - $\Pi_3^1(x)$ iff there is a $\Pi_3^1(x)$ set $B \subseteq u_\omega \times \mathcal{X}$ such that $A = \text{Diff } B$. A is β - Π_3^1 iff A is β - $\Pi_3^1(0)$. A is β - $\mathbf{\Pi}_3^1$ iff A is β - $\Pi_3^1(x)$ for some real x .

By Theorem 2.1, when β is a limit ordinal, $A \subseteq \mathcal{X}$ is β - $\Pi_3^1(x)$ iff there is a pair of Σ_1 -formulas (φ, ψ) such that

$$(\vec{n}, \vec{y}) = (n_1, \dots, n_k, y_1, \dots, y_l) \in A$$

iff

$$L_{\kappa_3^{x, \vec{y}}}[T_2, x, \vec{y}] \models \exists \alpha < \beta (\forall \eta < \alpha \varphi(\eta, \vec{n}, \vec{y}, T_2, x) \wedge \neg \psi(\alpha, \vec{n}, \vec{y}, T_2, x)).$$

Lemma 3.2. *Assume Δ_2^1 -determinacy. Suppose n, m are positive integers. If A is $(u_n)^m$ - $\Pi_3^1(x)$, then A is $\mathcal{O}^2(\omega n$ - $\Pi_1^1(x))$.*

Proof. Without loss of generality, we assume $x = 0$, $m = 1$, and $A \subseteq \mathbb{R}$. Let B be a Π_3^1 subset of $u_\omega \times \mathbb{R}$ such that $A = \text{Diff } B$. Let $B^* = \{(w, y) : (|w|, y) \in B\}$ be the Π_3^1 code set of B . Let $C \subseteq \mathbb{R}^3$ be a Σ_2^1 set such that

$$(w, y) \in B^* \leftrightarrow \forall r (w, y, r) \in C.$$

Consider the game $H(y)$, where I produces $w, r \in \mathbb{R}$, II produces $w', r' \in \mathbb{R}$. The game is won by I iff both of the following hold:

1. $w \in \text{WO}_n$, $|w|$ is odd, and $(w, y, r) \notin C$.

2. If $w' \in \text{WO}_n$, $|w'|$ is even, and $(w', y, r') \notin C$, then $|w| < |w'|$.

Therefore, $y \in A$ iff I has a winning strategy in $H(y)$.

Since $L[y, w, r, w', r']$ is Σ_2^1 -absolute, and since the relation $|w| \leq |w'|$ for $w, w' \in \text{WO}_n$ is definable over $L[y, w, r, w', r']$ from parameters u_1, \dots, u_{n-1} , the payoff set of the game $H(y)$ can be expressed as a first order statement over $L[y, \cdot]$ from parameters u_1, \dots, u_{n-1} . That is, there is a formula θ such that an infinite run

$$(w, r, w', r')$$

is won by I iff

$$L[y, w, r, w', r'] \models \theta(y, w, r, w', r', u_1, \dots, u_{n-1}).$$

It follows by Martin [22] that the payoff set of $H(y)$ is $\mathcal{D}(\omega n\text{-}\Pi_1^1(y))$, uniformly in y , hence determined. Hence A is in $\mathcal{D}^2(\omega n\text{-}\Pi_1^1)$. \square

Lemma 3.3. Assume Δ_2^1 -determinacy. Let $n < \omega$. If A is $\mathcal{D}^2(\omega n\text{-}\Pi_1^1(x))$, then A is $u_{n+2}\text{-}\Pi_3^1(x)$.

Proof. Without loss of generality, assume $x = 0$ and $A \subseteq \mathbb{R}$. We produce an effective transformation from a $\mathcal{D}^2(\omega n\text{-}\Pi_1^1)$ definition to the desired $u_{n+2}\text{-}\Pi_3^1$ definition. By Martin [22], if $(y, r) \in \mathbb{R}^2$, $C \subseteq \mathbb{R}$ is $\omega n\text{-}\Pi_1^1(y, r)$, then there is a formula φ such that Player I has a winning strategy in $G(C)$ iff

$$L[y, r] \models \varphi(y, r, u_1, \dots, u_n).$$

The transform from the $\omega n\text{-}\Pi_1^1(y, r)$ definition of C to φ is uniform, independent of (y, r) . Suppose $A = \mathcal{D}B$, where $B \subseteq \mathbb{R}^2$ is $\mathcal{D}(\omega n\text{-}\Pi_1^1)$. Suppose φ is a formula such that

$$(y, r) \in B \leftrightarrow L[y, r] \models \varphi(y, r, u_1, \dots, u_n).$$

To establish a $u_{n+2}\text{-}\Pi_3^1$ definition of A , we have to decide which player has a winning strategy in $G(B_y)$, for $y \in \mathbb{R}$. For ordinals $\xi_1 < \dots < \xi_n < \eta < \omega_1$, we say that M is a Kechris-Woodin non-determined set with respect to $(y, \xi_1, \dots, \xi_n, \eta)$ iff

1. M is a countable subset of \mathbb{R} ;
2. M is closed under join and Turing reducibility;
3. $\forall \sigma \in M \exists v \in M L_\eta[y, \sigma \otimes v] \models \neg \varphi(y, \sigma \otimes v, \xi_1, \dots, \xi_n)$;
4. $\forall \sigma \in M \exists v \in M L_\eta[y, v \otimes \sigma] \models \varphi(y, v \otimes \sigma, \xi_1, \dots, \xi_n)$.

In clause 3, “ $\forall \sigma \in M$ ” is quantifying over all strategies σ for Player I that is coded in some member of M ; $\sigma * v$ is Player I’s response to v according to σ , and $\sigma \otimes v = (\sigma * v) \oplus v$ is the combined infinite run. Similarly for clause 4, roles between two players being exchanged. Say that z is $(y, \xi_1, \dots, \xi_n, \eta)$ -stable iff z is not contained in any Kechris-Woodin non-determined set with respect to $(y, \xi_1, \dots, \xi_n, \eta)$. z is y -stable iff z is $(y, \xi_1, \dots, \xi_n, \eta)$ -stable for all $\xi_1 < \dots < \xi_n < \eta < \omega_1$. The set of (y, z) such that z is y -stable is Π_2^1 . By the proof of Kechris-Woodin [18], for all $y \in \mathbb{R}$, there is $z \in \mathbb{R}$ which is y -stable.

Note that if z is $(y, \xi_1, \dots, \xi_n, \eta)$ -stable and $z \leq_T z'$, then z' is $(y, \xi_1, \dots, \xi_n, \eta)$ -stable. Let $<_y^{\xi_1, \dots, \xi_n, \eta}$ be the following wellfounded relation on the set of z which is $(y, \xi_1, \dots, \xi_n, \eta)$ -stable:

$$\begin{aligned} z' <_y^{\xi_1, \dots, \xi_n, \eta} z &\leftrightarrow z \text{ is } (y, \xi_1, \dots, \xi_n, \eta)\text{-stable} \wedge z \leq_T z' \wedge \\ &\quad \forall \sigma \leq_T z \exists v \leq_T z' L_\eta[y, \sigma \otimes v] \models \neg \varphi(y, \sigma \otimes v, \xi_1, \dots, \xi_n) \\ &\quad \forall \sigma \leq_T z \exists v \leq_T z' L_\eta[y, v \otimes \sigma] \models \varphi(y, v \otimes \sigma, \xi_1, \dots, \xi_n). \end{aligned}$$

Wellfoundedness of $<_y^{\xi_1, \dots, \xi_n, \eta}$ follows from the definition of $(y, \xi_1, \dots, \xi_n, \eta)$ -stability. If z is $(y, \xi_1, \dots, \xi_n, \eta)$ -stable, then $<_y^{\xi_1, \dots, \xi_n, \eta} \{z' : z' <_y^{\xi_1, \dots, \xi_n, \eta} z\}$ is a Σ_1^1 wellfounded relation in parameters (y, z) and the code of $(\xi_1, \dots, \xi_n, \eta)$, hence has rank $< \omega_1$ by Kunen-Martin. If z is y -stable, let f_y^z be the function that sends $(\xi_1, \dots, \xi_n, \eta)$ to the rank of z in $<_y^{\xi_1, \dots, \xi_n, \eta}$. Then f_y^z is a function into ω_1 . By Σ_2^1 -absoluteness between V and $L[y, z]^{\text{Coll}(\omega, \eta)}$, we can see $f_y^z \in L[y, z]$. Furthermore, f_y^z is definable over $L[y, z]$ in a uniform way, so there is a $\{\subseteq\}$ -Skolem term τ such that for all $(y, z) \in \mathbb{R}^2$, if z is y -stable, then

$$f_y^z(\xi_1, \dots, \xi_n, \eta) = \tau^{L[y, z]}(y, z, \xi_1, \dots, \xi_n, \eta).$$

Let

$$\beta_y^z = \tau^{L[y, z]}(y, z, u_1, \dots, u_{n+1}).$$

The function

$$(y, z) \mapsto \beta_y^z$$

is Δ_3^1 in the sharp codes. We say that z is y -ultrastable iff z is y -stable and $\beta_y^z = \min\{\beta_y^w : w \text{ is } y\text{-stable}\}$.

Claim 3.4. *If z is y -ultrastable, then there is $\sigma \leq_T z$ such that σ is a winning strategy for either of the players in $G(B_y)$.*

Proof of Claim 3.4. Suppose otherwise. For any $\sigma \leq_T z$ which is a strategy for either player, pick w_σ which defeats σ in $G(B_y)$. Let w be a real

coding $\{(\sigma, w_\sigma) : \sigma \leq_T z\}$. By an indiscernability argument, for any (y, w) -indiscernibles $\xi_1 < \dots < \xi_n < \eta$, for any $\sigma \leq_T z$, if σ is a strategy for Player I, then

$$L_\eta[y, \sigma \otimes w_\sigma] \models \neg \varphi(y, \sigma \otimes w_\sigma, \xi_1, \dots, \xi_n);$$

if σ is a strategy for Player II, then

$$L_\eta[y, w_\sigma \otimes \sigma] \models \varphi(y, w_\sigma \otimes \sigma, \xi_1, \dots, \xi_n).$$

This exactly means

$$w <_{y, \xi_1, \dots, \xi_n, \eta} z,$$

and hence

$$f_y^w(\xi_1, \dots, \xi_n, \eta) < f_y^z(\xi_1, \dots, \xi_n, \eta).$$

Since z is y -stable and $z \leq_T w$, w is y -stable. Therefore, β_y^w is defined and $\beta_y^w < \beta_y^z$, contradicting to y -ultrastableness of z . \square

From Claim 3.4, if Player I (or II) has a winning strategy in $G(B_y)$, then for any y -ultrastable z , there is a winning strategy for Player I (or II) in $G(B_y)$ which is Turing reducible to z . Therefore, Player I has a winning strategy in $G(B_y)$ iff there is $\delta < u_{n+2}$ such that

$$\exists z (z \text{ is } y\text{-stable} \wedge \beta_y^z = \delta) \tag{1}$$

and

$$\begin{aligned} \forall \gamma \leq \delta \forall z ((z \text{ is } y\text{-stable} \wedge \beta_y^z = \gamma) \rightarrow \\ \exists \sigma \leq_T z (\sigma \text{ is a winning strategy for I in } G(B_y))) \end{aligned} \tag{2}$$

Note that in (1),

$$\{(\delta, y) : \exists z (z \text{ is } y\text{-stable} \wedge \beta_y^z = \delta)\}$$

is a Σ_3^1 subset of $u_\omega \times \mathbb{R}$, and in (2),

$$\begin{aligned} \{(\gamma, y) : \forall z ((z \text{ is } y\text{-stable} \wedge \beta_y^z = \gamma) \rightarrow \\ \exists \sigma \leq_T z (\sigma \text{ is a winning strategy for I in } G(B_y)))\} \end{aligned}$$

is a Π_3^1 subset of $u_\omega \times \mathbb{R}$. So $\exists \delta < u_{n+2}((1) \wedge (2))$ is a u_{n+2} - Π_3^1 definition of A . \square

Lemma 3.2 and Lemma 3.3 are concluded in a simple equality between pointclasses.

Theorem 3.5. *Assume Δ_2^1 -determinacy. Then for $x \in \mathbb{R}$,*

$$\mathfrak{O}^2(<\omega^2\text{-}\Pi_1^1(x)) = <u_\omega\text{-}\Pi_3^1(x).$$

Definition 3.6.

$$\mathcal{O}^{T_2, x} = \{(\ulcorner\varphi^\top, \alpha) : \varphi \text{ is a } \Sigma_1\text{-formula, } \alpha < u_\omega, L_{\kappa_3^x}[T_2, x] \models \varphi(T_2, x, \alpha)\}.$$

$\mathcal{O}^{T_2, x}$ is the u_ω -version of Kleene's \mathcal{O} relative to (T_2, x) . It is called \mathcal{P}_3^x in [14].

Definition 3.7.

$$x_n^{2\#} = \{(\ulcorner\varphi^\top, \ulcorner\psi^\top) : \exists \alpha < u_n((\ulcorner\varphi^\top, \alpha) \notin \mathcal{O}^{T_2, x} \wedge \forall \eta < \alpha(\ulcorner\psi^\top, \eta) \in \mathcal{O}^{T_2, x})\}.$$

$$x^{2\#} = \{(n, \ulcorner\varphi^\top, \ulcorner\psi^\top) : n < \omega \wedge (\ulcorner\varphi^\top, \ulcorner\psi^\top) \in x_n^{2\#}\}.$$

$\mathcal{O}^{T_2, x}$ splits into ω many parts $(\mathcal{O}^{T_2, x} \cap (\omega \times u_n))_{n < \omega}$. Each part is squeezed into a real $x_n^{2\#}$ by applying the difference operator on its second coordinate. The join of $(x_n^{2\#})_{n < \omega}$ is $x^{2\#}$. In particular, $x_0^{2\#}$ is Turing equivalent to the good universal Π_3^1 real, which is called the Δ_3^1 -jump of x . Each $x_n^{2\#}$ belongs to $L_{\kappa_3^x}[T_2, x]$, but $x^{2\#} \notin L_{\kappa_3^x}[T_2, x]$. The distinction between $x_0^{2\#}$ and $x^{2\#}$ does not have a lower level analog.

The expression of $0^{2\#}$ generalizes Kleene's \mathcal{O} to the higher level. Note that the transformations between $\mathfrak{O}^2(<\omega^2\text{-}\Pi_1^1(x))$ and $<u_\omega\text{-}\Pi_3^1(x)$ definitions in Theorem 3.5 are uniform. Applying Theorem 3.5 to the space $\mathcal{X} = \omega$, in combination with Theorem 2.1, we get the equivalence between $x^{2\#}$ and $M_1^\#(x)$.

Theorem 3.8. *Assume Δ_2^1 -determinacy. Then $x^{2\#}$ is many-one equivalent to $M_1^\#(x)$, the many-one reductions being independent of x .*

$0^{2\#}$ is essentially a fancy way of expressing y_3 , the leftmost real of \widehat{T}_2 which is used in the standard uniformization argument. T_2 and y_3 are used in [20] to show that every nonempty Σ_3^1 set of reals contains a member which is recursive in y_3 , or in our terminology, recursive in $0^{2\#}$. Basis theorems can also be proved with inner model theory. If $M_{2n-1}^\#$ exists, then every nonempty Σ_{2n+1}^1 set of reals contains a member recursive in $M_{2n-1}^\#$ (cf. [30, 34]). At higher levels, the leftmost real basis arguments are investigated in [16]. It is shown by Harrington (modulo Neeman [25, 26]) that under Δ_{2n}^1 -determinacy, there is a Δ_{2n+1}^1 -scale on a Δ_{2n+1}^1 set whose leftmost real y_{2n+1} is Δ_{2n+1}^1 -equivalent to $M_{2n-1}^\#$ and such that every nonempty Σ_{2n+1}^1 set contains a real recursive in y_{2n+1} . It is asked in [16, Conjecture 11.2] whether y_{2n+1} is Turing equivalent to $M_{2n-1}^\#$. Theorem 3.8 solves this conjecture in the $n = 1$ case in an effective manner.

Acknowledgements

The breakthrough ideas of this series of papers were obtained during the AIM workshop on Descriptive inner model theory, held in Palo Alto, and the Conference on Descriptive Inner Model Theory, held in Berkeley, both in June, 2014. The author greatly benefited from conversations with Rachid Atmai and Steve Jackson that took place in these two conferences. The final phase of this paper was completed whilst the author was a visiting fellow at the Isaac Newton Institute for Mathematical Sciences in the programme ‘Mathematical, Foundational and Computational Aspects of the Higher Infinite’ (HIF) in August and September, 2015 funded by NSF Career grant DMS-1352034 and EPSRC grant EP/K032208/1.

References

- [1] A list of problems that was produced at the AIM meeting “recent advances in core model theory”, organized by E. Schimmerling and J. Steel in Dec 13-17, 2004, available at <http://www.math.uni-muenster.de/u/rds>.
- [2] Rachid Atmai. *Contributions to descriptive set theory*. PhD thesis, University of North Texas, 2015.
- [3] Howard S. Becker and Alexander S. Kechris. Sets of ordinals constructible from trees and the third Victoria Delfino problem. In *Axiomatic set theory (Boulder, Colo., 1983)*, volume 31 of *Contemp. Math.*, pages 13–29. Amer. Math. Soc., Providence, RI, 1984.
- [4] Andrés Eduardo Caicedo and Benedikt Löwe. The fourteen Victoria Delfino problems and their status in the year 2015. submitted to The Cabal Seminar. Volume IV: Large Cardinals, Determinacy and Other Topics.
- [5] Leo Harrington. Analytic determinacy and 0^\sharp . *J. Symbolic Logic*, 43(4):685–693, 1978.
- [6] Greg Hjorth. Variations of the Martin-Solovay tree. *J. Symbolic Logic*, 61(1):40–51, 1996.
- [7] Greg Hjorth. A boundedness lemma for iterations. *J. Symbolic Logic*, 66(3):1058–1072, 2001.

- [8] Steve Jackson. A computation of δ_5^1 . *Mem. Amer. Math. Soc.*, 140(670):viii+94, 1999.
- [9] Steve Jackson. Structural consequences of AD. In *Handbook of set theory. Vols. 1, 2, 3*, pages 1753–1876. Springer, Dordrecht, 2010.
- [10] Steve Jackson. AD and the projective ordinals. In *Wadge degrees and projective ordinals. The Cabal Seminar. Volume II*, volume 37 of *Lect. Notes Log.*, pages 364–483. Association for Symbolic Logic, La Jolla, CA; Cambridge University Press, Cambridge, 2012.
- [11] Steve Jackson and Farid Khafizov. Descriptions and cardinals below δ_5^1 . To appear in the *Journal of Symbolic Logic*.
- [12] Steve Jackson and Benedikt Löwe. Canonical measure assignments. *J. Symbolic Logic*, 78(2):403–424, 2013.
- [13] A. S. Kechris and D. A. Martin. On the theory of Π_3^1 sets of reals. *Bull. Amer. Math. Soc.*, 84(1):149–151, 1978.
- [14] A. S. Kechris and D. A. Martin. On the theory of Π_3^1 sets of reals, II. In *Ordinal Definability and Recursion Theory. The Cabal Seminar. Volume III*, volume 43 of *Lect. Notes Log.*, pages 200–219. Cambridge University Press, Cambridge, 2016.
- [15] Alexander S. Kechris. AD and projective ordinals. In *Cabal Seminar 76–77 (Proc. Caltech-UCLA Logic Sem., 1976–77)*, volume 689 of *Lecture Notes in Math.*, pages 91–132. Springer, Berlin, 1978.
- [16] Alexander S. Kechris, Donald A. Martin, and Robert M. Solovay. Introduction to Q -theory. In *Ordinal Definability and Recursion Theory. The Cabal Seminar. Volume III*, volume 43 of *Lect. Notes Log.*, pages 126–199. Cambridge University Press, Cambridge, 2016.
- [17] Alexander S. Kechris and Robert M. Solovay. On the relative consistency strength of determinacy hypotheses. *Trans. Amer. Math. Soc.*, 290(1):179–211, 1985.
- [18] Alexander S. Kechris and W. Hugh Woodin. The equivalence of partition properties and determinacy. In *Games, scales, and Suslin cardinals. The Cabal Seminar. Vol. I*, volume 31 of *Lect. Notes Log.*, pages 355–378. Association for Symbolic Logic, Chicago, IL; Cambridge University Press, Cambridge, 2008.

- [19] Peter Koellner and W. Hugh Woodin. Large cardinals from determinacy. In *Handbook of set theory. Vols. 1, 2, 3*, pages 1951–2119. Springer, Dordrecht, 2010.
- [20] D. A. Martin and R. M. Solovay. A basis theorem for Σ_3^1 sets of reals. *Ann. of Math. (2)*, 89:138–159, 1969.
- [21] Donald A. Martin. Measurable cardinals and analytic games. *Fund. Math.*, 66:287–291, 1969/1970.
- [22] Donald A. Martin. The largest countable this, that, and the other. In *Games, scales, and Suslin cardinals. The Cabal Seminar. Vol. I*, volume 31 of *Lect. Notes Log.*, pages 121–129. Association for Symbolic Logic, Chicago, IL; Cambridge University Press, Cambridge, 2008.
- [23] Donald A. Martin and John R. Steel. A proof of projective determinacy. *J. Amer. Math. Soc.*, 2(1):71–125, 1989.
- [24] Yiannis N. Moschovakis. *Descriptive set theory*, volume 155 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, second edition, 2009.
- [25] Itay Neeman. Optimal proofs of determinacy. *Bull. Symbolic Logic*, 1(3):327–339, 1995.
- [26] Itay Neeman. Optimal proofs of determinacy. II. *J. Math. Log.*, 2(2):227–258, 2002.
- [27] Grigor Sargsyan. On the prewellorderings associated with the directed systems of mice. *J. Symbolic Logic*, 78(3):735–763, 2013.
- [28] Ralf Schindler, Sandra Uhlenbrock, and Hugh Woodin. Mice with finitely many Woodin cardinals from optimal determinacy hypotheses, in preparation.
- [29] Robert M. Solovay. A Δ_3^1 coding of the subsets of ω_ω . In *Wadge degrees and projective ordinals. The Cabal Seminar. Volume II*, volume 37 of *Lect. Notes Log.*, pages 346–363. Association for Symbolic Logic, La Jolla, CA; Cambridge University Press, Cambridge, 2012.
- [30] J. R. Steel. Projectively well-ordered inner models. *Ann. Pure Appl. Logic*, 74(1):77–104, 1995.

- [31] J. R. Steel and W. Hugh Woodin. HOD as a core model. In *Ordinal Definability and Recursion Theory. The Cabal Seminar. Volume III*, volume 43 of *Lect. Notes Log.*, pages 257–346. Cambridge University Press, Cambridge, 2016.
- [32] John R. Steel. $\text{HOD}^{L(\mathbb{R})}$ is a core model below Θ . *Bull. Symbolic Logic*, 1(1):75–84, 1995.
- [33] John R. Steel. Games and scales. Introduction to Part I. In *Games, scales, and Suslin cardinals. The Cabal Seminar. Vol. I*, volume 31 of *Lect. Notes Log.*, pages 3–27. Assoc. Symbol. Logic, Chicago, IL, 2008.
- [34] John R. Steel. An outline of inner model theory. In *Handbook of set theory. Vols. 1, 2, 3*, pages 1595–1684. Springer, Dordrecht, 2010.