

Supersimplicity and countable reducts of a unidimensional hypersimple theory

Ziv Shami

March 7, 2022

Abstract

We show that a hypersimple unidimensional theory that has a club of reducts, in the partial order of all countable reducts, that are coordinatized in finite rank, is supersimple.

1 Introduction

In this paper we suggest an approach to the problem on supersimplicity of unidimensional hypersimple theories. The problem has been answered in the affirmative in the following cases. In [H], for any stable theory, in [S1] for any countable theory (this improved an earlier result for the case of a countable theory with the wfc [P]) and in [S2] it has been proved for any (possibly uncountable) non-s-essentially 1-based theory (roughly, a theory that is far from being 1-based).

It is easy to see that supersimplicity of a theory is determined by (the supersimplicity of) the family of its countable reducts. Therefore, it is natural to try and reflect properties of the given unidimensional hypersimple theory to countable reducts. Clearly, unidimensionality is not preserved under reducts. On the other hand, easily any unidimensional hypersimple theory is coordinatized in finite rank (see Definition 5.1). In this paper we show that supersimplicity of any (possibly uncountable) unidimensional hypersimple theory follows from coordinatization in finite rank of sufficiently many countable reducts of it.

We thank Ehud Hrushovski for discussions on this topic and for allowing us to include a remark of him about elimination of hyperimaginaries in

reducts (section 3). We will assume basic knowledge of simple theories as in [K1],[KP],[HKP]. A good text book on simple theories that covers much more is [W]. The notations are standard, and throughout the paper we work in a large saturated model \mathcal{C} of a complete first-order theory T in a language L .

2 Preliminaries

In this section T is assumed to be simple. We quote several known facts that we will apply.

2.1 Almost internality, analyzability and unidimensionality

In this subsection we work in \mathcal{C} with hyperimaginaries unless otherwise stated; if T is hypersimple (i.e. simple and eliminates hyperimaginaries) and we work in \mathcal{C}^{eq} we get equivalent definitions.

In this subsection, \mathcal{P} denotes an A -invariant family of partial types and p a partial type over A . We say that p is (almost-) \mathcal{P} -internal for every realization a of p there exists b with $a \downarrow_A b$ such that for some tuple c of realizations of partial types in \mathcal{P} over Ab we have $a \in dcl(b, c)$ (respectively, $a \in acl(b, c)$). We say that p is analyzable in \mathcal{P} if for any $a \models p$ there exists a sequence $I = \langle a_i \mid i \leq \alpha \rangle$ such that $a_\alpha = a$ and $tp(a_i / \{a_j \mid j < i\} \cup A)$ is almost \mathcal{P} -internal for every $i \leq \alpha$.

First, the following fact is straightforward.

Fact 2.1 1) Assume $tp(a_i/A)$ are (almost) \mathcal{P} -internal for $i < \alpha$. Then $tp(\langle a_i \mid i < \alpha \rangle/A)$ is (respectively, almost) \mathcal{P} -internal. Thus, if $tp(a_i/A)$ are analyzable in \mathcal{P} for $i < \alpha$. Then $tp(\langle a_i \mid i < \alpha \rangle/A)$ is analyzable in \mathcal{P} .
 2) If $tp(a/A)$ almost \mathcal{P} -internal, so is $tp(a/B)$ for any set $B \supseteq A$.

T is said to be *unidimensional* if whenever p and q are complete non-algebraic types, p, q are non-orthogonal.

We will also need the following easy Fact.

Fact 2.2 *Work in \mathcal{C} (without hyperimaginaries). Let $p \in S(\emptyset)$ and let $\theta \in L$. Assume p is analyzable in θ . Then p is analyzable in θ in finitely many steps. In particular, if T is a hypersimple unidimensional theory and there exists a non-algebraic supersimple definable set, then T has finite SU-rank, i.e. every complete type has finite SU-rank. In fact, for every given sort there is a finite bound on the SU-rank of all types in that sort, equivalently the global D-rank of any sort is finite.*

Another useful fact is the following.

Fact 2.3 [S1] *Let T be any unidimensional simple theory. Then T eliminates \exists^∞ .*

2.2 The forking topology, EPFO and PCFT

The forking topology is introduced in [S0] and is a variant of Hrushovski's and Pillay's topologies from [H0] and [P], respectively. In this section T is assumed to be simple and we work in \mathcal{C} .

Definition 2.4 Let $A \subseteq \mathcal{C}$ and let x be a finite tuple of variables.

1) An invariant set \mathcal{U} over A is said to be a *basic τ^f -open set over A* if there is $\phi(x, y) \in L(A)$ such that

$$\mathcal{U} = \{a \mid \phi(a, y) \text{ forks over } A\}.$$

Note that the family of basic τ^f -open sets over A is closed under finite intersections, thus forms a basis for a unique topology on $S_x(A)$. An open set in this topology is called a τ^f -open set over A or a forking-open set over A .

2) An invariant set \mathcal{U} over A is said to be a *basic τ_∞^f -open set over A* if \mathcal{U} is a type-definable τ^f -open set over A . The family of basic τ_∞^f -open sets over A is a basis for a unique topology on $S_x(A)$. An open set in this topology is called a τ_∞^f -open set over A .

Remark 2.5 The τ_∞^f -topology and in particular the τ^f -topology on $S_x(A)$ refines the Stone-topology of $S_x(A)$ for all x, A .

We will apply the following Fact.

Fact 2.6 *Let \mathcal{U} be a τ^f -open set over \emptyset and let A be any set. Then \mathcal{U} is τ^f -open over A .*

Recall the following definition from [S0].

Definition 2.7 We say that *the τ^f -topologies over A are closed under projections (T is PCFT over A)* if for every τ^f -open set $\mathcal{U}(x, y)$ over A the set $\exists y \mathcal{U}(x, y)$ is a τ^f -open set over A . We say that *the τ^f -topologies are closed under projections (T is PCFT)* if they are such over every set A .

In [BPV, Proposition 4.5] the authors proved the following equivalence which, for convenience, we will use as a definition (their definition involves extension with respect to pairs of models of T).

Definition 2.8 We say that the extension property is first-order in T or T is EPFO iff for every formulas $\phi(x, y), \psi(y, z) \in L$ the relation $Q_{\phi, \psi}$ defined by:

$$Q_{\phi, \psi}(a) \text{ iff } \phi(x, b) \text{ doesn't fork over } a \text{ for every } b \models \psi(y, a)$$

is type-definable (here a can be an infinite tuple from \mathcal{C} whose sorts are fixed).

Fact 2.9 [S1, Corollary 3.13] *Suppose the extension property is first-order in T . Then T is PCFT.*

We say that an A -invariant set \mathcal{U} has *finite SU -rank* if $SU(a/A) < \omega$ for all $a \in \mathcal{U}$, and has *bounded finite SU -rank* if there exists $n < \omega$ such that $SU(a/A) \leq n$ for all $a \in \mathcal{U}$. The existence of a τ^f -open set of bounded finite SU -rank implies the existence of an SU -rank 1 formula (i.e. a weakly-minimal formula):

Fact 2.10 [S0, Proposition 2.13] *Let \mathcal{U} be an unbounded τ^f -open set over some set A . Assume \mathcal{U} has bounded finite SU -rank. Then there exist a set $B \supseteq A$ with $|B \setminus A| < \omega$ and $\theta(x) \in L(B)$ of SU -rank 1 such that $\theta^{\mathcal{C}} \subseteq \mathcal{U} \cup \text{acl}(B)$.*

Now, recall the following two facts and their corollary. First, let $\mathcal{P}^{SU \leq 1}$ denote the class of complete real types over sets of size $\leq |T|$, of SU -rank ≤ 1 .

Fact 2.11 [P1] *Let T be a simple theory that eliminates \exists^∞ . Moreover, assume every type is analyzable in $\mathcal{P}^{SU \leq 1}$. Then the extension property is first-order in T .*

For a more general statement, see [S1, Lemma 3.7].

2.3 Stable independence and stable SU-rank

In this subsection T is assumed to be simple and we work in \mathcal{C} .

First we recall the notion of stable independence.

Definition 2.12 Let $a \in \mathcal{C}$, $A \subseteq B \subseteq \mathcal{C}$. We say that a is *stably-independent* from B over A if for every stable $\phi(x, y) \in L$, if $\phi(x, b)$ is over B and $a' \models \phi(x, b)$ for some $a' \in dcl(Aa)$, then $\phi(x, b)$ doesn't divide over A . In this case we denote it by $a \downarrow^s_B A$.

The notion of stable SU -rank is defined via stable dependence.

Definition 2.13 1) For $a \in \mathcal{C}$ and $A \subseteq \mathcal{C}$ the SU_s -rank is defined by induction on α : if $\alpha = \beta + 1$, then $SU_s(a/A) \geq \alpha$ if there exists $B \supseteq A$ such that $a \not\downarrow^s_B A$ and $SU_s(a/B) \geq \beta$. For limit α , $SU_s(a/A) \geq \alpha$ if $SU_s(a/A) \geq \beta$ for all $\beta < \alpha$.

2) Let \mathcal{U} be an A -invariant set. We write $SU_s(\mathcal{U}) = \alpha$ (the SU_s -rank of \mathcal{U} is α) if $\text{Max}\{SU_s(p) \mid p \in S(A), p^{\mathcal{C}} \subseteq \mathcal{U}\} = \alpha$. We say that \mathcal{U} has *bounded finite SU_s -rank* if for some $n < \omega$, $SU_s(\mathcal{U}) = n$. Note that the SU_s -rank of \mathcal{U} might, a priori, depend on the choice of the set A over which \mathcal{U} is invariant.

The following rank is a variation of stable SU -rank; it is non-increasing in extensions.

Definition 2.14 1) For $a \in \mathcal{C}$ and $A \subseteq \mathcal{C}$ the SU_{se} -rank is defined by induction on α : if $\alpha = \beta + 1$, $SU_{se}(a/A) \geq \alpha$ if there exist $B_1 \supseteq B_0 \supseteq A$ such that $a \not\downarrow^s_{B_0} B_1$ and $SU_{se}(a/B_1) \geq \beta$. For limit α , $SU_{se}(a/A) \geq \alpha$ if $SU_{se}(a/A) \geq \beta$ for all $\beta < \alpha$.

2) Let \mathcal{U} be an A -invariant set. We write $SU_{se}(\mathcal{U}) = \alpha$ (the SU_{se} -rank of \mathcal{U} is α) if $\text{Max}\{SU_{se}(p) \mid p \in S(A), p^{\mathcal{C}} \subseteq \mathcal{U}\} = \alpha$. We say that \mathcal{U} has *bounded finite SU_{se} -rank* if for some $n < \omega$, $SU_{se}(\mathcal{U}) = n$.

Remark 2.15 Note that $SU_{se}(a/B) \leq SU_{se}(a/A)$ for all $a \in \mathcal{C}$ and $A \subseteq B \subseteq \mathcal{C}$ (this is the reason for introducing SU_{se}). Also, clearly $SU_s(a/A) \leq SU_{se}(a/A) \leq SU(a/A)$ for all a, A . Clearly $SU_{se}(a/A) = 0$ iff $SU_s(a/A) = 0$ iff $a \in acl(A)$ for all a, A .

We will apply the following easy fact.

Fact 2.16 *For $a \in \mathcal{C}$ and $A \subseteq B \subseteq \mathcal{C}$, assume $tp(a/B)$ doesn't fork over $acl(aA) \cap acl(B)$ and $a \not\downarrow_A B$. Then $a \not\downarrow_S B$.*

3 Elimination of hyperimaginaries in reducts

In this section we include a remark by Ehud Hrushovski that allowed us to remove the assumption that the reducts eliminate hyperimaginaries (in the main theorem). Here T denotes any complete theory in a language L and we work in \mathcal{C} .

Definition 3.1 *A reduct T^- of T to a sublanguage $L^- \subseteq L$ is said to be E -closed if for every L^- -definable sets $D_1 \vdash D_2$ on S^2 (for some sort S of L^-) there exists a definable equivalence relation $E^- \in L^-$ satisfying $D_1 \vdash E^- \vdash D_2$, provided that there exists such definable equivalence relation in L .*

For a partial order (P, \leq) , a subset $A \subseteq P$ is called a *club* in (P, \leq) , if A is unbounded in (P, \leq) , that is, above any element of P there is an element of A , and A is closed in (P, \leq) , that is, for any chain $C \subseteq A$, if $a \in P$ is the supremum of C (i.e. a is an upper bound of C and a is smaller than any other upper bound of C) then $a \in A$.

Notation 3.2 *Let T^- be a reduct of T to L^- . The size of the reduct T^- is just $|T^-|$. Let λ be any infinite cardinal (or ∞). Let $(\mathcal{R}_T^\lambda, \leq_T)$ be the partial order of all reducts of T of size $\leq \lambda$, where the order is just inclusion (of the sublanguages of the reducts, i.e. of both the set of sorts and the set of formulas). It will be convenient to consider the isomorphic partial order $(\mathcal{R}_C^\lambda, \leq_C)$ of all the (saturated) model reducts of \mathcal{C} to a sublanguage of L size $\leq \lambda$.*

Claim 3.3 *Let T be any complete L -theory that eliminates hyperimaginaries.*
 1) *Let T^- be an E -closed reduct of T . Then T^- eliminates hyperimaginaries.*
 2) *The set of E -closed reducts of T is a club in $(\mathcal{R}_T^\infty, \leq_T)$. Given any infinite $\lambda \leq |L|$, the set of E -closed reducts of T of size $\leq \lambda$ is a club in $(\mathcal{R}_T^\lambda, \leq_T)$.*

Proof: 1) Say T^- is the reduct of T to L^- and so $\mathcal{C}|L^-$ is a saturated model of T^- . We claim that the hyperimaginaries of T^- are eliminated, namely: for every type-definable equivalence relation E^- of T^- on a complete type p^- of T^- over \emptyset , there are definable equivalence relations $E_i^- \in L^-$ such that E^- is equivalent to $\bigwedge_i E_i^-$ on p^- . Indeed, let $E^- = E^-(x, x')$ and $p^- = p^-(x)$ be such. Let $\phi_i^-(x, x') \in L^-$ be such that $E^-(x, x') = \bigwedge_i \phi_i^-(x, x')$. Let p be any complete type of T over \emptyset that extends p^- . By elimination of hyperimaginaries in T , there are $E_j(x, x') \in L$ such that $\bigwedge_j E_j(x, x')$ is equivalent to $E^-(x, x')$ on $p^{\mathcal{C}}$. Now, by compactness, for any i there is $j(i)$ such that $E_{j(i)}(x, x') \vdash \phi_i^-(x, x')$ on $p^{\mathcal{C}}$, likewise, for every j there exists $k(j)$ such that $\phi_{k(j)}^-(x, x') \vdash E_j(x, x')$ on $p^{\mathcal{C}}$. As T^- is an E -closed reduct of T , for every i , there is a definable equivalence relation $E_i^- \in L^-$ such that $\phi_{k(j(i))}^-(x, x') \vdash E_i^-(x, x') \vdash \phi_i^-(x, x')$ on $p^{\mathcal{C}}$ (using compactness). We conclude that E^- is equivalent to $\bigwedge_i E_i^-(x, x')$ on $p^{\mathcal{C}}$ and thus on $p^- \mathcal{C}$ as well (as E^- and E_i^- are all invariant under automorphisms of $\mathcal{C}|L^-$). 2) is immediate.

4 Dichotomies for \emptyset -invariant families of rank 1 types

Here we verify the following extension of [S2, Corollary 2.13] to a general \emptyset -invariant family of SU -rank 1 types. In this section T is assumed to be a simple theory with elimination of imaginaries.

We first recall some basic definitions from [S1].

Definition 4.1 A family

$$\Upsilon = \{\Upsilon_{x,A} \mid x \text{ is a finite sequence of variables and } A \subset \mathcal{C} \text{ is small}\}$$

is said to be a *projection closed family of topologies* if each $\Upsilon_{x,A}$ is a topology on $S_x(A)$ that refines the Stone-topology on $S_x(A)$, this family is invariant under automorphisms of \mathcal{C} and change of variables by variables of the same sort, the family is closed under product by the full Stone spaces $S_y(A)$ (where y is a disjoint tuple of variables) and closed by projections, namely whenever $\mathcal{U}(x, y) \in \Upsilon_{xy,A}$, $\exists y \mathcal{U}(x, y) \in \Upsilon_{x,A}$.

We will be interested in the case $\Upsilon = \tau^f$, where T is a PCFT theory. From now on fix a general projection closed family Υ of topologies.

Definition 4.2 1) A type $p \in S(A)$ is said to be *s-essentially 1-based over* $A_0 \subseteq A$ (*essentially 1-based over* $A_0 \subseteq A$) *by means of* Υ if for every finite tuple \bar{c} from p and for every (respectively, type-definable) Υ -open set \mathcal{U} over $A\bar{c}$, with the property that a is independent from A over A_0 for every $a \in \mathcal{U}$, the set $\{a \in \mathcal{U} \mid Cb(a/A\bar{c}) \not\subseteq bdd(aA_0)\}$ is nowhere dense in the Stone-topology of \mathcal{U} . We say $p \in S(A)$ is *s-essentially 1-based (essentially 1-based) by means of* Υ if p is s-essentially 1-based (respectively, essentially 1-based) over A by means of Υ .

2) Let V be an A_0 -invariant set and let $p \in S(A_0)$. We say that p is *analyzable in* V *by s-essentially 1-based (by essentially 1-based) types by means of* Υ if there exists $a \models p$ and there exists a sequence $(a_i \mid i \leq \alpha) \subseteq dcl(A_0a)$ with $a_\alpha = a$ such that $tp(a_i/A_0 \cup \{a_j \mid j < i\})$ is V -internal and s-essentially 1-based (respectively, essentially 1-based) over A_0 by means of Υ for all $i \leq \alpha$.

Theorem 4.3 *Let T be any countable hypersimple theory with PCFT. Let \mathcal{P}_0 be an \emptyset -invariant family of SU-rank 1 partial types. Then, either there exists a weakly-minimal formula that is almost \mathcal{P}_0 -internal, or every complete type $p \in S(A)$ that is internal in \mathcal{P}_0 is essentially 1-based over \emptyset by means of τ^f . In particular, either there exists a weakly-minimal formula that is almost \mathcal{P}_0 -internal, or whenever $p \in S(A)$, where A is countable, and p is analyzable in \mathcal{P}_0 , p is analyzable in \mathcal{P}_0 by essentially 1-based types by means of τ^f .*

The most general dichotomy theorem of this type that we present is the following theorem that generalizes [S2, Theorem 2.3] to any \emptyset -invariant family of SU-rank 1 types. The proof of this theorem is almost identical to the proof of [S2, Theorem 2.3] (but note that the next version that we present with a proof contains all modifications that are needed for the proof of it).

Theorem 4.4 *Let T be any hypersimple theory. Let Υ be a projection-closed family of topologies. Let \mathcal{P}_0 be an \emptyset -invariant family of SU-rank 1 types. Then, either there exists an unbounded Υ -open set (over some small set A) that is almost \mathcal{P}_0 -internal (and in particular has finite SU-rank), or every complete type $p \in S(A)$ that is internal in \mathcal{P}_0 is s-essentially 1-based over \emptyset by means of Υ . In particular, either there exists an unbounded Υ -open set that is almost \mathcal{P}_0 -internal, or whenever $p \in S(A)$ and p is analyzable in \mathcal{P}_0 , p is analyzable in \mathcal{P}_0 by s-essentially 1-based types by means of Υ .*

The next theorem is a version of Theorem 4.4 for a countable language with a stronger consequence and is a generalization of [S2, Theorem 2.11] to

\emptyset -invariant family of SU -rank 1 types. We give the complete proof of this theorem.

Theorem 4.5 *Let T be any countable hypersimple theory. Let Υ be a projection-closed family of topologies such that $\{a \in \mathcal{C}^x \mid a \notin \text{acl}(A)\} \in \Upsilon_{x,A}$ for all x and set A . Let \mathcal{P}_0 be an \emptyset -invariant family of SU -rank 1 types. Then, either there exists an unbounded type-definable Υ -open set over some small set that is almost \mathcal{P}_0 -internal and has **bounded** finite SU -rank, or every complete type $p \in S(A)$ that is internal in \mathcal{P}_0 is essentially 1-based over \emptyset by means of Υ . In particular, either there exists an unbounded type-definable Υ -open set that is almost \mathcal{P}_0 -internal and has **bounded** finite SU -rank, or whenever $p \in S(A)$, where A is countable, and p is analyzable in \mathcal{P}_0 , p is analyzable in \mathcal{P}_0 by essentially 1-based types by means of Υ .*

Proof: Υ will be fixed and we'll freely omit the phrase "by means of Υ ". To see the "In particular" part, work over a countable A and assume that every $p' \in S(A')$, with countable $A' \supseteq A$, that is internal in \mathcal{P}_0 , is essentially 1-based over A . Moreover, assume $p \in S(A)$ is non-algebraic and every non-algebraic extension of p is non-foreign to \mathcal{P}_0 . Then, for $a \models p$ there exists $a' \in \text{dcl}(Aa) \setminus \text{acl}(A)$ such that $tp(a'/A)$ is \mathcal{P}_0 -internal and thus essentially 1-based over A by our assumption. Thus, by repeating this process we get that p is analyzable in \mathcal{P}_0 by essentially 1-based types.

We now prove the main part. Assume there exists $p \in S(A)$ that is internal in \mathcal{P}_0 , and p is not essentially 1-based over \emptyset . By the definition, there exist a finite tuple d of realizations of p and b that is independent from d over A , and a finite tuple \bar{c} of realizations of types from \mathcal{P}_0 over Ab such that $d \in \text{dcl}(Ab\bar{c})$, and there exists a type-definable Υ -open set \mathcal{U} over Ad such that a is independent from A for all $a \in \mathcal{U}$ and $\{a \in \mathcal{U} \mid Cb(a/Ad) \not\subseteq \text{acl}(a)\}$ is not nowhere dense in the Stone-topology of \mathcal{U} . So, since Υ refines the Stone-topology, by intersecting \mathcal{U} with a definable set, we may assume that $\{a \in \mathcal{U} \mid Cb(a/Ad) \not\subseteq \text{acl}(a)\}$ is dense in the Stone-topology of \mathcal{U} . Now, for each (finite) subsequence \bar{c}_0 of \bar{c} , let

$$F_{\bar{c}_0} = \{a \in \mathcal{U} \mid \exists b', \bar{c}'_0, \bar{c}'_1 \text{ s.t. } tp(b'\bar{c}'_0\bar{c}'_1/Ad) = tp(b\bar{c}_0(\bar{c} \setminus \bar{c}_0)/Ad) \text{ and } a \perp Ab'\bar{c}'_0\}.$$

Note that since d is independent from b over A , any $a \in \mathcal{U}$ is independent from Ab' whenever $tp(b'/Ad) = tp(b/Ad)$ and $a \perp_{Ad}^a b'$. Thus $F_{\emptyset} = \mathcal{U}$. Let \bar{c}_0^* be a maximal subsequence (with respect to inclusion) of \bar{c} such that $F_{\bar{c}_0^*}$ has

non-empty Stone-interior in \mathcal{U} over Ad (note that $F_{\bar{c}}$ has no Stone-interior relatively in \mathcal{U}). Let $\mathcal{U}^* = \bigcap_{\bar{c}_0^* \subset \bar{c}' \subseteq \bar{c}} \mathcal{U} \setminus F_{\bar{c}'}$. Note that each $F_{\bar{c}'}$ is Stone closed relatively in \mathcal{U} . Thus \mathcal{U}^* is Stone-dense and Stone-open in \mathcal{U} and therefore there exists a non-empty relatively Stone-open in \mathcal{U} set $W^* \subseteq F_{\bar{c}_0^*} \cap \mathcal{U}^*$. As \mathcal{U} is type-definable, we may assume W^* is type-definable.

Claim 4.6 W^* is a non-empty Υ -open set over Ad such that $\{a \in W^* \mid Cb(a/Ad) \not\subseteq acl(a)\}$ is dense in the Stone-topology of W^* and for every $a \in W^*$ we have: there exists $b'\bar{c}'_0\bar{c}'_1 \models tp(b\bar{c}_0^*(\bar{c} \setminus \bar{c}_0^*)/Ad)$ such that a is independent from $Ab'\bar{c}'_0$ over \emptyset and moreover, for every $b'\bar{c}'_0\bar{c}'_1 \models tp(b\bar{c}_0^*(\bar{c} \setminus \bar{c}_0^*)/Ad)$ such that a is independent from $Ab'\bar{c}'_0$ we necessarily have $\bar{c}'_1 \in acl(aAb'\bar{c}'_0)$.

Proof: The first part is immediate by the fact that $W^* \subseteq F_{\bar{c}_0^*}$. For the "moreover" part note that since $a \in W^* \subseteq \mathcal{U}^*$, we get that by the definition of \mathcal{U}^* , $c' \in acl(aAb'\bar{c}'_0)$ for every $b'\bar{c}'_0\bar{c}'_1 \models tp(b\bar{c}_0^*(\bar{c} \setminus \bar{c}_0^*)/Ad)$ and $c' \in \bar{c}'_1$ (as $SU(c'/Ab') \leq 1$ for every $c' \in \bar{c}'_1$). \square

Let us now define a set V over Ad by

$$V = \{(e', b', \bar{c}'_0, \bar{c}'_1, a') \mid \text{if } tp(b'\bar{c}'_0\bar{c}'_1/Ad) = tp(b\bar{c}_0^*(\bar{c} \setminus \bar{c}_0^*)/Ad) \text{ and } a' \perp Ab'\bar{c}'_0 \\ \text{then } e' \in acl(Cb(Ab'\bar{c}'_0\bar{c}'_1/a'))\}.$$

Let $V^* = \{e' \mid \exists a' \in W^* \forall b', \bar{c}'_0, \bar{c}'_1 V(e', b', \bar{c}'_0, \bar{c}'_1, a')\}$.

Claim 4.7 V^* is a Υ -open set over Ad .

Proof: Recall the following fact [S2, Proposition 2.4].

Fact 4.8 Let $q(x, y) \in S(\emptyset)$ and let $\chi(x, y, z)$ be an \emptyset -invariant set such that for all $(c, b, a) \models \chi(x, y, z)$ we have $b \succeq_a bc$. Then the set

$$\mathcal{U} = \{(e, c, b, a) \mid e \in acl(Cb(cb/a))\}$$

is relatively Stone-open inside the set

$$F = \{(e, c, b, a) \mid b \perp a, \models \chi(c, b, a), tp(cb) = q\}.$$

(where e is taken from a fixed sort too).

By Fact 4.8 and Claim 4.6, there exists a Stone-open set V' over Ad such that for all $a' \in W^*$ and for all $e', b', \bar{c}'_0, \bar{c}'_1$ we have $V'(e', b', \bar{c}'_0, \bar{c}'_1, a')$ if and only if $V(e', b', \bar{c}'_0, \bar{c}'_1, a')$. Thus, we may replace V by V' in the definition of V^* . As Stone-open sets are closed under the \forall quantifier, the Υ topology refines the Stone-topology and closed under product by a full Stone-space and closed under projections, we conclude that V^* is a Υ -open set. \square

Claim 4.9 *For appropriate sort for e' , the set V^* is unbounded and is almost \mathcal{P}_0 -internal (over Ad) and thus has finite SU -rank over Ad .*

Proof: First, note the following general observation.

Fact 4.10 *Assume $d \in dcl(c)$. Then $Cb(d/a) \in dcl(Cb(c/a))$ for all a .*

Let $a^* \in W^*$ be such that $Cb(a^*/Ad) \not\subseteq acl(a^*)$. Then $Cb(Ad/a^*) \not\subseteq acl(Ad)$. By Fact 4.10, there exists $e^* \notin acl(Ad)$ such that $e^* \in acl(Cb(Ab'\bar{c}'_0\bar{c}'_1/a^*))$ for all $b'\bar{c}'_0\bar{c}'_1 \models tp(b\bar{c}_0^*(\bar{c}\backslash\bar{c}_0^*)/Ad)$. In particular, $e^* \in V^*$. Thus, if we fix the sort for e' in the definition of V^* to be the sort of e^* , then V^* is unbounded. Now, let $e' \in V^*$. Then for some $a' \in W^*$, $\models V(e', \bar{c}'_0, \bar{c}'_1, b', a')$ for all $b', \bar{c}'_0, \bar{c}'_1$. By Claim 4.6, there exists $b'\bar{c}'_0\bar{c}'_1 \models tp(b\bar{c}_0^*(\bar{c}\backslash\bar{c}_0^*)/Ad)$ such that a' is independent from $Ab'\bar{c}'_0$ over \emptyset . Thus, by the definition of V^* and V , $e' \in acl(Cb(Ab'\bar{c}'_0\bar{c}'_1/a'))$. Since Ab' is independent from a' over \emptyset , $tp(e')$ is almost \mathcal{P}_0 -internal (as $Cb(Ab'\bar{c}'_0\bar{c}'_1/a')$ is in the definable closure of any Morley sequence of $Lstp(Ab'\bar{c}'_0\bar{c}'_1/a')$). In particular, $tp(e'/Ad)$ is almost \mathcal{P}_0 -internal by Fact 2.1 and therefore $tp(e'/Ad)$ has finite SU -rank. \square

Claim 4.11 *There exists $V^{**} \subseteq V^*$ that is unbounded, type-definable and Υ -open over Ad .*

Proof: By the definition of V^* and the proof of Claim 4.7 there exist a Stone open set V_0 over Ad such that $V^* = \{e' | \exists a' \in W^* (V_0(e', a'))\}$. By replacing V_0 by a definable set and using the fact that W^* is type-definable and that Υ is a projection-closed family of topologies we get the required set V^{**} \square

Now, by the proof of Claim 4.9 we know that for all $e' \in V^{**}$ we have $e' \in acl(Cb(Ab'\bar{c}'_0\bar{c}'_1/a'))$ for some $a' \in W^*$ and some $b', \bar{c}'_0, \bar{c}'_1$ such that a' is independent from $Ab'\bar{c}'_0$ over \emptyset and $b'\bar{c}'_0\bar{c}'_1 \models tp(b\bar{c}_0^*(\bar{c}\backslash\bar{c}_0^*)/Ad)$. Let $q = tp(Ab)$.

For every $\chi = \chi(x, y_0, \dots, y_n, \bar{z}_0, \bar{z}_1, \dots, \bar{z}_n) \in L$ (for some $n < \omega$) such that $\forall y_0 y_1 \dots y_n \bar{z}_0 \bar{z}_1 \dots \bar{z}_n \exists^{< \infty} x \chi(x, y_0, y_1, \dots, y_n, \bar{z}_0, \bar{z}_1, \dots, \bar{z}_n)$, let

$$F_\chi = \{e \in V^{**} \mid \models \chi(e, C_0, C_1, \dots, C_n, \bar{c}_0, \bar{c}_1, \dots, \bar{c}_n) \text{ for some } \bar{c}_0, \dots, \bar{c}_n \text{ and some } \\ \emptyset\text{-independent sequence } (C_i \mid i \leq n) \text{ of realization of } q \\ \text{ such that } tp(C_i, \bar{c}_i) = tp(Ab, \bar{c}) \text{ and } e \perp (C_i \mid i \leq n)\}.$$

Note that each F_χ is type-definable. By the aforementioned, we get that $V^{**} \subseteq \bigcup_\chi F_\chi$ (the union is over each χ as above). By the Baire category theorem applied to the Stone-topology of the Stone-closed set $V^{**} \setminus acl(Ad)$, there exists $\theta \in L(Ad)$ such that

$$\tilde{V} \equiv \theta^c \cap (V^{**} \setminus acl(Ad)) \neq \emptyset \text{ and } \tilde{V} \subseteq F_{\chi^*}$$

for some χ^* as above. Clearly, \tilde{V} is unbounded, type-definable and Υ -open (by the assumptions on Υ). Now, there exists a fixed $m^* < \omega$ such that for every $a \in \tilde{V}$, $SU(a/Ad) \leq m^*$ and $tp(a/Ad)$ is almost \mathcal{P}_0 -internal (as $tp(a)$ is almost \mathcal{P}_0 -internal). This completes the proof of the main part of the theorem. \square

The proof of the main result of this section now follows exactly in the same way as in [S2]. We write the proof for completeness.

Proof of Theorem 4.3 Our assumptions are clearly a special case of the assumptions of Theorem 4.5, thus we only need to prove the first part. By the conclusion of Theorem 4.5, we may assume that there exists a τ^f -open set \mathcal{U} of bounded finite SU -rank over some small set A that is almost \mathcal{P}_0 -internal. By Fact 2.10, there exists a weakly-minimal $\theta(x, b) \in L(B)$ for some small set $B \supseteq A$, such that $\theta^c \subseteq \mathcal{U} \cup acl(B)$. Now, $tp(a/B)$ is almost \mathcal{P}_0 -internal for every $a \in \theta^c$, and so $tp(a/b)$ (b is the parameter of $\theta(x, b)$) is almost \mathcal{P}_0 -internal for every $a \in \theta^c$ (by taking non-forking extensions). \square

5 Main result

From now on we assume T is an arbitrary simple theory with elimination of imaginaries unless stated otherwise. We work in \mathcal{C} .

Definition 5.1 1) We say that T is *analyzable in SU -rank 1 types* if every type is analyzable in the family of SU -rank 1 types.
2) We say that T is *coordinatized in finite rank* if for every $a \in \mathcal{C}$ and $A \subseteq \mathcal{C}$ such that $a \notin \text{acl}(A)$ there exists $a' \in \text{acl}(aA) \setminus \text{acl}(A)$ with $SU(a'/A) < \omega$.

Lemma 5.2 *Assume T is hypersimple. T is coordinatized in finite rank iff T is analyzable in SU -rank 1 types.*

Proof: If T is analyzable in SU -rank 1 types then clearly T is coordinatized in finite rank. Assume now that T is coordinatized in finite rank. We first note the following.

Claim 5.3 *Let T be any simple theory. Let $a \in \mathcal{C}$ be such that $SU(a) = n < \omega$ and such that for some $b \in \mathcal{C}$ with $SU(b) < \infty$ we have $SU(a/b) = n - 1$. Then $tp(a)$ is non-orthogonal to an SU -rank 1 hyperimaginary type.*

Proof: Let $e = Cb(Lstp(a/b))$ (e is a hyperimaginary). Since $SU(e) < \infty$ (as we assume $SU(b) < \infty$), there exists a set A such that $SU(e/A) = 1$. By extension we may clearly assume $a \downarrow_e A$. We claim that $e \in bdd(aA)$ (*).

Indeed, otherwise $e \not\downarrow_A a$ and so $e \in bdd(A)$ (as $tp(a/e)$ is canonical), a contradiction to $SU(e/A) = 1$. Now, $SU(a/eA) = SU(a/e) = n - 1$. By (*), $SU(a/A) = SU(ae/A) \geq SU(a/eA) + SU(e/A) = n$. Thus $a \not\downarrow A$ and so $tp(a)$ is non-orthogonal to $tp(e/A)$. \square

Now, let a, A be given such that $a \notin \text{acl}(A)$. By our assumption, there exists $a' \in \text{acl}(aA) \setminus \text{acl}(A)$ with $SU(a'/A) = n$ for some $n < \omega$. Let $b \in \mathcal{C}$ be such that $SU(a/Ab) = n - 1$ and let $(b_i | i < \alpha)$ be such that $b_i \in \text{acl}(bA)$ and $0 < SU(b_i/Ab_{<i}) < \omega$ for all $i < \alpha$ and such that $\text{acl}(Ab) = \text{acl}(A \cup \{b_i | i < \alpha\})$. As $a' \not\downarrow_A b$, there exists a minimal $i^* < \alpha$ such that $a' \not\downarrow_A \{b_i | i \leq i^*\}$.

Then $a' \not\downarrow_A \{b_i | i < i^*\}$. Now, a' satisfies the assumptions of Claim 5.3 when working over $A \cup \{b_i | i < i^*\}$. Thus $tp(a'/A \cup \{b_i | i < i^*\})$ is non-orthogonal to an SU -rank 1 type which we may clearly assume to be a type of an imaginary. Thus $tp(a'/A)$ is non-orthogonal an SU -rank 1 imaginary type and so is $tp(a/A)$ (as $a' \in \text{acl}(aA)$).

We start with the following proposition that generalizes the main result in [S1]; the proof is similar but applies Theorem 4.3.

Proposition 5.4 *Assume T is a countable hypersimple theory that is coordinatized in finite rank and eliminates \exists^∞ . Then there exists a weakly minimal formula.*

Proof: By Lemma 5.2, T is analyzable in SU-rank 1 types. As T eliminates \exists^∞ , T is EPFO by Fact 2.11. By Fact 2.9, T is PCFT. Let \mathcal{P}_0 be the family of all SU-rank 1 types. By Theorem 4.3, we may assume that every complete finitary type over a countable set is analyzable in \mathcal{P}_0 by essentially 1-based types by means of τ^f . We recall the following fact that for simplicity we state for a special case [S3, Corollary 4.5]. Infact, the result is valid for a large class of sets ($\tilde{\tau}_{low}^f$ -sets) instead of the specific set \mathcal{U}_0 that is fixed in the following statement.

Fact 5.5 *Let T be a countable simple theory with EPFO. Let $\mathcal{U}_0 = \mathcal{C}^s \setminus \text{acl}(\emptyset)$ for some non-algebraic sort s say. Assume for every $a \in \mathcal{U}_0$ there exists $a' \in \text{acl}(a) \setminus \text{acl}(\emptyset)$ such that $SU_{se}(a') < \omega$. Then there exists an unbounded τ_∞^f -open set \mathcal{U} over a finite set such that \mathcal{U} has bounded finite SU_{se} -rank.*

T satisfies the assumptions of Fact 5.5, so let \mathcal{U} be a set as in its conclusion. In particular, $SU_s(\mathcal{U}) = n$ for some $n < \omega$. Recall now the following easy lemma.

Lemma 5.6 [S1, Lemma 7.4] *Assume \mathcal{U} is an unbounded τ_∞^f -open set of bounded finite SU_s -rank over some finite set A . Then there exists a τ_∞^f -open set $\mathcal{U}^* \subseteq \mathcal{U}$ over some finite set $B^* \supseteq A$ of SU_s -rank 1.*

By Lemma 5.6, we may assume $SU_s(\mathcal{U}) = 1$, \mathcal{U} is a type-definable τ^f -open set over a finite set A_0 . We claim $SU(\mathcal{U}) = 1$. Indeed, otherwise there exists a and $d \in \mathcal{U}$ such that $d \not\downarrow_{A_0} a$ and $d \notin \text{acl}(aA_0)$. Since every finitary type over a countable set is analyzable in \mathcal{P}_0 , there exists $(a_i | i \leq \alpha) \subseteq \text{dcl}(aA_0)$ with $a_\alpha = a$ (where $\alpha < \omega_1$) such that $\text{tp}(a_i/A_0 \cup \{a_j | j < i\})$ is essentially 1-based over A_0 by means of τ^f for all $i \leq \alpha$. Now, let $i^* \leq \alpha$ be minimal such that there exists $d' \in \mathcal{U}$ satisfying $d' \not\downarrow_{A_0} \{a_i | i \leq i^*\}$ and

$d' \notin \text{acl}(A_0 \cup \{a_i | i \leq i^*\})$. Pick $\phi(x, a') \in L(A_0 \cup \{a_i | i \leq i^*\})$ that forks over A_0 and such that $\phi(d', a')$. Let

$$V = \{d \in \mathcal{U} \mid \phi(d, a') \text{ and } d \notin \text{acl}(A_0 \cup \{a_i | i \leq i^*\})\}.$$

By minimality of i^* , d is independent from $\{a_i | i < i^*\}$ over A_0 for all $d \in V$. Clearly V is type-definable and by Fact 2.6, V is a τ^f -open set over $A_0 \cup \{a_i | i \leq i^*\}$. Now, since $\text{tp}(a_{i^*}/A_0 \cup \{a_i | i < i^*\})$ is essentially 1-based over A_0 by means of τ^f , the set

$$\{d \in V \mid \text{Cb}(d/A_0 \cup \{a_i | i \leq i^*\}) \in \text{bdd}(dA_0)\}$$

contains a relatively Stone-open and Stone-dense subset of V . In particular, there exists $d^* \in V$ such that $\text{tp}(d^*/A_0 \cup \{a_i | i \leq i^*\})$ doesn't fork over $\text{acl}(A_0 d^*) \cap \text{acl}(A_0 \cup \{a_i | i \leq i^*\})$. Since we know $d^* \not\perp_{A_0} A_0 \cup \{a_i | i \leq i^*\}$,

Fact 2.16 implies $d^* \not\perp_{A_0}^s A_0 \cup \{a_i | i \leq i^*\}$. Hence $d^* \in V$ implies $SU_s(d^*/A_0) \geq$

2, which contradicts $SU_s(\mathcal{U}) = 1$. Thus we have proved $SU(\mathcal{U}) = 1$. Now, by Fact 2.10 there exists a definable set of SU -rank 1. \square

Before stating the main theorem, we give some terminology and easy remarks. Recall that we work in $\mathcal{C} = \mathcal{C}^{eq}$ and that $(\mathcal{R}_{\mathcal{C}}^{\lambda, \leq c})$ is the partial order of reducts of \mathcal{C} of size $\leq \lambda$.

Definition 5.7 Let $\mathcal{C}|L^- \in \mathcal{R}_{\mathcal{C}}^{\lambda}$. We will say that $\mathcal{C}|L^-$ is *eq-closed* if $T^- = \text{Th}(\mathcal{C}|L^-)$ has uniform elimination of imaginaries, i.e. for every definable equivalence relation $E \in L^-$ on $S_0 \times S_1 \times \dots \times S_k$, where S_i are sorts of L^- , there is a definable function $f_E \in L^-$ whose domain is $(S_0 \times S_1 \times \dots \times S_k)^{\mathcal{C}}$ such that for all \bar{a}, \bar{b} , we have $f_E(\bar{a}) = f_E(\bar{b})$ iff $E(\bar{a}, \bar{b})$.

Remark 5.8 For every reduct $\mathcal{C}|L^- \in \mathcal{R}_{\mathcal{C}}^{\lambda}$ there exists a reduct $\mathcal{C}|L^* \in \mathcal{R}_{\mathcal{C}}^{\lambda}$ that is eq-closed and is an expansion of $\mathcal{C}|L^-$. Thus for every infinite cardinal λ , the set of reducts in $\mathcal{R}_{\mathcal{C}}^{\lambda}$ that are eq-closed is a club in $(\mathcal{R}_{\mathcal{C}}^{\lambda, \leq c})$.

Proof: Expand the reduct $\mathcal{C}|L^-$ of $\mathcal{C} = \mathcal{C}^{eq}$ by adding for every definable equivalence relation E on $S_0 \times S_1 \times \dots \times S_k$, where S_i are sorts of L^- and $E \in L^-$, a definable function $f_E \in L$ whose domain is $(S_0 \times S_1 \times \dots \times S_k)^{\mathcal{C}}$ and is onto the interpretation of some sort of L such that $f_E(\bar{a}) = f_E(\bar{b})$ iff $E(\bar{a}, \bar{b})$. Now,

the resulting expansion will have uniform elimination of imaginaries. It is immediate that the set of eq-closed reducts in $\mathcal{R}_{\mathcal{C}}^\lambda$ is closed in $(\mathcal{R}_{\mathcal{C}}^\lambda, \leq_c)$.

Now, note the following easy general remark on clubs.

Remark 5.9 Let (P, \leq) be a directed partial order that is ω -closed (i.e. any increasing sequence $(a_i | i < \omega)$ has a supremum). Then the intersection of finitely many clubs in (P, \leq) is a club.

In the proof we will refer to the following notion.

Definition 5.10 We say that T is strongly non-supersimple if $D(\phi(x, a)) = \infty$ for every non-algebraic $\phi(x, a) \in L(\mathcal{C})$.

Remark 5.11 Note that T is strongly non-supersimple iff for every non-algebraic $\phi(x, a) \in L(\mathcal{C})$ there exists a non-algebraic $\psi(x, b) \in L(\mathcal{C})$ such that $\psi(x, b) \vdash \phi(x, a)$ and $\psi(x, b)$ forks over a iff there does not exist a weakly minimal formula.

Theorem 5.12 Let $T = T^{eq}$ be a hypersimple unidimensional theory. Assume there is a club of countable reducts of T in $(\mathcal{R}_{\mathcal{C}}^{\aleph_0}, \leq_c)$ that are coordinatized in finite rank. Then T is supersimple.

Proof: First, if T is not strongly non-supersimple then we are done by Fact 2.2. Therefore, we may assume T is strongly non-supersimple. By Fact 2.3, T eliminates \exists^∞ , thus every reduct of T eliminates \exists^∞ .

Claim 5.13 The set $\tilde{\mathcal{C}}_1$ of countable strongly non-supersimple reducts of \mathcal{C} is a club in $(\mathcal{R}_{\mathcal{C}}^{\aleph_0}, \leq_c)$.

Proof: First, we prove that $\tilde{\mathcal{C}}_1$ is unbounded in $(\mathcal{R}_{\mathcal{C}}^{\aleph_0}, \leq_c)$. Let $\mathcal{C}|L^- \in \mathcal{R}_{\mathcal{C}}^{\aleph_0}$. We construct by induction an increasing sequence of reducts $(\mathcal{C}_n | n < \omega)$, $\mathcal{C}_n \in \mathcal{R}_{\mathcal{C}}^{\aleph_0}$, where $\mathcal{C}_n = \mathcal{C}|L_n^-$ for some countable sublanguge L_n^- of L , $T_n^- = Th(\mathcal{C}_n)$ in the following way. Let $\mathcal{C}_0 = \mathcal{C}|L_0^-$, $L_0^- = L^-$ and assume L_k^- have already been defined for $k \leq n$. We define L_{n+1}^- . For any fixed $\phi(x, y) \in L_n^- \setminus L_{n-1}^-$ we define a finite set of formulas $\Delta_\phi = \{\psi_i | i \leq n(\phi)\}$, where $\psi_i = \psi_i(x, y_i) \in L = L(T)$, $n(\phi) < \omega$ in the following way. Since T is strongly non-supersimple, for every $a \in \mathcal{C}$ such that $\exists^\infty x \phi(x, a)$, there exists

$\psi_a(x, z) \in L$ and some $b \in \mathcal{C}$ such that $\psi_a(x, b) \vdash \phi(x, a)$, $\psi_a(x, b)$ forks over a and $\exists^\infty x \psi_a(x, b)$ (*). For every $\psi(x, z) \in L$ let

$$\theta_{\phi, \psi}(z, y) = \exists^\infty x \psi(x, z) \wedge \forall x (\psi(x, z) \rightarrow \phi(x, y)).$$

By elimination of \exists^∞ (in \mathcal{C}), $\theta_{\phi, \psi}(z, y)$ is definable. Now, let $Q_{\psi, \theta_{\phi, \psi}}(y)$ be the relation in Fact 2.8 defined for $\theta_{\phi, \psi}, \psi$. So, for every $a \in \mathcal{C}$, $\neg Q_{\psi, \theta_{\phi, \psi}}(a)$ iff there exists b such that $\psi(x, b)$ is not algebraic, $\psi(x, b) \vdash \phi(x, a)$ and $\psi(x, b)$ forks over a . Since T is EPFO, we know that each $\neg Q_{\psi, \theta_{\phi, \psi}}$ is Stone-open. By (*), in \mathcal{C} :

$$\exists^\infty x \phi(x, y) \vdash \bigvee_{\psi \in L} \neg Q_{\psi, \theta_{\phi, \psi}}(y).$$

By compactness, there are $\psi_0 = \psi_0(\phi), \dots, \psi_{n(\phi)} = \psi_{n(\phi)}(\phi) \in L$ such that in \mathcal{C} :

$$\exists^\infty x \phi(x, y) \vdash \bigvee_{i < n(\phi)} \neg Q_{\psi_i, \theta_{\phi, \psi_i}}(y) (**).$$

Now, let $\Delta_\phi = \{\psi_i(\phi) \mid i \leq n(\phi)\}$. Let L_{n+1}^- be the set of formulas generated by the set

$$\nu_{n+1} = L_n^- \cup \bigcup \{\Delta_\phi \mid \phi(x, y) \in L_n^- \setminus L_{n-1}^-\},$$

where the set of sorts attached to L_{n+1}^- is the set of all sorts of variables that appears in ν_{n+1} . Let $\mathcal{C}_{n+1} = \mathcal{C} \upharpoonright L_{n+1}^-$. Now, let $L_\omega^- = \bigcup_{n < \omega} L_n^-$ and let $\mathcal{C}_\omega = \mathcal{C} \upharpoonright L_\omega^-$, $T_\omega^- = Th(\mathcal{C}_\omega)$. We claim that T_ω^- is strongly non-supersimple. Indeed, given a formula $\phi_\omega(x, y) \in L_\omega^-$, let $a \in \mathcal{C}_\omega$ be such that $\models \exists^\infty x \phi_\omega(x, a)$. Then, by (***) there exists $\psi(x, z) \in \Delta_{\phi_\omega} \subseteq L_\omega^-$ such that for some b we have $\psi(x, b) \vdash \phi_\omega(x, a)$ and $\psi(x, b)$ forks over a in \mathcal{C} and thus in particular $\psi(x, b)$ forks over a in \mathcal{C}_ω . Thus T_ω^- is strongly non-supersimple. Now, to show that $\tilde{\mathcal{C}}_1$ is closed in $(\mathcal{R}_{\mathcal{C}}^{\aleph_0}, \leq_{\mathcal{C}})$, let $\tilde{\mathcal{C}}$ be a chain in $\tilde{\mathcal{C}}_1$. We claim that $\mathcal{C}^* = \bigcup \tilde{\mathcal{C}}$ (the universe of \mathcal{C}^* is the union of the interpretations of the sorts of all members of $\tilde{\mathcal{C}}$ and likewise for the definable sets of \mathcal{C}^*) is strongly non-supersimple. Indeed, let $\phi(x, a) \in L(\mathcal{C}^*)$ be non-algebraic. Then there exists $\mathcal{C}_0 = \mathcal{C} \upharpoonright L_0 \in \tilde{\mathcal{C}}$ for some countable sublanguage L_0 of L such that $\phi(x, a) \in L_0(\mathcal{C}_0)$. Since $Th(\mathcal{C}_0)$ is strongly non-supersimple, there exists a non-algebraic $\psi(x, b) \in L_0(\mathcal{C}_0)$ such that $\psi(x, b) \vdash \phi(x, a)$ and $\psi(x, b)$ forks over a in \mathcal{C}_0 . By Ramsey and compactness, there exists a formula $\psi(x, b')$ that is a -conjugate to $\psi(x, b)$ in \mathcal{C}_0 and that forks over a in the sense of \mathcal{C}^* . Thus $Th(\mathcal{C}^*)$ is strongly non-supersimple. \square

By Claim 5.13, Claim 3.3, Remark 5.8, the assumptions of the theorem and Remark 5.9, there is a club of reducts in $\mathcal{R}_C^{\aleph_0}$ that are strongly non-supersimple, hypersimple, eq-closed and coordinatized in finite rank. Any such reduct contradicts Proposition 5.4 (as elimination of \exists^∞ is clearly preserved under reducts). \square

References

- [BPV] I.Ben-Yaacov, A.Pillay, E.Vassiliev, Lovely pairs of models, *Annals of Pure and Applied Logic* 122 (2003), no. 1-3.
- [H0] E.Hrushovski, Countable unidimensional stable theories are superstable, unpublished paper.
- [H1] E.Hrushovski, Unidimensional theories are superstable, *Annals of Pure and Applied Logic*, 50 (1990), pgs 117-138.
- [HKP] B.Hart, B.Kim and A.Pillay, Coordinatization and canonical bases in simple theories, *Journal of Symbolic Logic*, 65 (2000), pgs 293-309.
- [HP] E. Hrushovski, A. Pillay, Groups definable in local fields and pseudo-finite fields, *Israel J. Math.* 85 (1994), 203 - 262.
- [K1] B.Kim, Forking in simple unstable theories, *Journal of London Math. Society*, 57 (1998), pgs 257-267.
- [KP] B.Kim and A.Pillay, Simple theories, *Annals of Pure and Applied Logic*, 88, 1997 pgs 149-164.
- [P] A.Pillay, On countable simple unidimensional theories, *Journal of Symbolic Logic* 68 (2003), no. 4.
- [P1] A.Pillay, The extension property is first order in unidimensional simple theories, unpublished note.
- [S0] Z.Shami, On analyzability in the forking topology for simple theories, *Annals of Pure Applied Logic* 142 (2006), no. 1-3, 115–124.
- [S1] Z.Shami, Countable hypersimple unidimensional theories, *J. London Math. Soc. Volume 83, Issue 2* (2011), pgs. 309-332.

- [S2] Z.Shami, On uncountable hypersimple unidimensional theories, Arch. Math. Logic 53 (2014), no. 1-2, 203-210.
- [S3] Z.Shami, A model theoretic Baire category theorem for simple theories, Fundamenta Mathematicae 220 (2013), no. 3, 191-206.
- [W] Frank O. Wagner, Simple Theories, Academic Publishers, Dordrecht, The Netherlands, 2000.

Ziv Shami, E-mail address: zivsh@ariel.ac.il.