

ALLOCATION SCHEMES OF RESSOURCES WITH DOWNGRADING

CHRISTINE FRICKER, FABRICE GUILLEMIN, PHILIPPE ROBERT,
AND GUILHERME THOMPSON

ABSTRACT. We consider a server with large capacity delivering video files encoded in various resolutions. We assume that the system is under saturation in the sense that the total demand exceeds the server capacity C . In such a case, requests may be rejected. For the policies considered in this paper, instead of rejecting a video request, it is downgraded. When the occupancy of the server is above some value $C_0 < C$, the server delivers the video at a minimal bit rate. The quantity C_0 is the bit rate adaptation threshold. For these policies, request blocking is thus replaced by bit rate adaptation. Under the assumptions of Poisson request arrivals and exponential service times, we show that, by rescaling the system, a process associated to the occupancy of the server converges to some limiting process whose invariant distribution is computed by a Wiener-Hopf technique. This allows us to derive an asymptotic expression of the key performance measure of such a policy, the equilibrium probability that a request is transmitted without downgrading. Numerical applications of these results are presented.

CONTENTS

1. Introduction	1
2. Model description	2
3. Scaling Results	4
4. Invariant Distribution	10
5. Applications	15
References	16

1. INTRODUCTION

Video streaming applications have become over the past few years the dominant applications in the Internet and generate the prevalent part of traffic in today's IP networks; see for instance [6] for an illustration of the application breakdown in a commercial IP backbone network. Video files are currently downloaded by customers from large data centers, like Google's data centers for YouTube files. In the future, it is very likely that video files will be delivered by smaller data centers located closer to end users, for instance cache servers disseminated in a national network. It is worth noting that as shown in [7], caching is a very efficient solution for YouTube traffic. While this solution can improve performances by reducing

Date: Version of December 2, 2023.

2010 Mathematics Subject Classification. 60K25; 60K30 (primary), and 90B18 (secondary).

Key words and phrases. Resource Allocation; Scaling Methods; Loss Systems; Wiener-Hopf Problems.

delays, the limited capacity of those servers in terms of bandwidth and computing can cause overload.

One possibility to reduce overload is to use bit rate adaptation. Video files can indeed be encoded at various bit rates (e.g, small and high definition video). If a data center cannot serve a file at a high bit rate, then the video can be transmitted at a smaller rate. It is remarkable that video bit rate adaptation has become very popular in the past few years with the development of MPEG-DASH standard where it is possible to downgrade the quality of a given transmission, see Schwarz et al. [13] and Sieber et al. [14], see also Añorga et al. [1] and Vadlakonda et al. [15]. Adaptive streaming is frequently used in mobile networks where bandwidth is highly varying.

In this paper, we investigate the effect of bit rate adaptation in a data center under saturation. We assume that customers request video files encoded at various rates, say, A_j for $j=1, \dots, J$ with $A_1 < A_2 < \dots < A_J$. When there is not enough capacity to accommodate a request at rate A_j , $j > 1$ then the basic rate A_1 may be used to accommodate the request. This bit rate adaption reduces but does not completely eliminate losses of requests. In this paper, we introduce a bit rate adaptation threshold C_0 smaller than the server capacity C at which bit rate adaptation is applied when the occupation of the link is above C_0 .

To study this allocation scheme, a scaling approach is used. It is assumed that the server capacity is very large, namely scaled up by a factor N . The bit rate adaptation threshold and the request arrival rates are scaled up accordingly. This scaling has been introduced by Kelly for loss networks. See Kelly [10]. Despite the scaling approach is also used to study loss networks, a significant difference in the mathematical analysis is that the invariant distribution of the Markov process describing the state of the network does not seem to have a closed form expression. See Kelly [9]. The equilibrium behavior of these policies is investigated by studying the limiting transient properties of the associated Markov process. We consider a system in overload, due the bit rate adaptation threshold bit rate, requests may be downgraded but not rejected. As it will be seen, the stability properties of this algorithm are linked to the behavior of a Markov process associated to the occupation of the link. Under exponential assumptions for inter-arrival and service times, this process turns out to be, when the scaling parameter N is large, a bilateral random walk instead of a reflected random walk as in the case of loss networks. By using a Wiener-Hopf factorization, an explicit expression of the equilibrium of this random walk is obtained. With this result, the asymptotic expression of the probability that, at equilibrium, a job is transmitted at its required rate (and therefore does not experience a bit rate adaptation) is derived. See Corollary 1.

This paper is organized as follows: In Section 2, we present the model used to study a data center under saturation. Convergence results when the scaling factor N tends to infinity are proved in Section 3. The invariant distribution of a limiting process associated to the occupation of the link is computed in Section 4 by means of a Wiener-Hopf factorization. Some applications are discussed in Section 5.

2. MODEL DESCRIPTION

One considers a service system where J classes of requests arrive at a server with bandwidth/capacity C . Requests of class j , $1 \leq j \leq J$, arrive according to a Poisson process \mathcal{N}_{λ_j} with rate λ_j . A class j request has a bandwidth requirement

A_j for a duration of time which is exponentially distributed with parameter μ_j . For the networks investigated in this paper, there is no buffering, requests have to be processed at their arrival otherwise they are rejected. Without any flexibility on the resource allocation, this is a classical loss network with one link. See Kelly [10] for example.

This paper investigates allocation schemes which consist in reducing the bandwidth allocation of arriving requests to a minimal value when the link has a high level of congestion. In other words the service is downgraded for new requests arriving during a saturation phase. If the system is correctly designed, it will reduce significantly the fraction of rejected transmissions and, hopefully, few jobs will in fact experience downgrading.

2.1. Downgrading policy $\mathcal{D}(C_0)$. We introduce $C_0 < C$, the parameter C_0 will indicate the level of congestion of the link. If the total occupancy of the link is less than C_0 , then any arriving request is accepted. If the occupancy is between C_0 and $C-1$, it is accepted but with a minimal allocation. More precisely, it is assumed that the vector of integers $A=(A_j)$ is such that $A_1 = 1 < A_2 < \dots < A_J$. The condition $A_1=1$ is used to simplify the presentation of the results and to avoid problems of irreducibility in particular but this is not essential.

Let, for $1 \leq j \leq J$, ℓ_j be the number of class j requests at some moment and suppose that a new request of class $j_0 \in \{1, \dots, J\}$ arrives, then it is accepted if

$$\langle A, \ell \rangle \stackrel{\text{def.}}{=} A_1 \ell_1 + A_2 \ell_2 + \dots + A_J \ell_J < C_0.$$

Otherwise, if $C_0 \leq \langle A, \ell \rangle < C$, it is accepted but as a class 1 job. Finally it is rejected if the link is fully occupied, i.e. $\langle A, \ell \rangle = C$.

It is assumed that $\mu_1 \leq \mu_j$, for all $1 \leq j \leq J$, i.e. class 1 jobs are served with the smallest service rate. Note that, in practice, the various service rates should not be different since it concerns the duration of a given transmission and downgrading does not change this quantity.

Mathematically, the stochastic model is close to a loss network with the restriction that a job may change its requirements depending on the state of the network. This is a controlled loss network, see Zachary and Ziedins [17]. It does not seem that, like in uncontrolled loss networks, the associated Markov process has reversibility properties, or that its invariant distribution has a product form expression. Related schemes with product form are trunk reservation policies for which requests of a subset of classes are systematically rejected when the level of congestion of the link is above some threshold. See Bean et al. [3] and Zachary and Ziedins [16] for example. Concerning controlled loss networks, mathematical results are more scarce, one can mention networks where jobs requiring congested links are redirected to less loaded links. Several mathematical approximations have been proposed to study these models. See the surveys Kelly [10] and Zachary and Ziedins [17]. In our model, in the language of loss networks, the control is on the change of capacity requirement in the link instead of a change of link.

2.2. Scaling Regime. The invariant distribution being, in general, not known, a scaling approach is used. The network is investigated under Kelly's regime, i.e. under heavy traffic regime with a scaling factor N . It has been introduced in Kelly [9] to study the equilibrium of uncontrolled networks. The arrival rates are scaled by N : λ_j is replaced by $\lambda_j N$ as well as the capacity C^N and the threshold

C_0^N which are such that

$$C^N = cN + o(N) \text{ and } C_0^N = c_0N + o(N),$$

for $0 < c_0 < c$. It will be assumed that the system is overloaded for capacity C_0^N when the jobs have their initial bandwidth requirements. The relation

$$(R_1) \quad \langle \rho, A \rangle = \sum_{j=1}^J \rho_j A_j > c_0,$$

holds, with $\rho_j = \lambda_j / \mu_j$, $1 \leq j \leq J$. If this condition is not satisfied, it is not difficult to see that the system is equivalent to a classical underloaded loss network with one link and multiple classes of jobs. There is, of course, no need to use downgrading policies since the system can accommodate the incoming requests without any loss when N is large. See Kelly [10] or Section 7 of Chapter 6 of Robert [11] for example.

Additionally, it is also assumed that the system can accommodate the flow of requests when they have the minimal allocation $A_1=1$, i.e.

$$(R_2) \quad \frac{\Lambda}{\mu_1} < c_0,$$

with $\Lambda = \lambda_1 + \dots + \lambda_J$.

It should be noted that, from the point of view of the design of algorithms, the constant c_0 , which is not a capacity parameter, has to be defined. To take into account only the basic parameters of the model, the conditions (R_1) and (R_2) should be in fact replaced by the conditions

$$\langle \rho, A \rangle = \sum_{j=1}^J \rho_j A_j > c \text{ and } \frac{\Lambda}{\mu_1} < c.$$

The first condition gives that, without any change on the bandwidth requirement of jobs, the system will lose jobs. The second condition implies that the network can accommodate all jobs without losses (with high probability) if all of them require the reduced bandwidth $A_1 = 1$. It is easily seen then that, in this situation, one can find $0 < c' < c$ such that Conditions (R_1) and (R_2) hold for any $c_0 \in (c', c)$.

3. SCALING RESULTS

In this section, we prove convergence results when the scaling parameter N goes to infinity. These results are obtained by studying the asymptotic behavior of the occupation of the link translated by C_0^N , $m^N(t) = \langle A, L^N(t) \rangle - C_0^N$. In the context of loss networks, the analogue of such quantity is the number of empty places. As it will be seen, in the limit $(m^N(t))$ is related to a family of partially homogeneous random walks on \mathbb{Z} . The distribution of the increments of one of these random walks depends on the half-line, \mathbb{N} or $\mathbb{Z}_^*$, where it is located.

The following proposition shows that, for the downgrading policy, the boundary C^N does not play a role after some time if Condition (R_2) holds.

Proposition 1. *Under Condition (R_2) and if the initial state is such that*

$$\lim_{N \rightarrow +\infty} \left(\frac{L_j^N(0)}{N} \right) = \ell(0) = (\ell_{j,0}) \in \mathbb{R}_+^J,$$

then, for any $\varepsilon > 0$, there exists some $t_0 \geq 0$ such that, for any $T > t_0$,

$$\lim_{N \rightarrow +\infty} \mathbb{P} \left(\sup_{t_0 \leq t \leq T} \langle A, L^N(t) \rangle < (c_0 + \varepsilon)N \right) = 1.$$

Proof. Define

$$\left(\tilde{L}_j^N(t) \right) \stackrel{\text{def.}}{=} (D_1^N(t) + X^N(t), D_2^N(t), \dots, D_J^N(t)),$$

where, for $1 \leq j \leq J$,

$$D_j^N(t) = \sum_{k=1}^{L_j^N(0)} \mathbb{1}_{\{E_{\mu_j, k} > t\}},$$

$D_j^N(t)$ is the number of initial class j jobs still present at time t . Let $(X^N(t))$ be the process of the number of jobs of an independent $M/M/\infty$ queue with $X^N(0) = 0$, service rate μ_1 and the arrival rate $\Lambda = \lambda_1 + \dots + \lambda_J$. By using Theorem 6.13 of Robert [11], one gets the convergence in distribution

$$\lim_{N \rightarrow +\infty} \left(\frac{X^N(t)}{N} \right) = \frac{\Lambda}{\mu_1} (1 - e^{-\mu_1 t}),$$

and, consequently

$$(1) \quad \lim_{N \rightarrow +\infty} \left(\frac{1}{N} \langle A, \tilde{L}^N(t) \rangle \right) = \left(\frac{\Lambda}{\mu_1} (1 - e^{-\mu_1 t}) + \sum_{j=1}^J A_j \ell_{j,0} e^{-\mu_j t} \right).$$

Since $\mu_1 \leq \mu_j$ for all $1 \leq j \leq J$,

$$\begin{aligned} \frac{\Lambda}{\mu_1} (1 - e^{-\mu_1 t}) + \sum_{j=1}^J A_j \ell_{j,0} e^{-\mu_j t} \\ \leq \frac{\Lambda}{\mu_1} (1 - e^{-\mu_1 t}) + e^{-\mu_1 t} \langle A, \ell(0) \rangle \leq \max(c_0, \langle A, \ell(0) \rangle), \end{aligned}$$

by Condition (R_2) . Note that the asymptotic occupancy, when N is large, remains below the initial occupancy.

If $\langle A, \ell(0) \rangle > c_0 + \varepsilon$, let

$$\tau^N = \inf \left\{ t > 0 : \langle A, L^N(t) \rangle \leq C_0^N + \frac{\varepsilon}{2} N \right\},$$

then, on the event $\{\tau^N > T\}$, the downgrading policy gives that the identity in distribution

$$(2) \quad ((L_j^N(t)), 0 \leq t \leq T) \stackrel{\text{dist.}}{=} ((\tilde{L}_j^N(t)), 0 \leq t \leq T)$$

holds. Condition (R_2) gives the existence of t_0 such that

$$\frac{\Lambda}{\mu_1} (1 - e^{-\mu_1 t_0}) + \sum_{j=1}^J A_j \ell_{j,0} e^{-\mu_j t_0} = c_0 + \frac{\varepsilon}{2}.$$

Convergence (1) shows that the sequence (τ_N) converges in distribution to t_0 .

Note that, as long as the process $(\langle A, L^N(t) \rangle)$ is above C_0^N , Relation (2) holds. By using again Convergence (1), one gets that, as N goes to infinity, the process $(\langle A, L^N(t) \rangle / N)$ remains below $c_0 + \varepsilon$ with probability close to 1 on any finite time interval of $[t_0, +\infty)$. The proposition is proved. \square

Denote by $(m^N(t))$ the quantity

$$(3) \quad m^N(t) = \langle A, L^N(t) \rangle - C_0^N.$$

This is the analogue, for classical loss networks, of the process of the number of empty places in the link. Here this is an indication if the network is operating in saturation, $m^N(t) \geq 0$, or not, $m^N(t) < 0$. In loss networks, when N is large, up to a change of time scale, the process of the number of empty places converges to a reflected random walk in \mathbb{N} . In our case, the corresponding process is in fact a two-sided random walk which we introduce.

Definition 1. For $\ell = (\ell_j) \in \mathbb{R}_+^J$, let $(m_\ell(t))$ be the Markov process on \mathbb{Z} whose Q -matrix Q_ℓ is defined by, for $x \in \mathbb{Z}$ and $1 \leq j \leq J$,

$$(4) \quad \begin{cases} Q_\ell(x, x - A_j) = \mu_j \ell_j, \\ Q_\ell(x, x + A_j) = \lambda_j, \text{ if } x < 0, \\ Q_\ell(x, x + 1) = \Lambda, \text{ if } x \geq 0, \end{cases}$$

with $\Lambda = \lambda_1 + \lambda_2 + \dots + \lambda_J$.

The following proposition summarizes the stability properties of the Markov process $(m_\ell(t))$.

Proposition 2. If $\ell = (\ell_j) \in \mathbb{R}_+^J$, the Markov process $(m_\ell(t))$ is ergodic if $\ell \in \Delta_0$ with

$$(5) \quad \Delta_0 \stackrel{\text{def.}}{=} \left\{ x = (x_j) \in \mathbb{R}_+^J : \sum_{j=1}^J (\lambda_j - \mu_j x_j) A_j > 0 \text{ and } \sum_{j=1}^J \lambda_j - \mu_j x_j A_j < 0 \right\}.$$

Let π_ℓ be the corresponding invariant distribution.

Otherwise, if $\ell \in \Delta_-$,

$$\Delta_- \stackrel{\text{def.}}{=} \left\{ x = (x_j) \in \mathbb{R}_+^J : \sum_{j=1}^J (\lambda_j - \mu_j x_j) A_j \leq 0 \right\},$$

then the process $(m_\ell(t))$ converges in distribution to $\pi_\ell = \delta_{-\infty}$, the Dirac measure at $-\infty$. If $\ell \in \Delta_+$,

$$\Delta_+ \stackrel{\text{def.}}{=} \left\{ x = (x_j) \in \mathbb{R}_+^J : \sum_{j=1}^J \lambda_j - \mu_j x_j A_j \geq 0 \right\},$$

the process $(m_\ell(t))$ converges in distribution to $\pi_\ell = \delta_{+\infty}$.

Proof. The Markov process $(m_\ell(t))$ on \mathbb{Z} behaves like a random walk on each of the two half-lines \mathbb{N} and \mathbb{Z}_-^* . Definition (5) implies that if $\ell \in \Delta_0$, then the drift of the random walk in \mathbb{Z}_-^* is positive and negative when in \mathbb{N} which ensures the ergodicity of the Markov process by using the Lyapounov function $F(x) = |x|$, for example. See Corollary 8.7 of Robert [11] for example.

If $\ell \in \Delta_-$, then

$$\sum_{j=1}^J \lambda_j - A_j \mu_j \ell_j < \sum_{j=1}^J (\lambda_j - \mu_j \ell_j) A_j \leq 0,$$

this implies that the random walk on the half-line \mathbb{N} is ergodic. In particular, the duration of the excursions in \mathbb{N} are integrable. If

$$\sum_{j=1}^J (\lambda_j - \mu_j \ell_j) A_j < 0$$

there is a positive probability that, starting from -1 , the process never returns to 0 . In this case it therefore behaves as a random walk with negative drift, almost surely it converges to $-\infty$. If the inequality is not strict, i.e.

$$\sum_{j=1}^J (\lambda_j - \mu_j \ell_j) A_j = 0,$$

in the half-line \mathbb{Z}_-^* , the random walk is null recurrent and its invariant distribution is the Dirac mass at $-\infty$. The case $\ell \in \Delta_+$ is similar. \square

Stochastic Evolution Equations. For $\xi > 0$, denote by $\mathcal{N}_\xi(dt)$ a Poisson process on \mathbb{R}_+ with rate ξ and $(\mathcal{N}_{\xi,i}(dt))$ an i.i.d. sequence of such processes. All Poisson processes are assumed to be independent. Classically, the process $(L^N(t))$ can be seen as the solution of the following stochastic differential equations (SDE),

$$(6) \quad dL_1^N(t) = - \sum_{k=1}^{L_1^N(t-)} \mathcal{N}_{\mu_1,k}(dt) + \mathbb{1}_{\{m^N(t-) < C^N - C_0^N\}} \mathcal{N}_{\lambda_1 N}(dt) + \sum_{j=2}^J \mathbb{1}_{\{0 \leq m^N(t-) < C^N - C_0^N\}} \mathcal{N}_{\lambda_j N}(dt),$$

$$(7) \quad dL_j^N(t) = - \sum_{k=1}^{L_j^N(t-)} \mathcal{N}_{\mu_j,k}(dt) + \mathbb{1}_{\{m^N(t-) < 0\}} \mathcal{N}_{\lambda_j N}(dt), \quad 2 \leq j \leq J,$$

with initial condition $(L_j^N(0)) \in \mathbb{N}^J$.

Theorem 1 (Limiting Dynamical System). *Under Condition (R_2) , if the initial conditions are such that $m^N(0) = m \in \mathbb{Z}$ and*

$$\lim_{N \rightarrow +\infty} \left(\frac{L_j^N(0)}{N} \right) = (\ell_j(0)) \in \mathbb{R}_+^J,$$

then there exists a continuous process $(\ell(t)) = (\ell_j(t))$ such that the convergence in distribution

$$(8) \quad \lim_{N \rightarrow +\infty} \left(\left(\frac{L_j^N(t)}{N} \right), \int_0^t f(m^N(u)) du \right) = \left((\ell_j(t)), \int_0^t \int_{\mathbb{Z}} f(x) \pi_{\ell(u)}(dx) du \right)$$

holds for any function f with finite support on \mathbb{Z} . Furthermore, there exists $t_0 > 0$ such that $(\ell(t), t \geq t_0)$ satisfies the differential equations

$$(9) \quad \begin{cases} d\ell_1(t) = -\mu_1 \ell_1(t) dt + \lambda_1 + \sum_{j=2}^J \lambda_j \pi_{\ell(t)}(\mathbb{N}) dt, \\ d\ell_j(t) = -\mu_j \ell_j(t) dt + \lambda_j \pi_{\ell(t)}(\mathbb{Z}_-^*), \quad 2 \leq j \leq J, \end{cases}$$

where, for $\ell \in \mathbb{R}_+^J$, π_ℓ is the distribution associated to the process $(m_\ell(t))$ of Definition 1.

Proof. By using the same method as Hunt and Kurtz [8], one gets that an analogous result as Theorem 3 of this reference holds. Fix some $\varepsilon > 0$ such that $c_0 + \varepsilon < c$, from Proposition 1, one gets that the existence of t_0 such that

$$\lim_{N \rightarrow +\infty} \mathbb{P} \left(\sup_{t_0 \leq t \leq T} \langle A, L^N(t) \rangle < (c_0 + \varepsilon)N \right) = 1,$$

which implies that the boundary condition $m^N(t) < C^N - C_0^N$ in Equations (6) and (7) can be removed. Consequently, only the boundary condition of $(m^N(t))$ at 0 plays a role which gives Relation (9) as in Hunt and Kurtz [8]. Note that, contrary to the general situation described in this reference, we have indeed a convergence in distribution because, for any $\ell \in \mathbb{R}_+^J$, $(m_\ell(t))$ has exactly one invariant distribution (which may be a Dirac mass at infinity) by Proposition 2. See Conjecture 5 of [8]. \square

The following proposition gives a characterization of the fixed point of the dynamical system $(\ell_j(t))$.

Proposition 3 (Fixed Point). *Under Conditions (R_1) and (R_2) , there exists a unique fixed point ℓ^* of the process $(\ell_j(t))$ defined by Equation (8) given by*

$$(10) \quad \begin{cases} \ell_1^* = c_0 - \pi^-(\rho_2 A_2 + \rho_3 A_3 + \cdots + \rho_J A_J), \\ \ell_j^* = \rho_j \pi^-, \quad 2 \leq j \leq J, \end{cases}$$

where

$$(11) \quad \pi^- = \frac{c_0 - \Lambda/\mu_1}{\langle \rho, A \rangle - \Lambda/\mu_1}$$

with $\Lambda = \lambda_1 + \cdots + \lambda_J$. The process $(m_{\ell^*}(t))$ is ergodic in this case.

Proof. Assume that there exists a fixed point $\ell^* = (\ell_j^*)$ of $(\ell_j(t))$ defined by Equation (8), it is also a fixed point of the dynamical system defined by Equation (9), then

$$(12) \quad \begin{cases} \mu_1 \ell_1^* = \lambda_1 + (\lambda_2 + \lambda_3 + \cdots + \lambda_J)(1 - \pi^-), \\ \mu_j \ell_j^* = \lambda_j \pi^-, \end{cases}$$

with $\pi^- = \pi_{\ell^*}(\mathbb{Z}_+^*)$. One gets

$$\sum_{j=1}^J \lambda_j = \sum_{j=1}^J \mu_j \ell_j^* < \sum_{j=1}^J \mu_j \ell_j^* A_j = \pi^- \sum_{j=1}^J \lambda_j A_j + (1 - \pi^-) \sum_{j=1}^J \lambda_j < \sum_{j=1}^J \lambda_j A_j,$$

hence $\ell^* \in \Delta_0$, the associated process $(m_{\ell^*}(t))$ is necessarily ergodic by Proposition 2. We now show that the vector ℓ^* is on the boundary, i.e.

$$(13) \quad \sum_{j=1}^J A_j \ell_j^* = c_0.$$

If we assume that

$$\lim_{N \rightarrow +\infty} \left(\frac{\mathbf{L}_j^N(0)}{N} \right) = (\ell_j^*),$$

from Theorem 1 and the definition of $(m^N(t))$, we know that, for the convergence of processes, the following relation holds

$$\lim_{N \rightarrow +\infty} \left(\frac{m^N(t)}{N} \right) = (\kappa_0), \text{ with } \kappa_0 \stackrel{\text{def.}}{=} \sum_{j=1}^J A_j \ell_j^* - c_0.$$

For $N_0 \in \mathbb{N}$, $\varepsilon > 0$ and $N \geq N_0$,

$$\int_0^1 \mathbb{1}_{\{|m^N(u)| \geq \varepsilon N\}} du \leq \int_0^1 \mathbb{1}_{\{|m^N(u)| \geq \varepsilon N_0\}} du.$$

By using again Theorem 1 and the fact that ℓ^* is a fixed point of the dynamical system, we have, for the convergence in distribution

$$\lim_{N \rightarrow +\infty} \int_0^1 \mathbb{1}_{\{|m^N(u)| \leq \varepsilon N_0\}} du = \pi_{\ell^*}([- \varepsilon N_0, \varepsilon N_0]).$$

The left-hand side of the above expression can be arbitrarily close to 1 when N_0 is large. By convergence of the sequence $(m^N(t)/N)$ to (κ_0) , one gets that, for the convergence in distribution, the relation

$$\lim_{N \rightarrow +\infty} \int_0^1 \mathbb{1}_{\{|m^N(u)|/N \geq \varepsilon\}} du = 0$$

holds for all $\varepsilon > 0$, which implies that $\kappa_0 = 0$. Thus Relation (13) holds. Finally, Relations (12) and (13) give Relation (10).

To prove that the ℓ^* defined by Relations (10) and (11) is indeed a fixed point of the dynamical system defined by Equation (9), one has to show that the right-hand side of Equation (11) is indeed equal to $\pi_{\ell^*}(\mathbb{Z}_+^*)$. This is proved in Proposition 6 of Section 4. \square

Convergence of Invariant Distributions.

Proposition 4. *The equilibrium distribution of $(m^N(t))$ converges to π_{ℓ^*} when N goes to infinity, where ℓ^* is defined in Proposition 3.*

Proof. Recall that $m^N(t) = \langle A, L^N(t) \rangle - C_0^N$ and let Π^N be the invariant distribution of $(L^N(t))$. It is assumed that the distribution of $L^N(0)$ is Π^N for the rest of the proof. In particular $(m^N(t))$ is a stationary process.

One first proves that $(L^N(0)/N)$ converges in distribution to ℓ^* . The boundary condition $L^N(0) \leq C^N$ gives that the sequence of random variables $(L^N(0)/N)$ is tight. If $(L^{N_k}(0)/N_k)$ is a convergent subsequence to some random variable ℓ^∞ , by Theorem 1, one gets that, for the convergence in distribution, the relation

$$\lim_{k \rightarrow +\infty} \left(\left(\frac{L_j^{N_k}(t)}{N_k} \right) \right) = (\ell_j(t))$$

holds, where $(\ell_j(t))$ is a solution of Equation (9) with initial point at ℓ^∞ . Since $(L_j^N(t)/N)$ is a stationary process, the same property holds for $(\ell_j(t))$, this implies that ℓ^∞ is a fixed point of Equation (9), i.e. ℓ^∞ is ℓ^* by Proposition 3. Hence $(L^N(0)/N)$ converges in distribution to ℓ^* .

By using again Theorem 1, one gets that, for the convergence in distribution,

$$\lim_{N \rightarrow +\infty} \int_0^1 f(m^N(u)) du = \int_{\mathbb{Z}} f(x) \pi_{\ell^*}(dx)$$

holds for any function f with finite support on \mathbb{Z} . By using the stationarity of $(m^N(t))$ and Lebesgue's Theorem, one obtains

$$\lim_{N \rightarrow +\infty} \mathbb{E}(f(m^N(0))) = \int_{\mathbb{Z}} f(x) \pi_{\ell^*}(dx).$$

The proposition is proved. \square

Since a job arriving at time t is not downgraded if $m^N(t) < 0$, one obtains the following corollary.

Corollary 1. *The equilibrium probability that a job is not downgraded by the allocation scheme is converging to*

$$\pi^- \stackrel{\text{def.}}{=} \frac{c_0 - \Lambda/\mu_1}{\langle \rho, A \rangle - \Lambda/\mu_1}$$

as N goes to infinity.

4. INVARIANT DISTRIBUTION

We assume in this section that $\ell \in \Delta_0$, as defined in Proposition 2, so that $(m_\ell(t))$ is an ergodic Markov process. The goal of this section is to derive an explicit expression of the invariant distribution π_ℓ on \mathbb{Z} of $(m_\ell(t))$. At the same time, Proposition 6 below gives the required argument to complete the proof of Proposition 3 on the characterization of the fixed point of the dynamical system.

4.1. Functional Equation. In the following we denote by Y_ℓ a random variable with distribution π_ℓ .

For $r > 0$, we will use the notation

$$D(r) = \{z \in \mathbb{C}, |z| < r\}, \quad D^c(r) = \{z \in \mathbb{C}, |z| > r\} \text{ and } \gamma(r) = \{z \in \mathbb{C}, |z| = r\}.$$

For sake of simplicity, we will use $D = D(1)$ and $D^c = D^c(1)$.

Lemma 1. *With the notation*

$$\varphi_+(z) = \mathbb{E}(z^{Y_\ell} \mathbb{1}_{\{Y_\ell \geq 0\}}), \quad \varphi_-(z) = \mathbb{E}(z^{Y_\ell} \mathbb{1}_{\{Y_\ell < 0\}}),$$

the random variable Y_ℓ is such that

$$(14) \quad P(z)\varphi_+(z) = Q(z)\varphi_-(z)$$

where P and Q are polynomials defined by

$$(15) \quad \begin{aligned} P(z) &= \sum_{j=1}^J ((\lambda_j + \mu_j \ell_j) z^{A_j} - \lambda_j z^{A_j+1} - \mu_j \ell_j z^{A_j-A_j}) \text{ and} \\ Q(z) &= \sum_{j=1}^J (\lambda_j z^{A_j+A_j} + \mu_j \ell_j z^{A_j-A_j} - (\lambda_j + \mu_j \ell_j) z^{A_j}). \end{aligned}$$

Proof. For $z \in \gamma(1)$ define $f_z : \mathbb{Z} \mapsto \mathbb{C}$ such that $f_z(x) = z^x$, for $x \in \mathbb{Z}$. Equilibrium equations for $(m_\ell(t))$ give the identity

$$\langle \pi_\ell, Q_\ell(f_z) \rangle = \sum_{x,y \in \mathbb{Z}} \pi_\ell(x) Q_\ell(x,y) (f_z(y) - f_z(x)) = 0,$$

where Q_ℓ is the Q -matrix of $(m_\ell(t))$ given by Equation (4). After some simple reordering, one gets the relation

$$(16) \quad \mathbb{E}(z^{Y_\ell} \mathbb{1}_{\{Y_\ell \geq 0\}}) \sum_{j=1}^J (\lambda_j(1-z) + \mu_j \ell_j(1-z^{-A_j})) = \\ - \mathbb{E}(z^{Y_\ell} \mathbb{1}_{\{Y_\ell < 0\}}) \sum_{j=1}^J (\lambda_j(1-z^{A_j}) + \mu_j \ell_j(1-z^{-A_j})).$$

By using the definition of $\varphi_+(z)$ and $\varphi_-(z)$, Equation (16) can be rewritten as Equation (14). \square

Relation (14) is valid on the unit circle, however the function φ_+ (resp. φ_-) is defined on D (resp. D^c). This can then be expressed as a Wiener-Hopf factorization problem analogous to the one used in the analysis of reflected random walks on \mathbb{N} . This is used in the analysis of the $GI/GI/1$ queue, see Chapter VIII of Asmussen [2] or Chapter 3 of Robert [11] for example. In a functional context, this is a special case of a Riemann's problem, see Gakhov [5]. In our case, this is a kind of doubly random walk in \mathbb{Z} , with a different drift depending on the half-line. The first (resp. second) condition in the definition of the set Δ_0 in Definition (5) implies that the drift of the random walk in \mathbb{Z}_+ (resp. in \mathbb{N}) is positive (resp. negative).

The first step in the analysis of Equation (14) is of determining the locations of the zeroes of P and Q . This is the purpose of the following lemma.

Lemma 2. *[Location of the Zeroes of P and Q] Let ℓ be in Δ_0 .*

- (i) *Polynomial Q has exactly two positive real roots 1 and $q_1 \in]0, 1[$. There are $A_J - 1$ roots in $D(q_1)$ and $A_J - 1$ roots with a modulus strictly greater than 1.*
- (ii) *Polynomial P has exactly two positive real roots 1 and $p_1 > 1$. The $A_J - 1$ remaining roots have a modulus strictly smaller than 1.*

Proof. One first notes that Q is a polynomial with the same form as the f defined by Equation (13) in Bean et al. [3] (with $e_j = A_j$, $\kappa_j = \lambda_j$ and $\hat{e} = A_J$). The roots of Q are exactly the roots of f . Lemma 2.2 of Bean et al. [3] gives assertion (i) of our lemma.

The proof of assertion (ii) uses an adaptation of the argument for the proof of Lemma 2.2 of Bean et al. [3]. Define the function f by $f(z) = z^{-A_J} P(z)$. Recall that P is a polynomial with degree $A_J + 1$. There are exactly two real positive roots for P . Indeed, $f(1) = 0$ and it is easily checked that f is strictly concave with

$$f'(1) = \sum_{j=1}^J (-\lambda_j + A_j \mu_j \ell_j) > 0,$$

since $\ell \in \Delta_0$, by the second condition in the definition of the set Δ_0 in Definition (5). Hence P has a real zero p_1 greater than 1.

Let $r \in]1, p_1[$ be fixed, note that $P(r) > 0$. Define

$$f_1(z) = K z^{A_J}, \text{ with } K = \sum_{j=1}^J (\lambda_j + A_j \mu_j \ell_j), \\ f_2(z) = \sum_{j=1}^J (\lambda_j z^{A_J+1} + \mu_j \ell_j z^{A_J-A_j}),$$

so that $P = f_1 - f_2$.

Fix some $z \in \gamma(r)$. By expressing these functions in terms of real and imaginary parts,

$$z^{A_J} = \alpha_1 + i\beta_1 \text{ and } f_2(z) = \alpha_2 + i\beta_2,$$

one gets

$$(17) \quad |f_1(z) - f_2(z) - bz^{A_J}|^2 = |K(\alpha_1 + i\beta_1) - b(\alpha_1 + i\beta_1) - (\alpha_2 + i\beta_2)|^2 \\ = (K\alpha_1 - \alpha_2)^2 + (K\beta_1 - \beta_2)^2 + H = |f_1(z) - f_2(z)|^2 + H,$$

with

$$H = (b\alpha_1)^2 - 2b\alpha_1(K\alpha_1 - \alpha_2) + (b\beta_1)^2 - 2b\beta_1(K\beta_1 - \beta_2) \\ = b(b - 2K)(\alpha_1^2 + \beta_1^2) + 2b(\alpha_1\alpha_2 + \beta_1\beta_2).$$

Cauchy-Schwarz's Inequality gives the relation

$$\alpha_1\alpha_2 + \beta_1\beta_2 \leq \frac{1}{K}|f_2(z)||f_1(z)| \leq \frac{1}{K}f_2(r)f_1(r),$$

since $|f_i(z)| \leq f_i(|z|)$ for $i = 1, 2$. Thus,

$$\frac{H}{b} = (b - 2K)(\alpha_1^2 + \beta_1^2) + 2(\alpha_1\alpha_2 + \beta_1\beta_2) \leq (b - 2K)\frac{f_1(r)^2}{K^2} + 2f_2(r)\frac{f_1(r)}{K} \\ = \frac{f_1(r)}{K^2} \left((b - 2K)f_1(r) + 2Kf_2(r) \right) = \frac{f_1(r)}{K^2} (bf_1(r) - 2KP(r)).$$

Since $P(r) > 0$, b can be chosen so that $bf_1(r) < 2KP(r)$. From the above relation and Equation (17), one gets that for $z \in \gamma(r)$, the relation

$$|f_1(z) - f_2(z) - bz^{A_J}| < |f_1(z) - f_2(z)|$$

holds. By Rouché's theorem, one obtains that, for any $r \in]1, p_1[$, P has exactly A_J roots in $D(r)$. One concludes that P has exactly A_J roots in \bar{D} . It is easily checked that if $|z|=1$ and $z \notin \mathbb{R}$ then the real part of $P(z)$ is positive, hence z cannot be a root of the polynomial P . Consequently, P has exactly $A_J - 1$ roots in D . The lemma is proved. \square

4.2. Wiener-Hopf factorization.

Definition 2. For $U \in \{P, Q\}$, denote by \mathcal{Z}_U the set of the zeroes of U different from 1.

Define

$$(18) \quad \Phi(z) = \begin{cases} -\varphi_+(z)\lambda_J^{-1}(z - p_1) \prod_{q \in \mathcal{Z}_Q \cap D^c} (z - q)^{-1}, & z \in D \\ \varphi_-(z)\Lambda^{-1} \prod_{q \in \mathcal{Z}_Q \cap D} (z - q) \prod_{p \in \mathcal{Z}_P \cap D} (z - p)^{-1}, & z \notin D \end{cases}$$

with $\Lambda = \lambda_1 + \dots + \lambda_J$ and the same notations as before. By definition, function Φ is holomorphic in D and D^c and, from Relation (14), is continuous on $\gamma(1)$. The analytic continuation theorem, Theorem 16.8 of Rudin [12] for example, gives that Φ is holomorphic on \mathbb{C} . For $z \in D^c$,

$$|\varphi_-(z)| \leq \mathbb{E} \left(\mathbb{1}_{\{Y_\ell < 0\}} |z|^{Y_\ell} \right) \leq \frac{1}{|z|},$$

since the cardinality of $\mathcal{Z}_P \cap D$ (resp. $\mathcal{Z}_Q \cap D$) is $A_J - 1$ (resp. A_J), the holomorphic function Φ is therefore bounded on \mathbb{C} . By Liouville's theorem, Φ is constant, equal to $\kappa \in \mathbb{C}$. Therefore

$$(19) \quad \begin{cases} \varphi_+(z) &= -\kappa \lambda_J (z - p_1)^{-1} \prod_{q \in \mathcal{Z}_Q \cap D^c} (z - q), & z \in D, \\ \varphi_-(z) &= \kappa \Lambda \prod_{q \in \mathcal{Z}_Q \cap D} (z - q)^{-1} \prod_{p \in \mathcal{Z}_P \cap D} (z - p), & z \in D^c. \end{cases}$$

Recall that $\varphi(z) = \varphi_+(z) + \varphi_-(z) = \mathbb{E}(z^{Y_\ell})$ is a generating function, in particular $\varphi(1) = 1$. Plugging the previous expressions for φ_+ and φ_- in $\varphi_+(1) + \varphi_-(1) = 1$, one gets the relation

$$1 = -\kappa \prod_{q \in \mathcal{Z}_Q \cap D} (1 - q)^{-1} \frac{1}{1 - p_1} (Q'(1) + P'(1)),$$

hence, using equation (15),

$$\kappa = \frac{p_1 - 1}{\langle A - e, \lambda \rangle} \prod_{q \in \mathcal{Z}_Q \cap D} (1 - q),$$

where $e = (1, 1, \dots, 1) \in \mathbb{N}^J$. Note that κ is positive. We can now state the main result of this section.

Proposition 5 (Invariant Measure). *Under the condition $\ell \in \Delta_0$ defined by Relation (5), the invariant measure π_ℓ can be expressed, for $n \in \mathbb{Z}$, as*

$$\pi_\ell(n) = \begin{cases} -\kappa \sum_{q \in \mathcal{Z}_Q \cap D} \frac{P(q)q^{-n-1}}{(q - p_1)(q - 1)Q_D(q)}, & n < 0, \\ \kappa \left(\alpha_n + \frac{Q(p_1)p_1^{-n-1}}{(p_1 - 1)Q_D(p_1)} \right), & 0 \leq n < A_J - 1, \\ \kappa \frac{Q(p_1)p_1^{-n-1}}{(p_1 - 1)Q_D(p_1)}, & n \geq A_J - 1, \end{cases}$$

where p_1 is defined in Lemma 2, P and Q by Relation (15),

$$Q_D(z) = \prod_{q \in \mathcal{Z}_Q \cap D} (z - q), \quad \kappa = \frac{(p_1 - 1)Q_D(1)}{\langle A - 1, \lambda \rangle},$$

and, for $0 \leq n < A_J - 1$, α_n is the coefficient of degree n of the polynomial

$$-\frac{1}{z - p_1} \left(\frac{Q(z)}{(z - 1)Q_D(z)} - \frac{Q(p_1)}{(p_1 - 1)Q_D(p_1)} \right).$$

Proof. Note that, for $z \in \mathbb{C}$,

$$\prod_{p \in \mathcal{Z}_P \cap D} (z - p) = -\frac{1}{\Lambda} \frac{P(z)}{(z - p_1)(z - 1)}.$$

For $z \in D^c$,

$$\varphi_-(z) = \kappa \Lambda \prod_{q \in \mathcal{Z}_Q \cap D} (z - q)^{-1} \prod_{p \in \mathcal{Z}_P \cap D} (z - p).$$

Since $|\mathcal{Z}_P \cap D| = A_J - 1 < A_J = |\mathcal{Z}_Q \cap D|$ by Lemma 2, φ_- has the following partial fraction decomposition

$$\begin{aligned}\varphi_-(z) &= -\kappa \sum_{q \in \mathcal{Z}_Q \cap D} \frac{P(q)}{(q-p_1)(q-1)Q'_D(q)} \frac{1}{z-q} \\ &= \sum_{i=0}^{\infty} -\kappa \sum_{q \in \mathcal{Z}_Q \cap D} \frac{P(q)q^i}{(q-p_1)(q-1)Q'_D(q)} \frac{1}{z^{i+1}}.\end{aligned}$$

Denote

$$Q_{D^c}(z) = \prod_{q \in \mathcal{Z}_Q \cap D^c} (z-q) = \frac{Q(z)}{\lambda_J(z-1)Q_D(z)},$$

then

$$\varphi_+(z) = -\kappa \lambda_J \frac{Q_{D^c}(z)}{z-p_1} = \kappa \left(-\lambda_J \frac{Q_{D^c}(z) - Q_{D^c}(p_1)}{z-p_1} + \frac{Q(p_1)}{(1-p_1)Q_D(p_1)} \frac{1}{z-p_1} \right).$$

One concludes by using the expression of κ obtained before. \square

Proposition 6. *If $\ell^* \in \mathbb{R}_+^J$ is given by Relation (10) then*

$$\pi_{\ell^*}(\mathbb{Z}_-^*) = \frac{c_0 - \Lambda/\mu_1}{\langle \rho, A \rangle - \Lambda/\mu_1}.$$

Note that the right hand side of the last relation is precisely π^- of Relation (11) which is the result necessary to complete the proof of Proposition 3.

Proof. With the same notations as before, from Relation (14),

$$\frac{\varphi_-(z)}{\varphi_+(z)} = \frac{P(z)}{Q(z)}$$

holds for $z \in \mathbb{C}$, with $|z| = 1$. By definition of $\varphi_-(z)$ and $\varphi_+(z)$,

$$\lim_{z \rightarrow 1} \varphi_-(z) = \pi_{\ell^*}(\mathbb{Z}_-^*) \text{ and } \lim_{z \rightarrow 1} \varphi_+(z) = 1 - \pi_{\ell^*}(\mathbb{Z}_-^*).$$

Since 1 is a zero of P and Q , this gives the relation

$$\frac{\pi_{\ell^*}(\mathbb{Z}_-^*)}{1 - \pi_{\ell^*}(\mathbb{Z}_-^*)} = \frac{P'(1)}{Q'(1)} = \frac{\sum_{j=1}^J (A_j \mu_j \ell_j^* - \lambda_j)}{\sum_{j=1}^J A_j (\lambda_j - \mu_j \ell_j^*)}.$$

By using the expression of (ℓ_j^*) , with some algebra, one gets

$$\pi_{\ell^*}(\mathbb{Z}_-^*) = \left(c_0 - \sum_{j=1}^J \frac{\lambda_j}{\mu_1} \right) \bigg/ \left(\sum_{j=1}^J \rho_j A_j - \sum_{j=1}^J \frac{\lambda_j}{\mu_1} \right) = \pi^-.$$

The proposition is proved. \square

4.3. Some Moments of (π_{ℓ^*}) . Using the probability generating function $\varphi(z)$ of π_{ℓ^*} from Equation (19), one can derive an explicit expression of the mean, the variance and the skewness of such distribution. The skewness of a random variable X is a measure of the asymmetry of the distribution of X ,

$$\text{Skew}(X) \stackrel{\text{def.}}{=} \frac{\mathbb{E}([X - \mathbb{E}(X)]^3)}{\text{Var}(X)^{3/2}}.$$

See Doane and Seward [4] for example.

Proposition 7. *If Y_{ℓ^*} is a random variable with distribution π_{ℓ^*} then*

$$\begin{aligned}\mathbb{E}(Y_{\ell^*}) &= A_J + \frac{\theta_2}{2\theta_1} - S(1), \\ \text{Var}(Y_{\ell^*}) &= \frac{\theta_2 + 2\theta_3}{6\theta_1} - \left(\frac{\theta_2}{2\theta_1}\right)^2 - (S(1) + S'(1)), \\ \text{Skew}(Y_{\ell^*}) &= \frac{\theta_2^3}{4\theta_1^3} + \theta_2 \frac{\theta_2 - 2\theta_3}{4\theta_1^2} + \frac{\theta_4 - \theta_3}{4\theta_1} - (S(1) + 3S'(1) + S''(1)),\end{aligned}$$

where, for $i \geq 1$,

$$\theta_i = \sum_{j=2}^J \lambda_j A_j^{i-1} (A_j - 1),$$

and

$$S(z) = \frac{1}{z - p_1} + \sum_{q \in \mathcal{Z}_Q \cap D} \frac{1}{z - q},$$

with $Q_D(z)$ defined in Proposition 5.

The proof is straightforward, modulo some tedious calculations of the successive derivatives of $\varphi(z)$ evaluated at 1. Figure 1 shows that the distribution of Y_{ℓ^*} is significantly asymmetrical. For this example $\mathbb{E}(Y_{\ell^*})=8.04819$, $\text{Var}(Y_{\ell^*})=77.2284$ and $\text{Skew}(Y_{\ell^*}) = 0.967069$.




FIGURE 1. The histogram of Y_{ℓ^*} with the parameters $J=5$, $A=(1, 2, 4, 8, 16)$, $\lambda=(0.25, 0.2, 0.15, 0.1, 0.05)$, $\mu=(1, 1, 1, 1, 1)$ and $c_0=0.97$.

5. APPLICATIONS

In this section, we consider a link with large bandwidth, $C = 10.0$ Gbps, in charge of video streaming. Requests that cannot be immediately served are lost. Video transmission is offered in two standard qualities, namely, *Low Quality* (LQ) and *High Quality* (HQ). From Añorga *et al.* [1], the bandwidth requirement for YouTube's videos at 240p is 1485.11 Kbps, and for 720p it is 2737.27 Kbps.

Using the values above, after renormalization, one takes $A_1 = 1$, $C = 7060$ and $A_2 = 2$. Jobs arrive into this system asking for HQ transmission, but clients accept to watch the video in LQ. In particular $\lambda_1 = 0$. Service times are assumed to be the same in both qualities and taken as the unity, $\mu_1 = \mu_2 = 1$. We define $C_0 = \alpha C$,

with $0 < \alpha < 1$ and the quantity α_ε as the largest value to start degradation, such that the blocking probability of a job is less than some $\varepsilon > 0$. With the notations of Section 4, we write

$$\alpha_\varepsilon = \sup \{ \alpha \in (0, 1) : \mathbb{P}(Y_{\ell^*} > (1 - \alpha)C) < \varepsilon \}.$$




FIGURE 2. Maximal threshold $\alpha_\varepsilon = C_0/C$ for a loss probability $\leq \varepsilon$.

Figure 2 plots the threshold α_ε that ensures a loss rate less than ε as a function of ε for several values of λ_2 . In the numerical example, a 2% reservation is sufficient to get a blocking probability less than 10^{-7} even when system load is above the saturation limit.

Now let π_ε^- be the value of π^- defined by Corollary 1 for $C_0 = \alpha_\varepsilon C$,

$$\pi_\varepsilon^- = \frac{\alpha_\varepsilon}{\lambda_2} - 1.$$

Note that π_ε^- is the asymptotic equilibrium probability that a job does not suffer degradation given that the system operates with loss probabilities inferior to ε . For comparison, β is the corresponding loss probability when no control is used in the system. We show in Figure 3 the relation between these quantities and the workload λ_2 , for fixed loss rates of 10^{-3} , 10^{-6} and 10^{-9} . We have $\beta = 1/(2\lambda_2)$, see [11, Section 6.7]. The curves plotting π_ε^- for $\varepsilon = 10^{-3}$, 10^{-6} and 10^{-9} are in fact very close. As in Figure 2, it implies that very low loss rates are achievable without expanding the degradation zone $[C_0, C]$. The quantity C_0 can be fixed very close to C with very few losses. Nevertheless, one remarks that, for high loads, the system cannot hold these demands, because our policy is no longer effective. Still, even with C less than 10^4 , π_ε^- is very close to the limiting value of π^- as C gets large, especially when the system is not working beyond its capacity.

REFERENCES

- [1] Javier Añorga, Saioa Arrizabalaga, Beatriz Sedano, Maykel Alonso-Arce, and Jaizki Mendizabal. YouTube's DASH implementation analysis. In *Proceedings of the 19th International Conference on Communications*, volume 50 of *Recent Advances in Electrical Engineering Series*, pages 61–66, 06 2015.




FIGURE 3. Fraction of non-downgraded jobs at equilibrium.

- [2] Søren Asmussen. *Applied probability and queues*, volume 51 of *Applications of Mathematics*. Springer-Verlag, New York, second edition, 2003.
- [3] N. G. Bean, R. J. Gibbens, and S. Zachary. Asymptotic analysis of single resource loss systems in heavy traffic, with applications to integrated networks. *Advances in Applied Probability*, 27(1):273–292, 1995.
- [4] David P. Doane and Lori E. Seward. Measuring skewness: A forgotten statistic? *Journal of Statistics Education*, 19(2):1–18, 2011.
- [5] F. D. Gakhov. *Boundary value problems*. Dover Publications Inc., New York, 1990. Translated from the Russian, Reprint of the 1966 translation.
- [6] Fabrice Guillemin, Thierry Houdoin, and Stéphanie Moteau. Volatility of YouTube content in Orange networks and consequences. In *Proceedings of IEEE International Conference on Communications, ICC 2013, Budapest, Hungary, June 9-13, 2013*, pages 2381–2385, 2013.
- [7] Fabrice Guillemin, Bruno Kauffmann, Stéphanie Moteau, and Alain Simonian. Experimental analysis of caching efficiency for YouTube traffic in an ISP network. In *25th International Teletraffic Congress, ITC 2013, Shanghai, China, September 10-12, 2013*, pages 1–9, 2013.
- [8] P.J. Hunt and T.G Kurtz. Large loss networks. *Stochastic Processes and their Applications*, 53:363–378, 1994.
- [9] F.P. Kelly. Blocking probabilities in large circuit-switched networks. *Advances in Applied Probability*, 18:473–505, 1986.
- [10] F.P. Kelly. Loss networks. *Annals of Applied Probability*, 1(3):319–378, 1991.
- [11] Philippe Robert. *Stochastic Networks and Queues*, volume 52 of *Stochastic Modelling and Applied Probability Series*. Springer, New-York, June 2003.
- [12] Walter Rudin. *Real and complex analysis*. McGraw-Hill Book Co., New York, third edition, 1987.
- [13] H. Schwarz, D. Marpe, and T. Wiegand. Overview of the scalable video coding extension of the H.264/AVC standard. *IEEE Trans. Cir. and Sys. for Video Technol.*, 17(9):1103–1120, September 2007.
- [14] Christian Sieber, Tobias Hofeld, Thomas Zinner, Phuoc Tran-Gia, and Christian Timmerer. Implementation and user-centric comparison of a novel adaptation logic for DASH with SVC. In *IM'13*, pages 1318–1323, 2013.
- [15] S. Vadlakonda, A. Chotai, B.D. Ha, A. Asthana, and S. Shaffer. System and method for dynamically upgrading / downgrading a conference session, April 6 2010. US Patent 7,694,002.
- [16] Stan Zachary and Ilze Ziedins. A refinement of the Hunt-Kurtz theory of large loss networks, with an application to virtual partitioning. *The Annals of Applied Probability*, 12(1):1–22, 02 2002.

- [17] Stan Zachary and Ilze Ziedins. Loss networks. In Richard J. Boucherie and Nico M. van Dijk, editors, *Queueing Networks*, volume 154 of *International Series in Operations Research & Management Science*, pages 701–728. Springer US, 2011.

(F. Guillemin) CNC/NCA ORANGE LABS2, AVENUE PIERRE MARZIN, 22300 LANNION, FRANCE
E-mail address: `Fabrice.Guillemin@orange.com`

(C. Fricker, Ph. Robert, G. Thompson) INRIA PARIS, 2 RUE SIMONE IFF, CS 42112, 75589
PARIS CEDEX 12, FRANCE
E-mail address: `Philippe.Robert@inria.fr`
URL: `http://www-rocq.inria.fr/~robert`