

Confidence Bands for the Logistic and Probit Regression Models Over Intervals

Lucy Kerns

Department of Mathematics and Statistics

Youngstown State University

Youngstown OH 44555

xlu@ysu.edu

Summary

This article presents methods for the construction of two-sided and one-sided simultaneous hyperbolic bands for the logistic and probit regression models when the predictor variable is restricted to a given interval. The bands are constructed based on the asymptotic properties of the maximum likelihood estimators. Past articles have considered building two-sided asymptotic confidence bands for the logistic model, such as Piegorsch and Casella (1988). However, the confidence bands given by Piegorsch and Casella are conservative under a single interval restriction, and it is shown in this article that their bands can be sharpened using the methods proposed here. Furthermore, no method has yet appeared in the literature for constructing one-sided confidence bands for the logistic model, and no work has been done for building confidence bands for the probit model, over a limited range of the predictor variable. This article provides methods for computing critical points in these areas.

Keywords: One-sided confidence bands; Two-sided confidence bands; Logistic regression; Probit regression; Simple linear regression

1 Introduction

Logistic and probit regression models are widely used for modeling dichotomous outcomes, and have been increasingly applied in medical research, public health research, environmental science, and many other fields, such as human behavior modeling (Chou, Lu, and Mao, 2002), environmental modeling (Pradhan and Lee, 2010), and biomedical research (Austin and Steyerberg, 2014). The logistic and probit regression models are statistical methods that allow one to estimate the response probability for a dichotomous response, that is, a response which is binary, taking values 1 (success, normal, positive, etc.) and 0 (failure, abnormal, negative, etc.). In this article, we consider the case where the binary response variable Y is determined by a predictor variable x , and the response probability is:

$$P(Y = 1) = p(x) = \begin{cases} 1/[1 + \exp(-\mathbf{c}'\boldsymbol{\beta})], & \text{logistic model} \\ \Phi(\mathbf{c}'\boldsymbol{\beta}), & \text{probit model} \end{cases}$$

where $\mathbf{c} = (1 \ x)'$, $\boldsymbol{\beta} = (\beta_0 \ \beta_1)'$, and Φ is the cumulative distribution function (cdf) of the standard normal distribution.

The construction of confidence bands on $p(x)$ is often of interest, and we will use the following link functions to transform the problem of constructing confidence bands for the response probability $p(x)$ to the problem of constructing confidence bands for the linear predictor $\mathbf{c}'\boldsymbol{\beta}$ on which the bands are defined.

$$\text{logit}(p(x)) = \log_e \left[\frac{p(x)}{1 - p(x)} \right] = \mathbf{c}'\boldsymbol{\beta}, \text{ logistic regression}$$

$$\text{probit}(p(x)) = \Phi^{-1}(p(x)) = \mathbf{c}'\boldsymbol{\beta}, \text{ probit regression}$$

There is a wealth of literature on building exact or conservative two-sided bands for the linear predictor $\mathbf{c}'\boldsymbol{\beta}$ with one or more than one predictor variable. This includes the work of Scheffé (1953), and Working & Hotelling (1929), among others. Many authors have improved earlier work by restricting the predictor variables to given intervals, including Wynn and Bloomfield (1971), Casella and Strawderman (1980), and Uusipaikka (1983). Liu *et al.* (2005) developed a simulation-based method for constructing two-sided confidence bands over a restricted region for the multiple regression model. Less work has

been done on the construction of one-sided (lower or upper) confidence bands for linear regression. With no restriction on the predictor variable, Hochberg and Quade (1975) developed a method for constructing one-sided bands in the multiple regression setting. When the predictor variable is constrained to a pre-specified interval, Bohrer and Francis (1972) gave exact one-sided hyperbolic confidence bands for the simple linear case. Pan *et al.* (2003) extended Uusipaikka's work (1983) to the computation of one-sided bands under a single interval restriction. Liu *et al.*'s article (2008) summarized several existing methods and also provided new methods for the construction of two-sided and one-sided confidence bands with a restricted predictor variable.

Much less work has been done on constructing confidence bands for the logistic regression model. Brand, Pinnock, and Jackson (1973) constructed confidence bands for both $p(x)$ and the inverse of $p(x)$ when there is one predictor variable and no constraints exist on the predictor variable. Hauck (1983) extended their work to more than one predictor variable yet still with no constraints. In the case of restricted predictor variables, Piegorsch and Casella (1988) extended the work of Casella and Strawderman (1980) from the multiple regression model to the logistic case. The method of Casella and Strawderman produced asymptotic two-sided bands over a particular constrained region of predictor variables. However, if the restricted region on predictor variables is rectangular, the resulting confidence bands are rather conservative even with large samples.

Wei Liu's book (2010, Chapter 8) presented confidence bands for the logistic model with more than one explanatory variable. The method utilized simulation-based confidence bands (Liu *et al.* (2005)) for the linear predictor $\mathbf{c}'\boldsymbol{\beta}$ in the multiple linear model, and the desired bands for the logistic model were then obtained via the logit link function. The new contribution of this paper is that we provide explicit expressions for determining the critical values in the logistic setting. Wei Liu's method is very broad, but relies on simulation. Our method is more focused but admits tractable forms.

Furthermore, no work has yet appeared for building one-sided confidence bands for the logistic regression model when the predictor variable is bounded on a given interval, and no methods have been developed for constructing both two-sided and one-sided bands for the probit regression model with a restricted predictor variable. In this paper, we center our

attention on building asymptotic two-sided and one-sided hyperbolic confidence bands for the logistic and probit models over a limited range of the predictor variable, say, between (a, b) , where a and b are constants.

For the logistic and probit models, we denote the ML estimator of $\boldsymbol{\beta}$ by $\hat{\boldsymbol{\beta}}$, and under certain regularity conditions (Kendall and Stuart (1979)), $\hat{\boldsymbol{\beta}}$ follows asymptotically as:

$$\hat{\boldsymbol{\beta}} \sim ASN_2(\boldsymbol{\beta}, \mathbf{F}^{-1}),$$

where \mathbf{F} is the Fisher information matrix. It is well known that the logistic model has an explicit formula for the Fisher information matrix, while the probit model does not, but the information matrix can be obtained through numerical methods.

For $x \in (a, b)$, where a and b are given constants, a $100(1 - \alpha)\%$ two-sided hyperbolic band for the linear function $\mathbf{c}'\boldsymbol{\beta}$ has the form

$$\mathbf{c}'\boldsymbol{\beta} \in \mathbf{c}'\hat{\boldsymbol{\beta}} \pm w(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2},$$

where $\mathbf{c} = (1 \quad x)'$ and w satisfies

$$P[\mathbf{c}'\boldsymbol{\beta} \in \mathbf{c}'\hat{\boldsymbol{\beta}} \pm w(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2}, \text{ for all } x \in (a, b)] = 1 - \alpha. \quad (1)$$

An upper $100(1 - \alpha)\%$ one-sided hyperbolic band for the linear function $\mathbf{c}'\boldsymbol{\beta}$ has the form

$$\mathbf{c}'\boldsymbol{\beta} \leq \mathbf{c}'\hat{\boldsymbol{\beta}} + w_u(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2},$$

where w_u satisfies

$$P[\mathbf{c}'\boldsymbol{\beta} \leq \mathbf{c}'\hat{\boldsymbol{\beta}} + w_u(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2}, \text{ for all } x \in (a, b)] = 1 - \alpha. \quad (2)$$

A $100(1 - \alpha)\%$ lower one-sided hyperbolic band for $\mathbf{c}'\boldsymbol{\beta}$ can be defined similarly.

Since both logit and probit functions are monotonically increasing functions, $100(1 - \alpha)\%$ two-sided confidence bands for $p(x)$ in the logistic regression model are given by

$$\{1 + \exp[-\mathbf{c}'\hat{\boldsymbol{\beta}} + w(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2}]\}^{-1} \leq p(x) \leq \{1 + \exp[-\mathbf{c}'\hat{\boldsymbol{\beta}} - w(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2}]\}^{-1},$$

and $100(1 - \alpha)\%$ two-sided confidence bands for $p(x)$ in the probit regression model are given by

$$\Phi(\mathbf{c}'\hat{\boldsymbol{\beta}} - w(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2}) \leq p(x) \leq \Phi(\mathbf{c}'\hat{\boldsymbol{\beta}} + w(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2}).$$

Similarly, $100(1 - \alpha)\%$ upper confidence bands for $p(x)$ are given by

$$p(x) \leq \begin{cases} \{1 + \exp[-\mathbf{c}'\hat{\boldsymbol{\beta}} - w_u(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2}]\}^{-1}, & \text{logistic model} \\ \Phi(\mathbf{c}'\hat{\boldsymbol{\beta}} + w_u(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2}), & \text{probit model} \end{cases}$$

In what follows, we consider the logistic model and focus on Equations (1) and (2). In particular, we propose methods to find the critical values w in Equation (1) for the two-sided case and w_u in Equation (2) for the one-sided case under the logistic model. The derivation of the critical values under the probit model is the same except \mathbf{F}^{-1} is replaced by $\boldsymbol{\Omega}^{-1}$.

Both the logistic and probit models are special cases of the generalized linear model (GLM), and the methodology proposed here can also be applied to other forms of GLM (the complementary-log-log model, for example), that can be transformed into the standard regression model via a link function. In each case, a set of simulations are required to confirm the validity of the method in small samples, thus in the interest of brevity only logistic and probit models are considered here. We devote most of our attention to the logistic model to illustrate the methodology.

Liu *et al.*'s article (2008) presents several methods for the construction of two-sided and one-sided confidence bands for the simple linear model with a restricted predictor variable. Since we can transform the logistic and probit regression setting to the simple linear regression setting via the link functions (logit and probit), we extend their work to the logistic and probit regression models and develop methods to derive asymptotic confidence bands for large samples. This paper is organized as follows. We give the general setting of the problem in Section 1. Two-sided confidence bands are discussed in Section 2, and Section 3 presents the results in the one-sided case. In Section 4, a Monte Carlo simulation is run to investigate how well the asymptotic approximation holds for small sample sizes. An example is given in Section 5 to illustrate the proposed methods.

2 Two-Sided Bands

Our construction of confidence bands are based on the methods given by Liu *et al.* (2008). There are undoubtedly some similarities between the methods presented here and in their review. The main difference between their method and ours lies at the beginning of the derivation. The formulation in their review relied on a standard bivariate t random variable (See Equations (13) and (14) in their review), while the derivation given here involves a standard z bivariate random variable, as will be soon shown. As a result of this reformulation, our critical values are based on a chi-square random variable, instead of an F variate. In the interest of completeness, we include most of the computational details regarding the critical values w and w_u .

The Fisher information matrix \mathbf{F}^{-1} is a positive semi-definite symmetric matrix, so there exists a positive semi-definite matrix \mathbf{B} such that $\mathbf{F}^{-1} = \mathbf{B}^2$. Then

$$\mathbf{z} = \mathbf{B}^{-1}(\boldsymbol{\beta} - \hat{\boldsymbol{\beta}}) \sim N_2(\mathbf{0}, \mathbf{I}).$$

Define the polar coordinates of $\mathbf{z} = (z_1, z_2)'$, (R_z, Q_z) , by

$$z_1 = R_z \cos Q_z, \quad z_2 = R_z \sin Q_z, \quad \text{for } R_z \geq 0 \text{ and } Q_z \in [0, 2\pi].$$

It is well known that R_z^2 has the χ_2^2 distribution and is statistically independent of Q_z , which has a uniform $[0, 2\pi]$ distribution. Write Equation (1) as

$$\begin{aligned} & P[\mathbf{c}'\boldsymbol{\beta} \in \mathbf{c}'\hat{\boldsymbol{\beta}} \pm w(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2}, \text{ for all } x \in (a, b)] \\ &= P \left[\sup_{x \in (a, b)} \frac{|\mathbf{c}'\boldsymbol{\beta} - \mathbf{c}'\hat{\boldsymbol{\beta}}|}{(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2}} < w \right] \\ &= P \left[\sup_{x \in (a, b)} \frac{|(\mathbf{B}\mathbf{c})'\mathbf{z}|}{\|\mathbf{B}\mathbf{c}\|} < w \right] \end{aligned} \tag{3}$$

Notice that for a given vector $\mathbf{u} \in R^2$ and a number $w > 0$, $\{\mathbf{z} : \mathbf{u}'\mathbf{z}/\|\mathbf{u}\| = w\}$ represents a straight line that is perpendicular to the vector \mathbf{u} and is on the same side of the origin as the vector \mathbf{u} . The perpendicular distance from the origin to this line is w . Therefore, the set defined by

$$\{\mathbf{z} : |\mathbf{u}'\mathbf{z}|/\|\mathbf{u}\| < w\} \subset R^2$$

consists of all points that are sandwiched between two parallel straight lines $\mathbf{u}'\mathbf{z}/\|\mathbf{u}\| = w$ and $\mathbf{u}'\mathbf{z}/\|\mathbf{u}\| = -w$. Therefore, letting $\mathbf{B}\mathbf{c} = \mathbf{u}$, Equation (3) can be further expressed as:

$$P \left[\sup_{x \in (a,b)} \frac{|(\mathbf{u})'\mathbf{z}|}{\|\mathbf{u}\|} < w \right] = P\{\mathbf{z} \in R_2\}, \quad (4)$$

where $R_2 = \cap_{x \in (a,b)} R_2(x)$, and $R_2(x) = \{\mathbf{z} : |\mathbf{u}'\mathbf{z}|/\|\mathbf{u}\| < w\}$. R_2 is depicted in Figure 1(a). We can rotate the region R_2 around the origin so that the angle ϕ between the two vectors $\mathbf{u}_a = \mathbf{B}(1-a)$ and $\mathbf{u}_b = \mathbf{B}(1-b)$ is equally divided by the z_1 -axis. The new region, denoted by R_2^* , is depicted in Figure 1(b). Because of the rotation invariance of the normal distribution, we have $P\{\mathbf{z} \in R_2\} = P\{\mathbf{z} \in R_2^*\}$, and R_2^* can be expressed as

$$R_2^* = \{\mathbf{z} : |\mathbf{u}'\mathbf{z}|/\|\mathbf{u}\| < w, \text{ for all } \mathbf{u} \in E(\phi)\},$$

where $E(\phi)$ is a cone depicted as the shaded area in Figure 1(b), and $E(\phi) = \{\mathbf{u} : u_2 > \|\mathbf{u}\|\cos(\phi/2)\}$. Therefore, the simultaneous confidence level is equal to

$$P \left[\sup_{\mathbf{u} \in E(\phi)} \frac{|\mathbf{u}'\mathbf{z}|}{\|\mathbf{u}\|} < w \right] = P\{\mathbf{z} \in R_2^*\}. \quad (5)$$




Figure 1: The Regions R_2 and R_2^*

2.1 Method 1: Finding Expression for the Supremum

This method is based on finding an exact expression for the supremum. Notice that

$$\frac{|\mathbf{u}'\mathbf{z}|}{\|\mathbf{u}\|} = \|\mathbf{z}\| \frac{|\mathbf{u}'\mathbf{z}|}{\|\mathbf{u}\|\|\mathbf{z}\|} = \|\mathbf{z}\|\cos(\theta_{uz}),$$

where θ_{uz} is the angle between \mathbf{u} and \mathbf{z} . Denote $\Psi_1 = [-\phi/2, \phi/2] \cup [\pi - \phi/2, \pi + \phi/2]$, $\Psi_2 = [\phi/2, \pi/2] \cup [\pi + \phi/2, 3\pi/2]$, and $\Psi_3 = [\pi/2, \pi - \phi/2] \cup [3\pi/2, -\phi/2]$. Since $\cos(\theta)$ is monotonically decreasing on $[0, \pi]$, we have the following results:

$$\sup_{\mathbf{u} \in E(\phi)} |\cos(\theta_{uz})| = \begin{cases} 1 & \text{if } \theta_z \in \Psi_1, \\ |\cos(\theta_z - \phi/2)| & \text{if } \theta_z \in \Psi_2, \\ |\cos(\theta_z + \phi/2)| & \text{if } \theta_z \in \Psi_3. \end{cases}$$

Hence, the probability on the left side of Equation (5) can be written as

$$\begin{aligned} & P\{\theta_z \in \Psi_1, \|\mathbf{z}\| < w\} + P\{\theta_z \in \Psi_2, \|\mathbf{z}\|\cos(\theta_z - \phi/2) < w\} \\ & \quad + P\{\theta_z \in \Psi_3, \|\mathbf{z}\|\cos(\theta_z + \phi/2) < w\} \\ & = \frac{\phi}{\pi} P\{\|\mathbf{z}\| < w\} \\ & \quad + 2 \int_{\phi/2}^{\pi/2} \frac{1}{2\pi} P\{\|\mathbf{z}\|\cos(\theta - \phi/2) < w\} d\theta \\ & \quad + 2 \int_{\pi/2}^{\pi - \phi/2} \frac{1}{2\pi} P\{\|\mathbf{z}\|\cos(\theta + \phi/2) < w\} d\theta \\ & = \frac{\phi}{\pi} P\{R_z < w\} \\ & \quad + \frac{2}{\pi} \int_{\phi/2}^{\pi/2} P\{R_z |\cos(\theta - \phi/2)| < w\} d\theta \\ & = \frac{\phi}{\pi} \chi_2^2(w^2) + \frac{2}{\pi} \int_{\phi/2}^{\pi/2} \chi_2^2\left(\frac{w^2}{\cos^2(\theta - \phi/2)}\right) d\theta, \end{aligned} \tag{6}$$

where $\chi_2^2(\cdot)$ is the cdf of a chi-squared distribution with 2 degrees of freedom.

2.2 Method 2: A Method Based on Wynn & Bloomfield's Approach

This method calculates $P\{\mathbf{z} \in R_2^*\}$ directly, and is based on the method given by Wynn & Bloomfield (1971). It can be seen from Figure 1(b) that the region R_2^* is made up of

the circle with radius w and the remaining area. The probability that \mathbf{z} lies in the circle is given by

$$P\{\|\mathbf{z}\| < w\} = P\{R_z < w\} = \chi_2^2(w^2) \quad (7)$$

The probability that \mathbf{z} lies in the remaining region is 4 times the probability that \mathbf{z} lies in the slanted-line shaded area in Figure 1(b), which is

$$\begin{aligned} & 4P\{\theta_z \in [\phi/2, \pi/2], \|\mathbf{z}\| > w, \|\mathbf{z}\|\cos[\theta_z - \phi/2] < w\} \\ &= 4P\{\theta_z \in [\phi/2, \pi/2], w < \|\mathbf{z}\| < w/\cos[\theta_z - \phi/2]\} \\ &= 4 \int_{\phi/2}^{\pi/2} \frac{1}{2\pi} P\{w < \|\mathbf{z}\| < w/\cos[\theta - \phi/2]\} d\theta \\ &= \frac{2}{\pi} \int_{\phi/2}^{\pi/2} \left[\chi_2^2\left(\frac{w^2}{\cos^2[\theta - \phi/2]}\right) - \chi_2^2(w^2) \right] d\theta \end{aligned} \quad (8)$$

Combining Equation (7) and Equation (8), we have the following:

$$P\{\mathbf{z} \in R_2^*\} = \frac{\phi}{\pi} \chi_2^2(w^2) + \frac{2}{\pi} \int_{\phi/2}^{\pi/2} \left[\chi_2^2\left(\frac{w^2}{\cos^2[\theta - \phi/2]}\right) \right] d\theta, \quad (9)$$

which is equal to Equation (6).

3 One-Sided Bands

Similar to the two-sided case, we can express Equation (2) as

$$\begin{aligned} & P[\mathbf{c}'\boldsymbol{\beta} \leq \mathbf{c}'\hat{\boldsymbol{\beta}} + w_u(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2}, \text{ for all } x \in (a, b)] \\ &= P\left[\sup_{x \in (a, b)} \frac{(\mathbf{Bc})'\mathbf{z}}{\|\mathbf{Bc}\|} < w_u \right] \\ &= P\{\mathbf{z} \in R_1\}, \end{aligned} \quad (10)$$

where $R_1 \subset R^2$ is given by $R_1 = \cap_{x \in (a, b)} R_1(x)$, and $R_1(x) = \{\mathbf{z} : \mathbf{u}'\mathbf{z}/\|\mathbf{u}\| < w\}$ with $\mathbf{u} = \mathbf{Bc}$. Rotating the region R_1 around the origin the same way as we did in the two-sided case, we obtain a new region R_1^* with the property $P\{\mathbf{z} \in R_1\} = P\{\mathbf{z} \in R_1^*\}$. The new region is depicted in Figure 2, and has the expression

$$R_1^* = \{\mathbf{z} : \mathbf{u}'\mathbf{z}/\|\mathbf{u}\| < w, \text{ for all } \mathbf{u} \in E(\phi)\},$$

where $E(\phi)$ is defined the same as before. Therefore, the simultaneous confidence level is equal to

$$P \left[\sup_{\mathbf{u} \in E(\phi)} \frac{\mathbf{u}'\mathbf{z}}{\|\mathbf{u}\|} < w_u \right] = P\{\mathbf{z} \in R_1^*\}. \quad (11)$$




Figure 2: Region R_1^*

3.1 Method 1: Finding Expression for the Supremum

Similar to the two-sided bands, this method is based on finding the supremum in Equation (11). As before, we write

$$\frac{\mathbf{u}'\mathbf{z}}{\|\mathbf{u}\|} = \|\mathbf{z}\| \cos(\theta_{uz}),$$

where θ_{uz} is the angle between \mathbf{u} and \mathbf{z} . Denote $\Psi_1 = [-\phi/2, \phi/2]$, $\Psi_2 = [\phi, -\pi/2]$, and $\Psi_3 = [\phi/2, \pi]$. Because of the monotonicity of $\cos(\theta)$ on $[0, \pi]$, we have

$$\sup_{\mathbf{u} \in E(\phi)} \cos(\theta_{uz}) = \begin{cases} 1 & \text{if } \theta_z \in \Psi_1, \\ \cos[\theta_z + \phi/2] & \text{if } \theta_z \in \Psi_2, \\ \cos[\theta_z - \phi/2] & \text{if } \theta_z \in \Psi_3. \end{cases}$$

Hence, the probability on the left side of Equation (11) can be written as

$$\begin{aligned}
& P\{\theta_z \in \Psi_1, \|\mathbf{z}\| < w_u\} + P\{\theta_z \in \Psi_2, \|\mathbf{z}\|\cos[\theta_z + \phi/2] < w_u\} \\
& \quad + P\{\theta_z \in \Psi_3, \|\mathbf{z}\|\cos[\theta_z - \phi/2] < w_u\} \\
& = \frac{\phi}{2\pi} P\{\|\mathbf{z}\| < w_u\} \\
& \quad + 2 \int_{\phi/2}^{\pi} \frac{1}{2\pi} P\{\|\mathbf{z}\|\cos[\theta - \phi/2] < w_u\} d\theta \\
& = \frac{\phi}{2\pi} P\{\|\mathbf{z}\| < w_u\} \\
& \quad + 2 \int_{\phi/2}^{(\pi+\phi)/2} \frac{1}{2\pi} P\{\|\mathbf{z}\|\cos[\theta - \phi/2] < w_u\} d\theta \\
& \quad + 2 \int_{(\pi+\phi)/2}^{\pi} \frac{1}{2\pi} P\{\|\mathbf{z}\|\cos[\theta - \phi/2] < w_u\} d\theta \\
& = \frac{\phi}{2\pi} \chi_2^2(w_u^2) + \frac{\pi - \phi}{2\pi} + \frac{1}{\pi} \int_0^{\pi/2} P\{\|\mathbf{z}\| < \frac{w_u}{\cos(\theta)}\} d\theta. \tag{12}
\end{aligned}$$

The last step follows from the fact that supremum of $\cos(\theta_{uz})$ is negative when $\theta_z \in [\pi, (3\pi - \phi)/2] \cup [(\pi + \phi)/2, \pi]$.

3.2 Method 2: A Method Based on Bohrer & Francis's Approach

This method is similar to that for the two-sided case in Section 2.2, and is based on the method given by Bohrer & Francis (1972). Notice that the region R_1^* can be partitioned into four sub-regions, C_1, C_2, C_3 , and C_4 . The probability $P\{\mathbf{z} \in R_1^*\}$ can be calculated by summing up the probabilities that \mathbf{z} lies in the four sub-regions. The probability that \mathbf{z} falls in C_1 is equal to

$$\begin{aligned}
P\{\mathbf{z} \in C_1\} & = P\{\theta_z \in [(\pi - \phi)/2, (\pi + \phi)/2], \|\mathbf{z}\| < w_u\} \\
& = \frac{\phi}{2\pi} P\{\|\mathbf{z}\| < w_u\} \\
& = \frac{\phi}{2\pi} \chi_2^2(w_u^2). \tag{13}
\end{aligned}$$

The probability that \mathbf{z} falls into the region C_2 is given by

$$P\{\mathbf{z} \in C_2\} = P\{\theta_z \in [(\pi + \phi)/2, (3\pi - \phi)/2]\} = \frac{\pi - \phi}{2\pi}. \tag{14}$$

Now, if we rotate the region C_4 counterclockwise by an angle ϕ , then $C_3 \cup C_4$ forms a strip of width w_u . By further rotating the resulting strip clockwise by an angle $\phi/2$, we obtain a new strip with two sides parallel to the z_2 axis. Because of the rotational invariance of the normal distribution, the probability that \mathbf{z} is in $C_3 \cup C_4$ is equivalent to the probability of \mathbf{z} in the final strip, which is given by

$$P\{\mathbf{z} \in C_2 \cup C_3\} = P\{0 < Z_1 < w_u\} = \frac{1}{2}\chi_1^2(w_u^2), \quad (15)$$

where $\chi_1^2(\cdot)$ is the cdf of a chi-squared distribution with 1 degree of freedom.

Combining Equations (13), (14), and (15) gives us

$$P\{\mathbf{z} \in R_1^*\} = \frac{\phi}{2\pi}\chi_2^2(w_u^2) + \frac{\pi - \phi}{2\pi} + \frac{1}{2}\chi_1^2(w_u^2). \quad (16)$$

It can be shown that Equation (12) and Equation (16) are equal.

4 Monte Carlo Simulation

Since the confidence bands, defined in Equations (1) and (2), are constructed based on the asymptotic properties of the maximum likelihood estimates for the logistic and probit model parameters assuming the sample size is large, we conducted Monte Carlo simulation studies to examine the small-sample performance of the bands. In our simulation, five different values of β were selected: $\beta = [-2, .3]'$, $[0, 1.5]'$, $[2, 5]'$, $[-.2, -.3]'$, $[-2, -4]'$, which represent five different forms of the probability response functions: slowly increasing, moderately increasing, fast increasing, slowly decreasing, and fast decreasing, respectively.

For the logistic model, we examined three different intervals, narrow, wide, and extremely wide ("unrestricted"). The endpoints for each interval were calculated based on the formula $x = (\log_e[p/(1-p)] - \beta_0)/\beta_1$. For the narrow interval, $p = .3, .7$ were selected for computing the endpoints, and $p = .1, .9$ were selected for computing the endpoints of the wide interval. We included the very wide interval in the study to compare the performance of the bands for the restricted and unrestricted cases, and $p = 10^{-10}, 1 - 10^{-10}$ were chosen for computing the endpoints of this interval. The intervals are presented in Table 1.

After the endpoints were calculated for each interval, we generated values for the predictor variable x that were equally spread inside each interval with four different sample sizes ($n = 25, 50, 100, 150$). A uniform $(0, 1)$ random variable was generated, and then the dichotomous response variable Y was generated based on the following: $Y = 1$ if the uniform random variable was less than $p(x) = 1/(1 + \exp(\beta_0 + \beta_1 x))$; $Y = 0$ otherwise.

Table 1: Restricted Intervals For Monte Carlo Simulation

β	Narrow Interval	Wide Interval	“Unrestricted Interval”
$[-2, .3]'$	(3.842, 9.491)	(-.657, 13.991)	(-70.086, 83.420)
$[0, 1.5]'$	(-.565, .565)	(-1.465, 1.465)	(-15.351, 15.351)
$[2, 5]'$	(-.569, -.231)	(-.839, -.039)	(-5.005, 4.205)
$[-.2, -.3]'$	(-3.491, 2.158)	(-7.991, 6.657)	(-77.420, 76.086)
$[-2, -4]'$	(-.712, -.288)	(-1.049, .049)	(-6.256, 5.256)

Three nominal error rates α were considered, .01, .05 and .10. The computer language **R**, version 3.2.1, was used to carry out all Monte Carlo simulations with $N = 5000$ iterations run in each simulation to estimate the Monte Carlo error (1 – the estimated coverage probability) for the two-sided confidence bands. The results are presented in Table 2. The results in the table suggest that the confidence bands are conservative for small samples, and the error approaches the nominal level as the sample size increases. Generally, the error reaches the nominal level when the sample size is 100, but in some cases it could be as small as 50. It is also noted from the table that there is not much difference in error between the three different intervals.

Table 2: Estimated Monte Carlo Errors For the Confidence Bands

β	Sample size n	Narrow			Wide			“Unrestricted ”		
		$\alpha = .01$.05	.10	.01	.05	.10	.01	.05	.10
$[-2, .3]'$	25	.001	.020	.061	.003	.018	.065	.002	.025	.054
	50	.004	.038	.076	.007	.032	.078	.004	.036	.065
	100	.006	.044	.086	.008	.040	.092	.005	.040	.080
	150	.007	.049	.097	.009	.046	.092	.006	.042	.090
$[0, 1.5]'$	25	.002	.022	.064	.002	.026	.058	.006	.029	.059
	50	.005	.036	.082	.006	.035	.072	.010	.035	.066
	100	.008	.042	.086	.009	.043	.094	.008	.042	.076
	150	.006	.044	.097	.008	.051	.092	.010	.045	.087
$[2, 5]'$	25	.002	.021	.069	.004	.027	.058	.005	.024	.057
	50	.004	.036	.080	.007	.044	.074	.010	.037	.060
	100	.008	.040	.092	.007	.042	.084	.009	.037	.076
	150	.010	.048	.101	.008	.049	.089	.010	.046	.085
$[-.3, -.2]'$	25	.001	.020	.065	.005	.024	.057	.008	.025	.070
	50	.005	.039	.073	.008	.034	.074	.005	.037	.067
	100	.008	.043	.092	.007	.036	.091	.009	.042	.077
	150	.007	.046	.102	.007	.041	.090	.010	.040	.084
$[-4, -2]'$	25	.002	.019	.063	.007	.025	.064	.008	.022	.058
	50	.005	.032	.063	.008	.037	.074	.007	.036	.059
	100	.008	.042	.095	.007	.039	.088	.011	.045	.090
	150	.011	.046	.091	.009	.045	.099	.012	.048	.087

In addition, we considered various other designs. One case was that instead of choosing values that were equally spaced inside the chosen intervals for the predictor variable x , we chose values for x that were more concentrated toward one of the endpoints of the interval. The values for predictor variable x were generated as follows: consider the narrow interval $(3.842, 9.491)$, with $n = 25$, we first generated 25 values, stored in a variable z , that were

equally spaced in $(0, 1)$. Then we used the transformation $y = z^6$ to make the values of y concentrate toward the lower end of the interval, 0. Finally, we let $x = 5.649y + 3.842$ to obtain the desired x values which fell inside the chosen interval and were clustered at the lower end, 3.842.

The other case examined was when most of the x values were located at the center of the chosen intervals, which were generated as follows: for the narrow interval $(3.842, 9.491)$ centered at 6.6665, with $n = 25$, we first generated 25 values that were equally spaced in $(-1, 1)$, which were stored in a variable called z . Then we used the transformation $y = z^5$ to make the values of y concentrate toward the center of the interval $(-1, 1)$, which is 0. Finally, we let $x = 5.649/2 * y + 6.6665$ to obtain the desired x values which fell inside the chosen interval and were concentrated at the center of the interval.

Since the estimated error rates presented in Table 2 are similar at three different α values, we chose to perform Monte Carlo simulations at $\alpha = .05$ for these two additional cases. The results are given in Table 3, and it is clear that the results are consistent with what we observed from Table 2. We also ran Monte Carlo simulations for the one-sided bands in the logistic model, and for both bands in the probit model, and again, the results agree with the two-sided case for the logistic model; those results are not presented here to avoid driving the size of the paper to cumbersome levels.

Table 3: Estimated Monte Carlo Errors For the Confidence Bands at $\alpha = .05$

β	Sample size n	Narrow		Wide		“Unrestricted ”	
		Endpoint ^a	Center ^b	Endpoint	Center	Endpoint	Center
[-2, .3]'	25	.015	.020	.021	.023	.021	.021
	50	.022	.031	.031	.038	.036	.040
	100	.042	.037	.040	.046	.041	.043
	150	.049	.045	.045	.044	.042	.045
[0, 1.5]'	25	.025	.018	.021	.023	.016	.022
	50	.032	.031	.039	.038	.025	.030
	100	.042	.037	.040	.046	.040	.048
	150	.045	.045	.046	.044	.041	.047
[2, 5]'	25	.022	.018	.015	.022	.016	.021
	50	.026	.031	.031	.037	.028	.032
	100	.041	.037	.039	.045	.043	.045
	150	.045	.045	.042	.043	.042	.048
[-.3, -.2]'	25	.018	.020	.022	.028	.021	.021
	50	.024	.028	.039	.035	.036	.030
	100	.041	.043	.038	.042	.045	.044
	150	.042	.043	.040	.046	.040	.049
[-4, -2]'	25	.018	.020	.016	.028	.021	.025
	50	.024	.028	.029	.035	.039	.042
	100	.041	.043	.034	.042	.040	.047
	150	.041	.044	.042	.046	.042	.049

^a The x values are concentrated toward the lower end of the interval

^b The x values are concentrated toward the center of the interval

5 Example

If we restrict the independent variable to an interval, the resulting confidence bands will be narrower than the unconstrained bands. Here, we use the data provided by LaVelle (1986) to illustrate the proposed method. The data are presented in Table 4, and these were the same data that were considered by Piegorsch and Casella (1988). The study in LaValle (1986) investigated the comutagenic effects of chromate on frameshift mutagenesis in bacterial assays. The results presented here are findings for the bacterium *E. coli*, strain 343/435. A control and five doses of the suspected mutagen, 9-Aminoacridine (9-AA) are reported in Table 4.

Table 4: Mutagenicity of 9-Aminoacridine in *E. coli* strains 343/435

Dose	-*	.8	2.4	8.0	24	80
Log-dose	-1.374	-.223	0.875	2.079	3.178	4.382
Response	7/96	28/96	64/96	54/96	81/96	96/96

*The first data pair corresponds to a zero-dose control. The log-dose for this datum was calculated using consecutive-dose average spacing (Margolin *et al.*, 1986).

As pointed out by Piegorsch and Casella, rather than report confidence bands over the whole real line, it is often of interest to report narrower bands over constrained intervals. Furthermore, it is often noted that human exposure to environmental toxins usually occurs at low dose levels. Restricting dose levels enables us to direct our attention to intervals toward the lower end, and hence improve the confidence limits greatly.

We fit a logistic model to the data using the log-dose level as our predictor variable x , and the ML estimators from the logistic fit are $\hat{\beta}_0 = -.789$, and $\hat{\beta}_1 = .854$. The inverse of the Fisher information matrix is $\mathbf{F}^{-1} = \begin{bmatrix} 0.017 & -0.005 \\ -0.005 & 0.005 \end{bmatrix}$. If we restrict the predictor variable to an interval $(-1.3, .8)$, then $\mathbf{u}_a = (0.163 \quad -0.109)$, $\mathbf{u}_b = (0.106 \quad 0.0234)$, and the angle between these two vectors are $\phi = 0.809$. The two methods for two-sided bands give the same value for the critical point, $w = 2.206$, at a 95% confidence level.

Therefore, a 95% two-sided confidence band for $p(x)$ when $x \in (-1.3, .8)$ is given by

$$p(x) \in \{1 + \exp[(.789 - .854x) \pm 2.206(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2}]\}^{-1}.$$

To make a comparison with the method of Piegorsch and Casella (1988), the same constrained intervals are considered here, and values of the critical points w are reported in Table 5, along with the critical points given by Piegorsch and Casella. Values of the critical point, based on Scheffé’s method, when there is no restriction on x , are also given here.

Table 5: Critical Values for Two-sided Bands with $\alpha = .05$

Restricted Intervals	Proposed methods	Piegorsch and Casella
$(-\infty, \infty)$	2.447	2.447
$(-1.3, 2.0)$	2.344	2.445
$(-1.3, 0.8)$	2.206	2.274
$(-1.3, -.2)$	2.067	2.170

The sample size in the example is $n = 576$, therefore it is sufficiently large for the asymptotic approximation to hold, based on the Monte Carlo simulation results in Section 4. It can be seen from the table that there is clear improvement in the width of confidence bands when the dose level is restricted to smaller intervals. It is also noted from the table that the proposed method outperforms Piegorsch and Casella’s procedure by giving smaller critical values and hence narrower confidence bands. This is what we expected, because as we pointed out earlier, the embedding procedure used in Piegorsch and Casella’s paper only gave conservative asymptotic confidence bands when the restricted region for predictor variables is rectangular. In this data set, the restricted region on the predictor variable, dose level, is a special case of the rectangular region. The critical values for one-sided confidence bands over the same intervals are also presented here in Table 6.

Table 6: Critical Values for one-sided bands with $\alpha = .05$

Restricted Intervals	Proposed methods
(-1.3, 2.0)	2.049
(-1.3, 0.8)	1.899
(-1.3, -.2)	1.754

Therefore, a 95% upper confidence band for $p(x)$ when $x \in (-1.3, .8)$ is given by

$$p(x) \leq \{1 + \exp[(.789 - .854x) - 1.899(\mathbf{c}'\mathbf{F}^{-1}\mathbf{c})^{1/2}]\}^{-1}.$$

References

- [1] Austin, P. C. and Steyerberg, E. W. (2014). Graphical Assessment of Internal and External Calibration of Logistic Regression Models by Using Loess Smoothers *Statistics in Medicine*, **Volume 33**, Issue 3, 517 - 535.
- [2] Brand, R.J., Pinnock, D.E., and Jackson, K. L. (1973). Large-sample Confidence Bands for the Logistic Response Curve and its Inverse. *American Statistician*, **27**, 157-160.
- [3] Bohrer, R. and Francis, G.K. (1972). Sharp One-sided Confidence Bounds for Linear Regression Over Intervals. *Biometrika*, **59**, 99-107.
- [4] Casella, G. and Strawderman, W.E. (1980). Confidence Bands for Linear Regression with Restricted Predictor Variables. *Journal of the American Statistical Association*, **75**, 862-868.
- [5] Chou, K., Lu, R., and Mao, W. (2002). Factors Relevant to Patient Assaultive Behavior and Assault in Acute Inpatient Psychiatric Units in Taiwan. *Archives of Psychiatric Nursing*, **Volume 16**, Issue 4, 187 - 195.
- [6] Hauck, W. W. (1983). A Note on Confidence Bands for the Logistic Response Curve. *American Statistician*, **37**, 158-160.

- [7] LaVelle, J. M. (1986). Potassium Chromate Potentiates Frameshift Mutagenesis in *E. coli*. and *S. typhimurium*. *Mutation Research*, **171**, 1-10.
- [8] Margolin, B.H., Resnick, M.A., Rimpo, J.Y., Archer, P., Galloway, S.M., Bloom, A.D., and Zeiger, E. (1986). Statistical Analysis for *in Vitro*. Cytogenic Assays Using Chinese Hamster Ovary Cells. *Environmental Mutagenesis*, **8**, 183-204.
- [9] Hochberg, Y. and Quade, D. (1975). One-sided Simultaneous Confidence Bands on Regression Surfaces with Intercepts. *Journal of the American Statistical Association*, **70**, 889-891.
- [10] Kendall, M. and Stuart, A. (1979). *The Advanced Theory of Statistics, Volume II: Inference and Relationship*,. New York: Macmillan.
- [11] Pradhan, B. and Lee, S. (2010). Delineation of Landslide Hazard Areas on Penang Island, Malaysia, By Using Frequency Ratio, Logistic Regression, and Artificial Neural Network Models. *Environmental Earth Science*, **60**, 1037-1054.
- [12] Piegorsch, W.W. and Casella, G. (1988). Confidence Bands for Logistic Regression with Restricted Predictor Variables. *Biometrics*, **44**, 739-750.
- [13] Scheffé, H. (1959). *The Analysis of Variance*. New York: John Wiley.
- [14] Uusipaikka, E. (1983). Exact Confidence Bands for Linear Regression Over Intervals. *Journal of the American Statistical Association*, **78**, 638-644.
- [15] Liu, W., Lin, S., and Piegorsch, W.W. (2008). Construction of Exact Simultaneous Confidence Bands for a Simple Linear Regression Model. *International Statistical Review*, **Volume 76, Issue 1**, 39-57.
- [16] Liu, W., Jamshidian, M., Zhang, Y., and Donnelly, J. (2005). Simulation-based Simultaneous Confidence Bands for a Multiple Linear Regression Model When the Covariates Are Constrained. *Journal of Computational and Graphical Statistics*, **14**, **No. 2**, 459-484.
- [17] Liu, W. (2010). *Simultaneous Inference in Regression*. Chapman & Hall.

- [18] Pan, W. Piegorsch, W.W., and West, R.W. (2003). Exact One-sided Simultaneous Confidence Bands Via Uusipaikka's Method. *Annals of the Institute of Statistical Mathematics.*, **55**, 243-250.
- [19] Working, H. and Hotelling, H. (1929). Application of the Theory of Error to the Interpretation of Trends. *Journal of the American Statistical Association*, Supplement, **24**, 73-85.
- [20] Wynn, H.P. and Bloomfield, P. (1971). Simultaneous Confidence Bands in Regression Analysis. *Journal of the Royal Statistical Society, Series B*, **33**, 202-217.