

Automated Selection of r for the r Largest Order Statistics Approach with Adjustment for Sequential Testing

Brian Bader · Jun Yan · Xuebin Zhang

Abstract The r largest order statistics approach is widely used in extreme value analysis because it may use more information from the data than just the block maxima. In practice, the choice of r is critical. If r is too large, bias can occur; if too small, the variance of the estimator can be high. The limiting distribution of the r largest order statistics, denoted by GEV_r , extends that of the block maxima. Two specification tests are proposed to select r sequentially. The first is a score test for the GEV_r distribution. Due to the special characteristics of the GEV_r distribution, the classical chi-square asymptotics cannot be used. The simplest approach is to use the parametric bootstrap, which is straightforward to implement but computationally expensive. An alternative fast weighted bootstrap or multiplier procedure is developed for computational efficiency. The second test uses the difference in estimated entropy between the GEV_r and GEV_{r-1} models, applied to the r largest order statistics and the $r - 1$ largest order statistics, respectively. The asymptotic distribution of the difference statistic is derived. In a large scale simulation study, both tests held their size and had substantial power to detect various misspecification schemes. A new

approach to address the issue of multiple, sequential hypotheses testing is adapted to this setting to control the false discovery rate or familywise error rate. The utility of the procedures is demonstrated with extreme sea level and precipitation data.

Keywords entropy · generalized extreme value · goodness-of-fit · multiplier bootstrap · score test · sequential testing

1 Introduction

The largest order statistics approach is an extension of the block maxima approach that is often used in extreme value modeling. The focus of this paper is (Smith, 1986, p.28–29): “Suppose we are given, not just the maximum value for each year, but the largest ten (say) values. How might we use this data to obtain better estimates than could be made just with annual maxima?” The r largest order statistics approach may use more information than just the block maxima in extreme value analysis, and is widely used in practice when such data are available for each block. The approach is based on the limiting distribution of the r largest order statistics which extends the generalized extreme value (GEV) distribution (e.g., Weissman, 1978). This distribution, denoted by GEV_r , has the same parameters as the GEV distribution, which makes it useful to estimate the GEV parameters when the r largest values are available for each block. The approach was investigated by Smith (1986) for the limiting joint Gumbel distribution and extended to the more general limiting joint GEV_r distribution by Tawn (1988). Because of the potential gain in efficiency relative to the block maxima only, the method has found many applications such as corrosion engineering (e.g., Scarf and Laycock,

This research was partially supported by an NSF grant (DMS 1521730), a University of Connecticut Research Excellence Program grant, and a contract with Environment and Climate Change Canada.

Brian Bader
University of Connecticut
E-mail: brian.bader@uconn.edu

Jun Yan
University of Connecticut
E-mail: jun.yan@uconn.edu

Xuebin Zhang
Environment and Climate Change Canada
E-mail: xuebin.zhang@canada.ca

1996), hydrology (e.g., Dupuis, 1997), coastal engineering (e.g., Guedes Soares and Scotto, 2004), and wind engineering (e.g., An and Pandey, 2007).

In practice, the choice of r is a critical issue in extreme value analysis with the r largest order statistics approach. In general r needs to be small relative to the block size B (not the number of blocks n) because as r increases, the rate of convergence to the limiting joint distribution decreases sharply (Smith, 1986). There is a trade-off between the validity of the limiting result and the amount of information required for good estimation. If r is too large, bias can occur; if too small, the variance of the estimator can be high. Finding the optimal r should lead to more efficient estimates of the GEV parameters without introducing bias. A much related but different problem is the selection of threshold or fraction of a sample extreme value analysis (see Scarrott and MacDonald, 2012, for a review). Our focus here is the selection of r for situations where a number of largest values are available each of n blocks. In contrast, the methods for threshold or fraction selection reviewed in Scarrott and MacDonald (2012) deal with a single block ($n = 1$) of a large size B .

The selection of r has not been as actively researched as the threshold selection problem in the one sample case. Smith (1986) and Tawn (1988) used probability (also known as PP) plots for the marginal distribution of the r th order statistic to assess its goodness of fit. The probability plot provides a visual diagnosis, but different viewers may reach different conclusions in the absence of a p-value. Further, the probability plot is only checking the marginal distribution for a specific r as opposed to the joint distribution. Tawn (1988) suggested an alternative test of fit using a spacings results in Weissman (1978). Let $D_{n:i}$ be the spacing between the i th and $(i + 1)$ th largest value in a sample of size B from a distribution in the domain of attraction of the Gumbel distribution. Then $\{iD_i : i = 1, \dots, r - 1\}$ is approximately a set of independent and identically distributed exponential random variables as $B \rightarrow \infty$. The connections among the three limiting forms of the GEV distribution (e.g., Embrechts, Klüppelberg, and Mikosch, 1997, p.123) can be used to transform from the Fréchet and the Weibull distribution to the Gumbel distribution. Testing the exponentiality of the spacings on the Gumbel scale provides an approximate diagnosis of the joint distribution of the r largest order statistics when B is large. A limitation of this method, however, is that prior knowledge of the domain of attraction of the distribution is needed. Lastly, Dupuis (1997) proposed a robust estimation method, where the weights can be used to detect inconsistencies with the GEV_r distribution and assess the fit of the data to the joint

Gumbel model. The method can be extended to general GEV_r distributions but the construction of the estimating equations is computing intensive with Monte Carlo integrations.

In this paper, two specification tests are proposed to select r through a sequence of hypothesis testing. The first is the score test (e.g., Rao, 2005), but because of the nonstandard setting of the GEV_r distribution, the usual χ^2 asymptotic distribution is invalid. A parametric bootstrap can be used to assess the significance of the observed statistic, but is computationally demanding. A fast, large sample alternative to parametric bootstrap based on the multiplier approach (Kojadinovic and Yan, 2012) is developed. The second test uses the difference in estimated entropy between the GEV_r and GEV_{r-1} models, applied to the r largest order statistics and the $r - 1$ largest order statistics, respectively. The asymptotic distribution is derived with the central limit theorem. Both tests are intuitive to understand, easy to implement, and have substantial power as shown in the simulation studies. Each of the two tests is carried out to test the adequacy of the GEV_r model for a sequence of r values. The very recently developed stopping rules for ordered hypotheses in G'Sell, Wager, Chouldechova, and Tibshirani (2015) are adapted to control the false discovery rate (FDR), the expected proportion of incorrectly rejected null hypotheses among all rejections, or familywise error rate (FWER), the probability of at least one type I error in the whole family of tests. All the methods are available in an R package `eva` (Bader and Yan, 2015).

The rest of the article is organized as follows. The problem is set up in Section 2 with the GEV_r distribution, observed data, and the hypothesis to be tested. The score test is proposed in Section 3 with two implementations: parametric bootstrap and multiplier bootstrap. The entropy difference (ED) test is proposed and the asymptotic distribution of the testing statistic is derived in Section 4. A large scale simulation study on the empirical size and power of the tests are reported in Section 5. In Section 6, the multiple, sequential testing problem is addressed by adapting recent developments on this application. The tests are applied to sea level and precipitation datasets in Section 7. A discussion concludes in Section 8. The Appendix contains the details of random number generation from the GEV_r distribution and a sketch of the proof of the asymptotic distribution of the ED test statistic.

2 Model and Data Setup

The limit joint distribution of the r largest order statistics of a random sample of size B as $B \rightarrow \infty$ is the

GEV $_r$ distribution with density function (Weissman, 1978)

$$f_r(x_1, x_2, \dots, x_r | \mu, \sigma, \xi) = \sigma^{-r} \exp \left\{ - (1 + \xi z_r)^{-\frac{1}{\xi}} - \left(\frac{1}{\xi} + 1 \right) \sum_{j=1}^r \log(1 + \xi z_j) \right\} \quad (1)$$

for some location parameter μ , scale parameter $\sigma > 0$ and shape parameter ξ , where $x_1 > \dots > x_r$, $z_j = (x_j - \mu)/\sigma$, and $1 + \xi z_j > 0$ for $j = 1, \dots, r$. When $r = 1$, this distribution is exactly the GEV distribution. The parameters $\theta = (\mu, \sigma, \xi)^\top$ remain the same for $j = 1, \dots, r$, $r \ll B$, but the convergence rate to the limit distribution reduces sharply as r increases. The conditional distribution of the r th component given the top $r - 1$ variables in (1) is the GEV distribution right truncated by x_{r-1} , which facilitates simulation from the GEV $_r$ distribution; see Appendix A.

The r largest order statistics approach is an extension of the block maxima approach in extreme value analysis when a number of largest order statistics are available for each one of a collection of independent blocks (Smith, 1986; Tawn, 1988). Specifically, let (x_{i1}, \dots, x_{ir}) be the observed r largest order statistics from block i for $i = 1, \dots, n$. Assuming independence across blocks, the GEV $_r$ distribution is used in place of the GEV distribution in the block maxima approach to make likelihood-based inference about θ . Let $l_i^{(r)}(\theta) = l^{(r)}(x_{i1}, \dots, x_{ir} | \theta)$, where

$$l^{(r)}(x_1, \dots, x_r | \theta) = -r \log \sigma - (1 + \xi z_r)^{-\frac{1}{\xi}} - \left(\frac{1}{\xi} + 1 \right) \sum_{j=1}^r \log(1 + \xi z_j) \quad (2)$$

is the contribution to the log-likelihood from a single block (x_1, \dots, x_r) . The maximum likelihood estimator (MLE) of θ using the r largest order statistics is $\hat{\theta}_n^{(r)} = \arg \max \sum_{i=1}^n l_i^{(r)}(\theta)$.

Model checking is a necessary part of statistical analysis. The rationale of choosing a larger value of r is to use as much information as possible, but not set r too high so that the GEV $_r$ approximation becomes poor due to the decrease in convergence rate. Therefore, it is critical to test the goodness-of-fit of the GEV $_r$ distribution with a sequence of null hypotheses

$H_0^{(r)}$: the GEV $_r$ distribution fits the sample of the r largest order statistics well

for $r = 1, \dots, R$, where R is the maximum, predetermined number of top order statistics to test. Two test

procedures for $H_0^{(r)}$ are developed for a fixed r first to help choose $r \geq 1$ such that the GEV $_r$ model still adequately describes the data. The sequential testing process and the multiple testing issue are investigated in Section 6.

3 Score Test

A score statistic for testing goodness-of-fit hypothesis $H_0^{(r)}$ is constructed in the usual way with the score function and the Fisher information matrix (e.g., Rao, 2005). For ease of notation, the superscript (r) is dropped. Define the score function

$$S(\theta) = \sum_{i=1}^n S_i(\theta) = \sum_{i=1}^n \partial l_i(\theta) / \partial \theta$$

and Fisher information matrix $I(\theta)$, which have been derived in Tawn (1988). The behaviour of the maximum likelihood estimator is the same as that derived for the block maxima approach (Smith, 1985; Tawn, 1988), which requires $\xi > -0.5$. The score statistic is

$$V_n = \frac{1}{n} S^\top(\hat{\theta}_n) I^{-1}(\hat{\theta}_n) S(\hat{\theta}_n).$$

Under standard regularity conditions, V_n would asymptotically follow a χ^2 distribution with 3 degrees of freedom. The GEV $_r$ distribution, however, violates the regularity conditions for the score test (e.g., Casella and Berger, 2002, pp. 516-517), as its support depends on the parameter values unless $\xi = 0$. For illustration, Figure 1 presents a visual comparison of the empirical distribution of V_n with $n = 5000$ from 5000 replicates, overlaid with the $\chi^2(3)$ distribution, for $\xi \in \{-0.25, 0.25\}$ and $r \in \{1, 2, 5\}$. The sampling distribution of V_n appears to be much heavier tailed than $\chi^2(3)$, and the mismatch increases as r increases as a result of the reduced convergence rate.

[Fig. 1 about here.]

Although the regularity conditions do not hold, the score statistic still provides a measure of goodness-of-fit since it is a quadratic form of the score, which has expectation zero under the null hypothesis. Extremely large values of V_n relative to its sampling distribution would suggest lack of fit, and, hence, possible misspecification of $H_0^{(r)}$. So the key to applying the score test is to get an approximation of the sampling distribution of V_n . Two approaches for the approximation are proposed.

3.1 Parametric Bootstrap

The first solution is parametric bootstrap. For hypothesis $H_0^{(r)}$, the test procedure goes as follows:

1. Compute $\hat{\theta}_n$ under H_0 with the observed data.
2. Compute the testing statistic V_n .
3. For every $k \in \{1, \dots, L\}$ with a large number L , repeat:
 - (a) Generate a bootstrap sample of size n for the r largest statistics from GEV_r with parameter vector $\hat{\theta}_n$.
 - (b) Compute the $\hat{\theta}_n^{(k)}$ under H_0 with the bootstrap sample.
 - (c) Compute the score test statistic $V_n^{(k)}$.
4. Return an approximate p-value of V_n as $L^{-1} \sum_{k=1}^L 1(V_n^{(k)} > V_n)$.

Straightforward as it is, the parametric bootstrap approach involves sampling from the null distribution and computing the MLE for each bootstrap sample, which can be very computationally expensive. This is especially true as the sample size n and/or the number of order statistics r included in the model increases.

3.2 Multiplier Bootstrap

Multiplier bootstrap is a fast, large sample alternative to parametric bootstrap in goodness-of-fit testing (e.g., Kojadinovic and Yan, 2012). The idea is to approximate the asymptotic distribution of $n^{-1/2}I^{-1/2}(\theta)S(\theta)$ using its asymptotic representation

$$n^{-1/2}I^{-1/2}(\theta)S(\theta) = \frac{1}{\sqrt{n}} \sum_{i=1}^n \phi_i(\theta),$$

where $\phi_i(\theta) = I^{-1/2}(\theta)S_i(\theta)$. Its asymptotic distribution is the same as the asymptotic distribution of

$$W_n(\mathbf{Z}, \theta) = \frac{1}{\sqrt{n}} \sum_{i=1}^n (Z_i - \bar{Z})\phi_i(\theta),$$

conditioning on the observed data, where $\mathbf{Z} = (Z_1, \dots, Z_n)$ is a set of independent and identically distributed multipliers (independent of the data), with expectation 0 and variance 1, and $\bar{Z} = \frac{1}{n} \sum_{i=1}^n Z_i$. The multipliers must satisfy $\int_0^\infty \{\Pr(|Z_1| > x)\}^{\frac{1}{2}} dx < \infty$. An example of a possible multiplier distribution is $N(0, 1)$.

The multiplier bootstrap test procedure is summarized as follows:

1. Compute $\hat{\theta}_n$ under H_0 with the observed data.
2. Compute the testing statistic V_n .
3. For every $k \in \{1, \dots, L\}$ with a large number L , repeat:

- (a) Generate $\mathbf{Z}^{(k)} = (Z_1^{(k)}, \dots, Z_n^{(k)})$ from $N(0, 1)$.
 - (b) Compute a realization from the approximate distribution of $W_n(\mathbf{Z}, \theta)$ with $W_n(\mathbf{Z}^{(k)}, \hat{\theta}_n)$.
 - (c) Compute $V_n^{(k)}(\hat{\theta}_n) = W_n^\top(\mathbf{Z}^{(k)}, \hat{\theta}_n)W_n(\mathbf{Z}^{(k)}, \hat{\theta}_n)$.
4. Return an approximate p-value of V_n as $L^{-1} \sum_{k=1}^L 1(V_n^{(k)} > V_n)$.

This multiplier bootstrap procedure is much faster than parametric bootstrap procedure because, for each sample, it only needs to generate \mathbf{Z} and compute $W_n(\mathbf{Z}, \hat{\theta}_n)$. The MLE only needs to be obtained once from the observed data.

4 Entropy Difference Test

Another specification test for the GEV_r model is derived based on the difference in entropy for the GEV_r and GEV_{r-1} models. The entropy for a continuous random variable with density f is (e.g., Singh, 2013)

$$E[-\ln f(y)] = - \int_{-\infty}^{\infty} f(y) \log f(y) dy.$$

It is essentially the expectation of negative log-likelihood. The expectation can be approximated with the sample average of the contribution to the log-likelihood from the observed data, or simply the log-likelihood scaled by the sample size n . Assuming that the $r-1$ top order statistics fit the GEV_{r-1} distribution, the difference in the log-likelihood between GEV_{r-1} and GEV_r provides a measure of deviation from $H_0^{(r)}$. Its asymptotic distribution can be derived. Large deviation from the expected difference under $H_0^{(r)}$ suggests a possible misspecification of $H_0^{(r)}$.

From the log-likelihood contribution in (2), the difference in log-likelihood for the i th block, $Y_{ir}(\theta) = l_i^{(r)} - l_i^{(r-1)}$, is

$$Y_{ir}(\theta) = -\log \sigma - (1 + \xi z_{ir})^{-\frac{1}{\xi}} + (1 + \xi z_{ir-1})^{-\frac{1}{\xi}} - \left(\frac{1}{\xi} + 1\right) \log(1 + \xi z_{ir}). \quad (3)$$

Let $\bar{Y}_r = \frac{1}{n} \sum_{i=1}^n Y_{ir}$ and $S_{Y_r}^2 = \sum_{i=1}^n (Y_{ir} - \bar{Y}_r)^2 / (n-1)$ be the sample mean and sample variance, respectively. Consider a standardized version of \bar{Y}_r as

$$T_n^{(r)}(\theta) = \sqrt{n}(\bar{Y}_r - \eta_r) / S_{Y_r}, \quad (4)$$

where $\eta_r = -\log \sigma - 1 + (1 + \xi)\psi(r)$, and $\psi(x) = d \log \Gamma(x) / dx$ is the digamma function. The asymptotic distribution of $T_n^{(r)}$ is summarized by Theorem 1 whose proof is relegated to Appendix B.

Theorem 1 Let $T_n^{(r)}(\theta)$ be the quantity computed based on a random sample of size n from the GEV_r distribution with parameters θ and assume that $H_0^{(r-1)}$ is true. Then $T_n^{(r)}$ converges in distribution to $N(0, 1)$ as $n \rightarrow \infty$.

Note that in Theorem 1, $T_n^{(r)}$ is computed from a random sample of size n from a GEV_r distribution. If the random sample were from a distribution in the domain of attraction of a GEV distribution, the quality of the approximation of the GEV_r distribution to the r largest order statistics depends on the size of each block $B \rightarrow \infty$ with $r \ll B$. The block size B is not to be confused with the sample size n . Assuming $\xi > -0.5$, the proposed ED statistic for $H_0^{(r)}$ is $T_n^{(r)}(\hat{\theta}_n)$, where $\hat{\theta}_n$ is the MLE of θ with the r largest order statistics for the GEV_r distribution. Since $\hat{\theta}_n$ is consistent for θ with $\xi > -0.5$, $T_n^{(r)}(\hat{\theta}_n)$ has the same limiting distribution as $T_n^{(r)}(\theta)$ under $H_0^{(r)}$.

To assess the convergence of $T_n^{(r)}(\hat{\theta}_n)$ to $N(0, 1)$, 1000 GEV_r replicates were simulated under configurations of $r \in \{2, 5, 10\}$, $\xi \in \{-0.25, 0, 0.25\}$, and $n \in \{50, 100\}$. Their quantiles are compared with those of $N(0, 1)$ via quantile-quantile plots (not presented). It appears that a larger sample size is needed for the normal approximation to be good for larger r and negative ξ . This is expected because larger r means higher dimension of the data, and because the MLE only exists for $\xi > -0.5$ (Smith, 1985). For r less than 5 and $\xi \geq 0$, the normal approximation is quite good; it appears satisfactory for sample size as small as 50. For r up to 10, sample size 100 seems to be sufficient.

5 Simulation Results

5.1 Size

The empirical sizes of the tests are investigated first. For the score test, the parametric bootstrap version and the multiplier bootstrap version are equivalent asymptotically, but may behave differently for finite samples. It is of interest to know how large a sample size is needed for the two versions of the score test to hold their levels. Random samples of size n were generated from the GEV_r distribution with $r \in \{1, 2, 3, 4, 5, 10\}$, $\mu = 0$, $\sigma = 1$, and $\xi \in \{-0.25, 0, 0.25\}$. All three parameters (μ, σ, ξ) were estimated.

When the sample size is small, there can be numerical difficulty in obtaining the MLE. For the multiplier bootstrap score and ED test, the MLE only needs to be obtained once, for the dataset being tested. However, in addition, the parametric bootstrap score test must obtain a new sample and obtain the MLE for each

bootstrap replicate. To assess the severity of this issue, 10,000 datasets were simulated for $\xi \in \{-0.25, 0, 0.25\}$, $r \in \{1, 2, 3, 4, 5, 10\}$, $n \in \{25, 50\}$, and the MLE was attempted for each dataset. Failure never occurred for $\xi \geq 0$. With $\xi = -0.25$ and sample size 25, the highest failure rate of 0.69% occurred for $r = 10$. When the sample size is 50, failures only occurred when $r = 10$, at a rate of 0.04%.

For the parametric bootstrap score test with sample size $n \in \{25, 50, 100\}$, Table 1 summarizes the empirical size of the test at nominal levels 1%, 5%, and 10% obtained from 1000 replicates, each carried out with bootstrap sample size $L = 1000$. Included only are the cases that converged successfully. Otherwise, the results show that the agreement between the empirical levels and the nominal level is quite good for samples as small as 25, which may appear in practice when long record data is not available.

[Table 1 about here.]

For the multiplier bootstrap score test, the results for sample sizes $n \in \{25, 50, 100, 200, 300, 400\}$ are summarized in Table 2. When the sample size is less than 100, it appears that there is a large discrepancy between the empirical and nominal level. For $\xi \in \{0, 0.25\}$, there is reasonable agreement between the empirical level and the nominal levels for sample size at least 100. For $\xi = -0.25$ and sample size at least 100, the agreement is good except for $r = 1$, in which case, the empirical level is noticeably larger than the nominal level. This may be due to different rates of convergence for various ξ values as ξ moves away from -0.5 . It is also interesting to note that, everything else being held, the agreement becomes better as r increases. This may be explained by the more information provided by larger r for the same sample size n , as can be seen directly in the fisher information matrix (Tawn, 1988, pp. 247–249). For the most difficult case with $\xi = -0.25$ and $r = 1$, the agreement gets better as sample size increases and becomes acceptable when sample size was 1000 (not reported).

[Table 2 about here.]

To assess the convergence of $T_n^{(r)}(\hat{\theta}_n)$ to $N(0, 1)$, 10,000 replicates of the GEV_r distribution were simulated with $\mu = 0$ and $\sigma = 1$ for each configuration of $r \in \{2, 5, 10\}$, $\xi \in \{-0.25, 0, 0.25\}$, and $n \in \{50, 100\}$. A rejection for nominal level α , is denoted if $|T_n^{(r)}(\hat{\theta}_n)| > |Z_{\frac{\alpha}{2}}|$, where $Z_{\frac{\alpha}{2}}$ is the $\alpha/2$ percentile of the $N(0, 1)$ distribution. Using this result, the empirical size of the ED test can be summarized, and the results are presented in Table 3.

[Table 3 about here.]

For sample size 50, the empirical size is above the nominal level for all configurations of r and ξ . As the sample size increases from 50 to 100, the empirical size stays the same or decreases in every setting. For sample size 100, the agreement between nominal and observed size appears to be satisfactory for all configurations of r and ξ . For sample size 50, the empirical size is slightly higher than the nominal size, but may be acceptable to some practitioners. For example, the empirical size for nominal size 10% is never above 12%, and for nominal size 5%, empirical size is never above 7%.

In summary, the multiplier bootstrap procedure of the score test can be used as a fast, reliable alternative to the parametric bootstrap procedure for sample size 100 or more when $\xi \geq 0$. When only small samples are available (less than 50 observations), the parametric bootstrap procedure is most appropriate since the multiplier version does not hold its size and the ED test relies upon samples of size 50 or more for the central limit theorem to take effect.

5.2 Power

The powers of the score tests and the ED test are studied with two data generating schemes under the alternative hypothesis. In the first scheme, 4 largest order statistics were generated from the GEV_4 distribution with $\mu = 0$, $\sigma = 1$, and $\xi \in \{-0.25, 0, 0.25\}$, and the 5th one was generated from a KumGEV distribution right truncated by the 4th largest order statistic. The KumGEV distribution is a generalization of the GEV distribution (Eljabri, 2013) with two additional parameters a and b which alter skewness and kurtosis. Defining $G_r(\mathbf{x})$ to be the distribution function of the $\text{GEV}_r(\mu, \sigma, \xi)$ distribution, the distribution function of the $\text{KumGEV}_r(\mu, \sigma, \xi, a, b)$ is given by $F_r(\mathbf{x}) = 1 - \{1 - [G_r(\mathbf{x})]^a\}^b$ for $a > 0$, $b > 0$. The score test and the ED test were applied to the top 5 order statistics with sample size $n \in \{100, 200\}$. When $a = b = 1$, the null hypothesis of GEV_5 is true. Larger difference from 1 of parameters a and b means larger deviation from the null hypothesis of GEV_5 .

[Table 4 about here.]

Table 4 summarizes the empirical rejection percentages obtained with nominal size 5%, for a sequence value of $a = b$ from 0.4 to 2.0, with increment 0.2. Both tests hold their sizes when $a = b = 1$ and have substantial power in rejecting the null hypothesis for other values of $a = b$. Between the two tests, the ED test demonstrated much higher power than the score test in the more difficult cases where the deviation from the null

hypothesis is small; for example, the ED test's power almost doubled the score test's power for $a = b \in \{0.8, 1.2\}$. As expected, the powers of both tests increase as $a = b$ moves away from 1 or as the sample sizes increase.

In the second scheme, top 6 order statistics were generated from the GEV_6 distribution with $\mu = 0$, $\sigma = 1$, and $\xi \in \{-0.25, 0, 0.25\}$, and then the 5th order statistic was replaced from a mixture of the 5th and 6th order statistics. The tests were applied to the sample of first 5 order statistics with sample sizes $n \in \{100, 200\}$. The mixing rate p of the 5th order statistic took values in $\{0.00, 0.10, 0.25, 0.50, 0.75, 0.90, 1.00\}$. When $p = 1$ the null hypothesis of GEV_5 is true. Smaller values of p indicate larger deviations from the null. Again, both tests hold their sizes when $p = 1$ and have substantial power for other values of p , which increases as p decreases or as the sample sizes increase. The ED test again outperforms the score test with almost doubled power in the most difficult cases with $p \in \{0.75, 0.90\}$. For sample size 100 with $p = 0.50$, for instance, the ED test has power above 93% while the score test only has power above 69%.

[Table 5 about here.]

6 Automated Sequential Testing Procedure

As there are R hypotheses $H_0^{(r)}$, $r = 1, \dots, R$, to be tested in a sequence in the methods proposed, the sequential, multiple testing issue needs to be addressed. Most methods for error control assume that all the tests can be run first and then a subset of tests are chosen to be rejected (e.g., Benjamini, 2010a,b). The errors to be controlled are either the FWER (Shaffer, 1995), or the FDR (Benjamini and Hochberg, 1995; Benjamini and Yekutieli, 2001). In contrast to the usual multiple testing procedures, however, a unique feature in this setting is that the hypotheses must be rejected in an ordered fashion: if $H_0^{(r)}$ is rejected, $r < R$, then $H_0^{(k)}$ will be rejected for all $r < k \leq R$. Despite the extensive literature on multiple testing and the more recent developments on FDR control and its variants, no definitive procedure has been available for error control in ordered tests until the recent work of G'Sell et al (2015).

Consider a sequence of null hypotheses H_1, \dots, H_m . An ordered test procedure must reject H_1, \dots, H_k for some $k \in \{0, 1, \dots, m\}$, which rules out the classical methods for FDR control (Benjamini and Hochberg, 1995). Let $p_1, \dots, p_m \in [0, 1]$ be the corresponding p-values of the m hypotheses such that p_j is uniformly distributed over $[0, 1]$ when H_j is true. The methods of G'Sell et al (2015) transform the sequence of p-values

to a monotone sequence and then apply the original Benjamini–Hochberg procedure on the monotone sequence. They proposed two rejections rules, each returning a cutoff \hat{k} such that $H_1, \dots, H_{\hat{k}}$ are rejected. The first is called ForwardStop,

$$\hat{k}_F = \max \left\{ k \in \{1, \dots, m\} : -\frac{1}{k} \sum_{i=1}^k \log(1 - p_i) \leq \alpha \right\},$$

and the second is called StrongStop,

$$\hat{k}_S = \max \left\{ k \in \{1, \dots, m\} : \exp \left(\sum_{j=k}^m \frac{\log p_j}{j} \right) \leq \frac{\alpha k}{m} \right\},$$

where α is a pre-specified level. Both rules were shown to control the FDR at level α under the assumption of independent p-values. ForwardStop sets the rejection threshold at the largest k at which the average of first k transformed p-values is small enough. As it does not depend on those p-values with later indices, this rule is robust to potential misspecification at later indices. StrongStop offers a stronger guarantee than ForwardStop. If the non-null p-values indeed precede the null p-values, it controls the FWER at level α in addition to the FDR. Thus, for ForwardStop, this α refers to the FDR and for StrongStop, α refers to the FWER. As the decision to stop at k depends on all the p-values after k , its power may be harmed if, for example, the very last p-values are slightly higher than expected under the null hypotheses.

To apply the two rules to our setting, note that our objective is to give a threshold \hat{r} such that the first \hat{r} of $m = R$ hypotheses are accepted instead of rejected. Therefore, we put the p-values in reverse order: let the ordered set of p-values $\{p_1, \dots, p_R\}$ correspond to hypotheses $\{H_0^{(R)}, \dots, H_0^{(1)}\}$. The two rules give a cutoff $\hat{k} \in \{1, \dots, R\}$ such that the hypotheses $H_0^{(R)}, \dots, H_0^{(R-\hat{k}+1)}$ are rejected. If no $\hat{k} \in \{1, \dots, R\}$ exists, then no rejection is made.

A caveat is that, unlike the setting of G'Sell et al (2015), the p-values of the sequential tests are dependent. Nonetheless, the ForwardStop and StrongStop procedures may still provide some error control. For example, in the non-sequential multiple testing scenario Benjamini and Yekutieli (2001) show that their procedure controls the FDR under certain positive dependency conditions, while Blanchard and Roquain (2009) implement adaptive versions of step-up procedures that provably control the FDR under unspecified dependence among p-values.

The empirical properties of the two rules for the tests in this paper are investigated in simulation studies. To check the empirical FWER of the StrongStop

rule, only data under the null hypotheses are needed. With $R = 10$, $\xi \in \{-0.25, 0.25\}$, $n \in \{30, 50, 100, 200\}$, $\mu = 0$, and $\sigma = 1$, 1000 GEV_{10} samples were generated. For the ED, multiplier bootstrap score, and parametric bootstrap score test, the observed FWER is compared to the expected rates at various nominal α control levels. The StrongStop procedure is used, as well as no error control (i.e. a rejection occurs any time the raw p-value is below the nominal level). The results of this simulation are presented in Figure 2.

It is clear that the StrongStop reasonably controls the FWER for the ED test and the agreement between the observed and expected rate increases as the sample size increases. For both the parametric and multiplier bootstrap versions of the score test however, the observed FWER is above the expected rate, at times 10% higher. Regardless, it is apparent that using no error control results in an inflated FWER, and this inflation can only increase as the number of tests increase.

[Fig. 2 about here.]

To check the empirical FDR of the ForwardStop rule, data need to be generated from a non-null model. To achieve this, consider the sequence of specification tests of GEV_r distribution with $r \in \{1, \dots, 6\}$, where the 5th and 6th order statistics are misspecified. Specifically, data from the GEV_7 distribution with $\mu = 0$ and $\sigma = 1$ were generated for n blocks; then the 5th order statistic is replaced with a 50/50 mixture of the 5th and 6th order statistics, and the 6th order statistic is replaced with a 50/50 mixture of the 6th and 7th order statistics. This is replicated 1000 times for each value of $\xi \in \{-0.25, 0.25\}$ and $n \in \{30, 50, 100, 200\}$. For nominal level α , the observed FDR is defined as the number of false rejections (i.e. any rejection of $r \leq 4$) divided by the number of total rejections.

The results are presented in Figure 3. The plots show that the ForwardStop procedure controls the FDR for the ED test, while for both versions of the score test, the observed FDR is slightly higher than the expected at most nominal rates. Here, sample size does not appear to effect the observed rates.

[Fig. 3 about here.]

Similarly, the observed FWER rate in this particular simulation setting can be found by taking the number of simulations with at least one false rejection (here, any rejection of $r \leq 4$) and dividing that number by the total number of simulations. This calculation is performed for a variety of nominal levels α , using the StrongStop procedure. The results are presented in Figure 3. In this particular simulation setting, the StrongStop

procedure controls the FWER for the ED test and both versions of the score test at all sample sizes investigated.

It is of interest to investigate the performance of the ForwardStop and StrongStop in selecting r for the r largest order statistics method. In the simulation setting described in the last paragraph, the correct choice of r should be 4, and a good testing procedure should provide a selection \hat{r} close to 4. The choice $\hat{r} \in \{0, \dots, 6\}$ is recorded using the ED test and bootstrap score tests with both ForwardStop and StrongStop. Due to space constraints, we choose to present one setting, where $\xi = 0.25$ and $n = 100$. The non-adjusted sequential procedure is also included, testing in an ascending manner from $r = 1$ and \hat{r} is chosen by the first rejection found (if any). The results are summarized in Table 6.

[Table 6 about here.]

In general, larger choices of α lead to a higher percentage of $\hat{r} = 4$ being correctly chosen with ForwardStop or StrongStop. Intuitively, this is not surprising since a smaller α makes it more difficult to reject the ‘bad’ hypotheses of $r \in \{5, 6\}$. A larger choice of α also leads to a higher probability of rejecting too many tests; i.e. choosing r too small. From the perspective of model specification, this is more desirable than accepting true negatives. A choice of 6, 5, or 0 is problematic, but choosing 1, 2, or 3 is acceptable, although some information is lost. When no adjustment is used and an ascending sequential procedure is used, both tests have reasonable classification rates. When $\alpha = 0.05$, the ED test achieves the correct choice of r 79.9% of the time, with the parametric bootstrap and multiplier bootstrap score tests achieving 68.1% and 59.6% respectively. Of course, as the number of tests (i.e., R) increase, with no adjustment the correct classification rates will go down and the ForwardStop/StrongStop procedures will achieve better rates. This may not be too big an issue here as R is typically small. In the case where rich data are available and R is big, the ForwardStop and StrongStop becomes more useful as they are designed to handle a large number of ordered hypothesis.

7 Illustrations

7.1 Lowestoft Sea Levels

Sea level readings in 60 and 15 minute intervals from a gauge at Lowestoft off the east coast of Britain during the years 1964–2014 are available from the UK Tide Gauge Network website. The readings are hourly from 1964–1992 and in fifteen minute intervals from 1993 to present. Accurate estimates of extreme sea levels are of

great interest. The current data are of better quality and with longer record than those used in Tawn (1988) — annual maxima during 1953–1983 and hourly data during 1970–78 and 1980–82.

Justification of the statistical model was considered in detail by Tawn (1988). The three main assumptions needed to justify use of the GEV_r model are: (1) The block size B is large compared to the choice of r ; (2) Observations within each block and across blocks are approximately independent; and (3) The distribution of the block maxima follows GEV_1 . The first assumption is satisfied, by letting $R = 125$, and noting that the block size for each year is $B = 365 \times 24 = 8760$ from 1964–1992 and $B = 365 \times 96 = 35040$ from 1993–2014. This ensures that $r \ll B$. The third assumption is implicitly addressed in the testing procedure; if the goodness-of-fit test for the block maxima rejects, all subsequent tests for $r > 1$ are rejected as well.

The second assumption can be addressed in this setting by the concept of independent storms (Tawn, 1988). The idea is to consider each storm as a separate event, with each storm having some storm length, say τ . Thus, when selecting the r largest values from each block, only a single contribution can be obtained from each storm, which can be considered the r largest independent annual events. By choosing τ large enough, this ensures both approximate independence of observations within each block and across blocks. The procedure to extract the independent r largest annual events is as follows:

1. Pick out the largest remaining value from the year (block) of interest.
2. Remove observations within a lag of $\tau/2$ from both sides of the value chosen in step 1.
3. Repeat (within each year) until the r largest are extracted.

A full analysis is performed on the Lowestoft sea level data using $\tau = 60$ as the estimated storm length (Tawn, 1988). Using $R = 125$, both the parametric bootstrap score (with bootstrap sample size $L = 10,000$) and ED test are applied sequentially on the data. The p-values of the sequential tests (adjusted and unadjusted) can be seen in Figure 4. Due to the large number of tests, the adjustment for multiplicity is desired and thus, ForwardStop is used to choose r . For this dataset, the score test is more powerful than the ED test. With ForwardStop and the score test, Figure 4 suggests that $r = 33$. The remainder of this analysis proceeds with the choice of $r = 33$. The estimated parameters and corresponding 95% profile confidence intervals for $r = 1$ through $r = 40$ are shown in Figure 5.

[Fig. 4 about here.]

[Fig. 5 about here.]

When $r = 33$, the parameters are estimated as $\hat{\mu} = 3.462$ (0.023), $\hat{\sigma} = 0.210$ (0.013), and $\hat{\xi} = -0.017$ (0.023), with standard errors in parenthesis. An important risk measure is the t -year return level z_t (e.g., Hosking, 1990; Ribereau, Guillo, and Naveau, 2008; Singo, Kundu, Odiyo, Mathivha, and Nkuna, 2012). It can be thought of here as the sea level that is exceeded once every t years on average. Specifically, the t -year return level is the $1 - 1/t$ quantile of the GEV distribution

$$z_t = \begin{cases} \mu - \frac{\sigma}{\xi} \{1 - [-\log(1 - \frac{1}{t})]^{-\xi}\}, & \xi \neq 0, \\ \mu - \sigma \log[-\log(1 - \frac{1}{t})], & \xi = 0. \end{cases}$$

The return levels can be estimated with parameter values replaced with their estimates, and confidence intervals can be constructed using profile likelihood (e.g., Coles, 2001, p.57).

The 95% profile likelihood confidence intervals for the 50, 100, and 200 year return levels (i.e. z_{50}, z_{100}, z_{200}) are given by (4.102, 4.461), (4.210, 4.641) and (4.312, 4.824), respectively. The benefit of using $r = 1$ versus $r = 33$ can be seen in the return level confidence intervals in Figure 5. For example, the point estimate of the 100 year return level decreases slightly as r increases and the width of the 95% confidence interval decreases drastically from 2.061 ($r = 1$) to 0.432 ($r = 33$), as more information is used. The lower bound of the interval however remains quite stable, shifting from 4.330 to 4.210 — less than a 3% change. Similarly, the standard error of the shape parameter estimate decreases by over two-thirds when using $r = 33$ versus $r = 1$.

7.2 Annual Maximum Precipitation: Atlantic City, NJ

The top 10 annual precipitation events (in centimeters) were taken from the daily records of a rain gauge station in Atlantic City, NJ from 1874–2015. The year 1989 is missing, while the remaining records are greater than 98% complete. This provides a total record length of 141 years. The raw data is a part of the Global Historical Climatology Network (GHCN-Daily), with an overview given by Menne, Durre, Vose, Gleason, and Houston (2012). The specific station identification in the dataset is USW00013724.

[Fig. 6 about here.]

Unlike for the Lowestoft sea level data, a rather small value is set for R at $R = 10$ because of the much lower frequency of the daily data. Borrowing ideas from Section 7.1, a storm length of $\tau = 2$ is used to ensure approximate independence of observations. Both the parametric bootstrap score (with $L = 10,000$) and ED test

are applied sequentially on the data. The p-values of the sequential tests (ForwardStop, StrongStop, and unadjusted) are shown in Figure 6. The score test does not pick up anything. The ED test obtains p-values 0.002 and 0.016, respectively, for $r = 9$ and $r = 10$, which translates into a rejection using ForwardStop. Thus, Figure 6 suggests that $r = 8$ be used for the analysis.

[Fig. 7 about here.]

With $r = 8$, the estimated parameters are given as $\hat{\mu} = 6.118$ (0.139), $\hat{\sigma} = 2.031$ (0.118), and $\hat{\xi} = 0.219$ (0.032). This suggests a heavy upper tail for the estimated distribution (i.e. $\hat{\xi} > 0$). The progression of parameters and certain return level estimates can be seen in Figure 7. The 50, 100, and 200 year return level 95% confidence intervals for $r = 8$ are calculated using the profile likelihood method and are given by (16.019, 22.411), (18.606, 26.979), and (21.489, 31.136), respectively. The advantages of using $r = 8$ versus the block maxima for analysis are quite clear from Figure 7. The standard error of the shape parameter decreases from 0.071 to 0.032, a decrease of over 50%. Similarly, the 50 year return level 95% confidence intervals decreases in width by over 25%.

8 Discussion

We proposed two model specification tests for a fixed number of largest order statistics as the basis for selecting r for the r largest order statistics approach in extreme value analysis. The score test has two versions of bootstrap procedure: the multiplier bootstrap method providing a fast, large sample alternative to the parametric bootstrap method, with a speedup of over 100 times. The ED test depends on asymptotic normal approximation of the testing statistic, which becomes acceptable for sample size over 50. It assumes that the $r - 1$ top order statistics included already fits the GEV_{r-1} distribution. Therefore, the initial hypothesis at $r = 1$ needs to be tested with the score tests. Both tests hold their size better when the shape parameter is further away from the lower limit of -0.5 or sample size is larger. When only small samples are available (50 observations or less), the parametric bootstrap score test is recommended.

Alternative versions of the ED test have been explored. One may define the testing statistics as the difference in entropy between GEV_1 and GEV_r , instead of between GEV_{r-1} and GEV_r . Nonetheless, it appeared to require a larger sample to hold its size from our simulation studies (not reported). In the calculation of $T_n^{(r)}$, the block maxima MLE $\hat{\theta}_n^{(1)}$ can be used as an estimate

for θ in place of $\hat{\theta}_n^{(r)}$. Again, in our simulation studies, this version of the ED test was too conservative, thus reducing the power, when the sample size was not large enough. This may be explained in that the resulting \hat{S}_{Y_r} underestimates S_{Y_r} .

Naively, the tests may be performed sequentially for each $r \in \{1, \dots, R\}$, for a prefixed, usually small R , at a certain significance level until $H_0^{(r)}$ is rejected. The issue of multiple, sequential testing is addressed in detail by adapting two very recent stopping rules to control the FDR and the FWER that are developed specifically for situations when hypotheses must be rejected in an ordered fashion (G'Sell et al, 2015). It is shown that these automated procedures reasonably control the error rate for the tests discussed in this paper. The naive unadjusted sequential testing procedure also appears to have decent performance at choosing the correct r and for this scenario is more conservative (selects a smaller r) than the stopping rules.

The tests can be extended to allow covariates in the parameters. For example, extremal precipitation in a year may be affected by large scale climate indexes such as the Southern Oscillation Index (SOI), which may be incorporated as a covariate in the location parameter (e.g., Shang, Yan, and Zhang, 2011). Both tests can be carried out with additional model parameters. When the underlying data falls into a rich class of dependence structures (such as time series), this dependence may be incorporated directly instead of using a procedure to achieve approximate independence (e.g. the storm length τ in Section 7). For example, take the GEV-GARCH model (Zhao, Scarrott, Oxley, and Reale, 2011) when $r = 1$. It may be extended to the case where $r > 1$ and the tests presented here may be applied to select r under this model assumption.

A Generating from GEV_r Distribution

The GEV_r distribution is closely connected to the GEV distribution. Let $X_1 > \dots > X_r$ follow a GEV_r distribution (1). It is obvious that the GEV_1 distribution is the GEV distribution with the same parameters, which is the marginal distribution of X_1 . More interestingly, note that, the conditional distribution of X_2 given $X_1 = x_1$ is simply the GEV distribution righted truncated by x_1 . In general, given $(X_1, \dots, X_k) = (x_1, \dots, x_k)$ for $1 \leq k < r$, the conditional distribution of X_{k+1} is the GEV distribution righted truncated at x_k . This property can be exploited to generate the r components in a realized GEV_r observation.

The pseudo algorithm to generate a single observation is the following:

- Generate the first value x_1 from the (unconditional) GEV distribution.
- For $i = 2, \dots, r$:
 - Generate x_i from the GEV distribution right truncated by x_{i-1} .

The resulting vector (x_1, \dots, x_r) is a single observation from the GEV_r distribution.

For $\xi \rightarrow 0$, caveat is needed in numerical evaluation. Using function `expm1` for $\exp(1+x)$ for $x \rightarrow 0$ provides much improved accuracy in comparison to a few implementations in existing R packages. For readability, here is a simplified version of our implemetation in R package `eva` (Bader and Yan, 2015).

```
## Quantile function of a GEVr(loc, scale, shape)
qgev <- function(p, loc = 0, scale = 1, shape = 0,
  lower.tail = TRUE, log.p = FALSE) {
  if (log.p) p <- exp(p)
  if(shape == 0) {
    loc - scale * log(-log(p))
  } else
    loc + scale * expm1(log(-log(p)) * -shape) / shape
}

## Random number generator of GEVr;
## Returns a matrix of n rows and r columns,
## each row a draw from GEVr
rgevr <- function(n, r, loc = 0, scale = 1, shape = 0) {
  umat <- matrix(runif(n * r), n, r)
  if (r > 1) {
    matrix(qgev(t(apply(umat, 1, cumprod)),
      loc, scale, shape),
      ncol = r)
  } else {
    qgev(umat, loc, scale, shape)
  }
}
```

B Asymptotic Distribution of $T_n^{(r)}(\theta)$

Proof (Theorem 1) Consider a random vector (X_1, X_2, \dots, X_r) which follows a $\text{GEV}_r(\theta)$ distribution. The following result given by Tawn (1988, pg. 248) will be used:

$$\begin{aligned} h(j|\theta, a, b, c) &\equiv E[Z_j^a (1 + \xi Z_j)^{-(\frac{1}{\xi} + b)} \log^c(1 + \xi Z_j)] \\ &= \frac{(-\xi)^{c-a}}{\Gamma(j)} \sum_{\alpha=0}^a (-1)^\alpha \binom{a}{\alpha} \Gamma^{(c)}(j + b\xi - \alpha\xi + 1) \end{aligned} \quad (5)$$

where $Z_j = (X_j - \mu)/\sigma$ and $\Gamma^{(c)}$ is the c th derivative of the gamma function, for $a \in \mathbb{Z}$, $b \in \mathbb{R}$, and $c \in \mathbb{Z}$, such that $(j + b\xi - \alpha\xi + 1) \notin \{0, -1, -2, \dots\}$, $\alpha = 0, 1, \dots, a$.

Assume that $\xi \neq 0$ and $1 + \xi Z_j > 0$ for $j = 1, \dots, r$. The difference in log-likelihoods for a single observation from the $\text{GEV}_r(\theta)$ and $\text{GEV}_{r-1}(\theta)$ distribution is given by (3) in Section 4. Thus, the first moment of Y_{ir} is

$$\begin{aligned} E[Y_{1r}] &= -\log \sigma - h(r|\theta, 0, 0, 0) + h(r-1|\theta, 0, 0, 0) \\ &\quad - \left(\frac{1}{\xi} + 1 \right) h(r|\theta, 0, -\xi^{-1}, 1) \\ &= -\log \sigma - 1 + (1 + \xi)\psi(r) \end{aligned}$$

where $\psi(x) = \frac{\Gamma^{(1)}(x)}{\Gamma(x)}$.

To prove that the second moment of Y_{ir} is finite, note that

$$|Y_{1r}| \leq 4 \max \left\{ \left| \log \sigma \right|, \left| (1 + \xi Z_{1r})^{-\frac{1}{\xi}} \right|, \right. \\ \left. \left| (1 + \xi Z_{1r-1})^{-\frac{1}{\xi}} \right|, \left| \left(\frac{1}{\xi} + 1 \right) \log(1 + \xi Z_{1r-1}) \right| \right\},$$

which implies

$$Y_{1r}^2 \leq 16 \left(\max \left\{ \left| \log \sigma \right|, \left| (1 + \xi Z_{1r})^{-\frac{1}{\xi}} \right|, \right. \right. \\ \left. \left. \left| (1 + \xi Z_{1r-1})^{-\frac{1}{\xi}} \right|, \left| \left(\frac{1}{\xi} + 1 \right) \log(1 + \xi Z_{1r-1}) \right| \right\} \right)^2.$$

The bound of $E(Y_{1r}^2)$ can be established by applying (5) to the last three terms in the max operator,

$$E[(1 + \xi Z_{1r})^{-\frac{2}{\xi}}] = h(r|\theta, 0, \xi^{-1}, 0) < \infty,$$

$$E[(1 + \xi Z_{1r-1})^{-\frac{2}{\xi}}] = h(r-1|\theta, 0, \xi^{-1}, 0) < \infty,$$

$$E[\log^2(1 + \xi Z_{1r-1})] = h(r-1|\theta, 0, -\xi^{-1}, 2) < \infty.$$

The desired result then follows from the central limit theorem and Slutsky's theorem.

The case where $\xi = 0$ in Theorem 1 can easily be derived by taking the limit as $\xi \rightarrow 0$ in (3) and in (5) by the Dominated Convergence Theorem.

Acknowledgements We would like to thank Dr. Zhiyi Chi for his comments and discussion on the multiple testing issues.

References

- An Y, Pandey MD (2007) The r largest order statistics model for extreme wind speed estimation. *Journal of Wind Engineering and Industrial Aerodynamics* 95(3):165–182
- Bader B, Yan J (2015) eva: Extreme Value Analysis with Goodness-of-Fit Testing
- Benjamini Y (2010a) Discovering the false discovery rate. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 72(4):405–416
- Benjamini Y (2010b) Simultaneous and selective inference: Current successes and future challenges. *Biometrical Journal* 52(6):708–721
- Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B* 57(1):289–300
- Benjamini Y, Yekutieli D (2001) The control of the false discovery rate in multiple testing under dependency. *The Annals of Statistics* 29(4):1165–1188
- Blanchard G, Roquain É (2009) Adaptive false discovery rate control under independence and dependence. *The Journal of Machine Learning Research* 10:2837–2871
- Casella G, Berger RL (2002) *Statistical Inference*, 2nd edn. Duxbury Pacific Grove, CA
- Coles S (2001) *An Introduction to Statistical Modeling of Extreme Values*, 1st edn. Springer
- Dupuis DJ (1997) Extreme value theory based on the r largest annual events: A robust approach. *Journal of Hydrology* 200(1):295–306
- Eljabri SSM (2013) *New statistical models for extreme values*. PhD thesis, The University of Manchester, Manchester, UK
- Embrechts P, Klüppelberg C, Mikosch T (1997) *Modelling Extremal Events*, vol 33. Springer Science & Business Media
- G'Sell MG, Wager S, Chouldechova A, Tibshirani R (2015) Sequential selection procedures and false discovery rate control. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* DOI 10.1111/rssb.12122, URL <http://dx.doi.org/10.1111/rssb.12122>, forthcoming
- Guedes Soares C, Scotto MG (2004) Application of the r largest-order statistics for long-term predictions of significant wave height. *Coastal Engineering* 51(5):387–394
- Hosking JRM (1990) L-moments: Analysis and estimation of distributions using linear combinations of order statistics. *Journal of the Royal Statistical Society Series B (Methodological)* 52(1):105–124
- Kojadinovic I, Yan J (2012) Goodness-of-fit testing based on a weighted bootstrap: A fast large-sample alternative to the parametric bootstrap. *Canadian Journal of Statistics* 40(3):480–500
- Menne MJ, Durre I, Vose RS, Gleason BE, Houston TG (2012) An overview of the global historical climatology network-daily database. *Journal of Atmospheric and Oceanic Technology* 29(7):897–910
- Rao CR (2005) Score test: Historical review and recent developments. In: Balakrishnan N, Kannan N, Nagaraja HN (eds) *Advances in Ranking and Selection, Multiple Comparisons, and Reliability*, Springer, pp 3–20
- Ribereau P, Guillou A, Naveau P (2008) Estimating return levels from maxima of non-stationary random sequences using the generalized PWM method. *Nonlinear Processes in Geophysics* 15(6):1033–1039
- Scarf PA, Laycock PJ (1996) Estimation of extremes in corrosion engineering. *Journal of applied statistics* 23(6):621–644
- Scarrott C, MacDonald A (2012) A review of extreme value threshold estimation and uncertainty quantification. *REVSTAT-Statistical Journal* 10(1):33–60
- Shaffer JP (1995) Multiple hypothesis testing. *Annual Review of Psychology* 46(1):561–584
- Shang H, Yan J, Zhang X (2011) El Niño–Southern Oscillation influence on winter maximum daily precipitation in California in a spatial model. *Water Resources Research* 47:W11,507–W11,515, DOI 10.1029/2011WR010415
- Singh VP (2013) *Entropy Theory and Its Application in Environmental and Water Engineering*. John Wiley & Sons
- Singo LR, Kundu PM, Odiyo JO, Mathivha FI, Nkuna TR (2012) Flood frequency analysis of annual maximum stream flows for Luvuvhu river catchment, Limpopo Province, South Africa. Tech. rep., University of Venda, Department of Hydrology and Water Resources
- Smith RL (1985) Maximum likelihood estimation in a class of nonregular cases. *Biometrika* 72(1):67–90
- Smith RL (1986) Extreme value theory based on the r largest annual events. *Journal of Hydrology* 86(1):27–43
- Tawn JA (1988) An extreme-value theory model for dependent observations. *Journal of Hydrology* 101(1):227–250
- Weissman I (1978) Estimation of parameters and large quantiles based on the k largest observations. *Journal of the American Statistical Association* 73(364):812–815
- Zhao X, Scarrott CJ, Oxley L, Reale M (2011) GARCH dependence in extreme value models with Bayesian inference. *Mathematics and Computers in Simulation* 81(7):1430–1440

Table 1: Empirical size (in %) for the parametric bootstrap score test under the null distribution GEV_r , with $\mu = 0$ and $\sigma = 1$ based on 1000 samples, each with bootstrap sample size $L = 1000$.

| Sample Size | r | 25 | | | 50 | | | 100 | | |
|---------------|-----|-----|-----|------|-----|-----|------|-----|-----|------|
| Nominal Size | | 1.0 | 5.0 | 10.0 | 1.0 | 5.0 | 10.0 | 1.0 | 5.0 | 10.0 |
| $\xi = -0.25$ | 1 | 0.4 | 2.8 | 6.0 | 1.1 | 4.8 | 9.3 | 0.6 | 4.1 | 8.0 |
| | 2 | 0.1 | 2.6 | 6.0 | 0.8 | 3.4 | 6.5 | 0.6 | 3.6 | 8.1 |
| | 3 | 0.3 | 2.5 | 5.0 | 0.8 | 4.3 | 7.7 | 1.1 | 4.8 | 8.1 |
| | 4 | 0.3 | 1.8 | 5.4 | 0.6 | 3.1 | 6.9 | 1.1 | 5.1 | 8.8 |
| | 5 | 0.4 | 2.4 | 6.7 | 0.4 | 3.3 | 8.3 | 0.6 | 3.1 | 6.5 |
| | 10 | 2.7 | 5.3 | 8.7 | 0.5 | 3.9 | 8.4 | 0.7 | 4.2 | 7.6 |
| $\xi = 0$ | 1 | 1.3 | 5.2 | 8.9 | 1.6 | 5.3 | 9.0 | 0.8 | 4.7 | 9.3 |
| | 2 | 1.4 | 5.1 | 9.4 | 2.0 | 4.9 | 10.0 | 1.0 | 4.3 | 9.9 |
| | 3 | 1.7 | 6.2 | 10.9 | 2.1 | 6.0 | 10.2 | 0.8 | 4.9 | 9.8 |
| | 4 | 1.5 | 4.5 | 8.5 | 1.3 | 6.0 | 10.2 | 1.0 | 4.4 | 9.8 |
| | 5 | 1.6 | 5.8 | 10.4 | 2.4 | 6.2 | 9.9 | 1.2 | 5.0 | 9.7 |
| | 10 | 1.5 | 4.0 | 7.3 | 1.5 | 4.3 | 8.9 | 0.7 | 4.6 | 8.2 |
| $\xi = 0.25$ | 1 | 1.7 | 4.5 | 9.7 | 2.6 | 7.1 | 11.5 | 1.1 | 4.6 | 9.1 |
| | 2 | 1.8 | 5.1 | 8.7 | 1.8 | 4.4 | 8.5 | 0.5 | 2.9 | 7.5 |
| | 3 | 1.5 | 4.4 | 9.4 | 1.5 | 3.7 | 8.1 | 1.0 | 4.2 | 9.4 |
| | 4 | 1.2 | 3.3 | 8.1 | 1.1 | 4.6 | 9.7 | 1.1 | 4.3 | 9.6 |
| | 5 | 1.7 | 4.4 | 9.4 | 1.1 | 4.2 | 8.6 | 0.6 | 4.8 | 9.6 |
| | 10 | 1.1 | 4.6 | 8.3 | 1.5 | 6.1 | 10.7 | 1.0 | 3.9 | 8.5 |

Table 2: Empirical size (in %) for multiplier bootstrap score test under the null distribution GEV_r , with $\mu = 0$ and $\sigma = 1$. 1000 samples, each with bootstrap sample size $L = 1000$ were used. Although not shown, the empirical size for $r = 1$ and $\xi = -0.25$ becomes acceptable when sample size is 1000.

| Sample Size | r | 25 | | | 50 | | | 100 | | |
|---------------|-----|-----|------|------|-----|------|------|-----|------|------|
| Nominal Size | | 1.0 | 5.0 | 10.0 | 1.0 | 5.0 | 10.0 | 1.0 | 5.0 | 10.0 |
| $\xi = -0.25$ | 1 | 7.0 | 13.4 | 18.9 | 6.3 | 13.8 | 19.6 | 5.4 | 11.4 | 16.3 |
| | 2 | 2.0 | 6.9 | 13.4 | 1.3 | 6.4 | 12.4 | 1.6 | 6.9 | 13.6 |
| | 3 | 2.1 | 5.8 | 11.7 | 1.1 | 5.9 | 11.1 | 1.1 | 5.0 | 10.8 |
| | 4 | 3.3 | 7.2 | 12.3 | 1.1 | 4.9 | 10.8 | 1.0 | 5.2 | 11.9 |
| | 5 | 3.6 | 9.0 | 14.0 | 2.3 | 6.8 | 11.2 | 1.1 | 6.2 | 10.6 |
| | 10 | 2.0 | 7.0 | 10.3 | 2.6 | 7.4 | 12.8 | 2.1 | 6.4 | 10.1 |
| $\xi = 0$ | 1 | 3.3 | 8.4 | 15.3 | 2.2 | 7.0 | 12.5 | 1.1 | 4.6 | 9.2 |
| | 2 | 2.8 | 8.7 | 14.4 | 1.8 | 7.5 | 13.0 | 0.9 | 5.7 | 10.3 |
| | 3 | 6.1 | 12.1 | 16.5 | 3.0 | 7.2 | 12.2 | 1.5 | 6.0 | 10.4 |
| | 4 | 5.1 | 10.4 | 14.5 | 3.6 | 10.1 | 14.9 | 1.0 | 5.6 | 10.3 |
| | 5 | 4.2 | 9.0 | 14.5 | 2.2 | 8.2 | 12.5 | 1.7 | 6.5 | 12.0 |
| | 10 | 3.1 | 9.2 | 14.4 | 2.4 | 6.4 | 9.8 | 0.6 | 4.6 | 9.0 |
| $\xi = 0.25$ | 1 | 1.8 | 6.7 | 13.7 | 1.3 | 4.7 | 10.4 | 0.8 | 4.4 | 11.5 |
| | 2 | 5.7 | 12.7 | 17.1 | 4.7 | 9.9 | 14.9 | 3.5 | 7.4 | 11.6 |
| | 3 | 7.1 | 12.2 | 16.5 | 5.3 | 9.4 | 14.8 | 4.2 | 8.4 | 12.5 |
| | 4 | 5.4 | 9.8 | 16.8 | 3.7 | 9.0 | 13.4 | 2.6 | 6.0 | 11.4 |
| | 5 | 4.4 | 10.1 | 15.8 | 3.5 | 8.2 | 13.6 | 2.4 | 7.4 | 11.4 |
| | 10 | 3.3 | 8.9 | 15.3 | 2.4 | 6.6 | 12.3 | 1.6 | 5.8 | 10.9 |

| Sample Size | r | 200 | | | 300 | | | 400 | | |
|---------------|-----|-----|------|------|-----|-----|------|-----|-----|------|
| Nominal Size | | 1.0 | 5.0 | 10.0 | 1.0 | 5.0 | 10.0 | 1.0 | 5.0 | 10.0 |
| $\xi = -0.25$ | 1 | 5.4 | 10.5 | 15.2 | 3.6 | 8.2 | 12.6 | 2.8 | 7.1 | 12.5 |
| | 2 | 1.4 | 6.7 | 12.8 | 1.4 | 6.4 | 11.4 | 1.4 | 5.1 | 10.9 |
| | 3 | 1.5 | 5.9 | 11.8 | 1.1 | 5.4 | 10.8 | 1.2 | 6.6 | 11.9 |
| | 4 | 1.1 | 5.6 | 10.6 | 1.0 | 5.6 | 11.5 | 1.0 | 4.7 | 9.0 |
| | 5 | 1.1 | 4.5 | 9.3 | 1.2 | 5.7 | 11.7 | 1.2 | 4.7 | 10.2 |
| | 10 | 1.4 | 6.4 | 11.6 | 1.7 | 6.2 | 11.3 | 0.8 | 5.0 | 10.1 |
| $\xi = 0$ | 1 | 1.3 | 6.1 | 11.2 | 0.8 | 5.2 | 10.0 | 1.0 | 5.1 | 11.4 |
| | 2 | 0.5 | 5.0 | 10.6 | 1.2 | 5.7 | 11.8 | 1.0 | 5.9 | 11.0 |
| | 3 | 1.4 | 4.5 | 9.8 | 1.3 | 6.0 | 9.6 | 0.9 | 4.4 | 8.3 |
| | 4 | 1.1 | 5.4 | 10.6 | 1.3 | 5.2 | 9.9 | 0.9 | 5.0 | 9.1 |
| | 5 | 1.8 | 6.2 | 12.5 | 0.9 | 4.6 | 9.8 | 1.2 | 4.6 | 9.0 |
| | 10 | 1.1 | 3.8 | 9.3 | 0.9 | 5.2 | 12.6 | 1.2 | 4.9 | 9.7 |
| $\xi = 0.25$ | 1 | 0.9 | 4.9 | 11.4 | 0.9 | 5.0 | 10.8 | 0.7 | 5.2 | 9.2 |
| | 2 | 3.2 | 7.9 | 11.7 | 2.3 | 7.1 | 11.2 | 2.5 | 6.6 | 12.1 |
| | 3 | 1.8 | 6.1 | 10.7 | 2.6 | 7.0 | 11.2 | 1.0 | 4.8 | 10.6 |
| | 4 | 1.2 | 4.9 | 11.2 | 1.2 | 6.0 | 9.9 | 1.2 | 5.8 | 11.8 |
| | 5 | 1.6 | 5.9 | 10.0 | 1.3 | 7.3 | 11.8 | 1.2 | 3.9 | 8.4 |
| | 10 | 1.7 | 6.6 | 12.4 | 0.9 | 4.4 | 9.8 | 1.6 | 5.7 | 10.4 |

Table 3: Empirical size (in %) for the entropy difference (ED) test under the null distribution GEV_r , with $\mu = 0$ and $\sigma = 1$ based on 10,000 samples.

| Sample Size | r | 50 | | | 100 | | |
|---------------|-----|-----|-----|------|-----|-----|------|
| Nominal Size | | 1.0 | 5.0 | 10.0 | 1.0 | 5.0 | 10.0 |
| $\xi = -0.25$ | 2 | 1.5 | 5.7 | 10.8 | 1.3 | 5.5 | 10.1 |
| | 5 | 2.4 | 6.8 | 11.9 | 1.6 | 5.9 | 10.6 |
| | 10 | 2.3 | 6.8 | 11.7 | 1.9 | 6.0 | 11.1 |
| $\xi = 0$ | 2 | 1.3 | 5.6 | 11.0 | 1.2 | 5.3 | 10.4 |
| | 5 | 1.6 | 5.9 | 11.2 | 1.5 | 5.7 | 10.6 |
| | 10 | 2.3 | 6.5 | 11.8 | 1.6 | 5.9 | 10.7 |
| $\xi = 0.25$ | 2 | 1.3 | 5.7 | 10.7 | 1.3 | 5.4 | 10.5 |
| | 5 | 1.6 | 5.8 | 11.5 | 1.3 | 5.6 | 10.2 |
| | 10 | 2.0 | 6.6 | 11.9 | 1.4 | 5.5 | 10.4 |

Table 4: Empirical rejection rate (in %) of the multiplier score test and the ED test in the first data generating scheme from 1000 replicates.

| Sample Size | ξ | Test | Value of $a=b$ | | | | | | | | |
|-------------|-------|-------|----------------|-------|------|-----|------|------|-------|-------|-------|
| | | | 0.4 | 0.6 | 0.8 | 1.0 | 1.2 | 1.4 | 1.6 | 1.8 | 2.0 |
| 100 | -0.25 | Score | 99.9 | 84.8 | 20.4 | 5.4 | 21.0 | 41.2 | 62.3 | 79.0 | 83.0 |
| | | ED | 100.0 | 99.0 | 46.5 | 4.6 | 48.7 | 89.5 | 99.2 | 100.0 | 99.8 |
| | 0 | Score | 100.0 | 87.0 | 21.6 | 7.4 | 24.2 | 48.9 | 67.8 | 79.6 | 89.4 |
| | | ED | 100.0 | 98.8 | 40.0 | 5.2 | 40.6 | 87.2 | 98.5 | 100.0 | 99.7 |
| | 0.25 | Score | 100.0 | 87.7 | 20.3 | 6.2 | 25.8 | 54.2 | 74.2 | 82.9 | 89.5 |
| | | ED | 100.0 | 97.5 | 37.7 | 4.8 | 34.8 | 78.1 | 96.1 | 99.5 | 99.7 |
| 200 | -0.25 | Score | 100.0 | 98.6 | 40.7 | 5.2 | 29.8 | 64.7 | 86.4 | 95.9 | 97.5 |
| | | ED | 100.0 | 100.0 | 78.4 | 6.2 | 70.0 | 99.2 | 100.0 | 100.0 | 100.0 |
| | 0 | Score | 100.0 | 99.4 | 44.6 | 6.1 | 34.9 | 75.0 | 92.4 | 97.3 | 98.6 |
| | | ED | 100.0 | 99.9 | 75.0 | 5.5 | 64.6 | 98.1 | 99.8 | 100.0 | 100.0 |
| | 0.25 | Score | 100.0 | 99.3 | 44.5 | 6.3 | 37.0 | 73.4 | 91.8 | 97.0 | 98.9 |
| | | ED | 100.0 | 100.0 | 71.0 | 5.2 | 57.2 | 95.9 | 100.0 | 100.0 | 100.0 |

Table 5: Empirical rejection rate (in %) of the multiplier score test and the ED tests in the second data generating scheme from 1000 replicates.

| Sample Size | ξ | Test | Mixing Rate p | | | | | | |
|-------------|-------|-------|-----------------|-------|-------|-------|------|------|------|
| | | | 0.00 | 0.10 | 0.25 | 0.50 | 0.75 | 0.90 | 1.00 |
| 100 | -0.25 | Score | 99.7 | 99.5 | 95.5 | 69.4 | 24.1 | 7.8 | 5.8 |
| | | ED | 100.0 | 100.0 | 100.0 | 97.7 | 51.8 | 10.9 | 6.2 |
| | 0 | Score | 100.0 | 99.7 | 97.8 | 72.4 | 22.7 | 6.2 | 6.8 |
| | | ED | 100.0 | 100.0 | 100.0 | 96.0 | 47.6 | 10.3 | 5.6 |
| | 0.25 | Score | 99.9 | 99.7 | 96.6 | 70.8 | 24.7 | 5.8 | 5.3 |
| | | ED | 100.0 | 100.0 | 99.9 | 93.6 | 43.4 | 9.8 | 5.2 |
| 200 | -0.25 | Score | 99.9 | 100.0 | 99.7 | 95.6 | 43.4 | 11.4 | 5.1 |
| | | ED | 100.0 | 100.0 | 100.0 | 100.0 | 83.6 | 20.0 | 5.8 |
| | 0 | Score | 100.0 | 100.0 | 100.0 | 96.5 | 44.4 | 11.2 | 5.4 |
| | | ED | 100.0 | 100.0 | 100.0 | 100.0 | 79.5 | 20.0 | 5.5 |
| | 0.25 | Score | 100.0 | 100.0 | 100.0 | 97.2 | 46.9 | 9.2 | 5.5 |
| | | ED | 100.0 | 100.0 | 100.0 | 99.7 | 72.5 | 17.9 | 4.2 |

Table 6: Percentage of choice of r using the ForwardStop and StrongStop rules at various significance levels or FDRs, under ED, parametric bootstrap (PB) score, and multiplier bootstrap (MB) score tests, with $n = 100$ and $\xi = 0.25$ for the simulation setting described in Section 6. Correct choice is $r = 4$.

| Test | r | Unadjusted | | | | | | ForwardStop | | | | | | StrongStop | | | | | |
|----------|---|------------|------|------|------|------|------|-------------|------|------|------|------|------|------------|------|------|------|------|------|
| | | 0.01 | 0.05 | 0.1 | 0.2 | 0.3 | 0.4 | 0.01 | 0.05 | 0.1 | 0.2 | 0.3 | 0.4 | 0.01 | 0.05 | 0.1 | 0.2 | 0.3 | 0.4 |
| ED | 6 | 19.0 | 3.4 | 1.5 | 0.5 | 0.0 | 0.0 | 86.6 | 69.8 | 58.5 | 43.0 | 30.0 | 22.4 | 52.4 | 22.2 | 13.1 | 5.4 | 1.7 | 1.0 |
| | 5 | 1.9 | 2.1 | 1.1 | 0.7 | 0.2 | 0.1 | 1.3 | 1.5 | 1.1 | 0.9 | 0.2 | 0.0 | 25.0 | 18.9 | 13.6 | 6.7 | 2.9 | 1.4 |
| | 4 | 76.2 | 79.9 | 70.2 | 50.5 | 35.2 | 22.3 | 12.0 | 25.1 | 31.7 | 33.4 | 31.6 | 26.3 | 22.6 | 58.9 | 72.9 | 84.9 | 89.0 | 85.7 |
| | 3 | 0.8 | 4.1 | 7.6 | 11.8 | 15.3 | 15.1 | 0.1 | 3.4 | 6.2 | 12.4 | 17.0 | 18.5 | 0.0 | 0.0 | 0.3 | 2.2 | 4.5 | 6.9 |
| | 2 | 1.1 | 5.3 | 9.5 | 16.2 | 19.9 | 22.8 | 0.0 | 0.1 | 1.9 | 5.4 | 9.4 | 11.5 | 0.0 | 0.0 | 0.1 | 0.7 | 1.8 | 4.4 |
| | 1 | 1.0 | 5.2 | 10.1 | 20.3 | 29.4 | 39.7 | 0.0 | 0.1 | 0.6 | 4.9 | 11.8 | 21.3 | 0.0 | 0.0 | 0.0 | 0.1 | 0.1 | 0.6 |
| PB Score | 0 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - |
| | 6 | 35.8 | 16.1 | 8.5 | 1.8 | 0.6 | 0.2 | 53.5 | 33.0 | 23.4 | 13.9 | 8.6 | 5.9 | 40.6 | 25.1 | 18.8 | 12.1 | 7.6 | 5.6 |
| | 5 | 2.5 | 1.8 | 0.8 | 0.5 | 0.2 | 0.1 | 1.9 | 1.3 | 0.7 | 0.8 | 0.4 | 0.4 | 29.8 | 37.7 | 29.2 | 17.5 | 10.5 | 6.6 |
| | 4 | 58.4 | 68.1 | 63.9 | 46.1 | 31.3 | 19.8 | 42.6 | 57.8 | 58.9 | 51.1 | 42.0 | 31.1 | 29.3 | 36.5 | 50.5 | 65.7 | 73.5 | 74.6 |
| | 3 | 0.5 | 2.3 | 4.5 | 6.8 | 7.1 | 8.0 | 1.4 | 4.2 | 9.0 | 15.5 | 16.0 | 17.1 | 0.0 | 0.4 | 1.1 | 3.3 | 4.8 | 5.6 |
| | 2 | 0.6 | 3.0 | 4.9 | 10.3 | 11.7 | 12.7 | 0.5 | 1.7 | 3.0 | 7.3 | 10.2 | 12.3 | 0.1 | 0.1 | 0.2 | 1.0 | 2.4 | 4.5 |
| MB Score | 1 | 0.8 | 3.7 | 6.9 | 13.9 | 18.3 | 18.5 | 0.1 | 1.3 | 2.3 | 4.6 | 8.3 | 9.6 | 0.2 | 0.2 | 0.2 | 0.4 | 1.2 | 2.7 |
| | 0 | 1.4 | 5.0 | 10.5 | 20.6 | 30.8 | 40.7 | 0.0 | 0.7 | 2.7 | 6.8 | 14.5 | 23.6 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.4 |
| | 6 | 49.9 | 16.9 | 6.9 | 1.3 | 0.2 | 0.0 | 71.7 | 40.3 | 24.7 | 12.6 | 7.8 | 5.5 | 51.6 | 27.3 | 16.9 | 10.4 | 6.2 | 4.3 |
| | 5 | 2.5 | 2.3 | 0.7 | 0.3 | 0.1 | 0.0 | 1.3 | 2.0 | 0.8 | 0.5 | 0.3 | 0.4 | 39.4 | 50.7 | 42.9 | 25.5 | 15.8 | 9.4 |
| | 4 | 38.3 | 59.6 | 59.1 | 44.0 | 31.2 | 18.4 | 26.6 | 53.3 | 59.3 | 49.9 | 40.1 | 28.0 | 6.2 | 18.5 | 35.0 | 55.4 | 62.3 | 64.8 |
| | 3 | 1.6 | 2.8 | 4.0 | 6.6 | 7.5 | 6.0 | 0.3 | 2.8 | 7.4 | 15.7 | 16.0 | 17.9 | 0.6 | 1.2 | 2.5 | 3.6 | 7.0 | 7.8 |
| | 2 | 2.7 | 4.4 | 7.1 | 11.0 | 10.0 | 10.6 | 0.1 | 0.6 | 3.4 | 8.6 | 9.7 | 9.5 | 0.7 | 0.8 | 1.2 | 2.8 | 5.3 | 7.0 |
| | 1 | 4.2 | 8.3 | 10.6 | 14.0 | 19.5 | 20.0 | 0.0 | 0.9 | 2.7 | 4.7 | 7.9 | 7.8 | 1.5 | 1.5 | 1.5 | 2.3 | 3.4 | 6.4 |
| | 0 | 0.8 | 5.7 | 11.6 | 22.8 | 31.5 | 45.0 | 0.0 | 0.1 | 1.7 | 8.0 | 18.2 | 30.9 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.3 |




Fig. 1: Comparisons of the empirical distribution based on 5000 replicates of the score test statistic and the $\chi^2(3)$ distribution (red solid curve). The number of blocks used is $n = 5000$ with parameters $\mu = 0$, $\sigma = 1$, and $\xi \in (-0.25, 0.25)$.




Fig. 2: Observed FWER for the ED, parametric bootstrap (PB) score, and multiplier bootstrap (MB) score tests (using No Adjustment and StrongStop) versus expected FWER at various nominal levels. The 45 degree line indicates agreement between the observed and expected rates under H_0 .




Fig. 3: Observed FDR (from ForwardStop) and observed FWER (from StrongStop) versus expected FDR and FWER, respectively, at various nominal levels. This is for the simulation setting described in Section 6, using the ED, parametric bootstrap (PB) score, and multiplier bootstrap (MB) score tests. The 45 degree line indicates agreement between the observed and expected rates.




Fig. 4: P-Values using ForwardStop, StrongStop, and no adjustment for the ED and PB Score tests applied to the Lowestoft sea level data. The horizontal dashed line represents the 0.05 possible cutoff value.




Fig. 5: Location, scale, and shape parameter estimates, with 95% profile confidence intervals for $r = 1, \dots, 40$ for the Lowestoft sea level data. Also included are the estimates and 95% profile likelihood confidence intervals for the 50 and 100 year return levels.




Fig. 6: P-Values using ForwardStop, StrongStop, and no adjustment for the ED and PB Score tests applied to the Atlantic City precipitation data. The horizontal dashed line represents the 0.05 possible cutoff value.




Fig. 7: Location, scale, and shape parameter estimates, with 95% delta confidence intervals for $r = 1$ through $r = 10$ for the Atlantic City precipitation data. Also included are the estimates and 95% profile likelihood confidence intervals for the 50, 100, and 200 year return levels.