

# Box-Cox symmetric distributions and applications to nutritional data

Silvia L.P. Ferrari\*

*Department of Statistics, University of São Paulo, São Paulo, Brazil*

Giovana Fumes

*Department of Exact Sciences, ESALQ, University of São Paulo, Piracicaba, Brazil*

## Abstract

We introduce and study the Box-Cox symmetric class of distributions, which is useful for modeling positively skewed, possibly heavy-tailed, data. The new class of distributions includes the Box-Cox t, Box-Cox Cole-Green (or Box-Cox normal), Box-Cox power exponential distributions, and the class of the log-symmetric distributions as special cases. It provides easy parameter interpretation, which makes it convenient for regression modeling purposes. Additionally, it provides enough flexibility to handle outliers. The usefulness of the Box-Cox symmetric models is illustrated in a series of applications to nutritional data.

*Key words:* Box-Cox transformation; Symmetric distributions; Box-Cox power exponential distribution; Box-Cox slash distribution; Box-Cox t distribution; Log-symmetric distributions; Nutrients intake.

## 1 Introduction

It is well-known that positive continuous data usually present positive skewness and outlier observations. This is the typical situation with survival times, nutrients intake and family income data, among many other examples. Since Box and Cox (1964) seminal paper, the Box-Cox power transformation has been routinely employed for transforming to normality. Let  $Y$  be a positive random variable. The Box-Cox transformation is defined as  $(Y^\lambda - 1)/\lambda$ , if  $\lambda \neq 0$ , and  $\log Y$ , if  $\lambda = 0$ . Despite its popularity and ease of implementation, this approach, however, has drawbacks, one of them being the fact that the model parameters cannot be

---

\*Corresponding author. Email: silviaferrari@usp.br

easily interpreted in terms of the original response. A conceptual shortcoming is that the support of the transformed variable is not the whole real line and hence a (non-truncated) normal distribution should not be assumed for the transformed data.

An alternative approach to the Box-Cox transformation that allows the parameters to be interpretable as characteristics of the original data is the Box-Cox Cole-Green distribution (or Box-Cox normal distribution); see Cole and Green (1992). It uses the Box-Cox approach, but the parameters are incorporated into the transformation. The Box-Cox Cole-Green distribution has support in  $\mathbb{R}^+$  and is defined from the transformation

$$Z \equiv h(Y; \mu, \sigma, \lambda) = \begin{cases} \frac{1}{\sigma\lambda} \left[ \left( \frac{Y}{\mu} \right)^\lambda - 1 \right], & \text{if } \lambda \neq 0, \\ \frac{1}{\sigma} \log \left( \frac{Y}{\mu} \right), & \text{if } \lambda = 0, \end{cases} \quad (1)$$

where  $\mu > 0$ ,  $\sigma > 0$ ,  $-\infty < \lambda < \infty$ , assuming that  $Z$  has a standard normal distribution truncated at a suitable interval of the real line; details are given in the next section. The Box-Cox symmetric (BCS) class of distributions defined in this paper replaces the normal distribution by the class of the continuous standard symmetric distributions. Replacing the normal distribution by the Student-t and the power exponential distributions results in the Box-Cox t (Rigby and Stasinopoulos; 2006) and the Box-Cox power exponential (Rigby and Stasinopoulos; 2004; Voudouris et al.; 2012) distributions. Additionally, it generalizes the class of the log symmetric distributions (Vanegas and Paula; 2016, 2015).

The paper unfolds as follows. In Section 2, the Box-Cox symmetric class of distributions is defined, some properties are stated and interpretation of the parameters in terms of quantiles is discussed. Tail heaviness of Box-Cox symmetric distributions is studied in Section 3. It is shown that the Box-Cox symmetric class of distributions allows much more tail flexibility than the log-symmetric distributions. Likelihood-based inference is discussed in Section 4. It is suggested that the choice of the symmetric distribution may lead to robust estimation against outliers. In Section 5, applications to 33 nutrients intake data are presented, and a comparison of alternative approaches is provided. Finally, concluding remarks (Section 6) close the paper. Technical details are left for the Appendix.

## 2 Box-Cox symmetric distributions

A continuous random variable  $W$  is said to have a symmetric distribution with location parameter  $\mu \in \mathbb{R}$ , scale parameter  $\sigma > 0$  and density generating function  $r$ , and we write  $W \sim S(\mu, \sigma; r)$ , if its probability density function (pdf) is given by

$$f_W(w) = \frac{1}{\sigma} r \left( \left( \frac{w - \mu}{\sigma} \right)^2 \right), \quad w \in \mathbb{R}, \quad (2)$$

where  $r(\cdot)$  satisfies  $r(u) > 0$ , for  $u \geq 0$ , and  $\int_0^\infty u^{-1/2} r(u) du = 1$ . The class of the symmetric distributions has a number of well-known distributions as special cases depending on the

choice of  $r$ . It includes the normal distribution as well as the Student-t, power exponential, type I logistic, type II logistic and slash distributions among others. Densities in this family have quite different tail behaviors, and some of them may have heavier or lighter tails than the normal distribution.

The symmetric distributions have some interesting properties. Some of them follow: (i) If  $W \sim S(\mu, \sigma; r)$ , its characteristic function is  $\psi_W(t) = e^{it\mu}\varphi(t^2\sigma^2)$ ,  $t \in \mathbb{R}$ , for some function  $\varphi$ , with  $\varphi(u) \in \mathbb{R}$ , for  $u > 0$ . Whenever they exist,  $E(W) = \mu$  and  $\text{Var}(W) = \xi\sigma^2$ , where  $\xi = -2\varphi'(0) > 0$ , with  $\varphi'(0) = d\varphi(u)/du|_{u=0}$  is a constant not depending on  $\mu$  and  $\sigma$  (Fang et al.; 1990). If  $u^{-(k+1)/2}r(u)$  is integrable, then the  $k$ th moment of  $W$  exist (Kelker; 1970). (ii) If  $W \sim S(\mu, \sigma; r)$ , then  $a + bW \sim S(a + b\mu, |b|\sigma; r)$ , where  $a, b \in \mathbb{R}$ , with  $b \neq 0$ . In particular, if  $W \sim S(\mu, \sigma; r)$ , then  $S = (W - \mu)/\sigma \sim S(0, 1; r)$ , and its pdf is  $f_S(s) = r(s^2)$ , for  $s \in \mathbb{R}$ .

Let  $Y$  be a positive continuous random variable, and consider  $Z \equiv h(Y; \mu, \sigma, \lambda)$  as in (1). Assume that  $Z$  has a standard symmetric distribution truncated at  $\mathbb{R} \setminus A(\sigma, \lambda)$ , where

$$A(\sigma, \lambda) = \begin{cases} \left(-\frac{1}{\sigma\lambda}, \infty\right), & \text{if } \lambda > 0, \\ \left(-\infty, -\frac{1}{\sigma\lambda}\right), & \text{if } \lambda < 0, \\ (-\infty, \infty), & \text{if } \lambda = 0, \end{cases} \quad (3)$$

i.e. the support of the truncated distribution is  $A(\sigma, \lambda)$ . We then say that  $Y$  has a Box-Cox symmetric distribution with parameters  $\mu > 0$ ,  $\sigma > 0$ , and  $\lambda \in \mathbb{R}$ , and density generating function  $r$ , and we write  $Y \sim BCS(\mu, \sigma, \lambda; r)$ . In other words,  $Y \sim BCS(\mu, \sigma, \lambda; r)$  if the transformed variable  $Z$  in (1) has the distribution of  $S \sim S(0, 1; r)$  truncated at  $\mathbb{R} \setminus A(\sigma, \lambda)$ .

The Box-Cox symmetric class of distributions reduces to the log-symmetric class of distributions (Vanegas and Paula; 2016) when  $\lambda$  is fixed at zero. Additionally, it leads to the Box-Cox Cole-Green (Stasinopoulos et al.; 2008), Box-Cox t (Rigby and Stasinopoulos; 2006) and Box-Cox power exponential (Rigby and Stasinopoulos; 2004; Voudouris et al.; 2012) distributions by taking  $Z$  as a truncated standard normal, Student-t and power exponential random variable, respectively. The density generating function,  $r(u)$ , for  $u \geq 0$ , for various distributions in the BCS class follows:

- (i) normal:  $r(u) = (2\pi)^{-1/2} \exp\{-u/2\}$ ;
- (ii) double exponential:  $r(u) = (\sqrt{2}/2) \exp\{-\sqrt{2}u^{1/2}\}$ ;
- (iii) power exponential:  $r(u) = [\tau/(p(\tau)2^{1+1/\tau}\Gamma(1/\tau))] \exp\{-u^{\tau/2}/(2p(\tau)^\tau)\}$ , where  $\tau > 0$  and  $p(\tau)^2 = 2^{-2/\tau}\Gamma(1/\tau)[\Gamma(3/\tau)]^{-1}$ ; when  $\tau = 1$  and  $\tau = 2$ ,  $r(u)$  coincides with the density generating function of the double exponential and normal, respectively;
- (iv) Cauchy:  $r(u) = \{\pi(1 + u)\}^{-1}$ ;
- (v) Student-t:  $r(u) = \tau^{\tau/2}\{B(1/2, \tau/2)\}^{-1}(\tau + u)^{-(\tau+1)/2}$ ,  $\tau > 0$ , where  $B(\cdot, \cdot)$  is the beta function;

- (vi) type I logistic:  $r(u) = c \exp\{-u\}(1 + \exp\{-u\})^{-2}$ , where  $c \approx 1.484300029$  is the normalizing constant, obtained from the relation  $\int_0^\infty u^{-1/2}r(u)du = 1$ ;
- (vii) type II logistic:  $r(u) = \exp\{-u^{1/2}\}(1 + \exp\{-u^{1/2}\})^{-2}$ ;
- (viii) canonical slash (Rogers and Tukey; 1972):  $r(u) = (1/\sqrt{2\pi}u)(1 - \exp\{-u/2\})$ , for  $u > 0$ , and  $r(u) = 1/(2\sqrt{2\pi})$ , for  $u = 0$ ;
- (ix) slash<sup>1</sup>:  $r(u) = \Psi((q+1)/2, u/2)q^{2q/2-1}/(\sqrt{\pi}u^{(q+1)/2})$ , for  $u > 0$ , and  $r(u) = q/[(q+1)\sqrt{2\pi}]$ , for  $u = 0$ ,  $q > 0$ , where  $\Psi(a, x) = \int_0^x t^{a-1}e^{-t}dt$  is the lower incomplete gamma function; when  $q = 1$  the slash distribution coincides with the canonical slash distribution.

Let  $f_Z(\cdot)$  be the pdf of  $Z$  with  $Z$  given in (1). We have

$$f_Z(z) = \frac{f_S(z)}{F_S\left(\frac{1}{\sigma|\lambda}\right)}, \quad z \in A(\sigma, \lambda), \quad (4)$$

where  $f_S(\cdot)$  and  $F_S(\cdot)$  are the pdf and the cumulative distribution functions (cdf) of  $S \sim S(0, 1; r)$ , respectively.<sup>2</sup> Now, let  $z = h(y; \mu, \sigma, \lambda)$  (see (1)). Because the Jacobian of the transformation from  $y$  to  $z$  is  $|\partial z/\partial y| = y^{\lambda-1}/\mu^\lambda\sigma$  the pdf of  $Y$  is given by

$$f_Y(y) = \frac{y^{\lambda-1}}{\mu^\lambda\sigma} f_Z(z), \quad y > 0. \quad (5)$$

Since  $f_S(s) = r(s^2)$ , we have from (1) and (4) that (5) can be written as

$$f_Y(y) = \begin{cases} \frac{y^{\lambda-1}}{\mu^\lambda\sigma} \frac{r(z^2)}{R\left(\frac{1}{\sigma|\lambda}\right)}, & \text{if } \lambda \neq 0, \\ \frac{1}{y\sigma} r(z^2), & \text{if } \lambda = 0, \end{cases} \quad (6)$$

for  $y > 0$ , where  $R(s) = \int_{-\infty}^s r(u^2)du$ , for  $s \in \mathbb{R}$ . The cdf of  $Y$  is given by

$$F_Y(y) = \begin{cases} \frac{R(z)}{R\left(\frac{1}{\sigma|\lambda}\right)}, & \text{if } \lambda \leq 0, \\ \frac{R(z) - R\left(-\frac{1}{\sigma|\lambda}\right)}{R\left(\frac{1}{\sigma|\lambda}\right)}, & \text{if } \lambda > 0, \end{cases}$$

for  $y > 0$ .

Rigby and Stasinopoulos (2006) and Voudouris et al. (2012) present figures of probability density functions of the Box-Cox t and the Box-Cox power exponential distributions for different values of the parameters. The figures suggest, as expected, that the transformation parameter  $\lambda$  controls the skewness of the distribution, while the right tail/kurtosis behavior is controlled by the extra parameter (degrees of freedom parameter of the Box-Cox t distribution

<sup>1</sup>It is the distribution of  $Z/U^{1/q}$ , where  $q > 0$  and  $Z$  and  $U$  are independent random variables with standard normal and uniform distribution, respectively.

<sup>2</sup>If  $\sigma|\lambda| = 0$ ,  $1/\sigma|\lambda|$  is interpreted as  $\lim_{\sigma|\lambda \rightarrow 0} (1/\sigma|\lambda|) = \infty$  and  $F(1/\sigma|\lambda|)$  is taken as 1.

and the shape parameter of the Box-Cox power exponential distribution). Figure 1 shows the pdf of the Box-Cox Cole-Green (BCCG), Box-Cox t (BCT), Box-Cox power exponential (BCPE) and Box-Cox slash (BCSlash) distributions for a particular choice of the parameters. It is apparent that the BCT and BCSlash distributions have heavier right tail than the other distributions. Figure 2 shows the pdf of the BCSlash distribution for different values of the parameters. Note that the extra parameter  $q$  controls for right tail heaviness; see Section 3 for a detailed discussion of tail heaviness of the Box-Cox symmetric distributions.




Figure 1: Probability density functions of BCCG, BCT ( $\tau = 4$ ), BCPE ( $\tau = 1.5$ ) and BCSlash ( $q = 4$ ) for  $\mu = 7$ ,  $\sigma = 0.5$  and  $\lambda = 0.5$ .

It is straightforward to verify that if  $Y \sim BCS(\mu, \sigma, \lambda; r)$ , for  $\mu > 0$ ,  $\sigma > 0$  and  $\lambda \in \mathbb{R}$ , the following properties hold:

- (i)  $dY \sim BCS(d\mu, \sigma, \lambda; r)$ , for all constant  $d > 0$ , and hence  $\mu$  is a scale parameter;
- (ii)  $(Y/\mu)^d \sim BCS(1, d\sigma, \lambda/d; r)$ , for all constant  $d > 0$ . In particular,  $Y/\mu \sim BCS(1, \sigma, \lambda; r)$ ,  $(Y/\mu)^{1/\sigma} \sim BCS(1, 1, \sigma\lambda; r)$ , and, for  $\lambda > 0$ ,  $(Y/\mu)^\lambda \sim BCS(1, \lambda\sigma, 1; r)$ ;
- (iii) if  $\lambda = 1$  then  $Y$  has a truncated symmetric distribution with parameters  $\mu$  and  $\mu\sigma$  and support  $(0, \infty)$ ;
- (iv) from (6), we have that, if  $\lambda = 0$ ,  $Y$  has a log-symmetric distribution with parameters  $\mu$  and  $\sigma^2$  and density generating function  $r$  (Vanegas and Paula; 2016); it is denoted here by  $LS(\mu, \sigma; r)$ .
- (v) for integer  $k$ ,

$$E(Y^k) = \begin{cases} \mu^k E\left((1 + \sigma\lambda S)^{\frac{k}{\lambda}} I_{A(\sigma, \lambda)}(S)\right), & \text{if } \lambda \neq 0, \\ \mu^k E(\exp(k\sigma S)), & \text{if } \lambda = 0, \end{cases}$$

where  $I_{A(\sigma, \lambda)}(\cdot)$  is the indicator function of the set  $A(\sigma, \lambda)$  given in (3), and  $S \sim S(0, 1; r)$ . Hence, when  $\lambda = 0$  (or the truncation set  $\mathbb{R} \setminus A(\sigma, \lambda)$  has negligible probability under the




Figure 2: Probability density functions of  $BCSlash(\mu, \sigma, \lambda, q)$ .

$S(0, 1; r)$  distribution) the moments of  $Y$  can be obtained from the characteristic function of a standard symmetric distribution.

The BCS class of distributions allows easy parameter interpretation from its quantiles. Let  $s_\alpha$  denote the  $\alpha$ -quantile of the  $S \sim S(0, 1; r)$ , and  $z_\alpha$  be the  $\alpha$ -quantile of the truncated  $S$ , i.e. of the standard symmetric distribution  $S(0, 1; r)$  truncated at  $\mathbb{R} \setminus A(\sigma, \lambda)$ . We have

$$z_\alpha = \begin{cases} R^{-1} \left[ \alpha R \left( \frac{1}{\sigma|\lambda|} \right) \right], & \text{if } \lambda \leq 0, \\ R^{-1} \left[ 1 - (1 - \alpha) R \left( \frac{1}{\sigma|\lambda|} \right) \right], & \text{if } \lambda > 0. \end{cases}$$

Recall that  $R(\cdot)$  is the cdf of  $S \sim S(0, 1; r)$ . If  $Y \sim BCS(\mu, \sigma, \lambda; r)$  its  $\alpha$ -quantile is given by

$$y_\alpha = \begin{cases} \mu(1 + \sigma\lambda z_\alpha)^{\frac{1}{\lambda}}, & \text{if } \lambda \neq 0, \\ \mu \exp(\sigma z_\alpha), & \text{if } \lambda = 0. \end{cases}$$

We note that all the quantiles are proportional to  $\mu$ . In particular, if  $\lambda = 0$  we have  $z_{1/2} = 0$  and hence  $\mu$  is the median of  $Y$ . Moreover, if the truncation set  $\mathbb{R} \setminus A(\sigma, \lambda)$  has

negligible probability under the  $S(0, 1; r)$  distribution (this happens when  $\sigma\lambda$  is small),  $\mu$  is approximately equal to the median of  $Y$ . In order to give an interpretation for  $\sigma$ , we consider the centile-based coefficient of variation (Rigby and Stasinopoulos; 2006)

$$CV_Y = \frac{3}{4} \frac{y_{0.75} - y_{0.25}}{y_{0.5}}.$$

When  $\lambda$  is not far from zero and ignoring the truncation region,  $CV_Y \approx 1.5 \sinh(\sigma s_{0.75})$ , which is an increasing function of  $\sigma$ . Here,  $\sinh(\cdot)$  is the hyperbolic sine function. The approximation is exact when  $\lambda = 0$ . Therefore,  $\sigma$  can be seen as a relative dispersion parameter. Finally,  $\lambda$  is regarded as a skewness parameter since it determines the power transformation to symmetry.

At this point it is informative to compare our approach with an alternative, closely related, approach that uses the original Box-Cox transformation. A usual strategy for dealing with positive continuous asymmetric data is to employ a Box-Cox transformation in the data and to assume that the transformed data follow a normal distribution. The normal distribution can be replaced by the class of the continuous symmetric distributions in the Box-Cox transformation approach; see Cordeiro and Andrade (2011). Formally, this approach does not correspond to assume a coherent distribution for the data because the support of the transformed variable is not the entire real line, unless the transformation parameter is zero; this is known as the truncation problem. In order to take the correct support of the Box-Cox transformed data into account, a truncated normal (or symmetric) distribution may be assumed for the transformed data; see e.g. Poirier (1978) and Yang (1996). However, the truncation point will depend on the three parameters, namely the location, dispersion and transformation parameters. In our approach the truncation point depends on  $\sigma$  and  $\lambda$  only. Hence, it does not depend on the regressor values if a regression model is assumed for  $\mu$ . Although the truncation problem is usually disregarded in the statistical literature, alternative transformations have been proposed to overcome this problem; see, e.g., Yeo and Johnson (2000) and Yang (2006). Furthermore, the model parameters are interpreted as characteristics of the transformed data, not the original data. Our approach does not have these two shortcomings: a genuine distribution is assumed for the data and the parameters are interpretable in terms of characteristics of the original data, not the transformed data.

In Section 5, we present a comparison of alternative approaches in real data applications.

### 3 Tail heaviness

Heavy-tailed distributions have been frequently used to model phenomena in various fields such as finance and environmental sciences; see, for instance, Resnick (2007). A usual criterion for evaluating tail heaviness in the extreme value theory is the tail index of regular variation functions. Informally, a regular variation function behaves asymptotically as a

power function. Formally, a Lebesgue measurable function  $M : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  is regularly varying at infinity with index of regular variation  $\alpha$  ( $M \in RV_\alpha$ ), if  $\lim_{t \rightarrow \infty} M(ty)/M(t) = y^\alpha$  for  $y > 0$ . If  $\alpha = 0$ ,  $M$  is said to be a slowly varying function. The function  $M$  varies rapidly at infinity or is regularly varying at infinity with index  $-\infty$  ( $\infty$ ), or  $M \in RV_{-\infty}$  ( $M \in RV_\infty$ ), if, for  $y > 0$ ,  $\lim_{t \rightarrow \infty} M(ty)/M(t) := y^{-\infty}$  ( $\lim_{t \rightarrow \infty} M(ty)/M(t) := y^\infty$ ); see de Haan (1970, p. 4).<sup>3</sup>

A continuous distribution with cdf  $F$  is said to have a heavy right tail whenever  $\bar{F} := 1 - F$  is a regularly varying at infinity function with negative index of regular variation  $\alpha = -1/\xi$ ,  $\xi > 0$ , i.e.,  $\lim_{t \rightarrow \infty} \bar{F}(ty)/\bar{F}(t) = y^{-1/\xi}$ . The parameter  $\xi$  is called the tail index. From the l'Hôpital rule, this limit can be written as  $y \lim_{t \rightarrow \infty} f(ty)/f(t)$ , for  $y > 0$ , where  $f$  is the pdf corresponding to  $F$ . When the limit equals  $y^{-\infty}$  we say that  $F$  has a light (non-heavy) right tail and that the tail index is zero. When the limit equals 1, i.e.  $\bar{F}$  is a slowly varying function, we will say that  $F$  has right heavy tail with tail index  $\infty$ .

From de Haan (1970, Corollary 1.2.1) it follows that the tail index is invariant to location-scale transformations. Hence, from (2) the tail index of a  $S(\mu, \sigma; r)$  distribution is independent of  $\mu$  and  $\sigma$  and is obtained from

$$\mathcal{L}_S(w; r) = w \lim_{t \rightarrow \infty} \frac{r(t^2 w^2)}{r(t^2)}. \quad (7)$$

It can be shown that the tail index of the  $BCS(\mu, \sigma, \lambda; r)$  distribution, for all  $\mu > 0$  and all  $\sigma > 0$ , can be obtained from

$$\mathcal{L}_{BCS}(y, \mu, \sigma, \lambda; r) = \begin{cases} \mathcal{L}_S(y^\lambda; r), & \text{if } \lambda > 0, \\ \mathcal{L}_{LS}(y, \mu, \sigma; r), & \text{if } \lambda = 0, \\ y^\lambda, & \text{if } \lambda < 0, \end{cases}$$

where  $\mathcal{L}_{LS}(y, \mu, \sigma; r)$  corresponds to the limit that defines the tail index of the log-symmetric distributions and is given by (7) with  $tw$  in the numerator replaced by  $\sigma^{-1} \log(tw/\mu)$  and  $t$  in the denominator replaced by  $\sigma^{-1} \log(t/\mu)$ .

Table 1 gives the tail index of some Box-Cox symmetric distributions.<sup>4</sup> When  $\lambda > 0$ , the Box-Cox t and Box-Cox slash distributions have heavy right tail with the extra parameter (the degrees of freedom parameter  $\tau$  and the shape parameter  $q$  in the case of the t and the slash distributions, respectively) controlling the tail weight for fixed  $\lambda$ ; the Box-Cox Cole-Green (normal), Box-Cox power exponential, Box-Cox type I logistic and Box-Cox type II logistic distributions are all right light-tailed distributions, i.e. the tail index for these distributions are zero. The results for  $\lambda = 0$  reveal that the log-normal, log-power exponential with  $\tau > 1$  and log-type I logistic distributions are right light-tailed while the log-double exponential and the log-type II logistic distributions have heavy right tail with tail index determined

<sup>3</sup> $y^{-\infty} = \infty$ , if  $0 < y < 1$ ,  $= 1$ , if  $y = 1$ ,  $= 0$ , if  $y > 1$ ;  $y^\infty = 0$ , if  $0 < y < 1$ ,  $= 1$ , if  $y = 1$ ,  $= \infty$ , if  $y > 1$ .

<sup>4</sup>The tail indices were obtained using Maple 13; see <http://www.maplesoft.com>. The tail index for the log-power exponential distribution with  $\tau > 1$  was obtained for  $\tau \in \mathbb{Q}$ , and for the slash distribution, for  $q \in \mathbb{N}^*$ .



by  $\sigma$ . All the others have right heavy tail with tail index  $\infty$ . It is noteworthy that the extra parameters have no influence on the tail index. This suggests that the class of the log-symmetric distributions (Vanegas and Paula; 2016) is much more restrictive than the Box-Cox symmetric distributions defined in this paper with respect to tail flexibility. Finally, when  $\lambda < 0$ , all the Box-Cox symmetric distributions have heavy right tail with tail index equal to  $|\lambda|^{-1}$ .

Table 1: Tail index ( $\xi$ ) of some symmetric and Box-Cox symmetric distributions.

distribution	symmetric	BCS ( $\lambda > 0$ )	BCS ( $\lambda = 0$ )	BCS ( $\lambda < 0$ )
normal	0	0	0	$1/ \lambda $
double exponential	0	0	$\sigma/\sqrt{2}$	$1/ \lambda $
power exponential				
$\tau > 1$	0	0	0	$1/ \lambda $
$\tau = 1$	0	0	$\sigma/\sqrt{2}$	$1/ \lambda $
$\tau < 1$	0	0	$\infty$	$1/ \lambda $
Cauchy	1	$1/\lambda$	$\infty$	$1/ \lambda $
t	$1/\tau$	$1/(\lambda\tau)$	$\infty$	$1/ \lambda $
type I logistic	0	0	0	$1/ \lambda $
type II logistic	0	0	$\sigma$	$1/ \lambda $
canonical slash	1	$1/\lambda$	$\infty$	$1/ \lambda $
slash ( $q \in \mathbb{N}^*$ )	$1/q$	$1/(\lambda q)$	$\infty$	$1/ \lambda $

An alternative approach to compare tail heaviness of statistical distributions is considered by Rigby et al. (2014, Chapter 12). Here, we focus on right tail heaviness only. If the random variables  $Y_1$  and  $Y_2$  have continuous pdf's  $f_{Y_1}(y)$  and  $f_{Y_2}(y)$  and  $\lim_{y \rightarrow \infty} f_{Y_1}(y) = \lim_{y \rightarrow \infty} f_{Y_2}(y) = 0$  then  $Y_2$  has heavier right tail than  $Y_1$  if and only if  $\lim_{y \rightarrow \infty} (\log f_{Y_2}(y) - \log f_{Y_1}(y)) = \infty$ . The authors classify the possible asymptotic (large  $y$ ) behavior of the logarithm of a pdf in three major forms:  $-k_2(\log y)^{k_1}$ ,  $-k_4 y^{k_3}$  or  $-k_6 \exp(k_5 y)$ , with positive  $k$ 's. The three forms are decreasing in order of tail heaviness and, for the first form, decreasing  $k_1$  results in a heavier tail while decreasing  $k_2$  for fixed  $k_1$  results in heavier tail. Similarly, for the two other forms.

Table 2 gives the coefficients of the right tail asymptotic form of the logarithm of pdf's for some symmetric and Box-Cox symmetric distributions. Following Rigby et al. (2014), the right tail heaviness of the distributions in Table 2 can be classified in the following four types (the corresponding tail index for each type is given in parentheses):

- non-heavy tail: type II with  $k_3 \geq 1$  ( $\xi = 0$ );
- heavy tail (i.e. heavier than any exponential distribution) but lighter than any 'Paretian type' tail: type I with  $k_1 > 1$  and type II with  $0 < k_3 < 1$  ( $\xi = 0$ );

- ‘Paretian type’ tail: type I with  $k_1 = 1$  and  $k_2 > 1$  ( $\xi = 1/(k_2 - 1)$ );
- heavier than any ‘Paretian type’ tail: type I with  $k_1 = 1$  and  $k_2 = 1$  ( $\xi = \infty$ ).

It should be noted that distributions with right tail index  $\xi = 0$ , which are classified as having light (non-heavy) right tail according to the regular variation theory, are split into two categories in Rigby’s criterion: non-heavy right tail and heavy right tail but lighter than any ‘Paretian type’ tail.

When  $\lambda > 0$ , the Box-Cox t and Box-Cox slash distributions have ‘Paretian type’ right tail with the extra parameter controlling the right tail heaviness for fixed  $\lambda$ ; the Box-Cox power exponential distributions have non-heavy right tail ( $\tau \geq 1/\lambda$ ) or heavy right tail but lighter than any ‘Paretian type’ tail ( $\tau < 1/\lambda$ ), with  $\tau$  determining the tail heaviness for fixed  $\lambda$ . Depending on the value of  $\lambda$ , the Box-Cox Cole-Green and Box-Cox type I logistic distributions may have a non-heavy right tail ( $\lambda \geq 1/2$ ) or a heavy right tail but lighter than any ‘Paretian type’ tail ( $0 < \lambda < 1/2$ ); similarly for the Box-Cox type II logistic distribution with  $\lambda \geq 1$  and  $0 < \lambda < 1$ , respectively. From the coefficients for  $\lambda = 0$  we note that, as expected, the log-normal, log-power exponential with  $\tau > 1$  and log-type I logistic distributions have non-heavy right tail while the log-double exponential and the log-type II logistic distributions have heavy right tail but lighter than any ‘Paretian type’ tail; all the others have right heavier than any ‘Paretian type’ tail. When  $\lambda < 0$ , all the Box-Cox symmetric distributions in Table 2 have right ‘Paretian type’ tail with the tail heaviness controlled by  $\lambda$  only.

Table 2: Coefficients of the right tail asymptotic form of the log of the pdf for some symmetric and Box-Cox symmetric distributions.

distribution	symmetric	BCS ( $\lambda > 0$ )	BCS ( $\lambda = 0$ )	BCS ( $\lambda < 0$ )
normal	$k_3 = 2, k_4 = 1/(2\sigma^2)$	$k_3 = 2\lambda, k_4 = 1/(2\mu^{2\lambda}\sigma^2\lambda^2)$	$k_1 = 2, k_2 = 1/(2\sigma^2)$	$k_1 = 1, k_2 =  \lambda  + 1$
double exponential	$k_3 = 1, k_4 = \sqrt{2}/\sigma$	$k_3 = \lambda, k_4 = \sqrt{2}/(\mu^\lambda\sigma\lambda)$	$k_1 = 1, k_2 = \sqrt{2}/\sigma + 1$	$k_1 = 1, k_2 =  \lambda  + 1$
power exponential				
$\tau > 1$	$k_3 = \tau, k_4 = 1/(2p(\tau)^\tau\sigma^\tau)$	$k_3 = \lambda\tau, k_4 = 1/(2p(\tau)^\tau\mu^{\lambda\tau}\sigma^\tau\lambda^\tau)$	$k_1 = \tau, k_2 = 1/(2p(\tau)^\tau\sigma^\tau)$	$k_1 = 1, k_2 =  \lambda  + 1$
$\tau = 1$	$k_3 = 1, k_4 = \sqrt{2}/\sigma$	$k_3 = \lambda, k_4 = \sqrt{2}/(\mu^\lambda\sigma\lambda)$	$k_1 = 1, k_2 = \sqrt{2}/\sigma + 1$	$k_1 = 1, k_2 =  \lambda  + 1$
$\tau < 1$	$k_3 = \tau, k_4 = 1/(2p(\tau)^\tau\sigma^\tau)$	$k_3 = \lambda\tau, k_4 = 1/(2p(\tau)^\tau\mu^{\lambda\tau}\sigma^\tau\lambda^\tau)$	$k_1 = 1, k_2 = 1$	$k_1 = 1, k_2 =  \lambda  + 1$
Cauchy	$k_1 = 1, k_2 = 2$	$k_1 = 1, k_2 = \lambda + 1$	$k_1 = 1, k_2 = 1$	$k_1 = 1, k_2 =  \lambda  + 1$
t	$k_1 = 1, k_2 = \tau + 1$	$k_1 = 1, k_2 = \lambda\tau + 1$	$k_1 = 1, k_2 = 1$	$k_1 = 1, k_2 =  \lambda  + 1$
type I logistic	$k_3 = 2, k_4 = 1/\sigma^2$	$k_3 = 2\lambda, k_4 = 1/(\mu^{2\lambda}\sigma^2\lambda^2)$	$k_1 = 2, k_2 = 1/\sigma^2$	$k_1 = 1, k_2 =  \lambda  + 1$
type II logistic	$k_3 = 1, k_4 = 1/\sigma$	$k_3 = \lambda, k_4 = 1/(\mu^\lambda\sigma\lambda)$	$k_1 = 1, k_2 = 1/\sigma + 1$	$k_1 = 1, k_2 =  \lambda  + 1$
canonical slash	$k_1 = 1, k_2 = 2$	$k_1 = 1, k_2 = \lambda + 1$	$k_1 = 1, k_2 = 1$	$k_1 = 1, k_2 =  \lambda  + 1$
slash	$k_1 = 1, k_2 = q + 1$	$k_1 = 1, k_2 = \lambda q + 1$	$k_1 = 1, k_2 = 1$	$k_1 = 1, k_2 =  \lambda  + 1$

The study presented in this section shows that the class of the Box-Cox symmetric distributions is very flexible for modeling positive data displaying different right tail behaviors. It covers from right light-tailed distributions to heavier than any ‘Paretian type’ tailed distribution. More importantly, some distributions in this class have an extra parameter that controls the right tail heaviness. Slightly heavy-tailed data may be modeled using a Box-Cox

power exponential distribution, which has a ‘slightly’ heavy right tail (heavy tail but lighter than any ‘Paretian type’ tail) when  $\lambda > 0$  and  $\lambda\tau < 1$  with the tail weight determined by  $\tau$ . Moderately or highly heavy-tailed data may require a Box-Cox t or Box-Cox slash distribution, because both are right ‘Paretian type’ tailed distributions with a parameter that controls the right tail heaviness.

## 4 Likelihood-based inference

The log-likelihood function for a single observation  $y$  taken from a BCS distribution is given by

$$\ell(\mu, \sigma, \lambda) = (\lambda - 1) \log y - \lambda \log \mu - \log \sigma + \log r(z^2) - \log R\left(\frac{1}{\sigma|\lambda|}\right),$$

where  $z = h(y; \mu, \sigma, \lambda)$  with  $h(y; \mu, \sigma, \lambda)$  given in (1); the last term in  $\ell$  is zero if  $\lambda = 0$ . The score vector and the Hessian matrix are obtained from the first and second derivatives of  $\ell$  with respect to the parameters; see the Appendix.

The maximum likelihood estimates of  $\mu$  and  $\sigma$ , for fixed  $\lambda$ , from a sample of  $n$  independent observations  $y_1, \dots, y_n$ , are solution of the system of equations

$$\mu = \begin{cases} \frac{1}{(n\sigma\lambda)^{1/\lambda}} \left( \sum_{i=1}^n \omega_i y_i^\lambda z_i \right)^{1/\lambda}, & \text{if } \lambda \neq 0, \\ \left( \prod_{i=1}^n y_i^{\omega_i} \right)^{1/\sum_{i=1}^n \omega_i}, & \text{if } \lambda = 0, \end{cases}$$

$$\sigma^2 = \begin{cases} \frac{1}{n\lambda^2(1-\delta)} \sum_{i=1}^n \omega_i \left[ \left( \frac{y_i}{\mu} \right)^\lambda - 1 \right]^2, & \text{if } \lambda \neq 0, \\ \frac{1}{n} \sum_{i=1}^n \omega_i \left( \log \frac{y_i}{\mu} \right)^2, & \text{if } \lambda = 0, \end{cases}$$

where

$$\delta = \frac{1}{\sigma|\lambda|} \frac{r\left(\frac{1}{\sigma|\lambda|}\right)}{R\left(\frac{1}{\sigma|\lambda|}\right)},$$

$z_i = h(y_i; \mu, \sigma, \lambda)$ , and  $\omega_i = \omega(z_i)$  with  $\omega(z) = -2r'(z^2)/r(z^2)$  being a weighting function that depends on  $r$ . We note that the maximum likelihood estimates of  $\mu$  and  $\sigma$  involve weighted arithmetic and geometric averages of the contributions of each observation  $y_i$  with weight  $\omega(z_i)$ . Table (3) gives  $\omega(z)$  for several BCS distributions. Note that some distributions in the BCS class produce robust estimation against outliers. For instance, for the Box-Cox t and Box-Cox power exponential (with  $\tau < 2$ ) distributions the weighting function is decreasing in the observation of  $Y$ . Hence, outlier observations have smaller weights in the estimation of the parameters.

The system of likelihood equations for  $(\mu, \sigma, \lambda)$  does not have analytical solution. Furthermore, it may involve an addition equation relative to an extra parameter (for instance, the degrees of freedom parameter of the BCT distribution). Maximization of the likelihood

Table 3: Weighting functions for some BCS distributions.

BCS distribution	$\omega(z)$
normal	1
double exponential	$\sqrt{2}/ z $
power exponential	$\tau(z^2)^{\tau/2-1}/(2p(\tau)^\tau)$
Cauchy	$2/(1+z^2)$
t	$(\tau+1)/(\tau+z^2)$
type I logistic	$-2(\exp\{-z^2\}-1)/(\exp\{-z^2\}+1)$
type II logistic	$(\exp\{- z \}-1)/[ z (\exp\{- z \}+1)]$
canonical slash	$2/z^2 - \exp\{-z^2/2\}/(1 - \exp\{-z^2/2\})$
slash	$2\Psi((q+3)/2, z^2/2)/(z^2\Psi((q+1)/2, z^2/2))$

function is implemented in the package `gamLSS` in R for the BCT, BCCG and BCPE distributions through the CG and the RS algorithms (Rigby and Stasinopoulos; 2005; Rigby et al.; 2014). It is noteworthy that `gamLSS` allows the fit of regression models with monotonic link functions relating the parameters ( $\mu$ ,  $\sigma$ ,  $\lambda$  and the possibly extra parameter) to explanatory variables through parametric or semi-parametric additive models.

It is of particular interest to test the null hypothesis  $H_0 : \lambda = 0$ ; recall that the BCS distributions reduce to the log-symmetric distributions when  $\lambda = 0$ . In order to evaluate the performance of the likelihood ratio test of  $H_0$  against a two sided alternative, we now present a small Monte Carlo simulation study. We set  $\mu = \sigma = 1$  and generate 10,000 samples of sizes  $n = 100, 200, 300, 500$  from a Box-Cox t distribution with two different values for the degrees of freedom parameter, namely  $\tau = 4, 10$ . The samples are generated under the null hypothesis. We assume that  $\tau$  is known, and we estimate the remaining parameters using the function `optim` in R with the analytical derivatives derived in the Appendix and with numerical derivatives. The type I error probability estimated from the simulated samples for a nominal level  $\alpha = 5\%$  are presented in Table 4. The figures in the table reveal that the likelihood ratio test performs well regardless of whether analytical or numerical derivatives are employed. As expected, the type I error probabilities are close to the nominal level of the test for the sample sizes considered here.

## 5 Applications and comparison of alternative approaches

In this section, we present applications of the Box-Cox distributions in the analysis of micro and macronutrients intake. The data refer to observations of nutrients intake based on the first 24-hour dietary recall interview for  $n = 368$  individuals. For each nutrient, we assume that the data  $Y_1, \dots, Y_n$  are independent.

Table 4: Type I error probability of the likelihood ratio test of  $H_0 : \lambda = 0$  using analytical and numerical derivatives; Box-Cox t distribution with  $\mu = \sigma = 1$  and  $\tau = 4$  and 10.

$n$	$\tau = 4$		$\tau = 10$	
	anal. der.	num. der.	anal. der.	num. der.
100	0.043	0.043	0.044	0.044
200	0.051	0.051	0.052	0.052
300	0.050	0.050	0.049	0.049
500	0.049	0.049	0.049	0.049

First, we fitted different models to each of all the 33 nutrients. All the models considered in the first analysis are constructed from the Student-t distribution and from its limiting case when the degrees of freedom parameter goes to infinity, i.e. the normal distribution. The following models were considered: Box-Cox t (BCT); Box-Cox Cole-Green (BCCG), which corresponds to the BCT model with  $\tau \rightarrow \infty$ ; skew-normal (SN) and skew-t (ST) (Azzalini; 2005); and transformed symmetric models with normal (TN) and t (TT) errors (Cordeiro and Andrade; 2011). The TN (TT) model assumes that the original Box and Cox (1964) transformed data follow a normal (Student-t) distribution. The unknown parameters (including the degrees of freedom parameter) were estimated by the maximum likelihood method. For the BCT, BCCG, SN and ST distributions, we used the `gamlss` package implemented in R, while for the TN and TT models we used both the function `optim` in R and the PROC NLP in SAS. Goodness-of-fit was evaluated using the following criteria: Akaike information criterion (AIC) and Anderson-Darling statistics (AD, ADR, and AD2R); see Luceño (2005, Tables 1, 2 and B.1). AD is a global measure of lack-of-fit, while both ADR and AD2R are more sensitive to the lack of fit in the right tail of the distribution; AD2R puts more weight in the right tail than ADR. For all the four criteria a lower value is preferred.

Tables 5-8 present the goodness-of-fit statistics for all the fitted models to 22 and 11 micro and macronutrients intakes data. Underlined numbers indicate the best fitting model. The blank cells in the tables indicate that the algorithm employed for maximum likelihood estimation did not achieve convergence or produced unrealistic estimates. The tables convey important information. First, the datasets cover a wide range of light-tailed to heavy-tailed data. This can be seen by the estimated values of the degrees of freedom parameter under the BCT model;  $\widehat{\tau}_{BCT}$  ranges from 2.2 to 196.0 (see Table 5). Second, no convergence problem was observed while fitting the BCT model, the TT model, and the TN model. The maximum likelihood estimation under the SN model and under the BCCG model did not achieve convergence in 10 cases. Under the ST model, convergence was not reached in two cases. Third, according to the AIC criterion, the BCT model achieved the best fit in most of the cases (26 out of 33 cases); the AICs of the BCT and TT fits coincided in 20 cases. The models derived from the normal distribution, namely the BCCG, SN, and TN models, did

not perform well in general, except for two cases in which the estimated degrees of freedom parameter under the BCT model was very large ( $\widehat{\tau}_{BCT} > 50$ ). In such cases, the BCCG and the TN models achieved the best fits. Forth, according to all the Anderson-Darling criteria, the BCCG, skew-normal and TN models did not outperform the BCT, ST and TT models in any of the cases. It suggests that the tail behavior of the nutrients data are better modeled by the distributions derived from the t distribution. The BCT and TT models were the best fitting models in most of the cases. Overall, we conclude that the Box-Cox t model performed better than the other models. The transformed t model behaved as well as the Box-Cox t model in many cases. However, as pointed out earlier, the transformed t model has some drawbacks that are overcome by the Box-Cox t model.

We now turn to a detailed analysis of the data on the intake of animal protein and energy. Adjusted boxplots (Hubert and Vandervieren; 2008) are presented in Figure 3 and show that the data sets are asymmetric and contain outlying observations. It is noteworthy that the data set on energy intake contains an outlier that is well above the second highest observed intake. We fitted the Box-Cox t, Box-Cox Cole-Green, Box-Cox power exponential and Box-Cox slash distributions to each data set. For the BCT, BCCG, and BCPE models, we used the `gamlss` package implemented in R, while for the BCSlash model we used the function `optim` in R. Tables 9 and 10 give descriptive statistics of the data, and parameter estimates and goodness-of-fit statistics for each fitted model. The descriptive statistics confirm the findings in the boxplots. For both data sets, the Akaike information criteria are similar for the BCT, BCPE, and BCSlash models, and both are smaller than that for the BCCG model. The Anderson-Darling statistics indicate that the BCT model gives the best fit. Note that the BCCG model gives a poor right tail fit for both data sets. This lack of fit is most pronounced for the energy intake data set, for which  $AD2R = 112.25$  for the BCCG model while  $AD2R = 1.78$ ,  $AD2R = 4.18$ , and  $AD2R = 2.15$  for the BCT, BCPE, and BCSlash models, respectively.

Figure 4 presents qq plots for quantile residuals  $\widehat{r} = \Phi^{-1}(\widehat{F}_Y(y))$ , where  $\Phi(\cdot)$  is the cdf of the standard normal distribution and  $\widehat{F}_Y(y)$  is the fitted cdf of  $Y$ . If the model is correct, the quantile residuals are expected to behave approximately as standard normal quantiles (Dunn and Smyth; 1996). The lack of fit of the BCCG model in the right tail is clear from the plots for both data sets. For the animal protein data, the residual plots are similar for the BCT and BCPE models and indicate reasonable fits. On the other hand, for the energy intake data set, which seems to require a right heavier tailed model, the BCT model provides a slightly better fit than the BCPE and BCSlash models.

Table 5: AIC for the fitted BCT, ST, TT, BCCG, SN, and TN models; micronutrients and macronutrients datasets.

micronutrient	$\widehat{\tau}_{BCT}$	BCT	ST	TT	BCCG	SN	TN
vitamin A (mcg)	7.2	<u>5807.4</u>	5809.2	<u>5807.4</u>			5822.7
vitamin D (mcg)	6.8	<u>1688.7</u>	1707.7	1690.5		1745.3	1698.9
vitamin E (mg)	6.9	<u>1812.8</u>	1818.4	<u>1812.8</u>	1824.3	1902.4	1824.3
vitamin K (mcg)	7.8	<u>4354.3</u>	4356.5	<u>4354.3</u>			4368.4
vitamin C (mg)	2.2	<u>4022.5</u>		4029.9			4120.9
vitamin B1 (mg)	6.8	<u>709.7</u>	711.6	<u>709.7</u>	720.9	778.4	720.9
vitamin B2 (mg)	6.5	<u>701.6</u>	702.2	<u>701.6</u>	716.1	769.6	716.1
vitamin B3 (mg)	6.8	<u>2722.4</u>	2726.3	<u>2722.4</u>	2730.8	2778.9	2730.8
vitamin B6 (mg)	51.3	853.2	858.5	853.2	<u>851.3</u>	866.9	<u>851.3</u>
vitamin B12 (mcg)	2.5	<u>1782.6</u>	1808.4	1782.8			1887.3
pantothenic acid (mg)	8.3	1466.0	<u>1465.5</u>	1466.0	1474.1	1509.3	1474.1
folate (mcg)	4.9	<u>4795.4</u>	4800.7	<u>4795.4</u>	4823.4		4823.4
calcium (mg)	13.3	<u>5311.3</u>	5312.6	<u>5311.3</u>	5312.5	5342.8	5312.5
phosphorus (mg)	14.7	5548.5	<u>5548.4</u>	5548.5	5549.0	5561.5	5549.0
magnesium (mg)	8.6	<u>4474.8</u>	4476.8	<u>4474.8</u>	4481.5	4522.8	4481.5
iron (mg)	5.9	2409.7	<u>2409.3</u>	2409.7	2431.9		2431.9
zinc (mg)	14.6	<u>2185.3</u>	2188.7	<u>2185.3</u>	2185.4	2203.4	2185.4
copper (mg)	5.5	<u>566.1</u>		<u>566.1</u>			593.6
selenium (mcg)	5.2	<u>3992.6</u>	4000.6	<u>3992.6</u>			4020.8
sodium (mg)	4.6	<u>6525.4</u>	6535.6	<u>6525.4</u>			6572.3
potassium (mg)	9.3	<u>6144.5</u>	6147.5	<u>6144.5</u>	6151.9	6195.2	6151.9
manganese (mg)	2.5	<u>1616.5</u>	1647.0	1623.7			1656.2
macronutrient	$\widehat{\tau}_{BCT}$	BCT	ST	TT	BCCG	SN	TN
protein (g)	10.2	<u>3659.5</u>	3660.9	<u>3659.5</u>	3662.5	3678.8	3662.5
energy (kcal)	6.1	<u>5861.9</u>	5863.7	<u>5861.9</u>	5876.1	5912.8	5876.1
fiber (g)	10.0	<u>2652.2</u>	2653.5	<u>2652.2</u>	2655.6	2669.2	2655.6
carbohydrate (g)	10.5	4360.0	<u>4359.9</u>	4360.0	4366.5	4382.3	4366.5
total fat (g)	13.9	<u>3587.0</u>	3593.7	<u>3587.0</u>	3587.5	3651.9	3587.5
animal protein (g)	4.9	<u>3514.4</u>	3532.5	3516.1	3526.3	3553.9	3526.5
vegetable protein (g)	6.6	<u>2963.3</u>	2964.4	<u>2963.3</u>	2972.6	2988.5	2972.6
saturated fat (g)	196.0	2819.6	2822.9	2819.6	<u>2817.7</u>	2844.2	<u>2817.7</u>
monounsaturated fat (g)	12.6	<u>2857.1</u>	2865.4	<u>2857.1</u>	2858.4	2920.1	2858.4
polyunsaturated fat (g)	7.0	2596.9	<u>2596.4</u>	2596.9	2612.4	2771.9	2612.4
cholesterol (mg)	5.8	<u>4724.7</u>	4753.7	4725.6		4782.9	4728.5

Table 6: Anderson-Darling statistics for the fitted BCT, ST, TT, BCCG, SN, and TN models; micronutrients datasets.

micronutrient	statistic	BCT	ST	TT	BCCG	SN	TN
vitamin A	AD	1.12	<u>0.94</u>	1.12			1.47
	ADR	0.64	<u>0.53</u>	0.64			0.94
	AD2R	<u>6.26</u>	8.46	6.27			> 100
vitamin D	AD	<u>0.25</u>	0.67	0.28		3.30	0.64
	ADR	<u>0.11</u>	0.32	0.13		2.27	0.38
	AD2R	<u>3.40</u>	13.95	3.80		> 100	> 100
vitamin E	AD	<u>0.21</u>	0.30	<u>0.21</u>	0.91	7.62	0.91
	ADR	<u>0.13</u>	0.18	<u>0.13</u>	0.52	5.17	0.52
	AD2R	<u>3.39</u>	8.58	<u>3.39</u>	61.43	> 100	61.43
vitamin K	AD	<u>0.60</u>	22.69	<u>0.60</u>			1.28
	ADR	<u>0.35</u>	2.22	<u>0.35</u>			0.86
	AD2R	55.60	> 100	<u>55.44</u>			> 100
vitamin C	AD	<u>0.70</u>		0.71			7.92
	ADR	<u>0.34</u>		0.39			4.23
	AD2R	<u>6.28</u>		6.35			> 100
vitamin B1	AD	0.20	<u>0.17</u>	0.20	1.02	6.29	1.02
	ADR	<u>0.10</u>	<u>0.10</u>	0.11	0.47	3.96	0.47
	AD2R	<u>2.83</u>	4.36	<u>2.83</u>	> 100	> 100	> 100
vitamin B2	AD	<u>0.24</u>	<u>0.24</u>	<u>0.24</u>	1.02	5.90	1.02
	ADR	<u>0.13</u>	0.14	<u>0.13</u>	0.46	3.79	0.47
	AD2R	<u>2.10</u>	4.36	<u>2.10</u>	> 100	> 100	> 100
vitamin B3	AD	<u>0.22</u>	0.30	<u>0.22</u>	1.07	6.54	1.07
	ADR	<u>0.15</u>	0.18	<u>0.15</u>	0.56	4.18	0.56
	AD2R	<u>3.46</u>	4.23	<u>3.46</u>	17.29	> 100	17.25
vitamin B6	AD	<u>0.37</u>	0.42	0.38	0.45	1.69	0.45
	ADR	<u>0.18</u>	0.21	<u>0.18</u>	0.21	1.04	0.21
	AD2R	4.13	5.00	<u>4.10</u>	4.44	27.4	4.45
vitamin B12	AD	<u>0.36</u>	1.23	0.37			8.69
	ADR	<u>0.26</u>	0.77	0.27			5.08
	AD2R	<u>8.88</u>	22.56	9.67			> 100
panthotenic acid	AD	<u>0.24</u>	0.26	<u>0.24</u>	0.65	4.59	0.65
	ADR	<u>0.10</u>	0.16	<u>0.10</u>	0.34	3.10	0.34
	AD2R	<u>2.38</u>	2.98	<u>2.38</u>	3.77	> 100	13.77
folate	AD	<u>0.19</u>	0.23	<u>0.19</u>	1.83		1.83
	ADR	<u>0.12</u>	0.13	<u>0.12</u>	0.94		0.95
	AD2R	4.61	14.68	<u>4.60</u>	> 100		> 100
calcium	AD	0.28	<u>0.23</u>	0.28	0.48	4.06	0.48
	ADR	<u>0.11</u>	<u>0.11</u>	<u>0.11</u>	0.18	2.38	0.18
	AD2R	<u>2.43</u>	2.52	<u>2.43</u>	7.49	> 100	7.50



Table 7: Anderson-Darling statistics for the fitted BCT, ST, TT, BCCG, SN, and TN models; micronutrients datasets (*cont.*).

micronutrient	statistic	BCT	ST	TT	BCCG	SN	TN
phosphorus	AD	0.25	<u>0.21</u>	0.25	0.36	1.50	0.36
	ADR	0.15	<u>0.13</u>	0.15	0.18	1.00	0.19
	AD2R	<u>2.42</u>	2.94	2.43	5.58	> 100	5.56
magnesium	AD	<u>0.31</u>	0.32	<u>0.31</u>	0.66	4.07	0.66
	ADR	<u>0.18</u>	0.20	<u>0.18</u>	0.36	2.77	0.36
	AD2R	<u>2.76</u>	5.08	<u>2.76</u>	25.34	> 100	25.36
iron	AD	0.25	<u>0.12</u>	0.25	1.21		1.21
	ADR	0.10	<u>0.07</u>	0.10	0.45		0.45
	AD2R	2.35	8.01	<u>2.34</u>	> 100		> 100
zinc	AD	<u>0.16</u>	0.19	<u>0.16</u>	0.38	1.58	0.38
	ADR	<u>0.09</u>	0.10	<u>0.09</u>	0.19	1.04	0.19
	AD2R	1.92	2.96	<u>1.91</u>	8.28	> 100	7.97
copper	AD	<u>0.37</u>		<u>0.37</u>			1.74
	ADR	<u>0.21</u>		<u>0.21</u>			0.99
	AD2R	<u>16.10</u>		16.25			> 100
selenium	AD	<u>0.21</u>	0.28	<u>0.21</u>			1.70
	ADR	<u>0.12</u>	0.17	<u>0.12</u>			0.83
	AD2R	7.22	> 100	<u>7.14</u>			> 100
sodium	AD	<u>0.18</u>	0.41	<u>0.18</u>			2.28
	ADR	<u>0.09</u>	0.20	<u>0.09</u>			1.24
	AD2R	8.28	99.13	<u>8.27</u>			> 100
potassium	AD	<u>0.34</u>	0.37	<u>0.34</u>	0.57	2.90	0.57
	ADR	<u>0.22</u>	0.25	<u>0.22</u>	0.34	2.07	0.34
	AD2R	10.71	24.71	<u>10.68</u>	> 100	> 100	> 100
manganese	AD	1.29	2.23	<u>0.92</u>			4.64
	ADR	1.03	1.67	<u>0.69</u>			2.85
	AD2R	> 100	74.60	<u>28.29</u>			56.04

Table 8: Anderson-Darling statistics for the fitted BCT, ST, TT, BCCG, SN, and TN models; macronutrients datasets.

macronutrient	statistics	BCT	ST	TT	BCCG	SN	TN
protein	AD	<u>0.23</u>	0.25	<u>0.23</u>	0.54	2.97	0.54
	ADR	<u>0.14</u>	0.15	<u>0.14</u>	0.30	2.02	0.30
	AD2R	3.60	4.62	<u>3.58</u>	10.93	67.43	10.93
energy	AD	<u>0.19</u>	0.20	<u>0.19</u>	1.13	4.13	1.10
	ADR	<u>0.10</u>	0.11	<u>0.10</u>	0.58	2.78	0.54
	AD2R	<u>1.78</u>	2.60	<u>1.78</u>	> 100	> 100	> 100
fiber	AD	<u>0.21</u>	<u>0.21</u>	<u>0.21</u>	0.51	1.72	0.51
	ADR	<u>0.11</u>	0.13	<u>0.11</u>	0.26	1.20	0.26
	AD2R	2.03	2.50	<u>2.02</u>	11.28	> 100	11.26
carbohydrate	AD	0.26	<u>0.20</u>	0.25	0.39	1.36	0.39
	ADR	0.13	<u>0.12</u>	0.13	0.17	0.91	0.17
	AD2R	5.22	16.34	<u>5.21</u>	> 100	> 100	> 100
total fat	AD	<u>0.41</u>	0.52	<u>0.41</u>	0.62	5.66	0.62
	ADR	<u>0.24</u>	0.27	0.25	0.36	3.77	0.36
	AD2R	5.70	11.12	<u>5.54</u>	14.46	> 100	14.46
animal protein	AD	<u>0.35</u>	0.42	<u>0.35</u>	1.46	2.84	1.43
	ADR	0.16	0.16	<u>0.14</u>	0.70	1.74	0.68
	AD2R	3.19	4.10	<u>3.03</u>	23.47	> 100	22.34
vegetable protein	AD	0.25	<u>0.10</u>	0.25	1.08	2.58	1.08
	ADR	0.10	<u>0.07</u>	0.10	0.43	1.62	0.43
	AD2R	1.98	<u>1.81</u>	1.98	39.96	> 100	39.88
saturated fat	AD	<u>0.16</u>	0.36	<u>0.16</u>	0.17	2.66	0.17
	ADR	<u>0.08</u>	0.14	<u>0.08</u>	<u>0.08</u>	1.72	<u>0.08</u>
	AD2R	3.07	3.76	<u>3.06</u>	3.27	> 100	3.27
monounsaturated fat	AD	<u>0.33</u>	0.63	<u>0.33</u>	0.62	5.17	0.62
	ADR	<u>0.11</u>	0.22	<u>0.11</u>	0.23	3.39	0.26
	AD2R	4.42	12.78	<u>4.40</u>	29.04	> 100	29.06
polyunsaturated fat	AD	0.57	<u>0.43</u>	0.57	1.02	21.93	1.02
	ADR	<u>0.26</u>	<u>0.26</u>	<u>0.26</u>	0.51	13.18	0.51
	AD2R	<u>3.52</u>	10.56	<u>3.52</u>	86.75	> 100	86.57
cholesterol	AD	0.32	0.35	<u>0.28</u>		4.22	1.16
	ADR	0.13	0.15	<u>0.12</u>		2.45	0.51
	AD2R	3.40	<u>2.98</u>	3.34		> 100	11.01

For the energy intake data, the estimate of the skewness parameter ( $\lambda$ ) is indistinguishable from zero for the four models. Its estimates are close to zero, and the standard errors are relatively large. It suggests that log-symmetric models may be appropriate. Table 11 presents the parameter estimates and the goodness-of-fit statistics for the log-t, log-normal, log-power exponential, and log-slash model fits. Comparing the figures in Tables 10 and 11 one may notice that AD and ADR, and the estimates for  $\mu$ ,  $\sigma$  and  $\tau$  do not change much and the

AICs are slightly smaller for the log-symmetric models. On the other hand, AD2R dropped from 112.25 (BCCG model) to 48.17 (log-normal model) and from 4.18 (BCPE model) to 2.70 (log-PE model). The change in AD2R is small when one moves from the BCT model to the log-t model or from the BCSlash model to the log-slash model. Also, the quantile residuals plots (not shown) are similar to those presented in Figure 4. Taking parsimony into account, we may conclude that the best fitting model is the log-t model.




Figure 3: Adjusted boxplots; animal protein data (left) and energy intake (right).

Table 9: Descriptive statistics, parameter estimates (standard error in parentheses) and goodness-of-fit measures; animal protein intake data.

statistics (mg)	min	.25-quantile	median	mean (s.d.)	.75-quantile	max
	0.02	30.51	44.26	52.27 (34.02)	67.59	207.10
distribution	BCT	BCCG	BCPE	BCSlash		
$\mu$	45.74 (1.46)	46.37 (1.50)	44.73 (1.36)	45.83 (1.47)		
$\sigma$	0.55 (0.03)	0.67 (0.02)	0.70 (0.03)	0.45 (0.02)		
$\lambda$	0.42 (0.07)	0.44 (0.06)	0.42 (0.07)	0.42 (0.10)		
$\tau$	4.90 (0.92)		1.24 (0.12)			
$q$					3.02 (0.35)	
AIC	3514.4	3526.30	3511.4	3517.0		
AD	0.35	1.46	0.51	0.43		
ADR	0.16	0.70	0.26	0.18		
AD2R	3.19	23.33	3.42	3.64		

Table 10: Descriptive statistics, parameter estimates (standard errors in parentheses) and goodness-of-fit measures; energy intake data.


statistics (kcal)	min	.25-quantile	median	mean (s.d.)	.75-quantile	max
		298.80	1356.00	1723.00	1868.00 (838.35)	2197.00


distribution	BCT	BCCG	BCPE	BCSlash
$\mu$	1725.00 (34.48)	1726.00 (36.86)	1724.00 (34.14)	1725.00 (34.69)
$\sigma$	0.34 (0.02)	0.41 (0.02)	0.41 (0.02)	0.29 (0.01)
$\lambda$	0.05 (0.12)	0.07 (0.10)	0.06 (0.11)	0.05 (0.14)
$\tau$	6.14 (1.37)		1.40 (0.14)	
$q$				3.71 (0.44)


AIC	5861.9	5876.1	5864.0	5863.3
AD	0.19	1.13	0.25	0.23
ADR	0.10	0.58	0.14	0.12
AD2R	1.78	112.25	4.18	2.15




(a) BCT




(b) BCCG




(c) BCPE




(d) BCSlash




(e) BCT



(f) BCCG



(g) BCPE



(h) BCSlash

Figure 4: qq plots for quantile residuals for the BCT, BCCG, BCPE and BCSlash model fits for the animal protein intake data (first row) and energy intake data (second row).

Table 11: Parameter estimates (standard errors in parentheses) and goodness-of-fit statistics for the log- $t$ , log-normal, log-power exponential and log-slash models; energy intake data.

distribution	log-t	log-normal	log-PE	log-slash
$\mu$	1722.00 (34.44)	1717.00 (36.74)	1721.00 (34.09)	1722.00 (34.63)
$\sigma$	0.34 (0.02)	0.41 (0.02)	0.41 (0.02)	0.28 (0.01)
$\tau$	6.09 (1.34)		1.40 (0.14)	
$q$				3.71 (0.44)
AIC	5860.01	5874.72	5862.27	5861.46
AD	0.19	1.16	0.26	0.23
ADR	0.10	0.55	0.15	0.12
AD2R	1.80	48.17	2.70	2.11

For the BCT model fitted to the animal protein intake data the estimates of  $\mu$ ,  $\sigma$ ,  $\lambda$ , and  $\tau$  reported in Table 9 are  $\widehat{\mu}_P = 45.74$ ,  $\widehat{\sigma}_P = 0.55$ ,  $\widehat{\lambda}_P = 0.42$ , and  $\widehat{\tau}_P = 4.90$ , respectively. For the log- $t$  model fitted to the energy intake data the estimates of  $\mu$ ,  $\sigma$ , and  $\tau$  are  $\widehat{\mu}_E = 1722.00$ ,  $\widehat{\sigma}_E = 0.34$ , and  $\widehat{\tau}_E = 6.09$ , respectively; see Table 11. Because  $\widehat{\sigma}_P \widehat{\lambda}_P$  is small,  $\widehat{\mu}_P$  may be seen as an estimate of the population median of the animal protein intake. As expected,  $\widehat{\mu}_P$  is close to the sample median (44.26). Similarly,  $\widehat{\mu}_E$ , which is an estimate of the population median of the energy intake, is close to the sample median (1723.00). Additionally,  $\widehat{\sigma}_P$  is considerably larger than  $\widehat{\sigma}_E$  indicating that the relative dispersion of the population intake of animal protein is larger than that of energy. The estimates of the degrees of freedom parameter  $\tau$ , both not large, reveal the need for right heavy tailed distributions for an adequate fit for the data.

## 6 Concluding remarks

This paper proposed a new class of distributions, the Box-Cox symmetric distributions. It contains some well known distributions as special cases and allows the definition of new distributions, such as the Box-Cox slash distribution. It is particularly suitable for inference on positively skewed, possibly heavy-tailed, data. It permits easy parameter interpretation, a desirable feature for modeling.

There is clear possibility for extension to regression models. Some or all the parameters of the BCS distributions may be modeled by a link function and a linear or nonlinear regression model structure. The GAMLSS framework (Rigby and Stasinopoulos; 2005) is a natural tool for implementing BCS regression models. It allows the regression structure to include parametric and nonparametric terms and random effects. Box-Cox  $t$ , Box-Cox Cole-Green, and Box-Cox power exponential models are already implemented in `gamlss` package in R.

Some BCS distributions include an extra parameter; e.g., the degrees of freedom parameter of the BCT distribution. We have not faced convergence problems or unrealistic estimation when the additional parameter is estimated simultaneously with the others. It should be noticed that the sample sizes in our applications were relatively large ( $n = 368$ ). In small samples, it may be advisable to set a grid of values for the extra parameter and choose the value that provides the best fit according to the chosen criteria.

Applications to data on intake of several nutrients illustrated that the BCS distributions are useful in practice. The data correspond to the first 24-hour dietary recall interview for the individuals in the sample. It is part of our current research to develop Box-Cox symmetric models with random and mixed effects to model nutrients intake data taken from repeated 24-hour recalls.

As a final remark, we recall that a comprehensive study on the right tail heaviness of the Box-Cox symmetric distributions was presented. For future possible investigation, it might be interesting to search for skewness-kurtosis boundaries allowing the existence of BCS distributions, as in Jondeau and Rockinger (2003).

## Acknowledgments

We thank José Eduardo Corrente for providing the data used in this study, and Eliane C. Pinheiro for helpful discussions. We acknowledge the financial support of FAPESP, CAPES and CNPq (Brazil). We are grateful to the associate editor and two anonymous referees for constructive comments and suggestions.

## Appendix

In this appendix we give the first and second derivatives of the log-likelihood function with respect to the parameters. Let  $z = h(y; \mu, \sigma, \lambda)$ , where  $h(y; \mu, \sigma, \lambda)$  is given in (1),  $\omega = -2r'(z^2)/r(z^2)$ , and  $\xi = r((\sigma\lambda)^{-2})/R((\sigma|\lambda|)^{-1})$ . We have

$$\frac{\partial z}{\partial \mu} = -\frac{1}{\mu\sigma} \left(\frac{y}{\mu}\right)^\lambda \xrightarrow{\lambda \rightarrow 0} -\frac{1}{\mu\sigma},$$

$$\frac{\partial z}{\partial \lambda} = \frac{1}{\sigma\lambda^2} \left\{ 1 + \left(\frac{y}{\mu}\right)^\lambda \left[ -1 + \lambda \log\left(\frac{y}{\mu}\right) \right] \right\} \xrightarrow{\lambda \rightarrow 0} \frac{1}{2\sigma} \left[ \log\left(\frac{y}{\mu}\right) \right]^2,$$

$$\frac{\partial^2 z}{\partial \mu^2} = \frac{(\lambda + 1)}{\mu^2\sigma} \left(\frac{y}{\mu}\right)^\lambda \xrightarrow{\lambda \rightarrow 0} \frac{1}{\mu^2\sigma},$$

$$\frac{\partial^2 z}{\partial \lambda^2} = \frac{1}{\sigma\lambda^3} \left\{ -2 + \left(\frac{y}{\mu}\right)^\lambda \left[ 2 - 2\lambda \log\left(\frac{y}{\mu}\right) + \lambda^2 \left( \log\left(\frac{y}{\mu}\right) \right)^2 \right] \right\} \xrightarrow{\lambda \rightarrow 0} \frac{1}{3\sigma} \left[ \log\left(\frac{y}{\mu}\right) \right]^3,$$

$$\frac{\partial^2 z}{\partial \mu \partial \lambda} = -\frac{1}{\mu \sigma} \left(\frac{y}{\mu}\right)^\lambda \log\left(\frac{y}{\mu}\right) \xrightarrow{\lambda \rightarrow 0} -\frac{1}{\mu \sigma} \log\left(\frac{y}{\mu}\right).$$

Let  $\ell$  denote the log-likelihood for a single observation  $y$ . We have

$$\ell = (\lambda - 1) \log y - \lambda \log \mu - \log \sigma + \log r(z^2) - \log R\left(\frac{1}{\sigma|\lambda|}\right),$$

if  $\lambda \neq 0$ ; the last term in  $\ell$  is zero if  $\lambda = 0$ . The first derivatives of  $\ell$  are given by

$$\begin{aligned} \frac{\partial \ell}{\partial \mu} &= -\frac{\lambda}{\mu} - \omega z \frac{\partial z}{\partial \mu}, \\ \frac{\partial \ell}{\partial \sigma} &= \begin{cases} -\frac{1}{\sigma} + \frac{\omega z^2}{\sigma} + \frac{\xi}{\sigma^2 |\lambda|}, & \text{if } \lambda \neq 0, \\ -\frac{1}{\sigma} + \frac{\omega z^2}{\sigma}, & \text{if } \lambda = 0, \end{cases} \\ \frac{\partial \ell}{\partial \lambda} &= \log\left(\frac{y}{\mu}\right) - \omega z \frac{\partial z}{\partial \lambda} + \text{sign}(\lambda) \frac{\xi}{\sigma \lambda^2}. \end{aligned}$$

The second derivatives of  $\ell$  are given by

$$\begin{aligned} \frac{\partial^2 \ell}{\partial \mu^2} &= \frac{\lambda}{\mu^2} - \left(z \frac{d\omega}{dz} + \omega\right) \left(\frac{\partial z}{\partial \mu}\right)^2 - \omega z \frac{\partial^2 z}{\partial \mu^2}, \\ \frac{\partial^2 \ell}{\partial \sigma^2} &= \begin{cases} \frac{1}{\sigma^2} - \frac{z^3}{\sigma^2} \frac{d\omega}{dz} - \frac{3\omega z^2}{\sigma^2} + \frac{1}{\sigma^2 |\lambda|} \frac{\partial \xi}{\partial \sigma} - \frac{2\xi}{\sigma^3 |\lambda|}, & \text{if } \lambda \neq 0, \\ \frac{1}{\sigma^2} - \frac{z^3}{\sigma^2} \frac{d\omega}{dz} - \frac{3\omega z^2}{\sigma^2}, & \text{if } \lambda = 0, \end{cases} \\ \frac{\partial^2 \ell}{\partial \lambda^2} &= -\left(z \frac{d\omega}{dz} + \omega\right) \left(\frac{\partial z}{\partial \lambda}\right)^2 - \omega z \frac{\partial^2 z}{\partial \lambda^2} + \text{sign}(\lambda) \left(\frac{1}{\sigma \lambda^2} \frac{\partial \xi}{\partial \lambda} - \frac{2\xi}{\sigma \lambda^3}\right), \\ \frac{\partial^2 \ell}{\partial \mu \partial \sigma} &= \frac{z}{\sigma} \frac{\partial z}{\partial \mu} \left(z \frac{d\omega}{dz} + 2\omega\right), \\ \frac{\partial^2 \ell}{\partial \mu \partial \lambda} &= -\frac{1}{\mu} - \left(z \frac{d\omega}{dz} + \omega\right) \frac{\partial z}{\partial \mu} \frac{\partial z}{\partial \lambda} - \omega z \frac{\partial^2 z}{\partial \mu \partial \lambda}, \\ \frac{\partial^2 \ell}{\partial \sigma \partial \lambda} &= \frac{z}{\sigma} \frac{\partial z}{\partial \lambda} \left(z \frac{d\omega}{dz} + 2\omega\right) + \frac{1}{\sigma^2 |\lambda|} \frac{\partial \xi}{\partial \lambda} - \text{sign}(\lambda) \frac{\xi}{\sigma^2 \lambda^2}. \end{aligned}$$

The first and second derivatives of  $\ell$  are obtained after plugging the derivatives of  $z$  given above.

Note that the first derivatives of  $\ell$  depend on the weighting function  $\omega$  ( $\omega$  is given in Table 3 for some distributions). Consequently,  $d\omega/dz$  appears in all the second derivatives of  $\ell$ . Note that  $\partial \ell / \partial \sigma$  and  $\partial \ell / \partial \lambda$  involve  $\xi$ , which in turn depends on the particular distribution in the BCS class and the truncation set. The first derivatives of  $\xi$  appear in  $\partial^2 \ell / \partial \sigma^2$ ,  $\partial^2 \ell / \partial \lambda^2$

and  $\partial^2 \ell / \partial \sigma \partial \lambda$ . The stability of the terms that involve  $\xi$  and its first derivatives around  $\lambda = 0$  may vary according to different distributions. For instance, they may be unstable for the Box-Cox t distribution with small degrees of freedom parameter. Yet, a simulation study of the type I error probability of the likelihood ratio test of  $H_0 : \lambda = 0$  in the Box-Cox t model for different values of the degrees of freedom parameter performed well; see Section 4.

## References

- Azzalini, A. (2005). The skew-normal distribution and related multivariate families, *Scandinavian Journal of Statistics* **32**(2): 159–188.
- Box, G. E. P. and Cox, D. R. (1964). An analysis of transformations, *Journal of the Royal Statistical Society. Series B* **26**(2): 211–252.
- Cole, T. and Green, P. J. (1992). Smoothing reference centile curves: the LMS method and penalized likelihood, *Statistics in Medicine* **11**(10): 1305–1319.
- Cordeiro, G. M. and Andrade, M. G. (2011). Transformed symmetric models, *Statistical Modelling* **11**(4): 371–388.
- de Haan, L. (1970). *On Regular Variation and Its Application to the Weak Convergence of Sample Extremes*, Mathematical Centre Tracts 32, Amsterdam: Mathematics Centre.
- Dunn, P. K. and Smyth, G. K. (1996). Randomized quantile residuals, *Journal of Computational and Graphical Statistics* **5**(3): 236–244.
- Fang, K. T., Kotz, S. and NG, K. W. (1990). *Symmetric Multivariate and Related Distributions*, Chapman and Hall, London.
- Hubert, M. and Vandervieren, E. (2008). An adjusted boxplot for skewed distributions, *Computational Statistics & Data Analysis* **52**(12): 5186–5201.
- Jondeau, E. and Rockinger, M. (2003). Conditional volatility, skewness, and kurtosis: existence, persistence, and comovements, *Journal of Economic Dynamics & Control* **27**: 1699–1737.
- Kelker, D. (1970). Distribution theory of spherical distributions and a location-scale parameter generalization, *Sankhya A* **32**(4): 419–430.
- Luceño, A. (2005). Fitting the generalized Pareto distribution to data using maximum goodness-of-fit estimators, *Computational Statistics & Data Analysis* **51**(2): 904–917.
- Poirier, D. J. (1978). The use of box-cox transformation in limited dependent variable models, *Journal of the American Statistical Association* **73**(362): 284–287.



- Resnick, S. I. (2007). *Heavy-Tail Phenomena Probabilistic and Statistical Modeling*, Springer.
- Rigby, R. A. and Stasinopoulos, D. M. (2004). Smooth centile curves for skew and kurtotic data modelled using the BoxCox power exponential distribution, *Statistics in Medicine* **23**(19): 3053-3076.
- Rigby, R. A. and Stasinopoulos, D. M. (2005). Generalized additive models for location, scale and shape, *Journal of the Royal Statistical Society: Series C (Applied Statistics)* **54**(3): 507–554.
- Rigby, R. A. and Stasinopoulos, D. M. (2006). Using the Box-Cox t distribution in GAMLSS to model skewness and kurtosis, *Statistical Modelling* **6**(3): 209–229.
- Rigby, R. A., Stasinopoulos, D. M., Heller, G. and Voudouris, V. (2014). *The Distribution Toolbox of GAMLSS*, London. <http://www.gamlss.org/wp-content/uploads/2014/10/distributions.pdf>.
- Rogers, W. H. and Tukey, J. W. (1972). Understanding some long-tailed symmetrical distributions, *Statistica Neerlandica* **26**(3): 211–226.
- Stasinopoulos, D. M., Rigby, R. A. and Akantziliotou, C. (2008). *Instructions on how to use the GAMLSS package in R*, London. <http://www.gamlss.org>.
- Vanegas, L. H. and Paula, G. A. (2015). A semiparametric approach for joint modeling of median and skewness, *Test* **24**(1): 110–135.
- Vanegas, L. H. and Paula, G. A. (2016). Log-symmetric distributions: statistical properties and parameter estimation, *Brazilian Journal of Probability and Statistics* **30**: 196–220.
- Voudouris, V., Gilchrist, R., Rigby, R. A., Sedgwick, J. and Stasinopoulos, D. M. (2012). Modelling skewness and kurtosis with the BCPE density in GAMLSS, *Journal of Applied Statistics* **39**(6): 1279–1293.
- Yang, Z. (2006). A modified family of power transformations, *Economics Letters* **92**(1): 14–19.
- Yang, Z. L. (1996). Some asymptotic results on Box-Cox transformation methodology, *Communications in Statistics, Theory and Methods* **25**(2): 403–415.
- Yeo, I. K. and Johnson, R. A. (2000). A new family of power transformation to improve normality or symmetry, *Biometrika* **87**(4): 954–959.