

Second Noether theorem for quasi-Noether systems

V. Rosenhaus and R. Shankar

Department of Mathematics and Statistics, California State University, Chico, USA

Abstract

Quasi-Noether differential systems are more general than variational systems and are quite common in mathematical physics. They include practically all differential systems of interest, at least those that have conservation laws. In this paper, we discuss quasi-Noether systems that possess infinite-dimensional (infinite) symmetries involving arbitrary functions of independent variables. For quasi-Noether systems admitting infinite symmetries with arbitrary functions of all independent variables, we state and prove an extension of the Second Noether Theorem. In addition, we prove that infinite sets of conservation laws involving arbitrary functions of all independent variables are trivial and that the associated differential system is under-determined. We discuss infinite symmetries and infinite conservation laws of two important examples of non-variational quasi-Noether systems: the incompressible Euler equations and the Navier-Stokes equations in vorticity formulation, and we show that the infinite sets of conservation laws involving arbitrary functions of all independent variables are trivial. We also analyze infinite symmetries involving arbitrary functions of not all independent variables, prove that the fluxes of conservation laws in these cases are total divergences on solutions, and demonstrate examples of this situation.

Keywords: second Noether theorem, infinite conservation laws, Navier-Stokes equations, vorticity equations, quasi-Noether systems, infinite symmetries

1 Introduction

Infinite-dimensional symmetry algebras parametrized by arbitrary functions and their relations to conservation laws have been studied considerably less extensively than finite-dimensional Lie symmetry groups and their corresponding conservation laws. For variational problems, according to the classical Noether result [15] (see also [16]), infinite variational symmetries with arbitrary functions of *all independent variables* lead to certain identity relationships between the equations of the original system of partial differential equations (differential system) and their derivatives. The most well known examples of such under-determined systems are gauge (invariant) theories.

The problem of a correspondence between variational infinite symmetries with arbitrary functions of *not all independent variables* and conservation laws was studied in [21]. It was demonstrated that these infinite variational symmetries lead to a finite number of essential

local conservation laws (those that give rise to non-vanishing conserved quantities) and that each essential conservation law is determined by a specific form of boundary condition. Using this method, conserved densities were found for a number of equations including the Zabolotskaya-Khokhlov equation [22] and the Kadomtsev-Petviashvili equation [23]. In [6], Noether's approach was extended to include infinite variational symmetries involving functions of independent variables with some constraints.

A cohomological consideration of Noether's theorems in terms of the variational bicomplex was given in [1, 14].

It was shown in [24] that the situation with infinite symmetry algebras parametrized by arbitrary functions of *dependent variables* is radically different, leading to an infinite set of local conserved densities. Two known examples of this situation [25] are equations of Liouville type, see e.g. [33], that can be integrated by the Darboux method (e.g. [10]), and hydrodynamic-type equations, [28].

However, Noether's approach is directly applicable only to the equations of a variational problem with a well-defined Lagrangian function. A main tool for our study is a method based on the Noether operator identity that establishes correspondences between symmetries and local conservation laws for differential systems without well-defined Lagrangian functions [19, 20]. *Quasi-Noether* systems are differential systems for which it is possible to establish a correspondence between symmetries and conservation laws based on the Noether operator identity, see [26]). Quasi-Noether differential systems are rather general and include all systems possessing conservation laws.

In the present paper, we apply an approach based on the Noether operator identity to quasi-Noether systems possessing infinite-dimensional symmetry algebras involving arbitrary functions of all independent variables. We show that, similarly to the Second Noether Theorem of variational systems, the existence of infinite symmetries with arbitrary functions of all independent variables results in differential identities relating equations of the original quasi-Noether system to their derivatives (showing that the original system is under-determined). We consider two important examples of such quasi-Noether systems, the incompressible Euler equations and the Navier-Stokes equations for vorticities, which are known to admit infinite sets of conservation laws. We show that these conservation laws are trivial.

We also analyze infinite symmetries involving arbitrary functions of not all independent variables. For these cases, we prove that the fluxes of the corresponding conservation laws are total divergences on solutions and demonstrate examples of this situation.

Section 2 discusses symmetries and conservation laws for variational systems, as well as the Noether identity and the two Noether theorems. Section 3 introduces quasi-Noether systems and discusses the correspondence between symmetries and conservation laws for such systems. We formulate and prove extensions of the two Noether theorems for quasi-Noether systems. Section 4 deals with the case when conservation laws of a general differential system involve arbitrary functions of all independent variables. We show that such conservation laws are trivial and lead to differential identities in extended space. Section 5 discusses a case of infinite conservation laws that involve arbitrary functions of *not all* independent variables. We demonstrate that the conservation laws here are, in general, not trivial and that their fluxes are total divergences on solutions. In Section 6, we consider two interesting examples of

quasi-Noether systems: the incompressible Euler equations and the Navier-Stokes equations in vorticity formulation. We analyze infinite sets of conservation laws for these equations and show that they are trivial.

2 Symmetries and conservation laws. Variational systems

Let us briefly outline the approach we follow; for details, see [21, 24].

Definition 1 (Conservation laws). By a conservation law of a differential system

$$\Delta_a(x, u, u_{(1)}, u_{(2)}, \dots, u_{(l)}) = 0, \quad a = 1, \dots, n, \quad (2.1)$$

we mean a divergence expression

$$D_i K^i(x, u, u_{(1)}, u_{(2)}, \dots) \doteq 0, \quad (2.2)$$

that vanishes on all solutions of the original system; we denote this type of equality by (\doteq) . Here, $x = (x^1, x^2, \dots, x^p)$ and $u = (u^1, u^2, \dots, u^m)$ are the tuples of independent and dependent variables, respectively; $u_{(r)}$ is the tuple of r th-order derivatives of u , and $r = 1, 2, \dots, l$. Each K^i is a differential function ([16]), i.e. a smooth function of x , u , and a finite number of derivatives of u . Each Δ_a , $a = 1, 2, \dots, n$ is a differential function of x , u , and all partial derivatives of each u^v , ($v = 1, \dots, m$) with respect to the x^i ($i = 1, \dots, p$) up to the l th order ([16]). Smooth functions $u^q = u^q(x)$, $q = 1, \dots, m$ are defined on a non-empty open subset D of p -dimensional space-time \mathbb{R}^p . Summation over repeated indices is assumed. \square

We use the following notations for the differential system:

$(\Delta_1, \Delta_2, \dots, \Delta_n) \equiv \Delta \equiv \Delta[u] \equiv \Delta(x, u, u_{(1)}, u_{(2)}, \dots, u_{(l)})$. In the multi-index notation of [16], $x_J = (x_{j_1}, x_{j_2}, \dots, x_{j_k})$, and u_J^v are partial derivatives, where $J = (j_1, j_2, \dots, j_k)$.

Definition 2 (Trivial conservation laws). A conservation law $D_i P^i \doteq 0$ is *trivial* [16] if a linear combination of *two kinds of triviality* is taking place: 1. The p -tuple P vanishes on the solutions of the original system: $P^i \doteq 0$. 2. The divergence identity is satisfied for any point $[u] = (x, u_{(p)})$ in the extended jet space (e.g. $\partial_t \operatorname{div} u + \operatorname{div}(-\partial_t u) = 0$). Two conservation laws K and \tilde{K} are equivalent if they differ by a trivial conservation law. \square

Remark 1. If a conservation law (2.2) exists, the following equality holds:

$$D_i J^i = \xi^a \Delta_a, \quad (2.3)$$

where each ξ^a is a *characteristic*, and $J^i \doteq K^i$. If $\xi^a = 0$, the conservation law (2.3) is trivial; if $\xi^a \neq 0$, it is nontrivial. \square

Definition 3 (Euler operator).

$$E_{u^a} = \frac{\partial}{\partial u^a} - \sum_i D_i \frac{\partial}{\partial u_i^a} + \sum_{i \leq j} D_i D_j \frac{\partial}{\partial u_{ij}^a} + \dots \quad (2.4)$$

is the Euler (Euler–Lagrange) operator (variational derivative). In the notation of [16], we could give the Euler operator the following form:

$$E_{u^a} = (-D)_J \frac{\partial}{\partial u_J^a}, \quad 1 \leq a \leq n, \quad (2.5)$$

the sum extending over all unordered multi-indices $J = (j_1, j_2, \dots, j_k)$ for $1 \leq k \leq p$. For $k = 0$ we set $u_J^a = u^a$ and $D_J = 1$. Here and in what follows, we make use of the multi-index notation [16] for total derivatives: $D_J = D_1^{J_1} \dots D_p^{J_p}$, where $J = (J_1, \dots, J_p)$ is a multi-index, $D_i = \partial_{x_i} + u_i^q \partial_{u^q} + \dots$ is a total derivative, and each J_i is non-negative. The adjoint operator to D_J is $(-D)_J = (-D_1)^{J_1} \dots (-D_p)^{J_p}$.

Remark 2. The operator E annihilates total divergences [16]. That is, $E_{u^a}(D_i J^i) \equiv 0$.

Definition 4 (Variational problems). Let

$$S = \int_D L(x, u, u_{(1)}, \dots) d^p x$$

be the action functional, where L is the Lagrangian density. The equations of motion are

$$E_{u^a}(L) \equiv \Delta_a(x, u, u_{(1)}, \dots) = 0, \quad (2.6)$$

where E is as in (2.4). □

Definition 5 (Variational symmetry). Consider a one-parameter transformation with a canonical (vertical) infinitesimal operator

$$\begin{aligned} X_\alpha &= \alpha^a \frac{\partial}{\partial u^a} + \sum_i (D_i \alpha^a) \frac{\partial}{\partial u_i^a} + \sum_{i \leq j} (D_i D_j \alpha^a) \frac{\partial}{\partial u_{ij}^a} + \dots, \\ \alpha^a &= \alpha^a(x, u, u_{(1)}, \dots). \end{aligned} \quad (2.7)$$

The variation of the functional S under the transformation with operator X_α is

$$\delta S = \int_D X_\alpha L d^p x. \quad (2.8)$$

X_α is a variational (Noether) symmetry if

$$X_\alpha L = D_i M^i, \quad (2.9)$$

where $M^i = M^i(x, u, u_{(1)}, \dots)$ are smooth functions of their arguments. □

Remark 3. We can write the symmetry operator (2.7) in the form

$$X_\alpha = (D_J \alpha^a) \partial_{u_J^a}, \quad (2.10)$$

where the sum is taken over all unordered multi-indices J . □

The Noether identity [18] (see also [8] or [19] for the version used here) relates a symmetry operator X_α to the Euler- Lagrange operator E_{u^q} .

Lemma 1 (Noether Identity). *Let $X_\alpha = (D_J \alpha^q) \partial / \partial u_J^q$ be a vertical symmetry operator with infinitesimal α . The following operator identity holds:*

$$X_\alpha = \alpha^q E_{u^q} + D_i R_\alpha^i, \quad (2.11)$$

where

$$R_\alpha^i = \alpha^q \frac{\partial}{\partial u_i^q} + \left\{ \sum_{k \geq i} (D_k \alpha^q) - \alpha^q \sum_{k \leq i} D_k \right\} \frac{\partial}{\partial u_{ik}^q} + \dots \quad (2.12)$$

□

Remark 4. The operator R_α^i can be given a general form using multi-index notation [26].

$$R_\alpha^i = \sum_J \sum_{m=1}^k [\delta^{ij_m} (-1)^{k+m} (D_{J_{m-1}} \alpha^a) D_{J-J_m}] \partial_{u_J^a}, \quad (2.13)$$

$$J_m = (j_1, j_2, \dots, j_m), \quad J - J_m = (j_{m+1}, j_{m+2}, \dots, j_k), \quad D_{J_0} = D_0 = 1.$$

□

Applying the identity (2.11) to L and using (2.9), we obtain

$$D_i(M^i - R^i L) = \alpha^a \Delta^a, \quad (2.14)$$

leading to the statement of the First Noether Theorem: any one-parameter variational symmetry transformation with infinitesimal operator X_α (2.7) gives rise to a conservation law

$$D_i(M^i - R_\alpha^i L) \doteq 0 \quad (2.15)$$

(on the solution manifold $\Delta = 0, D_i \Delta = 0, \dots$).

Theorem 1 (Noether 1). *Let $X_\alpha = (D_J \alpha^q) \partial / \partial u_J^q$ generate a finite variational symmetry group of a Lagrangian \mathcal{L} , such that $X_\alpha \mathcal{L} = D_i M_\alpha^i$ for some M_α . Then there exists the following conservation law associated to the symmetry X_α :*

$$D_i(M_\alpha^i - R_\alpha^i \mathcal{L}) = \alpha^q E_{u^q} \mathcal{L}. \quad (2.16)$$

Remark 5. Equation (2.16) is a conservation law since the right hand side is zero *on-shell*, or when $E_{u^q} \mathcal{L} = 0$ for each q . In general, we say that computations are taken *on-shell* if u solves our differential system ($\Delta = 0$). Otherwise, we say that we work *off-shell*.

Remark 6. The first Noether theorem was formulated for continuous *finite groups* of symmetry transformations. For *infinite groups* depending on arbitrary functions of all independent variables, we can formally generate the same type of continuity equations (conservation laws). However, these equations will not have the same meaning as their finite-dimensional counterparts and will lead to *differential identities* relating the equations of motion to their derivatives.

Theorem 2 (Second Noether Theorem). *Let X_{α_f} (2.7) be a variational symmetry of Lagrangian \mathcal{L} with infinitesimal $\alpha_f^q = \alpha^{qJ} D_J f$ involving an arbitrary function of all independent variables $f = f(x_1, \dots, x_n)$. Then X_{α_f} generates an identity involving the original equations and their derivatives. This off-shell identity takes the following form:*

$$(-D)_J [\alpha^{qJ} \Delta_q] = 0. \quad (2.17)$$

Remark 7. The identity (2.17) expresses the fact that the original system $\Delta = E(\mathcal{L})$ is under-determined.

We next consider differential systems without well-defined Lagrangian functions.

3 Symmetries and conservation laws of quasi-Noether systems

For a general differential system, a relationship between symmetries and conservation laws is unknown. In [19] and [20], an approach based on the Noether Identity (2.11) was suggested to relate symmetries to conservation laws for a large class of differential systems that may not have well-defined Lagrangian functions.

Consider a system (2.1)

$$\Delta_a(x, u, u_{(1)}, u_{(2)}, \dots, u_{(l)}) = 0, \quad a = 1, \dots, n, \quad (3.1)$$

of n l th order differential equations for functions u .

Applying the Noether identity (2.11) to a combination of Δ_a , we obtain

$$X_\alpha(\beta^a \Delta_a) = \alpha^v E_{u^v}(\beta^a \Delta_a) + D_i R^i(\beta^a \Delta_a), \quad a = 1, \dots, n, \quad v = 1, \dots, m, \quad i = 1, \dots, p. \quad (3.2)$$

If there exist coefficients β^a such that [20]

$$E_v(\beta^a \Delta_a) \doteq 0, \quad a = 1, \dots, n, \quad v = 1, \dots, m, \quad (3.3)$$

then each symmetry X_α of the system will lead to a local conservation law

$$D_i R^i(\beta^a \Delta_a) \doteq 0, \quad (3.4)$$

for any differential system of class (3.3). In [19], the quantity $\beta^a \Delta_a$ was referred to as an alternative Lagrangian.

Note that a condition similar to (3.3) (for a system to be quasi-Noether and possess a correspondence between symmetries and conservation laws; [19, 20]) was obtained earlier within an alternative approach based on the Lagrange identity; see [30, 29, 32, 3, 27, 11]. It was shown in [26] that, for a given symmetry transformation, both approaches give rise to the same conservation law.

Note also that the same condition (3.3) has played a key role in the later developed nonlinear self-adjointness approach [7] to a correspondence between symmetries and conservation laws. Using this condition and alternative Lagrangians (that were called "formal Lagrangians"; see also [9]), numerous applications and examples were given within this approach. Note in addition that the idea of alternative Lagrangians was suggested in [16] as Lagrangians in auxiliary variables.

Definition 6 (Quasi-Noether systems). We say that a system (2.1) Δ is *quasi-Noether* if it has an *alternative-Lagrangian* in the form of a linear combination of Δ_a that has zero variational derivative on-shell [26]. According to (3.3), $\mathcal{A} = \beta^a \Delta_a$ is an alternative-Lagrangian if

$$E_{u^q} \mathcal{A} = \Xi^{qaJ} D_J \Delta_a, \quad 1 \leq q \leq m, \quad (3.5)$$

where the β 's and Ξ 's are some functions. Clearly, $E_{u^q} \mathcal{A} = 0$ when $\Delta = 0$.

For a regular Lagrangian of a variational problem, we can take $\mathcal{A} = \mathcal{L}$, such that (3.5) reduces to $E_{u^q} \mathcal{L} = \Delta_q$.

Remark 8. Note that not all variational systems are quasi-Noether. However, those variational systems that are not quasi-Noether are unlikely to be of physical interest, since such systems necessarily lack conservation laws, and, hence, lack continuous symmetries. For example, the sine-Gordon-type Lagrangian

$$\mathcal{L} = -u_t u_x / 2 + f(t, x) \cos u$$

leads to the following Euler-Lagrange equation

$$E(\mathcal{L}) = u_{tx} - f(t, x) \sin u = 0.$$

This equation admits no local conservation laws if $f(t, x)$ does not satisfy the following linear equation

$$\partial_t(a(t)f) + \partial_x(b(x)f) = 0$$

for any choice of $a(t)$ and $b(x)$ not both zero. For example, $f(t, x) = \exp(-t^2 x^2) / [1 + (t + x)^2]$.

Remark 9. Quasi-Noether systems are quite common and include most known variational systems and all differential systems in the form $D_i N_a^i = 0$, such as the Euler and the Navier-Stokes equations; see [19] for other examples. Most, if not all, interesting differential systems are quasi-Noether.

Theorem 3 (Noether's First theorem for quasi-Noether systems). *Suppose that a quasi-Noether system (2.1) Δ possesses a finite continuous symmetry X_α . Then there exists conservation law (3.8) of the system Δ associated to the symmetry X_α .*

Proof. Applying X_α to an alternative-Lagrangian $\mathcal{A} = \beta^a \Delta_a$ and using (3.2), we obtain:

$$\begin{aligned} X_\alpha \mathcal{A} &= \alpha^q E_{u^q} \mathcal{A} + D_i R_\alpha^i \mathcal{A} \\ &= \alpha^q \Xi^{qaJ} D_J \Delta_a + D_i R_\alpha^i \mathcal{A}. \end{aligned} \quad (3.6)$$

Next, we rewrite the left hand side using the fact that X_α is a symmetry of Δ , or that $X_\alpha \Delta = 0$ on-shell. Since $X_\alpha \Delta_b = \tau_\alpha^{baJ} D_J \Delta_a$ for some coefficients τ_α , we have:

$$\begin{aligned} X_\alpha \mathcal{A} &= X_\alpha (\beta^a \Delta_a) \\ &= \beta^a X_\alpha \Delta_a + (X_\alpha \beta^a) \Delta_a \\ &= (\beta^b \tau_\alpha^{baJ} D_J + X_\alpha \beta^a) \Delta_a \\ &=: \gamma_\alpha^{aJ} D_J \Delta_a. \end{aligned} \quad (3.7)$$

Combining (3.6) with (3.7) yields the following:

$$\alpha^q \Xi^{qaJ} D_J \Delta_a + D_i R_\alpha^i \mathcal{A} = \gamma_\alpha^{aJ} D_J \Delta_a.$$

Rearranging, we obtain a conservation law for each symmetry α , ([19]):

$$D_i R_\alpha^i \mathcal{A} = (\gamma_\alpha^{aJ} - \alpha^q \Xi^{qaJ}) D_J \Delta_a. \quad (3.8)$$

Indeed, the left hand side is a total divergence, and the right hand side is zero on-shell. \square

Remark 10. The *characteristic* of the conservation law (3.8) corresponding to the symmetry X_α of a quasi-Noether system Δ is as follows:

$$\xi_\alpha^a = (-D)_J (\gamma_\alpha^{aJ} - \alpha^q \Xi^{qaJ}). \quad (3.9)$$

Example 1. Just to demonstrate an application of Theorem 3, consider the Korteweg de-Vries (KdV) equation:

$$\Delta = u_t + uu_x + u_{xxx} = 0. \quad (3.10)$$

Since this equation is, itself, a conservation law, the KdV equation is quasi-Noether: $\mathcal{A} = \Delta$, $E(\Delta) = 0$. The KdV equation (3.10) is known to admit a scaling symmetry (see e.g. [16]):

$$X_\alpha = (2u + xu_x + 3tu_t) \partial_u. \quad (3.11)$$

The form of the conservation law resulting from $X_\alpha \mathcal{A}$ is immediate after using the commutator identity $[X_\alpha, D_t] = [X_\alpha, D_x] = 0$:

$$X_\alpha \mathcal{A} = D_t \alpha + D_x(\alpha u) + D_x^3 \alpha.$$

Expanding this expression, we can find that its characteristic is $\xi = 1$. We obtain the following conservation law:

$$\begin{aligned} D_t[2u + 3tu_t + xu_x] + D_x[u(2u + 3tu_t + xu_x)] + D_x^3[2u + 3tu_t + xu_x] \\ = \Delta + D_t(3t\Delta) + D_x(x\Delta). \end{aligned} \quad (3.12)$$

See [19] for more examples and details of applying Theorem 3.

We now formulate Noether's second theorem for quasi-Noether systems.

Theorem 4 (Noether's Second theorem for quasi-Noether systems). *Suppose that X_{α_f} is an infinite symmetry of quasi-Noether system (2.1) Δ , where the infinitesimals $\alpha_f^q = \alpha^{qJ} D_J f$ depend on an arbitrary function $f(x_1, \dots, x_p)$ of all independent variables. Then X_{α_f} generates identity (3.16) involving the equations of the system Δ and their derivatives.*

Proof. For infinite symmetry group operator X_{α_f} , we can apply the same procedure as in Theorem 3 to formally obtain analogous conservation laws (3.8) with characteristic (3.9); here, however, the infinite symmetry X_{α_f} leads to an infinite set of conservation laws involving the arbitrary function f and its derivatives:

$$D_i J_{\alpha_f}^i = (\xi^{aJ} D_J f) \Delta_a \quad (3.13)$$

for some functions ξ^{aJ} . We follow Noether's approach and show that these conservation laws have a different meaning; they lead to differential identities involving the equations of the system Δ and their derivatives and, hence, express the under-determinacy of the system Δ . Our computations are performed *off-shell*. Integrating (3.13) over an arbitrary bounded connected subset S of \mathbb{R}^p , we obtain an integral identity:

$$\int_S D_i J_{\alpha_f}^i d^p x = \int_S (\xi^{aJ} D_J f) \Delta_a d^p x.$$

We integrate the right hand side by parts using the identity (4.9) with (4.10) and collect the divergences to the left hand side:

$$\int_S D_i \left(J_{\alpha_f}^i - \Phi_J^i[f, \xi^{aJ} \Delta_a] \right) d^p x = \int_S f(-D)_J [\xi^{aJ} \Delta_a] d^p x.$$

We observe that, inside the divergence operator, there are two vectors: $J_{\alpha_f}^i$ and $\Phi_J^i[f, \xi^{aJ} \Delta_a]$. The first one is:

$$J_{\alpha_f}^i = R_{\alpha_f}^i \mathcal{A} - \Phi_J^i[\Delta_a, \gamma_{\alpha_f}^{aJ} - \alpha_f^q \Xi^{qaJ}], \quad (3.14)$$

where $\mathcal{A} = \beta^a \Delta_a$, $R_{\alpha_f}^i$ is expressed through α_f and its derivatives according to (2.12), and Φ^i , expressed in (4.10), comes from integrating the right hand side (RHS) of (3.8) by parts to obtain (3.13). Therefore, the vector $J_{\alpha_f}^i$ is a linear combination of the arbitrary function $f(x_1, \dots, x_p)$ and its derivatives. The same conclusion can be made with respect to the second vector $\Phi_J^i[f, \xi^{aJ} \Delta_a]$. We choose the arbitrary function of all independent variables $f(x)$ so that it vanishes on ∂S together with all its derivatives. Then an application of Gauss's theorem to the left hand side reduces it to vanishing surface terms, and, therefore, we obtain:

$$0 = \int_S f(-D)_J [\xi^{aJ} \Delta_a] d^p x. \quad (3.15)$$

The equation (3.15) holds for arbitrary functions f that vanish on ∂S together with their derivatives, and, therefore, it is an identity. Since the bounded subset S is arbitrary, the integrand must vanish identically everywhere in the extended space:

$$(-D)_J [\xi^{aJ} \Delta_a] = 0. \quad (3.16)$$

The identity (3.16) is a functional relationship between the original equations of the system Δ and their derivatives. Thus, not all equations of Δ are independent, and the system Δ is *under-determined*. The existence of an infinite symmetry X_{α_f} and the infinite conservation laws (3.13) is a consequence of the additional degree(s) of freedom related to the description of an under-determined system Δ . \square

Remark 11. We express the identity (3.16) in terms of the symmetry coefficients $\alpha^q = \alpha^{qJ} D_J f$. In (3.7), we suppose that $X_{\alpha} \beta^a = \bar{\beta}^{aJ} D_J f$, and $X_{\alpha} \Delta_b = \bar{\tau}^{baJK} D_K f D_J \Delta_a$. Then the characteristic (3.9) in (3.13) is as follows:

$$\begin{aligned} \xi^a &= (-D)_J \left[\left(\bar{\beta}^{aK} \delta^{J0} + \beta^b \bar{\tau}^{baJK} - \alpha^{qK} \Xi^{qaJ} \right) D_K f \right] \\ &= (-D)_J \left[\bar{\beta}^{a0} \delta^{J0} + \beta^b \bar{\tau}^{baJ0} - \alpha^{q0} \Xi^{qaJ} \right] f + \dots \\ &=: \xi^{a0} f + \dots \end{aligned} \quad (3.17)$$

Remark 12. In terms of these coefficients, we can rewrite identity (3.16) as follows:

$$\xi^a \Delta_a - D_i(\xi^{ai} \Delta_a) + \sum_{i \leq j} D_i D_j(\xi^{aij} \Delta_a) - \sum_{i \leq j \leq k} D_i D_j D_k(\xi^{aijk} \Delta_a) + \dots = 0. \quad (3.18)$$

Remark 13. Alternative Lagrangians do not correspond to well-defined variational problems, and true variational formulas are not the same as the ones obtained with an alternative Lagrangian of quasi-Noether systems. For example, we can see that (2.17) is different from (3.16) since the functions ξ^{aJ} are, clearly, different from the functions α^{qJ} ; see (3.9).

4 Arbitrary functions, infinite conservation laws, and differential identities

4.1 Overview

In this section, we discuss a general case of conservation laws that contain arbitrary functions of *all* independent variables. We show that these conservation laws are necessarily trivial. To illustrate our conclusion, we consider the special case of a first order differential equation $\Delta(t, x, u, u_t, u_x) = 0$ for a single function u of two variables t and x . Suppose that Δ admits the following infinite set of conservation laws:

$$D_t M_f^t + D_x M_f^x = (f \xi^0 + \partial_t f \xi^{10} + \partial_x f \xi^{01}) \Delta, \quad (4.1)$$

where $f(t, x)$ is an arbitrary function of *all independent variables*, ξ^i are some functions of (t, x, u) , and the coefficients M_f^i involve f and its derivatives. Rewriting the right hand side, we obtain

$$D_t M_f^t + D_x M_f^x = \xi^0 \Delta f - D_t(\xi^{10} \Delta) f + D_t(\xi^{10} \Delta f) - D_x(\xi^{01} \Delta) f + D_x(\xi^{01} \Delta f). \quad (4.2)$$

Equivalently,

$$D_t(M_f^t - f \xi^{10} \Delta) + D_x(M_f^x - f \xi^{01} \Delta) = f[\xi^0 \Delta - D_t(\xi^{10} \Delta) - D_x(\xi^{01} \Delta)]. \quad (4.3)$$

We integrate the left hand side over the whole space t, x . Similarly to derivation of (3.16), we can choose the arbitrary function of all independent variables $f(t, x)$ to vanish at the boundaries together with all its derivatives. Then, using Gauss' theorem and the fact that the surface terms vanish, we obtain

$$\int_S f[(\xi^0 \Delta - D_t(\xi^{10} \Delta) - D_x(\xi^{01} \Delta))] dx dt = 0. \quad (4.4)$$

Equation (4.4) holds for an arbitrary function f that vanishes on ∂S together with its derivatives, so the integrand must vanish identically on S :

$$\xi^0 \Delta - D_t(\xi^{10} \Delta) - D_x(\xi^{01} \Delta) = 0. \quad (4.5)$$

The identity (4.5) is a functional relationship between Δ and its derivatives that holds for all smooth functions $u(t, x)$. Thus, the equation Δ for $u = u(t, x)$ is *under-determined*.

The final step to prove the triviality of (4.1) ((4.3)) is to substitute the identity (4.5) into (4.3). We rewrite (4.3):

$$D_t(M_f^t - f\xi^{10}\Delta) + D_x(M_f^x - f\xi^{01}\Delta) = 0. \quad (4.6)$$

Equation (4.6) is an identity that holds for all functions $u = u(t, x)$, not only solutions of $\Delta = 0$. Since the coefficient of Δ on the right hand side is zero, *this conservation law is trivial*.

4.2 Arbitrary functions of all independent variables, and differential identities

For general differential systems, we have the following result.

Theorem 5. *Consider a differential system (2.1) $\Delta \equiv (\Delta_1, \Delta_2, \dots, \Delta_n)$ for functions $u^q, 1 \leq q \leq m$. Suppose that Δ possesses an infinite conservation law*

$$D_i M_f^i = \xi_f^a \Delta_a, \quad (4.7)$$

where the fluxes M_f^i and characteristics ξ_f^a are given by

$$\begin{aligned} M_f^i &= M^{iJ} D_J f, \quad 1 \leq i \leq p, \\ \xi_f^a &= \xi^{aJ} D_J f, \quad 1 \leq a \leq n, \end{aligned} \quad (4.8)$$

and $f(x) = f(x_1, \dots, x_p)$ is an arbitrary function of all independent variables.

Then there exists differential identity (4.12) involving the equations of the system Δ and their derivatives (i.e. Δ is an under-determined system), and the infinite conservation law (4.7) is trivial.

We present an algebraic proof of this theorem using some tools implemented in [16]. We recall the Euler operator E_f in Definition 3 and the following integration by parts identity.

Lemma 2. *Let f and g be functions, $J = (J_1, J_2, \dots, J_p)$ be a multi-index, and $D_J = D_1^{J_1} \dots D_p^{J_p}$ be a total derivative. Then the following identity holds:*

$$g D_J f = f (-D)_J g + D_i \Phi_J^i[f, g], \quad (4.9)$$

where $(-D)_J = (-D_1)^{J_1} \dots (-D_p)^{J_p}$ is the adjoint operator to D_J , and

$$\Phi_J^i[f, g] = \sum_{j=0}^{J_i-1} \left(D_i^{J_i-1-j} D_{i+1}^{J_{i+1}} \dots D_n^{J_n} f \right) \left((-D_i)^j (-D_1)^{J_1} \dots (-D_{i-1})^{J_{i-1}} g \right), \quad (4.10)$$

for each $1 \leq i \leq p$.

Example 2.

$$g D_x^2 D_y f = (-1)^3 f D_x^2 D_y g + D_x (D_x D_y f g - D_y f D_x g) + D_y (f D_x^2 g).$$

The identity (4.9) helps us directly apply the Euler operator (2.5) instead of integrating over space, as in Noether's approach.

Proof of Theorem 5. Using the integration by parts identity (4.9) in the RHS of (4.7), we obtain:

$$D_i M_f^i = D_J f \xi^{aJ} \Delta_a = f(-D)_J [\xi^{aJ} \Delta_a] + D_i \Phi_J^i[f, \xi^{aJ} \Delta_a]. \quad (4.11)$$

Applying now the Euler operator E_f to (4.11), we obtain:

$$0 = (-D)_J [\xi^{aJ} \Delta_a]. \quad (4.12)$$

Equation (4.12) is a differential identity holding for all functions u . It relates Δ to its derivatives and shows that the system Δ is under-determined.

Substituting (4.12) into (4.11) gives an equivalent representation of (4.7):

$$D_i M_f^i = D_i \Phi_J^i[f, \xi^{aJ} \Delta_a]. \quad (4.13)$$

The right hand side is a total divergence. Thus, the characteristic of this conservation law is zero, and the infinite set of conservation laws (4.7) is trivial. \square

Remark 14. Any differential system that has conservation laws is necessarily quasi-Noether, according to Definition 3.3. Therefore, Theorem 5 is formulated for quasi-Noether systems.

Example 3. An example of Theorem 5 is a known fact that (infinitesimal) local gauge transformations lead to trivial conservation laws (continuity equations). Consider a scalar field φ interacting with an electromagnetic field (scalar electrodynamics with zero potential) with the Lagrangian:

$$\mathcal{L} = \mathcal{D}_\mu \bar{\varphi} \mathcal{D}^\mu \varphi - \frac{1}{4} F_{\mu\nu} F^{\mu\nu}, \quad (4.14)$$

where $\bar{\varphi}$ is the complex conjugate of φ , $\mathcal{D}_\mu = \partial_\mu - iA_\mu$, and $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, $\mu, \nu = 1, \dots, 4$.

The Euler-Lagrange equations for (4.14) are as follows:

$$\begin{aligned} E_{A^\mu} \mathcal{L} &= i(\bar{\varphi} \partial_\mu \varphi - \varphi \partial_\mu \bar{\varphi}) + 2A_\mu |\varphi|^2 - \partial^\nu F_{\mu\nu} = 0, \\ E_{\bar{\varphi}} \mathcal{L} &= -\partial_\mu \partial^\mu \varphi + 2iA^\mu \partial_\mu \varphi + i\varphi \partial_\mu A^\mu + 2A_\mu A^\mu \varphi = 0, \\ E_\varphi \mathcal{L} &= \overline{E_{\bar{\varphi}} \mathcal{L}}. \end{aligned} \quad (4.15)$$

Lagrangian (4.14) is invariant with respect to the following infinite symmetry (local gauge) transformation:

$$X_\theta = i\theta \varphi \partial_\varphi - i\theta \bar{\varphi} \partial_{\bar{\varphi}} + \partial^\mu \theta \partial_{A^\mu}, \quad (4.16)$$

where $\theta = \theta(x)$ is an arbitrary function of all independent variables x_μ , $X_\theta \mathcal{L} \equiv 0$.

Applying the Noether operator identity (2.11) to the Lagrangian (4.14), we obtain the following infinite set of conservation laws (2.15):

$$\begin{aligned} \partial^\mu M_\mu &\doteq 0, \\ M_\mu &= \theta [i(\bar{\varphi} \partial_\mu \varphi - \varphi \partial_\mu \bar{\varphi}) + 2A_\mu |\varphi|^2] + \partial^\nu \theta F_{\mu\nu}. \end{aligned} \quad (4.17)$$

Using equations (4.15), we can rewrite the fluxes M_μ in the following form:

$$M_\mu = \theta E_{A^\mu} \mathcal{L} + \partial^\nu (\theta F_{\mu\nu}).$$

The first term vanishes on the solutions of equations (4.15), and, therefore, corresponds to a trivial conservation law of the first type. The second term corresponds to a trivial conservation law of the second type; its divergence vanishes identically since the tensor $F_{\mu\nu}$ is anti-symmetric. Therefore, according to the second part of Theorem 5, the infinite conservation law (4.17) with an arbitrary function $\theta(x)$ of all independent variables x_μ is trivial.

Consistent with the first part of Theorem 5 (and the Second Noether Theorem), the following identity relationship (4.12) between equations (4.15) and their derivatives holds:

$$i\varphi E_\varphi \mathcal{L} - i\bar{\varphi} E_{\bar{\varphi}} \mathcal{L} - \partial^\mu (E_{A^\mu} \mathcal{L}) \equiv 0. \quad (4.18)$$

Examples of infinite conservation laws involving arbitrary functions of all independent variables for non-Lagrangian systems are demonstrated in Section 6.

5 Arbitrary functions of not all independent variables, and conservation laws

In this section, we discuss the general case of infinite conservation laws that contain arbitrary functions of *not all* independent variables. Theorem 5 is valid for $f(x_1, \dots, x_p)$, or for arbitrary functions f of *all* independent variables. It is not valid for arbitrary functions of *not all* independent variables, e.g. $f(x_1)$. We present an extension of Theorem 5 to this case.

Theorem 6. *Suppose that the arbitrary function f in Theorem 5 depends only on the variables $(x_r, x_{r+1}, \dots, x_p)$ for some $1 < r \leq p$. Then there exists another conservation law of the system $\Delta : (5.4)$, different from (4.7), where the fluxes $M_f^r, M_f^{r+1}, \dots, M_f^p$ (4.8) are total divergences on-shell ($\Delta = 0$).*

An application of the variational derivative gives rise, instead of to the differential identity (4.12), to *another conservation law that does not involve f* . We also find that the conservation law (4.7) here is not trivial, and its *fluxes are total divergences on solutions*. In particular, a conservation law involving $f(t)$ has a density M_f^t that is a spatial divergence on solutions.

Proof of Theorem 6.

Part 1. According to (4.11):

$$\sum_{i=1}^{r-1} D_i M_f^i + \sum_{i=r}^p D_i M_f^i = f(-D)_J [\xi^{aJ} \Delta_a] + \sum_{i=r}^p D_i \Phi_J^i[f, \xi^{aJ} \Delta_a], \quad (5.1)$$

where $J = (J_r, J_{r+1}, \dots, J_p)$ is a multi-index. Applying the variational derivative E_f (2.5) annihilates the second and fourth terms:

$$\sum_{i=1}^{r-1} E_f D_i M_f^i = (-D)_J [\xi^{aJ} \Delta_a]. \quad (5.2)$$

Using (4.8) and the integration by parts identity (4.9), we have

$$D_i M_f^i = D_J f D_i M^{iJ} = f (-D)_J D_i M^{iJ} + \sum_{j=r}^p D_j \Phi_J^j[f, D_i M^{iJ}], \quad 1 \leq i \leq r-1. \quad (5.3)$$

Substituting (5.3) into (5.2), we obtain:

$$\sum_{i=1}^{r-1} D_i [(-D)_J M^{iJ}] = (-D)_J [\xi^{aJ} \Delta_a]. \quad (5.4)$$

Equation (5.4) is a conservation law distinct from (4.7). It is nontrivial if $\xi^{a0} \neq 0$ for some a .

Part 2. To prove the second part of the Theorem, we multiply (5.4) by f and subtract it from (5.1), noting that $\Phi_J^j[f, D_i M^{iJ}] = D_i \Phi_J^j[f, M^{iJ}]$ in (5.3):

$$\sum_{i=r}^n D_i \sum_{j=1}^{r-1} D_j \Phi_J^i[f, M^{jJ}] + \sum_{i=p}^n D_i M_f^i = \sum_{i=p}^n D_i \Phi_J^i[f, \xi^{aJ} \Delta_a]. \quad (5.5)$$

The general solution of this equation for the M_f^i is as follows:

$$M_f^i = - \sum_{j=1}^{r-1} D_j \Phi_J^i[f, M^{jJ}] + \Phi_J^i[f, \xi^{aJ} \Delta_a], \quad r \leq i \leq n, \quad (5.6)$$

where it is understood that M_f^i is arbitrary up to $M_f^i \rightarrow M_f^i + \lambda^i + \sum_{j=r}^n D_j R_f^{ij}$, where λ^i is a constant, and each $R_f^{ij} = -R_f^{ji}$ is an arbitrary function (i.e. up to fluxes of trivial conservation laws).

From (4.9), we see that each $\Phi_J^i[f, \xi^{aJ} \Delta]$ is a linear function of Δ and vanishes when $\Delta = 0$. Thus, on solutions u of $\Delta = 0$, each flux M_f^i in (5.6) is a total divergence. \square

5.1 Examples of Theorem 6

Consider Liouville's equation for $u(t, x)$:

$$\Delta = u_{tx} - e^u = 0 \quad (5.7)$$

with the Lagrangian:

$$\mathcal{L} = -\frac{1}{2} u_t u_x - e^u. \quad (5.8)$$

This Lagrangian admits an infinite symmetry:

$$X = [f'(t) + f(t)u_t] \partial_u =: \xi_f \partial_u \quad (5.9)$$

and an infinite set of corresponding conservation laws:

$$\frac{1}{2} D_t [u_x f_t - 2f e^u] + \frac{1}{2} D_x [(f_t + f u_t) u_t - f_{tt} u] = \xi_f E_u \mathcal{L} = \xi_f \Delta. \quad (5.10)$$

We write

$$\xi_f \Delta = f(u_t \Delta - D_t \Delta) + D_t(f \Delta).$$

Applying the Euler operator identity (5.2), we obtain:

$$u_t \Delta - D_t \Delta = (-D)_J[\xi^{aJ} \Delta_a]. \quad (5.11)$$

Since

$$\xi_f^a = \xi^{aJ} D_J f = u_t f(t) + f'(t),$$

it follows that

$$\xi^0 = u_t, \quad \xi^t = 1.$$

Calculating the RHS of (5.11), we obtain:

$$(-D)_J[\xi^{aJ} \Delta_a] = u_t(u_{tx} - e^u) - D_t(u_{xt} - e^u) = u_t u_{xt} - u_{xtt} = D_x \left[\frac{1}{2} u_t^2 - u_{tt} \right], \quad (5.12)$$

and

$$u_t \Delta - D_t \Delta = D_x \left[\frac{1}{2} u_t^2 - u_{tt} \right], \quad (5.13)$$

which can be easily verified. This is the additional conservation law independent of f predicted by Theorem 6, since the left hand side is zero when $\Delta = 0$. In fact, this is a first integral of Liouville's equation.

We now demonstrate the second part of Theorem 6 and rewrite the density of (5.10) as a total divergence on shell:

$$u_x f_t - 2f e^u = u_x f_t + 2f \Delta - 2f u_{tx} = D_x(u f_t - 2f u_t) + 2f \Delta.$$

Thus, when $\Delta = 0$, we see that the density is a total derivative in x .

Consider another example, *the equation for non-stationary transonic gas flow*

$$\Delta = 2u_{xt} + u_x u_{xx} - u_{yy} \quad (5.14)$$

with the Lagrangian function

$$L = -u_x u_t - \frac{u_x^3}{6} + \frac{u_y^2}{2}. \quad (5.15)$$

The conservation law in question is:

$$D_t M_f^t + D_x M_f^x + D_y M_f^y = \xi_f \Delta, \quad (5.16)$$

where

$$\xi_f = 2x f'(t) + 2y^2 f''(t) - f(t) u_x, \quad (5.17)$$

and the density and fluxes are given by

$$\begin{aligned} M_f^t &= f(t)[-u_x^2] + f'(t)[2xu_x - 2u] + f''(t)[2y^2u_x], \\ M_f^x &= -Lf(t) - 2xuf''(t) - 2y^2uf'''(t) + \xi_f(u_t + u_x^2/2), \\ M_f^y &= 4yuf''(t) - \xi_fu_y, \end{aligned} \quad (5.18)$$

We find that the density can be rewritten as a total divergence on solutions of $\Delta = 0$, in agreement with Theorem 6:

$$M_f^t \equiv Q_{f\Delta}^t + D_x T_f^x + D_y T_f^y, \quad (5.19)$$

where

$$\begin{aligned} T_f^x &= f(t)[-4xu_t + 4y^2u_{tt} + 2y^2u_xu_{xt} - xu_x^2] + f'(t)[2xu - 4y^2u_t - y^2u_x^2] + f''(t)[2y^2u], \\ T_f^y &= f(t)[2xu_y + 4yu_t - 2y^2u_{yt}] + f'(t)[2y^2u_y - 4yu], \\ Q_{f\Delta}^t &= (2xf(t) + 2y^2f'(t))\Delta - 2y^2f(t)D_t\Delta. \end{aligned} \quad (5.20)$$

Let us show how Theorem 6 can be used to generate conserved densities. Integrating the conservation law (5.16) over the whole space on shell, we will get

$$D_t \int M_f^t dx dy + D_x \int M_f^x dx dy + D_y \int M_f^y dx dy \doteq 0, \quad (5.21)$$

where M_f^x, M_f^y are determined in (5.18), and M_f^t is given by (5.19) and (5.20). Equation (5.21) leads to a quantity conserved in time

$$D_t \int M_f^t dx dy \doteq 0 \quad (5.22)$$

if the contributions (surface terms) from the second and third integrals vanish, i.e.

$$M^x \Big|_{x \rightarrow \pm\infty} \rightarrow 0, \quad M^y \Big|_{y \rightarrow \pm\infty} \rightarrow 0. \quad (5.23)$$

It is easy to see that, for sufficiently strict boundary conditions, all surface terms vanish. But for these boundary conditions, all terms in T^x , and T^y also vanish, and we get no conserved densities of the equation (5.14).

However, for weaker boundary conditions, the situation changes. Consider a “regular” asymptotic behavior

$$\begin{aligned} u, u_i &\xrightarrow{x \rightarrow \pm\infty} 0, \\ u, u_i &\xrightarrow{y \rightarrow \pm\infty} 0. \end{aligned} \quad (5.24)$$

In this case, after eliminating vanishing terms, we obtain

$$\begin{aligned} M_f^x \Big|_{x \rightarrow \pm\infty} &\rightarrow f'(t)2x(u_t + u_x^2/2) - 2xuf''(t), \\ M_f^y \Big|_{y \rightarrow \pm\infty} &\rightarrow f''(t)(4yu - 2y^2u_y). \end{aligned} \quad (5.25)$$

Requiring these terms to vanish:

$$f(t) = a, \quad a = \text{constant}. \quad (5.26)$$

T_f^x , and T_f^y reduce to

$$\begin{aligned} T^x &= -4xu_t - xu_x^2 + 2y^2(2u_{tt} + u_x u_{xt}), \\ T^y &= 4yu_t - 2y^2 u_{ty} + 2xu_y. \end{aligned} \quad (5.27)$$

Calculating M_f^t for this case, taking into consideration equation (5.14), and noting that the last term in T^y vanishes when integrated over y , we obtain

$$M_f^t \doteq T_{,x}^x + T_{,y}^y = -u_x^2 - 4xu_{xt} - 2xu_x u_{xx}. \quad (5.28)$$

The last two remaining terms in M^t are:

$$-2x(2u_{xt} + u_x u_{xx}) \doteq -2xu_{yy} = D_y(-2xu_y). \quad (5.29)$$

The contribution of this term in the integral over y is, obviously, zero. Finally, we have

$$M_f^t \doteq -u_x^2. \quad (5.30)$$

Thus, for regular boundary conditions (5.24), we obtain the following conserved quantity

$$D_t \int u_x^2 dx dy \doteq 0. \quad (5.31)$$

In [21], this result was obtained using a different technique.

6 Arbitrary functions of all independent variables in vorticity-type systems

6.1 General systems

We consider a class of vorticity-type systems studied in [4] and [5]:

$$\begin{aligned} \vec{\Delta} &= \vec{\omega}_t + \nabla \times \vec{M}(t, x, \omega, \dots) = 0, \quad \text{or} \quad \Delta_i = \omega_t^i + \epsilon^{ijk} D_j M^k = 0, \quad i = 1, 2, 3, \\ \Delta_4 &= \nabla \cdot \vec{\omega} = 0, \quad \text{or} \quad \Delta_4 = \omega_i^i = 0. \end{aligned} \quad (6.1)$$

Here, ω is the vorticity, and \vec{M} is some vector function.

The following infinite set of conservation laws of (6.1) involving an arbitrary function $F(t, x)$ was presented in [4] and [5]:

$$\partial_t(\vec{\omega} \cdot \nabla F) + \nabla \cdot (\vec{M} \times \nabla F - \partial_t F \vec{\omega}) = \nabla F \cdot \vec{\Delta} - \partial_t F \Delta_4 = 0. \quad (6.2)$$

We show that the conservation laws (6.2) are trivial. Indeed, applying the procedure of Theorem 5 to (6.2) (i.e. taking the variational derivative E_F of (6.2)) leads us to the following off-shell identity:

$$D_t \Delta_4 - \nabla \cdot \vec{\Delta} = 0. \quad (6.3)$$

Substituting (6.3) into (6.2) yields:

$$\partial_t(\vec{\omega} \cdot \nabla F) + \nabla \cdot (\vec{M} \times \nabla F - \partial_t F \vec{\omega}) = \nabla \cdot (F \vec{\Delta}) - D_t(F \Delta_4). \quad (6.4)$$

Since the characteristic (coefficient of Δ) of this conservation law is zero, the conservation law (6.2) is trivial. It is a combination of two kinds of triviality: the divergence terms on the RHS have fluxes that vanish on solutions (on shell, $\Delta = 0$), and, therefore, do not contribute to the conserved densities or integrals. The rest is a differential identity that holds everywhere in the extended space (off-shell).

The relation (6.3) expresses an interdependence of equations in the governing differential system Δ , or the fact that the system is under-determined (abnormal, see [16]). The relation (6.3) was noted in [5] in the case of the Navier-Stokes equations, but not the fact that the infinite conservation laws (6.2) are trivial.

6.2 Symmetries of the Euler vorticity system

We now apply Theorem 4 to the vorticity system (6.1) for velocity u and vorticity ω :

$$\begin{aligned} \Delta_i &= \omega_t^i + \partial_j(\omega^i u^j - \omega^j u^i) = 0, \quad i = 1, 2, 3, \\ \Delta_4 &= \omega_i^i = 0, \\ \Delta_5 &= u_i^i = 0. \end{aligned} \quad (6.5)$$

This system is clearly quasi-Noether, and we can set alternative Lagrangians as $\mathcal{A} = \Delta_i$ for each $i = 1, 2, 3$. It admits several infinite sets of symmetries. If functions $g^k(t, x)$ are arbitrary, and $f^i = \epsilon^{ijk} \partial_j g^k$ ($\partial_i f^i = 0$), then the infinite symmetries are given by:

$$\begin{aligned} X_f &= \alpha_f^i \partial_{\omega^i} + \beta_f^i \partial_{u^i}, \\ \alpha_f^i &= \omega^j \partial_j f^i - f^j \partial_j \omega^i, \\ \beta_f^i &= \partial_t f^i + u^j \partial_j f^i - f^j \partial_j u^i, \end{aligned} \quad (6.6)$$

(the symmetries (6.6) were obtained with the use of the program *MathLie*TM in [2]). Applying X_f to the alternative Lagrangians $\mathcal{A} = \Delta_i$, $1 \leq i \leq 3$ and using the Noether identity (2.11), these infinite symmetries lead to the following infinite sets of conservation laws:

$$D_t \alpha_f^i + D_j (\alpha_f^i u^j - \alpha_f^j u^i + \omega^i \beta_f^j - \omega^j \beta_f^i) = \partial_j f^i \Delta_i - \partial_t f^i \Delta_4 - D_j (f^j \Delta_i). \quad (6.7)$$

If we apply now the procedure of Theorem 4 to the conservation laws (6.7), using the fact that functions g are arbitrary functions of all independent variables, we obtain the following (off-shell) differential identities:

$$D_i (D_t \Delta_4 - D_j \Delta_j) = 0, \quad 1 \leq i \leq 3. \quad (6.8)$$

Examining the interior expression, and the fact that $D_j \Delta_j = \partial_j \partial_t \omega^j = \partial_t \partial_j \omega^j = D_t \Delta_4$, we find a stronger off-shell identity:

$$D_t \Delta_4 - D_j \Delta_j = 0. \quad (6.9)$$

Thus, the existence of infinite symmetries (6.6) with arbitrary functions of all independent variables is an indication that the vorticity system (6.5) is under-determined.

6.3 Symmetries and Noether's second theorem

Let us try to relate the differential identity (6.9) (and trivial conservation laws (6.2)) to an infinite variational symmetry according to Theorem 2.

As shown in [16], any differential system can be given a variational formulation by introducing additional unknown variables and equations. If we start with a differential system Δ_a , $1 \leq a \leq n$ for variables u^q , $1 \leq q \leq m$, we can introduce additional “adjoint variables” v^a , $1 \leq a \leq n$ and define the following Lagrangian:

$$\mathcal{L}(u, v) = v^a \Delta_a(u). \quad (6.10)$$

Indeed, applying the Euler operator $E_{v^a} = \partial_{v^a} + \dots$ to \mathcal{L} recovers Δ_a . The adjoint variables v^a will satisfy their own differential systems, but these functions are, in general, non-physical.

We modify the approach of [16] and construct a variational formulation for (6.1). Letting $\vec{\omega} = \nabla \times u$ for “velocity potential” \vec{u} , we rewrite (6.1) equivalently as a higher order system for \vec{u} :

$$\vec{\Delta} = \partial_t (\nabla \times \vec{u}) + \nabla \times \vec{M}(t, x, \nabla \times \vec{u}, \dots) = 0. \quad (6.11)$$

We define the following Lagrangian:

$$\mathcal{L}(u, v) = -\vec{u} \cdot \partial_t (\nabla \times \vec{v}) + \vec{M} \cdot \nabla \times \vec{v}. \quad (6.12)$$

This Lagrangian recovers (6.11), since

$$\mathcal{L} - \vec{v} \cdot \vec{\Delta} = -\partial_t (\vec{v} \cdot \nabla \times \vec{u}) + \nabla \cdot (\vec{v} \times \vec{M} - \partial_t \vec{v} \times \vec{u}),$$

which means that $E_{v^a} \mathcal{L} = \Delta_a$.

Now, the Lagrangian (6.12) admits the following infinite set of (gauge) symmetries:

$$\vec{u} \rightarrow \vec{u} + \nabla F, \quad (6.13)$$

where $F(t, x)$ is an arbitrary function of all independent variables.

Applying the Noether identity (2.11), we associate to this symmetry the following set of conservation laws involving an arbitrary function F :

$$\partial_t (\nabla F \cdot \nabla \times \vec{u}) + \nabla \cdot (\vec{M} \times \nabla F + \nabla (\partial_t F) \times \vec{u}) = \nabla F \cdot \vec{\Delta}. \quad (6.14)$$

Rewriting the last term in the LHS of (6.14):

$$\nabla (\partial_t F) \times \vec{u} = \nabla \times (\partial_t F \vec{u}) - \partial_t F \nabla \times \vec{u},$$

and substituting it into (6.14):

$$\partial_t (\nabla F \cdot \nabla \times \vec{u}) + \nabla \cdot (\vec{M} \times \nabla F - \partial_t F \nabla \times \vec{u}) = \nabla F \cdot \vec{\Delta}, \quad (6.15)$$

for $\vec{\omega} = \nabla \times \vec{u}$ recovers the infinite conservation law (6.4) proposed in [4].

Finally, performing on (6.15) the integration procedure of Noether's second theorem (see e.g. Theorem 4) yields the following off-shell identity:

$$\nabla \cdot \vec{\Delta} = 0, \quad (6.16)$$

which verifies the fact that the vorticity system (6.11) is a total curl. This is unsurprising from a fluid dynamics perspective, since the vorticity system is obtained precisely by taking the curl of the Navier-Stokes system. We see that the infinite conservation laws (6.2) are expressions of this structural fact.

6.4 Navier-Stokes equations

In [4] were considered special cases of (6.2) for physical systems including Maxwell's equations, the Navier-Stokes equations, and the equations of magnetohydrodynamics. In [5] were considered infinite conservation laws (6.2) in the case of the Euler and Navier-Stokes equations of incompressible fluid dynamics; many more results were subsequently derived using these conservation laws. For Navier-Stokes, the system (6.1) for ω takes the following form:

$$\begin{aligned} \vec{\Delta} &= \vec{\omega}_t + \nabla \times (\vec{\omega} \times \vec{u} - \nu \nabla^2 \vec{u}) = 0, \\ \Delta_4 &= \nabla \cdot \vec{\omega} = 0. \end{aligned} \quad (6.17)$$

Here, \vec{u} is the velocity vector, and ν is the viscosity. The system (6.17) is the system (6.1) for $\vec{M} = \vec{\omega} \times \vec{u} - \nu \nabla^2 \vec{u}$; it comprises the *vorticity equations* of incompressible flow. Equation Δ_4 expresses the fact that $\vec{\omega} = \nabla \times \vec{u}$.

The infinite conservation laws (6.2) considered in these papers take the following form:

$$\begin{aligned} \partial_t (\vec{\omega} \cdot \nabla F) + \nabla \cdot [(\vec{\omega} \times \vec{u} - \nu \nabla^2 \vec{u}) \times \nabla F - (\partial_t F) \vec{\omega}] &= \nabla F \cdot \vec{\Delta} - (\partial_t F) \Delta_4 \\ &= \nabla \cdot (F \vec{\Delta}) - D_t (F \Delta_4). \end{aligned} \quad (6.18)$$

As we have shown, these conservation laws are trivial: the RHS is a divergence expression with fluxes that vanish on-shell (triviality of the first kind) and, therefore, do not contribute to the conserved densities or integrals. The rest is a differential identity that holds everywhere in the extended space (off-shell; triviality of the second kind).

Let us discuss the conserved charge of (6.18), namely $Q[\vec{\omega}] := \int \vec{\omega} \cdot \nabla F \, d^3x$, which was proposed in [4]. If we integrate (6.18) over space and assume that F and $\vec{\omega}$ are such that the surface terms of the fluxes (arguments of $\nabla \cdot ()$) vanish at the boundary (infinity), then we obtain the following relation for Q :

$$D_t Q[\vec{\omega}] = -D_t \int F \Delta_4 \, d^3x, \quad (6.19)$$

meaning that Q is conserved on-shell:

$$D_t Q[\vec{\omega}] \doteq 0. \quad (6.20)$$

Let us now construct an off-shell identity by subtracting the terms on the right hand side of (6.18) from both sides:

$$\partial_t(\vec{\omega} \cdot \nabla F) + D_t(F\Delta_4) + \nabla \cdot [(\vec{\omega} \times \vec{u} - \nu \nabla^2 \vec{u}) \times \nabla F - \partial_t F \vec{\omega}] - \nabla \cdot (F \vec{\Delta}) = 0, \quad (6.21)$$

or

$$\partial_t [\nabla \cdot (F \vec{\omega})] + \nabla \cdot [\nabla \times (F(\vec{\omega} \times \vec{u} - \nu \nabla^2 \vec{u})) - \partial_t (F \vec{\omega})] = 0. \quad (6.22)$$

The equation (6.22) is an equivalent statement of (6.18). Let $R[\vec{\omega}] := \int \nabla \cdot (F \vec{\omega}) d^3x$ be the conserved charge of this formulation. Integrating (6.22) over space and assuming that all surface terms vanish at the boundary (infinity), we see that R satisfies the following off-shell integral identity:

$$D_t R[\vec{\omega}] = 0,$$

meaning that R is conserved off-shell. In general, when $\vec{\omega}$ does not satisfy (6.17), $Q \neq R$. However, for those $\vec{\omega}$ that satisfy $\Delta[\vec{\omega}] = 0$, assuming the stricter of two boundary conditions, we can demonstrate that $Q = R$ on-shell. Indeed,

$$Q = \int \vec{\omega} \cdot \nabla F d^3x \doteq \int (\vec{\omega} \cdot \nabla F + F \nabla \cdot \vec{\omega}) d^3x = \int \nabla \cdot (F \vec{\omega}) d^3x = R.$$

If we call R the “trivial charge”, the charge that is conserved for every $\vec{\omega}$, we see that, although Q is a nontrivial charge off-shell, it is equal to the trivial charge R on-shell. This is a contrast to conventional conserved quantities for well-defined systems, which are equal to trivial charges neither on- nor off-shell.

Further study is needed to understand the role of trivial conservation laws and the nature of conserved charges for under-determined systems; see also [12], [16], [17].

6.5 Ertel’s Theorem

Infinite trivial conservation laws (6.18) also play a role in inviscid fluid dynamics, $\nu = 0$. They take the form of *Ertel’s theorem*, an important tool in atmospheric sciences (see e.g. [13]). The Euler vorticity equations for ω are as follows:

$$\begin{aligned} \vec{\Delta} &= \vec{\omega}_t + \nabla \times (\vec{\omega} \times \vec{u}) = 0, \\ \Delta_4 &= \nabla \cdot \vec{\omega} = 0, \end{aligned} \quad (6.23)$$

which is the system (6.1) for $\vec{M} = \vec{\omega} \times \vec{u}$, where, as earlier, \vec{u} is the velocity vector.

Ertel’s theorem [31] states that the following relationship holds on solutions of (6.23):

$$\partial_t(\vec{\omega} \cdot \nabla \psi) + \nabla \cdot [(\vec{\omega} \cdot \nabla \psi) \vec{u}] = 0. \quad (6.24)$$

Here, $\psi(t, x)$ is a scalar function advected with the flow of \vec{u} . This means that it solves the following differential equation:

$$\partial_t \psi + \vec{u} \cdot \nabla \psi = 0. \quad (6.25)$$

In fact, Ertel's theorem (6.24) follows directly from trivial conservation law (6.18) for $\nu = 0$. To see this, we start with (6.18) for $\nu = 0$ and $F = \psi$:

$$\partial_t(\vec{\omega} \cdot \nabla \psi) + \nabla \cdot [(\vec{\omega} \times \vec{u}) \times \nabla \psi - \partial_t \psi \vec{\omega}] = \nabla \cdot (\psi \vec{\Delta}) - D_t(\psi \Delta_4). \quad (6.26)$$

Since $(\vec{\omega} \times \vec{u}) \times \nabla \psi = (\vec{\omega} \cdot \nabla \psi) \vec{u} - (\vec{u} \cdot \nabla \psi) \vec{\omega}$, equation (6.26) becomes:

$$\partial_t(\vec{\omega} \cdot \nabla \psi) + \nabla \cdot [(\vec{\omega} \cdot \nabla \psi) \vec{u} - (\partial_t \psi + \vec{u} \cdot \nabla \psi) \vec{\omega}] = \nabla \cdot (\psi \vec{\Delta}) - D_t(\psi \Delta_4).$$

But since ψ satisfies (6.25), Ertel's theorem (6.24) follows:

$$\partial_t(\vec{\omega} \cdot \nabla \psi) + \nabla \cdot [(\vec{\omega} \cdot \nabla \psi) \vec{u}] = \nabla \cdot (\psi \vec{\Delta}) - D_t(\psi \Delta_4),$$

which is a trivial conservation law, since its characteristic is zero.

7 Conclusion

We considered quasi-Noether systems, a class of differential system that includes all equations possessing conservation laws. We extended an approach based on the Noether operator identity and formulated and proved an extension of a Second Noether theorem for quasi-Noether systems. As in the case of equations of a variational problem, the existence of an infinite symmetry group of the system with an arbitrary function of *all* independent variables generates a differential identity between the equations of the system and their derivatives. In addition, we showed that infinite conservation laws involving an arbitrary function of *all* independent variables are necessarily trivial. We analyzed recently obtained sets of infinite conservation laws (with an arbitrary function of all independent variables) for the Euler and Navier-Stokes equations in vorticity formulations and demonstrated that these conservation laws are trivial. We also showed that the existence of infinite conservation laws with an arbitrary function of *not all* independent variables leads to the conclusion that these conservation laws are, in general, non-trivial, but that their fluxes are total divergences on solutions.

References

- [1] Anderson, I. M., The variational bicomplex, Utah State University, 1989.
- [2] Baumann G. (2000). Symmetry Analysis of Differential Equations with Mathematica (MathLie), Springer, New York.
- [3] Caviglia G., Symmetry transformations, isovectors, and conservation laws, *J. Math. Phys* **27** (1986), 973–978.

- [4] Cheviakov, A. F. (2014) Conservation properties and potential systems of vorticity-type equations. *J. Math. Phys.* 92(14) **55**, no. 3, 033508, 16pp.
- [5] Cheviakov, A. F., and Oberlack, M. (2014). Generalized Ertel's theorem and infinite hierarchies of conserved quantities for three-dimensional time-dependent Euler and Navier-Stokes equations. *Journal of Fluid Mechanics* **760**, 368-386.
- [6] Hydon, P., and Mansfield, E. (2011). Extensions of Noether's Second Theorem: from continuous to discrete systems. *Proc. R. Soc. A* (2011) **467**, 3206-3221.
- [7] Ibragimov N.H., Nonlinear self-adjointness and conservation laws. *J Phys A Math Theor* **44**, (2011) :432002, 8pp.
- [8] Ibragimov, N.H., Transformation groups applied to mathematical physics. Translated from the Russian. Mathematics and its Applications (Soviet Series). D. Reidel Publishing Co., Dordrecht, 1985.
- [9] Ibragimov, N.H., A new conservation theorem, *J. Math. Anal. Appl.* **333**, (2007), 311–328.
- [10] Juras M., and Anderson I.M., Generalized Laplace invariants and the method of Darboux, *Duke Math. J.* **89**, (1997), 351–375.
- [11] Lunev, F. A., An analogue of the Noether theorem for non-Noether and nonlocal symmetries. *Theor. Math. Phys.* **84**, 2 (1991), 816–820.
- [12] D'Emilio, E., and Mintchev, M. Identically Conserved Currents and their Charges. *Lett. Math. Phys.* **12**, No 1 (1986), 77-86.
- [13] Muller, P. Ertel's potential vorticity theorem in physical oceanography. *Reviews of Geophysics* **33**, No 1 (1995), 67-97.
- [14] Navarro, J., and Sancho, J. B. Noether's second theorem on natural bundles. *Int. J. Geom. Methods Mod. Phys.* **11**, No. 9, (2014), 1460035, 8 pp.
- [15] Noether, E., Invariantenprobleme, Nachr. König. Gessell. Wissen. Göttingen, Math.-Phys. **K1**. (1918), pp. 235–257.
- [16] Olver, P. J. (2000). Applications of Lie groups to differential equations (Vol. 107). Springer Science, and Business Media.
- [17] Olver, P. J. (1983) Conservation laws and null divergences. *Math. Proc. Camb. Phil. Soc.* **94**, 529–540
- [18] Rosen J., Some properties of the Euler-Lagrange operators, *Preprint TAUP-269-72*, Tel-Aviv University, Tel-Aviv (1972).
- [19] Rosenhaus, V., and Katzin, G. H. (1994). On symmetries, conservation laws, and variational problems for partial differential equations. *J. Math. Phys.* **35** (1994), 1998–2012.

- [20] Rosenhaus, V., and Katzin, G. H. (1996). Noether operator relation and conservation laws for partial differential equations. in *Proc. of First Workshop: Nonlinear Physics. Theory and Experiment, June 29-July 7, 1995, Gallipoli, Italy*, World Scientific, 1996, 286–293.
- [21] Rosenhaus, V. (2002). Infinite Symmetries and Conservation Laws. *J. Math. Phys.* **43**, 6129–6150 (2002).
- [22] Rosenhaus V., Boundary conditions and conserved densities for potential Zabolotskaya–Khokhlov equation, *J. Nonlinear Math. Phys.* **13** (2006), 255–270.
- [23] Rosenhaus V., On conserved densities and asymptotic behavior for the potential Kadomtsev–Petviashvili equation, *J. Phys. A: Math. Gen.* **39** (2006), 7693–7703.
- [24] Rosenhaus V., On infinite set of conservation laws for infinite symmetries, *Theor. Math. Phys.* **151** (2007), 869–878.
- [25] Rosenhaus, V. (2013). On Differential Equations with Infinite Conservation Laws, Proceedings of the 6th Intern. Workshop Group Analysis of Differential Equations and Integrable Systems, pp. 192–202, University of Cyprus, Nicosia, 2013.
- [26] Rosenhaus, V., and Shankar R. Sub-symmetries and Infinite Conservation Laws for the Euler Equations, to be published.
- [27] Sarlet W., Cantrijn F., and Crampin M., Pseudo-symmetries, Noether’s theorem and the adjoint equation, *J. Phys. A* **20** (1987) 1365–1376.
- [28] Sheftel M.B., Symmetry group analysis and invariant solutions of hydrodynamic-type systems, *Int. J. Math. Math. Sci.* (2004), No 9–12, 487–534.
- [29] Vinogradov A.M., Local Symmetries and Conservation Laws, *Acta Appl. Math.* **2** (1984), 21–78.
- [30] Vladimirov, V.S., and Zharinov, V.V., Closed forms associated with linear differential operators. *Differ. Equations* **16** (1980), 534–552.
- [31] Webb G.M., and Mace R.L. Potential vorticity in magnetohydrodynamics. *J. Plasma Physics* **81**, (2015) 905810115, 1–18.
- [32] Zharinov, V. V. Conservation laws of evolution systems, *Theor. Math. Phys.* **68**(2) (1986), 745–751.
- [33] Zhiber A.V., and Sokolov V.V., Exactly integrable hyperbolic equations of Liouville type, *Russian Math. Surveys* **56**, (2001), 61–101.