

Local convergence analysis of Newton's method for solving strongly regular generalized equations

O. P. Ferreira* G. N. Silva †

September 26, 2016

Abstract

In this paper we study Newton's method for solving generalized equations in Banach spaces. We show that under strong regularity of the generalized equation, the method is locally convergent to a solution with superlinear/quadratic rate. The presented analysis is based on Banach Perturbation Lemma for generalized equation and the classical Lipschitz condition on the derivative is relaxed by using a general majorant function, which enables obtaining the optimal convergence radius, uniqueness of solution as well as unifies earlier results pertaining to Newton's method theory.

Keywords: Generalized equation, Newton's method, strong regularity, majorant condition.

Mathematical Subject Classification (2010): Primary 65K15; 49M15; 90C31

1 Introduction

In this paper we consider the Newton's method for solving the generalized equation of the form

$$f(x) + F(x) \ni 0, \quad (1)$$

where $f : \Omega \rightarrow Y$ is a continuously differentiable function, X and Y are Banach spaces, $\Omega \subseteq X$ is an open set and $F : X \rightrightarrows Y$ is a set-valued mapping with nonempty closed graph. As is well known, the model of the generalized equation (1) covers several class of problems, due to this important characteristic it has been studied in several works, having [2, 3, 10, 14, 15, 16, 23, 29] as part of a whole. For instance, as we can see, if F is the normal cone mapping N_C , of a convex set C in Y and $Y = X^*$ is the dual of X , the inclusion (1) is the variational inequality problem; for more details see [14].

Newton's method is undoubtedly one of the most popular methods for numerically solving nonlinear equation. This is because of its importance both theoretical and practical, and even more is due to its quadratic rate of convergence. Throughout the years, this method has been extended in many directions by several authors, one of the most studied currently is the generalization of this to solve (1), which has its origin in the works of N. H. Josephy [23]. Based on the work of N. H. Josephy [23], we study the local convergence of the following Newton's method for solving (1):

$$f(x_k) + f'(x_k)(x_{k+1} - x_k) + F(x_{k+1}) \ni 0, \quad k = 0, 1, \dots \quad (2)$$

*IME/UFG, CP-131, CEP 74001-970 - Goiânia, GO, Brazil (Email: orizon@ufg.br). The author was supported in part by FAPEG, CNPq Grants 305158/2014-7 and PRONEX–Optimization(FAPERJ/CNPq).

†CCET/UFOB, CEP 47808-021 - Barreiras, BA, Brazil (Email: gilson.silva@ufob.edu.br). The author was supported in part by CAPES .

This algorithm has been studied in several papers including but not limited to [2, 3, 6, 9, 10, 11, 16]; see also [14, Section 6C] and [26]. If $F \equiv 0$, the iteration (2) becomes the usual Newton method for solving the equation $f(x) = 0$. If $F = N_C$, the normal cone mapping of a convex set C in Y and $Y = X^*$, then (2) is the version of the Newton's method for solving variational inequality; see [6, 9, 23]. In particular, if (1) represents the Karush-Kuhn-Tucker optimality conditions for a mathematical programming problem, then (2) describes the well-known sequential quadratic programming method; see for example a detailed discussion in [14, pag. 334]; see also [12].

Under the assumption that f is Fréchet differentiable in some neighborhood of a solution \bar{x} of (1), S. M. Robinson in [30] obtained a condition on the linearization of (1) about \bar{x} , i.e., on the generalized equation

$$f(\bar{x}) + f'(\bar{x})(x - \bar{x}) + F(x) \ni 0,$$

which he called *strong regularity*, in order to guarantee unique solution of the generalized equation

$$f(\hat{x}) + f'(\hat{x})(x - \hat{x}) + F(x) \ni 0,$$

for all \hat{x} in a neighborhood \bar{x} . The classic local analysis of Newton's method for solving $f(x) = 0$ require invertibility of the derivative f' at the solution, which is actually critical for the well definition of the method. Therefore, for the local analysis of Newton's method for solving (1) we will need of a similar concept, namely, the *strong regularity* of $f + F$ at the solution $\bar{x} \in X$ for $0 \in Y$. If $X = Y$ and $F = \{0\}$, then strong regularity of $f + F$ at the solution $\bar{x} \in X$ for $0 \in X$ is equivalent to assumption that $f'(\bar{x})^{-1}$ is a continuous linear operator. An important case is when (1) represents the Karush-Kuhn-Tucker's system for the standard nonlinear programming problem with a strict local minimizer, see [14] pag. 232. In this case, strong regularity of this system is equivalent to the linear independence of the gradients of the active constraints and a strong form of the second-order sufficient optimality condition; for details see [13, Theorem 6]. The analysis presented in this paper will be made under strong regularity on the solution of (1).

It is well-known that, to obtain quadratic convergence rate of Newton's method (2), the Lipschitz continuity of f' in a neighborhood of a solution of (1) is required, see for example [3, 10, 14, 15]. Indeed, keeping control of the derivative is an important point in the convergence analysis of Newton's methods and its variations, as we can see in [7, 16, 2, 20, 36]. Recently, there has been an increased interest in the study of Newton's method and its variations for solving the equation $f(x) = 0$, by relaxing the hypothesis of Lipschitz continuity of f' . For instance, the majorant condition is one of those conditions that relax the Lipschitz condition which has several equivalent formulations, see for example [1, 4, 21, 20, 25, 27, 35, 36, 37]. The advantage of working with a majorant condition is that it makes us clearly see how big the radius of convergence is, besides allow us unify several convergence results pertaining to Newton's method; see [21, 35]. In this paper, under the majorant condition, we establish a local convergence analysis of Newton's method (2) by assuming strong regularity of $f + F$ at the solution $\bar{x} \in X$ for $0 \in X$. Before proving our main result, which establish the optimal convergence radius for the method with respect to the majorant condition and uniqueness of solution, a clear relationship between the majorant function and the function defining the generalized equation is obtained. As special cases, we present an analysis of this result under Lipschitz's and Smale's conditions. Up to our knowledge, this is the first time that the Newton method for solving generalized equations under a general majorant condition and, in particular, under Smale's condition is analyzed, similar studies has been done in [1, 8, 31, 32, 27, 24, 25, 36]. In addition, it is worth mentioning that our approach is based in the Banach Perturbation Lemma obtained by S. M. Robinson in [30, Theorem 2.4]. In this sense, our approach is related to the techniques used in [7, 9, 23].

The organization of the paper is as follows. In the following section we present background material and some technical results used throughout the paper. Section 3 is devoted to our main result

and in Section 3.1 properties of the majorant function, the main relationships between the majorant function and the generalized equation, the uniqueness of the solution and the optimal convergence radius are established. In Section 3.2 the main result is then proved and some applications of this result are given in Section 4. Finally, conclusions are stated in Section 5.

2 Preliminaries

We use the following notation. Let X, Y be Banach spaces. The *space consisting of all continuous linear mappings* $A : X \rightarrow Y$ will be denote by $\mathcal{L}(X, Y)$ and the *operator norm* of A will be defined by $\|A\| := \sup \{\|Ax\| : \|x\| \leq 1\}$. Let $\Omega \subseteq X$ and $h : \Omega \rightarrow Y$ a function with Fréchet derivative at all $x \in \text{int}(\Omega)$. The Fréchet derivative of h at x is the linear map $h'(x) : X \rightarrow Y$ which is continuous. We identify as the *graph* of the set-valued mapping $H : X \rightrightarrows Y$ the set $\text{gph } H := \{(x, y) \in X \times Y : y \in H(x)\}$. The *inverse* H^{-1} of a map $H : X \rightrightarrows Y$ is defined as $H^{-1}(y) := \{x \in X : y \in H(x)\}$. Define $B(x, \delta) := \{y \in X : \|x - y\| < \delta\}$ and $B[x, \delta] := \{y \in X : \|x - y\| \leq \delta\}$ as the *open* and *closed balls* centered at x with radius $\delta \geq 0$.

Definition 1. *Let Ω be a nonempty, open, convex subset of X . Let $h : \Omega \rightarrow Y$ be a function with continuous derivative h' and $H : X \rightrightarrows Y$ be a set-valued mapping. The partial linearization mapping of $h + H$ at x is the set-valued mapping $L_h(x, \cdot) : X \rightrightarrows Y$ given by*

$$L_h(x, z) := h(x) + h'(x)(z - x) + H(z).$$

Thus, the linearization of a generalized equation $h(z) + H(z) \ni 0$ at x is obtained by replacing $h + H$ with $L_h(x, \cdot)$. The inverse $L_h(x, \cdot)^{-1}$ of the map $L_h(x, \cdot)$ at $y \in Y$ is denoted by

$$L_h(x, y)^{-1} := \{z \in X : y \in h(x) + h'(x)(z - x) + H(z)\}. \quad (3)$$

An important element in the analysis of Newton's method, for solving the equation $f(x) = 0$, is the behavior of the inverse $f'(x)^{-1}$, for x in a neighborhood of a solution \bar{x} . The analogous element for the generalized equation (1) is the behavior of the inverse mapping $L_f(x, \cdot)^{-1}$, for x in a neighborhood of a solution \bar{x} . It is worth to point out that, N. H. Josephy in [23] was the first to consider Newton's method for solving the generalized equation $f(x) + N_C(x) \ni 0$, where N_C is the normal cone of a convex set $C \subset \mathbb{R}^n$, by defining the Newton iteration as

$$f(x_k) + f'(x_k)(x_{k+1} - x_k) + N_C(x_{k+1}), \quad k = 0, 1, \dots$$

For analyzing this method, was employed the important concept of strong regularity defined by S. M. Robinson [30], which assure "good behavior" of $L_f(x, \cdot)^{-1}$, for x in a neighborhood of a solution \bar{x} . Here we adopt the following definition due to S. M. Robinson; see [30].

Definition 2. *Let X, Y be Banach spaces, Ω be an open and nonempty subset of X , $h : \Omega \rightarrow Y$ be Fréchet differentiable with derivative h' and $H : X \rightrightarrows Y$ be a set-valued mapping. The mapping $h + H$ is said to be strongly regular at x for y with modulus $\lambda > 0$, when $y \in h(x) + H(x)$ and there exist neighborhoods U_x of x and V_y of y in Y such that $U_x \subset \Omega$, the mapping $z \mapsto L_h(x, z)^{-1} \cap U_x$ is a single-valued function from V_y to U_x , which is Lipschitzian on V_y with modulus λ , i.e.,*

$$\|L_h(x, u)^{-1} \cap U_x - L_h(x, v)^{-1} \cap U_x\| \leq \lambda \|u - v\|, \quad \forall u, v \in V_y.$$

Since the mapping $z \mapsto L_h(x, z)^{-1} \cap U_x$ is single-valued from V_y to U_x , for simplify the notation we are using in above definition $w = L_h(x, z)^{-1} \cap U_x$ instead of $\{w\} := L_f(x, z)^{-1} \cap U_x$. From now on we will use this simplified notation.

Remark 1. If in above definition $H(x) \equiv \{0\}$ then the property of $h + H \equiv h$ be strongly regular at the solution \bar{x} for 0, reduces to $h'(\bar{x})$ has an inverse $h'(\bar{x})^{-1}$. Moreover, in this case, $\lambda = \|h'(\bar{x})^{-1}\|$, $U_{\bar{x}} = X$ and $V_{\bar{x}} = Y$. An important particular instance is when (1) represents the Karush-Kuhn-Tucker's system for the standard nonlinear programming problem with a strict local minimizer. In this case, the strong regularity of this system is equivalent to the linear independence of the gradients of the active constraints and the strong second-order sufficient optimality condition; see [14, Example 6C.8], see also [13, Theorem 6].

For a detailed discussion about Definition 2 see [14, 30]. The next result is a type of implicit function theorem for generalized equations satisfying the condition of strong regularity, its proof is similar to [30, Theorem 2.1], it also can be seen as a particular instance of [14, Theorem 5F.4] on page 294.

Theorem 1. Let X, Y and Z be Banach spaces, $G : X \rightrightarrows Y$ be a set-valued mapping and $g : Z \times X \rightarrow Y$ be a continuous function, having partial Fréchet derivative with respect to the second variable $D_x g$ on $Z \times X$, which is also continuous. Let $\bar{p} \in Z$ and suppose that \bar{x} solves the generalized equation

$$g(\bar{p}, x) + G(x) \ni 0.$$

Assume that the mapping $g(\bar{p}, \cdot) + G$ is strongly regular at \bar{x} for 0, with modulus λ . Then, for any $\epsilon > 0$ there exist neighborhoods U_ϵ of \bar{x} and V_ϵ of \bar{p} and a single-valued function $s : V_\epsilon \rightarrow U_\epsilon$ such that for any $p \in V_\epsilon$, $s(p)$ is the unique solution in U_ϵ of the inclusion $g(p, x) + G(x) \ni 0$, and $s(\bar{p}) = \bar{x}$. Moreover, there holds

$$\|s(p') - s(p)\| \leq (\lambda + \epsilon) \|g(p', s(p)) - g(p, s(p))\|, \quad \forall p, p' \in V_\epsilon.$$

Indeed, the first version of the Theorem 1 was proved by S. M. Robinson; see [30, Theorem 2.1], to the particular case $F = N_C$, where N_C is the normal cone of a convex set $C \subset X$. As an application, a version of the Banach Perturbation Lemma involving the normal cone was obtained; see [30, Theorem 2.4]. N. H. Joseph in [23], used this version of Banach Perturbation Lemma, see [23, Corollary 1], for proving that the Newton iteration

$$f(x_k) + f'(x_k)(x_{k+1} - x_k) + N_C(x_{k+1}) \ni 0, \quad k = 0, 1, \dots,$$

where N_C is the normal cone of a convex set $C \subset \mathbb{R}^n$, is quadratically convergent to a solution of $f(x) + N_C(x) \ni 0$. In the next lemma we state a version of the Banach Perturbation Lemma involving a general set-valued mapping, its proof is similar to the correspondent one [30, Theorem 2.4].

Lemma 2. Let X, Y be Banach spaces, a_0 be a point of Y , $G : X \rightrightarrows Y$ be a set-valued mapping and $A_0 : X \rightarrow Y$ be a bounded linear operator. Suppose that \bar{x} is a point of X which satisfies the generalized equation

$$0 \in A_0 x + a_0 + G(x).$$

Assume that the mapping $A_0 + a_0 + G$ is strongly regular at \bar{x} for 0 with modulus λ . Then there exist neighborhoods M of A_0 in $\mathcal{L}(X, Y)$, N of a_0 and W of origin in Y , and U of \bar{x} , such that, for any $A \in M$ and $a \in N$, letting $T(A, a, \cdot) : U \rightrightarrows Y$ be defined as

$$T(A, a, x) := Ax + a + G(x),$$

then $y \mapsto T(A, a, y)^{-1} \cap U$ is a single-valued function from W to U . Moreover, letting a neighborhood \bar{M} of A_0 such that $\bar{M} \subset M$ and $\lambda \|A - A_0\| < 1$ for each $A \in \bar{M}$, then for each $A \in \bar{M}$ and $a \in N$ the function $y \mapsto T(A, a, y)^{-1} \cap U$ is Lipschitz on W as follows

$$\|T(A, a, y_1)^{-1} \cap U - T(A, a, y_2)^{-1} \cap U\| \leq \frac{\lambda}{1 - \lambda \|A - A_0\|} \|y_1 - y_2\|,$$

for each $y_1, y_2 \in W$.

Next we establish a corollary to Lemma 2, which will have important role in the sequel. A similar result has been obtained by S. P. Dokov and A. L. Dontchev, see lemma on pag. 119 of [9], for studying the local quadratic convergence of Newton's method for variational inequality.

Corollary 3. *Let X, Y be Banach spaces, Ω be an open nonempty subset of X , $f : \Omega \rightarrow Y$ be continuous with Fréchet differentiable f' continuous, and $F : X \rightrightarrows Y$ be a set-valued mapping. Suppose that $\bar{x} \in \Omega$ and $f + F$ is strongly regular at \bar{x} for 0 with modulus $\lambda > 0$. Then, there exist constants $r_{\bar{x}} > 0$ and $r_0 > 0$ such that, for any $x \in B(\bar{x}, r_{\bar{x}})$, there hold $\lambda \|f'(x) - f'(\bar{x})\| < 1$, the mapping $z \mapsto L_f(x, z)^{-1} \cap B(\bar{x}, r_{\bar{x}})$ is single-valued from $B(0, r_0)$ to $B(\bar{x}, r_{\bar{x}})$ and Lipschitzian on $B(0, r_0)$ as follows*

$$\|L_f(x, u)^{-1} \cap B(\bar{x}, r_{\bar{x}}) - L_f(x, v)^{-1} \cap B(\bar{x}, r_{\bar{x}})\| \leq \frac{\lambda \|u - v\|}{1 - \lambda \|f'(x) - f'(\bar{x})\|},$$

$\forall u, v \in B(0, r_0)$.

Proof. Since $f + F$ is strongly regular at \bar{x} for 0 with modulus $\lambda > 0$, thus \bar{x} is also solution of the generalized equation

$$0 \in L_f(\bar{x}, x) = f'(\bar{x})x + f(\bar{x}) - f'(\bar{x})\bar{x} + F(x),$$

and the mapping $x \mapsto f'(\bar{x})x + f(\bar{x}) - f'(\bar{x})\bar{x} + F(x)$ from X to Y is strongly regular at \bar{x} for 0 with the same modulus $\lambda > 0$. Thus, applying first part of Lemma 2 with $A_0 = f'(\bar{x})$, $a_0 = f(\bar{x}) - f'(\bar{x})\bar{x}$ and $G = F$, we conclude that there exist neighborhoods M of $f'(\bar{x})$ in $\mathcal{L}(X, Y)$, N of $f(\bar{x}) - f'(\bar{x})\bar{x}$ in X and $W = B(0, r_0)$ of origin in Y , and $U = B(\bar{x}, r_{\bar{x}}) \subset \Omega$ of \bar{x} , where $r_{\bar{x}} > 0$ and $r_0 > 0$, such that, for any $A \in M$ and $a \in N$, letting $T(A, a, \cdot) : B(\bar{x}, r_{\bar{x}}) \rightrightarrows Y$ be defined as

$$T(A, a, y) := Ay + a + F(y), \quad (4)$$

the mapping $z \mapsto T(A, a, z)^{-1} \cap B(\bar{x}, r_{\bar{x}})$ is a single-valued function from $B(0, r_0)$ to $B(\bar{x}, r_{\bar{x}})$. On the other hand, due to f be continuous with Fréchet differentiable f' continuous on Ω , we can shrink $r_{\bar{x}}$, if necessary, such that

$$\lambda \|f'(x) - f'(\bar{x})\| < 1, \quad f'(x) \in M, \quad f(x) - f'(x)x \in N, \quad \forall x \in B(\bar{x}, r_{\bar{x}}). \quad (5)$$

Since Definition 1 and (4) imply that $L_f(x, y) = T(f'(x), f(x) - f'(x)x, y)$, for all $y, x \in B(\bar{x}, r_{\bar{x}})$, after some manipulations we have, for each $z \in B(0, r_0)$,

$$L_f(x, z)^{-1} \cap B(\bar{x}, r_{\bar{x}}) = T(f'(x), f(x) - f'(x)x, z)^{-1} \cap B(\bar{x}, r_{\bar{x}}), \quad (6)$$

for each $x \in B(\bar{x}, r_{\bar{x}})$. Therefore, for each $x \in B(\bar{x}, r_{\bar{x}})$, the last equality and (5) imply that $z \mapsto L_f(x, z)^{-1} \cap B(\bar{x}, r_{\bar{x}})$ is single-valued from $B(0, r_0)$ to $B(\bar{x}, r_{\bar{x}})$, which prove the first part of corollary. Finally, taking into account (6) and second part of Lemma 2, we conclude that the mapping $z \mapsto L_f(x, z)^{-1} \cap B(\bar{x}, r_{\bar{x}})$ is Lipschitzian from $B(0, r_0)$ to $B(\bar{x}, r_{\bar{x}})$ with constant $\lambda/[1 - \lambda \|f'(x) - f'(\bar{x})\|]$, which conclude the proof. \square

3 Local Convergence of the Newton method

In this section, we state our main result. We present an analysis of the behavior of the sequence generated by Newton's method for solving the generalized equation (1). For this purpose, we suppose that $f + F$ is strongly regular at \bar{x} for 0 with modulus $\lambda > 0$, where \bar{x} is such that $f(\bar{x}) + F(\bar{x}) \ni 0$. Moreover, the Lipschitz continuity of f' is relaxed, i.e., we assume that f' satisfies the conditions of the next definition.

Definition 3. Let X, Y be Banach space, Ω be an open nonempty subset of X , $f : \Omega \rightarrow Y$ be continuous with Fréchet derivative f' continuous in Ω . Let $\bar{x} \in \Omega$ and $R > 0$ and $\kappa := \sup\{t \in [0, R) : B(\bar{x}, t) \subset \Omega\}$. A twice continuously differentiable function $\psi : [0, R) \rightarrow \mathbb{R}$ is a majorant function for f on $B(\bar{x}, R)$ with modulus $\lambda > 0$ if it satisfies

$$\lambda \|f'(x) - f'(\bar{x} + \tau(x - \bar{x}))\| \leq \psi'(\|x - \bar{x}\|) - \psi'(\tau\|x - \bar{x}\|), \quad (7)$$

for all $\tau \in [0, 1]$, $x \in B(\bar{x}, \kappa)$ and, moreover, there hold:

h1) $\psi(0) = 0$ and $\psi'(0) = -1$;

h2) ψ' is strictly increasing.

The statement of the our main result is:

Theorem 4. Let X, Y be Banach spaces, Ω be an open nonempty subset of X , $f : \Omega \rightarrow Y$ be continuous with Fréchet derivative f' continuous in Ω , $F : X \rightrightarrows Y$ be a set-valued mapping and $\bar{x} \in \Omega$. Suppose that $f + F$ is strongly regular at \bar{x} for 0 with modulus $\lambda > 0$. Let $R > 0$ and assume that $\psi : [0, R) \rightarrow \mathbb{R}$ is a majorant function for f on $B(\bar{x}, R)$ with modulus $\lambda > 0$. Let $\nu := \sup\{t \in [0, R) : \psi'(t) < 0\}$, $\rho := \sup\{t \in (0, \nu) : \psi(t)/(t\psi'(t)) - 1 < 1\}$ and $r := \min\{\kappa, \rho\}$. Then, there exists a convergence radius $r_{\bar{x}} > 0$ with $r_{\bar{x}} \leq r$ such that the sequences with starting point $x_0 \in B(\bar{x}, r_{\bar{x}}) \setminus \{\bar{x}\}$ and $t_0 = \|\bar{x} - x_0\|$, respectively,

$$0 \in f(x_k) + f'(x_k)(x_{k+1} - x_k) + F(x_{k+1}), \quad t_{k+1} = |t_k - \psi(t_k)/\psi'(t_k)|, \quad (8)$$

$k = 0, 1, \dots$, are well defined, $\{t_k\}$ is strictly decreasing, is contained in $(0, r)$ and converges to 0, $\{x_k\}$ is contained in $B(\bar{x}, r_{\bar{x}})$ and converges to the point \bar{x} which is the unique solution of the generalized equation $f(x) + F(x) \ni 0$ in $B(\bar{x}, \bar{\sigma})$, where $0 < \bar{\sigma} \leq \min\{r_{\bar{x}}, \sigma\}$ and $\sigma := \sup\{0 < t < \kappa : \psi(t) < 0\}$ and there hold

$$\lim_{k \rightarrow \infty} \frac{\|x_{k+1} - \bar{x}\|}{\|x_k - \bar{x}\|} = 0, \quad \lim_{k \rightarrow \infty} \frac{t_{k+1}}{t_k} = 0. \quad (9)$$

Moreover, given $0 \leq p \leq 1$ and assume that

h3) the function $(0, \nu) \ni t \mapsto [\psi(t)/\psi'(t) - t]/t^{p+1}$ is strictly increasing,

then the sequence $\{t_{k+1}/t_k^{p+1}\}$ is strictly decreasing and there holds

$$\|x_{k+1} - \bar{x}\| \leq \frac{t_{k+1}}{t_k^{p+1}} \|x_k - \bar{x}\|^{p+1}, \quad k = 0, 1, \dots \quad (10)$$

If, additionally, $\psi(\rho)/(\rho\psi'(\rho)) - 1 = 1$ and $\rho < \kappa$, then $r_{\bar{x}} = \rho$ is the biggest convergence radius.

Remark 2. The first equation in (9) means that $\{x^k\}$ converges superlinearly to \bar{x} . Note that always ψ has derivative ψ' convex, condition **h3** holds with $p = 1$. In this case, there holds

$$t_{k+1}/t_k^2 \leq [\psi''(t_0)]/[2|\psi'(t_0)|], \quad k = 0, 1, \dots$$

and $\{x^k\}$ converges quadratically. Indeed, convexity of ψ' is necessary to obtain the quadratic convergence; see Example 2 in [18]. Moreover, as $\{t_{k+1}/t_k^{p+1}\}$ is strictly decreasing we have $t_{k+1}/t_k^{p+1} \leq t_1/t_0^{p+1}$, for $k = 0, 1, \dots$. Thus, (10) implies $\|x_{k+1} - \bar{x}\| \leq [t_1/t_0^{p+1}] \|x_k - \bar{x}\|^{p+1}$, for $k = 0, 1, \dots$. Consequently, if $p = 0$ then $\|x_k - \bar{x}\| \leq t_0[t_1/t_0]^k$ for $k = 0, 1, \dots$ and if $0 < p \leq 1$ then

$$\|x_k - \bar{x}\| \leq t_0(t_1/t_0)^{[(p+1)^k - 1]/p}, \quad k = 0, 1, \dots$$

Remark 3. Note that throughout the proof of the above theorem, if we assume that $F \equiv \{0\}$ then the constant $r_{\bar{x}} = \nu$. In this case, Theorem (4) merges into Theorem 2 of [18].

From now on, we assume that the hypotheses of Theorem 4 hold, with the exception of **h3**, which will be considered to hold only when explicitly stated.

3.1 Preliminary results

In this section, our first goal is to prove all statements in Theorem 4 concerning the sequence $\{t_k\}$ associated to the majorant function ψ defined in (8). Moreover, we obtain some relationships between the majorant function ψ and the set-valued mapping $f + F$, which will play an important role throughout the paper. Furthermore, the results in Theorem 4 related to the uniqueness of the solution and the optimal convergence radius will be proved. We begin with some observations on the majorant function.

As proven in Proposition 2.5 of [17], the constants κ , ν and σ , defined in Definition 3 and Theorem 4, are all positives and $t - \psi(t)/\psi'(t) < 0$, for all $t \in (0, \nu)$. According to **h2** and definition of ν , we have $\psi'(t) < 0$, for all $t \in [0, \nu)$. Therefore, the Newton iteration map for ψ is well defined in $[0, \nu)$, namely, $n_\psi : [0, \nu) \rightarrow (-\infty, 0]$ is defined by

$$n_\psi(t) := t - \psi(t)/\psi'(t), \quad t \in [0, \nu). \quad (11)$$

The next proposition was proved in Proposition 4 of [18].

Proposition 5. $\lim_{t \rightarrow 0} |n_\psi(t)|/t = 0$ and the constant ρ is positive. As a consequence, $|n_\psi(t)| < t$ for all $t \in (0, \rho)$.

Using (11), it is easy to see that the sequence $\{t_k\}$ is equivalently defined as

$$t_0 = \|\bar{x} - x_0\|, \quad t_{k+1} = |n_\psi(t_k)|, \quad k = 0, 1, \dots \quad (12)$$

Next result, which is a consequence of above proposition, contains the main convergence properties of the above sequence and its prove can be found in Corollary 5 of [18].

Corollary 6. The sequence $\{t_k\}$ is well defined, is strictly decreasing and is contained in $(0, \rho)$. Moreover, $\{t_k\}$ converges to 0 with superlinear rate, i.e.,

$$\lim_{k \rightarrow \infty} t_{k+1}/t_k = 0.$$

If additionally **h3** holds, then the sequence t_{k+1}/t_k^{p+1} is strictly decreasing.

In the sequel we study the *linearization error of the function f* at a point in Ω , namely,

$$E_f(x, y) := f(y) - [f(x) + f'(x)(y - x)], \quad y, x \in \Omega. \quad (13)$$

We show that this error is bounded by the linearization error of the majorant function ψ , i.e.,

$$e_\psi(t, u) := \psi(u) - [\psi(t) + \psi'(t)(u - t)], \quad t, u \in [0, R),$$

and as consequence, we prove that the partial linearization of $f + F$ has a single-valued inverse, which is Lipschitz in a neighborhood of \bar{x} .

Lemma 7. There holds $\lambda \|E_f(x, \bar{x})\| \leq e_\psi(\|x - \bar{x}\|, 0)$, for all $x \in B(\bar{x}, \kappa)$.

Proof. Since $\bar{x} + (1 - u)(x - \bar{x}) \in B(\bar{x}, \kappa)$, for all $0 \leq u \leq 1$ and f is continuously differentiable in Ω , thus the definition of E_f and some simple manipulations yield

$$\lambda \|E_f(x, \bar{x})\| \leq \int_0^1 \lambda \|f'(x) - f'(\bar{x} + (1 - u)(x - \bar{x}))\| \|\bar{x} - x\| \, du.$$

Combining last inequality with (7) and then performing the integral obtained and using the definition of e_ψ , the statement follows. \square

The next result states that, if a generalized equation (1) is strongly regular at \bar{x} for 0 with modulus $\lambda > 0$ and (7) holds, then there exists a neighborhood of \bar{x} such that, for all x in this neighborhood, (1) is also strongly regular at x for 0 with modulus $\lambda/(\|\psi'(\|x - \bar{x}\|)\|)$. The result is a consequence of Corollary 3 and its statement is:

Lemma 8. *There exists a constant $r_{\bar{x}} \leq r$ such that, the mapping*

$$x \mapsto L_f(x, 0)^{-1} \cap B(\bar{x}, r_{\bar{x}})$$

is single-valued in $B(\bar{x}, r_{\bar{x}})$ and there holds

$$\|\bar{x} - L_f(x, 0)^{-1} \cap B(\bar{x}, r_{\bar{x}})\| \leq \frac{\lambda}{|\psi'(\|x - \bar{x}\|)|} \|E_f(x, \bar{x})\|, \quad \forall x \in B(\bar{x}, r_{\bar{x}}).$$

Proof. Take $x \in B(\bar{x}, r)$. Since $r < \nu$ we have $\|x - \bar{x}\| < \nu$. Thus, $\psi'(\|x - \bar{x}\|) < 0$ which, together (7) and **h1**, imply that

$$\lambda \|f'(x) - f'(\bar{x})\| \leq \psi'(\|x - \bar{x}\|) - \psi'(0) < -\psi'(0) = 1, \quad \forall x \in B(\bar{x}, r). \quad (14)$$

Due to $f + F$ be strongly regular at \bar{x} for 0 with modulus $\lambda > 0$, we can apply Corollary 3 to obtain $r_{\bar{x}} > 0$ and $r_0 > 0$ such that, for any $x \in B(\bar{x}, r_{\bar{x}})$, the mapping $z \mapsto L_f(x, z)^{-1} \cap B(\bar{x}, r_{\bar{x}})$ is single-valued from $B(0, r_0)$ to $B(\bar{x}, r_{\bar{x}})$. In particular, we conclude that the mapping $x \mapsto L_f(x, 0)^{-1} \cap B(\bar{x}, r_{\bar{x}})$ is single-valued in $B(\bar{x}, r_{\bar{x}})$. Moreover, Corollary 3 implies that $\forall u, v \in B(0, r_0)$

$$\|L_f(x, u)^{-1} \cap B(\bar{x}, r_{\bar{x}}) - L_f(x, v)^{-1} \cap B(\bar{x}, r_{\bar{x}})\| \leq \frac{\lambda \|u - v\|}{1 - \lambda \|f'(x) - f'(\bar{x})\|}.$$

If necessary, we shrink $r_{\bar{x}}$ such that $r_{\bar{x}} \leq r$, in order to combine the last inequality with the first inequality in (14) and **h1**, to conclude that, for all $x \in B(\bar{x}, r_{\bar{x}})$ there holds

$$\|L_f(x, u)^{-1} \cap B(\bar{x}, r_{\bar{x}}) - L_f(x, v)^{-1} \cap B(\bar{x}, r_{\bar{x}})\| \leq \frac{\lambda \|u - v\|}{|\psi'(\|x - \bar{x}\|)|}, \quad (15)$$

for each $u, v \in B(0, r_0)$. On the other hand, due to f be continuous with f' continuous in Ω , we have $\lim_{x \rightarrow \bar{x}} E_f(x, \bar{x}) = 0$. Thus, we can shrink $r_{\bar{x}}$, if necessary, such that

$$E_f(x, \bar{x}) \in B(0, r_0), \quad \forall x \in B(\bar{x}, r_{\bar{x}}).$$

Let $x \in B(\bar{x}, r_{\bar{x}})$. Note that, after some algebraic manipulations we obtain

$$\begin{aligned} 0 \in f(\bar{x}) + F(\bar{x}) &= f(x) + f'(x)(\bar{x} - x) - f(x) - f'(x)(\bar{x} - x) + f(\bar{x}) + F(\bar{x}) \\ &= f(x) + f'(x)(\bar{x} - x) + E_f(x, \bar{x}) + F(\bar{x}). \end{aligned}$$

Thus, $-E_f(x, \bar{x}) \in L_f(x, \bar{x}) = f(x) + f'(x)(\bar{x} - x) + F(\bar{x})$. Since $E_f(x, \bar{x}) \in B(0, r_0)$ and the mapping $z \mapsto L_f(x, z)^{-1} \cap B(\bar{x}, r_{\bar{x}})$ is single-valued from $B(0, r_0)$ to $B(\bar{x}, r_{\bar{x}})$ we conclude that

$$\bar{x} = L_f(x, -E_f(x, \bar{x}))^{-1} \cap B(\bar{x}, r_{\bar{x}}).$$

Therefore, substituting $u = -E_f(x, \bar{x})$ and $v = 0$ into (15) the desired inequality follows. \square

Let $r_{\bar{x}} > 0$ the constant given by Lemma 8. Lemma 8 guarantees, in particular, that the mapping $x \mapsto L_f(x, 0)^{-1} \cap B(\bar{x}, r_{\bar{x}})$ is single-valued in $B(\bar{x}, r_{\bar{x}})$ and consequently, the Newton iteration mapping is well-defined. Let us call N_{f+F} , the Newton iteration mapping for $f + F$ in that region, namely, $N_{f+F} : B(\bar{x}, r_{\bar{x}}) \rightarrow X$ is defined by

$$N_{f+F}(x) := L_f(x, 0)^{-1} \cap B(\bar{x}, r_{\bar{x}}), \quad \forall x \in B(\bar{x}, r_{\bar{x}}). \quad (16)$$

Using (3), the definition of Newton iteration mapping in (16) is equivalent to

$$0 \in f(x) + f'(x)(N_{f+F}(x) - x) + F(N_{f+F}(x)), \quad N_{f+F}(x) \in B(\bar{x}, r_{\bar{x}}), \quad (17)$$

for each $x \in B(\bar{x}, r_{\bar{x}})$. Therefore, since Lemma 8 guarantees that $N_{f+F}(x)$ is single-valued at $B(\bar{x}, r_{\bar{x}})$, see (16), we can apply a *single* Newton iteration for any $x \in B(\bar{x}, r_{\bar{x}})$ to obtain $N_{f+F}(x)$ which may not belong to $B(\bar{x}, r_{\bar{x}})$, or even may not belong to the domain of f . Thus, this allow us to guarantee the well-definedness of only one iteration of Newton's method. In particular, the next result shows that for any $x \in B(\bar{x}, r_{\bar{x}})$, the Newton iterations, see (17), may be repeated indefinitely.

Lemma 9. *If $\|x - \bar{x}\| \leq t < r_{\bar{x}}$ then $\|N_{f+F}(x) - \bar{x}\| \leq |n_\psi(\|x - \bar{x}\|)|$. As a consequence, $N_{f+F}(B(\bar{x}, r_{\bar{x}})) \subset B(\bar{x}, r_{\bar{x}})$. Moreover, if **h3** holds and $x \neq \bar{x}$ then $\|N_{f+F}(x) - \bar{x}\| \leq [|n_\psi(t)|/t^{p+1}]\|x - \bar{x}\|^{p+1}$.*

Proof. Since $0 \in f(\bar{x}) + F(\bar{x})$ we have $\bar{x} = N_{f+F}(\bar{x})$. Thus, the inequality of the lemma is trivial for $x = \bar{x}$. Now, assume that $0 < \|x - \bar{x}\| \leq t$. Hence, Lemma 8 implies that the mapping $x \mapsto L_f(x, 0)^{-1} \cap B(\bar{x}, r_{\bar{x}})$ is single-valued in $B(\bar{x}, r_{\bar{x}})$ and Lipschitz continuous with modulus $\lambda/|\psi'(\|x - \bar{x}\|)|$. Using (16) and Lemma 8, it is easy to conclude that

$$\|\bar{x} - N_{f+F}(x)\| \leq \frac{\lambda}{|\psi'(\|x - \bar{x}\|)|} \|E_f(x, \bar{x})\|.$$

Now, applying Lemma 7 we obtain

$$\|\bar{x} - N_{f+F}(x)\| \leq \frac{e_\psi(\|x - \bar{x}\|, 0)}{|\psi'(\|x - \bar{x}\|)|}.$$

On the other hand, taking into account that $\psi(0) = 0$, the definitions of e_ψ and n_ψ imply that

$$\frac{e_\psi(\|x - \bar{x}\|, 0)}{|\psi'(\|x - \bar{x}\|)|} = \frac{\psi(\|x - \bar{x}\|)}{\psi'(\|x - \bar{x}\|)} - \|x - \bar{x}\| = |n_\psi(\|x - \bar{x}\|)|.$$

Hence, the first statement follows by combining the above two expressions. For proving the inclusion of the lemma, take $x \in B(\bar{x}, r_{\bar{x}})$. Since $\|x - \bar{x}\| < r_{\bar{x}}$, $r_{\bar{x}} \leq \rho$ and $\|N_{f+F}(x) - \bar{x}\| \leq |n_\psi(\|x - \bar{x}\|)|$, thus using the second part of Proposition 5 we conclude that $\|N_{f+F}(x) - \bar{x}\| < \|x - \bar{x}\|$ which prove the inclusion.

In the following, we prove last inequality. Due $0 \in f(\bar{x}) + F(\bar{x})$, the inequality is trivial for $x = \bar{x}$. If $0 < \|x - \bar{x}\| \leq t$ then assumption **h3** and (11) yields

$$\frac{|n_\psi(\|x - \bar{x}\|)|}{\|x - \bar{x}\|^{p+1}} < \frac{|n_\psi(t)|}{t^{p+1}}.$$

Therefore, using the first part of Lemma 9 the inequality follows. \square

In the next result we obtain the uniqueness of the solution in the neighborhood $B[\bar{x}, \sigma]$.

Lemma 10. *There exists a constant $\bar{\sigma} \leq \min\{r_{\bar{x}}, \sigma\}$ such that \bar{x} is the unique solution of (1) in $B[\bar{x}, \bar{\sigma}]$.*

Proof. Let $r_{\bar{x}} > 0$ the constant given by Lemma 8. Thus, Corollary 3 implies that there exists $r_0 > 0$ such that, for any $x \in B(\bar{x}, r_{\bar{x}})$, the mapping $z \mapsto L_f(x, z)^{-1} \cap B(\bar{x}, r_{\bar{x}})$ is single-valued from $B(0, r_0)$ to $B(\bar{x}, r_{\bar{x}})$ and there holds

$$\|L_f(x, u)^{-1} \cap B(\bar{x}, r_{\bar{x}}) - L_f(x, v)^{-1} \cap B(\bar{x}, r_{\bar{x}})\| \leq \frac{\lambda \|u - v\|}{1 - \lambda \|f'(x) - f'(\bar{x})\|},$$

for each $u, v \in B(0, r_0)$. Now, due to f be continuous, we have

$$\lim_{x \rightarrow \bar{x}} E_f(\bar{x}, x) = 0.$$

Thus, we can take $\bar{\sigma} \leq \min\{r_{\bar{x}}, \sigma\}$, such that

$$E_f(\bar{x}, y) \in B(0, r_0).$$

Let $y \in B(\bar{x}, \bar{\sigma})$ and assume that $0 \in f(y) + F(y)$. Then, some manipulations yield

$$\begin{aligned} 0 \in f(y) + F(y) &= f(y) - f(\bar{x}) - f'(\bar{x})(y - \bar{x}) + f(\bar{x}) + f'(\bar{x})(y - \bar{x}) + F(y) \\ &= E_f(\bar{x}, y) + L_f(\bar{x}, y), \end{aligned}$$

which implies that $-E_f(\bar{x}, y) \in L_f(\bar{x}, y)$. Since the mapping $z \mapsto L_f(\bar{x}, z)^{-1} \cap B(\bar{x}, r_{\bar{x}})$ is single-valued from $B(0, r_0)$ to $B(\bar{x}, r_{\bar{x}})$, we have

$$y = L_f(\bar{x}, -E_f(\bar{x}, y))^{-1} \cap B(\bar{x}, r_{\bar{x}}), \quad \bar{x} = L_f(\bar{x}, 0)^{-1} \cap B(\bar{x}, r_{\bar{x}}).$$

Thus, substituting into above inequality $x = \bar{x}$, $u = 0$ and $v = -E_f(\bar{x}, y)$, we conclude that

$$\|\bar{x} - y\| = \|L_f(\bar{x}, 0)^{-1} \cap B(\bar{x}, r_{\bar{x}}) - L_f(\bar{x}, -E_f(\bar{x}, y))^{-1} \cap B(\bar{x}, r_{\bar{x}})\| \leq \lambda \|E_f(\bar{x}, y)\|.$$

Using definition on (13) and (7) with $x = \bar{x} + u(y - \bar{x})$ and $\tau = 0$, last inequality implies

$$\begin{aligned} \|\bar{x} - y\| &\leq \lambda \|f(y) - f(\bar{x}) - f'(\bar{x})(y - \bar{x})\| \\ &\leq \int_0^1 \lambda \|f'(\bar{x} + u(y - \bar{x})) - f'(\bar{x})\| \|y - \bar{x}\| du \\ &\leq \int_0^1 [\psi'(u\|y - \bar{x}\|) - \psi'(0)] \|y - \bar{x}\| du. \end{aligned}$$

Performing the integral of the right hand side of the above inequality we have $0 \leq \psi(\|y - \bar{x}\|)$, which implies that $\psi(\|y - \bar{x}\|) = 0$ due to $\psi(t) < 0$ for $t \in (0, \sigma)$ and $\|y - \bar{x}\| \leq \sigma$. Since $0 \leq \|y - \bar{x}\| \leq \sigma$ and 0 is the unique zero of ψ in $[0, \sigma]$, we conclude that $\|y - \bar{x}\| = 0$, i.e., $y = \bar{x}$ and \bar{x} is the unique solution of (1) in $B[\bar{x}, \bar{\sigma}]$. \square

The next result gives the biggest convergence radius, its proof is similar to the proof of Lemma 2.15 of [17].

Lemma 11. *If $\psi(\rho)/(\rho\psi'(\rho)) - 1 = 1$ and $\rho < \kappa$, then $r_{\bar{x}} = \rho$ is the biggest convergence radius.*

3.2 Proof of Theorem 4

First, note that the inclusion in (8) together (16) and (17) imply that the sequence $\{x_k\}$ satisfies

$$x_{k+1} = N_{f+F}(x_k), \quad k = 0, 1, \dots, \quad (18)$$

which is indeed an equivalent definition of this sequence.

Proof. All statements involving $\{t_k\}$ were proved in Corollary 6. Since Lemma 8 and (16) implies that there exist constants $r_{\bar{x}} > 0$ and $r_0 > 0$ such that $r_{\bar{x}} \leq r$ and, for any $x \in B(\bar{x}, r_{\bar{x}})$, the mapping N_{f+F} is single-valued in $B(\bar{x}, r_{\bar{x}})$. Thus, taking into account that $x_0 \in B(\bar{x}, r_{\bar{x}})$, we conclude by combining (18) and inclusion $N_{f+F}(B(\bar{x}, r_{\bar{x}})) \subset B(\bar{x}, r_{\bar{x}})$ in Lemma 9 that $\{x_k\}$ is well defined and remains in $B(\bar{x}, r_{\bar{x}})$. Now, we are going to prove that $\{x_k\}$ converges towards \bar{x} . Without lose generality we assume that the sequence $\{x_k\}$ is infinity. Since $0 < \|x_k - \bar{x}\| < r_{\bar{x}} \leq \rho$, for $k = 0, 1, \dots$, we obtain from (18), Lemma 9 and second part of Proposition 5 that

$$\|x_{k+1} - \bar{x}\| \leq |n_\psi(\|x_k - \bar{x}\|)| < \|x_k - \bar{x}\|, \quad k = 0, 1, \dots \quad (19)$$

Thus, $\{\|x_k - \bar{x}\|\}$ is strictly decreasing and convergent. Let $\bar{\alpha} = \lim_{k \rightarrow \infty} \|x_k - \bar{x}\|$. Because $\{\|x_k - \bar{x}\|\}$ rest in $(0, \rho)$ and it is strictly decreasing we have $0 \leq \bar{\alpha} < \rho$. Then, by continuity of n_ψ and (19) imply $0 \leq \bar{\alpha} = |n_\psi(\bar{\alpha})|$, and from second part of Proposition 5 we have $\bar{\alpha} = 0$. Therefore, the convergence of $\{x_k\}$ to \bar{x} is proved. Now we are going show that \bar{x} is a solution of the generalized equation $f(x) + F(x) \ni 0$. From inclusion in (8) we conclude

$$(x_{k+1}, -f(x_k) - f'(x_k)(x_{k+1} - x_k)) \in \text{gph } F, \quad k = 0, 1, \dots$$

By assumption the set-valued mapping F has closed graph and f is continuous with f' continuous, thus last inclusion implies that

$$\lim_{k \rightarrow \infty} ((x_{k+1}, -f(x_k) - f'(x_k)(x_{k+1} - x_k)) = (\bar{x}, -f(\bar{x})) \in \text{gph } F,$$

which implies $f(\bar{x}) + F(\bar{x}) \ni 0$. Now, we are going to show the first inequality in (9). Note that (19) implies

$$\frac{\|x_{k+1} - \bar{x}\|}{\|x_k - \bar{x}\|} \leq \frac{|n_\psi(\|x_k - \bar{x}\|)|}{\|x_k - \bar{x}\|}, \quad k = 0, 1, \dots$$

Since $\lim_{k \rightarrow \infty} \|x_k - \bar{x}\| = 0$, the desired equality follows from the first statement in Proposition 5. To prove (10), firstly we will show by induction that the sequences $\{x_k\}$ and $\{t_k\}$ defined in (8) satisfy

$$\|x_k - \bar{x}\| \leq t_k, \quad k = 0, 1, \dots \quad (20)$$

Since $t_0 = \|x_0 - \bar{x}\|$, the above inequality holds to $k = 0$. Now, we assume that $\|x_k - \bar{x}\| \leq t_k$ holds. Using (18), second part of Lemma 9, the induction assumption and (12) we have

$$\|x_{k+1} - \bar{x}\| = \|N_{f+F}(x_k) - \bar{x}\| \leq \frac{|n_\psi(t_k)|}{t_k^{p+1}} \|x_k - \bar{x}\|^{p+1} = \frac{t_{k+1}}{t_k^{p+1}} \|x_k - \bar{x}\|^{p+1} \leq t_{k+1},$$

and the proof by induction is complete. Thus, the inequality (10) follows by combination of (20) and second part of Lemma 9. Finally, the uniqueness follows from Lemma 10 and the last statement in the theorem follows from Lemma 11. \square

4 Particular cases

In this section, some special cases of Theorem 4 will be considered. For instance, if $F \equiv \{0\}$ and f' satisfies a Hölder-type condition, a particular instance of Theorem 4, which retrieves the classical convergence theorem on Newton's method under the Lipschitz condition will be obtained; see [28, 34]. We also obtain Theorem 1 of N. H. Josephy in [23] and, up to some minor adjustments, Theorem 1 of A. L. Dontchev [10]. To complete this section, a version of Smale's theorem on Newton's method for analytical functions is proved in Theorem 15.

4.1 Under Hölder-type condition

The next result, which is a consequence of our main result Theorem 4, is a version of classical convergence theorem for Newton's method under Hölder-type condition for solving generalized equations of the type (1). Classical versions for $F \equiv \{0\}$ have appeared in [22, 24, 28, 34].

Theorem 12. *Let X, Y be Banach spaces, $\Omega \subseteq X$ an open set and $f : \Omega \rightarrow Y$ be continuous with Fréchet derivative f' continuous in Ω , $F : X \rightrightarrows Y$ be a set-valued mapping with closed graph and $\bar{x} \in \Omega$. Suppose that $f + F$ is strongly regular at \bar{x} for 0 with modulus $\lambda > 0$ and there exist constants $K > 0$ and $0 < p \leq 1$ such that*

$$\lambda \|f'(x) - f'(\bar{x} + \tau(x - \bar{x}))\| \leq (K - \tau^p) \|x - \bar{x}\|^p, \quad x \in B(\bar{x}, \kappa), \quad \tau \in [0, 1]. \quad (21)$$

Let $r := \min\{\kappa, [(p+1)/((2p+1)K)]^{1/p}\}$, where $\kappa := \sup\{t > 0 : B(\bar{x}, t) \subset \Omega\}$. Then, there exists a convergence radius $r_{\bar{x}} > 0$ with $r_{\bar{x}} \leq r$ such that the sequences with starting point $x_0 \in B(\bar{x}, r_{\bar{x}}) \setminus \{\bar{x}\}$ and $t_0 = \|\bar{x} - x_0\|$, respectively,

$$f(x_k) + f'(x_k)(x_{k+1} - x_k) + F(x_{k+1}) \ni 0, \quad t_{k+1} = \frac{Kpt_k^{p+1}}{(p+1)[1 - Kt_k^p]}, \quad (22)$$

$k = 0, 1, \dots$, are well defined, $\{t_k\}$ is strictly decreasing, is contained in $(0, r)$ and converges to 0, $\{x_k\}$ is contained in $B(\bar{x}, r_{\bar{x}})$ and converges to the point \bar{x} which is a unique solution of $f(x) + F(x) \ni 0$ in $B(\bar{x}, \bar{\sigma})$, where $\bar{\sigma} \leq \min\{r_{\bar{x}}, [(p+1)/K]^{1/p}\}$. Moreover, $\{t_{k+1}/t_k^2\}$ is strictly decreasing, $t_{k+1}/t_k^2 < 1/[2/K - 2\|\bar{x} - x_0\|]$ and

$$\|\bar{x} - x_{k+1}\| \leq \frac{Kp\|x_k - \bar{x}\|^{p+1}}{(p+1)[1 - Kt_k^p]} \leq \frac{Kp\|x_k - \bar{x}\|^{p+1}}{(p+1)[1 - K\|x_0 - \bar{x}\|^p]}, \quad k = 0, 1, \dots$$

If, additionally, $[(p+1)/(2p+1)K]^{1/p} < \kappa$, then $r_{\bar{x}} = [(p+1)/(2p+1)K]^{1/p}$ is the biggest convergence radius.

Proof. Using condition in (21), we can immediately prove that f, \bar{x} and $\psi : [0, \kappa) \rightarrow \mathbb{R}$, defined by $\psi(t) = Kt^{p+1}/(p+1) - t$, satisfy the inequality (7) and the conditions **h1**, **h2** and **h3** in Theorem 4. In this case, it is easy to see that ρ and ν , as defined in Theorem 4, satisfy $\rho = [(p+1)/(2p+1)K]^{1/p} \leq \nu = [1/K]^{1/p}$ and, as a consequence, $r := \min\{\kappa, [(p+1)/((2p+1)K)]^{1/p}\}$. Moreover, $\psi(\rho)/(\rho\psi'(\rho)) - 1 = 1$, $\psi(0) = \psi([(p+1)/K]^{1/p}) = 0$ and $\psi(t) < 0$ for all $t \in (0, [(p+1)/K]^{1/p})$. Also, the sequence $\{t_k\}$ in Theorem 4 is given by (22) and satisfies

$$t_{k+1}/t_k^2 = \frac{Kp}{(p+1)[1 - Kt_k^p]} < \frac{Kp}{(p+1)[1 - K\|x_0 - \bar{x}\|^p]}, \quad k = 0, 1, \dots$$

Therefore, the result follows by invoking Theorem 4. \square

Remark 4. *Theorem 12 contain, as particular instance, some results on Newton's method; as we can see in, Rall [28] and Traub and Wozniakowski [34].*

We are going to study the variational inequality problem, namely, the generalized equation associated to $F = N_C$ the normal cone of C a nonempty, closed and convex subset of Y ,

$$f(x) + N_C(x) \ni 0. \quad (23)$$

The next result is a version of classical convergence theorem for Newton's method under Lipschitz-type condition for the variational inequality (23), it has been prove by N. H. Josephy in [23].

Theorem 13. *Let X, Y be Banach spaces, C a nonempty, closed and convex subset of Y , $\Omega \subseteq X$ an open set and $f : \Omega \rightarrow Y$ be continuous with Fréchet derivative f' continuous in Ω such that*

$$\|f'(x) - f'(y)\| \leq L\|x - y\|, \quad x, y \in \Omega,$$

where $L > 0$. Let $\bar{x} \in \Omega$ and suppose that $f + N_C$ is strongly regular at \bar{x} for 0 with modulus $\lambda > 0$. Let $r = \min\{\kappa, 2/(3\lambda L)\}$, where $\kappa = \sup\{t \in [0, R) : B(\bar{x}, t) \subset \Omega\}$. Then, there exists a convergence radius $r_{\bar{x}} > 0$ with $r_{\bar{x}} \leq r$ such that the sequences with starting point $x_0 \in B(\bar{x}, r_{\bar{x}}) \setminus \{\bar{x}\}$ and $t_0 = \|\bar{x} - x_0\|$, respectively,

$$f(x_k) + f'(x_k)(x_{k+1} - x_k) + N_C(x_{k+1}) \ni 0, \quad t_{k+1} = ((\lambda L/2)t_k^2)/(1 - \lambda L t_k),$$

$k = 0, 1, \dots$, are well defined, $\{t_k\}$ is strictly decreasing, is contained in $(0, r)$ and converges to 0, $\{x_k\}$ is contained in $B(\bar{x}, r_{\bar{x}})$ and converges to the point \bar{x} which is a unique solution of $f(x) + N_C(x) \ni 0$ in $B(\bar{x}, \bar{\sigma})$, where $0 < \bar{\sigma} \leq \min\{r_{\bar{x}}, 2/K\}$. Moreover, $\{t_{k+1}/t_k^2\}$ is strictly decreasing, $t_{k+1}/t_k^2 < 1/[2/(\lambda L) - 2\|\bar{x} - x_0\|]$ and

$$\|\bar{x} - x_{k+1}\| \leq \frac{\lambda L}{2} \frac{1}{1 - \lambda L t_k} \|x_k - \bar{x}\|^2 \leq \frac{\lambda L}{2} \frac{1}{1 - \lambda L \|x_0 - \bar{x}\|} \|x_k - \bar{x}\|^2,$$

$k = 0, 1, \dots$. If, additionally, $2/(3\lambda L) < \kappa$, then $r_{\bar{x}} = 2/(3\lambda L)$ is the biggest convergence radius.

Proof. The result follows by applying Theorem 12 with $\tau = 0$, $p = 1$, $K = \lambda L$ and $F = N_C$. \square

A. L. Dontchev [10] under Aubin continuity of the mapping $L_f(\bar{x}, \cdot)^{-1} : Y \rightrightarrows X$, defined by

$$L_f(\bar{x}, z)^{-1} := \{y \in X : z \in f(\bar{x}) + f'(\bar{x})(y - \bar{x}) + N_C(y)\}, \quad (24)$$

has shown that the Newton's method for solving (23) generates a sequence that converges Q -quadratically to a solution. Now, our purpose is to show that, if $X = Y = \mathbb{R}^n$, $F = N_C$ and $C \subset \mathbb{R}^n$ is a nonempty, polyhedral and convex set, then this particular instance of Theorem 1 of [10] follows from Theorem 12. We begin with the formal definition of Aubin continuity; for more details see [13, 14]. First we need the following definitions: The *distance* from a point $v \in \mathbb{R}^n$ to a set $U \subset \mathbb{R}^n$ is $d(v, U) := \inf\{\|v - u\| : u \in U\}$ and the *excess* from the set U to the set V is $e(V, U) := \sup\{d(v, U) : v \in V\}$.

Definition 4. *A mapping $H : \mathbb{R}^m \rightrightarrows \mathbb{R}^n$ is said to be Aubin continuous, at $\bar{y} \in \mathbb{R}^m$ for $\bar{x} \in \mathbb{R}^n$ with modulus $\alpha \geq 0$, if $\bar{x} \in H(\bar{y})$ and there exist constants $a > 0$ and $b > 0$ such that*

$$e(H(y_1) \cap B(\bar{x}, a), H(y_2)) \leq \alpha \|y_1 - y_2\|, \quad \forall y_1, y_2 \in B(\bar{y}, b).$$

It has been shown in [13, Theorem 1] that if $C \subset \mathbb{R}^n$ is a polyhedral convex set, then Aubin continuity of $\mathcal{L}_f(\bar{x}, \cdot)^{-1}$ is equivalent to strong regularity of $f + N_C$. Next we state, with some adjustment, Theorem 1 of [10]; see also [11].

Theorem 14. *Let $C \subset \mathbb{R}^n$ be a polyhedral convex set, $\Omega \subseteq \mathbb{R}^n$ an open set and $f : \Omega \rightarrow Y$ be continuous with Fréchet derivative f' continuous in Ω such that*

$$\|f'(x) - f'(y)\| \leq L\|x - y\|, \quad \forall x, y \in \Omega,$$

where $L > 0$. Let $\bar{x} \in \Omega$ and suppose that $\mathcal{L}_f(\bar{x}, \cdot)^{-1} : \mathbb{R}^m \rightrightarrows \mathbb{R}^n$ defined in (24) is Aubin continuous at $0 \in \mathbb{R}^m$ for $\bar{x} \in \mathbb{R}^n$ with modulus $\alpha \geq 0$. Let $r := \min\{\kappa, 2/(3\lambda L)\}$, where $\kappa = \sup\{t \in [0, R) : B(\bar{x}, t) \subset \Omega\}$. Then, there exists a convergence radius $r_{\bar{x}} > 0$ with $r_{\bar{x}} \leq r$ such that the sequences with starting point $x_0 \in B(\bar{x}, r_{\bar{x}})/\{\bar{x}\}$ and $t_0 = \|\bar{x} - x_0\|$, respectively,

$$f(x_k) + f'(x_k)(x_{k+1} - x_k) + N_C(x_{k+1}) \ni 0, \quad t_{k+1} = ((\lambda L/2)t_k^2)/(1 - \lambda L t_k),$$

$k = 0, 1, \dots$, are well defined, $\{t_k\}$ is strictly decreasing, is contained in $(0, r)$ and converges to 0, $\{x_k\}$ is contained in $B(\bar{x}, r_{\bar{x}})$ and converges to the point \bar{x} which is a unique solution of $f(x) + N_C(x) \ni 0$ in $B(\bar{x}, \bar{\sigma})$, where $0 < \bar{\sigma} \leq \min\{r_{\bar{x}}, 2/K\}$. Moreover, $\{t_{k+1}/t_k^2\}$ is strictly decreasing, $t_{k+1}/t_k^2 < 1/[2/(\lambda L) - 2\|\bar{x} - x_0\|]$ and

$$\|\bar{x} - x_{k+1}\| \leq \frac{\lambda L}{2} \frac{1}{1 - \lambda L t_k} \|x_k - \bar{x}\|^2 \leq \frac{\lambda L}{2} \frac{1}{1 - \lambda L \|x_0 - \bar{x}\|} \|x_k - \bar{x}\|^2,$$

$k = 0, 1, \dots$. If, additionally, $2/(3\lambda L) < \kappa$, then $r_{\bar{x}} = 2/(3\lambda L)$ is the biggest convergence radius.

Proof. Since $C \subset \mathbb{R}^n$ is a polyhedral convex set, [13, Theorem 1] implies that Aubin continuity of $\mathcal{L}_f(\bar{x}, \cdot)^{-1}$ at $0 \in \mathbb{R}^m$ for $\bar{x} \in \mathbb{R}^n$ with modulus $\alpha \geq 0$, is equivalent to strong regularity of $f + N_C$ at \bar{x} for 0 with modulus $\alpha \geq 0$. Thus, the result follows by applying Theorem 13. \square

4.2 Under Smale's-type condition

In this section, we assume that f is an analytic function and using the ideas listed in [1], we present a version of the classical convergence theorem for Newton's method for solving the generalized equation (1). The classical version has appeared in corollary of Proposition 3 pp. 195 of Smale [33], see also Proposition 1 pp. 157 and Remark 1 pp. 158 of Blum, Cucker, Shub, and Smale [5]; see also [17]. For stating the result we need of the following definition.

Let $\Omega \subseteq X$ and $f : \Omega \rightarrow Y$ be an analytic function. The n -th derivative of f at x is a n -th multilinear map $f^n(x) : X \times \dots \times X \rightarrow X$ and its norm is defined by

$$\|f^n(x)\| = \sup\{\|f^n(x)(v_1, \dots, v_n)\| : v_1, \dots, v_n \in X, \|v_i\| \leq 1, i = 1, \dots, n\}.$$

Theorem 15. *Let X, Y be Banach spaces, $\Omega \subseteq X$ an open set and $f : \Omega \rightarrow Y$ be an analytic function, $F : X \rightrightarrows Y$ be a set-valued mapping with closed graph and $\bar{x} \in \Omega$. Suppose that $0 \in f(\bar{x}) + F(\bar{x})$ and $f + F$ is strongly regular at \bar{x} for 0 with modulus $\lambda > 0$. Suppose that*

$$\gamma := \sup_{n>1} \left\| \frac{\lambda f^{(n)}(\bar{x})}{n!} \right\|^{1/(n-1)} < +\infty. \quad (25)$$

Let $r := \min\{\kappa, (5 - \sqrt{17})/(4\gamma)\}$ the convergence radius, where

$$\kappa := \sup\{t > 0 : B(\bar{x}, t) \subset \Omega\}.$$

Then, there exists a convergence radius $r_{\bar{x}} > 0$ with $r_{\bar{x}} \leq r$ such that the sequences with starting point $x_0 \in B(\bar{x}, r_{\bar{x}}) \setminus \{\bar{x}\}$ and $t_0 = \|\bar{x} - x_0\|$, respectively

$$0 \in f(x_k) + f'(x_k)(x_{k+1} - x_k) + F(x_{k+1}), \quad t_{k+1} = \gamma t_k^2 / [2(1 - \gamma t_k)^2 - 1],$$

$k = 0, 1, \dots$, are well defined, $\{t_k\}$ is strictly decreasing, contained in $(0, r)$ and converges to 0, and $\{x_k\}$ is contained in $B(\bar{x}, r_{\bar{x}})$ and converges to the point \bar{x} which is the unique solution of $f(x) + F(x) \ni 0$ in $B(\bar{x}, \bar{\sigma})$, where $0 < \bar{\sigma} \leq \min\{r_{\bar{x}}, 1/(2\gamma)\}$. Moreover, $\{t_{k+1}/t_k^2\}$ is strictly decreasing, $t_{k+1}/t_k^2 < \gamma/[2(1 - \gamma\|x_0 - \bar{x}\|)^2 - 1]$, for $k = 0, 1, \dots$ and

$$\|x_{k+1} - \bar{x}\| \leq \frac{\gamma}{2(1 - \gamma t_k)^2 - 1} \|x_k - \bar{x}\|^2 \leq \frac{\gamma}{2(1 - \gamma\|x_0 - \bar{x}\|)^2 - 1} \|x_k - \bar{x}\|^2,$$

$k = 0, 1, \dots$. If, additionally, $(5 - \sqrt{17})/(4\gamma) < \kappa$, then $r_{\bar{x}} = (5 - \sqrt{17})/(4\gamma)$ is the biggest convergence radius.

To proving Theorem 15 we will need of the following results. The first one, gives us a condition that is easier to check than condition (7), whenever the functions under consideration are twice continuously differentiable, and its proof is similar to Lemma 5.3 of [1]. The second one, gives a relationship between the second derivatives f'' and ψ'' , which allow us to show that f and ψ satisfy (7), and its proof follows the same path of Lemma 22 of [19].

Lemma 16. *If $f : \Omega \subset X \rightarrow Y$ is an analytic function, $\bar{x} \in \Omega$ and $B(\bar{x}, 1/\gamma) \subset \Omega$, where γ is defined in (25), then for all $x \in B(\bar{x}, 1/\gamma)$, it holds that $\|f''(x)\| \leq 2\gamma/(1 - \gamma\|x - \bar{x}\|)^3$.*

Lemma 17. *Let X, Y be Banach spaces, $\Omega \subseteq X$ be an open set, $f : \Omega \rightarrow Y$ be twice continuously differentiable. Let $\bar{x} \in \Omega$, $R > 0$ and $\kappa = \sup\{t \in [0, R) : B(\bar{x}, t) \subset \Omega\}$. Let $\lambda > 0$ and $\psi : [0, R) \rightarrow \mathbb{R}$ be twice continuously differentiable such that $\lambda\|f''(x)\| \leq \psi''(\|x - \bar{x}\|)$, for all $x \in B(\bar{x}, \kappa)$, then f and ψ satisfy (7).*

[Proof of Theorem 15]. Let $\psi : [0, 1/\gamma) \rightarrow \mathbb{R}$ be defined by $\psi(t) = t/(1 - \gamma t) - 2t$. It is easy to see that ψ is analytic and $\psi(0) = 0$, $\psi'(t) = 1/(1 - \gamma t)^2 - 2$, $\psi'(0) = -1$, $\psi''(t) = 2\gamma/(1 - \gamma t)^3$. Thence, ψ satisfies **h1** and **h2**. Now, we combine Lemma 17 with Lemma 16, to conclude that f and ψ satisfy (7). The constants, ν , ρ and r , as defined in Theorem 4, satisfy

$$\rho = \frac{5 - \sqrt{17}}{4\gamma} < \nu = \frac{\sqrt{2} - 1}{\sqrt{2}\gamma} < \frac{1}{\gamma}, \quad r = \min \left\{ \kappa, \frac{5 - \sqrt{17}}{4\gamma} \right\}.$$

Moreover, $\psi(\rho)/(\rho\psi'(\rho)) - 1 = 1$ and $\psi(0) = \psi(1/(2\gamma)) = 0$ and $\psi(t) < 0$ for $t \in (0, 1/(2\gamma))$. Also, $\{t_k\}$ satisfy

$$t_{k+1}/t_k^2 = \frac{\gamma}{2(1 - \gamma t_k)^2 - 1} < \frac{\gamma}{2(1 - \gamma\|x_0 - \bar{x}\|)^2 - 1}, \quad k = 0, 1, \dots$$

Therefore, the result follows by applying the Theorem 4. □

5 Final remarks

In this paper, under a general majorant condition, we present a new local convergence analysis of the Newton's method for solving the generalized equation (1). Our approach is based in the Banach Perturbation Lemma obtained by S. M. Robinson in [30, Theorem 2.4] and used by Josephy in his

Ph.D thesis [23]. The majorant condition allow to unify several convergence results pertaining to Newton’s method. Besides, following the same idea of this paper, as future works, we propose to study the inexact Newton’s method for solving the problem (1) described by

$$(f(x_k) + f'(x_k)(x_{k+1} - x_k) + F(x_{k+1})) \cap R_k(x_k, x_{k+1}) \neq \emptyset, \quad k = 0, 1, \dots,$$

where $R_k : X \times X \rightrightarrows Y$ is a sequence of set-valued mappings with closed graphs, in order to support computational implementations of the method. Furthermore, it would be interesting to study the approach of this paper under a weak assumption than strong regularity, namely, the regularity metric; see [14].

References

- [1] F. Alvarez, J. Bolte, and J. Munier. A unifying local convergence result for newton’s method in riemannian manifolds. *Foundations of Computational Mathematics*, 8(2):197–226, 2008.
- [2] F. J. Aragón Artacho, A. Belyakov, A. L. Dontchev, and M. López. Local convergence of quasi-Newton methods under metric regularity. *Comput. Optim. Appl.*, 58(1):225–247, 2014.
- [3] F. J. Aragón Artacho, A. L. Dontchev, M. Gaydu, M. H. Geoffroy, and V. M. Veliov. Metric regularity of Newton’s iteration. *SIAM J. Control Optim.*, 49(2):339–362, 2011.
- [4] I. K. Argyros and S. Hilout. Improved local convergence of Newton’s method under weak majorant condition. *J. Comput. Appl. Math.*, 236(7):1892–1902, 2012.
- [5] L. Blum, F. Cucker, M. Shub, and S. Smale. *Complexity and real computation*. Springer-Verlag, New York, 1998. With a foreword by Richard M. Karp.
- [6] J. F. Bonnans. Local analysis of Newton-type methods for variational inequalities and nonlinear programming. *Appl. Math. Optim.*, 29(2):161–186, 1994.
- [7] R. Cibulka, A. Dontchev, and M. H. Geoffroy. Inexact Newton Methods and Dennis–Moré Theorems for Nonsmooth Generalized Equations. *SIAM J. Control Optim.*, 53(2):1003–1019, 2015.
- [8] J.-P. Dedieu, P. Priouret, and G. Malajovich. Newton’s method on Riemannian manifolds: convariant alpha theory. *IMA J. Numer. Anal.*, 23(3):395–419, 2003.
- [9] S. P. Dokov and A. L. Dontchev. Robinson’s strong regularity implies robust local convergence of Newton’s method. In *Optimal control (Gainesville, FL, 1997)*, volume 15 of *Appl. Optim.*, pages 116–129. Kluwer Acad. Publ., Dordrecht, 1998.
- [10] A. L. Dontchev. Local analysis of a Newton-type method based on partial linearization. In *The mathematics of numerical analysis (Park City, UT, 1995)*, volume 32 of *Lectures in Appl. Math.*, pages 295–306. Amer. Math. Soc., Providence, RI, 1996.
- [11] A. L. Dontchev. Local convergence of the Newton method for generalized equations. *C. R. Acad. Sci. Paris Sér. I Math.*, 322(4):327–331, 1996.
- [12] A. L. Dontchev. Uniform convergence of the newton method for aubin continuous maps. *Serdica Math. J.*, 22:385–398, 1996.

- [13] A. L. Dontchev and R. T. Rockafellar. Characterizations of strong regularity for variational inequalities over polyhedral convex sets. *SIAM J. Optim.*, pages 1087–1105, 1996.
- [14] A. L. Dontchev and R. T. Rockafellar. *Implicit functions and solution mappings*. Springer Monographs in Mathematics. Springer, Dordrecht, 2009. A view from variational analysis.
- [15] A. L. Dontchev and R. T. Rockafellar. Newton’s method for generalized equations: a sequential implicit function theorem. *Math. Program.*, 123(1, Ser. B):139–159, 2010.
- [16] A. L. Dontchev and R. T. Rockafellar. Convergence of inexact Newton methods for generalized equations. *Math. Program.*, 139(1-2, Ser. B):115–137, 2013.
- [17] O. P. Ferreira. Local convergence of Newton’s method in Banach space from the viewpoint of the majorant principle. *IMA J. Numer. Anal.*, 29(3):746–759, 2009.
- [18] O. P. Ferreira. Local convergence of Newton’s method under majorant condition. *J. Comput. Appl. Math.*, 235(5):1515–1522, 2011.
- [19] O. P. Ferreira, M. L. N. Gonçalves, and P. R. Oliveira. Local convergence analysis of the Gauss-Newton method under a majorant condition. *J. Complexity*, 27(1):111–125, 2011.
- [20] O. P. Ferreira, M. L. N. Gonçalves, and P. R. Oliveira. Convergence of the Gauss-Newton method for convex composite optimization under a majorant condition. *SIAM J. Optim.* 23(3): 1757–1783, 2013.
- [21] O. P. Ferreira and B. F. Svaiter. Kantorovich’s majorants principle for Newton’s method. *Comput. Optim. Appl.*, 42(2):213–229, 2009.
- [22] Z. Huang. The convergence ball of Newton’s method and the uniqueness ball of equations under Hölder-type continuous derivatives. *Comput. Math. Appl.*, 47(2-3):247–251, 2004.
- [23] N. Josephy. *Newton’s Method for Generalized Equations and the PIES Energy Model*. University of Wisconsin–Madison., 1979.
- [24] C. Li and W. Shen. Local convergence of inexact methods under the Hölder condition. *J. Comput. Appl. Math.*, 222(2):544–560, 2008.
- [25] C. Li and J. Wang. Newton’s method on Riemannian manifolds: Smale’s point estimate theory under the γ -condition. *IMA J. Numer. Anal.*, 26(2):228–251, 2006.
- [26] C. Li and K. F. Ng. Convergence analysis of the GaussNewton method for convex inclusion and convex-composite optimization problems. *J. Math. Anal. Appl.* 389(1): 469-485, 2012.
- [27] C. Li, J.-H. Wang, and J.-P. Dedieu. Smale’s point estimate theory for Newton’s method on Lie groups. *J. Complexity*, 25(2):128–151, 2009.
- [28] L. B. Rall. A note on the convergence of newton’s method. *SIAM Journal on Numerical Analysis*, 11(1):pp. 34–36, 1974.
- [29] S. M. Robinson. Extension of Newton’s method to nonlinear functions with values in a cone. *Numer. Math.*, 19:341–347, 1972.
- [30] S. M. Robinson. Strongly regular generalized equations. *Math. Oper. Res.*, 5(1):43–62, 1980.

- [31] G. N. Silva. Kantorovich's theorem on Newton's method for solving generalized equations under the majorant condition. *Applied Mathematics and Computation*, 286: 178–188, 2016.
- [32] G. N. Silva. Local convergence of Newton's method for solving generalized equations with monotone operator *ArXiv-eprints*, 2016.
- [33] S. Smale. Newton's method estimates from data at one point. In R. Ewing, K. Gross, and C. Martin, editors, *The Merging of Disciplines: New Directions in Pure, Applied, and Computational Mathematics*, pages 185–196. Springer New York, 1986.
- [34] J. F. Traub and H. Woźniakowski. Convergence and complexity of Newton iteration for operator equations. *J. Assoc. Comput. Mach.*, 26(2):250–258, 1979.
- [35] X. Wang. Convergence of Newton's method and inverse function theorem in Banach space. *Math. Comp.*, 68(225):169–186, 1999.
- [36] X. Wang, C. Li, and M.-J. Lai. A unified convergence theory for Newton-type methods for zeros of nonlinear operators in Banach spaces. *BIT Numerical Mathematics*, 42(1):206–213, 2002. cited By (since 1996) 2.
- [37] P. P. Zabrejko and D. F. Nguen. The majorant method in the theory of Newton-Kantorovich approximations and the Pták error estimates. *Numer. Funct. Anal. Optim.*, 9(5-6):671–684, 1987.